A transgenic pig model expressing a ZsGreen1 reporter across an extensive array of tissues

Amy T. Desaulniers[1], Rebecca A. Cederberg[1], Elizabeth P. Carreiro[1], Channabasavaiah B. Gurumurthy[2] and Brett R. White[1]

[1]University of Nebraska- Lincoln, Department of Animal Science, 3940 Fair Street, Lincoln, NE 68583-0908.

[2]University of Nebraska Medical Center, Department of Genetics, Cell Biology and Anatomy, 985815 Nebraska Medical Center, Omaha, NE 68198-5815

Amy T. Desaulniers, A224b Animal Science Building, 3940 Fair Street, Lincoln, NE 68583-0908, amy.desaulniers@huskers.unl.edu.

Rebecca A. Cederberg, A115 Animal Science Building, 3940 Fair Street, Lincoln, NE 68583-0908, rcederberg2@unl.edu.

Elizabeth P. Carreiro, A115 Animal Science Building, 3940 Fair Street, Lincoln, NE 68583-0908, carreiroelizabeth@gmail.com.

Channabasavaiah B. Gurumurthy, 985815 Nebraska Medical Center, Omaha, NE 68198-5815, cgurumurthy@unmc.edu.

B.R. White*, A224j Animal Science Building, 3940 Fair Street, Lincoln, NE 68583-0908, bwhite2@unl.edu, Phone: 402-472-6438.
*Corresponding author

ABSTRACT

Background: The advent of genetically engineered pig production has revealed a wide array of opportunities to enhance both biomedical and agricultural industries. One powerful method to develop these models is transgenesis; however, selection of a suitable promoter to drive transgene expression is critical. The cytomegalovirus (CMV) promoter is the most commonly used viral promoter as it robustly drives transgene expression in a ubiquitous nature. However, recent reports suggest that the level of CMV promoter activity is tissue-dependent in the pig. Therefore, the objective of this study was to quantify the activity of the CMV promoter in a wide range of porcine tissues. Swine harboring a CMV-ZsGreen1 transgene with a single integration site were utilized for this study. Thirty five tissue samples were collected from neonatal hemizygous (n = 3) and homozygous (n = 3) transgenic piglets and analyzed for ZsGreen1 abundance via immunoblot.

Results: ZsGreen1 was detected in all tissues examined; however, quantification revealed that ZsGreen1 protein levels were tissue-specific. Within organs of the digestive system, for example, ZsGreen1 was most abundant in the salivary gland, moderately produced in the esophagus and levels were lowest in the stomach. Interestingly, abundance of ZsGreen1 also differed within organ. For instance, levels were highest in the right ventricle compared with other chambers of the heart. There was no effect of transgene dose as ZsGreen1 expression patterns were similar between homozygous and hemizygous piglets.

Conclusions: Ultimately, these results elucidate the tissue-specific activity of the CMV promoter in the neonatal pig. Moreover, this model can serve as a useful tool for research applications requiring reporter gene activity in mammalian organs.

Keywords: CMV promoter, transgenic swine, ZsGreen1 reporter gene, gene editing

Background

Use of genetically engineered pig models has grown dramatically in recent years [1]. Transgenic swine are increasingly being utilized as animal models for human disease since they are anatomically, physiologically and phylogenetically more similar to humans than rodents [1, 2]. Consequently, successful xenotransplantation of transgenic pig organs to humans is eminent [3]. In addition to biomedical applications, genetically engineered pigs are also being developed to better understand porcine physiology and enhance the agricultural industry [4-7].

The first genetically engineered pig was generated in an effort to enhance growth rates and harbored a transgene containing the mouse metallothionein-I promoter fused to the human growth hormone gene [8]. Since then, new methodologies (e.g., homologous recombination, gene editing) have enhanced the ease and efficiency of modifying the pig genome [9]. However, incorporation of a transgene remains a prominent method to produce genetically engineered pigs [1]. The most critical aspect of transgenic animal production is promoter selection [10]. Depending on the research interest and application, promoter types include tissue-specific, cell-specific, inducible/conditional or ubiquitous [10]. Since ubiquitous promoters drive transgene expression in all tissues, they are a popular choice to address whole-body questions [10].

The human cytomegalovirus (CMV) immediate early promoter is one example of a ubiquitous promoter. It was discovered by Boshart et al. [11] and has since been heavily utilized to drive recombinant gene expression due to its constitutive and promiscuous nature [12]. These characteristics are directly related to its role in viral infection. Cytomegalovirus is a human herpesvirus which typically lays dormant in infected host cells [13]. Cellular infection quickly (<

1 h) leads to CMV gene expression via the interaction of host transcription factors with the CMV

55   promoter [13]. Gene products of this initial interaction mediate the activation of other promoters

within the CMV genome that facilitate virus replication and invasion [13]. Therefore, the CMV

promoter has evolved to contain binding sites for many ubiquitous transcription factors in order to

replicate in a broad range of host cells [12]. This discovery has had a major impact on the scientific

community; the CMV promoter is now the most commonly used ubiquitous promoter and it is also

60   considered the most robust [14].

The CMV promoter has been utilized to drive transgene expression in numerous vertebrate

animal models including: sheep [15-17], fish [14, 18], mice [19-22], rats [23, 24], chickens [25]

and pigs [4, 26-28]. In the pig, the CMV promoter was used to drive expression of green

fluorescent protein (GFP) [27]. As anticipated, expression of the transgene was detected in a wide

65   range of organs, including those derived from each of the 3 germ layers: skin (ectoderm), pancreas

(endoderm), and kidney (mesoderm). Despite apparent universal transgene expression [27], the

CMV promoter was preferentially active in exocrine vs. non-exocrine cells of the pig [26]. Given

the growing interest in the development of transgenic swine models, the objective of this study

was to quantify activity of the CMV promoter in a wide range of porcine tissues.

Methods

Animals

All animal procedures were conducted in accordance with the University of Nebraska-Lincoln

(UNL) Institutional Animal Care and Use Committee. Experiments were performed in the UNL

75   Animal Science Building (Lincoln, NE) and utilized transgenic swine that express ZsGreen1

controlled by the human CMV immediate early promoter; these animals were generated as

previously described [4]. In hemizygous animals, one copy of the CMV-ZsGreen1 transgene is stably integrated on chromosome 14 [4], and has remained transcriptionally active throughout the lifetime of the founder animal and in 10 subsequent generations.

80   Adult animals were housed individually with *ad libitum* access to water and fed approximately 2.5 kg of feed daily. Hemizygous CMV-ZsGreen1 transgenic sows were artificially inseminated with semen from non-littermate hemizygous transgenic boars and allowed to gestate to term. Following Mendelian inheritance, these matings yielded hemizygous, homozygous (2 copies of the transgene), and non-transgenic (control) littermates. After farrowing, piglets

85   remained with their dam to suckle *ad libitum*.

At 1 d of age, 3 littermate female piglets (control, hemizygous transgenic and homozygous transgenic) were selected from 3 different litters; each litter was produced from a different dam and sire mating to maximize genetic diversity. Initially, transgene status was assessed via evaluation of ZsGreen1 expression using an ultraviolet (UV) light and a Roscolux #15 filter

90   (deep straw; Rosco, Port Chester, NY).

Tissue collection

Piglets were euthanized via intracardiac injection with Euthasol (1 mL/4.53 kg BW; Delmarva Laboratories, Midlothian, VA) and stored at 4°C overnight prior to dissection. The following

95   morning, 35 organ samples generally representing 5 anatomical regions (brain, thoracic, digestive, renal and reproductive) were collected from each transgenic piglet. Isolated tissues included: cerebrum, cerebellum, spinal cord, anterior pituitary gland, posterior pituitary gland, hypothalamus, thymus, lymph node, the four chambers of the heart, lung, muscle, thyroid, salivary gland, esophagus, stomach, duodenum, jejunum, ileum, colon, gall bladder, spleen, pancreas, liver,

100   kidney (medulla and cortex), bladder, adrenal gland (medulla and cortex), ovary, uterus, oviduct,

and skin. Skin samples were collected from control littermate piglets to serve as a negative control

for ZsGreen1 immunoblotting. In addition, tail samples were collected from all piglets for

genotyping. All samples were frozen and stored at -80°C until analysis.

105   **Deoxyribonucleic acid extraction**

Genomic deoxyribonucleic acid (DNA) was isolated from tail samples using an enzymatic

digestion and phenol/chloroform extraction. Briefly, a tail slice (approximately 1 mm) was

incubated overnight in a lysing solution [50 mM Tris pH 8.0, 150 mM NaCl, 10 mM EDTA, 1%

sodium dodecyl sulfate (SDS), 0.1 mg/ml Proteinase K (Fisher Bioreagents, Fairlawn, NJ), 1% β-

110   mercaptoethanol] at 55°C. The samples were extracted twice with a 25:24:1

phenol:chloroform:isoamyl alcohol solution and once with a 24:1 chloroform:isoamyl alcohol

solution before precipitating the DNA with isopropanol. DNA was subsequently resuspended in

nuclease-free water.

115   **Genotyping**

Since the transgene integration site was previously identified in this model [4], copy number was

evaluated via conventional polymerase chain reaction (PCR) using 3 primers. The forward (F) and

reverse (R) primers were designed to flank the insertion site of the transgene (T) and an additional

reverse primer (RT) aligning the transgene was included (F - GCA ACC TCT TCG ACA CTC

120   CA; R - AGC TAC CAG GGA ACA AAG CC; RT - GGT TTC CCT TTC GCT TTC AAG T).

An MJ Research PTC-200 Thermocycler (Waltham, MA) was used for PCR reactions with the

following conditions: 1 X *Taq* Buffer A (Fisher Bioreagents) with a final concentration of 2.0 mM

MgCl$_2$, 200 µM dNTPs, 300 nM of each primer and 1.25 U of *Taq* DNA Polymerase (Fisher Bioreagents). Cycling conditions were 95ºC for 2 min, followed by 35 cycles of 95ºC for 30 sec,

125   56ºC for 30 sec and 72ºC for 60 sec and a final extension of 72ºC for 10 min. The resultant PCR products were subjected to electrophoresis on a 1% agarose gel. Products generated from the F and R primer pair were 627 bp in length and reflected a wild type chromosome 14. Products from the F and RT primer pair resulted in an 847 bp product, and indicating that the transgene was present.

Protein preparation

Protein extraction and sample preparation was performed as described previously [29]. Protein was extracted by homogenization of tissue samples in radioimmunoprecipitation assay (RIPA) buffer (1 ml/100 mg; 20 mM Tris, 137 mM NaCl, 10% glycerol, 1% NP40, 0.1% SDS, 0.5%

135   deoxycholic acid, 2 mM EDTA, 1mM PMSF, 1% protease inhibitor cocktail and 1% phosphatase inhibitor cocktail) using a Biospec Tissue Tearor (Bartlesville, OK). Protein concentrations were quantified using a bicinchoninic acid (BCA) assay (Pierce Biotechnology, Rockford, IL) according to the manufacturer's instructions. Extracted protein was mixed with 4X loading dye [2% Tris (pH 6.8), 28% glycerol, 20% SDS and Orange G] containing dithiothreitol (DTT; 100 mM) and frozen

140   at -80°C until immunoblotting.

Immunoblotting

Immunoblotting was performed as described previously [29]. Tissue samples were segregated by anatomical region (brain, thoracic, digestive, renal and reproductive) and protein (10 or 20 µg)

145   was separated using SDS-PAGE (15%) before transfer to an Immobilon-FL polyvinylidene

difluoride (PVDF) membrane (Millipore, Billerica, MA) pre-soaked in methanol. After electrophoretic transfer, nonspecific binding was blocked by incubation of blots in Odyssey Blocking Buffer (cat #: 927-40100; LI-COR Biosciences, Lincoln, NE) for approximately 3 h at room temperature. Membranes were then incubated with a mouse monoclonal primary antibody

150 directed against ZsGreen1 (1:1000; cat #: 632598; Clontech, Mountain View, CA), diluted in Odyssey Blocking Buffer with 0.05% Tween-20 and shaking at 4ºC overnight. The following morning, membranes were rinsed in Tris-buffered saline with 0.05% Tween-20 (TBS-T) prior to incubation with a goat anti-mouse secondary antibody (1:15000; IRDye 680; #926-32220; LI-COR Biosciences) in Odyssey Blocking Buffer plus 0.05% Tween-20 and 0.025% SDS for 1 h at room

155 temperature. Blots were rinsed with TBS-T to remove excess secondary antibody and briefly rinsed in TBS to remove Tween-20 prior to imaging.

Quantification of ZsGreen1 abundance

Blots were scanned on an Odyssey Infrared Imager (700 channel; intensity 5.0; 169 micron

160 resolution; LI-COR Biosciences) and converted to greyscale for analysis. Band density was quantitated with Odyssey imaging software (version 2.1; LI-COR Biosciences). Since the abundance of common loading controls (e.g., β-actin) differs across tissue types, total protein levels within each lane were used as the loading control according to Eaton et al. [30]. After ZsGreen1 imaging, membranes were stained with GelCode Blue Safe Protein Stain

165 (ThermoScientific, Rockford, IL) for approximately 1 h and then rinsed with a 50% methanol and 1% acetic acid solution for 1 h. Blots were reimaged (700 channel; intensity 1.0; 169 micron resolution; LI-COR Biosciences), converted to greyscale and quantitated using Odyssey imaging software (version 2.1; LI-COR Biosciences) to determine the relative quantity of protein present

in each sample. Data are presented as the relative band density of ZsGreen1 normalized by the

170   total protein levels within the entire lane as described previously [30].

Statistical Analysis

Analyses were performed using the Statistical Analysis System (SAS; Cary, NC; version 9.4).

Analysis of variance (ANOVA) was performed via the MIXED procedure of SAS with tissue type

175   as the fixed effect. Blot and litter were random effects. $P < 0.05$ was considered significant,

whereas $P < 0.10$ was considered a tendency. Results are presented as least squares means

(LSMEANS) ± the standard error of the mean (SEM). No effect of genotype (hemizygous versus

homozygous) on ZsGreen1 abundance was observed ($P > 0.10$).

180   **Results**

Identification of hemizygous and homozygous CMV-ZsGreen1 transgenic piglets

Presence of ZsGreen1 in the skin of neonatal transgenic piglets was detected visually (Figure 1A)

and via immunoblot (Figure 1B), indicative of CMV promoter activity. Even without the UV light,

transgenic animals were identifiable via the yellow hue of their skin (data not shown), which has

185   been previously noted in GFP pigs [26]. The genotype of the animals was confirmed by

conventional PCR (Figure 1D); a schematic depicting the genotyping method is available in Figure

1C. ZsGreen1 was not detected in the skin of control piglets (Figure 1A & 1B).

ZsGreen1 fluorescence within tissues of CMV-ZsGreen1 transgenic piglets

190   ZsGreen1 fluorescence was visually detectable within numerous internal organs of transgenic pigs

under UV light; fluorescence was absent in wild-type pig organs (Figure 2A). Fluorescence

appeared robust in the intestines (Figure 2A; middle panel), cerebellum (Figure 2B; left panel), anterior pituitary gland (Figure 2B; middle panel) and heart (Figure 2B; right panel), but modest within the stomach (Figure 2A; middle panel). ZsGreen1 was not visually detectable in the cerebrum (Figure 2B; left panel), posterior pituitary gland (Figure 2B; middle panel), liver or spleen (Figure 2A; middle panel). Differential fluorescent patterns were observed within organs as well. In the brain, ZsGreen1 fluorescence was intense within the cerebellum and not visually apparent in the cerebrum (Figure 2B; left panel). In addition, expression of ZsGreen1 appeared greatest in the right ventricle compared with other regions of the heart (Figure 2B; right panel).

Abundance of ZsGreen1 in hemizygous and homozygous porcine organs

Neural and endocrine tissues of the brain

There was an effect of tissue type on ZsGreen1 production levels within the brain and nervous system of hemizygous piglets ($P = 0.0002$). Abundance of ZsGreen1 was greatest within the anterior pituitary gland, followed by the cerebellum ($P < 0.05$; Figure 3A). ZsGreen1 levels were lowest in the cerebrum, spinal cord, hypothalamus and posterior pituitary gland ($P < 0.05$), which were not different from each other ($P > 0.05$; Figure 3A). In homozygous piglets, there was also an effect of tissue type on ZsGreen1 abundance ($P = 0.0396$). The anterior pituitary gland expressed more ZsGreen1 than the cerebrum, spinal cord, hypothalamus or posterior pituitary gland ($P < 0.05$; Figure 3B), but was not different from the cerebellum ($P > 0.05$; Figure 3B). ZsGreen1 levels were similar between the cerebellum, cerebrum, spinal cord, hypothalamus and posterior pituitary gland ($P > 0.05$; Figure 3B).

Thoracic organs

215   An effect of tissue type on ZsGreen1 protein abundance was observed among 9 thoracic organs of hemizygous piglets ($P < 0.0001$). The right ventricle produced the most ZsGreen1 in the thoracic region, even when compared with the other chambers of the heart ($P < 0.05$; Figure 4A). ZsGreen1 levels in the right ventricle were greater ($P < 0.05$) than the thyroid, right atrium, left atrium, left ventricle, lung, thymus or lymph node, which were similar ($P > 0.05$; Figure 4A). ZsGreen1 levels

220   within skeletal muscle were intermediate ($P < 0.05$; Figure 4A). In homozygous piglets, there was no effect of tissue type on expression of ZsGreen1 among thoracic organs ($P = 0.3326$; Figure 4B). Although, values for ZsGreen1 abundance were numerically higher in the right ventricle and skeletal muscle compared with other tissues.

225   *Digestive Organs*

ZsGreen1 protein levels differed across 11 digestive organs of hemizygous piglets ($P < 0.0001$). Levels were greatest in the salivary gland and lowest in the stomach, pancreas, duodenum, jejunum, ileum, colon, spleen, gall bladder, and liver, which were not different ($P > 0.10$). Protein amounts were intermediate in the esophagus ($P < 0.05$; Figure 5A), with the exception of the

230   duodenum which was similar ($P > 0.05$). In homozygous piglets, there was also an effect of tissue type on ZsGreen1 protein abundance ($P = 0.0212$). Expression was greatest in the salivary gland ($P < 0.05$; Figure 5B) compared with the other 10 organs, which were similar ($P > 0.05$; Figure 5B).

235   *Renal Organs*

An effect of tissue type was observed between 5 renal tissues of hemizygous piglets ($P = 0.0083$). The adrenal medulla produced more ZsGreen1 than the adrenal cortex, kidney medulla, kidney

cortex or bladder ($P < 0.05$; Figure 6A), which were similar to each other ($P > 0.05$). In homozygous animals, there tended to be a difference in ZsGreen1 abundance across renal tissues

240 ($P = 0.0620$). Levels were highest in the adrenal medulla and lowest in the kidney medulla, kidney cortex, and bladder ($P < 0.05$; Figure 6B), which were not different from one another ($P > 0.05$). ZsGreen1 levels were similar between the adrenal cortex and all other organs ($P > 0.05$; Figure 6B).

245 *Reproductive organs*

No differences in ZsGreen1 expression were detected between the ovary, oviduct, and uterus of hemizygous or homozygous piglets ($P > 0.05$; Figure 7A and 7B).

Discussion

250 Our study provides a quantitative, spatial analysis of tissue-specific ZsGreen1 protein abundance in swine containing a CMV-ZsGreen1 transgene. In accord with the ubiquitous nature of the CMV promoter, ZsGreen1 protein was detected in every tissue examined via immunoblotting. Consistent with this, another report from our laboratory indicated that exosomes isolated from the milk of hemizygous females are labeled with ZsGreen1 [31]. Our results,

255 however, demonstrate that the level of CMV promoter activity was variable across organs of the neonatal piglet. Others have observed similar results in different animal models expressing a transgene driven by the CMV promoter [14, 25, 26, 32]. Based upon these results, several hypotheses regarding the differential activity of the CMV promoter *in vivo* have arisen. The activity of the CMV promoter within a given cell type may be influenced by integration site [32];

260 namely, CMV promoter activation may only occur in transcriptionally active regions of the

genome [12]. Stochastic silencing can occur when a transgene is integrated near a heterochromatic region, such as a centromere [33], rendering the promoter inaccessible to transcriptional machinery [34]. Stochastic silencing of transgenes can also occur when multiple copies of the transgene integrate in tandem [35], due to local formation of heterochromatin [34]. In support of this hypothesis, Furth et al. [32] observed that two different transgenic mouse lines had marked differences in CMV promoter activity across tissues of the body, despite harboring the same transgene. Likewise, transgenic animals harboring a transgene, but lacking expression of the gene, have been generated [36]. These results could also be due to promoter methylation, which has been shown to silence transgenes *in vitro* and *in vivo* [15, 16, 21, 37]. Conversely, Vasey et al. (2009) hypothesized that integration site does not readily affect CMV promoter activity because a similar activity profile was detected in pigs and chickens harboring the same transgene, albeit integrated in different chromosomal regions [26, 27]. The activity patterns were also similar in multiple transgenic chicken lines expressing different reporter genes directed by the CMV promoter [25]. Our data support these results as a similar expression pattern was observed in CMV-ZsGreen1 pigs, despite a different integration site and transgene.

In addition to chromatin status, the repertoire of transcription factors and co-regulators present within the cell dictate CMV promoter activity [12]. Mella-Alvarado et al. (2013) hypothesized that CMV promoter activity is robust in the organs that CMV infects most voraciously (brain, eye, spinal cord, pancreas, kidney, testis, ovary, skin, cartilage, and skeletal muscle), reflective of optimal transcription factor expression. Moreover, Furth et al. [32] suggested that certain transcription factors may only be able to interact with euchromatin, whereas others retain functionality with heterochromatin [32]. In exocrine glands, Vasey et al. [26] proposed that the transcription factor, nuclear factor kappa-light-chain-enhancer of activated B cells (NF-kB), is

a major driver of CMV promoter activity. NF-kB is highly expressed in exocrine cells, classically

285 regulates the expression of proinflammatory genes [38], activates in response to cellular insult

such as viral infection [38], and is a potent regulator of the CMV promoter [12, 39-41]. In addition

to NF-kB, numerous transcription factors have been implicated in the regulation of the CMV

promoter, including ying yang 1 [12], activator protein 1 and its major subunits (c-Fos and c-Jun)

[41], p53, cAMP response element binding nuclear transcription factor/activating transcription

290 factor, retinoic acid receptor [42], specificity protein 1 [11], nuclear factor I, serum response factor,

Elk-1 and CCAAT/enhancer-binding protein [13]. Thus, it is apparent that the CMV promoter has

adapted to respond to a variety of cell signals in order to allow for the promiscuous infection of

host cells [12].

Consistent with elevated activity of the CMV promoter in exocrine glands, ZsGreen1

295 protein levels were increased in the salivary glands of hemizygous and homozygous transgenic

piglets, as previously noted in CMV-GFP pigs [26]. Enhanced CMV promoter activity was

reported in other exocrine tissues of the pig, including the glandular epithelia of the snout,

sebaceous glands of the skin and Harderian gland of the eye [26]. In the exocrine pancreas,

however, we did not detect elevated ZsGreen1 levels in either hemizygous or homozygous

300 transgenic piglets, conflicting with other reports of increased CMV promoter activity within the

porcine pancreas [26, 27]. This discrepancy could be related to age-specific activity levels of the

CMV promoter, as noted in zebrafish [14]. In the current study, neonatal (1 d old) piglets were

utilized, whereas others examined CMV promoter activity in the pancreases of older (2 mo to 1

yr) pigs [26, 27]. Likewise, our laboratory detected robust ZsGreen1 abundance in the pancreas of

305 1 mo old piglets (unpublished data). Interestingly, the exocrine pancreas is largely quiescent in the

neonatal piglet [43-47]; production of pancreatic juices and enzymes increase dramatically after

weaning [43-45]. In the liver, another exocrine organ that secretes bile made by hepatocytes [48], we also detected little CMV promoter activity. Others have reported similar levels of CMV promoter activity in porcine hepatocytes [26] and chicken liver [25]. In contrast, GFP was robustly

310   driven by the CMV promoter in hepatocytes of zebrafish [14]. Thus, it appears that the CMV promoter is not uniformly effective in all exocrine organs and can vary in activity across species.

In the current study, abundance of ZsGreen1 was also compared between homozygous and hemizygous animals. Based on our visual observation of ZsGreen1 production in the skin of hemizygous and homozygous piglets (Figure 1A), we hypothesized that an additional transgene

315   would yield more ZsGreen1 protein in the organs of homozygous animals. However, no genotype effect was detected based upon quantification of immunoblots for ZsGreen1. These results may be due to monoallelic expression, which frequently occurs in autosomes [49] and has also been described in other transgenic animal models [34, 50]. Conversely, translation of ZsGreen1 transcripts may be impaired in homozygous animals. For example, post-transcriptional silencing

320   of high expressing transgenes is common in homozygous, but not hemizygous, plants [51-54] and appears to be mediated by short interfering RNA [51]. Many have hypothesized that doubling transcription of a transgene yields copious mRNA transcripts, which breaches an unknown threshold that triggers cellular processes to suppress transgene translation [52, 55-58]. Regardless of the mechanism, transgene copy number did not significantly affect production of ZsGreen1

325   protein in this study.

To our knowledge, these animals represent the first pig model to express ZsGreen1 (Clontech, Palo Alto, CA). This protein, originally discovered by Matz et al. [59], is derived from Anthozoa coral reef (*Zoanthus* species). The variant, ZsGreen1, is a modification of the original protein (zFP506) to maximize expression, solubility and prompt chromophore maturation

330   (Clontech, personal communication). Interestingly, it is both structurally and functionally distinct from other fluorescent proteins, like GFP [59, 60]. Although both fluorescent proteins are configured in a β-barrel structure [59], ZsGreen1 exists as a tetramer, whereas GFP (derived from jellyfish) is a monomer that frequently dimerizes for solubility [61]. Interestingly, ZsGreen1 is only 26% homologous to GFP and each ZsGreen1 monomer is 231 amino acids in length with a

335   predicted molecular weight of approximately 26 kDa [59]. Functionally, ZsGreen1 displays extremely bright fluorescence and high signal-to-noise ratio; in fact, yeast cells overexpressing ZsGreen exhibited 8.6-fold more fluorescence than those expressing GFP [60]. In another study, ZsGreen1 had the greatest relative fluorescence compared with 5 other fluorescent proteins, including GFP [59].

340   Notably, this ZsGreen1 swine model could serve as a useful tool for genome editing applications. This model is well suited for re-engineering the transgene locus since expression of the transgene is readily detectable, the transgene is present in a single copy and the integration site has been identified [4]. For example, this swine line could be used for testing somatic cell gene editing using CRISPR/Cas9 systems or base editor nucleases by targeting the ZsGreen1 transgene

345   sequence. In addition, the model can be a useful tool for testing germline genome editing approaches *in vivo* such as Genome editing via Oviductal Nucleic Acids Delivery (GONAD), initially established in the mouse [62]. Finally, this pig model can be used for re-engineering the transgenic CMV-ZsGreen1 locus to make it suitable for other applications. For example, the ZsGreen1 coding sequence can be disrupted to create a frameshift mutation in ZsGreen1, creating

350   a model to serve as a reporter for homology directed repair genome editing studies. Alternatively, a floxed red fluorescent protein could be inserted upstream of the ZsGreen1 cassette to serve as a Cre recombinase reporter, similar to what is popularly used in mouse Cre-LoxP studies [63]. Re-

engineering of the locus can be achieved via Easi-CRISPR, a highly efficient genome engineering method recently described in mice [64, 65].

Conclusions

This study demonstrates that the CMV promoter is ubiquitously active in all porcine tissues types examined; however, the level of activity is tissue-specific. The CMV promoter potently drives transgene expression in many exocrine (e.g., salivary gland) and non-exocrine organs (e.g., skin,

360 anterior pituitary gland and muscle) of the neonatal pig. In certain organs (e.g., liver, kidney), however, activity of the CMV promoter is less robust. Ultimately, this report elucidates the tissue-specific activity of the human CMV promoter in swine and provides critical information for researchers seeking an effective promoter to drive transgene expression in the neonatal pig. Moreover, these animals represent a novel resource for scientists in need of ZsGreen1-labeled

365 porcine organs. Lastly, this CMV-ZsGreen1 swine model will also serve as a valuable tool to advance genome editing research.

Abbreviations

CMV: cytomegalovirus promoter; DNA: deoxyribonucleic acid; GFP: green fluorescent protein;

370 NF-kB: nuclear factor kappa-light-chain-enhancer of activated B cells; LSMEANS: least squares means; PCR: polymerase chain reaction; SDS: sodium dodecyl sulfate; SEM: standard error of the mean; UV: ultraviolet.

Declarations

375 **Acknowledgements**

The authors wish to thank Ginger Mills for pig husbandry as well as Scott Kurz, Guilherme Cezar, Kyle Regan and Amanda Lambrecht for technical assistance. The authors would also like to express gratitude to Lelanya Yates for aiding in the dissections.

380 **Funding**

This project is based on research that was partially supported by the Nebraska Agricultural Experiment Station with funding from the Hatch Multistate Research capacity funding program (Accession Number 1011129) from the USDA National Institute of Food and Agriculture. The funding agency had no role in study design, data collection, interpretation, or manuscript
385 generation.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

AD, RC and BW conceived of the study and designed the experiment. AD, RC and EP collected the data. AD analyzed the data. AD, RC, GC and BW interpreted the data. AD and RC drafted the manuscript. All authors read, revised and approved the final manuscript.

Ethics approval and consent to participate

All animal procedures were conducted in accordance with the University of Nebraska-Lincoln Institutional Animal Care and Use Committee.

400 **Consent for publication**

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. E. M. Walters, K. D. Wells, E. C. Bryda, S. Schommer and R. S. Prather. Swine models, genomic tools and services to enhance our understanding of human health and diseases. Lab Anim (NY). 2017;46(4):167-172.

410 2. A. L. Archibald, L. Bolund, C. Churcher, M. Fredholm, M. A. Groenen, B. Harlizius, K. T. Lee, D. Milan, J. Rogers, M. F. Rothschild, H. Uenishi, J. Wang, L. B. Schook and Swine Genome Sequencing Consortium. Pig genome sequence--analysis and publication strategy. BMC Genomics. 2010;11:438-2164-11-438.

3. D. K. C. Cooper, R. Gaston, D. Eckhoff, J. Ladowski, T. Yamamoto, L. Wang, H. Iwase, H. 415 Hara, M. Tector and A. J. Tector. Xenotransplantation-the current status and prospects. Br Med Bull. 2017;125(1):5-14.

4. A. T. Desaulniers, R. A. Cederberg, G. A. Mills, C. A. Lents and B. R. White. Production of a gonadotropin-releasing hormone 2 receptor knockdown (GNRHR2 KD) swine line. Transgenic Res. 2017;26(4):567-575.

420 5. G. T. Bleck, B. R. White, D. J. Miller and M. B. Wheeler. Production of bovine alpha-lactalbumin in the milk of transgenic pigs. J Anim Sci. 1998;76(12):3072-3078.

6. K. M. Whitworth, R. R. Rowland, C. L. Ewen, B. R. Trible, M. A. Kerrigan, A. G. Cino-Ozuna, M. S. Samuel, J. E. Lightner, D. G. McLaren, A. J. Mileham, K. D. Wells and R. S. Prather. Gene-edited pigs are protected from porcine reproductive and respiratory syndrome 425 virus. Nat Biotechnol. 2016;34(1):20-22.

7. R. S. Prather, K. D. Wells, K. M. Whitworth, M. A. Kerrigan, M. S. Samuel, A. Mileham, L. N. Popescu and R. R. R. Rowland. Knockout of maternal CD163 protects fetuses from infection with porcine reproductive and respiratory syndrome virus (PRRSV). Sci Rep. 2017;7(1):13371.

8. R. E. Hammer, V. G. Pursel, C. E. Rexroad Jr, R. J. Wall, D. J. Bolt, K. M. Ebert, R. D. Palmiter and R. L. Brinster. Production of transgenic rabbits, sheep and pigs by microinjection. Nature. 1985;315(6021):680-683.

9. K. D. Wells and R. S. Prather. Genome-editing technologies to improve research, reproduction, and production in pigs. Mol Reprod Dev. 2017;84(9):1012-1017.

10. Y. Hakamata and E. Kobayashi. Inducible and conditional promoter systems to generate transgenic animals. Methods Mol Biol. 2010;597:71-79.

11. M. Boshart, F. Weber, G. Jahn, K. Dorsch-Hasler, B. Fleckenstein and W. Schaffner. A very strong enhancer is located upstream of an immediate early gene of human cytomegalovirus. Cell. 1985;41(2):521-530.

12. A. J. Brown, B. Sweeney, D. O. Mainwaring and D. C. James. NF-kappaB, CRE and YY1 elements are key functional regulators of CMV promoter-driven transient gene expression in CHO cells. Biotechnol J. 2015;10(7):1019-1028.

13. S. Landolfo, M. Gariglio, G. Gribaudo and D. Lembo. The human cytomegalovirus. Pharmacol Ther. 2003;98(3):269-297.

14. V. Mella-Alvarado, A. Gautier, F. Le Gac and J. J. Lareyre. Tissue and cell-specific transcriptional activity of the human cytomegalovirus immediate early gene promoter (UL123) in zebrafish. Gene Expr Patterns. 2013;13(3-4):91-103.

15. C. Liu, L. Wang, W. Li, X. Zhang, Y. Tian, N. Zhang, S. He, T. Chen, J. Huang and M. Liu. Highly efficient generation of transgenic sheep by lentivirus accompanying the alteration of methylation status. PLoS One. 2013;8(1):e54614.

16. B. Duan, L. Cheng, Y. Gao, F. X. Yin, G. H. Su, Q. Y. Shen, K. Liu, X. Hu, X. Liu and G. P. Li. Silencing of fat-1 transgene expression in sheep may result from hypermethylation of its driven cytomegalovirus (CMV) promoter. Theriogenology. 2012;78(4):793-802.

17. Y. Tian, W. Li, L. Wang, C. Liu, J. Lin, X. Zhang, N. Zhang, S. He, J. Huang, B. Jia and M. Liu. Expression of 2A peptide mediated tri-fluorescent protein genes were regulated by epigenetics in transgenic sheep. Biochem Biophys Res Commun. 2013;434(3):681-687.

18. M. Kinoshita, T. Morita, H. Toyohara, T. Hirata, M. Sakaguchi, M. Ono, K. Inoue, Y. Wakamatsu and K. Ozato. Transgenic medaka overexpressing a melanin-concentrating hormone exhibit lightened body color but no remarkable abnormality. Mar Biotechnol (NY). 2001;3(6):536-543.

19. P. Loser, G. S. Jennings, M. Strauss and V. Sandig. Reactivation of the previously silenced cytomegalovirus major immediate-early promoter in the mouse liver: involvement of NFkappaB. J Virol. 1998;72(1):180-190.

20. J. L. Ewart, M. F. Cohen, R. A. Meyer, G. Y. Huang, A. Wessels, R. G. Gourdie, A. J. Chin, S. M. Park, B. O. Lazatin, S. Villabon and C. W. Lo. Heart and neural tube defects in transgenic mice overexpressing the Cx43 gap junction gene. Development. 1997;124(7):1281-1292.

21. M. K. Dyck, M. Ouellet, M. Gagn, D. Petitclerc, M. A. Sirard and F. Pothier. Testes-specific transgene expression in insulin-like growth factor-I transgenic mice. Mol Reprod Dev. 1999;54(1):32-42.

22. G. Villuendas, A. Gutierrez-Adan, A. Jimenez, C. Rojo, E. R. Roldan and B. Pintado. CMV-driven expression of green fluorescent protein (GFP) in male germ cells of transgenic mice and its effect on fertility. Int J Androl. 2001;24(5):300-305.

23. B. Charreau, L. Tesson, J. Buscail, J. P. Soulillou and I. Anegon. Analysis of human CD59 tissue expression directed by the CMV-IE-1 promoter in transgenic rats. Transgenic Res. 1996;5(6):443-450.

24. E. S. Green, K. G. Rendahl, S. Zhou, M. Ladner, M. Coyne, R. Srivastava, W. C. Manning and J. G. Flannery. Two animal models of retinal degeneration are rescued by recombinant adeno-associated virus-mediated production of FGF-5 and FGF-18. Mol Ther. 2001;3(4):507-515.

25. M. J. McGrew, A. Sherman, F. M. Ellard, S. G. Lillico, H. J. Gilhooley, A. J. Kingsman, K. A. Mitrophanous and H. Sang. Efficient production of germline transgenic chickens using lentiviral vectors. EMBO Rep. 2004;5(7):728-733.

26. D. B. Vasey, S. G. Lillico, H. M. Sang, T. J. King and C. B. Whitelaw. CMV enhancer-promoter is preferentially active in exocrine cells in vivo. Transgenic Res. 2009;18(2):309-314.

27. C. B. Whitelaw, P. A. Radcliffe, W. A. Ritchie, A. Carlisle, F. M. Ellard, R. N. Pena, J. Rowe, A. J. Clark, T. J. King and K. A. Mitrophanous. Efficient generation of transgenic pigs using equine infectious anaemia virus (EIAV) derived vector. FEBS Lett. 2004;571(1-3):233-236.

28. Q. Kong, M. Wu, Y. Huan, L. Zhang, H. Liu, G. Bou, Y. Luo, Y. Mu and Z. Liu. Transgene Expression Is Associated with Copy Number and Cytomegalovirus Promoter Methylation in Transgenic Pigs. PLoS One. 2009;4(8):e6679.

29. A. T. Desaulniers, R. A. Cederberg, G. A. Mills, J. J. Ford, C. A. Lents and B. R. White. LH-independent testosterone secretion is mediated by the interaction between GnRH2 and its receptor within porcine testes. Biol Reprod. 2015;93(2):45.

30. S. L. Eaton, S. L. Roche, M. Llavero Hurtado, K. J. Oldknow, C. Farquharson, T. H. Gillingwater and T. M. Wishart. Total protein analysis as a reliable loading control for quantitative fluorescent Western blotting. PLoS One. 2013;8(8):e72457.

31. S. Manca, B. Upadhyaya, E. Mutai, A. T. Desaulniers, R. A. Cederberg, B. R. White and J. Zempleni. Milk exosomes are bioavailable and distinct microRNA cargos have unique tissue distribution patterns. Sci Rep. 2018;8(1):11321-018-29780-1.

500   32. P. A. Furth, L. Hennighausen, C. Baker, B. Beatty and R. Woychick. The variability in activity of the universally expressed human cytomegalovirus immediate early gene 1 enhancer/promoter in transgenic mice. Nucleic Acids Res. 1991;19(22):6205-6208.

33. K. W. Dobie, M. Lee, J. A. Fantes, E. Graham, A. J. Clark, A. Springbett, R. Lathe and M. McClenaghan. Variegated transgene expression in mouse mammary gland is determined by the
505   transgene integration locus. Proc Natl Acad Sci U S A. 1996;93(13):6659-6664.

34. S. P. Chang, M. L. Opsahl, C. B. Whitelaw, S. D. Morley and J. D. West. Relative transgene expression frequencies in homozygous versus hemizygous transgenic mice. Transgenic Res. 2013;22(6):1143-1154.

35. D. Garrick, S. Fiering, D. I. Martin and E. Whitelaw. Repeat-induced gene silencing in
510   mammals. Nat Genet. 1998;18(1):56-59.

36. R. D. Palmiter and R. L. Brinster. Germ-line transformation of mice. Annu Rev Genet. 1986;20:465-499.

37. S. Prosch, J. Stein, K. Staak, C. Liebenthal, H. D. Volk and D. H. Kruger. Inactivation of the very strong HCMV immediate early promoter by DNA CpG methylation in vitro. Biol Chem
515   Hoppe Seyler. 1996;377(3):195-201.

38. F. Chen, V. Castranova, X. Shi and L. M. Demers. New insights into the role of nuclear factor-kappaB, a ubiquitous transcription factor in the initiation of diseases. Clin Chem. 1999;45(1):7-17.

39. X. Liu, J. Yuan, A. W. Wu, P. W. McGonagill, C. S. Galle and J. L. Meier. Phorbol ester-
520   induced human cytomegalovirus major immediate-early (MIE) enhancer activation through PKC-delta, CREB, and NF-kappaB desilences MIE gene expression in quiescently infected human pluripotent NTera2 cells. J Virol. 2010;84(17):8495-8508.

40. M. Ramanathan, G. Hasko and S. J. Leibovich. Analysis of signal transduction pathways in macrophages using expression vectors with CMV promoters: a cautionary tale. Inflammation.
525   2005;29(2-3):94-102.

41. E. Isern, M. Gustems, M. Messerle, E. Borst, P. Ghazal and A. Angulo. The activator protein 1 binding motifs within the human cytomegalovirus major immediate-early enhancer are functionally redundant and act in a cooperative manner with the NF-kappaB sites during acute infection. J Virol. 2011;85(4):1732-1746.

530   42. M. Rodova, R. Jayini, R. Singasani, E. Chipps and M. R. Islam. CMV promoter is repressed by p53 and activated by JNK pathway. Plasmid. 2013;69(3):223-230.

43. T. Corring, A. Aumaitre and G. Durand. Development of digestive enzymes in the piglet from birth to 8 weeks. I. Pancreas and pancreatic enzymes. Nutr Metab. 1978;22(4):231-243.

535 44. D. Rantzer, P. Kiela, M. J. Thaela, J. Svendsen, B. Ahren, S. Karlsson and S. G. Pierzynowski. Pancreatic exocrine secretion during the first days after weaning in pigs. J Anim Sci. 1997;75(5):1324-1331.

45. S. G. Pierzynowski, B. R. Westrom, C. Erlanson-Albertsson, B. Ahre'n, J. Svendsen and B. W. Karlsson. Induction of exocrine pancreas maturation at weaning in young developing pigs. J Pediatr Gastroenterol Nutr. 1993;16(3):287-293.

540 46. M. D. Lindemann, S. G. Cornelius, S. M. el Kandelgy, R. L. Moser and J. E. Pettigrew. Effect of age, weaning and diet on digestive enzyme levels in the piglet. J Anim Sci. 1986;62(5):1298-1307.

47. W. F. Owsley, D. E. Orr Jr and L. F. Tribble. Effects of age and diet on the development of the pancreas and the synthesis and secretion of pancreatic enzymes in the young pig. J Anim Sci. 545 1986;63(2):497-504.

48. Pavelka M, Roth J. Liver Epithelium: Bile Canaliculi. In: Functional Ultrastructure. Anonymous , editors. Springer; 2010. p.216.

49. A. Gimelbrant, J. N. Hutchinson, B. R. Thompson and A. Chess. Widespread monoallelic expression on human autosomes. Science. 2007;318(5853):1136-1140.

550 50. M. L. Opsahl, A. Springbett, R. Lathe, A. Colman, M. McClenaghan and C. B. Whitelaw. Mono-allelic expression of variegating transgene locus in the mouse. Transgenic Res. 2003;12(6):661-669.

51. J. Velten, C. Cakir, E. Youn, J. Chen and C. I. Cazzonelli. Transgene silencing and transgene-derived siRNA production in tobacco plants homozygous for an introduced AtMYB90 555 construct. PLoS One. 2012;7(2):e30141.

52. F. de Carvalho, G. Gheysen, S. Kushnir, M. Van Montagu, D. Inze and C. Castresana. Suppression of beta-1,3-glucanase transgene expression in homozygous plants. EMBO J. 1992;11(7):2595-2602.

53. F. de Carvalho Niebel, P. Frendo, M. Van Montagu and M. Cornelissen. Post-transcriptional 560 cosuppression of beta-1,3-glucanase genes does not affect accumulation of transgene nuclear mRNA. Plant Cell. 1995;7(3):347-358.

54. H. Vaucheret, J. C. Palauqui, T. Elmayan and B. Moffatt. Molecular and genetic analysis of nitrite reductase co-suppression in transgenic tobacco plants. Mol Gen Genet. 1995;248(3):311-317.

565   55. M. Fagard and H. Vaucheret. (Trans)gene silencing in plants: How Many Mechanisms? Annu Rev Plant Physiol Plant Mol Biol. 2000;51:167-194.

56. H. Vaucheret, C. Beclin and M. Fagard. Post-transcriptional gene silencing in plants. J Cell Sci. 2001;114(Pt 17):3083-3091.

57. P. Brodersen and O. Voinnet. The diversity of RNA silencing pathways in plants. Trends
570   Genet. 2006;22(5):268-280.

58. D. Schubert, B. Lechtenberg, A. Forsbach, M. Gils, S. Bahadur and R. Schmidt. Silencing in Arabidopsis T-DNA transformants: the predominant role of a gene-specific RNA sensing mechanism versus position effects. Plant Cell. 2004;16(10):2561-2572.

59. M. V. Matz, A. F. Fradkov, Y. A. Labas, A. P. Savitsky, A. G. Zaraisky, M. L. Markelov and
575   S. A. Lukyanov. Fluorescent proteins from nonbioluminescent Anthozoa species. Nat Biotechnol. 1999;17(10):969-973.

60. Y. Nakamura, J. Ishii and A. Kondo. Bright fluorescence monitoring system utilizing Zoanthus sp. green fluorescent protein (ZsGreen) for human G-protein-coupled receptor signaling in microbial yeast cells. PLoS One. 2013;8(12):e82237.

580   61. R. N. Day and M. W. Davidson. The fluorescent protein palette: tools for cellular imaging. Chem Soc Rev. 2009;38(10):2887-2921.

62. G. Takahashi, C. B. Gurumurthy, K. Wada, H. Miura, M. Sato and M. Ohtsuka. GONAD: Genome-editing via Oviductal Nucleic Acids Delivery system: a novel microinjection independent genome engineering method in mice. Sci Rep. 2015;5:11406.

585   63. M. D. Muzumdar, B. Tasic, K. Miyamichi, L. Li and L. Luo. A global double-fluorescent Cre reporter mouse. Genesis. 2007;45(9):593-605.

64. R. M. Quadros, H. Miura, D. W. Harms, H. Akatsuka, T. Sato, T. Aida, R. Redder, G. P. Richardson, Y. Inagaki, D. Sakai, S. M. Buckley, P. Seshacharyulu, S. K. Batra, M. A. Behlke, S. A. Zeiner, A. M. Jacobi, Y. Izu, W. B. Thoreson, L. D. Urness, S. L. Mansour, M. Ohtsuka
590   and C. B. Gurumurthy. Easi-CRISPR: a robust method for one-step generation of mice carrying conditional and insertion alleles using long ssDNA donors and CRISPR ribonucleoproteins. Genome Biol. 2017;18(1):92-017-1220-4.

65. H. Miura, R. M. Quadros, C. B. Gurumurthy and M. Ohtsuka. Easi-CRISPR for creating knock-in and conditional knockout mouse models using long ssDNA donors. Nat Protoc.
595   2018;13(1):195-215.

Figure legends

Fig. 1 Production of CMV-ZsGreen1 hemizygous and homozygous transgenic pigs. **(A)** Representative photograph of control piglet (left) with homozygous (middle) and hemizygous (right) transgenic siblings. ZsGreen1 was visualized with an ultraviolet light and a Roscolux #15 filter. **(B)** Representative immunoblot of ZsGreen1 in the skin of control piglets (n = 3) as well as hemizygous (n = 3) and homozygous (n = 3) transgenic littermates. Lower panel represents total protein levels within each lane. **(C)** Schematic demonstrating the genotyping method utilized to distinguish between control, homozygous and hemizygous piglets. Detection of monoallelic or biallelic transgene integration was performed by conventional polymerase chain reaction (PCR) using 3 primers. The forward primer, denoted by the black arrow, amplifies a non-coding region upstream of the transgene integration site. A reverse primer (grey arrow) aligns a non-coding region downstream of the transgene and a second reverse primer, indicated in green, detects the transgene. If a transgene is present, the resulting PCR product will be 847 bp; however, in the absence of the transgene, the PCR product will be 627 bp. **(D)** PCR detection of transgene integration and copy number. The presence of only a 847 bp PCR product indicates a homozygous transgenic animal whereas a 627 bp product indicates a control animal. If both 847 and 627 bp products are detected, the genotype is hemizygous transgenic.

Fig. 2 Visualization of ZsGreen1 fluorescence within internal organs of transgenic piglets. **(A)** Representative image of ZsGreen1 fluorescence in the internal organs of control and transgenic pigs. Left panel represents internal organs from a transgenic piglet. Middle panel represents a fluorescent image of internal organs from a transgenic piglet; and right panel is a UV image of wild-type piglet organs. **(B)** Representative fluorescent images of a transgenic piglet brain (left panel), anterior and posterior pituitary glands (within the sella turcica of the skull; middle panel)

and heart (right panel). Anterior pituitary gland and posterior pituitary gland are indicated by white

and grey arrows, respectively. ZsGreen1 expression was abundant within the right ventricle of the

heart (right panel). Abbreviations: L, liver; St, stomach, Sp, spleen; I, intestines; Cr, cerebrum; Cl,

cerebellum, RA, right atrium; RV, right ventricle; LA, left atrium; LV, left ventricle. ZsGreen1

625   fluorescence was detected with UV light and a Roscolux #15 filter.

Fig. 3. Protein levels of ZsGreen1 within neural and endocrine tissues of CMV-ZsGreen1 pigs.

(A) Representative immunoblot of neural and endocrine tissue from hemizygous transgenic piglets

(n = 3) using an antibody directed against ZsGreen1 (upper panel). Middle panel represents total

630   protein levels that served as the loading control. Quantification of immunoblots revealed

differences in relative ZsGreen1 protein levels between tissue types ($P = 0.0002$; lower panel). **(B)**

Representative immunoblot of neural and endocrine tissue from homozygous transgenic piglets (n

= 3) using an antibody directed against ZsGreen1 (upper panel). Middle panel represents total

protein levels that served as the loading control. Quantification of immunoblots revealed a

635   difference in relative ZsGreen1 protein levels between tissue types ($P = 0.0396$; lower panel). [a,b,c]

Bars with alternate letters differ ($P < 0.05$).

Fig. 4. Protein levels of ZsGreen1 within thoracic tissues of CMV-ZsGreen1 pigs. **(A)**

Representative immunoblot of thoracic tissue from hemizygous transgenic piglets (n = 3) using an

640   antibody directed against ZsGreen1 (upper panel). Middle panel represents total protein levels that

served as the loading control. Quantification of immunoblots revealed differences in relative

ZsGreen1 protein levels between tissue types ($P < 0.0001$; lower panel). **(B)** Representative

immunoblot of thoracic tissue from homozygous transgenic piglets (n = 3) using an antibody

directed against ZsGreen1 (upper panel). Middle panel represents total protein levels that served

645   as the loading control. Quantification of immunoblots indicated that there were no differences in

relative ZsGreen1 abundance between tissue types (P = 0.3326; lower panel). [a,b,c] Bars with

alternate letters differ (P < 0.05).

Fig. 5. Protein levels of ZsGreen1 within digestive tissues of CMV-ZsGreen1 pigs. **(A)**

650   Representative immunoblot of digestive tissue from hemizygous transgenic piglets (n = 3) using

an antibody directed against ZsGreen1 (upper panel). Middle panel represents total protein levels

that served as the loading control. Quantification of immunoblots revealed differences in relative

ZsGreen1 protein levels between tissue types (P < 0.0001; lower panel). **(B)** Representative

immunoblot of digestive tissue from homozygous transgenic piglets (n = 3) using an antibody

655   directed against ZsGreen1 (upper panel). Middle panel represents total protein levels that served

as the loading control. Quantification of immunoblots revealed differences in relative ZsGreen1

protein levels between tissue types (P = 0.0212; lower panel). [a,b,c] Bars with alternate letters differ

(P < 0.05).

660   **Fig. 6.** Protein levels of ZsGreen1 within renal tissues of CMV-ZsGreen1 pigs. **(A)** Representative

immunoblot of renal tissue from hemizygous transgenic piglets (n = 3) using an antibody directed

against ZsGreen1 (upper panel). Middle panel represents total protein levels that served as the

loading control. Quantification of immunoblots revealed differences in relative ZsGreen1 protein

levels between tissue types (P = 0.0083; lower panel). **(B)** Representative immunoblot of renal

665   tissue from homozygous transgenic piglets (n = 3) using an antibody directed against ZsGreen1

(upper panel). Middle panel represents total protein levels that served as the loading control.

Quantification of immunoblots revealed a tendency for differences in relative ZsGreen1 protein levels between tissue types (P = 0.0620; lower panel). [a,b,c] Bars with alternate letters differ ($P <$ 0.05).

Fig. 7. Protein levels of ZsGreen1 within reproductive tissues of CMV-ZsGreen1 pigs. **(A)** Representative immunoblot of reproductive tissue from hemizygous transgenic piglets (n = 3) using an antibody directed against ZsGreen1 (upper panel). Middle panel represents total protein levels that served as the loading control. Quantification of immunoblots indicated that there were

675   no differences in relative ZsGreen1 abundance between tissue types (P = 0.3227; lower panel). **(B)** Representative immunoblot of reproductive tissue from homozygous transgenic piglets (n = 3) using an antibody directed against ZsGreen1 (upper panel). Middle panel represents total protein levels that served as the loading control. Quantification of immunoblots indicated that there were no differences in relative ZsGreen1 abundance between tissue types (P = 0.4797).

Figure 1

Figure 2

A

B

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7