Article

Phylogenetic evidence for independent origins of GDF1 and GDF3 genes in amphibians and mammals

Juan C. Opazo & Kattina Zavala
Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias,
Universidad Austral de Chile, Valdivia, Chile.

Corresponding author
Juan C. Opazo
Instituto de Ciencias Ambientales y Evolutivas
Facultad de Ciencias
Universidad Austral de Chile
Phone: +56 63 2221674
Email: jopazo@gmail.com

Abstract

Growth differentiation factors 1 (GDF1) and 3 (GDF3) are members of the

transforming growth factor superfamily (TGF-β) that is involved in

fundamental early-developmental processes that are conserved across

vertebrates. The evolutionary history of these genes is still under debate due

to ambiguous definitions of homologous relationships among vertebrates.

Thus, the goal of this study was to unravel the evolution of the GDF1 and

GDF3 genes of vertebrates, emphasizing the understanding of homologous

relationships and their evolutionary origin. Surprisingly, our results revealed

that the GDF1 and GDF3 genes found in amphibians and mammals are the

products of independent duplication events of an ancestral gene in the

ancestor of each of these lineages. The main implication of this result is that

the GDF1 and GDF3 genes of amphibians and mammals are not 1:1

orthologs. In other words, genes that participate in fundamental processes

during early development have been reinvented two independent times during

the evolutionary history of tetrapods.

Keywords: gene duplication; gene family evolution; tetrapods; left-right

identity; development

Introduction

Growth differentiation factors 1 (GDF1) and 3 (GDF3) are members of the

transforming growth factor superfamily (TGF-β) that were originally isolated

from mouse embryonic libraries [1,2]. They perform fundamental roles during

early development, GDF1 has been mainly associated to the regulation of the

left-right patterning, whereas GDF3 is mainly involved in the formation of the

anterior visceral endoderm, mesoderm and the establishment of anterior-

posterior identity of the body [3–20]. Deficiencies in GDF1/GDF3 give rise to a

broad spectrum of defects including right pulmonary isomerism, visceral *situs*

inversus, transposition of the great arteries, and cardiac anomalies among

others [17,21–25]. In addition to their developmental roles, GDF1 has been

described as a tumor suppressor gene in gastric cells, counteracting

tumorogenesis by stimulating the SMAD signaling pathway [26], and GDF3 has

been associated with the regulation of adipose tissue homeostasis and

energy balance during nutrient overload [27–29].

The evolutionary history of the GDF1 and GDF3 genes is still a matter

of debate due to the unclear definition of homologous relationships.

Understanding homology is a fundamental aspect of biology as it allows us to

comprehend the degree of relatedness between genes that are associated to

a given phenotype in a group of organisms. This is particularly important for

GDF1 and GDF3 as these genes perform biological functions during early

stages of development that define key aspects of the body plan of all

vertebrates [3,4,13–20,5–12]. Until now, most of the inferences of the homology

relationships of these genes have been based on functional information. For

example, based on the developmental processes that these genes regulate, it

has been suggested that the mammalian GDF1 gene is the true ortholog of

the Vg1 (GDF1) gene found in amphibians [15,17,21]. Furthermore, phylogenetic

analyses performed by Andersson et al., (2007) were not able to define

orthologous relationships between the GDF1 and GDF3 genes among

vertebrates. However after performing genomic comparisons, these authors

did propose that the GDF1 gene present in mammals is the true ortholog of

the Vg1 (GDF1) gene present in amphibians [30]. It was also suggested that the

GDF3 gene could be an evolutionary innovation of mammals, as Andersson

et al., (2007) did not find GDF3 sequences in the amphibian and bird

genomes [30,31]. Given this scenario, the single copy gene found in amphibians

and birds would be a co-ortholog of the mammalian genes [30]. More recently, a

second copy of a GDF gene (derrière) has been annotated at the 3′ side of

the Vg1 (GDF1) in the genome of the western clawed frog (*Xenopus*

tropicalis), thus further complicating the definition of homologous relationships

and the origin of genes that accomplish fundamental roles during early

development in vertebrates.

With emphasis on understanding homologous relationships and their

evolutionary origin, the goal of this study was to unravel the evolution of the

GDF1 and GDF3 genes of vertebrates. Surprisingly, our phylogenetic

analyses revealed that the GDF1 and GDF3 genes of amphibians and

mammals are the products of independent duplication events of an ancestral

gene in the ancestor of each of these groups. We also found the signature of

two chromosomal translocations, the first occurred in the ancestor of

tetrapods whereas the second was found in the ancestor of mammals. Thus,

our results support the hypothesis that in amphibians and mammals,

descendent copies of the same ancestral gene (GDF1/3) have independently

subfunctionalized to perform key developmental functions in vertebrates.

Results and Discussion

Phylogenetic analyses suggest an independent origin of the GDF1 and

GDF3 genes in mammals and amphibians

From an evolutionary perspective, the definition of homologous relationships

among GDF1 and GDF3 genes is still a matter of debate [30,31]. The resolution

of this homology is important as these genes are involved in fundamental

developmental processes which are conserved all across vertebrates [8,32–34].

Thus, if extant species inherited these genes from the vertebrate ancestor, the

developmental processes in which GDF1 and GDF3 are involved have a

single evolutionary origin and are comparable among species.

Based on evolutionary analyses and the developmental processes in

which GDF1 has been shown to participate, it has been suggested that GDF1

in mammals, birds, and amphibians are 1:1 orthologs [15,17,21,30,35]. Given that it

has not been possible to identify copies of the GDF3 gene in amphibians and

birds, it has been proposed that this gene is an evolutionary innovation of

mammals, and the single gene copy found in amphibians and birds is co-
ortholog to the mammalian duplicates [30]. However, a second copy of a GDF
gene has been annotated in the genome of the western clawed frog (*Xenopus*
tropicalis) [36]. This newly described gene is located at the 3´ side of the GDF1
gene and indicates that amphibians, like mammals, also have a repertoire of
two GDF genes (GDF1 (Vg1) and GDF3 (derrière)). This new finding further
complicates the resolution of homologous relationships and the origin of
genes involved in the early development of all vertebrates [8,32–34].
Our maximum likelihood and Bayesian reconstructions revealed that
the GDF1 and GDF3 genes of amphibians and mammals were both
reciprocally monophyletic (Fig. 1), indicating that these genes originated
independently in each of these two lineages (Fig. 1). In agreement with the
literature, we found only one gene in sauropsids (e.g. birds, turtles, lizards
and allies), suggesting that they retained the ancestral condition of a single
gene copy as found in non-tetrapod vertebrates (e.g. coelacanths, bony fish,
chondricthyes and cyclostomes)(Fig. 1). Dot-plot comparisons provided
further support for the presence of a single gene copy in sauropsids as no
traces of an extra GDF gene were present in the syntenic region of
representative species of the group (Fig. 2). However, we cannot rule out an
alternative scenario of duplication and subsequent gene loss in the ancestor
of sauropsids. Thus, our evolutionary analyses suggest that the GDF1 and
GDF3 genes present in amphibians and mammals diversified independently
in the ancestor of each of these lineages. In other words, genes that
participate in fundamental processes during early development have been
reinvented two independent times during the evolutionary history of tetrapods.
As a consequence of the independent origin of these genes in amphibians
and mammals, they are not 1:1 orthologs.
To further test our hypothesis of the independent origins of the GDF1
and GDF3 genes in amphibians and mammals we performed topology tests.
In these analyses we compared our phylogenetic tree (Fig. 1 and Fig. 3B) to
the topology predicted from a one-duplication model in which the duplication
event that gave rise to the GDF1 and GDF3 genes in amphibians and
mammals occurred in the ancestor of tetrapods (Fig. 3A). In the one-
duplication model, it is assumed that the ancestor of sauropsids had two gene

copies and that subsequently one of these was lost. Thus, according to the

one-duplication model (Fig. 3A), the predicted phylogeny would recover a

monophyletic group containing GDF1 sequences from mammals, sauropsids,

and amphibians sister to a clade containing GDF3 sequences from the same

groups (Fig. 3A). Alternatively, the predicted phylogeny from the two-

duplication model would retrieve a clade containing GDF1 and GDF3

sequences from mammals sister to a clade containing GDF1/3 sequences

from sauropsids; additionally a clade containing GDF1 and GDF3 sequences

from amphibians would be recovered sister to the mammalian/sauropsid clade

(Fig. 3B). Results of the topology tests rejected the phylogeny predicted by

the one-duplication model (Weighted Shimodaira and Hasegawa, $p < 10^{-4}$;

Weighted Kishino Hasegawa, $p < 10^{-4}$). Thus, this result provided additional

support to our hypothesis that the GDF1 and GDF3 genes of amphibians and

mammals are the product of lineage independent duplication events in the

ancestors of each of these groups.

In the literature there are other cases of groups of genes that perform

similar biological functions in a diversity of species but that have originated via

lineage independent duplication events [37–43]. Among these, the independent

origin of the β-globin gene cluster in all main groups of tetrapods (e.g. therian

mammals, monotremes, birds, crocodiles, turtles, squamates, amphibians)

represents a well-documented phenomenon [41–43]. This case is of particular

interest as the two β-globin subunits that come from a gene family that has

been reinvented several times during the evolutionary history of tetrapods are

assembled in a tetramer with two α-globin subunits that belong to a group of

genes that possess a single origin [37,41–43]. Besides this example, the

independent origin of gene families makes the task of comparison difficult, as

the repertoire of genes linked to physiological processes in different lineages

do not have the same evolutionary origin. Additionally, the fact that the

evolutionary process can give rise to similar phenotypes following different

mutational pathways makes the problem of comparing even more difficult

(Natarajan et al., 2016). This is particularly important when extrapolating the

results of physiological studies performed in model species to other

organisms.

Two translocation events during the evolutionary history of the GDF1
and GDF3 genes of tetrapods
Based on the chromosomal distribution of the GDF1 and GDF3 genes and the
conservation pattern of flanking genes, we propose that during the
evolutionary history of tetrapods, these genes underwent two chromosomal
translocation events (Fig. 4). According to our analyses the genomic region
that harbors the single gene copy of chondricthyes, bony fish, and
coelacanths is conserved, yet this region in these groups differs from the
regions in which GDF1 and GDF3 are located in tetrapods. Overall this
suggests that the first translocation event occurred in the ancestor of
tetrapods (Fig. 4). Interestingly, the genomic region where the single gene
copy is located in non-tetrapod vertebrates, which is defined by the presence
of upstream (BMP2, HAO1, TMX4 and PLCB1) and downstream genes
(FERMT1, LRRN4, CRLS1), is conserved in tetrapods. Given this, it is
possible to identify the chromosomal location where the gene was located in
the tetrapod ancestor before the first translocation event (Fig. 4). On the other
hand, the fact that the mammalian GDF1 and GDF3 genes are located in
different chromosomes suggests that the second translocation event occurred
in the ancestor of the group (Fig. 4). In humans, the GDF1 gene is located on
chromosome 19 while GDF3 is located on chromosome 12. In opossum
(*Didelphis virginiana*) the GDF1 and GDF3 genes are located on
chromosomes 3 and 8, respectively.
202
Evolution of GDF1 and GDF3 genes in vertebrates
In this study we present compelling evidence suggesting that genes involved
in the formation of the primitive streak, anterior visceral endoderm, mesoderm
and the establishment of the left-right identity [3,4,14–17,20,30,5–7,9–13] in amphibians
and mammals are the product of independent duplications events. As such,
these results indicate that the GDF1 and GDF3 genes of amphibians and
mammals are not 1:1 orthologs.
Thus, according to our results the last common ancestor of vertebrates
had a repertoire of one gene (GDF1/3), and this condition is maintained in
actual species of cylostomes, chondricthyes, bony fish, and coelacanths (Fig.
5). In the ancestor of tetrapods, the single gene copy was translocated from a

chromosomal region defined by the presence of the BMP2, HAO1, TMX4,

PLCB1, FERMT1, LRRN4, CRLS1 genes to a chromosomal region defined by

the presence of the CERS1, COPE, DDX49 and HOMER3 genes (Fig. 4).

After this translocation, the ancestral gene underwent a duplication event in

the amphibian ancestor, giving rise to the GDF1 (Vg1) and GDF3 (derrière)

genes as they are found in actual species. In the case of the western clawed

frog (*Xenopus tropicalis*) these genes are located in tandem on chromosome

28 (Fig. 4). Sauropsids, the group that includes birds, crocodiles, turtles,

lizards and snakes, inherited the ancestral condition of a single gene copy as

is seen in non-tetrapod vertebrates (Fig. 5). Finally, in the ancestor of

mammals, the ancestral gene also underwent a duplication event, giving rise

to the GDF1 and GDF3 genes found in extant species of mammals. After

duplication but before the radiation of the group, a second translocation event

occurred in the ancestor of the group (Fig. 5). Thus, all mammals inherited a

repertoire of two genes (GDF1 and GDF3) that are located on two

chromosomes.

Concluding remarks

This study provides a comprehensive evolutionary analysis of the GDF1 and

GDF3 genes in representative species of all main groups of vertebrates. The

main focus of this study was to unravel the duplicative history of the GDF1

and GDF3 genes and to understand homologous relationships among

vertebrates. Understanding homology in this case is particularly important as

these genes perform fundamental roles during early development that are

conserved across vertebrates [8,32–34]. Surprisingly, our results revealed that the

GDF1 and GDF3 genes present in amphibians and mammals are the product

of independent duplication events in the ancestor of each of these groups.

Subsequently, the GDF1 and GDF3 genes of amphibians and mammals are

not 1:1 orthologs. Our results also show that all other vertebrate groups - i.e

non-tetrapods and sauropsids – maintained the ancestral condition of a single

gene copy (GDF1/3).

From an evolutionary perspective the independent duplication events

that occurred in the ancestors of mammals and amphibians could have

resulted in the division of labor, with some degree of redundancy, of the

function performed by the ancestral gene. In support of this idea, it has been

shown that the GDF1 and GDF3 genes of mammals have partially redundant

functions during development where GDF1 can to some degree compensate

for the lack of GDF3 [30]. Additionally, it has also been shown that double

knockout animals (GDF1$^{-/-}$ and GDF3$^{-/-}$) present more severe phenotypes

than those of either single knockout (GDF1$^{-/-}$ or GDF3$^{-/-}$)[30]. Detailed

comparisons of the developmental roles of the GDF1 and GDF3 genes

between mammals and amphibians will shed light regarding the reinvention of

genes that possess fundamental roles during early development. On the other

hand, comparing mammals and amphibians with sauropsids will provide

useful information regarding the evolutionary fate of duplicated genes. Finally,

it would be interesting to study the evolutionary history of genes that

cooperate with GDF1 and GDF3 during early development (e.g. nodal) [8,33,44]

in order to understand the evolutionary nature of the entire developmental

network.

Material and Methods

DNA sequences and phylogenetic analyses

We annotated GDF1 and GDF3 genes in representative species of chordates.

Our study included representative species from mammals, birds, reptiles,

amphibians, coelacanths, holostean fish, teleost fish, cartilaginous fish,

cyclostomes, urochordates and cephalochordates (Supplementary dataset 1

and 2). We identified genomic pieces containing GDF 1 and GDF3 genes in

the Ensembl database using BLASTN with default settings or NCBI database

(refseq_genomes, htgs, and wgs) using tbalstn (Altschul et al., 1990) with

default settings. Conserved synteny was also used as a criterion to define the

genomic region containing GDF1 and GDF3 genes. Once identified, genomic

pieces were extracted including the 5´and 3´ flanking genes. After extraction,

we curated the existing annotation by comparing known exon sequences to

genomic pieces using the program Blast2seq with default parameters

(Tatusova and Madden 1999). Putatively functional genes were characterized

by an open intact reading frame with the canonical exon/intron structure

typical of vertebrate GDF1 and GDF3 genes. Sequences derived from shorter

records based on genomic DNA or cDNA were also included in order to attain

a broad and balanced taxonomic coverage. Amino acid sequences were

aligned using the L-INS-i strategy from MAFFT v.7 [45] (Supplementary Dataset

3). Phylogenetic relationships were estimated using maximum likelihood and

Bayesian approaches. We used the proposed model tool from IQ-Tree [46] to

select the best-fitting model (JTT+F+R5). Maximum likelihood analysis was

also performed in IQ-Tree [46] to obtain the best tree. Node support was

assessed with 1000 bootstrap pseudoreplicates using the ultrafast routine.

Bayesian searches were conducted in MrBayes v.3.1.2 [47]. Two independent

runs of six simultaneous chains for 10×10^6 generations were set, and every

2,500 generations were sampled using default priors. The run was considered

to have reached convergence once the likelihood scores formed an

asymptote and the average standard deviation of the split frequencies

remained < 0.01. We discarded all trees that were sampled before

convergence, and we evaluated support for the nodes and parameter

estimates from a majority rule consensus of the last 2,000 trees. Sea squirt

(*Ciona intestinalis*) and Florida lancelet (*Branchiostoma floridae*) BMP2

sequences were used as outgroups.

Assessment of conserved synteny

We examined genes found up- and downstream of GDF1 and GDF3 in

species representative of vertebrates. Synteny assessment were conducted

for human (*Homo sapiens*), chicken (*Gallus gallus*), western-clawed frog

(*Xenopus tropicalis*), coelacanth (*Latimeria chalumnae*), spotted gar

(*Lepisosteus oculatus*), elephant shark (*Callorhinchus milii*) and sea lamprey

(*Petromyzon marinus*). Initial ortholog predictions were derived from the

EnsemblCompara database [48] and were visualized using the program

Genomicus v91.01[49]. In other cases, the genome data viewer platform from

the National Center for Biotechnology information was used.

Acknowledgements

This work was supported by the Fondo Nacional de Desarrollo Científico y

Tecnológico (FONDECYT 1160627) grant to JCO.

Figure legends

Figure 1. Maximum likelihood tree depicting evolutionary relationships among

the GDF1 and GDF3 genes of chordates. Numbers on the nodes represent

maximum likelihood ultrafast bootstrap and Bayesian posterior probability

support values. BMP2 sequences from urochordates (*Ciona intestinalis*) and

cephalochordates (*Branchiostoma floridae*) were used as outgroups.

Figure 2. Dot-plots of pairwise sequence similarity between the GDF1 and

GDF3 genes of the western clawed frog (*Xenopus tropicalis*) and the

corresponding syntenic region in the American alligator (*Alligator*

mississippiensis), Burmese python (*Python bivittatus*), chicken (*Gallus gallus*)

and green turtle (*Chelonia mydas*).

Figure 3. Schematic representations of alternative hypotheses of the sister

group relationships among duplicated GDF genes in tetrapods. A) According

to the one-duplication model the predicted phylogeny recovers a monophyletic

group containing GDF1 sequences from mammals, sauropsids and

amphibians sister to a clade containing GDF3 sequences from the same

groups. B) The phylogenetic prediction from the two-duplication model

retrieves a clade containing GDF1 and GDF3 sequences from mammals

sister to a clade containing GDF1/3 sequences from sauropsids; additionally a

clade containing GDF1 and GDF3 sequences from amphibians is recovered

sister to the mammalian/sauropsid clade.

Figure 4. Structure of the chromosomal region containing the GDF1 and

GDF3 genes of vertebrates. Asterisks denote that the orientation of the

genomic piece is from 3' to 5', gray lines represent intervening genes that do

not contribute to conserved synteny.

Figure 5. An evolutionary hypothesis of the evolution of the GDF1 and GDF3

genes in vertebrates. According to this model the last common ancestor of

vertebrates had a repertoire of one gene (GDF1/3), a condition that has been

maintained in actual species of cylostomes, chondrichthyes, bony fish,

coelacanths and sauropsids. In the ancestor of tetrapods, the single gene

copy was translocated to a different chromosomal location. After that, in the

amphibian ancestor, the single gene copy underwent a duplication event

giving rise to the amphibian GDF1 (Vg1) and GDF3 (derrière) genes. In the

ancestor of mammals, the single gene copy also underwent a duplication

event, giving rise to the mammalian GDF1 and GDF3 genes. After the

duplication, but before the radiation of the group, a second translocation event

occurred in the ancestor of the group. Thus, all mammals inherited a

repertoire of two genes (GDF1 and GDF3) that were located on two

chromosomes.

References

1. Lee, S.-J. Identification of a Novel Member (GDF-1) of the Transforming Growth Factor-beta Superfamily. *Mol. Endocrinol.* **4,** 1034–1040 (1990).

2. Jones, C. M., Simon-Chazottes, D., Guenet, J.-L., Hogan, B. L. M. & Pasteur, L. Isolation of Vgr-2, a Novel Member of the Transforming Growth Factor-P- Related Gene Family. *Mol. Endocrinol.* **6,** 1961–1968 (1992).

3. Lee, S.-J. Expression of growth/differentiation factor 1 in the nervous system: Conservation of a bicistronic structure. *Biochemistry* **88,** 4250–4254 (1991).

4. Dale ', L., Matthews, G. & Colman2, A. Secretion and mesoderm-inducing activity of the TGF-j-related domain of Xenopus Vgl. *EMBO J.* **12,** 4471–4480 (1993).

5. Chen, C. The Vg1-related protein Gdf3 acts in a Nodal signaling pathway in the pre-gastrulation mouse embryo. *Development* **133,** 319–329 (2006).

6. Andersson, O., Reissmann, E., Jörnvall, H. & Ibáñez, C. F. Synergistic interaction between Gdf1 and Nodal during anterior axis development. *Dev. Biol.* **293,** 370–381 (2006).

7. Bengtsson, H. *et al.* Generation and characterization of a Gdf1 conditional null allele. *Genesis* **46,** 368–372 (2008).

8. Komatsu, Y. & Mishina, Y. Establishment of left – right asymmetry in vertebrate development : the node in mouse embryos. *Cell. Mol. Life Sci.* **70,** 4659–4666 (2013).

9. Arias, C. F., Herrero, M. A., Stern, C. D. & Bertocchini, F. A molecular mechanism of symmetry breaking in the early chick embryo. *Scietific Reports* **7,** 1–6 (2017).

10. Bisgrove, B. W., Su, Y.-C. & Yost, H. J. Maternal Gdf3 is an obligatory cofactor in Nodal signaling for embryonic axis formation in zebrafish. *Elife* **6,** e28534 (2017).

11. Montague, T. G. & Schier, A. F. Vg1-Nodal heterodimers are the endogenous inducers of mesendoderm. *Elife* **6,** e28183 (2017).

12.   PellicciaJ.L., Jindal, G. A. & Burdine, R. D. Gdf3 is required for robust
Nodal signaling during germ layer formation and left-right patterning.
Elife **6,** e28635 (2017).

13.   Thomsen, G. H. & Melton, D. A. Processed Vg1 protein is an axial
mesoderm inducer in xenopus. *Cell* **74,** 433–441 (1993).

14.   Hyatt, B. A., Lohr, J. L. & Yost, H. J. Initiation of vertebrate left–right
axis formation by maternal Vg1. *Nature* **384,** 62–65 (1996).

15.   Hyatt, B. A. & Yost, H. J. The left-right coordinator: The role of Vg1 in
organizing left-right axis formation. *Cell* **93,** 37–46 (1998).

16.   Söderström, S. & Ebendal, T. Localized expression of BMP and GDF
mRNA in the rodent brain. *J. Neurosci. Res.* **56,** 482–492 (1999).

17.   Rankin, C. T., Bunton, T., Lawler, A. M. & Lee, S. J. Regulation of left-
right patterning in mice by growth/differentiation factor-1. *Nat. Genet.*
24, 262–265 (2000).

18.   Wright, C. V. E. Mechanisms of Left-Right Asymmetry : What ' s Right
and What ' s Left ? A recent meeting at the Juan March Foundation in.
Cell **1,** 179–186 (2001).

19.   Hamada, H., Meno, C., Watanabe, D. & Saijoh, Y. Establishment of
vertebrate left-right asymmetry. *Nat. Rev. Genet.* **3,** 103–113 (2002).

20.   Somi, S., Houweling, A. C., Buffing, A. A. M., Moorman, A. F. M. & Van
Den Hoff, M. J. B. Expression of cVg1 mRNA during chicken embryonic
development. *Anat. Rec.* **273A,** 603–608 (2003).

21.   Wall, N. A., Craig, E. J., Labosky, P. A. & Kessler, D. S. Mesendoderm
induction and reversal of left-right pattern by mouse Gdf1, a Vg1-related
gene. *Dev. Biol.* **227,** 495–509 (2000).

22.   Karkera, J. D. *et al.* Loss-of-Function Mutations in Growth Differentiation
Factor-1 (GDF1) Are Associated with Congenital Heart Defects in
Humans. *Am. J. Hum. Genet.* **81,** 987–994 (2007).

23.   Kaasinen, E. *et al.* Recessively inherited right atrial isomerism caused
by mutations in growth/differentiation factor 1 (GDF1). *Hum. Mol. Genet.*
19, 2747–2753 (2010).

24.   Bao, M. W. *et al.* Cardioprotective role of growth/differentiation factor 1
in post-infarction left ventricular remodelling and dysfunction. *J. Pathol.*
236, 360–372 (2015).

25. Zhang, J. *et al.* Association of GDF1 rs4808863 with fetal congenital
heart defects: A case-control study. *BMJ Open* **5,** e009352 (2015).

26. Yang, W. *et al.* Epigenetic silencing of GDF1 disrupts SMAD signaling
to reinforce gastric cancer development. *Oncogene* **35,** 2133–2144
(2016).

27. Witthuhn, B. A. & Bernlohr, D. A. Upregulation of bone morphogenetic
protein GDF-3/Vgr-2 expression in adipose tissue of FABP4/aP2 null
mice. *Cytokine* **14,** 129–135 (2001).

28. Wang, W., Yang, Y., Meng, Y. & Shi, Y. GDF-3 is an adipogenic
cytokine under high fat dietary condition. *Biochem. Biophys. Res.*
Commun. **321,** 1024–1031 (2004).

29. Andersson, O. *et al.* Growth/differentiation factor 3 signals through
ALK7 and regulates accumulation of adipose tissue and diet-induced
obesity. *Proc. Natl. Acad. Sci. USA* **105,** 7252–7256 (2008).

30. Andersson, O., Bertolino, P. & Ibáñez, C. F. Distinct and cooperative
roles of mammalian Vg1 homologs GDF1 and GDF3 during early
embryonic development. *Dev. Biol.* **311,** 500–511 (2007).

31. Levine, A. J., Brivanlou, A. H., Suzuki, A. & Hemmati-Brivanlou, A.
GDF3, a BMP inhibitor, regulates cell fate in stem cells and early
embryos. *Development* **133,** 209–16 (2006).

32. Mercola, M. & Levin, M. Left-right assymetry determination in
vertebrates. *Annu. Rev. Cell Dev. Biol.* **17,** 779–805 (2001).

33. Levine, A. J. GDF3, a BMP inhibitor, regulates cell fate in stem cells and
early embryos. *Development* **133,** 209–216 (2005).

34. Nakamura, T. & Hamada, H. Left-right patterning: conserved and
divergent mechanisms. *Development* **139,** 3262 (2012).

35. Seleiro, E. A. P., Connolly, D. J. & Cooke, J. Early developmental
expression and experimental axis determination by the chicken Vg1
gene. *Curr. Biol.* **6,** 1476–1486 (1996).

36. Suzuki, A. *et al.* Genome organization of the vg1 and nodal3 gene
clusters in the allotetraploid frog Xenopus laevis. *Dev. Biol.* (2017).
doi:10.1016/j.ydbio.2016.04.014

37. Goodman, M., Czelusniak, J., Koop, B. F., Tagle, D. A. & Slightom, J. L.
Globins: a case study in molecular phylogeny. *Cold Spring Harb. Symp.*

Quant. Biol. **52,** 875–90 (1987).

38.   Kriener, K., O 'huigin, C. & Klein, J. Independent Origin of Functional

MHC Class II Genes in Humans and New World Monkeys. *Hum.*

Immunol. **62,** 1–14 (2001).

39.   Wallis, O. C. & Wallis, M. Characterisation of the GH gene cluster in a

new-world monkey, the marmoset (Callithrix jacchus). *J. Mol.*

Endocrinol. **29,** 89–97 (2002).

40.   Li, Y. *et al.* Independent origin of the growth hormone gene family in

New World monkeys and Old World monkeys/hominoids. *J. Mol.*

Endocrinol. **35,** 399–409 (2005).

41.   Opazo, J. C., Hoffmann, F. G. & Storz, J. F. Genomic evidence for

independent origins of $_{NL}$ -like globin genes in monotremes and therian

mammals. *Proc. Natl. Acad. Sci. USA* **105,** 1590–1595 (2008).

42.   Hoffmann, F. G., Storz, J. F., Gorr, T. A. & Opazo, J. C. Lineage-

specific patterns of functional diversification in the α-and β-globin gene

families of tetrapod vertebrates. *Mol. Biol. Evol.* **27,** 1126–1138 (2010).

43.   Hoffmann, F. G., Vandewege, M. W., Storz, J. F. & Opazo, J. C. Gene

Turnover and Diversification of the α- and β-Globin Gene Families in

Sauropsid Vertebrates. *Genome Biol. Evol.* **10,** 344–358 (2018).

44.   Ramsdell, A. F. & Yost, H. J. Molecular mechanisms of vertebrate left–

right development. *Trends Genet.* **14,** 459–465 (1998).

45.   Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment

Software Version 7: Improvements in Performance and Usability. *Mol.*

Biol. Evol. **30,** 772–780 (2013).

46.   Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A. & Minh, B. Q. W-IQ-

TREE: a fast online phylogenetic tool for maximum likelihood analysis.

Nucleic Acids Res. **44,** W232–W235 (2016).

47.   Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic

inference under mixed models. *Bioinformatics* **19,** 1572–4 (2003).

48.   Herrero, J. *et al.* Ensembl comparative genomics resources. *Database*

2016, baw053 (2016).

49.   Louis, A., Nguyen, N. T. T., Muffato, M. & Roest Crollius, H. Genomicus

update 2015: KaryoView and MatrixView provide a genome-wide

perspective to multispecies comparative genomics. *Nucleic Acids Res.*
43, D682–D689 (2015).
502
503
504

Contributions

506 Designed the research: JCO; carried out the research: JCO and KZ;
507 contributed materials/reagents/analysis tools: JCO; wrote the paper: JCO

Additional information

Competing interests

511 The authors declare no competing interests.
512
513

A) One-duplication model

B) Two-duplication model

● = gene duplication