# Reactivation of latent HIV-1 *in vitro* using an alcoholic extract from *Euphorbia umbellata* (Euphorbiaceae) latex

Ana Luiza Chaves Valadão[1], Paula Pezzuto[1], Viviane A. Oliveira Silva[2], Barbara Simonson Gonçalves[3], Átila Duque Rossi[1], Rodrigo Delvecchio da Cunha[1], Antonio Carlos Siani[4], João Batista de Freitas Tostes[4], Marcelo Trovó[5], Paulo Damasco[3], Gabriel Gonçalves[1], Rui Manuel Reis[6,7], Renato Santana de Aguiar[1], Cleonice Alves de Melo Bento[3], Amilcar Tanuri[1] *

[1]Laboratório de Virologia Molecular, Departamento de Genética, Universidade Federal do Rio de Janeiro, Brazil.

[2]Centro de Pesquisa em Oncologia Molecular, Hospital de Câncer de Barretos, Barretos, Brazil.

[3]Departamento de Microbiologia e Parasitologia, Universidade Federal do Estado do Rio de Janeiro, Brazil.

[4]Departamento de Produtos Naturais, Instituto de Tecnologia em Fármacos, Fundação Oswaldo Cruz, Brazil.

[5]Departamento de Botânica, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Brazil.

[6]Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho, Braga, Portugal.

[7]ICVS/3B's – PT Government Associate Laboratory, Braga/Guimarães, Portugal.

* **Corresponding Author**: Amilcar Tanuri

**Address for correspondence:** Laboratory of Molecular Virology, Department of Genetics, IB, Federal University of Rio de Janeiro. Av. Carlos Chagas Filho 373, CCS, Bloco A, Sala 121 Cidade Universitária, CEP: 21941-902 Rio de Janeiro - RJ, Brasil.
**Email:** atanuri1@gmail.com

**Abstract**

*Euphorbia umbellata (E. umbellata)* belongs to *Euphorbiaceae* family, popularly known as Janauba, and contains in its latex a combination of phorbol esters with biological activities described to different cellular protein kinase C (PKC) isoforms. Here, we identified deoxi-phorbol esters present in *E. umbellata* latex alcoholic extract able to increase HIV transcription and reactivate HIV from latency models. This activity was mediated by NF-kB activation followed by nuclear translocation and binding to HIV LTR promoter. In addition, *E. umbellate* latex extract induced the production of pro inflammatory cytokines together with IL21 in *in vitro* human PBMC cultures. Our latex extract activates latent HIV in human PBMCs isolated from HIV positive patients as well as latent SIV in non-human primate primary CD4$^+$ T lymphocytes. These results strongly indicate that the phorbol esters present in *E. umbellata* latex are promising candidate compounds for future clinical trials for *shock and kill* therapy to promote HIV cure and eradication.

**Keywords:** HIV-1, latency, reactivation, PKC, NF-κB.

**Running Title :** New natural extract for reactivation of latent HIV-1

## INTRODUCTION

The Acquired Immunodeficiency Syndrome (AIDS) is caused by the Human Immunodeficiency Virus (HIV), which was identified in 1983 **(1,2)** from patients with AIDS.  In those 34 years of pandemic, more than 35 million people have died from AIDS-related illnesses since the start of the epidemic until 2016 **(3).**

The pandemic of HIV infection is an important socioeconomic burden and is considered one of the largest documented epidemics in history. HIV pathogenesis involves infection and replication in CD4$^+$ T lymphocytes, macrophages and dendritic cells. Replication and destruction of CD4$^+$ T cells, which are key effector cells in the host immune response, leads to a clinical outcome of immunosuppression known as AIDS **(4)**.

The current strategy used in the treatment of HIV infection is named Antiretroviral Therapy (ART), which is based on a combination of antiretroviral drugs, which target different stages of HIV replicative cycle. The primary goal of ART is to block viral replication and slow down the progression of immunodeficiency, restoring immunity and improving the quality of life of the infected patient **(5)**. However, adverse effects have been reported as a consequence of the antiretroviral therapy, in conjunction with immunological, neurological and metabolic co-morbidities associated either with HIV infection or drugs interaction **(6)**. These facts triggered the search for a functional or sterilizing cure for HIV infection.

Even when an undetectable viral load is achieved with effective antiretroviral therapy, it is still possible to detect latent viral reservoirs where HIV proviral DNA is integrated in resting memory CD4+ T cells **(7)**. These cells containing latent viral genomes become viral reservoirs inaccessible to ART that can only act on actively replicating viruses. Once ART is stopped, viral load increases rapidly and systemic infection reestablishes **(8)**. Therefore, the complete eradication of HIV from an infected individual is one of the most challenging areas of HIV research today. The way to eliminate HIV from patient is to activate the virus in the presence of ART and or activate the immune response to destroy viral reservoir or make patient´s immune response to keep viral replication at the bay.

New latent virus reactivating compounds, also called latency reversing agents (LRA), when used conjointly with ART, are capable to eliminate transcriptionally inactive viruses, ultimately resulting in the eradication of HIV infection. Histone deacetylase (HDAC) inhibitors are promising classes of latency-reversing agents (LRAs) that are undergoing extensive testing in *in vitro* models and in initial pilot clinical trials to reactivate latent HIV-1 infection. HDAC inhibitors were developed as anticancer drugs as HDACs play important role in epigenetic and non-epigenetic transcriptional regulation, inducing apoptosis and cell cycle arrest **(9)**. In the context of HIV-1 reactivation, HDAC inhibitors induce transcription at the HIV-1 long terminal repeat (LTR) **(10-13)**. This strategy is called "shock and kill" and it postulates that memory CD4$^+$ T lymphocytes treated with LRA along with ART could purge totally or partially the viral reservoir while blocking replication of emergent latent virus, leading to a HIV eradication or at least a functional cure **(14)**. Another promising class of molecules with therapeutic potential has been reported as possible candidates for elimination of viral reservoirs are the Protein Kinase C (PKC) agonists **(7)**. This class includes: Prostratin **(15)**, Bryostatin **(16)**, Ingenol **(17)** and Phorbol 12-Myristate 13-Acetate (PMA) **(18)**. These compounds can reactivate latent HIV in lymphoid and myeloid cells. Although several transcriptional regulatory mechanisms for HIV latency have been described in the context of epigenetic silencing and transcription repression, recent work suggests that reactivation of latent HIV is positively correlated with NF-κB activation, demonstrating the importance of this factor in latency reactivation **(19, 20)**.

Many natural compounds have been investigated for their antiviral and latency-reversing properties, being indicated as candidates for clinical trials. Several of these compounds are derived from secondary metabolics of plants, such as terpenoids **(21)**, polyphenols **(22)**, alkaloids **(23)** and phorbol esters **(24)**. *Euphorbiaceae* family is a source of potent phorbol esters, which specimens are found  worldwide and havebeen used for the treatment of several diseases in the popular medicinal use **(25-29)**. *Euphorbia umbellate* is popularly known in Brazil as Janauba or *leiterinha* and is used, in traditional medicine, as a latex preparation (diluted in water) for the treatment of gastric disorders, such as peptic ulcer and gastritis, as well as neoplastic diseases, allergy, diabetes, influenza, leprosy, among other diseases (**30**). Several papers

have reported the isolation of lectins, phorbols, terpenes, flavonoids and other

phenolic compounds from *E. umbellata* latex **(31-33)**. This plant contains

phorbol esters in its latex similar to PMA isolated from the seeds of *Croton*

*tiglium*, another representative of the *Euphorbiaceae* family. These compounds

are efficient in inhibiting the cytopathic effect generated by HIV infection on

lymphocyte cells, as well as activating protein kinase C (PKC) that in turn

activates NF-κB, promoting its nuclear translocation and promoting HIVLTR

transcriptional reactivation reverting viral latency **(34)**.

In this work we evaluated the ability of an alcoholic extract obtained from

Janauba latex (JALEx) to reactivate latent HIV and investigated the molecular

mechanisms involved in this activity. The anti-latency property of JALEx was

demonstrated either in *ex vivo* HIV latent derived human PBMC cells or in CD4+

T Lymphocytes isolated from SIV infected non-human-primates. All these

properties makes JALEx a promising natural medicine to be used in HIV cure

strategies.

**MATERIAL AND METHODS**

**Latex isolation from *Euphorbia umbellata*.** *Euphorbia umbellata* (Pax) Bruyns, the current accepted name for *Synadenium grantii* Hook.f., was collected and identified with the respective voucher specimen kept at the RFA Herbarium under the number Trovó, M.L.O 764. The crude latex was collected directly from incision mechanically made in the trunk of cultivated specimens at Barretos county, São Paulo state, Brazil. The latex was straightly diluted in 80% ethanol (1:10 v/v) to produce a white rubber-like precipitate that was filtered off by gravity in filter papers. The filtrate was dried in a rotary evaporator to afford a fine white powder, which was diluted in DMSO to a final concentration of 20 mg/mL. This extracted material was labeled as JALEx (Janaúba Alcoholic Latex Extract). JALEx was chromatographed in a silica gel column using a solvent gradient (0-5%) of ethyl ether in chloroform to enrich the phorbol-containing fractions and favor their detection and characterization.

**Chemical profile of JALEx.** JALEx was analyzed by gas chromatography coupled to mass spectrometry (GC-MS) by using an Agilent 6890N chromatograph (Palo Alto, CA) equipped with a capillary column DB-H17HT (30 m × 0.25 mm × 0.15 μm film thickness). Helium was the carrier gas at 1.0 mL/min and inlet pressure of 3.14 psi, 10:1 split/splitless ratio. The oven temperature programmed from 150°C to 340°C at 4°C/min and then held for 17 minutes. The injection volume was 1 μL of a 2 mg/mL chloroform solution. Data were processed using MSD Productivity Chem Station Software operating with ion source at 250°C and electron impact ionization at 70 eV. Individual components were characterized by their fragmentation patterns in the mass spectrum.

**Viruses and cells.** Analysis of HIV reactivation *in vitro* in latent infected cells after treatment with JALEx. Human HIV latent lymphocyte cell models (J-Lat 8.4 and 10.6) were used **(35)**. These cells are derived from Jurkat cells containing copies of HIV genome in fusion with GFP reporter gene integrated into the nuclear. These cells have different levels of reactivation capacity, depending on the site of HIV genome integration. The J-Lat 10.6 clone is a more responsive one showing reactivation levels higher than those observed by the J-Lat 8.4

clone. Both cells are cultured in RPMI medium supplemented with 10% fetal bovine serum (FBS) and 1% penicillin / streptomycin antibiotics. J-Lat cells at density of $10^5$ cells per well of 24-well plates were treated with different concentrations of JALEx dissolved in DMSO for 24 hours. Positive controls for reactivation of HIV latent virus were ingenol B (0.32 μM) and PMA (1 μM). Cells were analyzed by flow cytometric (BD Accuri C6) to determine the level of GFP expression, in order to evaluate HIV transcription. A total of 10,000 gated live cells were collected and data represent the percentage of GFP-expressing cells in total gated events. All the experiments were performed in three independent replicates.

**Cell viability assay.** Evaluation of the cytotoxicity of the compounds was performed by incubating J-Lat cells with the serial dilutions of JALEx compounds previously tested in the reactivation assays. Cell viability assessment was performed in $10^4$ cells plated in 96-well plates, in three independent experiments, each in sextuplicates. After 5 days of culture, vital resazurin dye (CellTiter Blue, Promega) was added to the cells and absorbance levels were evaluated by spectrophotometer at 595 nm.

**Analysis of the activation of PKCs.** To assess the mechanism of action of HIV latency reactivation by Janauba compounds, $10^5$ J-Lat cells in 24-well plates were treated with different specific inhibitors of PKCα, PKCδ and PKCγ isoforms Go6976, Go6983, Ro-31-8220, respectively, for 24 hours **(36-38)**. After this time, cells were treated with the positive control of PKC activation (1 μM PMA) and two concentrations of JALEx previously determined in the reactivation and cytotoxicity assays. Untreated cells were referred as MOCK. The following concentrations of the alcoholic extract were tested: 0.01 μg/mL and 0.001 μg/mL and the percentage of GFP positive cells was assessed by flow cytometry analysis.

In addition, PKC activation was evaluated by immunofluorescence to detect cytosolic location of different PKC isoforms. For this, $10^4$ HeLa cells were plated in black 96-well glass bottom plates. These cells were transfected with three different plasmids encoding three PKC isoforms fused to GFP (PKCα-GFP, PKCδ-GFP and PKCγ-GFP) for 24 hours and then treated with 1 μg/mL and 0.1

μg/mL of the JALEx, for 10 minutes or 24 hours. Positive control was performed using PMA (1 μM). Cells were washed with PBS and DAPI stained. Cellular localization of PKC-GFP was evaluated by High Content Screening confocal microscope (Molecular Devices, Inc). The redistribution of PKC isoforms was quantified in ImageJ software.

Phosphorylation of PKC isoforms was also investigated. For this, Jurkat cells were treated with 0.01 μg/mL and 0.001 μg/mL of JALEx and with 1 μM PMA as a positive control for 10 and 30 minutes, 1, 6 and 24 hours. Cellular extractes were prepared in lysis buffer (50 mM Tris pH 7.6, 150 mM NaCl, 5 mM EDTA, 1 mM $Na_3VO_4$, 10 mM NaF, 10 mM NaPyrophosphate, 1% NP-40 supplemented with protease inhibitors cocktail) for 1 hour. Lysates were subjected to 10% SDS-PAGE gels and after transferred to nitrocellulose membranes the following antibodies were used for immunoblotting: anti-PKC phosphorylated PKC (pan, δ, θ) (Cell Signaling; 1:500) and β-tubulin (Cell Signaling; 1:2000). Proteins quantification was evaluated using band densitometry analysis, performed with the Image J software (version 1.41; National Institutes of Health).

**NF-κB activation assays**. In order to verify whether JALEx induces the activation and downstream nuclear internalization of NF-κB, HeLa cells were plated at the density of $2x10^4$ cells per well in 96-well plates. Cells were treated with 1 μg/mL and 0.1 μg/mL of JALEx for 6 or 24 hours. PMA (1 μM) was used as positive control. The NF-κB subunit p65 was visualized by immunofluorescence (anti-p65, 1:300, Santa Cruz Biotechnologies), and nuclear (DAPI) translocation was evaluated by fluorescence confocal microscopy (Molecular Devices, Inc).

In order to evaluate the role of NF-κB in reactivation of latent HIV, Jurkat cells ($10^6$ cells) were transiently transfected with 8 μg of pBlue30LTR-Luc Wild Type (NFkB WT) or p-LTR-MUT (NFkB MUT) using the Neon Transfection System kit (Thermo Fisher Scientific) according to the manufacturer's recommendations. The latter plasmid was generated by site-directed mutagenesis by altering the two NF-κB binding sites, rendering that plasmid nonfunctional for NF-κB response. Both plasmids were provided by Dr. Renato S. de Aguiar (UFRJ) and those constructs have the luciferase sequence under the control of the HIV viral

promoter. In the case of plasmid carrying NF-kB mutate sequence, the second
guanine at both NF-κB binding sites into LTR promoter was changed to a
cytosine, resulting in the plasmid designed here as *pBlue3'LTR NF-κB MUT-*
*Luc* (NF-κB BS⁻). This plasmid carries a LTR sequence lacking functional NF-
κB binding sites. After transfections, cells were equally divided into two groups:
treated or untreated with 1μg/mL of JALEx, for 24 hours. After this interval, cells
were lysed and luciferase activity was assessed in a luminometer (Glomax,
Promega).

**Analysis of surface markers expression and cytokines production in**
**primary cells treated with JALEx.** In order to expand the results obtained in
HIV latency model cells, we performed assays with JALEx in peripheral blood
mononuclear cells (PBMCs) from 4 healthy donors. After separation of buffy
coat by Ficoll centrifugation, CD4$^+$ T cells were isolated using the Dynabeads
CD4 Positive Isolation Kit (Thermo Fisher) kit to obtain 10$^7$ PBMCs, according
to the manufacturer's recommendations. TCD4$^+$ cells were treated with 1 μM
PMA, 10 μg/mL (10$^{-3}$) or 1 μg/mL (10$^{-4}$) of JALEx, in addition to MOCK
(untreated cells), for 24 hours. Expression of surface markers such as CD4,
CCR5 and CXCR4, as well as the expression of activation markers CD25,
CD38, HLA-DR and CD69 were evaluated by specific antibody labeling and flow
cytometric analysis (BD Accuri C6). Comparatively, this assay was also
performed on MT4 lymphocyte cells line at the same conditions.

Cell supernatants from the TDC4$^+$ cells were collected for analysis of the
cytokine production stimulated by JALEx treatment. A colorimetric multiplex
immunoassay Bio-Plex Pro Human Cytokine 17-plex Assay (Bio-Rad)
containing fluorescent microspheres conjugated to different monoclonal
antibodies specific for each target cytokine was chosen for this analysis. The
following cytokines were measured: IL-1β, IL-2, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10,
IL- IL-17, G-CSF, GM-CSF, MCP-1 / MCAF, MIP-1β, IFNγ and TNF-α. .

**Quantitative viral outgrowth assay.** To compare the ability of the compounds
to stimulate virus replication in latently infected cells we used a variation of the
quantitative viral outgrowth assay (QVOA) described by Laird et al. **(39)**. Cells
for the QVOA were isolated from viremic SIVmac239-infected cynomolgus

macques at the Wisconsin National Primate Research Center (WNPRC). PBMC
isolated by density gradient centrifugation were enriched for CD4$^+$ T cells using
the CD4 T cell enrichment kit for non-human primates (Miltenyi Biotec, San
Diego, CA). After enrichment, cells were plated at 1x10$^6$ cells per well. These
cells were co-cultured with "stimulator" CEMx174 cells that were irradiated with
10,500 rads prior to co-culture. The ratio of stimulator cells to CD4$^+$ T cells was
2:1. Purified CD4$^+$ T cells were stimulated with JALEx, concanavalin A, ingenol,
or no mitogen, overnight. After stimulation, cells were washed and re-plated in
serial dilutions, in triplicate. At this point, additional non-irradiated CEMx174
cells were added to each well to serve as targets for induced virus to infect. The
target cells were added at a 2:1 ratio to the CD4$^+$ T cells. Co-cultures were
maintained for 1 week. On the last day, supernatant was collected and RNA
was isolated to test for the presence of SIV. RNA was isolated using the Viral
Total Nucleic Acid kit for the Maxwell 16 MDx instrument (Promega, Madison,
WI). Viral RNA was quantified using a qRT-PCR assay targeting the *gag* gene
as described previously **(40)** except that analysis was performed using the
Lightcycler 480.

**Effect of JALEx in T cell activation from HIV positive patients.** In order to
analyze the ability of JALEx in regulating different T cell phenotypes in the
context of HIV-1 infection, peripheral blood of 15 adults ART-treated patients (8
women and 9 men), with a mean plasma viral load of 953 ± 1.924 copies of
RNA/mL and CD4$^+$ (845 ± 345/mm$^3$) and CD8$^+$ (845 ± 345/mm$^3$) higher than
350 cells/mm$^3$, were collected and PBMC were obtained from Ficoll-Paque
gradient. Viable PBMC were quantified by trypan blue exclusion and suspended
in RPMI-1640 medium supplemented with 2 µM of L-glutamine (GIBCO,
Carlsbad, CA, USA), 10% of fetal calf serum, 20 U/mL of penicillin, 20 µg/mL of
streptomycin and 20 mM of HEPES buffer. Approximately 1 x 10$^6$ PBMC/mL
were cultured for 24 hours in 24-well plates with 1mL of complete medium in the
presence or absence of different concentration of JALEx (1, 0.1 and 0.01
µg/mL). Phorbol-12-myristate-13-acetate (PMA 20 ng/mL) plus ionomycin (600
ng/mL) were used as positive control. To optimize intracellular cytokine staining,
brefeldin A (10 µg/mL) was added to the cultures in the last 4 hours of
incubation time. For the fluorescence labeling, mouse anti-human fluorescent

monoclonal antibodies (mAbs) directed against CD4-FITC, CD8-FITC were purchased from BD Bioscience (San Diego, CA, USA). Briefly, anti-CD4 and anti-CD8 mAbs were added to the PBMC ($2 \times 10^5$/tube) and incubated for 30 min at room temperature in the dark. The cells were washed with PBS and then permeabilized with Cytofix/Cytoperm (BD Pharmingen, San Diego, CA) at 4 °C for 20 min. After washing, the antibodies for intracellular staining (anti- IL-17-PECy7, anti-IFN-γ-PE-Cy5, anti-IL-10-PE-Cy7, anti-IL-21-PE) or the corresponding anti-IgG1 isotype control, were added in various combinations and incubated for 30 min at 4°C. Cells were analyzed in the Attune ® (Thermo Fischer, Co) and FlowJo software. Isotype control antibodies and single-stained samples were periodically used to check the settings and gates on the flow cytometer. After acquisition of 100,000 to 200,000 events, lymphocytes were gated based on forward and side scatter properties after the exclusion of dead cells, by using propidium iodide and doublets. The analysis of CD4$^+$ and CD8$^+$ T cells viability was determined using 7-aminoactinomycin D (7-AAD) (R&D systems). Further, the gated cells were negative for CD14 marker. The CD4$^+$ T lymphocyte count (CD4) was performed on a flow cytometer (FACS Count, Becton Dickson®). Plasma HIV RNA quantification was performed using the RT-PCR with Abbott M2000 Sq HIV VL technology.

**Statistical analysis**. The statistical analysis was performed using Prism 5.0 software (GraphPad Software). Statistical evaluations were performed by Kruskal-Wallis Test, Mann–Whitney U test and Two-way ANOVA. The analysis are indicated in the figures.

**RESULTS**

**JALEx isolation from *Euphorbia umbellate* and chemical analysis.**

The JALEx soluble in water and ethanol was characterized mainly by a mixture of triterpenes and phorbol ester diterpenes. The GC profile and MS data indicated that triterpenes was predominant throughout the whole extract, with two main constituents standing out, represented as T1 and T2 in figure 1A. The triterpenes profile is in accordance with data from the literature that report the isolation of euphol, lanosterol, friedelin and 3β-friedelinol in the latex, leaves

and stem of *E. umbellata* **(41)**, whilst studies with the latex have led to the characterization of euphol and the steroid citrostadienol **(42)**. A phorbol esters-rich fraction was obtained by chromatographing the crude JALEx in a silica gel open column using mixtures of chloroform-ether as eluent (Ph1 and Ph 2 - Fig. 1). The class of phorbol ester has been represented by the 12-O-tigloyl-4-deoxyphorbol-13-isobutirate **(43)**, 12-deoxyphorbol-13-(2-methylpropionate), phorbol 12,13,20-triacetate **(44)**, 4-deoxyphorbol-12,13-ditiglate and 3,4,12,13-tetraacetylphorbol-20-phenylacetate **(45)** distributed in several plant tissues (Fig. S1).

**Reactivation of latent HIV by JALEx and its cytotoxicity effects**

In this work, we used the alcoholic extracts from JALEx resuspended in DMSO. The reactivation of latent HIV was evaluated using the lymphocytic lineage J-Lat cells (clones 8.4 and 10.6) harboring the HIV integrated genome. The cells were exposed to increasing concentrations of JALEx, ranging from 10 to $10^{-7}$ µg/mL. The reactivation of latent HIV was evaluated by GFP reporter gene expression cloned in fusion with the virus integrated genome. In parallel, cytotoxicity assay was performed to evaluate whether the concentrations of JALEx would be toxic to these cells. JALEx treatment at concentrations of 10 to 0.01 µg/mL showed HIV activation percentage similar to positive controls (PMA 1 µM and ING-B 0.32 µM), dropping to less than 5% reactivation at dilutions of 0.001 to 0.0000001 µg/mL in the J-Lat 8.4 clone (Fig. 2A). Our results showed cytotoxic effects of JALEx at 10 µg/mL, while the other concentrations were well tolerated, showing a good cell viability (Fig. 2B). As expected, HIV reactivation by JALEx was more expressive in J-Lat cells clone 10.6, reaching values of 75% compared to 12% reactivation in 8.4 cells (Fig. 2C). In relation to the cytotoxicity, JALEx at 10 µg/mL was also toxic for J-Lat 10.6, whereas the other concentrations presented cells viability between 80% and 75% (Fig. 2D). These results demonstrated the ability of JALEx to stimulate HIV transcription and reactivate virus from latency models.

**JALEx reactivation of latent HIV is mediated by PKC**

In order to evaluate whether JALEx activation of HIV transcription is mediated by PKC pathway J-Lat 8.4 and 10.6 cells were treated for 24 hours with different PKC inhibitors (Gö6666, Gö6983 and Rö31-8220) to prevent JALEx stimulation. After this period, the cells were incubated with non cytotoxic concentrations of JALEx (0.01 and 0.001 µg/mL). PMA is a phorbol ester that activates different isoforms of PKCs and was used as positive control (1 µM). HIV reactivation was evaluated by flow cytometry for quantification of GFP positive cells. The pre-exposition of both clones of J-Lat cells (8.4 and 10.6) to the PKC inhibitors reverse the activation promoted by the PKC agonist PMA compared with PMA only treated cells (Fig. 3). The same results were found when J-Lat clone 8.4 was pre-incubated with inhibitors Gö6976, Gö6983 and Rö31-8220 before JALEx treatment either at concentration 0.01 or 0.001 µg/mL. This result suggests that JALEx is a pan activator of different isoforms of PKCs (PKCα, PKCδ and PKCγ) (Fig. 3 - left panel). The same profile was found with the J-Lat clone 10.6 indicating that PKC inhibitors block the reactivation of latent HIV by JALEx treatment. Rö31-8220 was the only inhibitor that could not totally prevent JALEx stimulation reaching values around 20% of activation compared with 60% of non-treated cells (Fig. 3 - right panel). All together, these results suggests that JALEx is a broad range PKC agonist, since the treatment with PKC inhibitors Gö 6976 (specific for PKC-α and β), Gö 6983 (specific for PKC-α, PKC-β, PKC-γ, PKC-δ and PKC-ζ) and Rö-31–8220 (specific for PKC-α, PKC-β, PKC-ε and PKC-γ) decreased the levels of HIV reactivation compared with non treated cells.

**JALEx induces activation of different isoforms of PKC**

PKC family is a citoplasmatic sensor of several signal transduction pathways by phosphorylation of hydroxyl groups in serine/threonine residues of other proteins. Moreover, the catalytic domain present in the structure of PKCs is essential for its auto-phosphorylation and, consequently, its cellular activation and relocation to the inner face of plasma membrane. PKC family comprises different isoforms spanning different regulatory and catalytic domains. The

isoforms are divided into three subgroups: the conventional isoforms α, βI, βII

and γ that are $Ca^{2+}$ and DAG dependente for activation; the novel class

encompassing the δ, ε, η that require DAG, but not $Ca^{2+}$ for activation and the

atypical isoforms ι and λ that require neither $Ca^{2+}$ nor diacylglycerol for

activation. For these reasons, we investigated whether phosphorylation of

different PKC isoforms was occurring in response to JALEx treatment. In these

experiments, two concentrations of JALEx (0.01 µg/mL and 0.001 µg/mL) were

used and PMA was utilized as positive control of PKC phosphorylation. Cells

were collected at different time points (10 and 30 minutes; 1, 6 and 24 hours).

We observed that the phosphorylation of conventional PKCs, identified by anti-

pan-PKC-phosphorylated antibody labeling, occurs quite rapidly, after 10

minutes of treatment with JALEx, in the two concentrations tested (Fig. S2A,

upper panels). This activation is long lasting, remaining phosphorylated for up to

24 hours after the treatments (Fig. S2A, upper panels). The same pattern was

observed at the same time points for cells treated with PMA (Fig. S2A, upper

panels). For the novel PKC isoforms, PKC θ exhibited a rapid phosphorylation

pattern (10 min) that was not sustained over time (1 to 24h) (Fig. S2A, bottom

panels). PKC δ showed an early phosphorylation pattern maintained for up to 6

hours with JALEx treatment in both concentrations (0.01 µg/mL and 0.001

µg/mL). However, only JALEx at 0.001 µg/mL induces PKC δ phosphorylation

after 24 hours (Fig. S2A, bottom panels). Densitometry analyses compared with

loading controls (α-tubulin) and normalized by DMSO treated cells showed that,

in general, JALEx at concentrations of 0.01 µg/mL and 0.001 µg/mL induces the

phosphorylation of conventional and novel isoforms of PKC that remains at

least up to 6 hours of stimulation (Fig. S2B). However, as suggested by the WB

analysis, the phosphorylation of PKC-δ is sustained until 24 hours post JALEx

treatment. All these results are in agreement with the previous observations that

JALEx is a pan activator of PKC isoforms.

activation, PKCs are translocated to the plasma membrane by RACK

proteins (membrane-bound receptor for activated protein kinase C proteins)

**(46)**. To verify if JALEx promotes the translocation of PKC isoforms to the

plasma membrane, HeLa cells were transfected with plasmids expressing three

PKC isoforms fused to GFP (PKCα-GFP, PKCδ-GFP and PKCγ-GFP). After

transfection, cells were treated with 1 and 0.1 μg/mL of JALEx for 10 minutes or
24 hours and the citoplasmatic location of PKC were evidenced by fluorescence
confocal microscopy images. We observed a diffuse pattern of PKC distribution
in untreated cells (mock). However, PMA or JALEx treatment at both
concentrations (1 and 0.1 μg/mL) induced the periplasmatic distribution of all
PKC isoforms (α, δ and γ) (Fig. 4A). The same results were obtained with
longer treatment scheme (up to 24h) (Fig. 4B). Finally, these results showed
that JALEx promotes activation of different classes of PKC through
phosphorylation followed by downstream relocation to the plasma membrane.

**430 JALEx promotes nuclear translocation of NF-κB and interaction with HIV**
**431 promoter.**

Phosphorylation of NF-kB inhibitor IκB by PKC leads to its ubiquitination
and proteasomal degradation, releasing NF-κB complexes in the cytoplasm to
translocate to the nucleus where they bind to HIV promoter (LTR region) and
induce virus gene expression **(47-49)**. For this reason, we evaluated the effect
of JALEx treatment on nuclear translocation of NF-κB by means of two
strategies. In the first one, we analyzed the occurrence of nuclear translocation
of NF-κB after treatment of HeLa cells with 1 and 0.1 μg/mL of JALEx for 6 or
24 hours, through immunofluorescence and confocal fluorescence microscopy
using anti-p55 antibody. We observed no significant translocation of NF-κB to
the nucleus of HeLa cells 6 hours post JALEx treatment with the same
cytoplasmic distribution as observed in untreated cells (mock - Fig. 5A).
However, we confirmed the NF-kB translocation to the nucleus after 24 hours of
JALEx treatment, once NF-kB staining was located in the nuclei (Fig. 5B). To
better quantify the translocation of NF-κB, we quantified the mean values of
fluorescence intensity between the nucleus and the cytoplasm of 100 cells, in
each experimental condition. We observed an increasing ratio of NF-kB
fluorescence into the nucleus 24 hours after JALEx treatment, and the same
was observed for the positive control (PMA - Fig. 5C). JALEx treatment at the
concentration of 1 μg/mL showed the highest rates of NF-kB internalization (Fig.
5C).

We also evaluated the recruitment of NF-κB to the HIV promoter by
JALEx treatment. For that, we transfected Jurkat T CD4[+] cells with LTR-
luciferase reporter plasmids containing either intact NF-κB binding sites (NF-kB
WT) or harboring mutations that prevent NF-κB interaction **(17, 50, 51)**.
Transfected cells were treated with 1 μg/mL of JALEx and luciferase activity
was assessed 24 hours later. Our results showed that JALEx treatment induced
a 5-fold increase in luciferase expression in cells transfected with pNF-kB WT.
However, no luciferase activity was observed in cells transfected with pNF-kB
MUT (Fig. 5D). Together, our results demonstrated that JALEx induces HIV
transcription through NF-kB release of its inhibitors, followed by nuclear
translocation and binding to HIV promoter.

**JALEx modulates surface markers expression and cytokines release in**
**primary human CD4[+] lymphocytes**

To analyze whether the effects of treatment with JALEx would have an
impact on primary human cells, we performed assays on CD4[+] T cells isolated
from PBMCs from 4 healthy donors. Isolated TCD4[+] cells were treated with
PMA (1 μM) or JALEx 10 μg/mL or 1 μg/mL for 24 hours. We evaluated the
expression of HIV receptor and co-receptors (CD4, CCR5 and CXCR4) and
activation markers (CD25, CD38, HLA-DR and CD69) in response to JALEx
treatment. We observed that JALEx treatment (1 μg/mL) downregulates the
surface receptors CD4 and CXCR4 expression levels without any alteration in
CCR5  expression levels (Fig. 6A). The same pattern was observed for cells
treated with PMA (Fig. 6A). Indeed, JALEx treatment also induced the
expression of surface activation markers CD25, CD38 and HLA-DR when
compared to non-treated cells, and with PMA control (Fig. 6B). Figure 6C shows
that CD69 activation marker expression is 75-fold higher relative to control cells
in both treatments. These results suggest that JALEx not only reactivate latent
HIV, but also could prevent new cycles of virus infection down modulating the
virus receptors and co-receptors.

In the light of the previous results of JALEx inducing PBMC activation, we
further analyzed cytokines release directly from the supernatant of human CD4[+]
T cells. This analysis was performed using a multiplex immunoassay to quantify

the production of 17 pro-inflammatory cytokines. JALEx treatment induced the

production of the following cytokines relative to DMSO treated cells: IL-1β, IL-2,

IL-4, IL-5, IL-6, IL-10, IL-12 (p70), IL-13, IL- 17, G-CSF, GM-CSF, MCP-

1/MCAF, IFNγ and TNF-α (Fig. 6D). Overall, the highest concentration of JALEx

(10 µg/mL) induced the same levels of cytokines as did PMA positive control.

IFN-γ and TNF-α were highly induced under JALEx treatment, 5,000 and 1,800

pg/mL, respectively (Fig. 6D). Those results showed that JALEx activate T

CD4$^+$ cells in order to secrete important pro-inflammatory cytokines involved in

macrophage recruitment (MCP-1) and T CD8 response (INF- γ) to block HIV

replication. MIP-β, IL-7 e IL-8 levels were undetectable.

**495    JALEx increases viral load in PBMC from HIV-1 positive patients and**

**496    induces  Th1 and Th17 immune response**

Progression for AIDS in HIV-1 positive patients is mainly characterized

by CD4$^+$ T cell depletion specifically Th1 and Th17 cells, which impact

negatively on immune protection against different opportunistic pathogens. We

first analyzed the virus activation on PBMC isolated from HIV-1 positive patients

under ART treatment. We found that JALEx at 1 µg/mL increases HIV viral load

up to 1.5 log compared with non treated PBMC cells (Fig. 7A). This data was

significant in all PBMC generated from the 15 HIV positive patients. This

increase in HIV viral load was followed by T CD4$^+$ cells depletion (up to 82%)

with no correspondence in T CD8$^+$ cells (Fig. 7B). In order to investigate the

immune response in those patients we evaluate cytokine production in T CD4

and T CD8 positive cells. JALEx treatment (0.1 µg/mL) increased the frequency

of CD4$^+$ and CD8$^+$ T cells secreting IL-17, IFN-γ and IL-21 cytokines. This

induction was higher compared to PMA positive controls (Fig. 7C). Moreover,

JALEx (0.1 µg/mL) enhanced not only the percentage of single IL-21-secreting

CD4$^+$ and CD8$^+$ T cells, but also Th1 and Th17-like subsets positives for IL-21

(Fig. 7D). In contrast, no difference was observed regarding either IFN-γ or IL-

17-secreting CD4$^+$ and CD8$^+$ T cells from PBMC cultures of HIV-1 positive

patients following JALEx treatment (0.1 µg/mL, Fig. 7D). These results

suggested that JALEx treatment could reactivate HIV from reservoirs in patients

with ART suppressive therapy and induces CD4 and CD8 positive cells
secreting protective levels of IL-21.

**JALEx reactivation effects in latently SIV-infected macaque cells**

Finally, we explored the capacity of JALEx to reactivate latent viruses in
CD4$^+$ lymphocytes from SIVmac239-infected *cynomolgus macaques*. We
purified resting memory CD4$^+$ T cells from a single viremic macaque and
prepared serial dilution replicates of these cells to seed into quantitative viral
outgrowth assays (QVOA). Cells in these assays were stimulated in triplicate at
each cell dilution with JALEx or other compounds known to reactivate latent
HIV/SIV through PKC pathway (ingenol and concanavalin A). In order to
compare the ability of these compounds to reactivate latent SIV, we measured
the frequency of infected cells (quantified as infectious units per million cells;
IUPM) across treatments. A higher frequency of infected cells after stimulation
with a given compound suggests its efficiency to reverse latency. The highest
frequency of infected cells was observed in cultures treated with JALEx,
although this frequency was not significantly different from those observed in
other cultures (Table S1). However, JALEx was the only compound capable of
reactivating SIV in all replicates tested showing its superiority compared with
others PKC agonists, such as Ingenol and Concanavalin A.

**DISCUSSION**

In this work we addressed the ability of an ethanolic extract obtained from *Euphorbia umbellata* latex (JALEx) to reactivate HIV/SIV latency in several *in vitro* experiments. The traditional medical uses of this Euphorbiaceae are well known in South America and this plant is vastly used diluting a small volume of crude latex in drinkable water and consuming this solution regularly **(24)**. JALEx contains a series of bioactive phorbol esters (see Supplemental Figure 1). These class of molecules such as PMA are known PKC agonist implicated in lymphocyte activation as well HIV reactivation in well conducted *ex vivo* studies and these activities requires both PKCα and PKCθ isoforms activation **(52)**. Modifications in phorbol ring can lead molecules able to induce carcinogenesis and others such as a deoxy in ring position 4 (4-deoxyphorbol) can lead to molecules unable to induce carcinogenesis but retaining its PKC activation capacity **(53)**. The presence of this modified phorbol ester was identified in our JALEx in our chemical analysis and ina previous study done with *Euphorbia umbellata*latex **(45)**. Therefore, our JALEx is not likely to induce cell carcinogenesis retaining the capacity to reactivate HIV/SIV latency. Many LRA act activating PKC and several compounds such as prostatin **(15)**, bryostatin **(16)**, ingenol B **(17)**, and phorbols such as PMA **(18)**. Fit in this class of molecules Prostratin and bryostratin, are strong PKC agonist and they are very toxic when injected intravenously. In fact, bryostatin was used in a phase II study with 17 patients with progressive non-Hodgkin's lymphoma of indolent type (NHL), previously treated with chemotherapy. These patients received a median of 6 intravenous infusions of 25 µg/m$^2$ bryostatin 1 given once weekly over 24 h and the principal toxicities were myalgia and phlebitis **(54)**. Ingenol B a semisynthetic compound isolated from *Euphorbia tirucalli* was able to reactivate HIV *ex vivo* with great potency using the PKC/NF-kB pathway inducing increased cellular levels of CycT1 and CDK9 in human CD4+lymphocytes in ex vivo experiments **(17)**.

In our study we used JALEx to explore its capability to reactivate HIV latency taking the advantage of the use of this plant in traditional medicine without serious side effects. First we used JALEx in J-Lat 8.4 and 10.6 cells and we could show the potency of this extract reactivates latent HIV transcription

present in J-Lat 8.4 and 10.6 cells in low concentrations (0.01 μg / mL).We also
demonstrate that JALEx act through PKC by inhibiting J-Lat activation with
three different PKC inhibitors (G6666, G6983 and Ro-31-8220). In our assays
Gö6983 was more powerful inhibitor totally blocking HIV reactivation in both J-
Lat cell lines. Go 6983 is a fast pan-PKC inhibitor against PKCα, PKCβ, PKCγ
and PKCδ with $IC_{50}$ in low nM range then JALEx could be reactivating latent
HIV through the activation of different PKC isoforms. Additionally, we showed
that JALEx activates the PKC pathway by promoting NF-κB internalization into
the nucleus and increasing HIV transcription, which is dependent on NF-κB
binding to LTR promoter.  Using image and western blot analysis we showed
that JALEx was able to activate classical and novel PKC isoforms such as: the
conventional isoforms α, βI, βII and γ as well the new ones encompassing the δ,
ε, η and θ isoforms in *in vitro* experiments using JURKAT cell lines. JALEx
could induce θ isoform, in high levels. The transcription factors NF-κB and AP-1
are the primary physiological targets of PKCθ, and efficient activation of these
transcription factors by PKCθ requires integration of TCR and CD28
costimulatory signals. PKCθ cooperates with the protein Ser/Thr phosphatase,
calcineurin, in transducing signals leading to activation of JNK, NFAT, and
the *IL-2* gene. PKCθ also promotes T cell cycle progression **(57)**.

We also evaluate the capacity of JALEx to induce cytokines production in
human PBMC cells *in vitro*. This pattern of alteration in the production of
cytokines from treatment with reactivating compounds is quite common in the
case of the shock and kills strategy, which aims to activate the cells and
promote the exit of HIV latency that is integrated into the cells T. This
phenomenon is known as "cytokine storm" and needs to be further studied in
animal models to define a dose that does not induce this toxic reaction.
Although JALEx induced a conjunction of pro inflammatory cytokines we could
observed a noticeable level of induction of IL-21 by different CD4+ and CD8+T
lymphocyte subsets from PBMC cultures from Aids patients, a noble cytokine
enrolled in the HIV virus load control in HIV+ individuals (**55, 58, 59**). For this
reason, the search for new, more selective compounds that target only a few
PKC isoforms or that have their mechanism of action in other stages of the
activation pathways to reduce these risks is necessary. Of note, leaf and latex

ethanolic extract of E. umbellata latex had low toxicity in adults Rat when dose
orally and acute doses of 2000mg/kg given orally was well tolerated (56).

In our experiments JALEx was efficient in reactivating HIV production
directly in resting T CD4+ cells in HIV+ individuals in ex vivo assays measuring
viral RNA in cell supernatant as well primary non-human primate T CD4+ cells
counterpart in SIV251 latency model using a QVOA assay in low
concentrations. The anti-HIV action of JALEx could occur by three synergistic
effects: a)reactivation of HIV from latency and its destruction by ART; b)
blocking infection of new cells through the down-modulation of CD4 on the
lymphocyte surface; and c) by inducing the secretion of antiviral cytokines such
as IL21. All these *in vivo* properties of JALEx in non-human primate cell models
added to the molecular mechanisms of reactivation presented here as well the
traditional use of *E. umbellata* latex, strongly recommends JALEx or other *E.*
*umbellata* latex preparations as a great candidate for future clinical trials for HIV
eradication
618

---

619 **REFERENCES**

620 1. Barré-Sinousi et al. Isolation of a T-lymphotropic retrovirus from a patient at risk for
621 acquired immune deficiency syndrome (AIDS). Science 1983:Vol. 220, Issue 4599, pp.
622 868-871; DOI: 10.1126/science.6189183.
623
624 2. Gallo RC, Salahuddin SZ, Popovic M, Shearer GM, Kaplan M, Haynes BF, Palker
625 TJ, Redfield R, Oleske J, Safai B, et al. Frequent detection and isolation of cytopathic
626 retroviruses (HTLV-III) from patients with AIDS and at risk for AIDS. Science. 1984 May
627 4;224(4648):500-3. doi: 10.1126/science.6200936.
628
629 3.          UNAIDS,          2017          see          in
630 http://www.unaids.org/sites/default/files/media_asset/UNAIDS_FactSheet_en.pdf
631
632 4. J.A. Levy. Virus-Host Interactions in HIV Pathogenesis. Adv Dent Res. 2011 Apr;
633 23(1): 13–18.doi: 10.1177/0022034511398874.
634
635 5. John Coffin and Ronald Swanstrom . HIV Pathogenesis: Dynamics and Genetics of
636 Viral Populations and Infected Cells. Cold Spring Harb Perspect Med. 2013 Jan; 3(1):
637 a012526.doi: 10.1101/cshperspect.a012526.
638
639 6. Steven G. Deeks, Sharon R. Lewin, and Diane V. Havlir. The End of AIDS: HIV
640 Infection as a Chronic Disease.  Lancet. 2013 Nov 2; 382(9903): 1525–1533..
641 Published online 2013 Oct 23. doi:  10.1016/S0140-6736(13)61809-7.
642
643 7. Robert F. Siliciano and Warner C. Greene. HIV Latency. Cold Spring Harb Perspect
644 Med. 2011 Sep; 1(1): a007096. doi:10.1101/cshperspect.a007096.
645

8. Davey RT Jr, Bhat N, Yoder C, Chun TW, Metcalf JA, Dewar R, Natarajan V, Lempicki RA, Adelsberger JW, Miller KD, Joseph A. Kovacs, Michael A. Polis, Robert E. Walker, Judith Falloon, Henry Masur, Dennis Gee, Michael Baseler, Dimiter S. Dimitrov, Anthony S. Fauci and H. Clifford Lane. 1999. HIV-1 and T cell dynamics after interruption of highly active antiretroviral therapy (HAART) in patients with a history of sustained viral suppression. Proc Natl Acad Sci 96: 15109–15114, https://doi.org/10.1073/pnas.96.26.15109.

9. Kim HJ, Bae SC. Histone deacetylase inhibitors: molecular mechanisms of action and clinical trials as anti-cancer drugs. Am J Transl Res (2011) 3(2):166–79.

10. Ylisastigui L, Archin NM, Lehrman G, Bosch RJ, Margolis DM. Coaxing HIV-1 from resting CD4 T cells: histone deacetylase inhibition allows latent viral expression. AIDS (2004) 18(8):1101–8.10.1097/00002030-200405210-00003.

11. Shirakawa K, Chavez L, Hakre S, Calvanese V, Verdin E. Reactivation of latent HIV by histone deacetylase inhibitors. Trends Microbiol (2013) 21(6):277–85.10.1016/j.tim.2013.02.005.

12. Manson McManamy ME, Hakre S, Verdin EM, Margolis DM. Therapy for latent HIV-1 infection: the role of histone deacetylase inhibitors. Antivir Chem Chemother (2014) 23(4):145–9.10.3851/imp2551
.
13. Wei DG, Chiang V, Fyne E, Balakrishnan M, Barnes T, Graupe M, et al. Histone deacetylase inhibitor romidepsin induces HIV expression in CD4 T cells from patients on suppressive antiretroviral therapy at concentrations achieved by clinical dosing. PLoS Pathog (2014) 10(4):e1004071.10.1371/journal.ppat.1004071).
14. Deeks SG , Nature. HIV: Shock and kill. 2012; 487(7408):439-40. doi:10.1038/487439a.

15. Williams SA, Chen LF, Kwon H, Fenard D, Bisgrove D, Verdin E, and Greene WC. Prostratin antagonizes HIV latency by activating NF-kappaB J Biol Chem. 2004 Oct 1; 279(40):42008-17. doi: 10.1074/jbc.M402124200.

16. Mehla R, Bivalkar-Mehla S, Zhang R, Handy I, Albrecht H, Giri S, Nagarkatti P, Nagarkatti M, and Chauhan A. Bryostatin modulates latent HIV-1 infection via PKC and AMPK signaling but inhibits acute infection in a receptor independent manner. PLoS One. 2010 Jun 16; 5(6):e11160. doi: 10.1371/journal.pone.0011160.

17. Pandeló José D, Bartholomeeusen K, da Cunha RD, Abreu CM, Glinski J, da Costa TB, Bacchi Rabay AF, Pianowski Filho LF, Dudycz LW, Ranga U, Peterlin BM, Pianowski LF, Tanuri A, Aguiar RS. Reactivation of latent HIV-1 by new semi-synthetic ingenol esters. Virology. 2014 Aug;462-463:328-39. doi: 10.1016/j.virol.2014.05.033.

18. Sergey A. Trushin, Gary D. Bren, Susana Asin, Kevin N. Pennington, Carlos V. Paya, and Andrew D. Badley. Human Immunodeficiency Virus Reactivation by Phorbol Esters or T-Cell Receptor Ligation Requires both PKCα and PKCθ. J. Virol. August 2005 vol. 79 no. 15 9821-9830. doi: 10.1128/JVI.79.15.9821-9830.2005.

19. Savarino A, Mai A, Norelli S, El Daker S, Valente S, Rotili D, Altucci L, Palamara AT, Garaci E. "Shock and kill" effects of class I-selective histone deacetylase inhibitors in combination with the glutathione synthesis inhibitor buthionine sulfoximine in cell line models for HIV-1 quiescence. Retrovirology. 2009 Jun 2;6:52. doi: 10.1186/1742-4690-6-52.

20. Jiang G, Dandekar S. Targeting NF-kappaB signaling with protein kinase C agonists as an emerging strategy for combating HIV latency. AIDS Res Hum Retroviruses (2015) 31(1):4–12.10.1089/aid.2014.0199

21. Kashiwada, Y., Wang, H.-K., Nagao, T., Kitanaka, S., Yasuda, I., Fujioka, T. Takashi Yamagishi, L. Mark Cosentino, Mutsuo Kozuka, Hikaru Okabe, Yasumasa Ikeshiro, Chang-Qi Hu, Eric Yeh, and Kuo-Hsiung Lee. (1998). Anti-AIDS agents. 30. anti-HIV activity of oleanolic acid, pomolic acid, and structurally related triterpenoids. J. Nat. Prod. 61, 1090–1095. doi: 10.1021/np9800710.

22. Ohtake, N., Nakai, Y., Yamamoto, M., Sakakibara, I., Takeda, S., Amagaya, S., Aburada M (2004). Separation and isolation methods for analysis of the active principles of Sho-saiko-to (SST) oriental medicine. J. Chromatogr. B 812, 135–148. doi: 10.1016/j.jchromb.2004.06.051.

23. Loya, S., Rudi, A., Kashman, Y., and Hizi, A. (1999). Polycitone A, a novel and potent general inhibitor of retroviral reverse transcriptases and cellular DNA polymerases. Biochem. J. 344, 85–92. doi: 10.1042/bj3440085.

24. Dutra RC, Campos MM, Santos AR, Calixto JB. Medicinal plants in Brazil: Pharmacological studies, drug discovery, challenges and perspectives. Pharmacological research. 2016.

25. Franco-Salla GB, Prates J, Cardin LT, Dos Santos AR, Silva WA, Jr., da Cunha BR, et al. Euphorbia tirucalli modulates gene expression in larynx squamous cell carcinoma. BMC complementary and alternative medicine. 2016;16(1):136.

26. Lin MW, Lin AS, Wu DC, Wang SS, Chang FR, Wu YC, et al. Euphol from Euphorbia tirucalli selectively inhibits human gastric cancer cell growth through the induction of ERK1/2-mediated apoptosis. Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association. 2012;50(12):4333-9.

27. Wang L, Wang G, Yang D, Guo X, Xu Y, Feng B, et al. Euphol arrests breast cancer cells at the G1 phase through the modulation of cyclin D1, p21 and p27 expression. Molecular medicine reports. 2013;8(4):1279-85.

28. Fatope MO, Zeng L, Ohayaga JE, Shi G, McLaughlin JL. Selectively cytotoxic diterpenes from Euphorbia poisonii. Journal of medicinal chemistry. 1996;39(4):1005-8.

29. Prakash E, Gupta DK. Cytotoxic Activities of Extracts of Medicinal Plants of Euphorbiacae Family Studied on Seven Human Cancer Cell lines. Universal Journal of Plant Science. 2013;1(4):113-7.

30. El-Mekkawy, S., Meselhy, M. R., Nakamura, N., Hattori, M., Kawahata, T., and Otake, T. (2000). Anti-HIV-1 phorbol esters from the seeds of Croton tiglium. Phytochemistry 53, 457–464. doi: 10.1016/S0031-9422(99)00556-7.

31. Ortencio, W.B. Medicina popular do Centro-Oeste. 2.ed. Brasília: Thesaurus, 1997. 59p.

32. R. Bagavathi, B. Sorg and E. Hecker (1988). Tigliane-type diterpene esters from Euphorbia umbellatai, Planta Med. 54, 506–510.

33. G. W. Olivier, M. G. Rowan, S. K. Branch, M. F. Mohan and K. C. Molloy (1992). Two esters of synadenol, a new lathyrane diterpenoid, from the latex of Synadenium compactum (Euphorbiaceae): a crystal structure analysis, J. Chem. Soc. Perkin Trans. 1, 1831–1835.

34. B. Uzabakilihoc, C. Largeau and E. Casadwall (1987). Latex constituents of Euphorbia candelabrum, E. grantii, E. tirucalli and Euphorbia umbellatai, Phytochemistry 26, 3041–3045.

35. Jordan A1, Bisgrove D, Verdin E. HIV reproducibly establishes a latent infection after acute infection of T cells in vitro. EMBO J. ;22(8):1868-77. 10.1093/emboj/cdg188.

36.  Martiny-Baron Kazanietz MG, Mischak H, Blumberg PM, Kochs G, Hug H, and Marmé D, Schächtele C. (1993) Selective inhibition of protein kinase C isozymes by the indolocarbazole Gö 6976 J.Biol.Chem. *268* 9194.

37. Gschwendt Dieterich S, Rennecke J, Kittstein W, Mueller HJ, and Johannes FJ. (1996) Inhibition of protein kinase C μ by various inhibitors. Differentiation from protein kinase c isozymes. FEBS Lett.

38. Sullivan JA[1], Merritt JE, Budd JM, Booth RF, andHallam TJ. Effect of a selective protein kinase C inhibitor, Ro 31-8425, on Mac-1 expression and adhesion of human neutrophils. Eur J Immunol. 1994 Mar;24(3):621-6.

39. Laird G. M. et al. . Rapid quantification of the latent reservoir for HIV-1 using a viral outgrowth assay. PLoS Pathog 9, e1003398 (2013).

40. Valentine LE, Loffredo JT, Bean AT, León EJ, MacNair CE, Beal DR, Piaskowski SM, Klimentidis YC, Lank SM, Wiseman RW, Weinfurter JT, May GE, Rakasz EG, Wilson NA, Friedrich TC, O'Connor DH, Allison DB, Watkins DI. 2009. Infection with "escaped" virus variants impairs control of simian immunodeficiency virus SIVmac239 replication in Mamu-B*08-positive macaques. J Virol. 83:11514-11527. PMC2772717.

41. Kinghorn AD. Major skin-irritant principle from Euphorbia umbellatai. Journal of pharmaceutical sciences. 1980;69(12):1446-1447. Doi: 10.1002/jps.2600691228.

42. Hartmann DB, Marim RA, da Silva YL, Zardeto G, de Azevedo Silva I, de Arruda Mattos D, et al. Letalidade do extrato de Euphorbia umbellatai Hook. f.(Euphorbiaceae) frente a caramujos *Biomphalaria glabrata* say, 1818 (Gastropoda, Planorbidae). Arquivos de Ciências Veterinárias e Zoologia da UNIPAR, 2011;14(1). doi: 10.25110/arqvet.v14i1.2011.3779.

43. Costa LL, VC, David R, Pinto BR, Minozzo VA, Kozlowski Jr LA, Campos RZ, et al., Anti-ulcer activity of Euphorbia umbellatai latex. Revista Brasileira de Farmacognosia, 2012;22(5):1070-1078. Doi: 10.1590/S0102-695X2012005000050.

44. Hassan EM, Mohammed MM, Mohamed SM. Two new phorbol-type diterpene esters from Euphorbia umbellatai Hook f. leaves. *Records of Natural Products*, 2012;6(2):255.

45. Campos A, Vendramini-Costa DB, Longato GB, Zermiani T, Ruiz ALTG, de Carvalho JE, et al. Antiproliferative effect of Euphorbia umbellatai Hook f. stems (Euphorbiaceae) and a rare phorbol diterpene ester. *International journal of toxicology*, 2016;35(6):666-671. doi: 10.1177/1091581816659660.

810   46. Mochly-Rosen D, Gordon AS. Anchoring proteins for protein kinase C: a means for
811   isozyme selectivity. FASEB J. 1998;12:35–42.
812

813   47. Spitaler M, Cantrell D. a. Protein kinase C and beyond. Nat. Immunol. 2004;5:785–
814   790. http://dx.doi.org/10.1038/ni1097
815

816   48. Steinberg SF. Structural basis of protein Kinase C isoform function. Physiol.
817   Rev. 2008;88:1341–1378. http://dx.doi.org/10.1152/physrev.00034.2007.
818

819   49. Williams SA, Chen L-F, Kwon H, Fenard D, Bisgrove D, Verdin E, Greene WC.
820   Prostratin antagonizes HIV latency by activating NF-kappaB. J. Biol.
821   Chem. 2004;279:42008–42017. http://dx.doi.org/10.1074/jbc.M402124200).
822

50. Bachu M, Yalla S, Asokan M, Verma A, Neogi U, Sharma S, Murali RV, Mukthey
AB, Bhatt R, Chatterjee S, Rajan RE, Cheedarla N, Yadavalli VS, Mahadevan A,
Shankar SK, Rajagopalan N, Shet A, Saravanan S, Balakrishnan P, Solomon S,
Vajpayee M, Satish KS, Kundu TK, Jeang K-T, Ranga U. Multiple NF-κB sites in HIV-1
subtype C long terminal repeat confer superior magnitude of transcription and thereby
the enhanced viral predominance. J. Biol. Chem. 2012;287:44714–
44735. http://dx.doi.org/10.1074/jbc.M112.397158.
830

831   51. Naghavi MH, Schwartz S, Sönnerborg a., Vahlne a. Long terminal repeat
832   promoter/enhancer activity of different subtypes of HIV type 1. AIDS Res. Hum.
833   Retrovir. 1999;15:1293–1303. http://dx.doi.org/10.1089/088922299310197).
834

835   52. Sergey A. Trushin1, Gary D. Bren, Susana Asin, Kevin N. Pennington, Carlos V.
836   Paya, and Andrew D. Badley, Human Immunodeficiency Virus Reactivation by Phorbol
837   Esters or T-Cell Receptor Ligation Requires both PKCα and PKCθ. Virol. 2005 vol. 79
838   no. 15 9821-9830. doi: 10.1128/JVI.79.15.9821-9830.2005.
839

840   53. Tanaka M , Irie K , Nakagawa Y , Nakamura Y , Ohigashi H , Wender PA . The C4
841   hydroxyl group of phorbol esters is not necessary for protein kinase C binding. Bioorg
842   Med Chem Lett. 2001 Mar 12;11(5):719-22doi.org/10.1016/S0960-894X(01)00045-2.
843

844   54. Blackhall FH, Ranson M, Radford JA, Hancock BW, Soukop M, McGown
845   AT, Robbins A, Halbert G, Jayson GC; Cancer Research Campaign Phase I/II
846   Committee. A phase II trial of bryostatin 1 in patients with non-Hodgkin's lymphoma. Br
847   J Cancer. 2001;84(4):465-9. doi: 10.1054/bjoc.2000.1624.
848

849   55. Paiardini M, Müller-Trutwin M. HIV-associated chronic immune activation. Immunol
850   Rev. 2013; 254(1): 78–101.
851

852   56. Luiz C. Cunha, Flaubertt S. Azeredo, Ana C. V. Mendonça, Marcelo S. Vieira,
853   Liuba L. Pucci, Marize C. Valadares, Helemi O. G. Freitas, Ângela A. S. Sena, and Ruy
854   de Souza Lino Junior. Evaluation of acute and subacute toxicity in rats of leaf and latex
855   ethanolic extract of Synadenium umbellatum Pax.Brazilian Journal of Pharmacognosy,
856   2009; 19(2A): 403-411.
857

858   57. Isakov N, Altman A. Protein kinase C(theta) in T cell activation. Annu Rev
859   Immunol. 2002;20:761-94.
860

861   58. Porichis F, Kaufmann DE. HIV-specific CD4 T cells and immune control of viral
862   replication. Curr Opin HIV AIDS. 2011; 6(3):174-80.

863  59. Kared H, Fabre T, Bedard N, Brunneau J, Shoukry NH. Galectin-9 and IL-21
864  mediate cross-regulation between Th17 and Treg cells during acute hepatitis C. PLoS
865  Pathog. 9 (2013) e1003422.

866
867
**Acknowledgments**

We thanks Dr. Thomas Friedrich from Virology Services Unit  at the Wisconsin National

Primate Research Center, Wisconsin, USA for QVOA assay using SIVmac239 in NHP

CD4$^+$ Lymphocytes. We are also grateful to Mr. Wilson Camargo Barros Filho and Mr.

Luis Antônio Nogueira for providing *E. umbellata*´s latex used in this work. This work

was sponsored by CNPq, FAPERJ, CAPES and FINEP.

 **FIGURES AND LEGENDS**

**Figure 1**. **Isolation and chemical structures of JALEx.** Gas chromatograms of (A) whole extract JALEx and (B) Phorbol esters-enriched JALEx fractions. T1 and T2 = major triterpenes constituents; Ph1 and Ph2 = major phorbol esters constituents. The chemical classes were indicated by the characteristic mass fragmentation in the GC-MS analysis and data from literature.

**Figure 2: JALEx reactivates virus transcription in HIV-1 latent models.** J-Lat 8.4 (A) and 10.6 (C) cells were exposed to increasing concentrations of JALEx and HIV-1 activation was assessed 24 hours post-treatment by GFP positive cells screening by flow cytometry. Mock stands for not treated. PMA (1 µM) and INGB (0,32 µM) were used as positive induction controls. The cytotoxicity of JALEx was evaluated using CellTiter Blue vital stain 5 days after treatment in J-Lat 8.4 (B) and 10.6 cells (D). GFP positivity and cytotoxicity were expressed as percentage of cells normalized by mock non-treated cells (n=3). The dashed lines in Figure 2 B and D  represents 100% of viability measures in MOCK. The bars in the figure represents 1 SD around the mean.

**Figure 3: JALEx -mediated HIV-1 reactivation occurs via PKC activation.** J-Lat cells 8.4 (left panel) and J-Lat 10.6 (right panel) were pretreated for 24 h with three different PKC inhibitors (G6666, G6363and Ro-31-8220) at the concentration of 1 µM each. The cells were then incubated with different concentrations of JALEx for an additional 24 hours and GFP expression was assessed by flow cytometry. PMA (1 µM) was used as a positive control for activation of PKC-dependent HIV-1 (n=1).

**Figure 4: JALEx is responsible for the cellular relocation of PKC isoforms.** Cellular location analysis of different PKC (alpha, delta and gamma) isoforms after treatment of HeLa cells with two concentrations of JALEx: 1 μg / mL and 0.1 μg / mL. (A) HeLa cells were transfected with 3 different PKC isoforms for 24 h. After this time the cells were treated for 10 minutes with the two concentrations of JALEx. (B) HeLa cells were transfected with 3 different PKC isoforms for 24 h. After this time the cells were treated for additional 24 hours with the two concentrations of JALEx. As a positive control, PMA (1 μM) was used (n=3).

**Figure 5: . JALEx -mediated transcriptional activation is dependent on NF-κB activation**.
HeLa cells were treated with two concentrations of the **JALEx** (1 and 0.1 µg/mL) for 6 hours (A)
and 24 hours (B). After these time intervals, the cells were subjected to immunofluorescence
for labeling of NF-κB, in red and cell nucleus, in blue. As a positive control, PMA (1 µM) was
used. (C) Graph of the quantification of cells showing NF-κB  translocation. An average value
of the fluorescence intensity between the nucleus and the cytoplasm of 100 HeLa cells was
calculated under all conditions, at the time intervals of 6 and 24 h after the treatments (n=3;
Two-way ANOVA and (***) indicate p<0,0001). (D) Jurkat cells were transfected with
pBlue30LTR-Luc (NF-κB WT) and the mutant construct, pBlue30LTR NF-κB MUT-Luc (NF-κB
MUT), absent from the NF-κB binding site, preventing its interaction with NFκB. After
transfections, the cells were treated with 1µg / mL of the **JALEx** for 24 hours and luciferase
expression was measured after lysis of the cells. The fold-change of the luciferase expression
between the wild-type plasmid and the mutant is shown in the graph (n=4, Mann-Whitney test
and (*) indicate p<0,05). The bars in Figure 5 C and D represents 1 SD around the mean.

**Figure 6: JALEx modulates the expression of cell surface receptors, activation markers and cytokine secretion in primary CD4 + T cells from HIV- individuals.** TCD4 + cells were isolated from PBMCs from 4 healthy donors and treated for 24 hours with JALEx. After this period, the expression of some surface molecules was evaluated. (A) Expression of the CD4 receptor and CCR5 and CXCR4 co-receptors in CD4 + T cells isolated after treatment with JALEx (n=4). (B) Expression of CD25, CD38 and HLA-DR activation markers in CD4 + T cells isolated after treatment with JALEx (n=4). (C) Expression of the CD69 activation marker on CD4 + T cells isolated after treatment with janauba extract. The black bar represents the positive control (PMA 1 µM) and the gray bar represents the treatment with 10 µg / mL of JALEx (n=4). (D) CD4+ T cells were treated with 1 µM PMA, 10 µg/mL (-3) or 1 µg/mL (-4) of JALEx for 24 hours. After, supernatants from the CD4+ T cells were collected and cytokine analysis were performed by Bio-plex Pro Human Cytokines 17-plex platform (n=5, Kruskal-Wallis test and (*) indicate p<0.05). The bars in the figure represents 1 SD around the mean.

**Figure 7. Effect of JALEx on T cell viability, HIV-1 replication in PBMC cultures and**
**cytokine secretion in T cells from AIDS patients.** Different concentrations of JALEx (0.01, 0.1
and 1 μg/mL) were added to PBMC cultures (1 x 10$^6$/mL) obtained from 15 AIDS patients, and
24h after, the (A) the number of HIV-1 into supernatants was determined through RT-qPCR, (B)
CD4$^+$ and CD8$^+$ T cells viability was analyzed by FACS after 7-AAD staining. AIDS-derived
PBMC cultures (1 x 10$^6$/mL, n=15) were maintained for 24h in the presence of medium alone or
with different JALEx ( 0.01 and 0.1 μg/mL). As positive control, some well receiving PMA plus
ionomycine (PMA/IO). The percentage of CD4+ and CD8+ T cells producers of IFN-γ, IL-17, IL-
21 and IL-10 was determined by cytometry (C). The mean values were compared and the p
values are shown in the figures. In (D), the proportion of different (CD4+ and CD8+) T cell
subsets of secreting different combinations of IL-17, IFN-γ and IL-21 in response to JALEx (0.1
μg/mL) was also evaluated by cytometry. In the figure, (*), (**), and (***) indicate p<0.05,
p<0.001 and p<0.0001, respectively. The bars in the figure represents 1 SD around the mean.
900

**SUPPLEMENTAL MATERIAL**

**Supplemental Figure 1**

**Figure S1:** Phorbol esters isolated from *Euphorbia umbellata*. I = 12-*O*-Tigloyl-4-deoxyphorbol-
13-isobutyrate; II = Phorbol-12,13,20-triacetate; III = 12-Deoxyphorbol-13-(12-
methylpropionate); IV = 3,4,12,13-Tetraacetylphorbol-20-phenylacetate (synagrantol A); V =
Deoxyphorbol-12,13-ditiglate (synagrantol B).

**Figure S2. Panel of phosphorylation of different PKC isoforms after treatment with JALEx.** (A) Jurkat cells were treated with two different concentrations of JALEx (0.01 μg / mL and 0.001 μg / mL), PMA (1 μM) as positive control, at different time intervals (10, 30 minutes, 1, 6 and 24 hours). Then the cells were lysed for western blotting with phosphorylated anti-PKC antibodies (pan, δ, θ) and anti-tubulin as the loading control. (A) The intensity of the western blotting bands corresponding to the Jurkat cells in (B) that were quantified by densitometry with the aid of the Image J program. Dashed lines correspond to the band intensity of MOCK in these experiments (n=1).

935 **Supplemental Table 1**

937 **Table S1- QVOA results using macaques CD4+ T Lymphocytes extracted from**
938 **SIV 251 infected Macaques treated with HAART therapy.**

| Cells/well[a] | No mitogen | Con A | Ingenol B | **JALEx** |
|---|---|---|---|---|
| 1.25e5 | + / - / -[b] | - / - / - | + / - / + | + / + / + |
| 6.25e4 | - / - / + | - / + / + | + / + / - | + / + / + |
| 3.1e4 | + / - / + | - / - / + | - / + / + | + / + / + |
| Maximum likelihood (IUPM) | 7.735675 | 5.229369 | 15.525127 | **34.986678** |
| 95% CI | {2.866090, 20.878852} | {1.680200, 16.275622} | {6.619641, 36.411279} | {12.619641, 66.411279} |

939 [a]- Number of CD4+ T Lymphocytes extracted from SIV 251 infected Macaques treated with
940 HAART therapy inoculated in each well. [b]- Number of well with p27 antigen positive. Con A: