A Stochastic Model for Cancer Metastasis: Branching Stochastic Process with Settlement

Christoph Frei

Department of Mathematical and Statistical Sciences
University of Alberta, Edmonton, Alberta, T6G2G1, Canada
cfrei@ualberta.ca

Thomas Hillen

Department of Mathematical and Statistical Sciences
University of Alberta
thillen@ualberta.ca

Adam Rhodes

Department of Mathematical and Statistical Sciences
University of Alberta
arhodes@ualberta.ca

This version: April 2, 2018

We introduce a new stochastic model for metastatic growth, which takes the form of a *stochastic branching process with settlement*, whose particles are interpreted as clusters of cancer cells. The analysis of expected particle location, their locational variance, the furthest particle distribution, and the extinction probability leads to a common type of differential equation, namely, a non-local integro-differential equation with distributed delay. We prove global existence and uniqueness results for this type of equation. The solutions' asymptotic behavior for long time is characterized by an explicit index, a *metastatic reproduction number* R_0, which divides between metastatic spread for $R_0 > 1$ versus extinction for $R_0 < 1$. Using metastatic data from mouse experiments, we show the suitability of our framework to model metastatic cancer.

Keywords: branching processes, birth-jump processes, cancer metastasis, integro-differential equations, reproduction number

AMS Subject Classification: 60J85, 35Q92, 92B05

1. Introduction

Metastasis is the leading causes of cancer related deaths. It is the process by which cancer spreads from a primary site to distant secondary sites. Because of its implication in over 90% of all cancer-related deaths,[11,31] it is recognized as one of the *hallmarks* of cancer.[12] The *metastatic cascade*[11,31,26] provides a mechanistic

description of the process, viewing it as an ordered sequence of biological events. Growth of the primary tumor leads to local invasion of surrounding normal tissue. Such local invasion eventually encounters a vessel of the lymphatic or circulatory system. Individual cancer cells — or small clusters of cells — can enter these vessels (intravasate), thereby gaining access to rapid transport throughout the body. If these circulating cells survive their journey and exit the vessel at some distant site (extravasate), they may be able to establish a micrometastasis. Evasion of local defenses and adjustment to the hostile foreign environment can see the *micro*metastasis grow into a *macro*metastasis. Further growth may eventually lead to a secondary tumor.

The stochastic nature of metastasis makes experimental studies difficult, meaning that a thorough understanding of the process is elusive. While some of the steps in the metastatic cascade are well understood — for example, growth and invasion of primary tumors are well studied both experimentally and theoretically (see the reviews[26,27]) — understanding of others remains elusive. Travel *to*, and establishment *at* secondary sites are particularly poorly understood, with many theoretical investigations ignoring these aspects altogether. Though recent results[21] have led to novel theories that metastasis may be more intricately orchestrated than previously thought,[28] the process is still believed to be largely stochastic.

Based on these observations, we introduce a new stochastic framework for metastatic spread in the form of a *branching stochastic process with settlement*. This model captures simultaneously temporal and locational dynamics. Stationary tumors emit small clusters of cells at random times, which randomly move through the body, modeled by a stochastic process generalizing Brownian motion. The moving clusters — also known as circulating tumor cells/clusters (CTCs) — either die or settle, both cases happening randomly according to given rates. If the CTCs successfully settle, they may establish a secondary tumor, which itself may shed new CTCs into the blood stream. Moving and stationary groups of cancer cells die at different rates.

In contrast to existing stochastic metastasis models, our framework accounts for both travel between primary and secondary sites and establishment at the secondary sites. Hartung and Gomez[13] propose a stochastic model with secondary metastatic emission as a cascade of Poisson point processes and link it to the deterministic model introduced by Iwata et al.[19]. Differently from us, Hartung and Gomez[13] model only the size development over time and not the location.

From a probabilistic viewpoint, our model generalizes branching Brownian motion to arbitrary dynamics and the inclusion of settlement and death of particles. Branching Brownian motion has been analyzed for more than fifty years, starting with seminal work[18,25] on the fundamental link between branching Brownian motion and the Fisher-Kolmogorov-Petrovsky-Piscounov equation. Since then branching Brownian motion has been intensively studied in its own right. In statistical mechanics, branching Brownian motion is used for models of spin glasses.[1] Recently, branching processes have been found useful in simulating semi-linear partial

differential equations.[14,15] To the best of our knowledge, this paper is the first to introduce branching processes with settlement and study their applications in the context of cancer metastasis.

For the branching stochastic process with settlement, we prove two sets of main results. The first set deals with the characterization of the following key quantities in our stochastic model: the expected location F of metastases, their locational variance V, the distribution H of the furthest invading metastasis, the metastatic extinction probability Q, and the ratio of moving versus stationary cancer cell groups. We show that each of the first four quantities — F, V, H, and Q — satisfies a non-local integro-differential equation with distributed delay. These non-local differential equations are of the same type for all cases and are non-linear in some cases. We present a detailed analysis of these non-local differential equations, including positivity, and local and global existence of solutions.

The second set of main results provides a simple classification of the asymptotic behavior. Despite the model complexity and stochasticity, we are able to find a *metastatic reproduction number* R_0, which is explicitly given in terms of the rates of metastatic shedding, settlement, and death. We prove a global stability result of extinction of metastases for $R_0 < 1$ and growth for $R_0 > 1$. By parametrizing the model on experimental mouse metastatic data, we numerically explore the resulting metastatic pattern and confirm the suitability of our framework to model metastatic cancer.

The remainder of this paper is organized as follows. Section 2 introduces the model, presents the main results and relates the stochastic model to integro-differential equations. In Section 3, we analyze these integro-differential equations in more detail and characterize their asymptotic behavior using the metastatic reproduction number. Section 4 shows how our framework can be applied in the context metastatic cancer while Section 5 concludes and discusses possible extensions.

2. Branching stochastic processes with settlement

We define the *branching stochastic process with settlement* as follows: We start with one tumor that is located at locational position 0. We assume that the tumor sheds individual cells and small groups of cells into the circulatory system. It is believed that such CTCs are most responsible for metastasis formation.[9] Hence, in this model, we focus on the shedding of cell clusters. At a random time ν, the primary tumor emits a cell cluster, which starts moving randomly. We model the movement of this cell cluster by a stochastic process $(B(t))_{t\geq 0}$ with cumulative distribution function $G(t, x) = P[B(t) \leq x]$ for $x \in \mathbb{R}$ and $t \geq 0$. We assume that for every fixed t, $G(t, .)$ is absolutely continuous so that there is a density function $g(t, .)$ with $G(t, x) = \int_{-\infty}^{x} g(t, y) \, dy$ for all $x \in \mathbb{R}$. The prime example for $(B(t))_{t\geq 0}$ is Brownian motion, in which case g is the heat kernel $g(t, y) = \frac{1}{\sqrt{2\pi t}} e^{-y^2/(2t)}$, but our model allows for general stochastic processes $(B(t))_{t\geq 0}$. The primary tumor stays at the same location until it dies. We assume that the shedding time ν is exponentially

distributed, $\nu \sim \mathrm{Exp}(\mu)$ for some parameter $\mu \geq 0$, using the convention that $\nu = \infty$ if $\nu \sim \mathrm{Exp}(0)$.

After an additional random time τ (namely, at time $\nu + \tau$), the cell cluster settles down. We assume that τ is exponentially distributed, $\tau \sim \mathrm{Exp}(\lambda)$ for some parameter $\lambda \geq 0$. When the cell cluster has settled down, it will take again a time which is exponentially distributed with parameter $\mu \geq 0$ until it has grown enough to emit another cell cluster. Also the first cell cluster will continue to emit further cell clusters after a time which is exponentially distributed with the same parameter $\mu \geq 0$, independently of the other random variables and processes. Cell clusters that are moving are destroyed at a rate $\delta_1 \geq 0$ while stationary clusters die at a rate $\delta_2 \geq 0$, again independently of the other clusters, the movement and the growth. The process is repeated ad infinitum. Figure 1 illustrates our model.

Figure 1. Illustration of a branching stochastic process with settlement.

We now extend the notion from cell clusters to particles since the branching stochastic process with settlement is also relevant to other applications such as seed dispersal, epidemic spread, and forest fire spread. In those cases, particles would refer to plant seeds, infectious agents, and burning branches, respectively. In this paper, our focus is on cell clusters and micrometastasis.

We denote by $N(t)$ the number of particles born before time t. Their positions at time t are $X_i(t)$ for $1 \leq i \leq N(t)$ where we enumerate the particles by their birthdates. For fixed t and i, $N(t)$ and $X_i(t)$ are random variables with values in $\mathbb{N}$ and $\mathbb{R}$, respectively. We denote by $M(t)$ the number of particles alive at time t. We are interested in the following quantities:

- expected location $F(t,x) = E\big[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t) \leq x}\big]$: the expected number of particles located in $(-\infty, x]$ at time t,

- locational variance $V(t,x) = \operatorname{Var}\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\right]$: the variation in the number of particles located in $(-\infty, x]$ at time t due to randomness,

- furthest particle distribution $H(t,x) = P[\max_{i=1,\ldots,N(t)} X_i(t) \leq x]$: the probability that all particles are located in $(-\infty, x]$ at time t,

- survival probability $Q(t) = P[M(t) > 0]$: the probability that there is at least one particle alive at time t,

where we use the convention that $\mathbb{1}_{X_i(t)\leq x}$ is zero for all x and times t after the death time of particle i. We set $H(t,x) = 0$ for all x if no more particle exists at time t.

2.1. *Characterization via integro-differential equations*

We use the above branching stochastic process with settlement to find a common type of equation for the expected location F, the variance V, the distribution of the furthest particle H and the survival probability Q.

Theorem 2.1. *Consider the branching stochastic process with settlement defined above. The quantities F, V, H, Q satisfy*

$$F_t(t,x) = -\delta_2 F(t,x) + \mu e^{-(\lambda+\delta_1)t} G(t,x)$$
$$+ \lambda\mu \int_0^t \int_{-\infty}^\infty F(t-u, x-z) g(u,z) e^{-(\lambda+\delta_1)u} \, dzdu, \qquad (2.1)$$

$$V_t(t,x) = -\delta_2 V(t,x) + \mu e^{-(\lambda+\delta_1)t} h(t,x)$$
$$+ \lambda\mu \int_0^t \int_{-\infty}^\infty V(t-u, x-z) g(u,z) e^{-(\lambda+\delta_1)u} \, dzdu, \qquad (2.2)$$

with

$$h(t,x) = G(t,x) + e^{(\lambda+\delta_1)t} \frac{\delta_2}{\mu} F^2(t,x)$$
$$+ \lambda \int_0^t \int_{-\infty}^\infty F^2(t-u, x-z) g(u,z) e^{(\lambda+\delta_1)(t-u)} \, dzdu,$$

$$H_t(t,x) = \left[\mu e^{-\lambda t}(1 - e^{-\delta_1 t}(1 - G(t,x))) - (\mu+\delta_1)\right] H(t,x)$$
$$+ \lambda\mu H(t,x) \int_0^t \int_{-\infty}^\infty H(t-u, x-z) g(u,z) e^{-\lambda u} \, dzdu, \qquad (2.3)$$

$$Q'(t) = -\left(\delta_2 + \mu e^{-(\lambda+\delta_1)t}\right) Q(t) + \mu e^{-(\lambda+\delta_1)t}$$
$$+ \lambda\mu(1 - Q(t)) \int_0^t Q(t-s) e^{-(\lambda+\delta_1)s} \, ds. \qquad (2.4)$$

Case	r_1	r_2	$q(t,x)$	$p(t,x)$	$k(u,z)$
F	1	0	δ_2	$\mu e^{-(\lambda+\delta_1)t}G(t,x)$	$\lambda\mu e^{-(\lambda+\delta_1)u}g(u,z)$
V	1	0	δ_2	$\mu e^{-(\lambda+\delta_1)t}h(t,x)$	$\lambda\mu e^{-(\lambda+\delta_1)u}g(u,z)$
H	0	1	$q_H(t,x)$	0	$\lambda\mu e^{-\lambda u}g(u,z)$
Q	1	-1	$\delta_2 + \mu e^{-(\lambda+\delta_1)t}$	$\mu e^{-(\lambda+\delta_1)t}$	$\lambda e^{-(\lambda+\delta_1)u}$

Table 1. Parameter values for the different cases of expected location F, locational variance V, distribution of the furthest particle H and survival probability Q, using the abbreviation $q_H(t,x) = \mu + \delta_1 - \mu e^{-\lambda t}(1 - e^{-\delta_1 t}(1 - G(t,x)))$.

Before we present the proof of Theorem 2.1 in Section 2.3, we bring the above four equations (2.1)–(2.4) into a common form using a convolution notation. We encounter integral kernels that depend on one variable and on two variables. We use the same convolution symbol for both cases; given two kernels $k_1(t,x)$ and $k_2(t)$ and two test functions $f(t,x)$ and $g(t)$, we denote

$$k_1 * f(t,x) := \int_0^t \int_{-\infty}^{\infty} f(t-u, x-z)k(u,z)\,\mathrm{d}z\mathrm{d}u$$

$$k_2 * g(t) := \int_0^t g(t-u)k(u)\,\mathrm{d}u.$$

Then we combine equations (2.1)–(2.4) in the compact form

$$f_t = -qf + p + (r_1 + r_2 f)\, k * f, \tag{2.5}$$

where we identify parameters and functions as shown in Table 1.

In Section 3, we will analyze the general non-local integro-differential equation with distributed delay (2.5) and show positivity as well as local and global existence of solutions using a fixed-point argument. For the qualitative analysis, we encounter an index R_0, which can be understood as the *metastatic reproduction number*:

$$R_0 = \frac{\lambda}{\delta_1 + \lambda}\, \frac{\mu}{\delta_2}. \tag{2.6}$$

The fist factor $\lambda/(\delta_1 + \lambda)$ denotes the settlement rate divided by the sum of the settlement rate and the death rate during movement. It describes the odds of being still alive while settling. The second factor in (2.6), $\frac{\mu}{\delta_2}$, which refers to stationary particles, can be explained similarly, but here the effect of births of new particles needs to be considered. Indeed, when the ratio between shedding rate μ and total change rate $\delta_2 + \mu$ (stationary deaths and shedding) is summed over all potential offsprings of the particles, we obtain

$$\sum_{j=1}^{\infty}\left(\frac{\mu}{\delta_2+\mu}\right)^j = \sum_{j=0}^{\infty}\left(\frac{\mu}{\delta_2+\mu}\right)^j - 1 = \frac{1}{1-\mu/(\delta_2+\mu)} - 1 = \frac{\mu}{\delta_2},$$

regaining the second term in (2.6).

Before we go into the details, we paraphrase the main results, assuming $\mu > 0$ and $\lambda > 0$:

- ***Main result 1:*** *Under suitable assumptions (A1), defined later, the solutions to the above integro-differential equations (2.1)–(2.4) exist as mild solutions globally in time, whereby the two probabilities H and Q are bounded by 1. See Theorem 3.1 for global existence and uniqueness.*

- ***Main result 2:*** *The asymptotic behavior is characterized by R_0:*

  - *$R_0 < 1$: The expected number of particles shrinks exponentially and the particles die out asymptotically with certainty.*

  - *$R_0 > 1$: The expected number of particles grows exponentially and the asymptotic survival probability is given by $1 - 1/R_0 \in (0, 1)$.*

  - *$R_0 = 1$: The particles die out asymptotically with certainty while their number explodes on a set with vanishing probability so that their expected number converges to a finite limit $\frac{1 + \lambda/\delta_2}{\delta_2/\mu + \lambda/\delta_2}$.*

  See Lemma 3.3 for a threshold result of the survival probability Q and Corollary 3.1 for the asymptotic behavior of the expected particle number $E[M(t)]$.

Note that the survival probability in case $R_0 > 1$ equals $1 - 1/R_0$, which corresponds to the critical immunization threshold in common models for infectious diseases.[16]

2.2. *Instantaneous relocation*

In many biological situations, the time scale of movement is very different from the time scale of reproduction or death. Most mammals, for example, explore their environment on a daily basis, while they reproduce once per year or less. And even cells in tissues can travel relative large distances in minutes, while typical cell cycles are about 8–12 h or longer. Hence on the macroscopic time scale of birth and death, relocation appears to be instantaneous. To model this situation, we consider the limit $\lambda \to \infty$ in (2.5). When individuals are moving, they will jump immediately to a new location with density function of the distances travelled given by $g(0, x)$. We will then ignore the time in $g(0, x)$ and simply use a spatial relocation density $g(x)$.

For fixed t, we consider a common term in (2.5)

$$\lim_{\lambda \to \infty} \int_0^t \int_{-\infty}^{\infty} f(t - u, x - z) g(u, z) \lambda e^{-(\lambda + k_1)u} \, dz \, du = \int_{-\infty}^{\infty} f(t, x - z) g(0, z) \, dz.$$

Using integration by parts yields

$$\int_0^t \int_{-\infty}^\infty f(t-u, x-z)g(u,z)\lambda e^{-(\lambda+k_1)u}\,\mathrm{d}z\,\mathrm{d}u$$

$$= -\underbrace{\frac{\lambda}{\lambda+k_1}e^{-(k_1+\lambda)t}\int_{-\infty}^\infty h_0(x-z)g(t,z)\,\mathrm{d}z}_{\to 0 \text{ for } \lambda\to\infty} + \int_{-\infty}^\infty f(t, x-z)g(0,z)\,\mathrm{d}z$$

$$-\underbrace{\int_0^t \int_{-\infty}^\infty f_t(t-u, x-z)g(u,z)\frac{\lambda}{\lambda+k_1}e^{-(\lambda+k_1)u}\,\mathrm{d}z\,\mathrm{d}u}_{\to 0 \text{ for } \lambda\to\infty \text{ by monotone convergence}}.$$

Applying this scaling to the four equations (2.1)–(2.4) from Theorem 2.1 and using $g(x)$ instead of $g(0, x)$, we find

$$F_t(t, x) = -\delta_2 F(t, x) + \mu \int_{-\infty}^\infty F(t, x-z)g(z)\,\mathrm{d}z, \tag{2.7}$$

$$V_t(t, x) = -\delta_2 V(t, x) + \mu \int_{-\infty}^\infty V(t, x-z)g(z)\,\mathrm{d}z, \tag{2.8}$$

$$H_t(t, x) = -(\mu+\delta_1)H(t, x) + \mu H(t, x)\int_{-\infty}^\infty H(t, x-z)g(z)\,\mathrm{d}z, \tag{2.9}$$

$$Q'(t) = (\mu-\delta_2)Q(t)\left(1 - \frac{\mu Q(t)}{\mu-\delta_2}\right). \tag{2.10}$$

The equations for the instantaneous relocation case (2.7)–(2.9) are linear integral equations as they have been studied in the classical literature[22]. They are also of the form of *birth-jump processes* as introduced in Hillen et al.[17]. Hence the branching stochastic process with settlement appears as a generalization of birth-jump processes.

The metastatic reproduction number (2.6) becomes in this case

$$\lim_{\lambda\to\infty} R_0 = \frac{\mu}{\delta_2}$$

The survival probability Q satisfies a logistic differential equation (2.10), which can be analyzed quickly. If $R_0 < 1$, then the origin is asymptotically stable, and solutions converge to 0, metastasis die out. If $R_0 > 1$, then Q converges to a limit

$$\lim_{t\to\infty} Q(t) = 1 - \frac{1}{R_0}.$$

If $R_0 = 1$, then $Q(t) = Q(0)$ is constant.

This also corresponds to the classical result for the asymptotic extinction probability of a birth-death process; see, for example, Section 4.4 of Durrett[6]. Indeed, in the limit $\lambda \to \infty$, the number of particles evolves according to a Markov chain starting at 1 and with jump rates $q(i, i+1) = \mu i$ and $q(i, i-1) = \delta_2 i$ when there are i particles.

2.3. *Derivation of the integro-differential equations (proof of Theorem 2.1)*

The main idea to derive equations (2.1)–(2.4) from Theorem 2.1 is to use a renewal argument based on the recursive structure of branching stochastic processes with settlement. Since this goes similarly for the different functions, $F, V, H,$ and Q, we give a detailed proof in the case of F and mention afterwards how to start the proofs for V, H and Q, with the subsequent analysis being done analogously to that of F. We start by conditioning on time ν of the birth of the second particle, which means writing

$$F(t,x) = E\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\right] = \int_0^\infty E\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\Bigg|\nu = s\right]\mu e^{-\mu s}\,\mathrm{d}s. \qquad (2.11)$$

In the event $\{\nu > t\}$, the birth of the second particle happens only after time t so that at time t either only the first particle exists, which is located at zero, or no particle at all exists if the first particle has died before time t. We denote by $A_2(t)$ the event that the first particle has not died before time t, which has probability $P[A_2(t)] = \int_t^\infty \delta_2 e^{-\delta_2 s}\,\mathrm{d}s = e^{-\delta_2 t}$. For $s > t$, we have

$$E\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\Bigg|\nu = s\right] = E\left[\mathbb{1}_{A_2(t)}\mathbb{1}_{X_1(t)\leq x}\big|\nu = s\right] = e^{-\delta_2 t}\mathbb{1}_{0\leq x}.$$

On the other hand, in the event $\{\nu \leq t\}$, the second particle was born before time t if the first particle was then alive. We need to distinguish two cases on $\{\nu \leq t\}$. The first case is when the second particle has not yet settled down at time t. This happens when $\nu + \tau > t$ and then the distribution of $\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}$ conditional on $\nu = s$ and the first particle being alive at time s is equal to the sum of $\mathbb{1}_{A_1(t-s)}\mathbb{1}_{B(t-s)\leq x}$ and $\sum_{i=1}^{N(t-s)} \mathbb{1}_{X_i(t-s)\leq x}$, where $B(t-s)$ is an independent stochastic process at time $t-s$ (movement of second particle) with $A_1(t-s)$ the event that the second particle born at time s is alive at time t. The term $\sum_{i=1}^{N(t-s)} \mathbb{1}_{X_i(t-s)\leq x}$ arises because we can start again at time $\nu = s$ in the same way as at time zero thanks to the recursive structure and the memoryless property. The probability of the first particle being alive at time s is $e^{-\delta_2 s}$ while the survival probability of the second particle is

$$P[A_1(t-s)] = \int_{t-s}^\infty \delta_1 e^{-\delta_1 u}\,\mathrm{d}u = e^{-\delta_1(t-s)}.$$

Therefore, we have for $s \leq t$ and $u > t - s$ that

$$E\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\Bigg|\nu = s, \tau = u\right]$$

$$= e^{-\delta_2 s-\delta_1(t-s)}E\left[\mathbb{1}_{B(t-s)\leq x}\right] + e^{-\delta_2 s}E\left[\sum_{i=1}^{N(t-s)} \mathbb{1}_{X_i(t-s)\leq x}\right]$$

$$= e^{-\delta_2 s-\delta_1(t-s)}G(t-s,x) + e^{-\delta_2 s}F(t-s,x).$$

For the other case where $\nu + \tau \leq t$, the second particle has already settled down before time t and behaves in the same way at $\nu + \tau$ as the first particle did at time 0, but with starting point $B(\tau)$ and conditional on being then still alive. Therefore, the distribution of $\sum_{i=1}^{N(t)} X_i(t)$ conditional on $\nu = s$, $\tau = u$ and the first particle being alive at time s is then equal to the sum of independent copies of the random variables $\mathbb{1}_{A_1(u)} \sum_{i=1}^{N(t-s-u)} \mathbb{1}_{X_i(t-s-u)+B(u)\leq x}$ and $\sum_{i=1}^{N(t-s)} \mathbb{1}_{X_i(t-s)\leq x}$. This yields for $s + u \leq t$ that

$$
E\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x} \middle| \nu = s, \tau = u \right]
$$

$$
= e^{-\delta_2 s - \delta_1 u} E\left[\sum_{i=1}^{N(t-s-u)} \mathbb{1}_{X_i(t-s-u)+B(u)\leq x} \right] + e^{-\delta_2 s} E\left[\sum_{i=1}^{N(t-s)} \mathbb{1}_{X_i(t-s)\leq x} \right]
$$

$$
= e^{-\delta_2 s - \delta_1 u} E\left[F\big(t - s - u, x - B(u)\big) \right] + e^{-\delta_2 s} F(t - s, x)
$$

$$
= e^{-\delta_2 s - \delta_1 u} \int_{-\infty}^{\infty} F\big(t - s - u, x - z\big) g(u, z)\, \mathrm{d}z + e^{-\delta_2 s} F(t - s, x).
$$

Combining the terms, we obtain

$$
F(t, x) = \int_0^{\infty} E\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x} \middle| \nu = s \right] \mu e^{-\mu s}\, \mathrm{d}s
$$

$$
= e^{-\delta_2 t} e^{-\mu t} \mathbb{1}_{0\leq x} + \int_0^t \big(e^{-\delta_2 s - \delta_1 (t-s)} G(t - s, x)
$$

$$
+ e^{-\delta_2 s} F(t - s, x) \big) e^{-\lambda(t-s)} \mu e^{-\mu s}\, \mathrm{d}s
$$

$$
+ \int_0^t \int_0^{t-s} \left(e^{-\delta_2 s - \delta_1 u} \int_{-\infty}^{\infty} F(t - s - u, x - z) g(u, z)\, \mathrm{d}z \right. \tag{2.12}
$$

$$
\left. + e^{-\delta_2 s} F(t - s, x) \right) \lambda e^{-\lambda u}\, \mathrm{d}u\, \mu e^{-\mu s}\, \mathrm{d}s
$$

$$
= e^{-(\mu+\delta_2) t} \mathbb{1}_{0\leq x} + \int_0^t \big(e^{-(\lambda+\delta_1) s} G(s, x) + F(s, x) \big) \mu e^{-(\mu+\delta_2)(t-s)}\, \mathrm{d}s
$$

$$
+ \int_0^t \int_0^s \int_{-\infty}^{\infty} F(s - u, x - z) g(u, z)\, \mathrm{d}z \lambda e^{-(\lambda+\delta_1) u}\, \mathrm{d}u\, \mu e^{-(\mu+\delta_2)(t-s)}\, \mathrm{d}s.
$$

We deduce by differentiating with respect to t that

$$
F_t(t, x) = -(\mu + \delta_2) F(t, x) + G(t, x) e^{-(\lambda+\delta_1) t} \mu + F(t, x)\mu
$$

$$
+ \int_0^t \int_{-\infty}^{\infty} F(t - u, x - z) g(u, z) \lambda e^{-(\lambda+\delta_1) u} \mu\, \mathrm{d}z\, \mathrm{d}u,
$$

which shows that F satisfies (2.5) for the choices mentioned in Table 1.

The proof for the integro-differential equations of V, H and Q go analogously.

For V, we start by decomposing it as

$$V(t,x) = \mathrm{Var}\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\right] = \underbrace{E\left[\left(\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\right)^2\right]}_{=:\tilde{F}(t,x)} - \underbrace{\left(E\left[\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\right]\right)^2}_{= F(t,x)}.$$

This implies $V_t(t,x) = \tilde{F}_t(t,x) - 2F(t,x)F_t(t,x)$ so that the integro-differential equation of V follows from such equations for F and $\tilde{F}$. Similarly to (2.11), we write

$$\tilde{F}(t,x) = E\left[\left(\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\right)^2\right] = \int_0^\infty E\left[\left(\sum_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\right)^2 \middle| \nu = s\right] \mu e^{-\mu s}\,\mathrm{d}s,$$

$$H(t,x) = P\left[\max_{i=1,\ldots,N(t)} X_i(t) \leq x\right] = P\left[X_i(t) \leq x \text{ for all } i = 1,\ldots,N(t)\right]$$

$$= E\left[\prod_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x}\right] = \int_0^\infty E\left[\prod_{i=1}^{N(t)} \mathbb{1}_{X_i(t)\leq x} \middle| \nu = s\right] \mu e^{-\mu s}\,\mathrm{d}s,$$

$$Q(t) = P[M(t) > 0] = \int_0^\infty P[M(t) > 0 | \nu = s]\mu e^{-\mu s}\,\mathrm{d}s.$$

The subsequent steps for $\tilde{F}$, H and Q go analogously to the analysis done for F by using decompositions similar to (2.12). □

3. Analysis of the integro-differential equations

In this section, we first prove existence and uniqueness results for the integro-differential equations (2.5). We then analyze the mean numbers of moving and stationary particles, where we already see that R_0 has a threshold of 1 that distinguishes between metastatic growth or decay. Finally, we prove the main stability theorem.

3.1. *Local and global existence*

For the general equation (2.5), we make the following assumptions:

Assumptions (A1):

- We assume $r_1 \geq 0$, $r_2 \in \mathbb{R}$,
- $q(t,x) \geq \delta > 0$ is uniformly bounded and Lipschitz continuous in t and x.
- For each $T > 0$ $p(t,x) \geq 0$ is uniformly bounded on $[0,T]$ by P, absolute continuous in x and continuous in t.
- For the cases F, H, V: $k(u,z) \geq 0$ is continuous in x and satisfies for some constant $K > 0$ that

$$\int_0^t \int_{-\infty}^\infty k(u,z)\,\mathrm{d}z\mathrm{d}u < K.$$

- For the case Q: $k(u) \geq 0$ satisfies for some constant $K > 0$ that

$$\int_0^t k(u)\,\mathrm{d}u < K.$$

- The initial condition $f_0(x) = f(0, x) \geq 0$ is bounded in $L^\infty(\mathbb{R})$.

3.1.1. *Mild solutions*

To prove local existence and uniqueness, we consider $T > 0$ and use as phase space

$$\mathcal{X} = L^\infty(0, T; \mathbb{R})$$

with the essential supremum norm $||\cdot||_\infty$. To find a mild formulation of (2.5), we define an integrating factor

$$W(t, s, x) = \exp\left(-\int_s^t q(u, x)\,\mathrm{d}u\right) \tag{3.1}$$

where the x-dependence arises only in the case of $f = H$. Since q is Lipschitz continuous and $q > 0$ we have that $W(t, s, x)$ is a non-negative evolution family with $0 < W(t, s, x) \leq 1$ for all $0 \leq s \leq t \leq T$. We use the variation of constant formula to formally solve (2.5) as

$$f(t, x) = f_0(x)W(t, 0, x) + \int_0^t W(t, s, x)\Big[p(s, x) + (r_1 + r_2 f)\,k * f(s, x)\Big]\mathrm{d}s, \tag{3.2}$$

with $f(0, x) = f_0(x)$, which is our mild formulation.

Based on the various interpretations of f as probability or variance, we are interested in non-negative solutions.

Lemma 3.1. *Assume (A1) and let $f \in \mathcal{X}$ be a mild solution of (2.5).*

(1) Then $f(t, x) \geq 0$ as long as the solution exists.
(2) If $f(t_0, x_0) = 0$ for a point (t_0, x_0), then this implies that

$$f_0(x_0) = 0 \quad and \quad p(t, x_0) = 0 \quad for\ all \quad 0 \leq t \leq t_0. \tag{3.3}$$

(3) If in addition $p(t, x) > 0$ for all $(t, x) \in (0, T) \times \mathbb{R}$ then $f(t, x) > 0$ for all $(t, x) \in (0, T) \times \mathbb{R}$.

Proof.

(1) Consider any time t_0 such that $f(t, x) \geq 0$ for all $0 \leq t \leq t_0$ and all $x \in \mathbb{R}$. Define

$$\Omega^0 := \{x \in \mathbb{R} : f(t_0, x) = 0\}.$$

Then at each of the points (t_0, x), $x \in \Omega^0$, we have the following signs of the terms in (2.5):

$$f(t_0, x) = 0,$$

$$p(t_0, x) \geq 0,$$

$$\int_0^{t_0} \int_{-\infty}^{\infty} k(u, z) f(t_0 - u, x - z) \, dz du \geq 0,$$

$$r_1 + r_2 f(t_0, x) = r_1 \geq 0,$$

hence from (2.5), we find $\frac{\partial}{\partial t} f(t_0, x) \geq 0$ for all $x \in \Omega^0$, and the solution does not decay further.

(2) Taking a point (t_0, x_0) with $f(t_0, x_0) = 0$, we find from the mild formulation (3.2) that

$$0 = f_0(x_0) W(t_0, 0, x_0) + \int_0^{t_0} W(t_0, s, x_0)(p(s, x_0) + r_1 k * f(s, x_0)) \, ds, \quad (3.4)$$

where all terms are non-negative. Hence (3.3) follows.

(3) If $p > 0$, we see that (3.4) is not possible, hence $f > 0$. □

Proposition 3.1. *Assume (A1). Then there exists a time $T > 0$ and a unique mild solution $f \in \mathcal{X}$ which satisfies (3.2).*

Proof. Given a function $\tilde{f} \in \mathcal{X}$, we define an iteration operator $A : \tilde{f} \to f$ where f is the unique solution of

$$f_t = -qf + p + (r_1 + r_2 \tilde{f}) \, k * \tilde{f}. \quad (3.5)$$

Using the integrating factor (3.1) we solve (3.5) as

$$f(t, x) = f_0(x) W(t, 0, x) + \int_0^t W(t, s, x) \left[p(s, x) + (r_1 + r_2 \tilde{f}) \, k * \tilde{f}(s, x) \right] ds. \quad (3.6)$$

Step 1: Given an initial condition $f_0 \in \mathcal{X}$ with norm $\|f_0\|_{\mathcal{X}} = M_0$. We show that for each constant $M > 0$ there exists a time $T_1 > 0$ such that $A : B_M(f_0) \to B_M(f_0)$ for all $T \leq T_1$, where $B_M(f_0) \subset \mathcal{X}$ denotes the closed ball of radius M in $\mathcal{X}$ with center f_0. Consider $\tilde{f} \in B_M$ then $\tilde{f}$ has norm

$$\|\tilde{f}\|_{\mathcal{X}} \leq M_0 + M.$$

From (3.6) we find

$$f(t, x) - f_0(x) = (W(t, 0, x) - 1) f_0(x)$$
$$+ \int_0^t W(t, s, x) \left[p(s, x) + (r_1 + r_2 \tilde{f}) \, k * \tilde{f}(s, x) \right] ds.$$

Since $W(t, 0, x) < 1$ for all $t > 0$, we can ignore the first term and estimate

$$\|f(t, .) - f_0\|_{\infty} \leq tP + r_1 Kt \|\tilde{f}\|_{\mathcal{X}} + |r_2| Kt \|\tilde{f}\|_{\mathcal{X}}^2,$$

which implies that

$$\|f - f_0\|_{\mathcal{X}} \leq T\Big(P + r_1 K \|\tilde{f}\|_{\mathcal{X}} + |r_2| K \|\tilde{f}\|_{\mathcal{X}}^2\Big)$$
$$\leq T\left(P + r_1 K(M_0 + M) + |r_2| K(M_0 + M)^2\right).$$

To ensure $\|f - f_0\|_{\mathcal{X}} \leq M$, we need

$$T\left(P + r_1 K(M_0 + M) + |r_2| K(M_0 + M)^2\right) \leq M,$$

which is true for all

$$T \leq T_1 := \frac{M}{P + r_1 K(M_0 + M) + |r_2| K(M_0 + M)^2}.$$

Hence, for $T \leq T_1$, we have $A : B_M(f_0) \to B_M(f_0)$.

Step 2: We show that A is a contraction for $T \leq T_2$ for some $T_2 > 0$.
Given $\tilde{f}, \tilde{g} \in B_M(f_0)$ with $\tilde{f}_0(x) = \tilde{g}_0(x) = f_0(x)$ Then from (3.6), we find

$$\|f(t,.) - g(t,.)\|_{\infty} \leq \left\|\int_0^t \left((r_1 + r_2 \tilde{f})\, k * \tilde{f} - (r_1 + r_2 \tilde{g})\, k * \tilde{g}\right) \mathrm{d}s\right\|_{\infty}$$
$$\leq r_1 \left\|\int_0^t k * (\tilde{f} - \tilde{g})\, \mathrm{d}s\right\|_{\infty} + |r_2| \left\|\int_0^t \left((\tilde{f} - \tilde{g}) k * \tilde{f} + \tilde{g}\,(\tilde{f} - \tilde{g})\right) \mathrm{d}s\right\|_{\infty}$$
$$\leq r_1 K t \|\tilde{f} - \tilde{g}\|_{\mathcal{X}} + |r_2| K t \|\tilde{f}\|_{\mathcal{X}} \|\tilde{f} - \tilde{g}\|_{\mathcal{X}} + |r_2| K t \|\tilde{f} - \tilde{g}\|_{\mathcal{X}}$$
$$= K t \big(r_1 + |r_2| \|\tilde{f}\|_{\mathcal{X}} + |r_2| \|\tilde{g}\|_{\mathcal{X}}\big) \|\tilde{f} - \tilde{g}\|_{\mathcal{X}}.$$

Hence

$$\|f - g\|_{\mathcal{X}} \leq K T \big(r_1 + |r_2| \|\tilde{f}\|_{\mathcal{X}} + |r_2| \|\tilde{g}\|_{\mathcal{X}}\big) \|\tilde{f} - \tilde{g}\|_{\mathcal{X}}$$
$$\leq K T \left(r_1 + 2|r_2|(M_0 + M)\right) \|\tilde{f} - \tilde{g}\|_{\mathcal{X}}.$$

Then A is a contraction for all

$$T < T_2 := \frac{1}{K(r_1 + 2|r_2|(M_0 + M))}.$$

Step 3: For $T < \min\{T_1, T_2\}$ the Banach fixed point theorem applies, and A has a unique fixed point in $\mathcal{X}$, which satisfies the mild formulation (3.2). □

The local existence times T from the above proof depend on the norm of the initial condition, hence we cannot simply repeat the argument to obtain global existence. Nevertheless, we show next that solutions are global.

3.1.2. *Global existence*

Theorem 3.1. *The unique mild solutions from Proposition 3.1 exist for all times. The probabilities H and Q are globally bounded by 1 (as solutions of their corresponding integro-differential equation).*

Proof.

- For the cases F and V we have $r_2 = 0$, hence (2.5) becomes a linear equation in f. We estimate

$$\|f(t,.)\|_\infty \le \|f_0\|_\infty + tP + r_1 K \int_0^t \|f(s,.)\|_\infty \, \mathrm{d}s$$

and Gronwall's lemma implies that $\|f(t,.)\|_\infty$ grows at most exponentially, hence proving global existence.

- In the case Q the parameter $r_2 = -1$ is negative. Hence in the global estimate, we can remove the quadratic term. Again Gronwall's lemma applies and global existence follows. To show that the survival probability Q is bounded by 1, we assume that $Q(t_0) = 1$ and $Q(t) < 1$ for all $0 \le t < t_0$, or $t_0 = 0$ and $Q(0) = 1$. Then at t_0 we find from (2.4) that

$$Q_t(t_0) = - \left(\delta_2 + \mu e^{-(\lambda + \delta_1)t_0} \right) Q(t_0) + \mu e^{-(\lambda + \delta_1)t_0}$$

$$+ \lambda \mu (1 - Q(t_0)) \int_0^t Q(t_0 - s) e^{-(\lambda + \delta_1)s} \mathrm{d}s$$

$$= -\delta_2.$$

Hence Q decays near 1. In our specific application we have $Q(0) = 1$, hence Q decays initially and as survival probability, $Q(t)$ is non-increasing.

- A similar argument is applied for H. We assume that $H(t, x) \le 1$ for all $t \le t_0$ and that there exists a point x_0 with $H(t_0, x_0) = 1$. Then at (t_0, x_0) we observe

$$H_t(t_0, x_0) = \left[\mu e^{-\lambda t_0}(1 - e^{-\delta_1 t_0}(1 - G(t_0, x_0))) - (\mu + \delta_1) \right] H(t_0, x_0)$$

$$+ \lambda \mu H(t_0, x_0) \int_0^{t_0} \int_{-\infty}^{\infty} H(t_0 - u, x_0 - z) g(u, z) e^{-\lambda u} \, \mathrm{d}z \mathrm{d}u$$

$$\le \mu e^{-\lambda t_0} - (\mu + \delta_1) + \mu(1 - e^{-\lambda t_0}) = -\delta_1$$

and $H(t_0, x_0) = 1$ decays. $\qquad\square$

3.2. *Metastatic reproduction number R_0*

To introduce the metastatic reproduction number, we reduce the above model by looking at the expected numbers of moving and stationary particles. We denote by $a(t)$ and $b(t)$ the expected numbers of moving and stationary particles, respectively. Moving particles die at rate δ_1 and become stationary at rate λ while new moving particles are born at rate μ from the stationary particles. This reasoning leads to

$$a'(t) = -(\delta_1 + \lambda)a(t) + \mu b(t). \tag{3.7}$$

Formally, we can prove (3.7) analogously to the first part of the proof of Theorem 3.2. Similarly, stationary particles die at rate δ_2 and moving particles become stationary at rate λ, leading to

$$b'(t) = -\delta_2 b(t) + \lambda a(t). \tag{3.8}$$

The differential equations for a and b form a linear system with coordinate matrix

$$A_1 = \begin{pmatrix} -(\delta_1 + \lambda) & \mu \\ \lambda & -\delta_2 \end{pmatrix}.$$

The matrix A_1 has trace and determinant as

$$\mathrm{tr}A_1 = -(\delta_1 + \delta_2 + \lambda), \qquad \det A_1 = \delta_2(\delta_1 + \lambda) - \lambda\mu.$$

The trace is negative, hence the origin is asymptotically stable for $\det A_1 > 0$ and unstable for $\det A_1 < 0$. Using

$$R_0 = \frac{\lambda}{\delta_1 + \lambda}\frac{\mu}{\delta_2},$$

we find the following result:

Lemma 3.2. *Consider (3.7), (3.8).*

- *If $R_0 < 1$ then $(0,0)$ is globally asymptotically stable.*
- *If $R_0 > 1$ then $(0,0)$ is unstable (it is a saddle).*
- *If $R_0 = 1$ then $(0,0)$ is non-hyperbolic and we have a continuum of steady states in direction $(\delta_2, \lambda)^T$.*

Proof. For the first two statements note that we can write

$$\det A_1 = \delta_2(\delta_1 + \lambda)(1 - R_0).$$

For the third statement we find that for $R_0 = 1$ the vector $(\delta_2, \lambda)^T$ is an eigenvector of A_1 with eigenvalue 0. □

It should be noted that $a(t) + b(t) = E[M(t)]$ with the formula for the expected total number $E[M(t)]$ of particles.

Using specific initial conditions $(a(0), b(0)) = (0, 1)$ we can explicitly solve equations (3.7), (3.8) and we find the asymptotic ratio of moving versus stationary particles

$$\lim_{t\to\infty} \frac{a(t)}{b(t)} = \frac{\delta_2 - \delta_1 - \lambda + \sqrt{(\delta_2 - \delta_1 - \lambda)^2 + 4\lambda\mu}}{2\lambda}.$$

3.3. Survival probability $Q(t)$

The survival probability Q is a special case of (2.5), where there is no spatial variable. In this case, (2.5) becomes

$$Q'(t) = -\underbrace{\delta_2 Q(t)}_{A} + \underbrace{(1 - Q(t))\mu e^{-(\delta_1 + \lambda)t}\left(1 + \int_0^t Q(s)\lambda e^{(\delta_1 + \lambda)s}\,\mathrm{d}s\right)}_{B}. \tag{3.9}$$

where

A: death rate of original particle times the survival probability at time t. Note that this is the only term if $\mu = 0$ (no births), in which case, the survival probability equals $Q(t) = \exp(-\delta_2 t)$.

B: correction term because the birth of particles leads to a higher survival probability than in the case $\mu = 0$. If the original particle has offsprings, all of the original particle, the offsprings of the original particle and further offsprings must die to extinct all particles, which is reflected in the term B.

Based on (3.9), we can derive a second-order ODE for Q, namely,

$$
\begin{aligned}
Q''(t) &= \big((\delta_1 + \lambda)\big(Q(t) - 1\big) - Q'(t)\big)\mu e^{-(\delta_1 + \lambda)t}\left(1 + \int_0^t Q(s)\lambda e^{(\delta_1 + \lambda)s}\,\mathrm{d}s\right) \\
&\quad - \delta_2 Q'(t) + \mu\lambda Q(t)\big(1 - Q(t)\big) \\
&= \left(\frac{Q'(t)}{Q(t) - 1} - \delta_1 - \lambda\right)\big(Q'(t) + \delta_2 Q(t)\big) - \delta_2 Q'(t) + \mu\lambda Q(t)\big(1 - Q(t)\big)
\end{aligned}
$$

This equation can be transformed into a system of first-order ODEs as

$$
\begin{aligned}
Q' &= P, \\
P' &= \left(\frac{P}{Q - 1} - \delta_1 - \lambda\right)\big(P + \delta_2 Q\big) - \delta_2 P + \mu\lambda Q\big(1 - Q\big).
\end{aligned}
\tag{3.10}
$$

We show the following result:

Lemma 3.3. *Consider system (3.10).*

- *If $R_0 < 1$ then $(0,0)$ is locally asymptotically stable.*
- *If $R_0 > 1$ then $(0,0)$ is locally unstable (it is a saddle).*

Proof. The linearization of (3.10) at $(0,0)$ is

$$
\begin{pmatrix}
0 & 1 \\
-\delta_2(\delta_1 + \lambda) + \mu\lambda & -\delta_1 - \delta_2 - \lambda
\end{pmatrix}.
$$

This matrix has the same trace and determinant as the matrix A_1 that we encountered in Lemma 3.2. Hence for $R_0 < 1$, the origin is locally asymptotically stable, and for $R_0 > 1$, it is a saddle. □

3.4. *Asymptotic behavior as $t \to \infty$*

In this section, we study how the average number of particles and the survival probability behave in the limit $t \to \infty$. We first provide an explicit formula for the average number of particles at an arbitrary time t.

Theorem 3.2. *For $\mu \neq 0$, and $\lambda \neq 0$ or $\delta_1 \neq \delta_2$, the average number of particles alive at time t is given by*

$$
E[M(t)] = \frac{\mu - \delta_2 - \alpha_-}{\alpha_+ - \alpha_-}e^{\alpha_+ t} + \frac{-\mu + \delta_2 + \alpha_+}{\alpha_+ - \alpha_-}e^{\alpha_- t},
$$

where

$$\alpha_{\pm} = \frac{-\delta_2 - \delta_1 - \lambda \pm \sqrt{(\delta_2 + \delta_1 + \lambda)^2 + 4(R_0 - 1)\delta_2(\lambda + \delta_1)}}{2}. \tag{3.11}$$

For $\mu = 0$ or both $\lambda = 0$ and $\delta_1 = \delta_2$, the average number of particles at time t equals $E[M(t)] = (1 + \mu t)e^{-\delta_1 t}$.

Proof. Similarly to the proof of Theorem 2.1, we find

$$E[M(t)] = \int_0^{\infty} E[M(t)|\nu = s]\mu e^{-\mu s} \, ds$$

$$= e^{-(\delta_2 + \mu)t} + \int_0^t \left(e^{-\delta_2 s - \delta_1(t-s)} + e^{-\delta_2 s}E[M(t-s)]\right)e^{-\lambda(t-s)}\mu e^{-\mu s} \, ds$$

$$+ \int_0^t \int_0^{t-s} \left(e^{-\delta_2 s - \delta_1 u}E[M(t-s-u)] + e^{-\delta_2 s}E[M(t-s)]\right)\lambda e^{-\lambda u} \, du \, \mu e^{-\mu s} \, ds$$

$$= e^{-(\delta_2 + \mu)t} + \int_0^t \left(e^{-(\delta_1 + \lambda)s} + E[M(s)]\right)\mu e^{-(\delta_2 + \mu)(t-s)} \, ds$$

$$+ \int_0^t \int_0^s e^{-(\delta_2 + \mu)(t-s) - (\delta_1 + \lambda)u}E[M(s-u)]\lambda\mu \, du \, ds.$$

We deduce that

$$\frac{d}{dt}E[M(t)] = -(\delta_2 + \mu)E[M(t)] + \mu e^{-(\delta_1 + \lambda)t} + \mu E[M(t)]$$

$$+ \int_0^t e^{-(\delta_1 + \lambda)u}E[M(t-u)]\lambda\mu \, du$$

$$= -\delta_2 E[M(t)] + \mu e^{-(\delta_1 + \lambda)t} + \int_0^t e^{-(\delta_1 + \lambda)(t-u)}E[M(u)]\lambda\mu \, du,$$

$$\frac{d^2}{dt^2}E[M(t)] = -\delta_2\frac{d}{dt}E[M(t)] - (\delta_1 + \lambda)\left(\frac{d}{dt}E[M(t)] + \delta_2 E[M(t)]\right) + E[M(t)]\lambda\mu$$

$$= -(\delta_2 + \delta_1 + \lambda)\frac{d}{dt}E[M(t)] + \left(\lambda\mu - (\delta_1 + \lambda)\delta_2\right)E[M(t)],$$

which shows that $E[M(t)]$ satisfies a linear ordinary differential equation, whose solution is of the form $E[M(t)] = c_+ e^{\alpha_+ t} + c_- e^{\alpha_- t}$ with

$$\alpha_{\pm} = \frac{-\delta_2 - \delta_1 - \lambda \pm \sqrt{(\delta_2 + \delta_1 + \lambda)^2 + 4\lambda\mu - 4(\delta_1 + \lambda)\delta_2}}{2}$$

$$= \frac{-\delta_2 - \delta_1 - \lambda \pm \sqrt{(\delta_2 + \delta_1 + \lambda)^2 + 4(R_0 - 1)\delta_2(\lambda + \delta_1)}}{2}$$

if $\delta_2 \neq \delta_1 + \lambda$ or $\lambda\mu \neq 0$. To find the constants c_- and c_+, we use the boundary conditions

$$E[M(0)] = M(0) = 1 \quad \text{and} \quad \frac{d}{dt}E[M(t)]\Big|_{t=0} = \mu - \delta_2,$$

where we applied that at time 0 there is one particle which will be either eliminated (at rate δ_2) or doubled (at rate μ). This yields

$$c_- + c_+ = 1 \quad \text{and} \quad \alpha_- c_- + \alpha_+ c_+ = \mu - \delta_2$$

which gives $c_- = \frac{\mu - \delta_2 - \alpha_+}{\alpha_- - \alpha_+}$ and $c_+ = \frac{-\mu + \delta_2 + \alpha_-}{\alpha_- - \alpha_+}$. In the case $\lambda = 0$ and $\delta_1 = \delta_2$, we have

$$\frac{\mathrm{d}}{\mathrm{d}t} E[M(t)] = -\delta_1 E[M(t)] + \mu \mathrm{e}^{-\delta_1 t},$$

which satisfies

$$\frac{\mathrm{d}}{\mathrm{d}t}\left(\mathrm{e}^{\delta_1 t} E[M(t)]\right) = \mu,$$

hence $E[M(t)] = (1 + \mu t)\mathrm{e}^{-\delta_1 t}$. In the case $\mu = 0$, we have $\frac{\mathrm{d}}{\mathrm{d}t} E[M(t)] = -\delta_2 E[M(t)]$ so that $E[M(t)] = \mathrm{e}^{-\delta_1 t}$. $\qquad\square$

For $\mu \neq 0$, and $\lambda \neq 0$ or $\delta_1 \neq \delta_2$, we obtain from Theorem 3.2 that

$$\frac{E[M(t)]}{\mathrm{e}^{\alpha_+ t}} = \frac{\mu - \delta_2 - \alpha_-}{\alpha_+ - \alpha_-} + \frac{-\mu + \delta_2 + \alpha_+}{\alpha_+ - \alpha_-} \mathrm{e}^{-\sqrt{(\delta_2 - \delta_1 - \lambda)^2 + 4\lambda\mu}\, t}$$

$$= \frac{2\mu - \delta_2 + \delta_1 + \lambda + \sqrt{(\delta_2 - \delta_1 - \lambda)^2 + 4\lambda\mu}}{2\sqrt{(\delta_2 - \delta_1 - \lambda)^2 + 4\lambda\mu}} + o(1)$$

for $t \to \infty$ so that $E[M(t)]$ has exponential growth with rate α_+.

Note that $\alpha_+ > 0$ in (3.11) if and only if $R_0 > 1$ and $\alpha^+ < 0$ whenever $R_0 < 1$. Also note that α^+ is real for all $R_0 \in [0,1]$. From this reasoning, we deduce the following result.

Corollary 3.1. $E[M(t)]$ *has the following asymptotic behavior: it ...*

$$\begin{cases} \textit{equals } (1 + \mu t)\mathrm{e}^{-\delta_1 t} & \textit{if } \mu = 0 \textit{ or both } \lambda = 0 \textit{ and } \delta_1 = \delta_2, \\ \textit{grows exponentially at rate } \alpha_+ & \textit{if } R_0 > 1 \textit{ and not case 1,} \\ \textit{converges to } \frac{1 + \lambda/\delta_2}{\delta_2/\mu + \lambda/\delta_2} & \textit{if } R_0 = 1 \textit{ and not case 1,} \\ \textit{shrinks exponentially to 0 at rate } \alpha_+ & \textit{if } R_0 < 1 \textit{ and not case 1.} \end{cases}$$

In Lemma 3.3 we have already shown the corresponding threshold phenomenon for the survival probability Q. In more detail, we can also compute the asymptotic value as $t \to \infty$.

Lemma 3.4. *The asymptotic survival probability is*

$$\lim_{t \to \infty} Q(t) = \begin{cases} \max\{1 - \frac{1}{R_0}, 0\} & \textit{if } \mu > 0 \textit{ and } \lambda > 0, \\ 1 & \textit{if } \{\mu = 0 \textit{ or } \lambda = 0\} \textit{ and } \delta_2 = 0, \\ \frac{\mu}{\mu + \delta_2} & \textit{if } \lambda = 0 \textit{ and } \delta_1 = 0 \textit{ and } \delta_2 > 0, \\ 0 & \textit{otherwise.} \end{cases}$$

Proof. We first observe that if $\delta_2 = 0$, then $Q(t) = 1$ for all t and hence $\lim_{t \to \infty} Q(t) = 1$ because the first particle then lives forever. Now consider the case where $\delta_2 > 0$ and $\delta_1 = \lambda = 0$ so that (3.9) becomes

$$Q'(t) = -\delta_2 + \left(1 - Q(t)\right)(\mu + \delta_2), \quad Q(0) = 1,$$

whose solution is $Q(t) = \frac{\mu + \delta_2 e^{-(\mu+\delta_2)t}}{\mu+\delta_2}$, which converges to $\lim_{t\to\infty} Q(t) = \frac{\mu}{\mu+\delta_2}$. Let us now consider $\delta_1 + \lambda > 0$ and $\delta_2 > 0$. Because $M(t) = 0$ implies $M(s) = 0$ for all $s \geq t$ (after particles died out, there will always be zero particles), we obtain

$$Q(t) = P[M(t) > 0] = 1 - P[M(t) = 0] \geq 1 - P[M(s) = 0] = Q(s)$$

for all $s \geq t$, which shows that Q is a non-increasing function. Since Q is also bounded from below by 0, it converges to some limit $Q(\infty) := \lim_{t\to\infty} Q(t)$. Using $\lim_{t\to\infty} Q'(t) = 0$ and taking limits in (3.9) gives

$$0 = -Q(\infty)\delta_2 + \big(1 - Q(\infty)\big) \lim_{t\to\infty} \int_0^t Q(t-u)\mu\lambda e^{-(\delta_1+\lambda)u}\,\mathrm{d}u.$$

From L'Hôpital's rule, it follows that

$$\lim_{t\to\infty} \int_0^t Q(t-u)\mu\lambda e^{-(\delta_1+\lambda)u}\,\mathrm{d}u = \lim_{t\to\infty} \frac{\int_0^t Q(u)\mu\lambda e^{(\delta_1+\lambda)u}\,\mathrm{d}u}{e^{(\delta_1+\lambda)t}}$$

$$= \lim_{t\to\infty} \frac{Q(t)\mu\lambda e^{(\delta_1+\lambda)t}}{(\delta_1+\lambda)e^{(\delta_1+\lambda)t}}$$

$$= Q(\infty)\frac{\mu\lambda}{\delta_1+\lambda}.$$

Therefore, $Q(\infty)$ satisfies the quadratic equation

$$0 = \Big(\frac{\mu\lambda}{\delta_1+\lambda} - \delta_2\Big)Q(\infty) - \frac{\mu\lambda}{\delta_1+\lambda}Q(\infty)^2.$$

If $\lambda = 0$ or $\mu = 0$, its solution is $Q(\infty) = 0$ while otherwise, the solutions are zero or

$$Q(\infty) = \frac{\mu\lambda - \delta_2(\delta_1+\lambda)}{\mu\lambda} = 1 - \frac{1}{R_0}.$$

Note that $Q(t)$ cannot cross the level $1 - \frac{1}{R_0}$ so that it will converge at $1 - \frac{1}{R_0}$ if this quantity is greater than 0. □

4. Example: metastatic cancer

In this section, we apply the modeling framework developed above in the context of metastatic cancer. We first discuss the appropriate interpretations of the model parameters, and provide biologically reasonable estimates obtained from the literature. We then make a brief note on the numerical implementation, followed by a presentation of the results themselves.

4.1. *Murine data*

Because of the scarcity of quantitative data for metastatic cancer in humans, the majority of the values discussed below have come from experimental models of metastasis in mice. Many of the studies mentioned below follow a similar procedure (those in Refs. 2, 8, 24, 30 for example), and we provide a brief outline of

their methods. Tumor-free mice are injected with radio-labeled cancer cells (B16 melanoma[8], M19 Fibrosarcoma[30], B16F1(0) melanoma[2,24]), and observed and/or sacrificed at various time points ranging from 1 minute to 14 day post-injection. Organs of interest (multiple organs[8], lungs[2,30], and liver[24]) are removed and analyzed for the number and location of cancer cells, cancer cell clusters, and metastases. In addition to the radio-labeled cancer cells, the Chambers group[2,24] injected inert microspheres that become lodged within the microvasculature of the target organ in order to accurately determine the change in cell numbers over time. Details of specific experimental models can be found in the cited references.

4.2. *Modeling metastatic cancer*

In order to apply our model to the metastatic dissemination of cancer, we must first carefully define what is meant by 'stationary' and 'mobile' particles in this context. 'Stationary' particles will play the role of established tumors capable of shedding mobile particles without exhausting themselves. 'Mobile particles', therefore, will represent small clusters of individual cancer cells that are actively circulating through the vasculature. This interpretation necessitates different scales for the two classes of particles, with established tumors consisting of at least 10^8 cells — corresponding to a tumor volume of approximately 1cm^3 (Ref. 5) — and CTCs consisting of anything between a single cell to several dozen.[9] Such a distinction requires careful attention when parameterizing the model. Below we discuss our approach to address this concern.

First we consider the shedding rate, μ. Assuming that an established tumor consists of 10^8 cells[5] and that the number of CTCs shed per day range between $0.0001\% - 0.01\%$ of the cells available within the established tumor,[32] we may choose $\mu \in [100, 10000]$ cells per day. For the simulations presented herein, we chose $\mu = 346$ cells/day. This choice was made in order to have an average of 10 established tumors by the end of 14 days[30] (determined using the asymptotic expected ratio of moving to stationary particles from the comment after Lemma 3.2).

Second, we estimate the stationary particle death rate, δ_2. With the interpretation of a stationary particle as an established tumor, δ_2 corresponds to the rate of spontaneous tumor remission. We use Jessy's estimate of $p = 10^{-5}$ (Ref. 20) for the probability of spontaneous remission and assume that $p = \delta_2/\lambda$ to obtain the value of δ_2 reported in Table 2.

Third, we consider the rate of mobile particle settlement, λ. Experimental murine models of metastasis suggest that nearly 80% of the CTCs shed from the primary tumor into the vasculature will survive through the circulation and successfully extravasate at a secondary site.[2,24] For the purposes of our model however, successful extravasation alone does not represent 'settlement'. Indeed, because of the property that 'settled' particles are immediately capable of shedding mobile particles at the same rate as the original stationary particle, 'settlement' in our model includes not only successful extravasation at a secondary site, but survival and growth to

Parameter	Meaning	Value	Units	Reference(s)
μ	shedding rate	346	cells/day	[32,5]
λ	establishment rate	0.0035	tumors/day	[24]
δ_1	mobile death rate	7	cells/day	[8,23,3]
δ_2	stationary death rate	3.46×10^{-8}	tumors/day	[20]

Table 2. Model Parameters and the values used in presented simulations. The units can also be interpreted as simply 'particles/day', where it is understood that the meaning of 'particles' can be either an established tumor or a CTC. See text for further details.

a palpable secondary tumor as well. For this reason, we use the more suggestive terminology 'establishment' instead of 'settlement'. Moreover, we assume that the establishment rate, λ, is related to the shedding rate, μ, via $\lambda = \mu q$ where q denotes the probability (per cell) of establishment. The probability q has been estimated by several investigators to range between 0.0001 and 0.00001.[3,8,24,30] In the results presented below, we have used the lower estimate of $q = 0.00001$.[24]

Finally, we require an estimate for the mobile particle death rate, δ_1. While approximately 80% of cancer cells released into circulation will survive in the vasculature and successfully extravasate at a secondary location[2,24], the fraction of these extravasated cells that will grow and form a metastasis is very small[2]. Consequently, our estimate for the mobile particle death rate must also include the death rate of successfully extravasated cells that do not become metastases, and will be much larger than if we included only deaths during transit. Additionally, the time that circulating cancer cells spend traveling through the circulation has been estimated to be between 1 and 3 hours.[24,30] Therefore, assuming that 'mobile particle' means 'CTC', we expect these particles to be short-lived. Combining these observations, and based on previous results[23,8,3], we chose to have 99.9% of all the cells that are shed over the course of a day perish that day. Under this assumption, the mobile death rate becomes $\delta_1 = 7$ cells/day.

4.3. Numerical results

The stochastic model described at the beginning of Section 2 was simulated using a variant of the Gillespie algorithm[7,10]. The key feature of this algorithm is its non-uniform time stepping which is ideally suited for our model, as this model can see (for example) the total rate of mobile particle creation double upon successful establishment of a secondary tumor because of the assumption of independence amongst the stationary particles.

The implemented algorithm assumes a finite spatial domain, whereas the theoretical work presented in the previous sections does not. In order to simulate an in-

Figure 2. Total number of particles (stationary and mobile) as a function of time. Comparison of the theoretical expected value from Theorem 3.2 (black) and the average of 2250 realizations of the stochastic model (blue). Parameters as in Table 2.

finite domain, we have chosen the finite domain to be sufficiently large so that there are no collisions between our mobile particles and the domain boundaries within the time of interest. For the simulations presented here (using diffusion coefficient of $D = 19\text{cm}^2/\text{day}$; see Ref. 30), it was sufficient to consider the spatial domain $[-75\text{cm}, 75\text{cm}]$ divided into $K = 1500$ bins each of width 0.1cm. In Figures 2–4, we present the average results of 2250 distinct realizations of the stochastic model simulated over a period of 14 days. None of these realizations had particle-boundary collisions or complete disease extinction.

Theorem 3.2 provides an exact description of the expected number of particles as a function of time. We have therefore used this description to verify the accuracy of our numerical implementation of the model. Figure 2 shows a comparison between the exact dynamics from Theorem 3.2 (black curve) and the average dynamics over 2250 individual realizations of the stochastic model (blue curve). The average percent error over the 14 days simulated is 0.62%, with a maximum value of 1.86%.

While the simulations all begin with a single stationary particle, the expected number of particles increases to approximately 50 within the first day. This rapid increase is due to the relatively high shedding rate, resulting in the rapid creation of mobile particles. The slow-down upon reaching 50 particles reflects the expected asymptotic ratio of moving to stationary particles (comment after Lemma 3.2) which is approximately 48 with the parameters from Table 2. With the metastatic reproduction number $R_0 \approx 4.94 \times 10^6 > 1$, we expect the total particle number to grow exponentially at rate $\alpha_+ \approx 0.167$ (Corollary 3.1). After an initial period of transience, we do see exponential growth, both in the exact and simulated results.

Figure 3. Average results of 2250 individual realizations of the stochastic model. Left column: spatio-temporal dynamics of the stationary (top) and mobile (bottom) particles. The horizontal axis is time (in days) while the vertical axis denotes space (in cm from location of primary tumor). Number of particles indicated by the coloring (color version online). Note the different scales in the top plot. Right column: average spatial distribution of the stationary (top) and mobile (bottom) particles at the end of the 14 day simulations. Note the difference in scales from top to bottom. Parameters used as in Table 2.

Figure 3 illustrates the average spatio-temporal dynamics of the stochastic model. The left column presents the full spatio-temporal dynamics of the stationary (top) and mobile (bottom) particles, while the right column shows the spatial distribution of the stationary (top) and mobile (bottom) particles at time $t = 14$ days. In no individual simulation did we see the original established tumor perish. This result is not unexpected given the probability of a tumor perishing over the 14 days considered in our simulations is $1 - \exp(-14\delta_2) = 4.48 \times 10^{-7}$. Consequently, we always have at least one stationary particle located at position $x = 0$. This explains both the horizontal line in the top left plot, as well as the tall bar at the origin in the top right plot. The histograms on the right side of Figure 3 show relatively symmetric distributions of both stationary and mobile particles centered around the origin. While the individual location of each particle is given by a normal distribution as a result of the Brownian dynamics, the distribution of the aggregate particles (both stationary and mobile) is not normal. The reason is that shedding, settlement and death cause additional randomness. Even when $\lambda = 0$ (no settlement) and $\delta_1 = \delta_2 = 0$ (no deaths) so that only one stationary particle sheds moving particles, the distribution of the aggregate moving particles will not be normal. This can be seen from (2.1), which becomes $F_t = \mu G$, hence the density of the aggregate

Figure 4. Manipulation of the plots from the left column of Figure 3 showing the areas that may be significantly affected by the primary tumor and its metastatic spread. More specifically, areas that have, on average, at least 0.025 particles over all 2250 simulations are colored white. Areas in which this is not the case are colored black. The top shows results for stationary particles, and the bottom for mobile particles.

moving particles in this case is an integral of normal densities and not a normal density itself.

In order to more clearly see the interface between empty space and invading cancer cells, we have taken the data in the left column of Figure 3, and simplified them to be either 1 if there was, on average, at least 0.025 particles in that location across all 2250 simulations, or 0 otherwise. The results of this simplification are presented in Figure 4. We can see that for stationary particles, it takes close to

four days before we see any significant establishment events. This result mirrors the observations made by Cameron et al.[2] that no metastases established in the first four days post-injection.

Following a rapid initial jump, the mobile particle boundary appears to invade at a more or less constant speed. These mobile 'boundary' dynamics (Figure 4) are in stark contrast with the 'interior' dynamics (bottom left in Figure 3, in particular, the blue-teal interface) where the level sets form triangular regions with edges whose slopes are increasing as we advance through time.

5. Conclusion and discussion

We introduced and analyzed a *branching stochastic process with settlement* and we applied it to metastatic cancer growth. The fact that the expected number F of particles, the variance V, the distribution H of the furthest particle and the extinction probability Q satisfy the same type of integro-differential equation with distributed delay (2.5) reveals the recursive structure of this process. Methods from differential equations theory become available to analyze the qualitative behavior of this stochastic process. A recurring quantity was identified to play the role of a basic reproductive number, similar to epidemic models[16], which we call the *metastatic reproductive number*

$$R_0 = \frac{\lambda}{\delta_1 + \lambda} \frac{\mu}{\delta_2}.$$

For the mouse data that we analyzed as an example, we found $R_0 \sim 10^6$, which, of course, is huge. This is expected, as cell lines for metastasis studies are chosen specifically to generate metastases efficiently and reliably.

The value of R_0 for a typical human cancer will be quite different, and we leave a detailed estimate of R_0 for human cancers for further studies. Still, we can already see the impact of various possible treatment strategies. To shrink R_0, we like to reduce the shedding rate μ and the settlement rate λ while increasing the death rates δ_1 and δ_2 for moving and stationary particles, respectively. For example, the death rate δ_1 for CTCs could be increased through platelet inhibitors. Platelets are known to shield cancer cells from the immune surveillance, and less platelets can make cancer cells more exposed and more vulnerable.[4,28,26] The settlement rate λ might be reduced through decreasing the availability of metastatic niches [26]. This can be achieved through very simple means such as reduced pH-levels of tissue[29] to very advanced means such as novel immunotherapies designed to disrupt the preparation of the pre-metastatic niche[21]. However, removing 90% of cancer sites would not change the final outcome since the reproductive number is unchanged. A partial removal would significantly delay cancer spread, but metastasis would recur over time. Overall, the index R_0 has the potential to become a useful quantity in treatment planning.

We see various extensions and limitations of the model as we discuss now.

(1) While we started with a stationary individual, we could have started with a particle that is moving randomly, but this would make the analysis more difficult. Furthermore, if we know the function F of our model, we can find the corresponding function $\tilde{F}$ in a model with randomly moving first particle by

$$
\tilde{F}(t,x) = E\left[\sum_{i=1}^{N(t-\min\{\tau,t\})} \mathbb{1}_{X_i(t-\min\{\tau,t\})+B(\min\{\tau,t\})\leq x}\right]
$$

$$
= \int_0^t E\left[\sum_{i=1}^{N(t-s)} \mathbb{1}_{X_i(t-s)+B(s)\leq x}\right]\lambda\mathrm{e}^{-\lambda s}\mathrm{e}^{-\delta_1 s}\,\mathrm{d}s + \int_t^\infty P[B(t)\leq x]\lambda\mathrm{e}^{-\lambda s}\mathrm{e}^{-\delta_1 t}\,\mathrm{d}s
$$

$$
= \int_0^t \int_{-\infty}^\infty F(t-s,x-y)g(s,y)\lambda\mathrm{e}^{-(\lambda+\delta_1)s}\,\mathrm{d}y\,\mathrm{d}s + G(t,x)\mathrm{e}^{-(\lambda+\delta_1)t},
$$

where $B(\min\{\tau,.\})$ is a random process describing the movement of the first particle up to time τ, and the factor $\mathrm{e}^{-\delta_1 s}$ is the survival probability of the first particle at time s.

(2) A classical related model is branching Brownian motion[1], where all particles move according to Brownian motion and there are no deaths, no settlements and births occur independently at some exponentially distributed time $\nu \sim \mathrm{Exp}(\mu)$. Similarly to the proof of Theorem 2.1, it can be shown that the expected location function $\bar{F}$ satisfies

$$
\bar{F}_t(t,x) = G_t(t,x) + \mu G(t,x).
$$

Note that this is different from our model even when assuming no deaths ($\delta_1 = \delta_2 = 0$) and no settlement ($\lambda = 0$). Indeed, for $\delta_1 = \delta_2 = \lambda = 0$, F in our model satisfies $F_t(t,x) = \mu G(t,x)$. The reason for this difference to the case of branching Brownian motion is that, in our model with $\delta_1 = \delta_2 = \lambda = 0$, there is always a particle located at zero, and only this particle emits further particles.

(3) In the linear case $r_2 = 0$ and for constant decay rate $q(t) = \mathrm{const.}$ we can use Fourier and Laplace transforms to find an explicit solution of the general non-local model (2.5). Let $\mathcal{F}, \mathcal{F}^{-1}$ denote the Fourier transform and its inverse and let $\mathcal{L}, \mathcal{L}^{-1}$ denote the Laplace transform and its inverse, respectively. We use the hat $\hat{f}$ to denote the Fourier transform of a function and the tilde $\tilde{g}$ for the Laplace transform, i.e.

$$
\hat{f}(\omega) := \mathcal{F}(f(x))(\omega), \qquad \tilde{g}(s) := \mathcal{L}(g(t))(s).
$$

Transforming the general model (2.5) in the case of $r_2 = 0$ and $q(t) = \mathrm{const.}$ leads to

$$
s\tilde{\hat{f}} - \hat{f}(0) = -q\tilde{\hat{f}} + \tilde{\hat{p}} + r_1\sqrt{2\pi}\tilde{\hat{k}}\tilde{\hat{f}}
$$

for an unknown function $\tilde{\hat{f}}(s,\omega)$ in Laplace-Fourier space. We can solve this algebraic equation and find

$$\tilde{\hat{f}}(s,\omega) = \frac{\tilde{\hat{p}}(s,\omega) + \hat{f}(0)}{s + q + \sqrt{2\pi}r_1\tilde{\hat{k}}(s,\omega)}$$

Hence the solution of (2.5) for $r_2 = 0$ and $q(t) = $ const. is:

$$f(t,x) = \mathcal{F}^{-1}\mathcal{L}^{-1}\left(\frac{\tilde{\hat{p}}(s,\omega) + \hat{f}(0)}{s + q + \sqrt{2\pi}r_1\tilde{\hat{k}}(s,\omega)}\right).$$

It has to be seen if this formula can shed valuable information about the stochastic branching process with settlement.

(4) The spatial transport has been described by a general spatial process $(B(t))_{t\geq0}$, which we assumed to be Brownian motion in our example. Applied to human cancers we know that cell clusters are transported by the circulatory system, hence we need to develop a more realistic body-wide distribution process $(B(t))_{t\geq0}$. This is a complex issue and left for future research.

Acknowledgements

CF and TH gratefully acknowledge financial support by the Natural Sciences and Engineering Research Council of Canada through Discovery Grants. AR gratefully acknowledges funding from an Alberta Innovates Graduate Student Scholarship.

Bibliography

1. A. Bovier (2016): *Gaussian Processes on Trees: From Spin Glasses to Branching Brownian Motion*. Cambridge University Press
2. M.D. Cameron et al. (2000): Temporal Progression of Metastasis in Lung: Cell Survival, Dormancy, and Location Dependence of Metastatic Inefficiency. *Cancer Research*, 60(9): 2541–2546
3. A.F. Chambers, A.C. Groom, and I.C. MacDonald (2002): Dissemination and Growth of Cancer Cells in Metastatic Sites. *Nature Reviews Cancer*, 2(8): 563–572
4. L.A. Coupland, B.H. Chong, and C.R. Parish (2012): Platelets and P-Selectin Control Tumor Metastasis in an Organ-Specific Manner and Independently of NK Cells. *Cancer Research*, 72(18): 4662–4671
5. U. Del Monte (2009): Does the Cell Number 10^9 Still Really Fit One Gram of Tumor Tissue? *Cell Cycle*, 8(3): 505–506
6. R. Durrett (2012): *Essentials of Stochastic Processes*. Second edition. Springer
7. R. Erban, J. Chapman and P. Maini (2008): A Practical Guide to Stochastic Simulations of Reaction-Diffusion Processes. Technical Report, Mathematical Institute, University of Oxford, Oxford, United Kingdom.
8. I. J. Fidler (1970): Metastasis: Quantitative Analysis of Distribution and Fate of Tumor Emboli Labeled With 125 I-5-iodo-2'-deoxyuridine. *Journal of the National Cancer Institute*, 45(773): 773–782
9. P. Friedl and R. Mayor (2017): Tuning Collective Cell Migration by Cell-Cell Junction Regulation. *Cold Spring Harbor Perspectives in Biology*, 9: a029199

10. D.T. Gillespie (1976): A General Method for Numerically Simulating the Stochastic Time Evolution of Coupled Chemical Reactions. *Journal of Computational Physics*, 22: 403–434

11. G.P. Gupta and J. Massague (2006): Cancer Metastasis: Building a Framework. *Cell*, 127(4): 649–695

12. D. Hanahan and R.A. Weinberg (2011): Hallmarks of Cancer: The Next Generation. *Cell*, 144(5): 646–674

13. N. Hartung and C. Gomez (2014): A Stochastic Framework for Secondary Metastatic Emission. Available at hal.archives-ouvertes.fr/hal-01052961

14. P. Henry-Labordère, N. Oudjane, X. Tan, N. Touzi, and X. Warin (2017): Branching Diffusion Representation of Semilinear PDEs and Monte Carlo Approximation. Available at arXiv:1603.01727

15. P. Henry-Labordère, X. Tan and N. Touzi (2014): A Numerical Algorithm for a Class of BSDE via Branching Process. *Stochastic Processes and their Applications*, 124(2): 1112–1140

16. H.W. Hethcote (2000): The Mathematics of Infectious Diseases. *SIAM Review* 42(4): 599–653

17. T. Hillen, B. Greese, J. Martin and G. de Vries (2015): Birth-jump Processes and Application to Forest Fire Spotting. *Journal of Biological Dynamics*, Vol. 9, Suppl. 1, 104–127

18. N. Ikeda, M. Nagasawa, and S. Watanabe (1968): Markov Branching Processes I, *Journal of Mathematics of Kyoto University*, 8:233–278

19. K. Iwata, K. Kawasaki, N. Shigesada (2000): A Dynamical Model for the Growth and Size Distribution of Multiple Metastatic Tumors. *Journal of Theoretical Biology*, 203(2): 177–186

20. T. Jessy (2011): Immunity Over Inability: The Spontaneous Regression of Cancer. *Journal of Natural Science, Biology and Medicine*, 2(1): 43–49

21. R.N. Kaplan, R.D. Riba, S. Zacharoulis et al. (2005): VEGFR1-positive Haematopoietic Bone Marrow Progenitors Initiate the Pre-metastatic Niche, *Nature*, vol. 438: 820–827

22. M.A. Krasnoselskii (1964): *Positive Solutions of Operator Equations*. P. Noordhoff, Groningen, The Netherlands. English translation of Russian original.

23. L. A. Liotta and C. DeLisi (1977): Method for Quantitating Tumor Cell Removal and Tumor Cell-invasive Capacity in Experimental Metastases. *Cancer Research*, 37: 4003–4006

24. K. J. Luzzi et al. (1998): Multistep Nature of Metastatic Inefficiency: Dormancy of Solitary Cells After Successful Extravasation and Limited Survival of Early Micrometastases. *American Journal of Pathology*, 153(865), DOI: 10.1016/S0002-9440(10)65628-3

25. H.P. McKean (1975): Application of Brownian Motion to the Equation of Kolmogorov-Petrovskii-Piskunov. *Communications on Pure and Applied Mathematics*, 28(3): 323–331

26. N. Riggi, M. Aguet and I. Stamenkovic (2018): Cancer Metastasis: a Reappraisal of its Underlying Mechnisms and Their Relevance to Treatment. *Annual Rev. of Pathology: Mechanisms of Disease*, 13:117-140

27. J.G. Scott, P. Gerlee, D. Basanta, A.G. Fletcher, P.K. Maini, and A.R. Anderson (2013): Mathematical Modeling of the Metastatic Process. *In* A. Malek (Ed.), *Experimental Metastasis: Modeling and Analysis*, Springer: 189–208

28. L. Shahriyari (2016): A New Hypothesis: Some Metastases are the Result of Inflammatory Processes by Adapted Cells, Especially Adapted Immune Cells at Sites of

Inflammation. *F1000 Research*, 5(175), DOI: 10.12388/f1000research.8055.1

29. AS. Silva, JA. Yunes, RJ. Gillies, RA. Gatenby (2009): The Potential Role of Systemic Buffers in Reducing Intratumoral Extracellular pH and Acid-mediated Invasion. *Cancer Research*, 69(6): 2677–2684.

30. W.F. Sindelar et al. (1975): Electron Microscopic Observations on Formation of Pulmonary Metastases. *Journal of Surgical Research*, 18(2): 137–161

31. S. Valastyan and R.A. Weinberg (2011): Tumor Metastasis: Molecular Insights and Evolving Paradigms. *Cell*, 147(2): 275–292

32. L. Weiss (1990): Metastatic Inefficiency. *Advances in Cancer Research*, 54: 159–211