1

Title: Differential carbohydrate utilization and organic acid production by honey bee symbionts

[1,2]Fredrick J. Lee, [1]Kayla I. Miller, [1]James B. McKinlay, and [1]Irene L. G. Newton*
1=Department of Biology, Indiana University, Bloomington, IN
2=Current address: Department of Biology, Tufts University, Medford, MA
*corresponding author
1001 E 3[rd] St
Bloomington, IN 47405
812-855-3883
irnewton@indiana.edu

Abstract:

The honey bee worker gut is host to a community of bacteria that primarily comprises 8-10 bacterial species. Collectively, these microbes break down and ferment saccharides present in the host's diet. The model of metabolism for these gut symbionts is rooted in previous analyses of genomes, metagenomes, and metatranscriptomes of this environment. Importantly, there is a correlation between the composition of the gut microbiome and weight gain in the honey bee, suggesting that bacterial production of organic acids might contribute to the observed phenomenon. Here we identify potential metabolic contributions of symbionts within the honey bee gut. We show significant variation in the metabolic capabilities of these microbes, highlighting the fact that although the microbiota appears simple and consistent based on 16S rRNA gene profiling, strains are highly variable in their ability to use specific carbohydrates and produce organic acids. Finally, we confirm that the honey bee core microbes, especially a clade of γ-proteobacteria (i.e. *Gilliamella*), are highly active *in vivo*, expressing key enzymatic genes critical for utilizing plant-derived molecules and producing organic acids. These results suggest that *Gilliamella*, and other core taxa, may contribute significantly to weight gain in the honey bee, specifically through the production of organic acids.

Introduction:

Insects are some of the most abundant and diverse species on earth, inhabiting a plethora of unique environments, ranging from the arid sands of the hottest desert to frigid temperatures of glaciers (1). In order to thrive in such unusual environments, many species of insects have evolved specialized symbioses with microbes capable of synthesizing essential nutrients (*e.g.,* amino acids and vitamins) or degrading complex macromolecules (2). The world's most important agricultural pollinator, the honey bee (*Apis mellifera*), is no exception, as it relies on microbes to aid in the digestion of its plant-derived diet of honey and bee bread (3). After consumption, food ingested by the bee is metabolized by the microbes that inhabit the honey bee gut, which include a small number of dominant taxa that constitute bacterial species considered to be "core" to the honey bee gut microbiota, including the proteobacteria *Snodograssella alvi* (Betaproteobacteria), *Parasaccharibacter apium* (Alphaproteobacteria)*, Frischella perrara* and *Gilliamella apicola* (Gammaproteobacteria), two *Lactobacillus* species (Firmicutes), a *Bifidobacterium* species (Actinobacteria), and a rarer *Bacteroidetes* (4). Three of these phyla, the Gammaproteobacteria, Firmicutes, and Bifidobacteria, dominate predicted functional capacity of the honey bee gut microbiome, based on a metatranscriptomic survey on the honey bee gut (5). The current proposed model of bacterial community metabolism in the bee gut, based on annotated gene presence and expression, suggests that these core microbes participate in the breakdown and fermentation of host dietary macromolecules into an array of alcohols, gases, short chain fatty acids (SCFAs), and other organic acids, some of which likely serve to sustain the host (5-10). Complementing this work, some studies have demonstrated, *in vitro*, the ability of some strains to ferment a variety of substrates (5, 9, 11-14) and produce organic acids (i.e. lactate and acetate)(11, 14). In addition to these observations, a separate study also recently demonstrated that gnotobiotic, mono-colonized bees

possessed guts containing significant depletion in plant derivatives (i.e. flavonoids, ω-

hydroxy acids, and phenolamides), likely originating from pollen consumed by the host,

and an enrichment in fermentative products in the form of organic acids and SCFAs,

compared to bees with a depleted microbiota (15). The production of these fatty acids

may contribute to honey bee health. Indeed, a recent study identified a correlation

between host weight gain and organic acids in the digestive tract of bees (3). Bees

supplemented with gut bacteria showed an increase in the variety and amount of organic

acids present within the host digestive system, as well as a significant increase in host

weight gain and the expression of host genes related to insect growth and nutrient

homeostasis, compared to controls. Collectively, these results link the honey bee

symbionts to the production of organic acids and then to host growth and nutrient

acquisition. It has been proposed that honey bee core bacteria likely convert

carbohydrates in the bee diet to a variety of organic acids, which can either act as a

signaling molecules eliciting a host response, or act as building blocks for host derived

anabolism.

In light of these recent findings, we sought to determine the metabolic

contribution of the Gammaproteobacteria, Firmicutes, and Bifidobacteria, within the bee

gut, as these phyla dominate the metatranscriptomic profile of the digestive tract of the

honey bee. Although there are well-defined species within these groupings, there is still

a large diversity of honey bee strains found within individual bees and colonies (16). The

strain diversity within each of the honey bee core clades is reflected in differences in

genomic content (7, 16), suggesting that significant functional diversity may exist within

each of the core 8-10 clades associated with the honey bee. In this study, we address

fundamental questions regarding metabolic characteristics of the honey bee gut

microbiota, such as: 1) What is the pattern of anaerobic utilization of environmentally

relevant carbon sources across dominant symbionts, and what are the direct byproducts

produced, per taxa, *in vitro*? 2) Are there functional differences in the patterns of

utilization between isolates of the same species? 3) What contribution does an individual

taxon make to the utilization and the subsequent fermentation of environmentally

relevant substrate throughout the digestive tract of the honey bee? Through utilizing a

combination of molecular, biochemical, and physiological approaches, both *in vitro* and

in vivo, we perform an in-depth metabolic analysis on 17 isolates from the honey bee

gut. Our results suggest that overall, bee-associated microbes are primed to metabolize

certain components of the honey bee diet, but that significant variation exists within

clades, suggesting that each member contributes differentially to honey bee metabolism.

Materials and Methods:

Bee Sampling and Bacterial Culture

Worker honey bees were collected from colonies located at an apiary in

Bloomington, IN and transported to lab in sterile vials on ice. Hindgut dissections were

performed using aseptic technique and sterile equipment. Whole digestive tracts of 2-3

bees were homogenized in 1X phosphate buffered saline (PBS) with a sterile disposable

pestle and a dilution series was plated onto Brain-Heart Infusion (BHI), de Man, Rogosa

and Sharpe (MRS), and Luria-Bertani (LB) agar (BD Difco™). All agar plates were

incubated under anaerobic conditions and supplemented with CO_2 using the GasPak EZ

system (BD, New Jersey) at $37^{o}C$ for 4 – 5 days. Isolated colonies were subcultured on

agar before genotyping (below).

16S rRNA genotyping of bacterial isolates

For each isolate, genomic DNA was extracted using the QIAGEN DNeasy Blood &

Tissue kit with one modification: the inclusion of a bead-beating step before lysates were

loaded onto the Qiagen column. DNA was quantified using a spectrophotometer (BioTek

instruments) and used in PCR using Phusion® High Fidelity PCR mix with HF buffer

(New England BioLabs INC.), with 27F (5'-AGA-GTT-TGA-TCC-TGG-CTC-AG-3') and

1492R (5'–ACG-GCT-ACC-TTG-TTA-CGA-CTT-3') primers. A reaction consisted of 0.4

mM primers, 1x Phusion mix, PCR grade water, and 1 µL of sample DNA, under the

following conditions: 5 min at 98^{o}C; 35 cycles-10 sec at 98^{o}C, 30 sec at 55^{o}C, 30 sec

72^{o}C; Extension-10 min at 72^{o}C. PCR products were visualized using gel

electrophoresis and all isolates with PCR products of the correct size were used for

sequencing (Beckman Coulter Genomics). Resulting ab1 files were concatenated and

exported to Fasta. To align, trim, and classify sequences, Mothur v.1.33.3 software was

used with the following commands (align.seqs, trim.seqs, and classify.seqs), using a

custom reference database (Greengeen + honey bee specific dataset) (Newton and

Roeselers, 2011) for classification. To demonstrate the evolutionary relationship

between isolates, a 16S rRNA gene phylogeny was constructed, using the concatenated

16S rRNA gene sequence used for classification. The tree was rooted with *Aquifex*

aeolicus (Accession: AJ309733.1) and to serve as representatives of the four dominant

bacterial families within the honey bee (*Gilliamella apicola* - Accession: NR_118433.1

(Gilliamella), *Lactobacillus* sp. wkB8 - Accession: NZ_CP009531 - Locus Tag:

LACWKB8_RS00480 (Firm-5), Uncultured *Lactobacillus* sp. - Accession: HM113352.1

(Firm-4), and *Bifidobacterium asteroides* strain Hma3 - Accession: EF187236.1 (Bifido)

were used. Sequences were uploaded to SINA 1.2.11 aligner (ARB SILVA), and aligned

using the default settings, after which the FASTA formatted alignment was converted to

MEGA format (MEGA v6.06) and a maximum likelihood tree (GTR + G + I) was

constructed using 1000 bootstrap replicates. Divergence estimates were performed in

BLASTClust (Alva et al, 2016).

Carbohydrate Utilization & Byproduct Production of Isolates

To prepare cell suspensions to assay substrate utilization and product formation,

all isolates classified as actinobacteria or bacilli were grown on MRS agar, while gamma-

proteobacteria isolates were grown on BHI; these media correspond to the medium

under which the microbes were isolated. Broth cultures were grown anaerobically at

30°C for 6 days, after which cultures were pelleted via centrifugation and the supernatant

removed. The pellet was washed with salt solution (137.0 mM NaCl, 2.7 mM KCl, pH

6.0) to remove remaining media. Bacilli and actinobacteria, grown in MRS, were washed

twice to decrease the transfer of nutrients present in the spent MRS media. Isolates

were normalized for growth based on OD, across taxonomic classes, such that isolates

with more robust growth were diluted in a larger amount of salt solution.

To test the ability of isolates to utilize carbon substrates, we used the MT2

MicroPlate™ assay. In the assay, each well in a 96-well plate contained an aliquot of a

TTC dye. To each well, 150 µL of cell suspension from each isolate was added, with or

without 15 µL of a filter-sterilized carbon source (either 1.1 M of D-(+)-glucose (SIGMA

G8270), D-(-)-fructose (ACROS Organics 57-48-7), D-sucrose (Fischer Scintific BP220-

1), D-(+)-mannose (SIGMA M6020), L-rhamnose (SIGMA W373011), D-(+)-galactose

(SIGMA G0625), D-(+)-xylose (SIGMA X1500); or 13.1% D-(+)-cellobiose (SIGMA

22150), or 2.1% pectin (SIGMA – Pectin from apple – SIGMA 76282) solutions).

Negative controls received 15 µL of salt solution. To verify that substrate solutions were

void of microbes and that cross-contamination between wells was not occurring, an

additional negative control was filled with 15 µL of a sole carbon source with 150 µL of

salt solution. Each experimental condition and control was analyzed in triplicate within a

single plate. 96 well plates were covered with an optical adhesive film, and incubated

anaerobically at 30°C for 3 days. To measure reduction of TTC (and utilization of a

substrate), absorbance readings at 590 nm were taken, in triplicate, for each well.

Significant utilization was determined by performing a t-Test comparing 590 nm readings

between experimental (cell suspension + 15 µL substrate) and negative control (cell

suspension + 15 µL of salt solution) wells. All statistics were performed in IBM SPSS

Statistics Version 24 on datasets tested for a normal distribution. Biological triplicates for

each condition or control were then combined in a screw-top cryovial and stored at -80^{o}C

until high performance liquid chromatography (HPLC) analysis (below).

To determine the soluble fermentation products, pooled samples were thawed,

vortexed, and then centrifuged at 13,000 RPM for 5 minutes to pellet bacterial debris.

After centrifugation, samples were filtered through a 0.45 µm HAWP membrane

(Millipore Corp., Bedford, Mass.) and acetate, lactate, formate, and succinate were

quantified by using a Shimadzu HPLC (Kyoto, Japan) as previously described (17).

Assaying the expression of enzymatic genes

For each dissection, the crop, midgut, and hindgut were aseptically incised from

the full digestive tract, and placed in a 2.0 mL Lysing matrix E tube with 500 uL of

Trizol® reagent. Samples were homogenized using a FastPrep®-24 instrument, twice, at

6.0 m/s for 40 seconds. RNA were extracted from samples using the standard protocol

provided by the manufacturer. After the final washing, RNA was resuspended in RNA

Storage Solution (Bioline) and stored at -80^{o}C for downstream analysis. For each gut

chamber, each biological replicate consisted of three separate individuals. All analyses

were performed in triplicate with the same biological samples.

Primer sets specific for a particular enzymatic gene and group of honey bee

microbes (Gilliamella, Firm-5, and Bifido), were designed for qRT-PCR analyses

(Supplementary Table 1, Supplementary Figure 2). Selected primer sets fulfilled the

following requirements: 1) they amplified a product of the expected size, 2) the resulting

sequenced product had a top BLAST hit(s) to the intended bacterial taxa and the gene of

interest, or for the reference sequence used to design the primer set and 3) they showed

specificity for the bacterial group of interest in laboratory experiments (Supplemental

Table 1, Supplemental Figures 2).   RNA extracted from individual gut chambers

(foregut, midgut, and hindgut) were used as a template for qRT-PCR reactions. Prior to

running reactions, all RNA samples were treated with RNase-free DNase I (Ambion) as

prescribed by the manufacture. RT-qPCR was performed using SensiMix ™ SYBER R

Hi-ROX One-Step kit and protocol. As a reference locus in these analyses, we used the

16S rRNA gene, using a universal bacterial primer set (S-*-Univ-05150a0S-19/S-D-Bact-

0787-b-A-20 (18)). Reactions were run on an Applied Biosystems StepOne RT-PCR

system: Initial denaturation-5 min at 98^{o}C; 40 cycles -15 sec at 98^{o}C, 1 min at 60^{o}C, 30

sec 72^{o}C; Extension-10 min at 72^{o}C. A standard curve was generated for each primer

set separately, to account for primer efficiencies in our analyses. All statistics were

performed using IBM SPSS Statistics Version 24.

To quantify the expression of metabolic genes from honey bee specific clades of

bacteria, across gut chambers of the honey bee, Ct values acquired from the

quantification analyses of individual genes were initially normalized to the mean

amplification of the reference locus (16S rRNA gene) across gut sections. The

advantage of normalizing the data to 16S RNA gene is that it demonstrates the presence

of genes, relative to a proxy for the whole bacterial community present throughout the

digestive tract of the honey bee. Next, for each metabolic gene prescribed to a taxa of

interest, we standardized the data to the gut section with the lowest observed

amplification (i.e the crop) for calculating the relative expression (RNA). To determine

the relative ratio of expression of metabolic loci attributed to a particular taxa, across gut

sections, normalized datasets were standardized to the gut section and locus with the

lowest level of expression. To determine variation in the relative abundance, relative

expression, or relative ratio of expression of loci across gut sections, we performed an

ANOVA or Kruskal-Wallis test on the standardized data acquired for each individual

gene analyzed, which was dependent on results obtained from normality test of

individual gene datasets. Results with statistical significance (p ≤ 0.05) were subjected to

pairwise comparisons using one-way ANOVA/post hoc or Mann-Whitney test.

Results:

The honey bee core microbiota utilizes carbohydrates from the host diet

To determine the ability of gut isolates to utilize environmentally relevant

substrates, we isolated 17 honey bee associated bacteria, spanning three

transcriptionally dominant clades: *Gilliamella*, *Lactobacillus* Firm-5, and *Bifidobacterium*

(Figure 1, Supplemental Figure 1). Based on the construction of a 16S rRNA gene

phylogeny, including data from the 17 cultured isolates and 145 sequences previously

obtained from bee-associated microbiota, some of the microbial cultures used herein are

closely related to core honey bee bacteria while others fall outside of the core clades

(Supplemental Figure 1). When comparing the diversity within the clades (based on 16S

rRNA genes), all isolates are unique at 98% and 99% divergence, and some are unique

at 97% divergence (Table 1), indicating that these microbes represent different bacterial

strains and species.

To test the ability of gut bacteria to utilize sole carbon sources relevant to the

honey bee diet, we used Biolog MT2™ plates, containing an aliquot of an indicator

tetrazolium chloride dye (TTC), that when combined with a sole carbon source, along

with an aliquot of culture, detects substrate utilization via a concomitant color change

due to reduction of the dye. Each isolate was examined for the sole utilization of 9

carbon sources (glucose, fructose, sucrose, xylose, pectin, cellobiose, mannose,

rhamnose, and galactose) in pure culture under anaerobic conditions. Honey, which

provides the host with simple carbohydrates (e.g. glucose, fructose, and sucrose) and

trace amounts of secondary sugars (*e.g.* mannose, rhamnose, and galactose) (19-21).

Matured pollen (beebread), provides the host with essential nutrients (i.e. amino acids,

minerals, lipids), and is also rich in complex plant-derived substrates (e.g. cellobiose,

xylose, and pectin)(22, 23). Therefore, the compounds selected for analysis have

previously been identified, through biochemical assays, as molecules that constitute part

of the honey bee diet.

When examining the individual and collective data for the utilization of glucose,

fructose, and sucrose, nearly all isolates were able to utilize at least one or more of

these substrate(s), with the exception of 4 taxa: *Staphylococcaceae* EBHJ3 and

Lactobacillales MAH G07-3 within the Firmicutes and *Enterobacteriaceae* LB Tet D09

and LB Tet C11 (Figure 2, Supplemental Figure 2). At the phylum-level, at least 3

isolates within each phylum were capable of utilizing glucose, fructose, and sucrose,

with varying degrees of consistency (Figure 2A, Supplemental Figure 2A). Collectively,

when comparing isolates representative of taxa specific to the honey bee gut microbiota

(i.e. *Gilliamella*, *Lactobacillus* Firm-5, and *Bifidobacterium*), the general trend for

substrate utilization suggested that the honey bee bacteria were primed to utilize these

substrates (Figure 2). Interestingly, however, we did observe differences among strains

within clades in their ability to use different compounds. For example, although all

Bifidobacteria and *Lactobacillus* Firm-4 strains used glucose, fructose, and sucrose, the

Lactobacillus Firm-5 strain MAH H05-3 did not metabolize either monosaccharaide to a

significant extent. Similarly, *Gilliamella* strains differentially processed sucrose; although

both strains tested could utilize glucose and fructose, only Gamma-1 LB D06 could

utilize sucrose, although to a modest extent (Figure 2).

Acetate, lactate, succinate, and formate are the primary fermentation products produced

by honey bee gut microbiota

To determine the soluble fermentation products produced by each individual

isolate, post anaerobic metabolism, we analyzed the contents within each experimental

and control well of the MT2™ plates described in the previous section using HPLC. For

this analysis, we focused on the products in pure cultures with glucose, fructose, and

sucrose. We chose to use simple sugars abundant in honey, rather than complex

polymeric carbon sources, because complex carbon sources are degraded to simple

sugars and funneled into the same catabolic pathways. With the exception of pectin, all

other saccharides analyzed for utilization are mono- or di- saccharides and are only a

few enzymatic reactions away from glycolysis or the pentose phosphate and Entner-

Doudoroff pathways. While the ratio of fermentation products is expected to change

during growth on different substrates, it is less likely that different products would be

produced from different substrates. We used known standards (*e.g.,* acetate, lactate,

formate, succinate) and identified peaks at the appropriate retention times within each

biological sample.

HPLC analysis revealed a variety of organic acids produced by honey bee

associated microbes, under anaerobic conditions. When comparing the supernatant of

experimental and control wells of isolates, almost all isolates that were positive for

utilizing a substrate exhibited the presence of fermentative products (as derived from

retention times identical to known substrate standards, *i.e.* lactate, acetate, succinate,

and formate; Figure 2). We expected to observe ethanol, acetoin, propanoate, 2,3-

butanediol, based on the metabolic pathways identified in the metatranscriptomic data

but did not identify other significant peaks beyond the four described above, even though

the instrument is capable of detecting these compounds.

When comparing the general pattern of products produced across taxa, bacteria

classified within the *Bifidobacterium* clade unanimously produced lactate and acetate,

regardless of substrate. However, production of succinate and formate by this same

clade was variable both across the phylogeny and across substrates (Figure 2). The

Firm-5 and *Gilliamella* isolates produced these same organic acids, but with an even

larger degree of variability within the clades (Figure 2). Altogether, the data for the 17

isolates supports the hypothesis that honey bee gut microbiota are primed to consume

and ferment the components of the honey bee diet, but significant variation exists within

strains such that no uniform profile could be ascribed to a clade or species group. For

example, even if isolates of the same species used a substrate, they did not produce the

same organic acids; formate was detected from some strains of *Bifidobacterium* under

glucose but not under fructose, while all produced formate when given sucrose (Figure

2).

Microbes utilize more complex carbohydrates and sugars toxic to the honey bee

We extended our analysis to plant derivatives of pollen (cellobiose, xylose, and

pectin), as well as secondary sugars commonly found in honey (mannose, rhamnose,

and galactose) and known to be toxic to the host in high quantities. For the plant derived

carbohydrates, we observed the same trend noted for the utilization of simple sugars,

with at least 2 isolates within each major taxon capable of utilizing cellobiose, xylose,

and pectin. Interestingly, all honey bee specific bacteria tested could utilize pectin while

more distantly related bacteria (*Staphilococcaceae, Lactobacillales, Enterobacteriaceae*)

could not (Figure 3). However, we found variation in the ability of strains within honey

bee associated clades in their use of other complex carbohydrates; xylose was only

utilized by half of the *Bifidobacterium* strains tested and only 1 out of 4 *Lactobacillus*

strains. In contrast, we observed no difference between honey bee specific and more

distantly related bacteria in their ability to utilize sugars toxic to the bee (Figure 3). Again,

however, we observed variation in the ability of strains to utilize these toxic sugars with

only some core members able to use all of these toxic sugars and some core members

unable to use any (*Bifidobacteriales* LB C04 and *Lactobacillus* Firm-4 MaH G10-2,

Figure 3). Overall, these results suggesting the honey bee associated bacteria utilize

mannose, rhamnose, and galactose, secondary sugars common in the honey bee diet,

but toxic to the host.

The honey bee gut compartments are associated with altered gene expression by the

microbiota

We determined that isolates from the honey bee gut microbiota ferment sugars

common to the honey bee diet into organic acids *in vitro* (Figure 2). Recent work by

Zheng et al. (2017) demonstrated *in vivo*, the presence of a variety of organic acids

across distinct gut chambers of the honey bee digestive tract. Based on the

observations described above, we hypothesized that symbionts of the honey bee gut are

likely contributing to presence of organic acids *in vivo*, and since each gut chamber

houses a specific compilation of symbionts (24, 25), perhaps functional differences exist

in the metabolic activity of symbionts across individual gut chambers. To address this

hypothesis, we quantified the expression of genes that encode for fermentative enzymes

throughout the three gut chambers of the honey bee digestive tract (i.e. crop, midgut,

and hindgut (anterior to posterior)). We designed taxon-specific primers for two genes:

acetate kinase (*ackA*) and L-lactate dehydrogenase (*ldh*), encoding for enzymes

responsible for the production of acetate and lactate, respectively. In addition, we also

designed primers for clade specific enzymes (that is, the metabolic function was only

identified in one of the three phyla involved in cleaving glycosidic bonds within plant-

derived compounds (*e.g.* cellulase and cellodextrinase for *Bifidobacteria*, pectate lyase

for *Gilliamella*, and glucosidase for the *Lactobacillus* Firm-5 clade; Supplementary Table

1)). We used qRT-PCR on RNA extracted from each gut section to quantify the

expression of these loci. For the *Gilliamella* and *Lactobacillus* Firm-5 clades, the midgut

and hindgut chambers were found to be the most active gut sections (*Gilliamella* (df = 2;

x^2 = 7.05; p = 0.029), Firm-5 (df = 2; F = 3.95; p = 0.040); Figure 4A and C). In contrast,

while the activity of the *Bifidobacterium* clade increases across gut sections from

posterior to anterior, the trend is not significant ((df = 2; 2 = 4.36; p = 0.113); Figure 4B).

To obtain a better understanding of genes responsible for the trends seen in the

Gilliamella and *Lactobacillus* Firm-5 clades (which both resulted in statistically significant

contrasts across the digestive tract), we performed a one-way ANOVA/Tukey-HSD to

analyze the differences in variation of relative gene expression across gut sections. For

Gilliamella, the analysis of ackA and ldh expression indicated that expression levels

significantly increased across gut sections in the following order: crop<midgut<hindgut

(Figure 4D). Pectate lyase was detected at equal levels of expression in the midgut and

hindgut, but no amplification was detected in the crop (Figure 4D). In contrast to

Gilliamella, we saw no pattern to the expression of ldh or ackA by *Lactobacillus* Firm-5

across the gut sections (Figure 4E). Importantly, we were not able to detect β-

glucosidase and cellulase gene expression from Firm-5 and *Bifidobacterium* clades, in

any gut section, suggesting that they are not active in the honey bee gut.

Discussion:

The honey bee gut microbiota is a deceptively simple community – although

dominated by 8-10 species (26) and transcriptionally dominated by only 3 phyla (5), the

strain diversity found within the gut is high (16). Here we show that strains within three

transcriptionally dominant clades associated with the bee are functionally distinct and

that there is indeed, no strict metabolic rule that describes each clade. Although overall,

the microbes seem primed to utilize carbohydrates in the honey bee diet, expressing

enzymes capable of cleaving glycosidic bonds and fermenting these substrates, strains

within each clade of core microbes investigated here differ in their ability to use different

simple sugars, sugars toxic to the bee, or complex carbohydrates found in the bee diet

(Figures 2, 3). They also differ in their production of organic acids, the production of

which are suspected to contribute to honey bee metabolism. The results presented in

this study reveal that bacterial taxa associated with the honey bee gut are differentially

present and metabolically active throughout the three major gut chambers, with

particular enrichment in the hindgut. Of the three taxa assayed for gene expression, the

Gilliamella clade of bacteria was the most active, showing exceptional expression of its

acetate kinase gene in the mid- and hindgut, a result supported by previous

metatranscriptomic analyses.

The utilization results presented in this study are supported by previous

observations of gross pH changes when honey bee isolates are cultured (9, 11-14) and

previous measures of acetic and lactic acid production from a few pure cultures (27, 28).

Although our results are well supported by predicted models based on the genomic

evidence, we did not detect the production of specific fermentative waste products

predicted (e.g. ethanol, acetoin, propanoate, 2,3-butanediol)(5). The lack of detection of

these products in our analysis could be due to the following: 1) a complete lack of

production under the *in vitro* conditions provided 2) or production below the detectable

range of the UV and refractive index detectors used.

In addition to analyzing the utilization of the substrates mentioned above, we also

examined the ability of isolates to utilize pectin, which is a heteropolysaccharide,

commonly found in the primary cell wall of terrestrial plants, such as pollen grains. In our

analysis, we identified multiple isolates within the *Gilliamella, Lactobacillus* Firm-5, and

Bifidobacterium clades able to utilize pectin (Figure 3). Importantly all tested core

microbiome strains utilized pectin in our assays – it was the only consistently utilized

carbohydrate tested – and more distantly related organisms (e.g. *Staphylococcaceae,*

Lactobacillales, Enterobacteriaceae) were unable to use it. Previous work has shown

that *Gilliamella* isolates are capable of degrading pectin, possessing a pectate lyase

gene capable of cleaving de-esterified pectin (7). In addition to *Gilliamella*, we also

report utilization of pectin and/or its derivatives by isolates within the Firm-4/-5 and

Bifidobacterium clades, observations supported by a recent publication that observed

growth of several core gut isolates in minimal media supplemented with pollen extract

(15). These results suggest that multiple constitutes of the honey bee microbiota may be

capable breaking-down and utilizing plant material abundant in the host's diet, an idea

that has previously been hypothesized (10). Indeed, the genome of *Bifidobacterium*

asteroides has an annotations for glycosidases and a pectinesterase, enzymes capable

of degrading pollen (6). Contrary to *Bifidobacterium*, genomes from bacteria in the

Lactobacillus Firm-5 clade currently have no annotated pectinases or glycosidases (8),

but these and many of the genomes of honey bee specific bacteria possess a plethora of

hypothetical genes of unknown function that might aid in the degradation and

metabolism of pectin (9). Importantly, although the pectin used in our *in vitro* assays was

of high quality, it is not a completely uniform nor pure substance and therefore it is

possible that trace amounts of other compounds might have been present and

metabolized by the honey bee specific microbes. Further experiments need to be

conducted to confirm the pectinase activity of the isolates analyzed. It should also be

noted, that the conditions provided in the utilization assay likely do not support the

growth of all 17 isolates as the salt solution lacks several elements required for growth.

However, although the assay does not rely on growth, it is possible that the utilization

patterns observed for individual isolates could potentially alter under alternative

conditions conducive for growth.

Recent important work has identified a link between the presence of canonical

gut symbionts and weight gain in honey bees; organic acids, which are presumed to be

bacterially derived, are thought to contribute to the observed influence symbionts to host

health (28). The fact that these microbes live within the bee gut and are consistently

associated with honey bees has led to the assumption that microbes isolated from the

bee are mutualists in this context – that they benefit the bee in some way. However,

supplementation of the bee diet with one of these core members (*Snodgrasella)* results

in an increased susceptibility to parasites (29) and another (*Frischella*) results in scab

formation in the intestine (7). The converse can also be true; non-core members of the

bee can provide a benefit as well. *Lactobacillus kunkeii*, an environmentally acquired

microbe, protects from pathogens (30) and although *Gilliamella* strains have been found

to use sugars that are toxic to the bee (31), here we show that both non-core bacteria

and core microbes can utilize these sugars.

The difference in the enzymatic activity pattern between members of the honey

bee microbiota suggests they might occupy different niches in the honey bee food,

digestive tract, or built environment. The idea that niche partitioning might exist in the

gut of the honey bee is physically plausible, as the alimentary system in the honey bee is

subdivided into three distinct chambers. Indeed, our analysis of gene expression (qRT-

PCR) suggests that expression of different enzymes can differ across gut sections,

genes, and taxa. It should be noted, however, that we could not detect the expression of

genes responsible for the breakdown of complex plant-derived molecules (i.e.

glucosidase, cellodextrinase, and cellulase). These results indicate that these genes are

either completely inactive, or are expressed at levels below detection. In either scenario,

the data suggest that perhaps complex carbohydrates are either present at minimal

levels in honey bee gut, or that the taxa analyzed in this study thrive on other resources

abundant in the environment (i.e. glucose, fructose, sucrose, and/or other plant

derivatives) by expressing other enzymatic loci capable of contributing to catabolism.

Our work suggests a dominant profile for *Gillamella in vivo.* Of the three analyzed

taxa, the *Gilliamella* clade was the most active in our analysis, with the *Bifidobacterium*

and *Lactobacillus* Firm-5 clades being roughly equal; this result fits well with previously

published transcriptomic data, where the gamma-proteobacteria dominated the

transcriptional profile of the gut (5). When assaying the relative expression of certain

genes (i.e. ackA and ldh) for the *Gilliamella* clade (Figure 4A, D), we saw highest levels

of expression in the hindgut, suggesting that *Gilliamella* may function differently in each

gut chamber or may be particularly active in the hindgut. Additionally, within each gut

section, *Gilliamella* consistently expresses more ackA than ldh, which might suggest a

preference for this taxon to produce acetate, a byproduct produced *in vitro* by *Gilliamella*

isolates under nearly all tested conditions (Figure 2). It is intriguing that *Gilliamella*

dominates transcriptomic analyses of the honey bee and related species are found in

bumble bees (8). *Gilliamella* and other proteobacterial species in the honey bee require

direct social contact between bees beyond mere trophallaxis (i.e. fecal exposure) to

achieve transmission (32). The fact that the honey bee workers acquire this important

symbiont through their social interactions, and that *Gilliamella* dominates the functional

profile of the honey bee worker gut suggests it to be an important, if not critical, part of

the honey bee core microbiota.

Recent work elegantly demonstrated through metabolomics that honey bees

primarily colonized by individual bacterial strains of taxa core to the gut microbiota show

depletion in plant derived substrates and an enrichment in fermentation product present

in the digestive system, compared to bees with a depleted gut microbiota (33). These

results are promising, as they demonstrate substrate utilization by individual core taxa

and enrichment of metabolites within the gut *in vivo.* However, our work here suggests

that the choice of strain used in colonization experiments may substantially influence

results; not all *Gilliamella apicola* strains produce the same organic acids under the

same conditions. It is perhaps the case that the diversity of strains found within the

honey bee gut complement each other, providing a range of organic acids under most

conditions and diets. This hypothesis awaits testing. Nevertheless, it is obvious that

individual taxa found in the core microbiome of bees might not accurately reflect the

metabolic dynamics occurring in the gut of a honey bee with a natural microbiota, where

microbe-microbe and microbe-host interactions likely govern niche partitioning and

microbiome function.

Acknowledgements:

We thank anonymous reviewers for their feedback on early stages of this manuscript.

Funding: FJL was supported on an NSF Predoctoral Fellowship.

Conflicts of interest statement:
486
487   The authors declare no conflict of interest.
488

References:

1.   **Kohshima S.** 1984. A novel cold-tolerant insect found in a Himalayan glacier.
2.   **Moran NA.** 2007. Symbiosis as an adaptive process and source of phenotypic
complexity. Proceedings of the National Academy of Sciences **104:**8627-
8633.
3.   **Zheng H, Powell JE, Steele MI, Dietrich C, Moran NA.** 2017. Honeybee gut
microbiota promotes host weight gain via bacterial metabolism and
hormonal signaling. Proc Natl Acad Sci U S A **114:**4775-4780.
4.   **Moran NA, Hansen AK, Powell JE, Sabree ZL.** 2012. Distinctive gut
microbiota of honey bees assessed using deep sampling from individual
worker bees. PLoS One **7:**e36393.
5.   **Lee FJ, Rusch DB, Stewart FJ, Mattila HR, Newton IL.** 2015. Saccharide
breakdown and fermentation by the honey bee gut microbiome. Environ
Microbiol **17**.
6.   **Bottacini F, Milani C, Turroni F, Sánchez B, Foroni E, Duranti S, Serafini**
F, Viappiani A, Strati F, Ferrarini A. 2012. Bifidobacterium asteroides
PRL2011 genome analysis reveals clues for colonization of the insect gut.
PLoS One **7:**e44229.

7.   **Engel P, Martinson VG, Moran NA.** 2012. Functional diversity within the
simple gut microbiota of the honey bee. Proceedings of the National Academy
of Sciences **109:**11002-11007.
8.   **Kwong WK, Mancenido AL, Moran NA.** 2014. Genome sequences of
lactobacillus sp. Strains wkB8 and wkB10, members of the Firm-5 Clade,
from honey bee guts. Genome Announc **2.**
9.   **Ellegaard KM, Tamarit D, Javelind E, Olofsson TC, Andersson SG,**
Vásquez A. 2015. Extensive intra-phylotype diversity in lactobacilli and
bifidobacteria from the honeybee gut. BMC Genomics **16:**1-22.
10.   **Kwong WK, Moran NA.** 2016. Gut microbial communities of social bees.
Nature Reviews Microbiology **14:**374-384.
11.   **Olofsson TC, Vásquez A.** 2008. Detection and identification of a novel lactic
acid bacterial flora within the honey stomach of the honeybee Apis mellifera.
Current microbiology **57:**356-363.
12.   **Kwong WK, Moran NA.** 2013. Cultivation and characterization of the gut
symbionts of honey bees and bumble bees: description of Snodgrassella alvi
gen. nov., sp. nov., a member of the family Neisseriaceae of the
Betaproteobacteria, and Gilliamella apicola gen. nov., sp. nov., a member of
Orbaceae fam. nov., Orbales ord. nov., a sister taxon to the order
'Enterobacteriales' of the Gammaproteobacteria. International journal of
systematic and evolutionary microbiology **63:**2008-2018.
13.   **Engel P, Kwong WK, Moran NA.** 2013. Frischella perrara gen. nov., sp. nov.,
a gammaproteobacterium isolated from the gut of the honeybee, Apis
mellifera. International journal of systematic and evolutionary microbiology
63:3646-3651.
14.   **Olofsson TC, Alsterfjord M, Nilson B, Butler E, Vasquez A.** 2014.
Lactobacillus apinorum sp. nov., Lactobacillus mellifer sp. nov., Lactobacillus
mellis sp. nov., Lactobacillus melliventris sp. nov., Lactobacillus kimbladii sp.
nov., Lactobacillus helsingborgensis sp. nov. and Lactobacillus kullabergensis
sp. nov., isolated from the honey stomach of the honeybee Apis mellifera. Int J
Syst Evol Microbiol **64.**
15.   **Kešnerová L, Mars RA, Ellegaard KM, Troilo M, Sauer U, Engel P.** 2017.
Disentangling metabolic functions of bacteria in the honey bee gut. PLoS
biology **15:**e2003467.
16.   **Engel P, Stepanauskas R, Moran NA.** 2014. Hidden diversity in honey bee
gut symbionts detected by single-cell genomics. PLoS Genet **10.**
17.   **McKinlay JB, Zeikus JG, Vieille C.** 2005. Insights into Actinobacillus
succinogenes fermentative metabolism in a chemically defined growth
medium. Applied and environmental microbiology **71:**6651-6656.
18.   **Alm EW, Oerther DB, Larsen N, Stahl DA, Raskin L.** 1996. The
oligonucleotide probe database. Applied and Environmental Microbiology
62:3557.
19.   **Winston ML.** 1991. The biology of the honey bee. Harvard University Press.
20.   **Val A, Huidobro J, Sánchez M, Muniategui S, Fernandez-Muino M, Sancho**
M. 1998. Enzymatic determination of galactose and lactose in honey. Journal
of Agricultural and Food Chemistry **46:**1381-1385.

21.   **Ball DW.** 2007. The chemical composition of honey. Journal of Chemical
556   Education **84:**1643.
557   22.   **De la Fuente E, Sanz M, Martínez-Castro I, Sanz J.** 2006. Development of a
558   robust method for the quantitative determination of disaccharides in honey
559   by gas chromatography. Journal of Chromatography A **1135:**212-218.
560   23.   **Burton RA, Gidley MJ, Fincher GB.** 2010. Heterogeneity in the chemistry,
561   structure and function of plant cell walls. Nature Chemical Biology **6:**724-
562   732.
563   24.   **Corby-Harris V, Maes P, Anderson KE.** 2014. The bacterial communities
564   associated with honey bee (Apis mellifera) foragers. PLoS One **9:**e95056.
565   25.   **Anderson KE, Sheehan TH, Mott BM, Maes P, Snyder L, Schwan MR.** 2013.
566   Microbial ecology of the hive and pollination landscape: bacterial associates
567   from floral nectar, the alimentary tract and stored food of honey bees (Apis
568   mellifera). PLoS One **8**.
569   26.   **Martinson VG, Danforth BN, Minckley RL, Rueppell O, Tingek S, Moran
570   NA.** 2011. A simple and distinctive microbiota associated with honey bees
571   and bumble bees. Molecular Ecology **20:**619-628.
572   27.   **Olofsson TC, Vasquez A.** 2008. Detection and identification of a novel lactic
573   acid bacterial flora within the honey stomach of the honeybee Apis mellifera.
574   Curr Microbiol **57:**356-363.
575   28.   **Li H, Tan J, Song X, Wu F, Tang M, Hua Q, Zheng H, Hu F.** 2017. Sublethal
576   doses of neonicotinoid imidacloprid can interact with honey bee
577   chemosensory protein 1 (CSP1) and inhibit its function. Biochem Biophys
578   Res Commun **486:**391-397.
579   29.   **Schwarz RS, Moran NA, Evans JD.** 2016. Early gut colonizers shape parasite
580   susceptibility and microbiota composition in honey bee workers. Proc Natl
581   Acad Sci U S A **113:**9345-9350.
582   30.   **Vasquez A, Forsgren E, Fries I, Paxton RJ, Flaberg E, Szekely L.** 2012.
583   Symbionts as major modulators of insect health: lactic acid bacteria and
584   honeybees. PLoS One **7**.
585   31.   **Zheng H, Nishida A, Kwong WK, Koch H, Engel P, Steele MI, Moran NA.**
586   2016. Metabolism of Toxic Sugars by Strains of the Bee Gut Symbiont
587   Gilliamella apicola. MBio **7**.
588   32.   **Powell JE, Martinson VG, Urban-Mead K, Moran NA.** 2014. Routes of
589   acquisition of the gut microbiota of Apis mellifera. Appl Environ Microbiol
590   doi:10.1128/AEM.01861-14.
591   33.   **Kesnerova L, Mars RAT, Ellegaard KM, Troilo M, Sauer U, Engel P.** 2017.
592   Disentangling metabolic functions of bacteria in the honey bee gut. PLoS Biol
593   **15:**e2003467.
594
595
596

Tables:

Table 1. Cluster analysis of the 16S rRNA gene sequences of 17 cultured honey bee gut isolates

used for metabolic analysis. A FASTA file with near full-length 16S rRNA sequences for each

isolate was uploaded to BLASTClust (Alva et al., 2016). Standard settings were used for the

analysis, using 97% or 98% identity thresholds. The total number of isolates analyzed for each

class of bacteria is provided, along with the number of phylotype clusters at 97% and 98% ID.

Bacterial Class	Total Number of isolates	Number of Phylotypes (%ID)	
		97	98
γ-Proteobacteria	5	5	5
Bacilli	6	5	6
Actinobacteria	6	1	6

7
8

9
10

Figures:

Figure 1. 16S rRNA gene phylogeny of gut isolates selected for metabolic analyses, in relationship to reference sequences of honey bee specific taxa. Sequences were aligned using SINA Aligner 1.2.11, the output file was used to construct a Maximum likelihood phylogeny, in MEGA 6.06 with GTR (G+I) and 1000 bootstraps. Bootstrap support is provided at nodes. Reference sequences indicated by black text. Cultured isolates indicated by grey text. Taxonomic class highlighted by arrow.

Figure 2. Phenotypic profile of substrate utilization by gut isolates. On the left of the figure are consensus phylogenetic trees derived from the 16S rRNA gene phylogeny. Utilization column is a heat map representation of OD590 measurements (background subtracted) from 1.4 (most purple) to zero (white). Asterisks indicate statistically significant results, compared to negative control, for glucose (A), fructose (B), or sucrose (C). Similarly, the detection of organic acids is denoted by boxes filled will the color that corresponds to a given compound detected: formate (blue), acetate (green), lactate (orange), and succinate (red), based on retention peaks on HPLC chromatograms.

Figure 3. Phenotypic profile of substrate utilization by honey bee isolates. On the left are cladograms derived from the 16S rRNA gene phylogeny. Isolates highlighted in grey indicate taxa considered core to the honey bee gut, while all other isolates are representative of non-core members. The utilization of other plant-derived sugars (i.e. cellobiose, xylose, and pectin) and dietary saccharides toxic to the host (mannose, rhamnose, and galactose) is a heat map representation of OD590 measurements

(background subtracted) from 1.4 (most purple) to zero (white). Asterisks indicate statistically significant results, compared to negative control.

Figure 4. Expression of metabolic genes derived from honey bee specific taxa. Reverse transcriptase quantitative PCR (RT-qPCR) expression shown for each enzymatic genes, relative to the average 16S rRNA gene expression, across chambers of the honey bee gut. Standardized to data from the crop, the overall relative ratio of expression of enzymes for *Gilliamella* (A), *Bifidobacterium* (B), and *Lactobacillus* clades(C) were determined across gut chambers. (D-F) The relative expression data of individual genes, for *Gilliamella* (D), *Bifidobacterium* (E), and *Lactobacillus* (F), across the gut sections. For each dataset, relative expression data were compared across gut sections, using ANOVA or Kruskal-Wallis test in SPSS. For statistically significant results (p ≤ 0.05), pairwise comparisons were conducted using one-way ANOVA/post hoc or Mann-Whitney test in SPSS, significant difference (p ≤ 0.05) between gut sections are denoted with letters.

	Plant-Derivatives				Sugars Toxic To The Host		
	Cellobiose	Xylose	Pectin		Mannose	Rhamnose	Galactose
Bifidobacteriales LB C04	*		*				
Bifidobacteriales MAH G07-1	*	*	*		*		
Bifidobacteriales LB F10-1	*		*		*		
Bifidobacteriales MAH H05-1			*				*
Bifidobacteriales LB A02	*	*	*		*	*	*
Bifidobacteriales LB A06	*	*	*		*		*
Staphylococcaceae EBHJ3-1	*	*			*	*	*
Lactobacillales MAH G07-3							
Lactobacillus Firm-4 MAH G10-2	*		*				
Lactobacillus Firm-5 MAH E12-2	*	*	*		*	*	*
Lactobacillus Firm-5 MAH E11-1	*		*		*	*	*
Lactobacillus Firm-5 MAH H05-3			*		*		*
Gilliamella LB D06	*	*	*		*		
Gilliamella LB Tet D01	*	*	*		*	*	
Enterobacteriaceae LB Tet D09		*			*	*	*
Enterobacteriaceae LB Tet C11		*			*	*	*
Enterobacteriaceae LB Tet C07		*			*	*	*