RNA sequencing (RNA-seq) reveals extremely low levels of reticulocyte-derived globin gene transcripts in peripheral blood from horses (*Equus caballus*) and cattle (*Bos taurus*)

Carolina N. Correia[1], Kirsten E. McLoughlin[1], Nicolas C. Nalpas[1, †], David A. Magee[1],
John A. Browne[1], Kevin Rue-Albrecht[1, §], Stephen V. Gordon[2, 3], David E. MacHugh[1, 3]*

[1] Animal Genomics Laboratory, UCD School of Agriculture and Food Science, UCD College
of Health and Agricultural Sciences, University College Dublin, Belfield, Dublin, D04
V1W8, Ireland.

[2] UCD School of Veterinary Medicine, UCD College of Health and Agricultural Sciences,
University College Dublin, Belfield, Dublin, D04 V1W8, Ireland.

[3] UCD Conway Institute of Biomolecular and Biomedical Research, University College
Dublin, Belfield, Dublin, D04 V1W8, Ireland.

[†] Current address: Quantitative Proteomics and Proteome Centre Tübingen, Interfaculty
Institute for Cell Biology, University of Tübingen, Tübingen, 72076, Germany.

[§] Current address: Kennedy Institute of Rheumatology, University of Oxford, Oxford OX3
7FY, United Kingdom.

* Correspondence: David E. MacHugh, Animal Genomics Laboratory, UCD School of

Agriculture and Food Science, UCD College of Health and Agricultural Sciences, University

College Dublin, Belfield, Dublin D04 V1W8, Ireland. Email: david.machugh@ucd.ie

Keywords: biomarker, blood, cattle, globin, horses, pigs, reticulocyte, RNA-seq,
transcriptome

23
24
25   **Note:** British English language style preferred for publication.

Abstract

RNA-seq has emerged as an important technology for measuring gene expression in peripheral blood samples collected from humans and other vertebrate species. In particular, transcriptomics analyses of whole blood can be used to study immunobiology and develop novel biomarkers of infectious disease. However, an obstacle to these methods in many mammalian species is the presence of reticulocyte-derived globin mRNAs in large quantities, which can complicate RNA-seq library sequencing and impede detection of other mRNA transcripts. A range of supplementary procedures for targeted depletion of globin transcripts have, therefore, been developed to alleviate this problem. Here, we use comparative analyses of RNA-seq data sets generated from human, porcine, equine and bovine peripheral blood to systematically assess the impact of globin mRNA on routine transcriptome profiling of whole blood in cattle and horses. The results of these analyses demonstrate that total RNA isolated from equine and bovine peripheral blood contains very low levels of globin mRNA transcripts, thereby negating the need for globin depletion and greatly simplifying blood-based transcriptomic studies in these two domestic species.

1.   Introduction

It is increasingly recognised that new technological approaches are urgently required for infectious disease diagnosis, surveillance and management in burgeoning domestic animal populations as livestock production intensifies across the globe (Thornton, 2010; Nabarro and Wannous, 2014; Animal Task Force, 2016). In this regard, new strategies have emerged that leverage peripheral blood gene expression to study host immunobiology and to identify panels of RNA transcript biomarkers that can be used as specific biosignatures of infection by particular pathogens for both animal and human infectious disease (Ramilo and Mejias, 2009; Mejias and Ramilo, 2014; Chaussabel, 2015; Ko et al., 2015; Holcomb et al., 2017). For example, we and

others have applied this approach to bovine tuberculosis (BTB) caused by infection with

Mycobacterium bovis (Meade et al., 2007; Killick et al., 2011; Blanco et al., 2012; Churbanov and

Milligan, 2012; McLoughlin et al., 2014; Cheng et al., 2015). It is also important to note that

peripheral blood transcriptomics using technologies such as microarrays or RNA-sequencing

(RNA-seq) can be used to monitor changes in the physiological status of domestic animals due to

reproductive status, diet and nutrition or stress (O'Loughlin et al., 2012; Takahashi et al., 2012;

Song et al., 2013; Kolli et al., 2014; Shen et al., 2014; de Greeff et al., 2016; Elgendy et al., 2016;

Jegou et al., 2016).

High-throughput transcriptomic profiling of whole blood samples can be performed using total

RNA purified from conventional venous blood collection tubes containing anticoagulants (*e.g.*

K_3EDTA, lithium heparin or sodium citrate) or with total RNA generated from dedicated RNA

stabilisation blood collection tubes such as PAXgene® (PreAnalytiX/Qiagen Ltd.) or Tempus™

(Thermo Fisher Scientific). However, a major hindrance to whole blood transcriptomics studies is

the presence of large quantities of globin mRNA transcripts in peripheral blood from many

mammalian species (Wu et al., 2003; Fan and Hegde, 2005; Liu et al., 2006). This is a consequence

of abundant α globin and β globin mRNA transcripts in circulating reticulocytes, which in humans,

may account for more than 95% of the total cellular mRNA content in these immature erythrocytes

(Debey et al., 2004). Reticulocytes, in turn, account for 1–4% of the erythrocytes in healthy adult

humans, which corresponds to between 5×10^7 and 2×10^8 cells per ml compared to 7×10^6 cells

per ml for leukocytes (Greer et al., 2013). Hence, globin transcripts can account for a substantial

proportion of total detectable mRNAs in peripheral blood samples collected from humans and many

other mammals (Bruder et al., 2010; Winn et al., 2010; Schwochow et al., 2012; Choi et al., 2014;

Shin et al., 2014; Bowyer et al., 2015; Huang et al., 2016; Morey et al., 2016). In particular, for

humans, more than 70% of peripheral blood mRNA transcripts are derived from the haemoglobin

subunit alpha 1, subunit alpha 2 and subunit beta genes (*HBA1*, *HBA2* and *HBB*) (Wu et al., 2003;

Field et al., 2007; Mastrokolias et al., 2012).

The emergence of massively parallel transcriptome profiling for clinical applications in human

peripheral blood—initially with gene expression microarrays, but more recently using RNA-seq—

has prompted development of methods for the systematic reduction of globin mRNAs in total RNA

samples purified from peripheral blood samples, including: oligonucleotides that bind to globin

mRNA molecules with subsequent digestion of the RNA strand of the RNA:DNA hybrid (Wu et al.,

2003); peptide nucleic acid (PNA) oligonucleotides that are complementary to globin mRNAs and

block reverse transcription of these targets (Liu et al., 2006); the GLOBINclear™ system, which

uses biotinylated oligonucleotides that hybridise with globin transcripts followed by capture and

separation using streptavidin-coated magnetic beads (Field et al., 2007); and the recently introduced

GlobinLock method that uses a pair of modified oligonucleotides complementary to the 3' portion

of globin transcripts and that block enzymatic extension (Krjutskov et al., 2016).

In the present study we use RNA-seq data generated from globin-depleted and non-depleted total

RNA purified from human and porcine peripheral blood, in conjunction with non-depleted total

RNA isolated from equine and bovine peripheral blood, for a comparative investigation of the

impact of reticulocyte-derived globin mRNA transcripts on routine transcriptome profiling of blood

in domestic cattle and horses.

2. Materials and Methods

2.1. Data sources

RNA-seq data sets from human peripheral whole blood samples used for assessment of globin

depletion and with parallel non-depleted controls (Shin et al., 2014) were obtained from the NCBI

Gene Expression Omnibus (GEO) database (accession number GSE53655). A comparable RNA-

seq data set from globin-depleted and non-depleted porcine peripheral whole blood was obtained

directly from the study authors (Choi et al., 2014). A published RNA-seq data set (Ropka-Molik et

al., 2017) from equine non-depleted peripheral whole blood was obtained from the NCBI GEO

database (accession number GSE83404). Finally, bovine RNA-seq data from peripheral whole

blood were generated by us as described below and can be obtained from the NCBI GEO database

(<mark>accession number to be determined</mark>). A summary overview of the methodology used for the current

study is shown in **Figure 1**.

2.2.   Human, porcine and equine sample collection, globin depletion and RNA-seq libraries

Detailed information concerning ethics approval, sample collection, total RNA extraction, and

RNA-seq library preparation and sequencing for the human, porcine, and equine data sets is

provided in the original publications (Choi et al., 2014; Shin et al., 2014; Ropka-Molik et al., 2017).

Supplementary Table 1 provides summary information on the human, porcine and equine samples

and RNA-seq libraries.

In brief, for the human samples, peripheral blood from six healthy subjects (three females and

three males) was collected into PAXgene blood RNA tubes (PreAnalytiX/Qiagen Ltd., Manchester,

UK). Total RNA, including small RNAs, was purified from the collected blood samples using the

PAXgene Blood miRNA Kit (PreAnalytiX/Qiagen Ltd.) as described by Shin et al. (Shin et al.,

2014). Human *HBA1*, *HBA2* and *HBB* mRNA transcripts were depleted from a subset of the total

RNA samples using the GLOBINclear kit (Invitrogen™/Thermo Fisher Scientific, Loughborough,

UK). RNA-seq data was then generated using 24 paired-end (PE) RNA-seq libraries (12 undepleted

and 12 globin-depleted) generated from the six biological replicates and six technical replicates

created from pooled total RNA across all six biological samples. The multiplexing and sequencing

was then performed such that data for the 12 samples in each treatment group (undepleted and

globin depleted) was generated from two separate lanes of a single flow cell twice, for a total of

four sequencing lanes (Shin et al., 2014).

Porcine peripheral blood samples were collected from 12 healthy crossbred pigs (Duroc ×

(Landrace × Yorkshire)) using Tempus™ blood RNA tubes (Applied Biosystems™/Thermo Fisher

Scientific, Warrington, UK) and total RNA was purified using the MagMAX™ for Stabilized

Blood Tubes RNA Isolation Kit (Invitrogen™/Thermo Fisher Scientific) (Choi et al., 2014). Porcine

HBA and *HBB* mRNA transcripts were subsequently depleted from a subset of the total RNA

samples using a modified RNase H globin depletion method with custom porcine-specific antisense

oligonucleotides for *HBA* and *HBB*. RNA-seq data was then generated from 24 PE RNA-seq

libraries (12 undepleted and 12 globin-depleted).

Equine peripheral blood samples were collected using Tempus™ blood RNA tubes from 12

healthy Arabian horses (five females and seven males) at three different time points during flat

racing training (Ropka-Molik et al., 2017). In addition, peripheral blood samples were collected

from six healthy untrained Arabian horses (two females and four males). Total RNA was purified

using the MagMAX™ for Stabilized Blood Tubes RNA Isolation Kit and 37 of the 42 total RNA

samples were used to generate single-end (SE) libraries for RNA-seq data generation. Globin

depletion for the equine samples was not performed prior to RNA-seq library preparation

(Katarzyna Ropka-Molik, pers. comm.).

2.3.   Bovine peripheral blood collection and RNA extraction

Approximately 3 ml of peripheral blood from ten age-matched healthy male Holstein-Friesian

calves were collected into Tempus™ blood RNA tubes. The Tempus™ Spin RNA Isolation Kit

(Applied Biosystems™/Thermo Fisher Scientific) was used to perform total RNA extraction and

purification, following the manufacturer's instructions. RNA quantity and quality checking were

performed using a NanoDrop™ 1000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA,

USA) and an Agilent 2100 Bioanalyzer using an RNA 6000 Nano LabChip kit (Agilent

Technologies Ltd., Cork, Ireland). The majority of samples displayed a 260/280 ratio > 1.8 and an

RNA integrity number (RIN) > 8.0 (Supplementary Table 2). Globin mRNA depletion was not

performed on the total RNA samples purified from bovine peripheral blood samples.

2.4. Bovine RNA-seq library generation and sequencing

Individually barcoded strand-specific RNA-seq libraries were prepared with 1 µg of total RNA

from each sample. Two rounds of poly(A)$^+$ RNA purification were performed for all RNA samples

using the Dynabeads$^®$ mRNA DIRECT™ Micro Kit (Thermo Fisher Scientific) according to the

manufacturer's instructions. The purified poly(A)$^+$ RNA was then used to generate strand-specific

RNA-seq libraries using the ScriptSeq™ v2 RNA-Seq Library Preparation Kit, the ScriptSeq™

Index PCR Primers (Sets 1 to 4) and the FailSafe™ PCR enzyme system (all sourced from

Epicentre$^®$/Illumina$^®$ Inc., Madison, WI, USA), according to the manufacturer's instructions.

RNA-seq libraries were purified using the Agencourt$^®$ AMPure$^®$ XP system (Beckman Coulter

Genomics, Danvers, MA, USA) according to the manufacturer's instructions for double size

selection (0.75× followed by 1.0× ratio). RNA-seq libraries were quantified using a Qubit$^®$

fluorometer and Qubit$^®$ dsDNA HS Assay Kit (Invitrogen™/Thermo Fisher Scientific), while library

quality checks were performed using an Agilent 2100 Bioanalyzer and High Sensitivity DNA Kit

(Agilent Technologies Ltd.). Individually barcoded RNA-seq libraries were pooled in equimolar

quantities and the quantity and quality of the final pooled libraries (three pools in total) were

assessed as described above. Cluster generation and high-throughput sequencing of three pooled

RNA-seq libraries were performed using an Illumina$^®$ HiSeq™ 2000 Sequencing System at the

MSU   Research   Technology   Support   Facility   (RTSF)   Genomics   Core

(https://rtsf.natsci.msu.edu/genomics; Michigan State University, MI, USA). Each of the three

pooled libraries were sequenced independently on five lanes split across multiple Illumina$^®$ flow

cells. The pooled libraries were sequenced as PE 2×100 nucleotide reads using Illumina® version

5.0 sequencing kits.

Deconvolution (filtering and segregation of sequence reads based on the unique RNA-seq library

barcode index sequences; Supplementary Table 2) was performed by the MSU RTSF Genomics

Core using a pipeline that simultaneously demultiplexed and converted pooled sequence reads into

discrete FASTQ files for each RNA-seq sample with no barcode index mismatches permitted. The

RNA-seq FASTQ sequence read data for the bovine samples were obtained from the MSU RTSF

Genomics Core FTP server.

2.5. RNA-seq data quality control and filtering/trimming of reads

Bioinformatics procedures and analyses were performed as described below for the human,

porcine, equine, and bovine samples, except were specifically indicated. All of the bioinformatics

workflow scripts were developed using GNU bash (version 4.3.48) (Free Software Foundation,

2013), Python (version 3.5.2) (Python Software Foundation, 2017), and R (version 3.4.0) (R Core

Team, 2017). The scripts and further information are available at a public GitHub repository

(https://github.com/carolcorreia/Globin_RNA-sequencing).   Computational   analyses   were

performed on a 32-core Linux Compute Server (4× AMD Opteron™ 6220 processors at 3.0 GHz

with 8 cores each), with 256 GB of RAM, 24 TB of hard disk drive storage, and with Ubuntu Linux

OS (version 14.04.4 LTS). Deconvoluted FASTQ files (generated from SE equine RNA-seq

libraries and PE RNA-seq libraries for the other species) were quality-checked with FastQC

(version 0.11.5) (Andrews, 2016).

Using the ngsShoRT software package (version 2.2) (Chen et al., 2014), filtering/trimming

consisted of: (1) removal of SE or PE reads with adapter sequences (with up to three mismatches);

(2) removal of SE or PE reads of poor quality (i.e., at least one of the reads containing $\geq 25\%$ bases

with a Phred quality score below 20); (3) for porcine samples only, 10 bases were trimmed at the 3'

end of all reads; (4) removal of SE or PE reads that did not meet the required minimum length (70

nucleotides for human and equine, 80 nucleotides for porcine and 100 nucleotides for bovine).

Filtered/trimmed FASTQ files were then re-evaluated using FastQC. Filtered FASTQ files were

transferred to a 36-core/64-thread Compute Server (2× Intel® Xeon® CPU E5-2697 v4 at 2.30 GHz

with 18 cores each), with 512 GB of RAM, 96 TB SAS storage (12× 8 TB at 7200 rpm), 480 GB

SSD storage, and with Ubuntu Linux OS (version 16.04.2 LTS).

2.6. Transcript quantification

The Salmon software package (version 0.8.2) (Patro et al., 2017) was used in quasi-mapping-

mode for transcript quantification. Sequence-specific and fragment-level GC bias correction was

enabled and transcript abundance was quantified in transcripts per million (TPM) for each filtered

library (multiple lanes from the same library were processed together) was estimated after mapping

of SE or PE reads to their respective reference transcriptomes.

As summarised in **Table 1**, the current annotations for equine and porcine haemoglobin subunit

alpha genes are inconsistent. Therefore, transcript quantification was divided into two parallel

analyses as follows: (1) a primary analysis that used NCBI RefSeq reference transcript models for

the human, porcine, and bovine data sets, and where the Table Browser data retrieval tool for the

UCSC Genome Browser Database was used for the equine data set (Karolchik et al., 2004; Casper

et al., 2018); (2) a secondary analysis where the equine data set was analysed using annotated

transcripts from the NCBI RefSeq database. Detailed information about the reference

transcriptomes used is provided in Supplementary Table 3.

2.7. Gene annotations and summarisation of TPM estimates at the gene level

Using R within the RStudio IDE (version 1.0.143) (RStudio Team, 2015) and Bioconductor

(version 3.5 using BiocInstaller 1.26.0) (Gentleman et al., 2004), the GenomicFeatures (version

1.28.0) (Lawrence et al., 2013) and AnnotationDbi (version 1.38.0) (Pagès et al., 2017) packages

were used to obtain corresponding gene and transcript identifiers from the relevant annotation

source for each species as detailed in **Table 1**. Using these identifiers, the tximport (version 1.4.0)

package (Soneson et al., 2015) was used to import into R and summarise at gene level the TPM

estimates obtained from the Salmon tool.

2.8.   Filtering of lowly expressed genes

For both the primary and secondary analyses, non-expressed genes (i.e. the sum of TPM in all

samples is equals to zero) were removed, and genes were considered to be expressed above a

threshold set to an average TPM $\geq$ 1.

2.9.   Data exploration, plotting and summary statistics

Data from the primary and secondary analyses were processed using the following R packages

tidyverse (version 1.1.1) (Wickham, 2017b), dplyr (version 0.7.2) (Wickham et al., 2017), tidyr

(version 0.7.0) (Wickham and Henry, 2017), reshape2 (version 1.4.2) (Wickham, 2017a), and

magrittr (version 1.5) (Bache and Wickham, 2017). The ggplot2 (version 2.2.1) (Wickham and

Chang, 2017), ggjoy (version 0.3.0) (Wilke, 2017b), and cowplot (version 0.7.0) (Wilke, 2017a)

packages were used for figure generation.

For the primary and secondary analyses, summary statistics were calculated for the undepleted

and globin-depleted groups in each species using the skimr (version 0.9000) R package (McNamara

et al., 2017).

3.   Results and Discussion

3.1.   Status of human, porcine, equine and bovine haemoglobin gene annotations

Annotation of the haemoglobin subunit alpha 1 and 2 genes (*HBA1* and *HBA2*, respectively) is

well established for the human genome; however, annotations for the porcine, equine and bovine

genomes are inconsistent. The equine *HBA1* and *HBA2* genes were either absent from the current

NCBI annotation release (NCBI *Equus caballus* Annotation Release 102, November 2015) or were

not annotated in the reference assembly (EquCab2.0, October 2007), despite both being assigned an

NCBI Entrez Gene ID. In addition, equine *HBA1* and *HBA2* were also absent from the Ensembl

database. To the best of our knowledge, and at the time this analysis was performed, the UCSC

Table Browser (equCab2.refGene track, last updated on August 2017) was the only source of

annotation for equine *HBA1* and *HBA2* genes (**Table 1**). Other genes lacking annotation were

porcine *HBA* and bovine *HBA1* in the UCSC Table Browser and equine *HBA1* and *HBA2* in

Ensembl (**Table 1**). Although it contained both the *HBA1* and *HBA2* genes, the UCSC

equCab2.refGene annotation track contained only 1,875 entries (August 2017). In contrast to this,

UCSC hg38.refGene, susScr3.refGene and bosTau8.refGene had 73,426, 4,548, and 14,254 genes

respectively (August 2017).

For the NCBI RefSeq genome annotations, porcine *HBA* has been assigned to two loci

(*LOC110259958* and *LOC100737768*) that have similar descriptions (haemoglobin subunit alpha

and haemoglobin subunit alpha-like). Therefore these locus symbols were used (**Table 1**). Equine

HBA is described as haemoglobin subunit alpha 1 and is referred to here as *HBA1*; whereas bovine

HBA is described as haemoglobin subunit alpha 2 and is referred to here as *HBA2*. In contrast to

this, the haemoglobin subunit beta (*HBB*) genes for the four species are well annotated in Ensembl,

NCBI RefSeq and UCSC Genome Browser databases (**Table 1**).

3.2.   Basic RNA-seq data outputs

Unfiltered SE (equine libraries) or PE (human, porcine, and bovine libraries) RNA-seq FASTQ

files were quality-checked, adapter- and quality-filtered prior to transcript quantification. As shown

in **Table 2**, the human and porcine undepleted groups each had approximately 40 million (M) raw

reads per library. After globin depletion, total raw read pairs were reduced by an average of 3.3 M

(8.3%) in human and 7.7 M (19.7%) in porcine libraries. Equine and bovine libraries, which did not

include a globin depletion step had an average of 24.3 M raw reads and 20.5 M raw read pairs,

respectively.

After adapter- and quality-filtering of RNA-seq libraries, an average of 20.4% and 29.0% read

pairs were removed from the human undepleted and globin-depleted libraries, respectively.

Conversely, 11.8% and 12.4% of read pairs were removed from the porcine undepleted and globin-

depleted libraries, respectively. For the undepleted equine (primary and secondary analyses) and

bovine RNA-seq libraries, an average of 0.2% reads and 17% read pairs were removed,

respectively. Detailed information for RNA-seq libraries from all species, including technical

replicates from libraries sequenced over multiple lanes, is presented in Supplementary Table 4.

3.3.   RNA-seq mapping

In the primary analysis, human, porcine and bovine data sets all exhibited a mean mapping rate

greater than 70% (**Table 2**). Conversely, for the equine data set, a mean mapping rate of only 11.5%

was observed when the UCSC refGene resource was used as the reference transcriptome (**Table 2**).

This can be explained by the smaller number of entries—a total of 1,875 RNA sequences—in the

UCSC equCab2 reference transcriptome file (Supplementary Table 3). This problem was resolved

for the secondary analysis by using the NCBI RefSeq reference file, which had 42,199 entries and

yielded a mean of 71.4% mapped reads (**Table 2**). Supplementary Tables 5 and 6 contain sample-

specific RNA-seq mapping statistics for the primary and secondary analyses, respectively. Notably,

the UCSC reference mRNA data is based on weekly GenBank updates; therefore, the difference in

the numbers of RNA sequences between UCSC and NCBI RefSeq reference files is due to the

former resource including only accession numbers from known RefSeq records (NM_ and NR_

prefixes), whereas the latter includes both known and model RefSeq accessions (NM_, NR_, XM_,

and XR prefixes). More information about the NCBI annotation pipelines is available at the

following link: https://www.ncbi.nlm.nih.gov/refseq/about.

3.4.   Transcript quantification

Transcript-level TPM estimates generated using the Salmon tool were imported into the R

environment and summarised at gene level with the package tximport. Gene-level TPM estimates

represent the sum of corresponding transcript-level TPMs and provide results that are more accurate

and comprehensible than transcript-level estimates (Soneson et al., 2015). In the current study gene-

level TPM estimates are referred as TPM and genes were considered expressed when mean TPM ≥

1.

In the primary analysis, undepleted and globin-depleted human samples exhibited mean numbers

of 11,577 and 11,946 expressed genes, respectively—corresponding to a 3.2% increase after globin

depletion. Porcine samples showed a substantially larger increase of 19.6% in expressed genes after

globin depletion with a mean of 9,167 genes, compared to 7,666 genes before depletion. The equine

and bovine samples, which did not undergo globin depletion, exhibited mean numbers of 860 and

12,275 expressed genes, respectively. However, for the secondary analysis that used the NCBI

RefSeq reference file, the equine samples exhibited a mean of 11,587 expressed genes. The

numbers of expressed genes in individual samples are provided in Supplementary Table 7 for the

primary analysis (human, porcine, equine and bovine), and Supplementary Table 8, for the

secondary equine analysis.

The density distribution of TPM values for the human and porcine samples improved after

globin depletion; this is evident by the shift of gene detection levels towards greater $\log_{10}$ TPM

values for the globin-depleted samples in panel A **Figure 2**. In this regard, it is noteworthy that the

undepleted bovine samples from the primary analysis and the undepleted equine samples from the

secondary analysis (panel B) also exhibited similar TPM density distributions to the human and

porcine globin-depleted samples. As might be expected equine samples from primary analysis,

which used the limited UCSC refGene annotation source, had the widest range of TPM densities

(panel A, **Figure 2**).

3.5. Proportions of human and porcine haemoglobin gene transcripts in undepleted and
depleted peripheral blood

In line with previous reports (Field et al., 2007; Mastrokolias et al., 2012), the proportion of

haemoglobin gene transcripts (*HBA1*, *HBA2*, and *HBB*) detected in undepleted human peripheral

blood samples for the current study averaged 68% (**Figure 3** and Supplementary Table 7), which is

substantially lower than the mean proportion of 81% reported by Shin et al. (2014). On the other

hand, after depletion the human samples exhibited an identical reduction to a 17% proportion of

globin sequence reads in both the present study and that of Shin et al. (2014) (**Figure 3** and

Supplementary Table 7).

In the current study, for the undepleted porcine peripheral blood samples, the percentage of

haemoglobin gene transcripts (*LOC110259958* [*HBA*], *LOC100737768* [*HBA*], and *HBB*) observed

as a proportion of the total expressed genes was 72% (**Figure 3** and Supplementary Table 7), which

is considerably larger than the mean of 46.1% reported in the original study (Choi et al., 2014).

Similarly, after depletion, the porcine samples in the present study contained a mean proportion of

22% globin transcripts (**Figure 3** and Supplementary Table 7) compared to a mean proportion of

8.9% reported by Choi et al. (2014).

A number of possible explanations, including the different approaches used for read mapping

and transcript quantification, may account for the different proportions of haemoglobin gene

transcript detected in human and porcine samples for the present study compared to the original

studies (Choi et al., 2014; Shin et al., 2014). For the present study, a lightweight alignment method

was adopted (Salmon and tximport), in contrast to the older methodologies used in the original

publications. Shin and colleagues (2014) used the TopHat and Cufflinks software tools (Trapnell et

al., 2012), while Choi et al. (2014) implemented TopHat with Htseq-count (Anders et al., 2015). In

addition to this, different gene annotations were used: NCBI *Homo sapiens* Annotation Release 108

and NCBI *Sus scrofa* Annotation Release 106 were used for the present study, while UCSC hg18

(*Homo sapiens*) and Ensembl release 71 (*Sus scrofa*) were used by Shin et al. (2014) and Choi et al.

(2014), respectively.

3.6. Equine and bovine peripheral blood contains extremely low levels of haemoglobin gene transcripts

The equine and bovine peripheral blood samples, which did not undergo globin depletion, had

extremely low proportions of haemoglobin gene transcripts (*HBA1*, *HBA2*, and *HBB*) to total

expressed genes: 0.79% and 0.17%, respectively (**Figure 3** and Supplementary Table 7). In this

regard, expression of *HBA1* in four bovine samples was actually so low that it was excluded by the

filtering threshold of TPM $\geq$ 1 (**Table 3** and Supplementary Table 7). Notably, similar results have

been reported in a transcriptomics study of bovine peripheral blood in response to vaccination

against neonatal pancytopenia. In that study, 12 cows were profiled before and after vaccination (24

peripheral blood samples in total), and a mean proportion of 1.0% of RNA-seq reads were observed

to map to the bovine α haemoglobin gene cluster on BTA25 or to the β haemoglobin gene cluster on

BTA15 (Demasius et al., 2013).

Finally, it is important to note that $\log_2$ TPM values for haemoglobin gene transcripts in the

undepleted equine and bovine peripheral blood RNA samples are substantially lower than $\log_2$ TPM

values for the globin-depleted human and porcine peripheral blood RNA samples (**Figure 4**). This

is a direct consequence of extremely low levels of circulating reticulocytes in equine and bovine

peripheral blood (Tablin and Weiss, 1985; Harper et al., 1994; Hossain et al., 2003; Cooper et al.,

2005).

3.7. Conclusion

In light of our RNA-seq data analyses, we propose that globin mRNA transcript depletion is not a pre-requisite for transcriptome profiling of bovine and equine peripheral blood samples. This observation greatly simplifies the laboratory and bioinformatics workflows required for RNA-seq studies of whole blood collected from domestic cattle and horses. It will also be directly relevant to future work on blood-based biomarker and biosignature development in the context of infectious disease, reproduction, nutrition and animal welfare. For example, transcriptomics of peripheral blood has been used extensively in development of new diagnostic and prognostic modalities for human tuberculosis (HTB) disease caused by infection with *Mycobacterium tuberculosis* (for reviews see: Blankley et al., 2014; Haas et al., 2016; Weiner and Kaufmann, 2017; Goletti et al., 2018). Therefore, as a consequence of this HTB research, comparable transcriptomics studies in cattle (Meade et al., 2007; Killick et al., 2011; Blanco et al., 2012; Churbanov and Milligan, 2012; McLoughlin et al., 2014; Cheng et al., 2015), and the ease with which RNA-seq can be performed in bovine peripheral blood, it should be feasible to develop transcriptomics-based biomarkers and biosignatures for BTB caused by *M. bovis* infection.

4.   Data Accessibility

The RNA-seq data generated for this study using peripheral blood from ten age-matched healthy male Holstein-Friesian calves can be obtained from the NCBI GEO database (accession number to be determined).

5.   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

6. Ethics Statement

Animal experimental work for the present study (cattle samples) was carried out according to the UK Animal (Scientific Procedures) Act 1986. The study protocol was approved by the Animal Health and Veterinary Laboratories Agency (AHVLA–Weybridge, UK), now the Animal & Plant Health Agency (APHA), Animal Use Ethics Committee (UK Home Office PCD number 70/6905).

7. Author Contributions

DEM, SVG, CNC, and KEM conceived and designed the project and organised bovine sample collection; KEM, NCN, DAM, and JAB performed RNA extraction and RNA-seq library generation; CNC, KEM, NCN, KRA, and DEM performed the analyses; CNC and DEM wrote the manuscript and all authors reviewed and approved the final manuscript.

8. Funding

This work was supported by Investigator Grants from Science Foundation Ireland (Nos: SFI/08/IN.1/B2038 and SFI/15/IA/3154), a Research Stimulus Grant from the Department of Agriculture, Food and the Marine (No: RSF 06 405), a European Union Framework 7 Project Grant (No: KBBE-211602- MACROSYS), a Brazilian Science Without Borders – CAPES grant (No: BEX-13070-13-4) and the UCD Wellcome Trust funded Computational Infection Biology PhD Programme (Grant no: 097429/Z/11/Z).

9. Acknowledgements

The authors wish to express their gratitude to Prof Martin Vordermeier and Dr Bernardo Villarreal-Ramos (Animal and Plant Health Agency, UK) for provision of bovine peripheral blood

409   samples, Prof Graham Plastow (University of Alberta, Canada) for provision of porcine peripheral

410   blood RNA-seq data. We also thank Drs Gabriella Farries and Kerri Malone (University College

411   Dublin) for stimulating discussion and advice concerning genome annotations, equine genetics and

412   data visualisation.

10. References

Aken, BL, Achuthan, P, Akanni, W, Amode, MR, Bernsdorff, F, Bhai, J, Billis, K, Carvalho-Silva, D, Cummins, C, Clapham, P, Gil, L, Giron, CG, Gordon, L, Hourlier, T, Hunt, SE, Janacek, SH, Juettemann, T, Keenan, S, Laird, MR, Lavidas, I, Maurel, T, McLaren, W, Moore, B, Murphy, DN, Nag, R, Newman, V, Nuhn, M, Ong, CK, Parker, A, Patricio, M, Riat, HS, Sheppard, D, Sparrow, H, Taylor, K, Thormann, A, Vullo, A, Walts, B, Wilder, SP, Zadissa, A, Kostadima, M, Martin, FJ, Muffato, M, Perry, E, Ruffier, M, Staines, DM, Trevanion, SJ, Cunningham, F, Yates, A, Zerbino, DR, and Flicek, P (2017). Ensembl 2017. *Nucleic Acids Res* 45**,** D635-d642. doi: 10.1093/nar/gkw1104.

Anders, S, Pyl, PT, and Huber, W (2015). HTSeq–a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31. doi: 10.1093/bioinformatics/btu638.

Andrews, S. 2016. FastQC: a quality control tool for high throughput sequence data. Available: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Animal Task Force. 2016. A strategic research and innovation agenda for a sustainable livestock sector in Europe. Available: http://www.animaltaskforce.eu.

Bache, SM, and Wickham, H. 2017. magrittr: a forward-pipe operator for R. Available: https://github.com/tidyverse/magrittr.

Blanco, FC, Soria, M, Bianco, MV, and Bigi, F (2012). Transcriptional response of peripheral blood mononuclear cells from cattle infected with *Mycobacterium bovis*. *PLoS ONE* 7**,** e41066. doi: 10.1371/journal.pone.0041066.

Blankley, S, Berry, MP, Graham, CM, Bloom, CI, Lipman, M, and O'Garra, A (2014). The application of transcriptional blood signatures to enhance our understanding of the host response to infection: the example of tuberculosis. *Philos Trans R Soc Lond B Biol Sci* 369**,** 20130427. doi: 10.1098/rstb.2013.0427.

Bowyer, JF, Tranter, KM, Hanig, JP, Crabtree, NM, Schleimer, RP, and George, NI (2015). Evaluating the stability of RNA-seq transcriptome profiles and drug-induced immune-related expression changes in whole blood. *PLoS ONE* 10**,** e0133315. doi: 10.1371/journal.pone.0133315.

Bruder, CE, Yao, S, Larson, F, Camp, JV, Tapp, R, McBrayer, A, Powers, N, Granda, WV, and Jonsson, CB (2010). Transcriptome sequencing and development of an expression microarray platform for the domestic ferret. *BMC Genomics* 11**,** 251. doi: 10.1186/1471-2164-11-251.

Casper, J, Zweig, AS, Villarreal, C, Tyner, C, Speir, ML, Rosenbloom, KR, Raney, BJ, Lee, CM, Lee, BT, Karolchik, D, Hinrichs, AS, Haeussler, M, Guruvadoo, L, Navarro Gonzalez, J,

Gibson, D, Fiddes, IT, Eisenhart, C, Diekhans, M, Clawson, H, Barber, GP, Armstrong, J,
Haussler, D, Kuhn, RM, and Kent, WJ (2018). The UCSC Genome Browser database: 2018
update. *Nucleic Acids Res* 46**,** D762-d769. doi: 10.1093/nar/gkx1020.

Chaussabel, D (2015). Assessment of immune status using blood transcriptomics and potential
implications for global health. *Semin Immunol* 27**,** 58-66. doi: 10.1016/j.smim.2015.03.002.

Chen, C, Khaleel, SS, Huang, H, and Wu, CH (2014). Software for pre-processing Illumina next-
generation sequencing short read sequences. *Source Code Biol Med* 9**,** 8. doi: 10.1186/1751-
0473-9-8.

Cheng, Y, Chou, C-H, and Tsai, H-J (2015). In vitro gene expression profile of bovine peripheral
blood mononuclear cells in early *Mycobacterium bovis* infection. *Exp Ther Med* 10**,** 2102-
2118. doi: 10.3892/etm.2015.2814.

Choi, I, Bao, H, Kommadath, A, Hosseini, A, Sun, X, Meng, Y, Stothard, P, Plastow, GS, Tuggle,
CK, Reecy, JM, Fritz-Waters, E, Abrams, SM, Lunney, JK, and Guan le, L (2014).
Increasing gene discovery and coverage using RNA-seq of globin RNA reduced porcine
blood samples. *BMC Genomics* 15**,** 954. doi: 10.1186/1471-2164-15-954.

Churbanov, A, and Milligan, B (2012). Accurate diagnostics for bovine tuberculosis based on high-
throughput sequencing. *PLoS ONE* 7**,** e50147. doi: 10.1371/journal.pone.0050147.

Cooper, C, Sears, W, and Bienzle, D (2005). Reticulocyte changes after experimental anemia and
erythropoietin treatment of horses. *J Appl Physiol* 99**,** 915-921. doi:
10.1152/japplphysiol.00438.2005.

de Greeff, A, Bikker, P, Smit-Heinsbroek, A, Bruininx, E, Zwolschen, H, Fijten, H, Zetteler, P,
Vastenhouw, S, Smits, M, and Rebel, J (2016). Increased fat and polyunsaturated fatty acid
content in sow gestation diet has no effect on gene expression in progeny during the first 7
days of life. *J Anim Physiol Anim Nutr (Berl)* 100**,** 127-135. doi: 10.1111/jpn.12345.

Debey, S, Schoenbeck, U, Hellmich, M, Gathof, BS, Pillai, R, Zander, T, and Schultze, JL (2004).
Comparison of different isolation techniques prior gene expression profiling of blood
derived cells: impact on physiological responses, on overall expression and the role of
different cell types. *Pharmacogenomics J* 4**,** 193-207. doi: 10.1038/sj.tpj.6500240.

Demasius, W, Weikard, R, Hadlich, F, Muller, KE, and Kuhn, C (2013). Monitoring the immune
response to vaccination with an inactivated vaccine associated to bovine neonatal
pancytopenia by deep sequencing transcriptome analysis in cattle. *Vet Res* 44**,** 93. doi:
10.1186/1297-9716-44-93.

Elgendy, R, Giantin, M, Castellani, F, Grotta, L, Palazzo, F, Dacasto, M, and Martino, G (2016).
Transcriptomic signature of high dietary organic selenium supplementation in sheep: A

nutrigenomic insight using a custom microarray platform and gene set enrichment analysis.

J Anim Sci 94**,** 3169-3184. doi: 10.2527/jas.2016-0363.

Fan, H, and Hegde, PS (2005). The transcriptome in blood: challenges and solutions for robust

expression profiling. *Curr Mol Med* 5**,** 3-10. doi: 10.2174/1566524053152861.

Field, LA, Jordan, RM, Hadix, JA, Dunn, MA, Shriver, CD, Ellsworth, RE, and Ellsworth, DL

(2007). Functional identity of genes detectable in expression profiling assays following

globin mRNA reduction of peripheral blood samples. *Clin Biochem* 40**,** 499-502. doi:

10.1016/j.clinbiochem.2007.01.004.

Free Software Foundation. 2013. Bash (4.3.48) Unix shell program. Available:

http://ftp.gnu.org/gnu/bash/.

Gentleman, RC, Carey, VJ, Bates, DM, Bolstad, B, Dettling, M, Dudoit, S, Ellis, B, Gautier, L, Ge,

Y, Gentry, J, Hornik, K, Hothorn, T, Huber, W, Iacus, S, Irizarry, R, Leisch, F, Li, C,

Maechler, M, Rossini, AJ, Sawitzki, G, Smith, C, Smyth, G, Tierney, L, Yang, JY, and

Zhang, J (2004). Bioconductor: open software development for computational biology and

bioinformatics. *Genome Biol* 5**,** R80. doi: 10.1186/gb-2004-5-10-r80.

Goletti, D, Lee, MR, Wang, JY, Walter, N, and Ottenhoff, THM (2018). Update on tuberculosis

biomarkers: From correlates of risk, to correlates of active disease and of cure from disease.

Respirology. doi: 10.1111/resp.13272.

Greer, JP, Arber, DA, Glader, B, List, AF, Means, RT, Paraskevas, F, Rodgers, GM, and Foerster, J

(2013). *Wintrobe's Clinical Hematology*. Philadelphia, PA, USA: Lippincott, Williams and

Wilkins.

Haas, CT, Roe, JK, Pollara, G, Mehta, M, and Noursadeghi, M (2016). Diagnostic 'omics' for active

tuberculosis. *BMC Med* 14**,** 37. doi: 10.1186/s12916-016-0583-9.

Harper, SB, Hurst, WJ, Ohlsson-Wilhelm, B, and Lang, CM (1994). The response of various

hematologic parameters in the young bovine subjected to multiple phlebotomies. *ASAIO J*

40**,** M816-825.

Hestand, MS, Kalbfleisch, TS, Coleman, SJ, Zeng, Z, Liu, J, Orlando, L, and MacLeod, JN (2015).

Annotation of the protein coding regions of the equine genome. *PLoS ONE* 10**,** e0124375.

doi: 10.1371/journal.pone.0124375.

Holcomb, ZE, Tsalik, EL, Woods, CW, and McClain, MT (2017). Host-based peripheral blood

gene expression analysis for diagnosis of infectious diseases. *J Clin Microbiol* 55**,** 360-368.

doi: 10.1128/jcm.01057-16.

Hossain, MA, Yamato, O, Yamasaki, M, Otsuka, Y, and Maede, Y (2003). Relation between
reticulocyte count and characteristics of erythrocyte 5'-nucleotidase in dogs, cats, cattle and
humans. *J Vet Med Sci* 65**,** 193-197. doi: 10.1292/jvms.65.193.

Huang, Z, Gallot, A, Lao, NT, Puechmaille, SJ, Foley, NM, Jebb, D, Bekaert, M, and Teeling, EC
(2016). A nonlethal sampling method to obtain, generate and assemble whole blood
transcriptomes from small, wild mammals. *Mol Ecol Resour* 16**,** 150-162. doi:
10.1111/1755-0998.12447.

Jegou, M, Gondret, F, Vincent, A, Trefeu, C, Gilbert, H, and Louveau, I (2016). Whole blood
transcriptomics is relevant to identify molecular changes in response to genetic selection for
feed efficiency and nutritional status in the pig. *PLoS ONE* 11**,** e0146550. doi:
10.1371/journal.pone.0146550.

Karolchik, D, Hinrichs, AS, Furey, TS, Roskin, KM, Sugnet, CW, Haussler, D, and Kent, WJ
(2004). The UCSC Table Browser data retrieval tool. *Nucleic Acids Res* 32**,** D493-496. doi:
10.1093/nar/gkh103.

Kent, WJ, Sugnet, CW, Furey, TS, Roskin, KM, Pringle, TH, Zahler, AM, and Haussler, D (2002).
The human genome browser at UCSC. *Genome Res* 12**,** 996-1006. doi: 10.1101/gr.229102.

Killick, KE, Browne, JA, Park, SD, Magee, DA, Martin, I, Meade, KG, Gordon, SV, Gormley, E,
O'Farrelly, C, Hokamp, K, and MacHugh, DE (2011). Genome-wide transcriptional
profiling of peripheral blood leukocytes from cattle infected with *Mycobacterium bovis*
reveals suppression of host immune genes. *BMC Genomics* 12**,** 611. doi: 10.1186/1471-
2164-12-611.

Ko, ER, Yang, WE, McClain, MT, Woods, CW, Ginsburg, GS, and Tsalik, EL (2015). What was
old is new again: using the host response to diagnose infectious disease. *Expert Rev Mol*
Diagn 15**,** 1143-1158. doi: 10.1586/14737159.2015.1059278.

Kolli, V, Upadhyay, RC, and Singh, D (2014). Peripheral blood leukocytes transcriptomic signature
highlights the altered metabolic pathways by heat stress in zebu cattle. *Res Vet Sci* 96**,** 102-
110. doi: 10.1016/j.rvsc.2013.11.019.

Krjutskov, K, Koel, M, Roost, AM, Katayama, S, Einarsdottir, E, Jouhilahti, EM, Soderhall, C,
Jaakma, U, Plaas, M, Vesterlund, L, Lohi, H, Salumets, A, and Kere, J (2016). Globin
mRNA reduction for whole-blood transcriptome sequencing. *Sci Rep* 6**,** 31584. doi:
10.1038/srep31584.

Lawrence, M, Huber, W, Pages, H, Aboyoun, P, Carlson, M, Gentleman, R, Morgan, MT, and
Carey, VJ (2013). Software for computing and annotating genomic ranges. *PLoS Comput*
Biol 9**,** e1003118. doi: 10.1371/journal.pcbi.1003118.

Liu, J, Walter, E, Stenger, D, and Thach, D (2006). Effects of globin mRNA reduction methods on
gene expression profiles from whole blood. *J Mol Diagn* 8**,** 551-558. doi:
10.2353/jmoldx.2006.060021.

Mastrokolias, A, den Dunnen, JT, van Ommen, GB, t Hoen, PA, and van Roon-Mom, WM (2012).
Increased sensitivity of next generation sequencing-based expression profiling after globin
reduction in human blood RNA. *BMC Genomics* 13**,** 28. doi: 10.1186/1471-2164-13-28.

McLoughlin, KE, Nalpas, NC, Rue-Albrecht, K, Browne, JA, Magee, DA, Killick, KE, Park, SD,
Hokamp, K, Meade, KG, O'Farrelly, C, Gormley, E, Gordon, SV, and MacHugh, DE
(2014). RNA-seq transcriptional profiling of peripheral blood leukocytes from cattle
infected with *Mycobacterium bovis*. *Front Immunol* 5**,** 396. doi:
10.3389/fimmu.2014.00396.

McNamara, A, Arino de la Rubia, E, Zhu, H, Lowndes, J, Ellis, S, Waring, E, Quinn, M, McLeod,
H, Wickham, H, and Müller, K. 2017. skimr: compact and flexible summaries of data.
Available: https://github.com/ropenscilabs/skimr.

Meade, KG, Gormley, E, Doyle, MB, Fitzsimons, T, O'Farrelly, C, Costello, E, Keane, J, Zhao, Y,
and MacHugh, DE (2007). Innate gene repression associated with *Mycobacterium bovis*
infection in cattle: toward a gene signature of disease. *BMC Genomics* 8**,** 400. doi:
10.1186/1471-2164-8-400.

Mejias, A, and Ramilo, O (2014). Transcriptional profiling in infectious diseases: ready for prime
time? *J Infect* 68 Suppl 1**,** S94-99. doi: 10.1016/j.jinf.2013.09.018.

Morey, JS, Neely, MG, Lunardi, D, Anderson, PE, Schwacke, LH, Campbell, M, and Van Dolah,
FM (2016). RNA-seq analysis of seasonal and individual variation in blood transcriptomes
of healthy managed bottlenose dolphins. *BMC Genomics* 17**,** 720. doi: 10.1186/s12864-016-
3020-8.

Nabarro, D, and Wannous, C (2014). The potential contribution of Iivestock to food and nutrition
security: the application of the One Health approach in livestock policy and practice. *Rev Sci*
Tech 33**,** 475-485. doi: 10.20506/rst.33.2.2292.

O'Leary, NA, Wright, MW, Brister, JR, Ciufo, S, Haddad, D, McVeigh, R, Rajput, B, Robbertse, B,
Smith-White, B, Ako-Adjei, D, Astashyn, A, Badretdin, A, Bao, Y, Blinkova, O, Brover, V,
Chetvernin, V, Choi, J, Cox, E, Ermolaeva, O, Farrell, CM, Goldfarb, T, Gupta, T, Haft, D,
Hatcher, E, Hlavina, W, Joardar, VS, Kodali, VK, Li, W, Maglott, D, Masterson, P,
McGarvey, KM, Murphy, MR, O'Neill, K, Pujar, S, Rangwala, SH, Rausch, D, Riddick, LD,
Schoch, C, Shkeda, A, Storz, SS, Sun, H, Thibaud-Nissen, F, Tolstoy, I, Tully, RE, Vatsan,
AR, Wallin, C, Webb, D, Wu, W, Landrum, MJ, Kimchi, A, Tatusova, T, DiCuccio, M,

Kitts, P, Murphy, TD, and Pruitt, KD (2016). Reference sequence (RefSeq) database at

583   NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res*

584   44**,** D733-745. doi: 10.1093/nar/gkv1189.

585   O'Loughlin, A, Lynn, DJ, McGee, M, Doyle, S, McCabe, M, and Earley, B (2012). Transcriptomic

586   analysis of the stress response to weaning at housing in bovine leukocytes using RNA-seq

587   technology. *BMC Genomics* 13**,** 250. doi: 10.1186/1471-2164-13-250.

588   Pagès, H, Carlson, M, Falcon, S, and Li, N. 2017. AnnotationDbi: Annotation Database Interface. R

589   package version 1.38.0. Available: https://doi.org/doi:10.18129/B9.bioc.AnnotationDbi.

590   Patro, R, Duggal, G, Love, MI, Irizarry, RA, and Kingsford, C (2017). Salmon provides fast and

591   bias-aware quantification of transcript expression. *Nat Methods* 14**,** 417-419. doi:

592   10.1038/nmeth.4197.

593   Python Software Foundation. 2017. Python (version 3.5.2). Available:

594   https://www.python.org/downloads/release/python-352/.

595   R Core Team. 2017. R: A Language and Environment for Statistical Computing. Available:

596   https://www.R-project.org.

597   Ramilo, O, and Mejias, A (2009). Shifting the paradigm: host gene signatures for diagnosis of

598   infectious diseases. *Cell Host Microbe* 6**,** 199-200. doi: 10.1016/j.chom.2009.08.007.

599   Ropka-Molik, K, Stefaniuk-Szmukier, M, Zukowski, K, Piorkowska, K, Gurgul, A, and Bugno-

600   Poniewierska, M (2017). Transcriptome profiling of Arabian horse blood during training

601   regimens. *BMC Genet* 18**,** 31. doi: 10.1186/s12863-017-0499-1.

602   RStudio Team. 2015. RStudio: Integrated Development for R. Available: http://www.rstudio.com.

603   Schwochow, D, Serieys, LE, Wayne, RK, and Thalmann, O (2012). Efficient recovery of whole

604   blood RNA--a comparison of commercial RNA extraction protocols for high-throughput

605   applications in wildlife species. *BMC Biotechnol* 12**,** 33. doi: 10.1186/1472-6750-12-33.

606   Shen, J, Zhou, C, Zhu, S, Shi, W, Hu, M, Fu, X, Wang, C, Wang, Y, Zhang, Q, and Yu, Y (2014).

607   Comparative transcriptome analysis reveals early pregnancy-specific genes expressed in

608   peripheral blood of pregnant sows. *PLoS ONE* 9**,** e114036. doi:

609   10.1371/journal.pone.0114036.

610   Shin, H, Shannon, CP, Fishbane, N, Ruan, J, Zhou, M, Balshaw, R, Wilson-McManus, JE, Ng, RT,

611   McManus, BM, and Tebbutt, SJ (2014). Variation in RNA-seq transcriptome profiles of

612   peripheral whole blood from healthy individuals with and without globin depletion. *PLoS*

613   *ONE* 9**,** e91041. doi: 10.1371/journal.pone.0091041.

Soneson, C, Love, MI, and Robinson, MD (2015). Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Res* 4**,** 1521. doi:
10.12688/f1000research.7563.2.

Song, KD, Dowd, SE, Lee, HK, and Kim, SW (2013). Long-term dietary supplementation of
organic selenium modulates gene expression profiles in leukocytes of adult pigs. *Anim Sci J*
84**,** 238-246. doi: 10.1111/j.1740-0929.2012.01060.x.

Tablin, F, and Weiss, L (1985). Equine bone marrow: a quantitative analysis of erythroid
maturation. *Anat Rec* 213**,** 202-206. doi: 10.1002/ar.1092130212.

Takahashi, J, Waki, S, Matsumoto, R, Odake, J, Miyaji, T, Tottori, J, Iwanaga, T, and Iwahashi, H
(2012). Oligonucleotide microarray analysis of dietary-induced hyperlipidemia gene
expression profiles in miniature pigs. *PLoS ONE* 7**,** e37581. doi:
10.1371/journal.pone.0037581.

Thornton, PK (2010). Livestock production: recent trends, future prospects. *Philos Trans R Soc*
Lond B Biol Sci 365**,** 2853-2867. doi: 10.1098/rstb.2010.0134.

Trapnell, C, Roberts, A, Goff, L, Pertea, G, Kim, D, Kelley, DR, Pimentel, H, Salzberg, SL, Rinn,
JL, and Pachter, L (2012). Differential gene and transcript expression analysis of RNA-seq
experiments with TopHat and Cufflinks. *Nat Protoc* 7**,** 562-578. doi:
10.1038/nprot.2012.016.

Tyner, C, Barber, GP, Casper, J, Clawson, H, Diekhans, M, Eisenhart, C, Fischer, CM, Gibson, D,
Gonzalez, JN, Guruvadoo, L, Haeussler, M, Heitner, S, Hinrichs, AS, Karolchik, D, Lee,
BT, Lee, CM, Nejad, P, Raney, BJ, Rosenbloom, KR, Speir, ML, Villarreal, C, Vivian, J,
Zweig, AS, Haussler, D, Kuhn, RM, and Kent, WJ (2017). The UCSC Genome Browser
database: 2017 update. *Nucleic Acids Res* 45**,** D626-d634. doi: 10.1093/nar/gkw1134.

Weiner, J, and Kaufmann, SH (2017). High-throughput and computational approaches for
diagnostic and prognostic host tuberculosis biomarkers. *Int J Infect Dis* 56**,** 258-262. doi:
10.1016/j.ijid.2016.10.017.

Wickham, H. 2017a. reshape2: flexibly reshape data: a reboot of the reshape package. Available:
https://github.com/hadley/reshape.

Wickham, H. 2017b. tidyverse. Available: http://tidyverse.tidyverse.org.

Wickham, H, and Chang, W. 2017. ggplot2: create elegant data visualisations using the grammar of
graphics. Available: http://ggplot2.tidyverse.org.

Wickham, H, Francois, R, Henry, L, and Müller, K. 2017. dplyr: a grammar of data manipulation.
Available: http://dplyr.tidyverse.org.

Wickham, H, and Henry, L. 2017. tidyr: easily tidy data with 'spread()' and 'gather()' functions.

Available: http://tidyr.tidyverse.org.

Wilke, CO. 2017a. cowplot: streamlined plot theme and plot annotations for 'ggplot2'. Available:

https://github.com/wilkelab/cowplot.

Wilke, CO. 2017b. ggjoy: joyplots in 'ggplot2'. Available: https://github.com/clauswilke/ggjoy.

Winn, ME, Zapala, MA, Hovatta, I, Risbrough, VB, Lillie, E, and Schork, NJ (2010). The effects of

globin on microarray-based gene expression analysis of mouse blood. *Mamm Genome* 21**,**

268-275. doi: 10.1007/s00335-010-9261-y.

Wu, K, Miyada, G, Martin, J, and Finkelstein, D. 2003. Technical note: globin reduction protocol: a

method for processing whole blood RNA samples for improved array results. Available:

https://tools.thermofisher.com/content/sfs/brochures/blood2_technote.pdf.

658
659

11. Figures

FIGURE 1 | Schematic of the bioinformatics workflow for RNA-seq data acquisition, quality

control, analysis and interpretation.

FIGURE 2 | Ridge plots showing density of sample gene-level transcripts per million (TPM). (A)

Primary analysis (NCBI RefSeq: human, porcine, and cattle; UCSC: equine) results from

undepleted (purple) or globin-depleted (green) treatments. (B) Secondary analysis [NCBI RefSeq:

equine] from undepleted (purple) libraries.

FIGURE 3 | Average proportions of haemoglobin genes to total expressed genes from the primary

analysis of peripheral blood RNA-seq data in humans, pigs, horses and cattle.

FIGURE 4 | Distribution of haemoglobin gene-level transcripts per million (TPM). (A) Primary

analysis [NCBI RefSeq: human, porcine, and cattle; UCSC: equine] results from undepleted

(purple) or globin-depleted (green) treatments. (B) Secondary analysis [NCBI RefSeq: equine] from

undepleted (purple) libraries.

12. Tables

Table 1: Status of human, porcine, equine and bovine haemoglobin gene annotations in the
Ensembl, NCBI RefSeq, and UCSC database releases used.

Homo sapiens			
	Ensembl	**NCBI RefSeq**	**UCSC Table Browser**
Annotation release	Human release 90 (August 2017)[1]	NCBI *Homo sapiens* Annotation Release 108 (June 2016)[2, *]	hg38.refGene annotation track (last updated on August 28th 2017)[3, 4]
HBA1	Annotated with gene ID ENSG00000206172	Annotated with Entrez Gene ID 3039	Annotated with Entrez Gene ID 3039
HBA2	Annotated with gene ID ENSG00000188536	Annotated with Entrez Gene ID 3040	Annotated with Entrez Gene ID 3040
HBB	Annotated with gene ID ENSG00000244734	Annotated with Entrez Gene ID 3043	Annotated with Entrez Gene ID 3043
Sus scrofa			
	Ensembl	**NCBI RefSeq**	**UCSC Table Browser**
Annotation release	Pig release 90 (August 2017)[1]	NCBI *Sus scrofa* Annotation Release 106 (May 2017)[2, *]	susScr3.refGene annotation track (last updated on August 16th 2017)[2, 4, 5]
LOC110259958 (*HBA*)	Imported from RefSeq with Ensembl version ID 110259958.1	Annotated with Entrez Gene ID 110259958	Absent
LOC100737768 (*HBA*)	Annotated with gene ID ENSSSCG00000007978	Annotated with Entrez Gene ID 100737768	Absent
HBB	Annotated with gene ID ENSSSCG00000014725	Annotated with Entrez Gene ID 407066	Annotated with Entrez Gene ID 407066
Equus caballus			
	Ensembl	**NCBI RefSeq**	**UCSC Table Browser**

Annotation release	Horse release 90 (August 2017)[1]	NCBI *Equus caballus* Annotation Release 102 (November 2015)[2, 6, ¶]	equCab2.refGene annotation track (last updated on August 10th 2017)[2, 4, 5, *]
HBA (also known as *HBA1*)	Absent	Absent from current annotation release, but accessible via online search with Entrez Gene ID 100036557	Annotated with Entrez Gene ID 100036557
HBA2	Absent	Not annotated on current reference assembly, but accessible via online search with Entrez Gene ID 100036558	Annotated with Entrez Gene ID 100036558
HBB	Annotated with gene ID ENSECAG00000010020	Annotated with Entrez Gene ID 100054109	Annotated with Entrez Gene ID 100054109
Bos taurus			
	Ensembl	**NCBI RefSeq**	**UCSC Table Browser**
Annotation release	Cow release 90 (August 2017)[1]	NCBI *Bos taurus* Annotation Release 105 (January 2016)[2, *]	bosTau8.refGene annotation track (last updated on August 20th 2017)[2, 4, 5]
HBA1	Annotated with gene ID ENSBTAG00000026417	Annotated with Entrez gene ID 100140149	Absent
HBA (also known as *HBA2*)	Annotated with gene ID ENSBTAG00000026418	Annotated with Entrez Gene ID 512439	Annotated with Entrez Gene ID 512439
HBB	Annotated with gene ID ENSBTAG00000038748	Annotated with Entrez Gene ID 280813	Annotated with Entrez Gene ID 280813

677
678 [1] (Aken et al., 2017); [2] (O'Leary et al., 2016); [3] (Kent et al., 2002); [4] (Karolchik et al., 2004); [5] (Tyner et al., 2017); [6]
679 (Hestand et al., 2015).
680
681 [*]Source of reference transcriptome in the primary analysis; [¶]Source of reference transcriptome in the secondary
682 analysis.

683   **Table 2:** Summary of RNA-seq filtering/trimming and mapping statistics.

Species	Treatment	RNA-seq library type	Sequencing mode	Mean no. of reads (SE) or pairs (PE)	Mean no. of reads (SE) or pairs (PE) removed	Mean proportion of reads (SE) or pairs (PE) removed	Mean no. of observed fragments*	Mean no. of mapped fragments*	Average mapping rate	Reference source
					Primary analysis					
Homo sapiens	Undepleted	Inward unstranded	PE	40,218,886	8,203,217	20.4%	32,015,669	25,583,548	80.9%	NCBI RefSeq
Homo sapiens	Globin depleted	Inward unstranded	PE	36,874,759	10,704,088	29.0%	26,170,671	20,362,333	75.8%	NCBI RefSeq
Sus scrofa	Undepleted	Inward unstranded	PE	39,036,515	4,613,991	11.8%	28,685,437	25,129,140	87.7%	NCBI RefSeq
Sus scrofa	Globin depleted	Inward unstranded	PE	31,339,886	3,899,427	12.4%	22,867,049	19,959,995	87.3%	NCBI RefSeq
Equus caballus	Undepleted	Unstranded	SE	24,271,141	38,892	0.2%	14,850,797	1,705,235	11.5%	UCSC refGene
Bos taurus	Undepleted	Inward stranded forward	PE	20,495,983	3,474,597	17.0%	17,021,386	12,039,012	70.7%	NCBI RefSeq
					Secondary analysis					
Equus caballus	Undepleted	Unstranded	SE	24,271,141	38,892	0.2%	14,850,797	10,618,507	71.4%	NCBI RefSeq

684
685   *The Salmon tool categorises fragments as one read (for SE RNA-seq libraries) or a read pair (for PE RNA-seq libraries).

686 **Table 3:** Summary statistics for haemoglobin gene-level transcripts per million (TPM).

Species	Gene symbol	Treatment	Maximum TPM	Minimum TPM	No. of samples	Mean TPM
Primary analysis						
Homo sapiens	*HBA1*	Undepleted	211,012	163,459	12	182,686
Homo sapiens	*HBA1*	Globin depleted	122,509	29,972	12	66,109
Homo sapiens	*HBA2*	Undepleted	334,003	237,796	12	286,052
Homo sapiens	*HBA2*	Globin depleted	162,301	41,680	12	79,464
Homo sapiens	*HBB*	Undepleted	292,039	113,654	12	204,718
Homo sapiens	*HBB*	Globin depleted	34,622	10,933	12	19,789
Sus scrofa	*LOC110259958* (*HBA*)	Undepleted	135	44	12	86
Sus scrofa	*LOC110259958* (*HBA*)	Globin depleted	39	2	12	13
Sus scrofa	*LOC100737768* (*HBA*)	Undepleted	274,472	185,949	12	243,864
Sus scrofa	*LOC100737768* (*HBA*)	Globin depleted	232,884	12,183	12	84,021
Sus scrofa	*HBB*	Undepleted	540,083	369,556	12	476,284
Sus scrofa	*HBB*	Globin depleted	367,310	18,269	12	136,172
Equus caballus	*HBA1*	Undepleted	8,767	32	37	1,600
Equus caballus	*HBA2*	Undepleted	12,382	80	37	2,364
Equus caballus	*HBB*	Undepleted	15,778	63	37	3,786
Bos taurus	*HBA1*	Undepleted	81	7	6	29
Bos taurus	*HBA2*	Undepleted	3,191	80	10	1,070
Bos taurus	*HBB*	Undepleted	1,599	109	10	526
Secondary analysis						
Equus caballus	*HBB*	Undepleted	4,594	17	37	1,060

688 Figure 1

695
696

697 Figure 2

699   Figure 3

705
706

707 Figure 4