Deciphering human ribonucleoprotein regulatory networks

Neelanjan Mukherjee[1], Hans-Hermann Wessels[2], Svetlana Lebedeva[2], Marcin Sajek[3,4], Mahsa

Ghanbari[2], Aitor Garzia[4], Alina Munteanu[2], Thalia Farazi[4*], Jessica I Hoell[4,5], Kemal Akat[4],

Thomas Tuschl[4], Uwe Ohler[2]

[1]Department of Biochemistry and Molecular Genetics, University of Colorado School of
Medicine, Aurora, Colorado 80045, USA.

[2]Berlin Institute for Medical Systems Biology, Max Delbrück Center for Molecular Medicine,
Berlin, Germany.

[3]Institute of Human Genetics, Polish Academy of Sciences, Poznan, Poland

[4]Howard Hughes Medical Institute and Laboratory for RNA Molecular Biology, The Rockefeller
University, 1230 York Ave, Box 186, New York, NY 10065

[5]Department of Pediatric Oncology, Hematology and Clinical Immunology, Center for Child and
Adolescent Health, Medical Faculty, Heinrich Heine University of Dusseldorf, Dusseldorf,
Germany

[*]Current address, Juno Therapeutics, 400 Dexter Avenue North, Seattle, WA 98109

Correspondence to: ttuschl@rockefeller.edu, uwe.ohler@mdc-berlin.de,
neelanjan.mukherjee@ucdenver.edu

RNA-binding proteins (RBPs) control and coordinate each stage in the life cycle of RNAs.

Although *in vivo* binding sites of RBPs can now be determined genome-wide, most studies

typically focused on individual RBPs. Here, we examined a large compendium of 114 high-

quality transcriptome-wide *in vivo* RBP-RNA cross-linking interaction datasets generated by the

same protocol in the same cell line and representing 64 distinct RBPs. Comparative analysis of

categories of target RNA binding preference, sequence preference, and transcript region

specificity was performed, and identified potential posttranscriptional regulatory modules, i.e.

specific combinations of RBPs that bind to specific sets of RNAs and targeted regions. These

regulatory modules encoded functionally related proteins and exhibited distinct differences in

RNA metabolism, expression variance, as well as subcellular localization. This integrative

investigation of experimental RBP-RNA interaction evidence and RBP regulatory function in a

human cell line will be a valuable resource for understanding the complexity of post-

transcriptional regulation.

Introduction

Of the 20,345 annotated protein-coding genes in human, at least 1,542 are RNA-binding proteins (RBPs) (Gerstberger et al., 2014). RBPs interact with RNA regulatory elements within RNA targets to control splicing, nuclear export, localization, stability, and translation (Moore, 2005). RBPs have specificity to bind one or multiple RNA categories, including messenger RNA (mRNA) and diverse categories of non-coding RNA such as ribosomal RNA (rRNA), transfer RNA (tRNA), small nuclear and nucleolar RNA (snRNA/snoRNA), microRNA (miRNA), and long non-coding RNA (lncRNA). Mutations in RBPs or RNA regulatory elements can result in defects in RNA metabolism that cause human disease (Cooper et al., 2009; Fredericks et al., 2015).

A standard technique for *in vivo* global identification of RBP-RNA interaction sites consists of immunoprecipitating the ribonucleoprotein (RNP) complex, isolating the bound RNA, and quantifying the RNA targets by microarrays or deep sequencing (Tenenbaum et al., 2000; Zhao et al., 2010). The introduction of cross-linking prior to immunoprecipitation (CLIP) as well as RNase digestion enabled the biochemical mapping of individual interaction sites (Ule et al., 2003). Subsequent modifications to CLIP increased the resolution of the interaction sites (Hafner et al., 2010; König et al., 2010). One of these methods, photoactivatable ribonucleoside-enhanced cross-linking and immunoprecipitation (PAR-CLIP), utilizes 4-thiouridine or 6-thioguanosine combined with 365 nm UV crosslinking to produce single-nucleotide RBP-RNA interaction evidence that is utilized to define binding sites (Corcoran et al., 2011; Garzia et al., 2017b; Hafner et al., 2010).

Experimentally-derived RBP binding sites provide valuable functional insights. First, they can

reveal the rules for regulatory site recognition by the RBP, whether due to sequence and/or

structural characteristics. Second, the region and position of the interaction sites of an RBP

within transcripts provides insights into its role in RNA metabolism and its subcellular

localization. For example, if most of the mapped interaction sites are intronic and adjacent to

splice sites, the RBP is highly likely to be a nuclear splicing factor rather than a cytoplasmic

translation factor. Finally, these data reveal the target transcripts and therefore the potential

biological role for the RBP.

Throughout the life of an RNA, interactions with many different RBPs determine the ultimate

fate of the transcript. Even though profiling of the interaction sites of a single RBP is clearly

powerful, it does not provide information on other RBPs potentially targeting the same RNA or

on other regulatory elements within the RNA. Small comparative efforts focusing on the

regulation of splicing, 3' end processing, RNA stability by AU-rich elements, and miRNA-

mediated silencing have demonstrated the value of integrating interaction sites from multiple

RBPs (Martin et al., 2012; Mukherjee et al., 2014; Pandit et al., 2013; Zhang et al., 2010).

Therefore, a large-scale comparative examination of interaction sites for many RBPs will yield

valuable knowledge regarding the architecture and determinants of RNA regulatory networks.

At least 173 PAR-CLIP experiments have been performed in HEK293 cells to date, laying the

groundwork for a large-scale integrative analysis and complementing efforts of ENCODE, which

focused on other cell types and utilized other CLIP protocols (Van Nostrand et al., 2016). We

describe a concerted effort to identify and uniformly process all high-quality PAR-CLIP data sets

by evaluating the characteristic T-to-C transitions induced by photocrosslinking. Using the

resulting compendium of high-quality in vivo RBP interaction maps from the same cell line

enabled us to determine the relationship between RBPs with respect to their preferred category of

target RNA and any underlying sequence specificity. We uncovered regulatory modules reflected

by combinatorial binding events, and assessed their role and functional implications on RNA

metabolism. Finally, our results support the role of RBPs in buffering gene expression variance.

Results

A high-quality map of in vivo RBP-RNA interactions across 64 proteins

In order to generate a comprehensive quantitative resource of RBP-RNA interactions within a

human cell line, we identified 166 published PAR-CLIP data sets performed predominantly in

HEK293 cells, and added 7 new libraries generated in our laboratories (Sup Table 1). Typically,

these datasets were generated using transgenic HEK293 cell lines in which each individual RBP

was FLAG-tagged and recombined into the same chromosomal locus containing a strong

promoter. In this way, the expression of each RBP as well as the strength of its

immunoprecipitation were generally comparable. Furthermore, the availability of orthogonal

transcriptome-wide datasets quantifying individual steps of RNA metabolism made HEK293

cells ideal for examining the functional characteristics of RNA targets (Mukherjee et al., 2017).

Each of the 173 PAR-CLIP libraries generated in HEK293 were subject to a stringent analysis

strategy to retain high-quality datasets (Supplemental Table 1). First, each library was analyzed

using the PAR-CLIP Suite v1.0 (https://rnaworld.rockefeller.edu/PARCLIP_suite) (Garzia et al.,

2017b) to discriminate significant target RNA categories from non-crosslinked background RNA

categories populated by fragments of abundant cellular RNAs (see Methods, Supplemental Fig.

1A). Next, we defined binding sites based on the local density of T-to-C transitions using

PARpipe (https://github.com/ohlerlab/PARpipe) (Corcoran et al., 2011) and only retained those

libraries with sufficiently high read counts and T-to-C transition specificity compared to a deeply

sequenced background reference library (Supplemental Fig 1b) (Friedersdorf and Keene, 2014).

Since the immunoprecipitation step was omitted in this reference library it served as an effective

comparison point to score read count and T-to-C transition for all RBPs. Finally, for RBPs with

more than 3 libraries available, outlier libraries exhibiting poor correlation of 6-mer frequencies

were excluded (Supplemental Fig 1d, e). This resulted in 114 libraries corresponding to 64 RBPs

that were the basis for downstream analysis. There were eight RBP families represented by two

or more RBPs.

Grouping RBPs by annotation category and positional binding site preferences

As first step to describe RBP-RNA regulatory networks, we determined the relative binding

preference of each RBP for specific target RNA annotation categories (Supplemental Table 2).

For each library, we calculated an RNA annotation category preference value, defined as the

difference in the fraction of T-to-C reads per annotation category between each RBP library and

the reference library. We performed hierarchical clustering of RBPs by annotation category

preference, using Ward's method and Euclidean distances. This yielded eight clusters of binding

preference (Figure 1a – orange line demarcates cluster definitions) with varying enrichment or

depletion for individual or combinations of specific annotation categories. For each of these

clusters, we compiled a detailed table summarizing the reported functions for each of the RBPs

(Table 1). Taken together, clustering by RNA annotation category separated RBPs into groups

according to their known subcellular localization and functions.

Three of the eight clusters (clusters 2, 4, and 5) contained nine RBPs that exhibited preference

for categories of non-coding RNA (rRNA, snRNA, snoRNA, and tRNA), but not mRNA,

precursor mRNA (pre-mRNA), or lncRNA. The remaining five clusters contained 55 RBPs

exhibiting preference for binding to mRNA, pre-mRNA and long-noncoding RNA (lncRNA)

annotation categories. The RBPs in clusters 1, 6, 7, and 8 exhibited strong preferences for

various mRNA annotation categories. The RBPs in cluster 3 did not exhibiting strong preference

for specific mRNA annotation categories. Additionally, for each of the RBPs in the cluster, we

performed a positional meta-analysis of binding sites with respect to major transcript landmarks

within target mRNAs. Many of the RBPs also showed strong preferences for binding to specific

positions within mRNAs relating to their role in specific steps of mRNA processing (Table 1).

We hypothesized that target annotation category preferences and positional binding preferences

should reflect subcellular localization of the RBP and its role(s) in mRNA processing. Cluster 6

contained twelve RBPs and exhibited strong preference for intronic regions and to a lesser

degree 3' UTRs of mRNAs and lncRNAs. The intronic preference was consistent with the

predominantly nuclear localization of these RBPs and the pre-mRNA splicing process. ELAVL1,

which is the sole member of the ELAVL1 family of RBPs that is predominantly localized in the

nucleus but capable of shuttling to the cytoplasm, exhibited positional binding flanking the end

of the 3' UTR and for 5' and 3' splice sites. Cluster 8 contained fourteen RBPs and exhibited

distinct preference for 3' UTR regions. This included the unpublished and predominantly

cytoplasmic ELAVL1 family members, ELAVL2, ELAVL3, and ELAVL4, which exhibited a

strong positional preference for binding in the distal region of the 3' UTR and acting

predominantly on mature mRNA (Mansfield and Keene, 2012). In summary, the annotation

category preferences and positional binding preferences implicated the specific steps of mRNA

processing the RBPs potentially regulate.

The spectrum of RNA sequence specificity

RBPs exist on a spectrum of specificity depending on a variety of primary and secondary

structure features (Jankowsky and Harris, 2015). Here, our goal was to identify the RBPs with

substantial primary sequence specificity and then examine their sequence preference. For each of

the 55 RBPs, we counted all possible 6-mers using Jellyfish (Marçais and Kingsford, 2011) for

the reads contributing to PARalyzer-defined binding sites. We observed 6-mer frequencies

ranging as high as 512-fold to as low as 5-fold over a uniform distribution of 6-mers

(Supplemental figure 2a). In contrast, our reference background library exhibited 16-fold

enrichment of at least one 6-mer compared to uniform. AGO1-4 libraries were excluded from 6-

mer analysis due to the overwhelming sequence contribution from crosslinked miRNAs. Twenty-

seven RBPs did not have a single 6-mer found at higher frequency than present in the reference

sample. Amongst these RBPs established or expected to display low sequence-specificity were

the RNA helicase MOV10, the nuclear exosome component DIS3, and the EIF3 complex

translation initiation factors.

For each of the 24 RBPs with stronger sequence enrichment than the reference library, we

clustered the top 5 sequences enriched over the reference library (Figure 2). Our results

recapitulated the sequence preference for the RBPs in this group with well-characterized

sequence motifs (detailed in Table 2). The ELAVL1 family proteins, which bound to different

regions and positions of mRNA, showed similar preference for U- and AU-rich 6-mers, while

ZFP36 only enriched a subset of the AU-rich 6-mers (Mukherjee et al., 2014). Complementing

the 6-mer enrichment analysis, we performed motif analysis for each RBP library with the motif

finding algorithm SSMART (sequence-structure motif identification for RNA-binding

proteins, (Munteanu et al., 2018)) (Supplemental Fig 2b). For most RBPs, we observed strong

concordance between the two analyses. RBM20 was a clear exception, for which we observed

the established UCUU-containing motifs (Maatz et al., 2014) with SSMART, but a GA-rich

sequence in the 6-mer enrichment analysis. However, we do observe UCUU-containing motifs in

the top 15, but not top5 6-mers for RBM20. Altogether, our analysis was remarkably consistent

with previously reported motifs in spite of differences in data processing and analysis (detailed

Table 2).

Identification of RNA regulatory modules

To understand the functional impact of co-regulation by multiple RBPs, we analyzed the co-

variation in binding patterns of all 55 RBPs across 13,299 target RNA encoding genes to probe

for the existence of regulatory modules, i.e., specific subsets of RNAs implicated in similar

function bound by subsets of RBPs. To this end, we employed Factor Analysis (FA), which

reduces a large number of observed variables to a smaller number of latent *factors*. Here, our

observed variables represented the normalized RBP binding (see methods) for each of the 55

RBPs across all target RNA encoding genes (n=13,299). The latent *factors* represented similar

binding patterns to RNA targets by one or more of the 55 RBPs. RBPs exhibiting high loadings

for the same *factor* would have very similar binding patterns to RNA targets. Importantly in this

framework, a single RBP could be assigned to multiple *factors*, just as a single RBP can

participate in multiple RNPs and regulate different aspects of RNA metabolism.

The FA model decomposed the 55 x 13,299 normalized RBP binding matrix into a 55 x 10 factor

loading matrix (representing the strength of the dependence of each of the 55 RBP target RNA

binding pattern on each of the 10 *factors*), a 13,299 x 10 factor score coefficient matrix

(representing the dependence between the binding of the 13,299 target RNA encoding gene and

each of the 10 *factors*), and residual error (Supplemental Fig 3a and methods). Cumulatively, the

FA model explained ~60% of the variance in the observed data. The remaining unexplained

variance was expected due to the challenges of integrating data sets of varying depth and quality,

in spite of our efforts to control these aspects. The communality, which is the amount of variance

explained by the model for each RBP-binding variable, varied drastically for all 55 RBPs; the

model explained at least 80% of the variance in enrichment scores for 12 RBPs, and at least 50%

of the variance in enrichment scores for 30 RBPs (Supplemental Figure 3b). RBPs with lower

communality often coincided with shallow depth of their PAR-CLIP libraries.

The FA model also uncovered interesting parallels between the similarity in the binding of target

RNA encoding genes and the target annotation category preferences (from Figure 1a). We

observed that individual *factors* contained RBPs that preferred binding to either mature (Factors

1, 3, 4, 5, 8) or precursor transcripts (Factors 2, 6), reflecting involvement in different stages of

RNA metabolism (Figure 3a). Furthermore, individual *factors* contained RBPs exhibiting similar

patterns of binding to specific regions of the mRNA (i.e., intron, coding, 3' UTR). Indeed, RBPs

from the same family, or known to regulate a specific aspect of RNA processing, had high

loadings for the same *factors*. For example, the ELAVL1 family members were associated with

Factor 1; the AGO1 family were associated with Factor 3; the IGF2BP1 family were associated

with Factor 4; the FMR1 family had were associated with Factor 5 and Factor 8; LINE-1

encoded proteins were associated with Factor 7. One of the unanticipated associations was that

of HNRNPC with Factor 2, which contained mainly cleavage and polyadenylation factors.

Interestingly, HNRNPC was shown to interact with U-rich sequences downstream of a viral

poly-adenylation signal nearly three decades ago (Wilusz et al., 1988), and more recently, to

repress cleavage and poly-adenylation in humans (Gruber et al., 2016). These examples highlight

the specific testable hypotheses generated by an integrative analysis that are not necessarily

obvious when examining a single RBP in isolation.

By clustering the factor score coefficients, i.e. the specific linear combination of RBP binding for

that target RNA, we identified target RNA encoding genes constituting putative regulatory

modules associated with a given *factor*. Therefore, each regulatory module was associated with

an RBP component (the subset of RBPs exhibiting similar binding pattern) and a RNA

component (the subsets of target RNA encoding genes bound by those RBPs). These regulatory

modules did not imply physical interactions between RBPs; rather, it identified RBPs that may

cooperate in controlling RNA metabolism for specific subsets of RNA targets, possibly across

cellular compartments. Almost a quarter of the target RNA encoding genes (3,180/13,299) were

assigned to regulatory modules by exhibiting high factor score coefficients for a single *factor*

(Supplemental figure 3c). We did not identify target RNA encoding genes with high factor score

coefficients for Factor 9 or 10. The remaining target RNA encoding genes did not exhibit high

factor score coefficients for any specific *factor* in our analysis, suggesting that the targets were

either not bound by specific combinations of these RBPs, bound broadly by all RBPs, or not

bound by the subset of RBPs in the analysis. As such, we labeled this target RNA encoding gene

category as "non-specific". The RNA regulatory modules encoding genes were enriched for

different GO categories. Factor 1 RNA regulatory modules were enriched for 'AU-rich element

binding' and Factor 3 RNA regulatory modules were enriched for 'gene silencing by miRNA';

AU-rich RBPs and AGO proteins were strongly associated with Factor 1 and Factor 3,

respectively. This was consistent with the recurrent observation that RBPs target the mRNAs

encoding themselves (Pullmann et al., 2007; Tenenbaum et al., 2000). In turn, the RNAs

encoding "non-specific" genes contained ribosomal proteins and mitochondrial electron-

transport proteins.

RNA regulatory modules underlie distinct patterns of RNA metabolism

In order to test the functional relevance of these RNA regulatory modules, we reasoned that

perturbation (change of protein abundance or activity) of an RBP will lead to pronounced effects

only for the RNA regulatory modules assigned to the specific *factor(s)* that RBP is associated

with. We examined mature and precursor RNA expression changes induced by siRNA

knockdown of ELAVL1 (Kishore et al., 2011). ELAVL1 was strongly associated with both

Factor 1 and Factor 2, which exhibited RNA targeting patterns for mature or precursor RNAs,

respectively. Concordantly, Factor 1 associated RNA regulatory modules, but not Factor 2 RNA

regulatory modules, exhibited ELAVL1-dependent stabilization of mature RNA (Figure 4a).

Likewise, Factor 2 RNA regulatory modules exhibited a more pronounced ELAVL1-dependent

stabilization of precursor RNA than Factor 1 RNA regulatory modules (Figure 4b). Each human

ELAV1 family protein contains three RRM domains (>90% sequence identity), but the hinge

region between the second and third RRM of ELAVL1 contains a shuttling sequence responsible

for its nuclear localization (Fan and Steitz, 1998). Due to the lack of this shuttling sequence,

ELAVL2/3/4 are predominantly cytoplasmic and were strongly associated with Factor 1, but not

Factor 2. Taken together, the model was able to correctly identify and distinguish ELAVL1-

dependent stabilization of both precursor and mature RNA (Lebedeva et al., 2011; Mukherjee et

al., 2011).

We hypothesized that the subsets of RNAs assigned to the different regulatory module would

exhibit differences in RNA metabolism driven by the RBPs in the *factor* associated with the

regulatory module. Therefore, we compared six aspects of RNA metabolism previously

quantified in HEK293 cells (Mukherjee et al., 2017), for each of the RNA regulatory modules

associated with each of the *factors*. The *factor*-associated RNA regulatory modules exhibited

very distinct RNA metabolic profiles compared to each other and to non-specific category

(Figure 4c, Supplemental Figure 4a). Factor 2 RNA regulatory modules, which was the only

factor associated with RBPs binding to precursor mRNA and lncRNA, had low processing rates,

high degradation rates and their encoded RNAs were preferentially localized in the nucleus

versus the cytoplasm. Factor 2 RNA regulatory modules were strongly enriched for lncRNAs

(Figure 4d). Indeed, these genes strongly overlapped with a set of lncRNAs likely to be

functional (Supplemental figure 4b) (Mukherjee et al., 2017).

We also examined regulatory differences in RNA metabolism for genes associated with

cytoplasm-enriched factors. For example, factor 1 RNA regulatory modules were more stable

than Factor 3 RNA regulatory modules (Figure 4c). Factor 1 was strongly associated with

ELAVL1 family proteins, which stabilize target mRNAs. Factor 3 was strongly associated with

AGO1 family proteins, which execute miRNA-mediated degradation of target mRNAs.

Additionally, Factor 4 RNA regulatory modules, which are bound by IGF2BP1 family proteins,

were highly synthesized, processed, stabilized, and translated (Figure 4c). The RNA targets of

IGF2BP1 family RBPs were strongly localized to the ER (Supplemental Figure 4c) (Jønson et

al., 2007), which is also consistent with the proposed role of IGF2BP1 family proteins for RNA

localization and translation (Farina et al., 2003; Nielsen et al., 2001). Although correlative, these

results indicate that different RBP binding patterns beget different consequences for RNA

metabolism.

Specific RNA regulatory modules also exhibited preferential localization to processing bodies

(P-bodies), which are cytoplasmic granules associated with translational repression (Sheth and

Parker, 2003). Namely, Factor 3 RNA regulatory modules, which were strongly associated with

the AGO1 family, were the most strongly enriched for localizing to P-bodies according to a

recent study characterizing the transcriptome and proteome of P-bodies, and the AGO2 protein

itself was 90-fold enriched (Hubstenberger et al., 2017). Similarly, Factor 5 RNA regulatory

modules, which were strongly associated with the FMR1 family, were also enriched for

localizing in P-bodies, along with the FMR1 protein (16-fold enriched). In contrast, the non-

specific category was depleted from P-bodies.

Fine-tuning of gene expression has been postulated to be an important function of post-

transcriptional regulation by RBP and miRNAs. Therefore, we examined the cell-to-cell

variability in gene expression across 25 individual HEK293 cells with respect to the RNA

regulatory modules. The single-cell RNA-seq data was very deeply sequenced and generated

using the massively parallel single-cell RNA-sequencing (MARS-Seq) protocol (Guillaumet-

Adkins et al., 2017). Most RNA regulatory modules exhibited lower expression variability than

the non-specific category (Figure 4e). In particular, Factor 4 RNA regulatory modules exhibited

the lowest variation and highest median expression across the 25 cells (Supplemental Figure 4d).

These results supported the broad notion that post-transcriptional gene regulation generally

confers robustness and fine-tuning of gene expression.

Conclusion

Our study presents a curation of existing datasets, followed by systematic analysis of high-

quality and high-resolution RBP-RNA interaction data. We focused on the RBPs that

preferentially bound to mRNA and lncRNA and examined their sequence specificity and

sequence motif preferences. Our survey of the RBP regulatory landscape identified the most

prevalent subsets of RNAs targeted by a specific subset of RBPs, which we refer to as RNA

regulatory modules.

We utilized high quality PAR-CLIP datasets for which the immunoprecipitation was generally

comparable due to fact most RBPs were FLAG-tagged. Nevertheless, several caveats associated

with the interpretation of this analysis need to be pointed out. Despite several measures of quality

control to decide which datasets to include in our analysis, the libraries varied greatly in depth,

quality, digestion biases and potentially other confounding variables with respect to the protocol.

The FA model quantitatively assessed the degree to which we could explain the full complement

of RBP-RNA target binding patterns. These confounders undoubtedly contributed to the ~40%

of variance not explained by the FA model. In comparison, the ENCODE eCLIP datasets (Van

Nostrand et al., 2016) are likely to suffer from different confounders: they were generated using

one consistent experimental protocol but used antibodies against endogenous proteins expressed

at varying levels, and for which IP efficiency can vary greatly in spite of the quality control

performed (Sundararaman et al., 2016). Essentially, this represents the trade-offs in experimental

design between analyzing the endogenous protein compared to an epitope-tagged protein.

Modifying the genomic loci of the protein to engineer an endogenous epitope tagged RBP is

a very promising strategy.

Assuming the RBPs investigated here are a representative sample of the ~1,542 RBPs encoded in

the human genome, there may be an astounding number of RBPs with substantial primary

sequence specificity. However, the degree of sequence specificity is determined by the nature of

the RBP-RNA interaction, which can be quite extensive and specific, as in the case of Pumilio,

or minimal and non-sequence specific, as in the case of an RNA-helicase. An interesting

exception were the A-rich sequences enriched by UPF1, which is an RNA helicase and therefore

unlikely to exhibit strong sequence specificity. One possible explanation is that such sequences

may represent pre-mature polyA tail recognition involved in aspects of ribosome quality control

demonstrated in yeast (Koutmou et al., 2015) and human cells (Garzia et al., 2017a). Likewise,

more examples of unanticipated sequence enrichments may shed light on novel RNA regulatory

mechanisms.

Our FA model was able to identify distinct RBP-RNA target regulatory modules. At the very

minimum, 25% of target RNA encoding genes were assigned to RNA regulatory modules. This

is very likely an underestimation due to noisy data and a biased, far from complete sampling of

RBPs. However, there is likely to be a subset of genes for which post-transcriptional gene

regulation indeed plays a negligible role, at least in HEK293 cells. Furthermore, a small number

of RBPs in our analysis are not endogenously expressed in HEK293 and their natural expression

is tissue-specific and/or context-dependent. The approach presented here can scale to binding

data for all ~700 RBPs experimentally shown to be associated with poly-adenylated RNA in

HEK293 cells or even ~1,542 known RBPs (Baltz et al., 2012).

The RNA regulatory modules exhibited different patterns of RNA processing, degradation,

localization, and translation. We speculate that these differences in RNA metabolism were driven

by individual RBPs or the combination of RBPs associated with that regulatory module. This

was supported by the response of specific RNA regulatory modules to ELAVL1 knockdown

(Figure 4A, B). Additionally, the RNA regulatory modules encoded functionally related proteins

and similarly localized proteins. The enrichments were for proteins with similar molecular

functions or multi-component complexes rather than signaling pathways (Supplemental Fig 3b).

Altogether, these lines of evidence provide support for the coordinate regulation of 'functionally

coherent' RNA regulatory modules as proposed by the post-transcriptional operon/regulon model

(Keene, 2007). The ultimate test of this model would involve manipulating specific combinations

of binding sites and RBPs. Our study provides the rationale for such experiments, which

unfortunately remain technically challenging.

Our observations have important implications for RBP-RNA regulatory networks and their

importance in gene expression. The mRNA targets within specific regulatory modules encoded

the RBP themselves, a generalization of a commonly made observation that RBPs bind to the

mRNAs encoding them (Mesarovic et al., 2004). Our analysis lends support for this frequently

observed potential auto-regulatory feedback. These feedback loops may in fact buffer the

expression range of the targeted mRNAs, including those of the RBP. In this context, the

observation that the RNA regulatory modules exhibited lower cell-to-cell gene expression

variance, provides more evidence for the importance of post-transcriptional regulation in

buffering transcriptional noise (Bahar Halpern et al., 2015; Battich et al., 2015). Systematic

perturbation of individual and combinations of RBPs will be quite powerful in revealing

fundamental properties of RNA regulatory networks such as auto-regulatory feedback and

buffering.

The binding preference and targets of the vast majority of human RBPs remains unknown. The

insights gained from this study demonstrate the value of large-scale efforts by ENCODE and

others in the community to globally identify RBP binding sites. Of the 64 RBPs in this study, 44

were not represented in the ENCODE cell lines. Cumulatively these efforts interrogate ~10% of

human RBPs with known RNA-binding domains. Thus, these two large scale efforts offer the

potential to complement one another in our continuing attempts to understanding RBP-RNA

regulatory networks, for which we have only glimpsed the tip of the iceberg.

Methods

Processing, filtering, and quality control of PAR-CLIP libraries

Each PAR-CLIP library was subject to two rounds of quality control. First, all PAR-CLIP

libraries generated in HEK293 cells were subject to the quality control pipeline PAR-CLIP Suite

v1.0 (https://rnaworld.rockefeller.edu/PARCLIP_suite/). Using raw Illumina sequencing data,

this pipeline identified the predominant target RNA category or categories for each RBP and

provided the T-to-C conversion frequency resolved by read length and RNA category

(Supplemental Fig 1). The mapped reads of each RNA category were resolved by error distance

0 (d0), error distance 1 (d1; split in T-to-C and d1 other than T-to-C), and error distance 2 (d2).

This process discriminated for each library true target RNA categories from non-crosslinked

background RNA categories populated by fragments of abundant cellular RNAs. In order to

disqualify experiments comprising too many non-crosslinked RBP-specifically bound RNAs or

co-purified non-crosslinked background RNAs, we pursued only datasets which collect at least

10,000 redundant d1 reads $\geq$ 20 nt in at least one of major RNA annotation categories with d1(T-

to-C)/(d0 + d1) $\geq$ 30%, and d1(T-to-C)/(d1-total) $\geq$ 65%.

For the libraries passing the first threshold, we defined and annotated binding sites using

PARpipe, which is a pipeline wrapper for PARalyzer (Corcoran et al., 2011; Mukherjee et al.,

2014). The threshold for additional filtering were determined by comparisons with the reference

library (Friedersdorf and Keene, 2014). This reference library was generated using a modified

PAR-CLIP protocol in which there was no immunoprecipitation and the addition of an rRNA

depletion step after proteinase K digestion, followed by a partial digestion using RNase T1. We

required libraries had to have an average fraction T-to-C over remaining reads greater than 0.32

(the average fraction T-to-C over remaining reads greater of the reference library), an average

conversion specificity greater than 0, more than 20000 aligned reads, not be digested only with

micrococcal nuclease, a redundant read copy fraction less than .98 (Supplemental Fig 1b,c and

Sup Table 1). For RBPs with three or more libraries, we removed outlier based on correlation of

6-mer frequency calculated from PARalyzer-utilized reads.

Annotation category preference and positional analysis of binding density

For calculating the annotation category preference, we calculated the difference in the fraction of

T-to-C reads per annotation category between each RBP library and the reference library. For

example, if the fraction of miRNA annotated reads with T-to-C transitions in a specific RBP

library was 0.20 compared to 0.05 in the reference library, the miRNA preference value for this

specific RBP is 0.15. For the positional binding analysis, we selected genes (n=15120) using

GENCODE v19 as annotation based on our earlier work on HEK293 RNA processing and

turnover dynamics (Mukherjee et al., 2017). Isoform expression was calculated using RSEM (Li

and Dewey, 2011). For each gene, we selected the transcript isoform with the highest isoform

percentage or chose one randomly in case of ties (n=8298). The list of selected transcript

isoforms was used to calculate the median 5' UTR, CDS and 3' UTR length proportions (5'

UTR=0.06, CDS=0.53, 3' UTR=0.41) using R Bioconductor packages GenomicFeatures and

GenomicRanges. For regions downstream annotated transcription ends (TES) and adjacent to

splice sites, we chose windows of fixed sizes (TES 500nt, 5' and 3' splice sites 250nt each). We

generated coverage tracks from the PARalyzer output alignment files and intersected those with

the filtered transcripts. Each annotation category was binned according to its relative coverage

averaged according to each bin. For intronic coverage, we averaged across all introns per gene,

given a minimal intron length of 500nt. All bins were stitched to one continuous track per

transcript. Altogether 6632 intron containing transcripts showed coverage in at least one

PARCLIP library. For each library, we required transcripts to have a minimal coverage

maximum of > 2. For each transcript, we scaled the binned coverage dividing by its maximal

coverage (min-to-1 scaling) to emphasize spatial patterns independent from transcript expression

levels. Replicate RBP PARCLIP libraries were combined at this point. Transcripts targeted in

more than one replicate library were aggregated using the average of their binned coverage.

RBPs with less than 50 filtered target transcripts (after aggregation) were not considered. Next,

we split transcript coverage in two parts, separating 5' UTR to TES regions and intronic regions.

To generate the scaled meta coverage across all targeted transcripts per RBP, we used the

heatMeta function from the Genomation package. For the 5'UTR to TES, we scaled each RBP

meta-coverage track independent of other RBPs. For each RBP, we subtracted the scaled meta

coverage of PARCLIP reference library (Friedersdorf and Keene, 2014). For intronic sequences,

we scaled each RBP relative to all other RBPs to highlight RBPs with more substantial intronic

binding patterns. Finally, we visualized the density using pheatmap.

Sequence analysis

We calculated 6-mer frequencies with Jellyfish from all reads that generated a PARalyzer

binding site for each library. For each RBP, we selected the library with the lowest percent of

duplicated sequences (see supplemental table 1) to serve as a representative library for the

sequence analysis and factor analysis. For each RBP, we counted the number of 6-mers with a

frequency of x or higher, where x was from 1/4096 to 1/4. To evaluate the 6-mers enriched by a

given RBP relative to the reference library, we regressed the RBP 6-mer frequency against the

the reference library 6-mer frequency and collected the residuals (the unexplained variance).

Next, identified all 6-mers that were found as the top 5 enriched over the reference library for

any of the analyzed RBPs. We clustered the enrichment scores for the 6-mers across all RBPs

and generated a heatmap using the 'aheatmap' function in NMF R package. We ran SSMART

using all binding sites found in mRNA-derived annotation categories ranked by the library size

normalized enrichment over the reference library.

Factor analysis

For each site identified we calculated a library size normalized enrichment compared to the the

reference library. We calculated the sum of all enrichment scores for all sites annotated as

mRNA and lncRNA. Next, we normalized for expression levels (collected the residuals) to

create the final matrix of values. The number of factors, 10, was determined using the majority

result of numerous methods to estimate the number of factors. Clustering of the score matrix was

performed using the most stable results from numerous iterations of k-means clustering.

Gene ontology analysis

Multiple-test corrected gene ontology enrichment values were calculated using the TOPGO R

package. For each set of genes, we used all 13,299 genes in the factor analysis as the background

or gene universe. Enrichment was calculated using the 'parent-child' approach on the top 100

enriched terms. This metric accounts for the hierarchical organization of gene ontology terms to

minimize false-positive enrichments. We performed a Bonferonni multiple test correction on the

enrichment p-values.

Premature and mature RNA quantification

Mature- and premature-transcript expression, transcripts per million (TPM), was quantified with

RSEMv1.2.11   (http://deweylab.biostat.wisc.edu/rsem/src/rsem-1.2.11.tar.gz)   as   described

previously (Mukherjee et al., 2017). Briefly, for each gene we included an additional isoform

corresponding to the sequence of the full gene locus. Specifically, we modified the

GENCODEv19 gtf and used this as the input for the 'rsem-prepare-reference' function to

generate a modified index used for quantification. For each gene, we calculated the expression of

'mature' RNA as the sum of all isoforms for that gene excluding the 'primary' transcript. For

intronless genes, premature and mature expression values were summed. We performed this

analysis on the ELAVL1 knockdown RNA-seq experiments (Kishore et al., 2011).

Cell-to-cell expression variability

RNA-seq gene expression data for 25 individual HEK293 cells were downloaded from

(Guillaumet-Adkins et al., 2017). We calculated the coefficient of variation (100*standard

deviation/mean) for each gene across all 25 cells.

Figure 1. RBP analyzed and binding preferences by RNA category. A) Heatmap of reference

normalized annotation category preference for each RBP clustered into 8 branches by color

(left). The heatmap represents the difference in the proportion of sites for a given annotation

category in the RBP library versus the reference library. Heatmap of the reference library

normalized relative positional binding preference of the 55 RBPs with enriched binding in at

least one mRNA-relevant annotation category per branch (right). RBP-specific binding

preferences were averaged across selected transcripts (see methods). The relative spatial

proportion of 5'UTR, coding regions and 3'UTR were averaged across all selected transcript

isoforms. For TES (regions beyond transcription end site), 5' splice site, and 3' splice site, we

chose fixed windows (250nt for TES and 500nt for splice sites). For each RBP, meta-coverage

was scaled between 5'UTR to TES. The 5' and 3' intronic splice site coverage was scaled

separately from other regions but relative to each other.

Figure 2. RBP binding sequence specificity and elements. A) Heatmap of reference

normalized 6-mer enrichment for top 5 enriched 6-mers for each RBP in the set of RBPs

exhibiting more sequence specificity than the reference.

Figure 3. RNA regulatory modules. A) Factor analysis of target RNA encoding genes binding

normalized by the reference library and expression for the 55 RBPs binding to mRNAs and

lncRNAs for 13,299 genes (see 'factor analysis' section in methods for details). Spring-

embedded graph of the factor loading matrix, indicating the association between each of the 55

RBPs and one of the 10 factors. Nodes color-coded by RNA annotation category preference

cluster membership from figure 1. Edge width scales with factor loadings (thicker edge = higher

factor loading = stronger association). Only edges with a factor loading > 0.2 (positive values in

black) or < -.2 (negative values in green) depicted.

Figure 4. Functional characterization of RNA regulatory modules. A) The difference in

either A) primary or B) mature RNA expression (transcripts per million) upon ELAVL1

knockdown by siRNA treatment (y-axis), specifically the $\log_2$[siRNA EGFP TPM]-$\log_2$[siRNA

ELAVL1 TPM], for each gene set. C) Heatmap of the median value of synthesis rate, processing

rates, degradation rates, cytoplasmic versus nuclear localization, polyribosomal versus

cytoplasmic localization, and translational status from ribosome profiling data for each gene set

(top). Heatmap of the odds-ratio of the overlap between factor associated gene sets with

annotation (bottom). D) Box-and-whisker plot for each gene set of the enrichment in P-bodies.

E) Box-and-whisker plot for each gene set of the coefficient of variation across 25 individual

HEK293 cells.

Supplemental Figure 1. QC filtering of libraries. A) Description of PAR-CLIP suite to assess

library quality control per annotation category (left). Example of number of reads mapping to

each RNA category with up to 2 mismatches resolved by length of adapter-extracted sequence

reads for an ELAVL1 library (middle). Sequencing read composition of the most abundant RNA

category fir the ELAVL1 library. Reads were assigned as d0 (white), d1 T-to-C (red), d1 other

than T-to-C, (light gray), and d2 (black) (right). B) Libraries had to have > 20,000 aligned reads

and a mean conversion specificity > 0, and a higher mean T-to-C fraction than the the reference

library (red lower, blue higher). C) Number of libraries analyzed and their quality control status.

D) Count of libraries passing QC per RBP. E) Examples of outlier library removal (libraries

labeled with red text were removed) based on correlation of read 6-mer frequency for RBPs with

3 or more libraries.

Supplemental Figure 2. Grouping RBPs by sequence specificity. A) Heatmap of the number

of 6-mers enriched per RBP at different specificity thresholds. The color scale represents the $\log_2$

[number of 6-mers] that are enriched at a given threshold (y-axis). The thresholds are represented

as $\log_2$ [6-mer frequency]. There are 4096 different 6-mers and if they were uniformly present

this would represent a value of -12 $=\log_2$ [1/4096]. The horizontal dashed lines at -8, represents

16-fold enrichment over a uniform background. For reference, the vertical dashed lines indicate

the behavior of the reference library. B) Top 3 SSMART motif results using all binding sites

found in mRNA-derived annotation categories ranked by the library size normalized enrichment

over reference library.

Supplemental Figure 3. Factor analysis model selection and performance. A) Plot

of eigenvalues versus number of factors to determine the optimal number of factors using four

methods (different colors). B) Barplot of the communality, or the variance in a given RBP

cumulatively explained by the all factors. C) Heatmap of the median factor score coefficient

value for all genes that clustered together. The number of genes assigned to a specific factor and

the top two most significant enriched GO annotations for each ontology class: molecular

function (MF), cellular component (CC), and biological process (BP).

Supplemental Figure 4. RNA metabolism profiles for factor-associated gene sets. A) Box-and-whisker plot for each gene set of the synthesis rates, processing rates, degradation rates, cytoplasmic versus nuclear localization (Cyt vs Nuc), polyribosomal versus cytoplasmic localization (Poly vs Cyt), and translational status from ribosome profiling data. B) Heatmap of the odds-ratio of the overlap between factor associated gene sets with RNA categories based on similar metabolic profiles from (Mukherjee et al., 2017). C) Heatmap of the odds-ratio of the overlap between factor associated gene sets and protein localization annotation. D) Box-and-whisker plot for each gene set of the median expression across 25 HEK293 cells.

Bahar Halpern, K., Caspi, I., Lemze, D., Levy, M., Landen, S., Elinav, E., Ulitsky, I., Itzkovitz, S., 2015. Nuclear Retention of mRNA in Mammalian Tissues. Cell Rep 13, 2653–2662. doi:10.1016/j.celrep.2015.11.036

Baltz, A.G., Munschauer, M., Schwanhäusser, B., Vasile, A., Murakawa, Y., Schueler, M., Youngs, N., Penfold-Brown, D., Drew, K., Milek, M., Wyler, E., Bonneau, R., Selbach, M., Dieterich, C., Landthaler, M., 2012. The mRNA-bound proteome and its global occupancy profile on protein-coding transcripts. Mol. Cell 46, 674–690. doi:10.1016/j.molcel.2012.05.021

Battich, N., Stoeger, T., Pelkmans, L., 2015. Control of Transcript Variability in Single Mammalian Cells. Cell 163, 1596–1610. doi:10.1016/j.cell.2015.11.018

Cooper, T.A., Wan, L., Dreyfuss, G., 2009. RNA and Disease. Cell 136, 777–793. doi:10.1016/j.cell.2009.02.011

Corcoran, D.L., Georgiev, S., Mukherjee, N., Gottwein, E., Skalsky, R.L., Keene, J.D., Ohler, U., 2011. PARalyzer: definition of RNA binding sites from PAR-CLIP short-read sequence data. Genome Biol. 12, R79. doi:10.1186/gb-2011-12-8-r79

Fan, X.C., Steitz, J.A., 1998. HNS, a nuclear-cytoplasmic shuttling sequence in HuR. Proc. Natl. Acad. Sci. U.S.A. 95, 15293–15298.

Farina, K.L., Hüttelmaier, S., Musunuru, K., Darnell, R., Singer, R.H., 2003. Two ZBP1 KH domains facilitate β-actin mRNA localization, granule formation, and cytoskeletal attachment. The Journal of Cell Biology 160, 77–87. doi:10.1083/jcb.200206003

Fredericks, A.M., Cygan, K.J., Brown, B.A., Fairbrother, W.G., 2015. RNA-Binding Proteins: Splicing Factors and Disease. Biomolecules 5, 893–909. doi:10.3390/biom5020893

Friedersdorf, M.B., Keene, J.D., 2014. Advancing the functional utility of PAR-CLIP by quantifying background binding to mRNAs and lncRNAs. Genome Biol. 15, R2. doi:10.1186/gb-2014-15-1-r2

Garzia, A., Jafarnejad, S.M., Meyer, C., Chapat, C., Gogakos, T., Morozov, P., Amiri, M.,

Shapiro, M., Molina, H., Tuschl, T., Sonenberg, N., 2017a. The E3 ubiquitin ligase and
RNA-binding protein ZNF598 orchestrates ribosome quality control of premature
polyadenylated mRNAs. Nat Commun 8, 16056. doi:10.1038/ncomms16056
Garzia, A., Meyer, C., Morozov, P., Sajek, M., Tuschl, T., 2017b. Optimization of PAR-CLIP
for transcriptome-wide identification of binding sites of RNA-binding proteins. Methods
118-119, 24–40. doi:10.1016/j.ymeth.2016.10.007
Gerstberger, S., Hafner, M., Tuschl, T., 2014. A census of human RNA-binding proteins. Nature
Reviews Genetics 15, 829–845. doi:10.1038/nrg3813
Gruber, A.J., Schmidt, R., Gruber, A.R., Martin, G., Ghosh, S., Belmadani, M., Keller, W.,
Zavolan, M., 2016. A comprehensive analysis of 3' end sequencing data sets reveals novel
polyadenylation signals and the repressive role of heterogeneous ribonucleoprotein C on
cleavage and polyadenylation. Genome Res. 26, 1145–1159. doi:10.1101/gr.202432.115
Guillaumet-Adkins, A., Rodríguez-Esteban, G., Mereu, E., Mendez-Lago, M., Jaitin, D.A.,
Villanueva, A., Vidal, A., Martinez-Marti, A., Felip, E., Vivancos, A., Keren-Shaul, H.,
Heath, S., Gut, M., Amit, I., Gut, I., Heyn, H., 2017. Single-cell transcriptome conservation
in cryopreserved cells and tissues. Genome Biol. 18, 45. doi:10.1186/s13059-017-1171-9
Hafner, M., Landthaler, M., Burger, L., Khorshid, M., Hausser, J., Berninger, P., Rothballer, A.,
Ascano, M., Jungkamp, A.-C., Munschauer, M., Ulrich, A., Wardle, G.S., Dewell, S.,
Zavolan, M., Tuschl, T., 2010. Transcriptome-wide identification of RNA-binding protein
and microRNA target sites by PAR-CLIP. Cell 141, 129–141.
doi:10.1016/j.cell.2010.03.009
Hubstenberger, A., Courel, M., Bénard, M., Souquere, S., Ernoult-Lange, M., Chouaib, R., Yi,
Z., Morlot, J.-B., Munier, A., Fradet, M., Daunesse, M., Bertrand, E., Pierron, G.,
Mozziconacci, J., Kress, M., Weil, D., 2017. P-Body Purification Reveals the Condensation
of Repressed mRNA Regulons. Mol. Cell 68, 144–157.e5. doi:10.1016/j.molcel.2017.09.003
Jankowsky, E., Harris, M.E., 2015. Specificity and nonspecificity in RNA-protein interactions.
Nat. Rev. Mol. Cell Biol. 16, 533–544. doi:10.1038/nrm4032
Jønson, L., Vikesaa, J., Krogh, A., Nielsen, L.K., Hansen, T.V., Borup, R., Johnsen, A.H.,
Christiansen, J., Nielsen, F.C., 2007. Molecular composition of IMP1 ribonucleoprotein
granules. Mol. Cell Proteomics 6, 798–811. doi:10.1074/mcp.M600346-MCP200
Keene, J.D., 2007. RNA regulons: coordination of post-transcriptional events. Nature Reviews
Genetics 8, 533–543. doi:10.1038/nrg2111
Kishore, S., Jaskiewicz, L., Burger, L., Hausser, J., Khorshid, M., Zavolan, M., 2011. A
quantitative analysis of CLIP methods for identifying binding sites of RNA-binding proteins.
Nat. Methods 8, 559–564. doi:10.1038/nmeth.1608
Koutmou, K.S., Schuller, A.P., Brunelle, J.L., Radhakrishnan, A., Djuranovic, S., Green, R.,
2015. Ribosomes slide on lysine-encoding homopolymeric A stretches. Elife 4, 446.
doi:10.7554/eLife.05534
König, J., Zarnack, K., Rot, G., Curk, T., Kayikci, M., Zupan, B., Turner, D.J., Luscombe, N.M.,
Ule, J., 2010. iCLIP reveals the function of hnRNP particles in splicing at individual
nucleotide resolution. Nat. Struct. Mol. Biol. 17, 909–915. doi:10.1038/nsmb.1838
Lebedeva, S., Jens, M., Theil, K., Schwanhäusser, B., Selbach, M., Landthaler, M., Rajewsky,
N., 2011. Transcriptome-wide analysis of regulatory interactions of the RNA-binding protein
HuR. Mol. Cell 43, 340–352. doi:10.1016/j.molcel.2011.06.008
Li, B., Dewey, C.N., 2011. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics 12, 323. doi:10.1186/1471-2105-12-323

Maatz, H., Jens, M., Liss, M., Schafer, S., Heinig, M., Kirchner, M., Adami, E., Rintisch, C.,
Dauksaite, V., Radke, M.H., Selbach, M., Barton, P.J.R., Cook, S.A., Rajewsky, N.,
Gotthardt, M., Landthaler, M., Hubner, N., 2014. RNA-binding protein RBM20 represses
splicing to orchestrate cardiac pre-mRNA processing. J. Clin. Invest. 124, 3419–3430.
doi:10.1172/JCI74523
Mansfield, K.D., Keene, J.D., 2012. Neuron-specific ELAV/Hu proteins suppress HuR mRNA
during neuronal differentiation by alternative polyadenylation. Nucleic Acids Res. 40, 2734–
2746. doi:10.1093/nar/gkr1114
Marçais, G., Kingsford, C., 2011. A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. Bioinformatics 27, 764–770. doi:10.1093/bioinformatics/btr011
Martin, G., Gruber, A.R., Keller, W., Zavolan, M., 2012. Genome-wide analysis of pre-mRNA
3" end processing reveals a decisive role of human cleavage factor I in the regulation of 3"
UTR length. Cell Rep 1, 753–763. doi:10.1016/j.celrep.2012.05.003
Mesarovic, M.D., Sreenath, S.N., Keene, J.D., 2004. Search for organising principles:
understanding in systems biology. Syst Biol (Stevenage) 1, 19–27.
Moore, M.J., 2005. From birth to death: the complex lives of eukaryotic mRNAs. Science 309,
1514–1518. doi:10.1126/science.1111443
Mukherjee, N., Calviello, L., Hirsekorn, A., de Pretis, S., Pelizzola, M., Ohler, U., 2017.
Integrative classification of human coding and noncoding genes through RNA metabolism
profiles. Nat. Struct. Mol. Biol. 24, 86–96. doi:10.1038/nsmb.3325
Mukherjee, N., Corcoran, D.L., Nusbaum, J.D., Reid, D.W., Georgiev, S., Hafner, M., Ascano,
M., Tuschl, T., Ohler, U., Keene, J.D., 2011. Integrative regulatory mapping indicates that
the RNA-binding protein HuR couples pre-mRNA processing and mRNA stability. Mol.
Cell 43, 327–339. doi:10.1016/j.molcel.2011.06.007
Mukherjee, N., Jacobs, N.C., Hafner, M., Kennington, E.A., Nusbaum, J.D., Tuschl, T.,
Blackshear, P.J., Ohler, U., 2014. Global target mRNA specification and regulation by the
RNA-binding protein ZFP36. Genome Biol. 15, R12. doi:10.1186/gb-2014-15-1-r12
Munteanu, A., Mukherjee, N., Ohler, U., 2018. SSMART: Sequence-structure motif
identification for RNA-binding proteins. bioRxiv 287953. doi:10.1101/287953
Nielsen, F.C., Nielsen, J., Christiansen, J., 2001. A family of IGF-II mRNA binding proteins
(IMP) involved in RNA trafficking. Scand. J. Clin. Lab. Invest. Suppl. 234, 93–99.
Pandit, S., Zhou, Y., Shiue, L., Coutinho-Mansfield, G., Li, H., Qiu, J., Huang, J., Yeo, G.W.,
Ares, M., Fu, X.-D., 2013. Genome-wide analysis reveals SR protein cooperation and
competition in regulated splicing. Mol. Cell 50, 223–235. doi:10.1016/j.molcel.2013.03.001
Pullmann, R., Kim, H.H., Abdelmohsen, K., Lal, A., Martindale, J.L., Yang, X., Gorospe, M.,
2007. Analysis of turnover and translation regulatory RNA-binding protein expression
through binding to cognate mRNAs. Mol. Cell. Biol. 27, 6265–6278.
doi:10.1128/MCB.00500-07
Sheth, U., Parker, R., 2003. Decapping and decay of messenger RNA occur in cytoplasmic
processing bodies. Science 300, 805–808. doi:10.1126/science.1082320
Sundararaman, B., Zhan, L., Blue, S.M., Stanton, R., Elkins, K., Olson, S., Wei, X., Van
Nostrand, E.L., Pratt, G.A., Huelga, S.C., Smalec, B.M., Wang, X., Hong, E.L., Davidson,
J.M., Lécuyer, E., Graveley, B.R., Yeo, G.W., 2016. Resources for the Comprehensive
Discovery of Functional RNA Elements. Mol. Cell 61, 903–913.
doi:10.1016/j.molcel.2016.02.012
Tenenbaum, S.A., Carson, C.C., Lager, P.J., Keene, J.D., 2000. Identifying mRNA subsets in

messenger ribonucleoprotein complexes by using cDNA arrays. Proc. Natl. Acad. Sci.
U.S.A. 97, 14085–14090. doi:10.1073/pnas.97.26.14085
Ule, J., Jensen, K.B., Ruggiu, M., Mele, A., Ule, A., Darnell, R.B., 2003. CLIP identifies Nova-
regulated RNA networks in the brain. Science 302, 1212–1215.
doi:10.1126/science.1090095
Van Nostrand, E.L., Pratt, G.A., Shishkin, A.A., Gelboin-Burkhart, C., Fang, M.Y.,
Sundararaman, B., Blue, S.M., Nguyen, T.B., Surka, C., Elkins, K., Stanton, R., Rigo, F.,
Guttman, M., Yeo, G.W., 2016. Robust transcriptome-wide discovery of RNA-binding
protein binding sites with enhanced CLIP (eCLIP). Nat. Methods 13, 508–514.
doi:10.1038/nmeth.3810
Wilusz, J., Feig, D.I., Shenk, T., 1988. The C proteins of heterogeneous nuclear
ribonucleoprotein complexes interact with RNA sequences downstream of polyadenylation
cleavage sites. Mol. Cell. Biol. 8, 4477–4483.
Zhang, C., Frias, M.A., Mele, A., Ruggiu, M., Eom, T., Marney, C.B., Wang, H., Licatalosi,
D.D., Fak, J.J., Darnell, R.B., 2010. Integrative modeling defines the Nova splicing-
regulatory network and its combinatorial controls. Science 329, 439–443.
doi:10.1126/science.1191150
Zhao, J., Ohsumi, T.K., Kung, J.T., Ogawa, Y., Grau, D.J., Sarma, K., Song, J.J., Kingston, R.E.,
Borowsky, M., Lee, J.T., 2010. Genome-wide identification of polycomb-associated RNAs
by RIP-seq. Mol. Cell 40, 939–953. doi:10.1016/j.molcel.2010.12.011
716

A)

preference vs reference

−0.6 −0.4 −0.2 0 0.2 0.4 0.6

binding density relative to reference

−0.4 −0.2 0 0.2 0.4

A)

A)

Legend:
- miRNA, 3' UTR
- 3' UTR
- CDS, 3' UTR
- intron, 3' UTR, lncRNA
- mixed