BayesENproteomics: Bayesian elastic nets for quantification of proteoforms in complex samples

Venkatesh Mallikarjun[1, 2], Stephen M. Richardson[2] and Joe Swift[1, 2, *]

(1) Wellcome Trust Centre for Cell-Matrix Research;

(2) Division of Cell Matrix Biology and Regenerative Medicine, School of Biological Sciences, Faculty of Biology, Medicine and Health, Manchester Academic Health Science Centre, University of Manchester, M13 9PL, UK.

  * Correspondence: joe.swift@manchester.ac.uk

Running title: *Label-free proteoform quantification*

Key words: Label-free proteomics / post translational modification / Bayesian linear regression / primary cells and tissue / Markov chain Monte Carlo

Abstract

Multivariate regression modelling provides a statistically powerful means of quantifying the effects of a given treatment while compensating for sources of variation and noise, such as variability between human donors and the behaviour of different peptides during mass spectrometry. However, methods to quantify endogenous post-translational modifications (PTMs) are typically reliant on summary statistical methods that fail to consider sources of variability such as changes in levels of the parent protein. Here, we compare three multivariate regression methods, including a novel Bayesian elastic net algorithm (BayesENproteomics) that enables assessment of relative protein abundances while also quantifying identified PTMs for each protein. We tested the ability of these methods to accurately quantify expression of proteins in a mixed-species benchmark experiment, and to quantify synthetic PTMs induced by stable isotope labelling. Finally, we extended our regression pipeline to calculate fold changes at the pathway level, providing a complement to commonly used enrichment analysis. Our results show that BayesENproteomics can quantify changes to protein levels across a broad dynamic range while also accurately quantifying PTM and pathway-level fold changes.

Introduction

Spiraling costs of drug/therapeutic development and a low probability of success have driven an increase in the use of models based on patient-derived material in an attempt to determine translational potential prior to clinical studies. However, unlike samples from genetically homogeneous, completely inbred model organisms, patient samples can possess substantial differences between individuals. This variability between samples can make discerning a statistically significant effect extremely difficult using the standard statistical methods commonly used in biology. The lack of statistical power caused by poor signal-to-noise ratios in primary samples and the difficulty in obtaining large donor cohorts can compromise potentially promising biological studies, resulting in wasted time and prohibitive expenditure.

The problem of high inter-individual differences is rarely more acute than in high-throughput omics experiments. The difference between patient samples is particularly evident in omics experiments where subtle differences in the levels of individual features within samples (e.g. transcripts, peptides derived from proteins, etc.) are compounded over many such features, demonstrating quantifiable inter-individual differences even between genetically identical twins (Brodin *et al*, 2015). This is especially relevant to typical mass spectrometry (MS) proteomic methods wherein quantities associated with multiple features (i.e. peptides) are used in the calculation of a fold change for a single protein.

In bottom-up MS-based proteomics, proteins are enzymatically digested into peptides and the abundances of these individual peptides are used to derive protein quantifications. A core assumption is that peptide abundance is proportional to protein abundance. However, each peptide has different physico-chemical properties that mean they each behave differently during sample preparation or within the mass spectrometer itself. For instance, residues surrounding cleavage sites can affect efficiency of enzymatic digestion (Rodriguez *et al*, 2008; Lawless & Hubbard, 2012); some peptides ionise less efficiently than others (Abaye *et al*, 2011) and in data-dependent analysis, peptides of similar mass-to-charge (m/z) ratios may compete during co-elution or ionisation (Schliekelman & Liu, 2014), potentially biasing the peptides finally detected. These discrepancies can be exacerbated by

biologically relevant post-translational modifications (PTMs) that alter the behaviour of individual peptides, resulting in substantial differences in measured intensities for peptides that belong to the same protein, even in purified protein samples. Even with normalisation, differences in the behaviour of individual features between donors and experimental treatments are difficult to account for.

It was recently shown that combining MS datasets from multiple fractions and extraction conditions could achieve near-complete coverage of the human cell proteome, including >14000 protein isoforms (Bekker-Jensen *et al*, 2017). Interestingly, the same study also identified ~10000 phosphorylation sites and ~7000 acetylation sites without specific enrichment. New sample preparation techniques and the increasing sensitivity of MS instrumentation have increased coverage of PTMs. However, with this increase in sensitivity comes the problem of resolving differential regulation of different forms of the same protein (proteoforms), as not all peptides belonging to a given protein respond in the same way to a given treatment, an assumption made by many commonly used quantification methods.

Common statistical methods for dealing with variability include the summarisation of peptide intensities using averages, sums or medians to give subject-level protein quantification (Goeminne *et al*, 2015). However, these subject-level summary models fail to account for differences in behaviour of peptides belonging to the same protein during analysis, and thus are prone to perturbation by outlier peptides. They are also highly dependent on obtaining large numbers of peptides from each protein and thus can suffer from low statistical power when analysing less extreme fold changes in small or low abundance proteins (e.g. transcription factors). To address this, previous statistical models have focused on exclusion of outlier peptides to improve protein quantification (Forshed, 2013; Swift *et al*, 2013a), at the cost of a loss of information relating to different proteoforms for a given protein; or have attempted to deduce differences in the abundance of peptides belonging to different sub-populations of the same protein (Henao *et al*, 2012, 2013; Webb-Robertson *et al*, 2014; Zeng *et al*, 2017).

For proteomics, linear regression models represent a powerful means of teasing apart different, known sources of variability in peptide intensities, with the aim of returning the variability caused by a given experimental condition. Linear modelling has become the standard for transcriptomic analysis

thanks to the widely used LIMMA package (Smyth, 2004) and has been successfully adapted for MS data analysis. For example, in the peptide-based model proposed by Choi et al. (2014) with donor effects added:

$$y_{fgd} = \beta_0 + X_f\beta_f + X_g\beta_g + X_d\beta_d + \varepsilon_{fgd}. \qquad (1)$$

Where, for a given protein, y_{fgd} (the **response variable**) corresponds to the observed log$_2$ intensity for peptide *f* of that protein under experimental condition *g* from donor *d*. **Predictor variables** β_f, β_g and β_d correspond to the effect sizes of peptide *f*, group *g* and donor *d*, respectively. β_0 represents the intercept term. ε corresponds to a Gaussian error term centred on 0 with a variance (σ^2) specific to each protein. The purpose of the regression calculation was to find how much the slope of the resulting line differed for a given experimental treatment (i.e. the effect size β_g caused by the experimental treatment of interest *g*). In this instance the design matrix *X* consisted solely of categorical variables encoded as binary variables of size $n \times p$ (where n = number of observations and p = number of βs).

However, peptide-based linear models can be prone to overfitting, wherein predictive power is lost as the model attempts to fit the noise within the data rather than any overall trends. This may be due to the effects of outlier peptides, such as those that possess a biologically relevant PTM with a fold change different to that of its parent protein, or simply due to peptide mis-identification. The effects of outlier peptides on the final linear model fit can be mediated by various weighting strategies (e.g. see Goeminne et al. (2016)). However, for modelling changes in PTM abundance, it would be useful to obtain effect sizes for peptides that specifically interact with a given treatment and separate them from effect sizes for other parameters that may be interacting with these peptides. Linear modelling is more flexible than the name suggests and can also handle interactions between predictor variables; for example, where an experimental condition has an effect on the intensity of an individual peptide – independent of its effect on the protein as a whole – possibly due to some biologically important PTM of that specific peptide. For example, we can extend the model in (1) to allow separation of

peptide:treatment and peptide:donor effects in studies using primary human samples, as in (2) (see also the model in (Clough *et al*, 2009)).

$$y_{fgd} = \beta_0 + X_f\beta_f + X_g\beta_g + X_d\beta_d + X_fX_g\beta_{f:g} + X_fX_d\beta_{f:d} + \varepsilon_{fgd}. \tag{2}$$

Wherein y_{fgd} corresponds to the intensity of peptide *f*, in experimental group *g*, from protein *i*, extracted from technical replicate *r* of donor *d*. The βs correspond to the fitted coefficients for each predictor variable. Note the interactions coefficients $\beta_{f:g}$, and $\beta_{f:d}$ in (2), denoting variability in individual peptide intensities, *f*, caused by the experimental treatment *g* or largely unknown features of the individual donor *d*. From here on we refer to (1) as linear and (2) as non-linear.

Unfortunately, overfitting becomes particularly acute in models that attempt to deal with many potential sources of variability, some of which may interact or correlate with one-another, and where the number of parameters to be estimated exceeds the number of observations. The inclusion of interaction terms in a model has potential to provide important biological insight regarding different isoforms and PTMs of a given protein (collectively called proteoforms) – vital mechanisms of post-translational regulation of cellular function. However, interaction terms may not be necessary for all proteins in a dataset and the decision whether or not to include them would need to be made on a protein-by-protein basis.

Simpler models (e.g. those used by (Choi et al. (2014) and (Goeminne et al. (2016)) minimise the potential for overfitting found in more complex models. However, when many potential sources of variability exist it may be necessary to fit increasingly complex models wherein the choice of terms included, particularly any interaction terms, may need to be decided on a protein-by-protein basis. Failure to select the correct terms could result in models failing to fit correctly, or even at all, especially if complex interaction terms are included when not needed. However, not including these terms may mean that subtle differences in peptide behaviour that occur only in specific proteins may go unaccounted for, resulting in a loss of accuracy. Thus, a means of automatically selecting appropriate features from a more complex model for use in fitting a final, sparse model is necessary.

Regularisation can be employed to minimise overfitting and perform feature selection. Regularisation involves attaching penalty weights to each coefficient (β), minimising its contribution to the final fit. The nature of the penalty can differ depending on the type of Regularisation used. In the ridge regression algorithm used by Goeminne et al. (2016) an "L_2" penalty is applied wherein βs are shrunk according to their squared norm, minimising the contribution of less important coefficients in explaining the total model variance. Although ridge regression shrinks coefficient estimates down to emphasise the most important βs, all βs remain non-zero. Another form of Regularisation is LASSO (least absolute shrinkage and selection operator), which shrinks βs according to their "L_1" absolute norm. With LASSO, some βs can be set to exactly zero, excluding them from the model and giving rise to so-called "sparse" solutions. Several variations of these methods have been proposed, including an "elastic net" variation that combines properties of both ridge regression and LASSO to perform both model selection and shrinkage of remaining non-zero βs (Zou & Hastie, 2005). These frequentist Regularisation methods have Bayesian parallels depending on the prior distributions the β posteriors are sampled from, with L_2 Regularisation being equivalent to sampling of βs from a Gaussian distribution, L_1 being equivalent to sampling from a Laplace (double exponential) prior and the elastic net being equivalent to β sampling from an intermediate distribution (Bornn *et al*, 2010).

Here we develop a Bayesian elastic net algorithm to provide regularised protein-specific models (BayesENproteomics), taking into account potential donor variability and interactions between specific peptides and experimental treatments and/or donor effects. This has the advantage in implicitly allowing for any individual peptide to behave differently from its identified protein group in response to treatment or donor effects. This means that BayesENproteomics does not assume that peptides have been identified correctly – a point we enforce by incorporating observation weights based on peptide identification confidence. However, unlike previous Bayesian regression models that attempt to accommodate variable peptide behaviour within a protein (Henao *et al*, 2012, 2013; Webb-Robertson *et al*, 2014), BayesENproteomics can still provide fold change estimates for protein groups containing peptides with specific identifications, maintaining interpretability of subsequent results by presenting relative quantification of the "dominant" proteoform for that protein (separating different isoforms if unique peptides are available), whilst also presenting values for differentially abundant PTMs for

further investigation. Finally, we extend our linear modelling pipeline to encompass pathway analysis as a complement to standard enrichment analysis. This enables complete analysis of a dataset from MS1 intensities to functional interpretation of observed proteoform changes.

Results

BayesENproteomics shows increased sensitivity and accuracy in estimating protein fold changes compared to other regression models

To produce a benchmark dataset with known (ground truth) fold changes, peptides prepared from primary human mesenchymal stem cell (MSC) lysates and female C57BL/6J mouse skin were mixed in ratios of 3:1, 1:1 and 1:3 and analysed with a Q Exactive HF mass spectrometer (Figure 1A). Mixed species datasets have been used previously to validate protein quantification methods (Swift *et al*, 2013a), but the method has practical applications, such as when quantifying protein content in xenograft models (Swift *et al*, 2013b; Ivanovska *et al*, 2017). Mixed mouse/human samples provide a difficult problem for prospective quantification algorithms as approximately 70% of tryptic peptides are shared between mouse and humans and thus have to be discarded to obtain estimates for specific proteins. This meant that the quantification algorithms examined here had a limited number of observations with which to work. Secondly, the dynamic range of absolute protein abundances was derived from actual biological samples and so provided an accurate report of the error associated with estimating fold changes in low abundance proteins. Finally, this experiment yielded twin datasets (mouse and human); one with pronounced donor-donor variability (human cells) and one where all animals had the same genotype and environment (mouse skin), and so could be adequately explained by the model in equation (2) without donor effects. We reasoned that an algorithm that could account for donor-donor variability should be as accurate on the human dataset as on the mouse dataset.

Logged signals from species-unique peptides were normalised to the medians of the opposite species-unique peptides to provide the observed $\log_2$ ratios used in quantification. To automatically

select appropriate β parameters it was possible to employ Regularisation strategies such as ridge, LASSO or elastic net regression to select the best set of βs by shrinking unimportant βs to zero (Figure 1B). As LASSO has been shown to lack consistency when selecting from several correlated βs, we chose to use elastic net-based Regularisation as the introduction of L_2 Regularisation has been shown to introduce consistency to LASSO estimates by inducing grouping of correlated βs (Zou & Hastie, 2005). However, the standard formula for calculating standard errors (SE) of β estimates ($\sigma^2 (X^TX)^{-1}$, where σ^2 is the residual variance and X is the design matrix, with superscript T denoting the transpose of X and SEs for each β, lie on the main diagonal of the resulting square matrix) was not suitable for sparse regression models due to shrinking of variance estimates by regularisation. This resulted in underestimation of variance and incorrectly lower SEs and p-values. While there were no simple ways to calculate SEs of coefficients in sparse regularised models, bootstrapping has commonly been used, however this is computationally intensive to do for hundreds of proteins. Bootstrapped SE estimates also differ from iteration to iteration, due to the inherent randomness of bootstrapping, thus compromising reproducibility. Bayesian regression methods addressed this limitation by estimating parameters as distributions for which summary statistics (e.g. means, standard deviations and confidence intervals) were trivial to calculate. Bayesian regression methods were also more flexible in how the model parameters and hyperparameters could be defined. This made Bayesian regression methods generally better suited to fit non-linear models, such as in equation (2), which frequentist methods may have struggled to fit. To implement this non-linear regression we used a Monte Carlo Markov Chain (MCMC)-based Gibbs sampler to repeatedly sample from and update the prior conditional distributions for the parameters we wished to estimate (Figure 1C). Examples of the resulting chains are shown in Figure 1D and demonstrate good mixing and exploration of the posterior distributions.

MCMC-based methods can be used to deduce any unknown value, including missing values in proteomics experiments, provided appropriate conditional distributions can be provided. We used this property to sample from two prior distributions depending on whether a missing value was determined to be missing at random (MAR) or missing non-randomly (MNR) by logistic regression (adaptive multiple imputation (AMI); see Materials and Methods section for more details). As the majority of

missing values in bottom-up proteomics are assumed to be left censored, it is common to perform imputation by sampling from a down-shifted Gaussian distribution (DGD). Missing values that were imputed separately for each protein by AMI possessed a similar overall distribution to those imputed from a DGD as in Tyanova *et al* (2016), but with a greater number of values closer to the observed MS1 intensity mean, as expected (Figure 2A-C). To interrogate the accuracy of AMI further we created a synthetic dataset consisting of a single protein with 10 peptides measured across 2 treatment groups (G1 and G2) with 6 replicates each. Values for G1 and G2 were populated from normal distributions with means of 0 and -2, respectively, and variance = 1, such that the ground-truth (log-scale) fold change, G2-G1 = -2. 30% of these values were deleted randomly (simulating MAR missingness) along with all values < -2 (simulating MNR missingness) and we asked which of the two imputation methods tested could most accurately recapitulate the deleted values (Figure 2D-E). While neither method was capable of fully approximating the deleted values, single imputation from a DGD (Figure 2D) was markedly worse (R = 0.16) than AMI (Figure 2E; R = 0.36) with a more positive correlation coefficient versus actual values from the complete dataset and an estimated fold change closer to ground-truth, albeit with some variance compression in MNR values. These results showed that AMI, as implemented in BayesENproteomics, even when given the ability to decide otherwise, identified most missing values as MNR, but possessed greater accuracy, particularly when imputing MAR values in a mixed-missingness-type dataset.

Figure 3A shows estimation of $\log_2$ fold changes for non-differentially abundant proteins in pairwise comparisons between three technical replicates from the same (mouse skin) sample, comparing the BayesENproteomics algorithm with ordinary least-squares (OLS) and linear mixed-effects (with Huber residual weights, LME-H) models (see Structured Methods section). All methods correctly determined $\log_2$ fold change values clustered around zero and that almost no changes reached statistical significance (p-value of false discovery, with Benjamini-Hochberg correction, BHFDR < 0.05). In contrast, estimation of differentially abundant mouse (Figure 3B) and human (Figure 3C) proteins in our mixed human:mouse dataset showed larger variation in observed fold change values, with a number of proteins giving consistently incorrect fold change directions, regardless of the regression method. OLS and LME-H models had difficulty detecting any significantly differentially abundant

proteins for expected fold changes < 3. In contrast, BayesENproteomics correctly identified more proteins as significantly differentially abundant in all comparisons (Figure 3B, C). Importantly, fold change estimates calculated by BayesENproteomics possessed lower mean squared deviation from ground truth compared to those from OLS and LME-H (Figure 3D). Regularisation and weighting of residuals in LME-H did not markedly improve accuracy of fold change estimates compared to standard OLS models, although this may be due to the more complex model (2) that LME-H was required to fit (Figure 3D). BayesENproteomics showed decreased coefficient of variance (CV) values for differentially abundant proteins in the mixed human:mouse dataset compared to both OLS and LME-H (Figure 3E).

BayesENproteomics correctly detects increases in stable isotope-labelled proteoforms following PNGase F treatment in $H_2^{18}O$

Protein function and activity is strongly determined by diverse PTMs, giving rise to different proteoforms within a given protein population. MS-based proteomics represents a powerful means to systemically interrogate changes in PTM abundance. However, determining relative PTM fold changes from PTM'd peptide abundance is dependent on accurate protein quantification (Wu *et al*, 2011). While it is possible to calculate protein abundance from only unmodified peptides, this ignores the fact that some proteins are constitutively modified as part of their normal maturation pathway (e.g. many extracellular matrix proteins are heavily glycosylated prior to secretion) and excluding these PTM-containing peptides from protein quantification would exclude a large proportion of all proteoforms present for that protein. While methods have been developed to analyse individual proteoform abundance in multiplexed, labelled experiments (Malioutov *et al*, 2017), labelling is not always feasible and methods for quantifying differentially abundant PTMs in label-free experiments lag behind. Furthermore, while an ideal scenario would be to directly compare modified to unmodified peptide ratios (Tsai *et al*, 2017), this is not always achievable in label-free proteomics of complex samples. While linear regression modelling has been shown to impart greater power and accuracy to protein quantification compared to summary statistical methods (Goeminne *et al*, 2015), applying it to

study PTMs is difficult due to the requirement for complex non-linear models that may be subject to overfitting.

To interrogate how well the three models implemented here were able to detect differentially abundant PTMs, we developed a benchmark dataset using PNGase F to introduce a stable isotope (^{18}O) label on N-linked glycosylated residues (Figure 4A). We reasoned that PNGase F-treated peptides should show an increase in N-linked ^{18}O labelling compared to spontaneous deamidation observed in control peptides incubated in $H_2^{18}O$ buffer alone. To test if our PNGase F treatment of peptides had worked correctly (independently of any subsequent quantification methods) we first searched treated and control samples separately and counted the total number of N- or Q-linked, ^{18}O-labelled peptides identified (Figure 4B). This analysis showed that more ^{18}O-labelled peptides were identified in the PNGase F-treated samples than controls, which contained 79 N-linked peptides presumably from spontaneous non-enzymatic deamidation. We expected that most spontaneously deamidated peptides would be unaffected by PNGase F treatment and that we would only observe an increase in relative ^{18}O-peptide abundance or parent ^{18}O-protein abundance for PNGase F-responsive peptides. Furthermore, as spontaneous deamidation can occur to both Q and N residues, resulting in ^{18}O-E and ^{18}O-D residues, respectively, while PNGase F catalyses only deamidation of N-linked glycosylation sites, comparison to fold changes of Q-linked deamidation provided an ideal negative control.

To detect differentially abundant PTMs we employed three statistical methods. The first was a simple summary statistical extension to the OLS protein quantification method wherein log$_2$ fold changes for donor-normalised PTM'd peptide intensities were normalised to the log$_2$ fold changes of their parent protein (calculated by OLS). The second involved using the ridge regression performed by LME-H to fit the non-linear model described in equation (2) where peptide:group and peptide:donor interaction βs were entered as random intercept terms to limit overfitting. Finally, we used BayesENproteomics to fit the non-linear model described in (2) to calculate PTM fold changes. From simple cell lysates, following PNGase F treatment without enrichment we found 261 ^{18}O-labelled sites (180 N-linked, 81 Q-linked) in 172 proteins when samples from all groups were aligned and search together. Histograms in Figure 4C showed that OLS (top-left) and LME-H (middle-left) could not discern an

increase in N-linked ^{18}O-protein abundance following PNGase F treatment compared to Q-linked (all p-values > 0.05). In contrast, BayesENproteomics was able to correctly discern a positive skewing of the N-linked ^{18}O-protein histogram compared to that of the Q-linked proteins (all p-values < 0.05, Figure 4C, bottom-left).

Similarly, when measuring differentially abundant PTMs there was great disparity between the three methods. The summary statistical extension to OLS produced noisy estimates of PTM fold changes and was unable to discern an increase in N-linked ^{18}O-PTMs following PNGase F treatment compared to Q-linked (Figure 4C, top-right; 4D, top). LME-H displayed overly strong Regularisation of interaction terms, resulting in the majority of PTM fold change estimates being approximately zero (Figure 4C, mid-right; 4D, middle). BayesENproteomics estimates of N-linked ^{18}O-PTM fold changes were correctly positively skewed (Figure 4C, bottom-right; 4D, bottom) and significantly different from Q-linked ^{18}O sites. This showed that BayesENproteomics could quantify PTM fold changes more accurately than either normalised summary statistics or mixed-effects models.

Pathway analysis using linear modelling (PALM)

Theoretically, modelling pathway activity as a function of protein abundances (PALM; Figure 5A) can have several advantages over classical pathway *over-representation* analysis. Firstly, the PALM approach proposed here utilised the entire dataset and so was not affected by potentially arbitrary statistical cut-offs used in defining 'interesting' and 'background' sets for over-representation analysis of proteins, similar to more recent *enrichment* analysis tools (reviewed in Huang et al., 2009). Secondly, PALM used additional information regarding the direction and magnitude of fold changes and the error regarding their estimation to provide an overall estimate of how a given pathway behaved based on information available in a given dataset. Finally, PALM provides pathway-level fold change and error estimates, facilitating pathway-level clustering in complex, multi-treatment or time-series datasets to gain a better "bird's-eye view" of what's happening in a dataset. Figures 5B and 5C show volcano plots assembled from Reactome (Fabregat *et al*, 2018; Milacic *et al*, 2012) pathway quantification from the human:mouse mixed species dataset based on protein-level output from OLS, LME-H and BayesENproteomics. While the magnitude of pathway-level log$_2$ fold changes was

typically not equal to their protein components due to being inferred from relatively fewer observations and down-weighted according to uncertainty in initial protein fold change estimates (always > 0), directionality was conserved leading to an estimate of whether that pathway significantly increased or decreased. Pathway significance rose in accordance with the magnitude of their component protein fold changes (Figures 5B, C).

Discussion

Multivariate regression methods represent a powerful means to deconvolute sources of variance in complex experiments. Here we attempted to design a statistical method that was able to accurately quantify protein and PTM fold changes in label-free proteomics experiments performed on heterogeneous primary human samples. Others have previously shown the superiority of linear regression-based effect size estimates over summary statistics (Goeminne *et al*, 2015; Clough *et al*, 2009). Here we outlined a weighted Bayesian regression algorithm with elastic net regularisation (to cope with differentially abundant PTMs as well as the high variability (and generally low availability) of samples from human donors, while still maintaining high interpretability of the resulting models (Figure 6). In particular, BayesENproteomics incorporates an "adaptive" missingness mechanism that attempts to determine whether a value is MAR or MNR. Missing value imputation is performed as part of the Gibb's sampler and is thus unfortunately not available as a standalone component. In addition, we utilised a novel weighing method that took into account confidence in peptide identification to improve quantification accuracy. We demonstrated the accuracy and specificity of the algorithm using serial mixed species datasets and stable isotope labelling using peptides obtained from multiple human donors.

Characterisation of complex samples is notoriously complex due to ionisation suppression and detection masking by peptides with similar m/z ratios often leading to missing values and underestimation of larger fold changes (this study and Goeminne et al. (2016)). The mixed species dataset used here represented an especially difficult case with most (~70%) peptides rendered unusable due to being shared between the two species. While all methods tended to underestimate

fold change magnitude (likely due to conflicting peptides in data-dependent analysis, exacerbated in mixed species analysis), we showed that BayesENproteomics was better able to confidently identify differentially abundant proteins and PTMs than simpler regression methods with greater accuracy in fold change estimates compared to OLS and LME-H regression models.

We used the iterative nature of MCMC processes to incorporate multiple imputation of missing values into protein fold change quantification. This eliminates the need for separate imputation and quantification steps and allows final fold change error estimates to take into account the intrinsic uncertainty regarding imputation of missing values (Schafer and Olsen, 1998). Here we employ fairly standard random imputation from prior distributions, but this could easily be substituted for any desired method.

Currently, BayesENproteomics does not utilise shared peptides, an approach that others have shown improves accuracy (Blein-Nicolas *et al*, 2012) at the cost of increased computation time. Use of empirical Bayes correction of variances (as in all methods tested here) can increase power by borrowing information from all proteins/PTMs without having to fit a model for all proteins simultaneously (as would be required to allow the use of shared peptides) for the purposes of calculating significance. MCMC samplers are known to be particularly computationally intensive. On an especially rich dataset, such as those generated by a modern Q Exactive HF mass spectrometer, analysis using BayesENproteomics can take several hours to process on a standard desktop computer. This is approximately three times longer than LME-H and several times longer than OLS regression. Importantly, the use of Gaussian/Laplacian priors for estimating β parameters presumed that logged fold changes lay on an approximately normal distribution. If it was suspected that this was not the case, BayesENproteomics may not be a suitable method of analysis.

Linear models are extremely customisable, and indeed, can require significant tailoring to best analyse complex multivariate experiments. Without sufficient background knowledge it can be very difficult to anticipate all potential sources of variability that need to be accounted for in any putative statistical model. Nevertheless, BayesENproteomics showed negligible perturbation by inclusion of

potentially extraneous parameters and so is well-suited for complex experimental setups that may include different batches, fractions, serial extractions, instruments and donors. Regularisation hyperparameters for additional effects can be sampled from the conditional distributions detailed here.

Finally, we extended our linear modelling pipeline to encompass pathway analysis. In transcriptomics experiments it is common to achieve coverage of the entire transcriptome (> 20000 transcripts in a given human cell/tissue), where hundreds to thousands may be differentially regulated in response to any given treatment. These large numbers make over-representation analysis using χ^2 or Fisher's exact tests well-suited to analysing differentially regulated transcriptomic processes. In contrast, proteomics experiments typically cover only a fraction of the total proteome (< 3000 proteins out of a potential > 20000 in a given human cell/tissue, increased further when considering multiple proteoforms) where only a hundred or fewer proteins may be differentially regulated, making it difficult to discern statistically significant process enrichment and erroneously giving the impression that "nothing is happening", often despite obvious morphological or metabolic changes assessed by other methods. Attempting to model pathway-level changes as a function of component parts for transcriptomics has previously shown that incorporating fold change estimates can increase specificity for discriminating pathway-level changes (Ozerov *et al*, 2016). Although, it should be noted that the aim with PALM is not biomarker discovery (for which more stringent statistical cut-offs are needed) but rather to simply distil a complex dataset with many proteins down to fewer, more readily understandable pathways by assessing what processes are represented and what they do in response to a given treatment. This works similar to previous enrichment tools (e.g. PANTHER (Mi *et al*, 2017)) and is similar to previous implementations of Bayesian regression in proteomics (Henao *et al*, 2013), however here we opt to maintain interpretability by utilizing curated protein-pathway annotations when picking proteins to use in pathway-level model fitting. Here we chose to use Reactome pathway annotations (Fabregat *et al*, 2018), but alternative categorical annotations could be used if desired. We believe that PALM provides a way of compressing information from many protein/proteoform fold changes into fewer annotated pathway fold changes, aiding interpretation of complex, multi-dimensional datasets. PALM can be used as a standalone pathway analysis tool;

though it does assume protein fold changes are formatted according to the BayesENproteomics protein-level output format (see examples on the GitHub page).

With the improvement of MS technology, and the production of highly efficient unconstrained searching algorithms (Kong *et al*, 2017; Devabhaktuni *et al*, 2018), identification and quantification of increasingly rare PTM'd peptides, and thus their parent proteoforms, has become possible even without enrichment. Many quantification methods have discarded or otherwise diminished the influence of these potentially relevant peptides in bottom-up protein quantification. Here, the use of regularised interaction coefficients facilitated the reporting of differentially abundant PTMs and the exclusion of potentially misidentified peptides from protein (and subsequent pathway) fold change estimates. Global quantification of proteoforms represents an exciting avenue for understanding how cell function is regulated post-translationally. BayesENproteomics was able to account for differences in PTM'd peptide behaviour whilst increasing accuracy of protein quantification in benchmark datasets with limited observations and inter-donor variability.

Structured Methods

Reagents and Tools Table

Reagent/resource	Reference or source	Identifier or catalog number
Experimental models		
C57BL/6J (*M. musculus*) skin	Skin provided by Hardman group, mice originally from Jackson Lab	000664
Primary human mesenchymal stromal cells (isolated from patient bone marrow)	Wrightington Hospital, Wigan, UK	N/A
Chemicals, enzymes and other reagents		
PNGase F from Elizabethkingia meningoseptica	Sigma-Aldrich	P7367
$H_2^{18}O$	Sigma	329878
low-glucose DMEM with pyruvate	Thermo Fisher Scientific	11885084
fetal bovine serum	Labtech.com	
penicillin/streptomycin cocktail	Sigma-Aldrich	P4333
ammonium bicarbonate	Sigma-Aldrich	09830
sodium dodecyl sulfate	Sigma	L3771
sodium deoxycholate	Sigma	D6750
protease inhibitor cocktail	Sigma	P8340
phosphatase inhibitor cocktail	Sigma	P0044
dithiothretol	Sigma	D0632

immobilized trypsin beads	Perfinity (via Thermo Fisher Scientific)	60109-101
urea	Thermo Fisher Scientific	29700
calcium chloride dihydrate	Sigma	C3306
iodoacetamide	Sigma	I1149
trifluoroacetic acid	Sigma	61030
ethyl acetate	Sigma	34858
POROS R3 beads	Thermo Fisher Scientific	1133903
acetonitrile	Thermo Fisher Scientific	TS-51101
formic acid	Thermo Fisher Scientific	28905
Software		
Matlab (2012a or higher) with Bioinformatics and Statistics and Machine Learning toolboxes	The MathWorks Inc.	
BayesENproteomics (Matlab code)	This manuscript	https://github.com/VenkMallikarjun/BayesENproteomics
Progenesis QI for proteomics (v3.0)	Non-linear Dynamics	
Mascot Daemon	Matrix Science	http://www.matrixscience.com/
Scaffold (v3.0)	Proteome Software, Inc.	http://www.proteomesoftware.com/products/scaffold/

Methods and Protocols

Cell culture

For the mixed species benchmark experiment, primary human mesenchymal stem cells (MSCs) were isolated from the bone marrow (knee and hip) of a single donor with informed consent and ethical approval, using established protocols (Strassburg et al, 2010). All experiments were performed in accordance with relevant guidelines and regulations, and with National Research Ethics Service and University of Manchester approvals. MSCs were cultured on tissue culture treated polystyrene (TCTP) in low-glucose DMEM with pyruvate (Thermo Fisher Scientific) supplemented with 10% fetal bovine serum (FBS, Labtech.com) and 1% penicillin/streptomycin cocktail (PS, Sigma-Aldrich).

Preparation of primary human MSCs lysates

Cells were lysed in 25 mM ammonium bicarbonate (AB) buffer containing 1.1% sodium dodecyl sulfate (SDS, Sigma), 0.3% sodium deoxycholate (Sigma), protease inhibitor cocktail (Sigma), phosphatase inhibitor cocktails (Sigma). Six 1.6 mm steel beads (Next Advance) were added to the tube and samples were homogenised with a Bullet Blender (Next Advance) at maximum speed for 2 minutes. Homogenates were cleared by centrifugation (12 °C, 10000 rpm, 5 minutes).

Preparation of mouse skin lysates

Skin from 3 month old female C57BL/6J mice was a gift from the Hardman group (University of Manchester; all animal work was performed under UK Home Office and local ethics committee approval). Ventral skin was shaved and scraped, then dissected into 1 mm^3 pieces. 100 μL of 8 M urea (Fisher Scientific) in 25 mM AB containing 25 mM dithiothretol (DTT, Sigma), protease and phosphatase inhibitor cocktails (Sigma) was added to the tissue sections with six 1.6 mm steel beads (Next Advance). Samples were then homogenised with a Bullet Blender (Next Advance) at maximum speed for 3 minutes. Resulting homogenates were cleared by centrifugation (12 °C, 10000 rpm, 5 minutes).

Preparation of tryptic peptides

Immobilised trypsin beads (Perfinity Biosciences) were suspended in 150 μL of digest buffer (1.33 mM CaCl$_2$, Sigma, in 25 mM AB) with 50 μL of cell or tissue lysate and shaken at 1400 rpm overnight at 37 °C in a thermocycler. The resulting digest was then reduced by the addition of 4 μL of 500 mM DTT (Sigma, in 25 mM AB; 10 min. shaking at 1400 rpm at 60 °C) and alkylated by the addition of 12 μL 500 mM iodoacetamide (Sigma, in 25 mM AB; 30 min. shaking in the dark at room temperature). Immobilised trypsin beads were removed by centrifugation at 10000 rpm for 10 min. Supernatant containing reduced, alkylated peptides were transferred to 1.5 mL 'LoBind' Eppendorf tubes and acidified by addition of 5 μL 10% trifluoroacetic acid (TFA, Sigma) in water, and cleaned by two-phase extraction (3 x addition of 200 μL ethyl acetate, Sigma, followed by vortexing and aspiration of the organic layer). Peptides were desalted using POROS R3 beads (Thermo Fisher). Briefly, R3 beads were thoroughly mixed in 50 % (v/v) acetonitrile (Sigma; prepared in milliQ H$_2$O; elute solution) by vortexing, to a concentration of 10 mg/ml. For each sample, 120 μl of bead suspension was pipetted into a well of a 96-well filtration membrane plate. The plate was then centrifuged (200 x g, 1 minute, room temperature) and the run-through was collected into a separate 96-well plate. The beads were then resuspended in 50 μl elute solution and centrifuged again twice before repeating the procedure with 0.1 % TFA (prepared in milliQ H$_2$O; wash solution). The run-through from the collection plate was discarded and the R3 beads were resuspended in 100 μl of sample before centrifugation. This process was repeated until all sample had been added. Bead-bound peptides were washed twice in wash solution. The run-through was discarded and the collection plate was washed with elute solution. Peptides were then eluted by two rounds of suspension in 50 μl elute solution and centrifugation. Eluted peptides were then lyophilized. Peptide concentrations (measured by Direct Detect spectrophotometer, Millipore) in injection solution (5% HPLC grade acetonitrile, Fisher Scientific, 0.1% TFA in deionized water) were adjusted to 200 ng/μL prior to MS analysis.

Preparation of serial mixtures of human and mouse peptides

Primary human MSCs and mouse skin sections were prepared separately as described above. Tryptic peptides from each preparation were mixed in ratios of 3:1, 1:1 and 1:3 (human:mouse, one series for each human donor:mouse), as described previously in (Swift et al, 2013a).

Stable isotope labelling of peptides

Desalted peptides were resuspended in 100 mM AB dissolved in $H_2^{18}O$ with or without 5 U PNGase F (Sigma), to a final volume of 10 μL. Peptides were then incubated in a thermocycler (1400 rpm, 37 °C) for 5 hours before drying down to a minimal volume in a vacuum centrifuge and resuspension in injection solution prior to MS analysis.

Mass spectrometry (MS)

Digested samples were analysed by liquid chromatography (LC) coupled tandem MS (LC-MS/MS) using an UltiMate® 3000 Rapid Separation LC system (RSLC, Dionex Corporation) coupled to a Q Exactive HF (Thermo Fisher), peptide mixtures were separated using a multistep gradient from 95% A (0.1% formic acid, FA, Thermo Fisher, in water) and 5% B (0.1% FA in acetonitrile) to 7% B at 1 min, 18% B at 58 min, 27% B at 72 min and 60% B at 74 min at 300 nL/min, using a 75 mm x 250 μm i.d. 1.7 μm CSH C18, analytical column (Waters). Peptides were selected for fragmentation automatically by data dependent analysis.

Preliminary data analysis using Progenesis QI (Nonlinear Dynamics)

Spectra from multiple samples were automatically aligned using Progenesis QI with manual placement of vectors where necessary. Peak picking sensitivity was set to 4/5 and features with a charge greater than +4 or fewer than 3 isotopes were excluded from further analysis. In the mouse/human mixed-species experiment, remaining features were searched using Mascot (Matrix Science), against the SwissProt and TREMBL pan-mammalian database (non-human and non-mouse identifications were removed). The peptide database was modified to search for alkylated cysteine residues (monoisotopic mass change, +57.021 Da) as a fixed modification, with oxidized methionine

(+15.995 Da), hydroxylation of asparagine, aspartic acid, proline or lysine (+15.995 Da) and phosphorylation of serine, tyrosine, threonine (+79.966 Da) as variable modifications. For the PNGase F-treated experiment, the peptide database was modified to search for alkylated cysteine residues (+57.021 Da) as a fixed modification, with oxidized methionine (+15.995 Da), ^{18}O-deamidation of asparagine/glutamine (+2.988 Da) and ^{16}O-deamidation of asparagine/glutamine (+0.984 Da). A maximum of one missed cleavage was allowed. Peptide detection intensities were exported from Progenesis QI as Excel (Microsoft) spread sheets for further processing. Deamidated spectrum counts were performed in Scaffold (version 4, Proteome Software) using .DAT files from Mascot, following pre-processing in Progenesis QI as detailed above. We applied quantification calculations to proteins detected with >2 unique (by sequence) peptides. Peptides assigned to proteins with 'unreviewed' status in the UniProt database were reassigned to the most abundant 'reviewed' protein with sequence identity in the dataset. Peptides shared between different protein groups were excluded from subsequent analysis.

Multi-variate regression modelling

To compare different types of regression analysis we used three different types of linear regression algorithms: (**A**) ordinary least squares (OLS); (**B**) Linear mixed effects models with Huber residual weights (LME-H); and (**C**) BayesENproteomics. Further details and references for each method are provided in the following sections. All analysis was performed using code written for Matlab R2015a (The MathWorks Inc).

Imputation of missing values

For OLS and LME-H methods, imputation of missing values was performed using a single imputation strategy whereby missing values are randomly sampled from a Gaussian distribution, downshifted to accommodate previous observations showing that missing values are more likely to be missing due to low abundance (Goeminne *et al*, 2016). Briefly, for each peptide, minimum non-missing peptide intensities were averaged to model the centre of a normal distribution with a width equal to 0.3 times that peptide's σ (standard deviation). The mean (i.e. centre) of this normal distribution was negatively

shifted by 1.6σ to model low abundance missingness, as described by Tyanova et al. (2016) (albeit for protein-level imputation). Missing values were then randomly imputed from this downshifted normal distribution.

Previous observations, such as by Tyanova et al. (2016), have shown that imputation from a down-shifted Gaussian distribution may be more suitable for imputation of missing peptide intensities from low abundance proteins, which are likely to be "missing non-randomly" (MNR), than for missing values from high abundance proteins, which are more likely to be "missing at random" (MAR) (Goeminne *et al*, 2015). However, others have suggested that the type of "missingness" (MAR or MNR) that predominates within a given dataset should decide what imputation strategy is used. For BayesENproteomics, we employed a logistic regression-based model similar to that employed by Li et al. (2011) to discern whether a given missing value was MAR or MNR. Briefly, a model of "missingness" was first created to differentiate observations that were MAR from those that were MNR as in (3).

$$Z_{qi} = \log\left(\frac{R_{qi}}{1-R_{qi}}\right) = \theta_0 + X_f\theta_f + X_g\theta_g + \varepsilon_{qfg}. \tag{3}$$

Where R_q represents a binary vector for protein q with elements $r_1, \dots r_n$ denoting whether an observation was missing ($r_i = 1$) or not ($r_i = 0$) and Z_q represents the logit transform of R_q to enable estimation of regressor coefficients θ using linear regression. As $Z_{qi} \in \{-\infty, +\infty\}$, we set minimum and maximum values for Z_q to -10 (corresponding to a probability for observation i being missing, $r_i < 0.00005$) and 10 ($r_i > 0.99995$). X_f and X_g represent binary design matrices denoting the peptide (*f*) and treatment group (*g*) from which a given observation is derived. θ_j ($j = 1\dots,(p_f + p_g + 1)$) represents a $1 \times (p_f + p_g + 1)$ vector of regressor coefficients; p_f and p_g represent the number of elements in θ_f and θ_g, respectively. θ_f and θ_g denote whether observations from a given peptide or treatment correlate with values in R_q (i.e. whether probability of "missingness" increases when looking at intensities from particular peptides or from particular experimental treatment groups). The intercept term, θ_0 denotes the intrinsic probability of missingness for that protein. We assume that most missing

values in any proteomics experiment are MNR, an assumption that others have shown is valid (Karpievitch *et al*, 2012; Goeminne *et al*, 2016). Therefore, if $\theta_j > \theta_0$ and $\theta_j > 0$ we inferred that missing observations associated with θ_j were MNR, and MAR otherwise. MAR and MNR missing values were imputed as part of each Gibbs sampler iteration as in (4) and (5).

$$\frac{\widetilde{\boldsymbol{y}}_{(MAR)iq}}{w_{(MAR)i-1}} \sim MVN\left(\boldsymbol{X}_{(MAR)}\beta_{i-1}^T , \ \sigma_{i-1}^2\big(\boldsymbol{X}_{(MAR)}\boldsymbol{X}_{(MAR)}^T\big)^{-1}\right); \tag{4}$$

$$\frac{\widetilde{\boldsymbol{y}}_{(MNR)iq}}{w_{(MNR)i-1}} \sim TGauss\left(\begin{matrix}\boldsymbol{X}_{(MNR)}\beta_{i-1}^T , \ \sigma_{i-1}^2\big(\boldsymbol{X}_{(MNR)}\boldsymbol{X}_{(MNR)}^T\big)^{-1}, \\ a = min\{\boldsymbol{y}_{observed}\} - 2, b = \boldsymbol{y}_{\left\lceil\frac{MNR}{n}\right\rceil}\end{matrix}\right). \tag{5}$$

Where $\widetilde{\boldsymbol{y}}_{(MAR)iq}$ represents a vector of score-weighted (see below) MAR $\log_2$(intensity) values for protein *q* at the i^{th} iteration of the Gibbs sampler. β_{i-1}^T is the vector of parameter estimates and σ_{i-1}^2 is the residual variance from the *i-1*th iteration of the Gibbs sampler, respectively. Missing values in $\widetilde{\boldsymbol{y}}_{iq}$ are sampled according to the multivariate normal distribution (MVN) described in (4) (see also section 3.1 in Zeng et al. (2017)). MNR missing values were imputed from a truncated Gaussian (TGauss) with an upper limit determined by the percentile of observed $\log_2$ intensity values corresponding to the fraction of values that are deemed MNR ($\boldsymbol{y}_{\left\lceil\frac{MNR}{n}\right\rceil}$) from the missingness regression model in (3). In (5), a and b represent the lower and upper limits of the TGauss distribution. Thus, if 5% of observations for a given protein are determined to be MNR according to (3), then the cut-off for the truncated Gaussian is equal to the 5th percentile of observed $\log_2$ intensity values. New missing values were imputed for each iteration of the Gibbs sampler. The multiple imputation strategy implemented in BayesENproteomics thus accounted for the inherent uncertainty in imputation by basing the resulting β estimates (and subsequent hypothesis testing) on distributional estimates of missing values (Schafer & Olsen, 1998) rather than fixed point estimates as in OLS and LME-H, where single random samples could strongly influence individual protein fold change estimates.

Weighting of residuals based on confidence of peptide identification

Identification of PTM'd peptides was performed by the inclusion of variable modifications during peptide database searching. Inclusion of multiple variable modifications was found to increase the

peptide false discovery rate (FDR). The number of false-positive identifications was reduced by discarding peptides with low Mascot scores using a standard FDR cut-offs based on identification p-values. We employ a Benjamini-Hochberg FDR (Benjamini & Hochberg, 1995) cut-off of < 0.2.

BayesENproteomics also employed a novel heuristic outlier weighting scheme that weighted against outlier peptides (which may possess biologically relevant PTMs), particularly if confidence in their identification was low. In this case we used Mascot scores as our indicator for peptide identification confidence. Firstly, Mascot scores, $S_1...,S_n$ were scaled by dividing them by a modified Bonferroni-like cut-off (similar to that described on the Mascot website, http://www.matrixscience.com/help/interpretation_help.html, accessed 23/10/17) and adjusted so that all the highest scoring peptides were weighted equally, as in (6) with values between 0 and 1.

$$\tilde{S} = min\left\{1, \frac{S}{\left(-10\left(\log_{10}\left(1/(20N)\right)\right)-13\right)}\right\}, \text{ N = total peptides in dataset.} \tag{6}$$

During each iteration of the Gibbs sampler, an n-dimensional vector of weights, w with elements $w_1...,w_n$, was calculated using a variation of the automatic outlier detection and weighting method in (Ting *et al*, 2007). Initially, there was no *a priori* reason to exclude - or diminish the influence of - peptides that have passed the initial FDR screen. Instead, we opted to weight in favour of those peptides that either have high Mascot scores or low residuals (7). Transformed Mascot scores in $\tilde{S}$ were used to parameterize a binomial distribution giving $\hat{S}$ that would determine if observations from that were favourably weighted each Gibbs sampler iteration (8).

$$w_i - 1 - \hat{S_i}|\sigma^2, \ \hat{S_i} \sim Gamma\left(\hat{S_i} + \frac{1}{2}, \ \frac{1}{2} + \frac{1}{2\sigma^2}R_i^2\right), i = 1 \dots n; \tag{7}$$

$$\hat{S_i}|\tilde{S} \sim Binom\left(1, \ \tilde{S}\right). \tag{8}$$

Where $\boldsymbol{R}$ is a vector of residuals with elements $R_1 \dots R_n$, $R = \boldsymbol{y} - \boldsymbol{X\beta}$. Observations were then weighted by multiplying each row of $\boldsymbol{X}$ and each value of $\boldsymbol{y}$ by their respective weight calculated in (7), (9) and (10).

$$\widetilde{X}_{ij} = X_{ij}w_i, \ i = 1 \dots n, \ j = 1 \dots p; \tag{9}$$

$$\widetilde{y}_i = y_iw_i, \ i = 1 \dots n. \tag{10}$$

Where $\widetilde{X}$ and $\widetilde{y}$ are the weighted design matrix and response vector, respectively.

Linear regression implementation

Peptide-based linear regression modelling has previously been shown to possess greater statistical power and accuracy than summarization models when detecting differentially abundant proteins (Goeminne *et al*, 2015). Here we compare three different models for calculating differentially abundant proteins and PTMs:

Ordinary least squares (OLS):

Differential protein abundance was calculated using the simple linear model (Choi *et al*, 2014; Clough *et al*, 2009) shown in (1), using the fitlm Matlab function. For calculating differential PTM abundance, the $\log_2$ fold change for each PTM'd peptide was normalised to the $\log_2$ fold change calculated for parent protein abundance. In cases where a single PTM site was shared by 2 or more peptides (i.e. missed cleavages), the most abundant one was used.

Linear mixed-effects models with Huber residual weights (LME-H):

The ridge regression/mixed-effects algorithm developed by Goeminne et al. (2016), wherein peptides were modelled as random effects (i.e. they were assumed to be randomly sampled from a larger population and that they accurately modelled the variance of that population) using the more complex model shown in equation (2). Goeminne et al. (2016) exploited the link between ridge regression and mixed effects models to assign each β_j a specific penalty, λ_j (where $\lambda_j = \sigma_i^2/\sigma_{ij}^2$, σ_i^2 = residual variance of protein i, σ_{ij}^2 = variance of coefficient β_j for protein $i, j = 1, 2 \dots p$). We recapitulated this algorithm using the fitglme Matlab function, including Huber weighting of residuals. To calculate changes in PTM abundances, peptide:group and peptide:donor interaction effects ($\beta_{f:g}$, and $\beta_{f:d}$ in (2), respectively) were included as random effects with the size of the resulting interaction βs denoting changes in the abundance of that peptide in response to treatment or donor effects, respectively.

BayesENproteomics:

This method utilises a novel Bayesian linear regression algorithm with elastic net Regularisation based on the hierarchical model detailed by Kyung et al. (2010). Bayesian methods employ Regularisation based on the prior distribution parameters were estimated from, with elastic net Regularisation being equivalent to sampling from an intermediate Gaussian/Laplacian prior. Sampling was performed using a Gibbs sampler with the maximum number of iterations for each protein set at $25 \times runs + p_f$ (*runs* = total number of MS runs to be analysed, p_f = number of peptides), or 1,000 – whichever was higher – with half of these as burn-in iterations. The full hierarchical model for a single protein is detailed in equations (11)-(15).

$$\beta | \sigma^2, \tau^2, \lambda_2, \widetilde{\boldsymbol{X}}, \widetilde{\boldsymbol{y}} \sim MVN \left(\begin{array}{c} \left(\widetilde{\boldsymbol{X}}^T \widetilde{\boldsymbol{X}} + \boldsymbol{I}_p (\tau^{-2} + \lambda_2) \right)^{-1} \widetilde{\boldsymbol{X}}^T \widetilde{\boldsymbol{y}}, \\ \sigma^2 \left(\widetilde{\boldsymbol{X}}^T \widetilde{\boldsymbol{X}} + \boldsymbol{I}_p (\tau^{-2} + \lambda_2) \right)^{-1} \end{array} \right); \tag{11}$$

$$\sigma^2 | \beta, \tau^2 \sim IG \left(\frac{(n-1+p)}{2}, \ \frac{1}{100} + \frac{R^T R}{2} + \frac{(\lambda_1 \beta) D_{\tau^2}^{-1} \beta}{2} + \frac{\lambda_2 \beta^T \beta}{2} \right); \tag{12}$$

$$\tau_j^{-2} | \beta_j, \sigma^2, \lambda_1 \sim GIG \left(-\frac{1}{2}, \ \sqrt{\frac{\lambda_1^2 \sigma^2}{\beta_j^2}}, \ \lambda_1^2 \right). \tag{13}$$

Let τ^{-2} represent a vector of latent variables $(\tau_1^{-2} \dots \tau_p^{-2})$ – sampled from the generalised inverse Gaussian (GIG) distribution in (13) using the sampler in Makalic and Schmidt (2016) – for each β_j such that larger values of τ_j^{-2} (*j* = 1…, *p*) result in β_j being shrunk towards zero. $D_{\tau^2}^{-1}$ denotes a diagonal matrix with elements τ_j^{-2}, *j* = 1…, *p*. Residual variance (σ^2) is sampled from an inverse gamma (IG) distribution in (11). To enforce sparsity, we employ two Regularisation hyperparameters, λ_1 and λ_2, with different conditional distributions, specifying LASSO (14) and ridge hyperparameters (15), respectively. Notably, while overall covariance is controlled by λ_1 through its effect on τ_j^{-2}, each β_j is given its own L$_2$ Regularisation hyperparameter, λ_{2_j}, similar to the LASSO-like "horseshoe" estimator (Makalic & Schmidt, 2016). This leads to smaller coefficients (i.e. "noise") being more aggressively shrunk towards zero compared to larger coefficients (i.e. "signal"), compared to regression using scalar Regularisation hyperparameters that lead to constant shrinkage across all

βs. λ_1^2 and λ_2 are sampled from gamma distributions of the form $Gamma(a, b)$, with a posterior mean of $\frac{a}{b}$.

$$\lambda_1^2 | \tau^2 \sim Gamma\left(p, \ 1 + \frac{1}{2}\sum_{j=1}^{p} \tau_j^2\right); \tag{14}$$

$$\lambda_{2\,j} | \beta_j, \sigma^2 \sim Gamma\left(1, \ 3 + \frac{\beta_j^2}{2\sigma^2}\right). \tag{15}$$

Estimates of parameters were taken as means of Gibbs-sampled posterior distributions made from post-burn-in iterations.

Pathway analysis using linear modelling (PALM)

PALM using Reactome (Fabregat *et al*, 2018; Milacic *et al*, 2012) pathway annotations was performed as described above using logged fold changes (calculated from the three methods detailed above) as the response variable (y) according to the model for a given Reactome pathway:

$$y_{gp} = \beta_0 + X_g\beta_g + X_p\beta_p + \varepsilon_{gp}. \tag{16}$$

Where β_g and β_p denote effect sizes due to experimental treatment g and protein p. Residual weights (w) were set to $min\left(1, \frac{|y_i|}{SE_i}\right)$, where SE_i ($i = 1\dots n$) is the standard error of that protein fold change estimate, y_i. The code allows the user to limit estimation to only those pathways with a minimum observed number of proteins present in a given dataset; here, we set the minimum to be five.

Empirical Bayes correction of protein variances

All methods included Empirical Bayes correction for stabilising variance estimation (Kammers *et al*, 2015; Smyth, 2004). Empirical Bayes-modified t-tests were used to calculate p-values for determining significance of differentially abundant proteins and PTMs.

Data availability

Progenesis QI peptide ion outputs used in this study, along with BayesENproteomics Matlab code used here (with protein/pathway-level output files as .csv) are freely available from www.github.com/VenkMallikarjun/BayesENproteomics.

Acknowledgements

VM was supported by a studentship from the Sir Richard Stapley Educational Trust. JS was funded by a Biotechnology and Biological Sciences Research Council (BBSRC) David Phillips Fellowship (BB/L024551/1). Mass spectrometry was carried out at the Wellcome Trust Centre for Cell-Matrix Research (WTCCMR; 203128/Z/16/Z) Biological Mass Spectrometry Core Research Facility. The authors thank: Prof. Tim Board (Wrightington Hospital) for provision of tissue samples for MSC isolation; Dr. Charis Saville (University of Manchester) for provision of mouse tissue and Dr. Craig Lawless and Mr. Jack Llewellyn for constructive criticism on the manuscript.

Author contributions

VM designed and implemented the algorithm and developed the benchmark datasets used in this manuscript. VM, SMR and JS wrote the paper.

Conflicts of interest

The authors declare no competing interests.

References

Abaye DA, Pullen FS & Nielsen B V. (2011) Peptide polarity and the position of arginine as sources of selectivity during positive electrospray ionisation mass spectrometry. *Rapid Commun. Mass Spectrom.* **25:** 3597–3608

Bekker-Jensen DB, Kelstrup CD, Batth TS, Larsen SC, Haldrup C, Bramsen JB, Sørensen KD, Høyer S, Ørntoft TF, Andersen CL, Nielsen ML & Olsen J V (2017) An optimized shotgun strategy for the rapid generation of comprehensive human proteomes. *Cell Syst.* **4:** 587–599.e4

Benjamini Y & Hochberg Y (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B* **57:** 289–300

Blein-Nicolas M, Xu H, de Vienne D, Giraud C, Huet S & Zivy M (2012) Including shared peptides for estimating protein abundances: A significant improvement for quantitative proteomics. *Proteomics* **12:** 2797–2801

Bornn L, Gottardo R & Doucet A (2010) Grouping priors and the Bayesian elastic net. *Arxiv Prepr.***:** 1001.4083

Brodin P, Jojic V, Gao T, Bhattacharya S, Angel CJL, Furman D, Shen-Orr S, Dekker CL, Swan GE, Butte AJ, Maecker HT & Davis MM (2015) Variation in the human immune system is largely driven by non-heritable influences. *Cell* **160:** 37–47

Choi M, Chang C, Clough T, Broudy D, Killeen T, MacLean B & Vitek O (2014) MSstats: an R package for statistical analysis of quantitative mass spectrometry-based proteomic experiments. *Bioinformatics* **30:** btu305

Clough T, Key M, Ott I, Ragg S, Schadow G & Vitek O (2009) Protein quantification in label-free LC-MS experiments. *J. Proteome Res.* **8:** 5275–5284

Devabhaktuni A, Olsson N, Gonzalez CG, Rawson K, Swaminathan K & Elias JE (2018) Measuring proteomes with long strings: A new, unconstrained paradigm in mass spectrum interpretation. *bioRxiv Prepr.***:** 10.1101/282624

Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, Haw R, Jassal B, Korninger F, May B, Milacic M, Roca CD, Rothfels K, Sevilla C, Shamovsky V, Shorser S, Varusai T, Viteri G, Weiser J, Wu G, et al (2018) The Reactome pathway knowledgebase. *Nucleic Acids Res.* **46:** D649–D655

Forshed J (2013) Protein quantification by peptide quality control (PQPQ) of shotgun proteomics data. *Methods Mol. Biol.* **1023:** 149–58

Goeminne LJE, Argentini A, Martens L & Clement L (2015) Summarization vs peptide-based models in label-free quantitative proteomics: Performance, pitfalls, and data analysis guidelines. *J. Proteome Res.* **14:** 2457–2465

Goeminne LJE, Gevaert K & Clement L (2016) Peptide-level robust ridge regression improves estimation, sensitivity, and specificity in data-dependent quantitative label-free shotgun proteomics. *Mol. Cell. Proteomics* **15:** 657–68

Henao R, Thompson JW, Moseley MA, Ginsburg GS, Carin L & Lucas JE (2012) Hierarchical factor modeling of proteomics data. In *2012 IEEE 2nd International Conference on Computational Advances in Bio and medical Sciences (ICCABS)* pp 1–6. IEEE

Henao R, Thompson JW, Moseley MA, Ginsburg GS, Carin L & Lucas JE (2013) Latent protein trees. *Ann. Appl. Stat.* **7:** 691–713

Huang DW, Sherman BT & Lempicki RA (2009) Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* **37:** 1–13

Ivanovska IL, Swift J, Spinler K, Dingal D, Cho S & Discher DE (2017) Cross-linked matrix rigidity and soluble retinoids synergize in nuclear lamina regulation of stem cell differentiation. *Mol. Biol. Cell* **28:** 2010–2022

Kammers K, Cole RN, Tiengwe C & Ruczinski I (2015) Detecting significant changes in protein abundance. *EuPA Open Proteomics* **7:** 11–19

Karpievitch Y V, Dabney AR & Smith RD (2012) Normalization and missing value imputation for label-free LC-MS analysis. *BMC Bioinformatics* **13:** S5

Kong AT, Leprevost F V., Avtonomov DM, Mellacheruvu D & Nesvizhskii AI (2017) MSFragger: Ultrafast and comprehensive peptide identification in mass spectrometry-based proteomics. *Nat. Methods* **14:** 513–520

Kyung M, Gilly J, Ghoshz M & Casellax G (2010) Penalized regression, standard errors, and Bayesian lassos. *Bayesian Anal.* **5:** 369–412

Lawless C & Hubbard SJ (2012) Prediction of missed proteolytic cleavages for the selection of surrogate peptides for quantitative proteomics. *OMICS* **16:** 449–456

Li F, Nie L, Wu G, Qiao J & Zhang W (2011) Prediction and characterization of missing proteomic data in Desulfovibrio vulgaris. *Comp. Funct. Genomics* **2011:** 780973

Makalic E & Schmidt DF (2016) A simple sampler for the horseshoe estimator. *IEEE Signal Process. Lett.* **23:** 179–182

Malioutov D, Chen T, Jaffe J, Airoldi E, Carr S, Budnik B & Slavov N (2017) Quantifying homologous proteins and proteoforms. *bioRxiv Prepr.***:** 10.1101/168765

Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D & Thomas PD (2017) PANTHER version 11: Expanded annotation data from Gene Ontology and Reactome pathways, and data analysis tool enhancements. *Nucleic Acids Res.* **45:** D183–D189

Milacic M, Haw R, Rothfels K, Wu G, Croft D, Hermjakob H, D'Eustachio P & Stein L (2012) Annotating cancer variants and anti-cancer therapeutics in Reactome. *Cancers (Basel).* **4:** 1180–1211

Ozerov I V., Lezhnina K V., Izumchenko E, Artemov A V., Medintsev S, Vanhaelen Q, Aliper A, Vijg J, Osipov AN, Labat I, West MD, Buzdin A, Cantor CR, Nikolsky Y, Borisov N, Irincheeva I, Khokhlovich E, Sidransky D, Camargo ML, Zhavoronkov A, et al (2016) In silico Pathway Activation Network Decomposition Analysis (iPANDA) as a method for biomarker development. *Nat. Commun.* **7:** 13427

Rodriguez J, Gupta N, Smith RD & Pevzner PA (2008) Does trypsin cut before proline? *J. Proteome Res.* **7:** 300–305

Schafer JL & Olsen MK (1998) Multiple imputation for multivariate missing-data problems: A data analyst's perspective. *Multivariate Behav. Res.* **33:** 545–571

Schliekelman P & Liu S (2014) Quantifying the effect of competition for detection between coeluting peptides on detection probabilities in mass-spectrometry-based proteomics. *J. Proteome Res.* **13:** 348–361

Smyth GK (2004) Linear models and empirical Bayes methods for assessing differential expression in microarray experiments. *Stat. Appl. Genet. Mol. Biol.* **3:** 1–25

Swift J, Harada T, Buxboim A, Shin J-W, Tang H-Y, Speicher DW & Discher DE (2013a) Label-free mass spectrometry exploits dozens of detected peptides to quantify lamins in wildtype and knockdown cells. *Nucleus* **4:** 450–9

Swift J, Ivanovska IL, Buxboim A, Harada T, Dingal PCDP, Pinter J, Pajerowski JD, Spinler KR, Shin J-W, Tewari M, Rehfeldt F, Speicher DW & Discher DE (2013b) Nuclear lamin-A scales with tissue stiffness and enhances matrix-directed differentiation. *Science* **341:** 1240104–1240104

Ting JA, D'Souza A & Schaal S (2007) Automatic outlier detection: A Bayesian approach. *Proc. - IEEE Int. Conf. Robot. Autom.*: 2489–2494

Tsai T-H, Hao Z, Hong Q, Moore B, Stella C, Zhang JH, Chen Y, Kim M, Koulis T, Ryslik G a., Verschueren E, Jacobson F, Haskins WE & Vitek O (2017) Statistical characterization of therapeutic protein modifications. *Sci. Rep.* **7:** 7896

Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M & Cox J (2016) The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat. Methods* **13:** 731–40

Webb-Robertson B-JM, Matzke MM, Datta S, Payne SH, Kang J, Bramer LM, Nicora CD, Shukla AK, Metz TO, Rodland KD, Smith RD, Tardiff MF, McDermott JE, Pounds JG & Waters KM (2014) Bayesian proteoform modeling improves protein quantification of global proteomic measurements. *Mol. Cell. Proteomics* **13:** 3639–3646

Wu R, Dephoure N, Haas W, Huttlin EL, Zhai B, Sowa ME & Gygi SP (2011) Correct interpretation of comprehensive phosphorylation dynamics requires normalization by protein expression changes. *Mol. Cell. Proteomics* **10:** M111.009654

Zeng IS, Lumley T, Ruggiero K & Middleditch M (2017) A Bayesian approach to multivariate and multilevel modelling with non-random missingness for hierarchical clinical proteomics data. *bioRxiv Prepr.*: 10.1101/153049

Zou H & Hastie T (2005) Regularization and variable selection via the elastic-net. *J. R. Stat. Soc.* **67:** 301–320

Figures

Figure 1:

Figure 1. Extraction and preparation of mouse and human proteins for use in label-free quantification using multiple regression models.

(**A**) Schematic diagram of the protocol used to prepare samples for the mixed species benchmark experiment.

(**B**) Flow diagram showing the effect of Regularisation on estimated parameters used in fitting models.

(**C**) Schematic representation of how the Gibbs sampler arrives at the final (posterior) distribution for a single parameter. As the sampler proceeds, it will ideally converge on a single posterior distribution, as progressive iterations are biased towards areas of higher likelihood. Iterations from before this point are typically discarded ("burn-in" iterations) as those samples will not come from the target posterior distribution. Each parameter is then summarized from its respective estimated distribution (green and blue distributions) and all parameters lie on the target posterior distribution (grey distribution).

(**D**) Example traces from post-burn-in samples to build posterior β distributions corresponding to $\log_2$ (protein abundance) estimates. Traces show good mixing with no bias towards particular regions of their respective posterior distributions.

Figure 2:

Figure 2. Comparison of missing value imputation strategies.

(**A**) Histograms allowing a comparison of distributions of observed peptide intensities from both mouse and human proteins.

(**B**) Peptide intensities from presumed non-random missing value imputation (Tyanova et al, 2016).

(**C**) Peptide intensities from adaptive multiple imputation.

(**D**) Imputation on a synthetic incomplete dataset using DGD single imputation, compared to known values from the complete dataset.

(**E**) Imputation on a synthetic incomplete dataset using AMI, compared to known values from the complete dataset.

Estimated $\log_2$ fold changes in (**D**) and (**E**) calculated using BayesENproteomics, actual $\log_2$ fold change = -2. R indicates Pearson's correlation coefficients.

Figure 3:

Figure 3. BayesENproteomics increases specificity compared to other multi-variate regression methods.

(**A**) Volcano plots showing quantification of fold change estimates (x-axis) vs. significance (y-axis) observed in pairwise comparisons between three technical replicates.

(**B**) Volcano plots showing quantification of mouse proteins across all comparisons in mixed human:mouse benchmark experiment.

(**C**) Volcano plots showing quantification of human proteins across all comparisons in mixed human:mouse benchmark experiment.

(**D**) Bar chart showing squared deviation from ground truth across all contrasts for the indicated dataset. Bars represent means ± standard error of squared deviations.

(**E**) Boxplots showing coefficients of variation (CVs) for all differentially abundant proteins in the mixed human:mouse benchmark experiment.

Dotted lines in (**A**)-(**C**) indicate Benjamini-Hochberg false discovery rate corrected p-value (BHFDR) = 0.05. OLS fit using simple model in equation (1), black; LME-H according to equation (2), red; BayesENproteomics as in (2), blue. n = 4 human donors, 4 mice. Donor effects in (1) and (2) were excluded in analysis of mouse proteins. Numbers of significantly differentially abundant proteins are indicated in the top right of each plot, with number in brackets denoting those following the direction of the expected fold change.

Figure 4:

Figure 4. PNGase-catalysed stable isotope labelling quantified by summarization with normalisation and multivariate regression methods.

(**A**) Schematic diagram illustrating experimental setup for the variable PTM benchmark dataset.

(**B**) The number of deamidated peptides was increased following 5 hours PNGase F treatment. Only peptides from proteins identified by >1 peptide were counted.

(**C**) Histograms showing fold changes in N-linked ^{18}O labelled N-linked deamidation (yellow) and ^{18}O labelled Q-linked deamidation (green).

(**D**) Scatter plots comparing protein and PTM $\log_2$ fold changes for the indicated quantification methods. n = 5 human donors. Indicated p-values show significance from comparison of N-linked (yellow) and Q-linked (green) histograms calculated by Welch's two-sample t-test (p^{Welch}) or Kruskal-Wallis test (p^{KW}).

Figure 5:

Figure 5. Pathway analysis using linear models (PALM) on mixed-species dataset.

(**A**) Schematic diagram showing hierarchical tree structure for linear model assembly from MS1 intensities (far left) to pathways (far right).

(**B**) and (**C**) Volcano plots showing significantly differentially regulated pathways calculated from mouse:human mixed species datasets from OLS (black), LME-H (red) and BayesENproteomics (blue) protein-level output with subsequent PALM pathway analysis. Each dot represent a single Reactome pathway. Pathways follow the expected direction of the component proteins, though confidence in original protein-level fold change estimates may affect exact magnitudes of pathway-level fold changes. Only pathways with ≥5 proteins present in the dataset are shown.

Figure 6:

Pathway Analysis using Linear Modeling (PALM)

Figure 6. Summary schematic of complete BayesENproteomics and PALM workflow, including missing value imputation.