Inversion frequencies and phenotypic effects are

modulated by the environment

Emma Berdan[1], Hanna Rosenquist[2], Keith Larson[3], Maren Wellenreuther[4,*]

[1]Department of Marine Sciences, University of Gothenburg, Gothenburg,

Sweden

[2] MEMEG, Ecology Building, Lund University, Lund, Sweden

[3]Department of Ecology and Environmental Sciences, Climate Impacts Research

Centre, Umeå University, Umeå, Sweden

[4]The New Zealand Institute for Plant & Food Research Ltd, Auckland, New

Zealand

*Corresponding author: Maren Wellenreuther, The New Zealand Institute for

Plant & Food Research Limited, Box 5114, Port Nelson, Nelson 7043, New

Zealand.

tel.: +64 3 989 7658; fax: +64 3 546 7049; e-mail:

maren.wellenreuther@plantandfood.co.nz

Running title: karyotype x environment effects

Word count: 4,511

Abstract

Understanding how environmental variation drives phenotypic diversification within species is a major objective in evolutionary biology. The seaweed fly *Coelopa frigida* provides an excellent model for the study of genetically driven phenotypes because it carries an α/β inversion polymorphism that affects body size. *Coelopa frigida* inhabits highly variable beds of decomposing seaweed on the coast in Scandinavia thus providing a suitable test ground to investigate the genetic effects of substrate on both the frequency of the inversion (directional selection) and on the phenotype (genotype x environment effects). Here we use a reciprocal transplant experiment to test the effect of the α/β inversion on body size traits and development time across four suitable natural breeding substrates from the clinal distribution. We show that while development time is unaffected by GxE effects, both the frequency of the inversion and the relative phenotypic effects of the inversion on body size differ between population x substrate combinations. This indicates that the environment modulates the fitness as well as the phenotypic effects of the inversion karyotypes. It further suggests that the inversion may have accumulated qualitatively different mutations in different populations that interact with the environment. Together our results are consistent with the idea that the inversion in *C. frigida* likely evolves via a combination of local mutation, GxE effects, and differential fitness of inversion karyotypes in heterogeneous environments.

Keywords: Coelopa frigida, inversion polymorphism, adaptation, GxE interactions, frequency effects

Introduction

One of the central goals of evolutionary biology is to understand how organisms cope with environmental heterogeneity (Savolainen *et al*, 2013). Phenotypic plasticity can solve this problem but the phenotypic response to variation in an environmental variable (i.e. phenotypic plasticity) may vary among species as well as among genotypes of the same species. This is commonly referred to as a Genotype-by-Environment interaction (GxE, Falconer, 1990; Lynch and Walsh, 1998; Via and Lande, 1985). Abundant experimental data and theoretical work now provide strong support for the idea that the relative performance of different genotypes across environmental gradients may be involved in the maintenance of phenotypic plasticity and genetic variation as well as the evolution of fitness related phenotypes (Carreira *et al*, 2006; Ungerer *et al*, 2003).

The study of environmental heterogeneity has been championed by work on geographic patterns along clines, with classic examples coming from genetic clines in flies almost a century ago (Dobzhansky, 1948; Sturtevant, 1921). Other examples include the genetic underpinnings of traits involved in adaptation to high altitude (Storz, 2002) and pigmentation (Hoekstra and Coyne, 2007). Clines hold a large attraction as natural Darwinian laboratories because they allow researchers to quantify the gradual effects of changing environments on

genotype, phenotype, and the interaction between them (Endler, 1977). These

studies have shown that while many environmental tolerance traits that vary

clinally are quantitative and polygenic (Wellenreuther and Hansson, 2016), they

can also be located within chromosomal segments that segregate as a single

locus (Tigano and Friesen, 2016). If environmentally adaptive variants are

chromosomally linked, either because they are found on chromosomal inversions

or are in areas of low recombination, their selective coefficients act in an additive

manner and they may be less likely to be lost via genetic drift and more likely to

contribute to rapid adaptation (Kirkpatrick and Barton, 2006). In the extreme

case, complex adaptations may be accomplished through the fixation of linked

sets of adaptive variants (Yeaman, 2013), often called supergenes, a case in

which polygenic adaptation may be more accurately modeled with single-locus

models (Schwander *et al*, 2014). Good examples for such single-locus

supergene adaptations come from *Drosophila melanogaster* where inversion

polymorphisms produce latitudinally varying phenotypes for traits including heat

and cold tolerance and body size (Reinhardt *et al*, 2014; Schrider *et al*, 2016).

Chromosomal inversion polymorphisms are being identified at a fast rate from an

increasing number of populations of organisms, e.g. insects, plants, bacteria and

humans, and are commonly linked to trait and fitness variation (Hoffmann and

Rieseberg, 2008; Kirkpatrick and Barton, 2006). Inversions are particularly

common in flies, and an intriguing mystery is why there are huge differences in

levels of polymorphism and rates of fixation for inversions between closely

related species and even between chromosomes in *Drosophila* spp. (Kirkpatrick,

2010). The seaweed fly *Coelopa frigida* (family Coelopidae) and its sister species

C. nebularum have a large chromosomal α/β inversion polymorphism on

chromosome I encompassing about 10% of the genome and containing ~100

genes (Aziz, 1975). The inversion polymorphism is known to affect a wide range

of life history and reproductive traits, such as development time (Day and

Buckley, 1980), body size (Butlin *et al*, 1982a), longevity and female fecundity

(Butlin and Day, 1984a; Butlin *et al*, 1982c), female resistance to male mating

attempts (Gilburn and Day, 1994; Gilburn *et al*, 1992) and also male's ability to

overcome female resistance (Butlin *et al*, 1982a).

The most striking of these phenotypic traits is a three-fold size difference in

males: male homokaryotypes for the inversion type α are much larger (~9mm)

than β- homokaryotypes (~3mm), whereas the heterokaryotype αβ is

intermediate in size (~6mm) (Wellenreuther pers. obs., Butlin and Day, 1984b).

Female size is not significantly affected by karyotype (Butlin and Day, 1984b).

Maintenance of the polymorphism is secured by strong heterosis (Collins, 1978),

since αβ-heterokaryotypes have higher fitness than either homokaryotype (Butlin

and Day, 1984a; Butlin and Day, 1989). The heterokaryotype αβ has a higher

viability during development than either homokaryotype, both in experiments

(Butlin and Day, 1984a; Leggett *et al*, 1996) and in the field, as indicated by

heterokaryotype excess (Butlin *et al*, 1982c; Day *et al*, 1983; Day *et al*, 1982b;

Gilburn and Day, 1994). Heterokaryotype excess increases under interspecific

competition with the congener *C. pilipes* and under intense intraspecific

competition (Butlin and Day, 1984a; Leggett *et al*, 1996). The effect is

disproportionally strong in males, especially decreasing the frequency of αα-

homokaryotypes (Butlin and Day, 1984a), possibly indicating lower viability of

males of this karyotype. However, inversion frequencies also form a cline from

the North Sea to the Baltic Sea, with the α inversion becoming more rare towards

the Baltic Sea (Day *et al*, 1983).

Here we explore if the environment modulates inversion frequencies and

phenotypic effects in *C. frigida*. For this, we study *C. frigida* populations in

Scandinavia that show variation in karyotype frequencies (Day *et al*, 1983).

Specifically, we test the possible effects of the α/β inversion polymorphism on

body traits and development time across four suitable natural breeding

substrates using a fully balanced reciprocal transplant experiment. We also

elucidate the relative fitness of the inversion karyotypes on these varying

substrates by examining the frequencies of the three karyotypes in different

environments.

Materials and methods

Study species and sites

Coelopa frigida belongs to the group of acalyptrate flies which exclusively forage

on decomposing seaweed (Cullen *et al*, 1987) and is found all along the

seashores of Scandinavia (Mcalpine, 1991). This species plays a vital role in

coastal environment health by accelerating the decomposition of algae, allowing

for faster release of nutrients (Cullen *et al*, 1987), and are an important food

source for predatory coastal invertebrates and seabirds (Fuller, 2003; Summers

et al, 1990). *Coelopa frigida* has 5 chromosomes with inversions occuring on 3 of

them (Aziz, 1975). The largest of these is the α/β which occurs on chromosome I.

The α/β inversion occurs at stable frequencies in all populations so far studied in

Northern Europe (Butlin *et al*, 1982a), although there may be seasonal (Butlin,

1983) and clinal variation (Day *et al*, 1983).

Inversion frequencies of *C. frigida* form a cline from the North Sea to the Baltic

Sea, with the α inversion becoming more rare towards the Baltic Sea (Day *et al*,

1983). Such clines are classic signs of selection and this cline is associated with

a decrease in salinity and tidal range and associated changes in the composition

of the wrackbed (i.e. the species of algae present). As salinity decreases from

the North Sea to the Skagerrak and Kattegat regions (Figure 1) there is a

corresponding decrease of marine species of brown seaweeds such as *Fucus*

spp. and *Laminaria spp.* moving into the Baltic Sea, *Zostera* eelgrass and red

algae gradually displace the brown seaweeds. We selected four populations

(Kämpinge, Kåseberga, Mölle and Steninge) from the southern part of the cline

that differ in tidal range, salinity, and wrackbed composition (Figure 1, Table 1).

Reciprocal transplant experiment

We conducted a fully balanced reciprocal transplant experiment involving all four

locations, where each fly location was allowed to grow on its own wrackbed

substrate and the substrates of the remaining three locations. For each location,

fly larvae and wrackbed substrate were collected between February and May

2014. The wrackbed substrate was placed in labelled plastic bags and then

frozen to ensure that no viable eggs or larvae remained. Fly larvae from were

reared on 150g of their own substrate in aerated plastic containers until a

puparium was formed, and then kept in individual Eppendorf tubes (to ensure

virginity) containing a small piece of cotton soaked in 5% sucrose solution. When

a fly eclosed, the Eppendorf was moved to the fridge to slow down development

time. Over a maximum of ten days, adult flies were collected until a minimum of

25 couples could be set up per location. Eggs were collected from 14-15 mating

pairs per population and subsequently mixed to populate three replicates of each

population x substrate combination with 30 eggs each (4 populations x4

substrates x 3 replicates, each one with 30 eggs, total eggs for the experiment:

1440). Replicate pots were kept at 25° C with a 12/12 hour light:dark cycle and

each pot (190x110x70 mm) was aerated and contained 150 g of representative

seaweed. Pots were checked daily and all eclosing flies were sexed and then

stored at -20° C on the same day.

Morphological measurements

Morphological measurements were carried out on adult flies only. All flies were

weighed (Sartorius Quintix 124-1S microbalance) to the nearest 0.0001 grams.

The thorax width of each fly was measured with a digital calliper (Cocraft Digital

Vemier Caliper) at the widest place and to the nearest 0.01 mm. This

measurement was repeated twice and the average was used. Then, the left wing

and right middle leg were removed (if one side was damaged the other side was

used). The wing and leg were placed under a thin glass slide and placed under

the microscope and a picture was the taken using a Nikon Digital Sight DS-Fi2

mounted on a Nikon SM2800 microscope set to a magnification of two power.

The resulting image was labelled with an individual identifier. Measurements

were conducted blind to location and substrate type to avoid bias.

Molecular determination of inversion types

In *C. frigida* the two most common alcohol dehydrogenase (Adh) allozymes, Adh-

B and Adh-D are in complete linkage disequilibrium with the α and β forms of the

inversion (Day *et al*, 1982a). Gel electrophoresis was used to visualise the

alcohol-dehydrogenase allozymes associated with the two inversion types and

this allowed for straightforward genotyping of the inversion types. Starch gels

(12%) were prepared by combining 24.5 g potato starch (Sigma-Aldrich, S4251)

with 90 ml TEB-buffer, which was then shaken in a 500 ml Buchner flask until

dissolved. Heated water (90 ml dH$_2$O) was added to the Buchner flask, and

shaken until homogenized. The Buchner flask was then heated in the microwave

at intervals of 5-10 seconds, and shaken in between to evenly distribute the heat.

The heating continued until the mixture was boiling, after which the flask was

closed with a rubber plug and de-gassed using a tap-based vacuum pump until

the mixture was without visible air bubbles. It was poured onto an 18.5x10 cm

glass plate with a plastic frame and left to set for 30 minutes, after which it was

left in a 4^{O}C refrigerator overnight.

Fly samples were taken from storage in -20° C freezer and kept on ice during

sample preparation. Porcelain plates with 12 wells were placed on ice packs. On

each well, a fly was placed together with a small amount of carborundum powder

(Fisher Chemical, #C192-500) and one drop of cold dH_2O. The flies were then

homogenized using a glass rod. Once homogenized, 3x4 mm filter paper wicks

were placed in the wells to soak up the homogenate. The wicks were then placed

in the starch gel.

Starch gels were put in the electrophoresis chamber, which was connected to a

power supply (Consort E863). The two chambers of the electrophoresis machine

were filled with TEB-buffer. Two cloth wicks were soaked in TEB-buffer and

placed so that they touched evenly along each of the exposed sides of the gel,

while still connected to the buffer. An ice pack was placed on top of the gel and

the electrophoresis (350 V, 50mA) took 1.5 hours.

After electrophoresis, the gel was cut horizontally with a metal thread and only

the bottom half was used for staining. The agar overlay used for staining

consisted of 20 ml of 2% agar, 10 ml HCl-Tris buffer(pH 8.6), 6 ml propan-2-ol,

1.5 ml 3-(4,5-dimethylthiazolyl-2)-2,5- diphenyltetrazolium bromide (MTT), 1 ml

Phenazine methosulphate (PMS), and 10 ml Nicotinamide-adenine dinucleotide

(NAD). The stain was mixed and poured onto the cut surface of the gel. The

covered box with the gel was then carefully put in a dark heating cabinet at 37° C

for 40 minutes, or until the banding pattern was clear and then photographed.

Statistical analysis

We analyzed weight and development time (in days) in R (R Development Core

Team, 2015) using general linear models. Weight was highly correlated with all

other morphological characters (Supplemental table S1) so we only analyzed

weight data and development time in a statistical framework. Weight data was

normally distributed and so was analyzed using a Gaussian distribution.

Development time is essentially count data and right skewed, so we analyzed the

data using a Poisson distribution. For both data sets we considered individual

flies to be our unit of replication and used the lme4 package in R (Bates *et al*,

2014) to construct a model with all factors (population, substrate, sex, and

karyotype) and all two and three way interactions. We also included pot

(replicate) nested within population*substrate as a random factor. We ran the

'Anova' function in the R package car (Fox and Weisberg, 2011) on the full

models and then examined the F-values for each term. Starting with a model with

only an intercept we built up a series of nested models adding a single main

effect, two or three way interaction at a time by decreasing F-value (i.e. starting

with the term with the highest F-value). We then used the 'Anova' function with

the χ^2 distribution to compare the models in a nested fashion (i.e. each model is

only compared to the one above it) and used a χ^2 test to determine if they were

significantly different. We selected the model with the lowest AIC whose nested

models were not significantly better.

To get an overview of our results we also visualized the morphological

measurements with a Principal Components Analysis (PCA). The PCA was run

on the four highly correlated morphological traits: weight, thorax width, leg and

wing length using the 'prcomp' function in R (R Development Core Team, 2015).

Visual inspection of our morphological data indicated that males were more

variable than females. We formally tested for differences in variability in a

statistical framework using Krishnamoorthy and Lee's (2014) modified signed-

likelihood ratio test (MS-LRT) implemented in the R package cvequality (Marwick

and Krishnamoorthy, 2016). We used a Bonferroni correction to account for

multiple testing (Bland and Altman, 1995).

We further examined the relative proportions of the three karyotypes (αα, αβ, and

ββ) in all population x substrate combinations. For these analyses we used pot

as our unit of replication. We tested for differences in two ways. First, we tested

for karyotype differences using a χ^2 test on a 3 karyotypes x 16 population-

substrate combination matrix. Second, we used a Dirichlet regression model in

the R package DirichletReg (Maier, 2014). The Dirichlet regression model allows

us to analyze karyotypes (variables with a bounded interval that sum up to a

constant - in our case frequencies of karyotypes in a population) exhibiting

skewness and heteroscedasticity, without having to transform the data (Maier,

2014). We used a nested approach starting with a simple model and then adding

factors. Models were tested against the one above them in a nested fashion

using the 'Anova' function in the R package car (Fox and Weisberg, 2011) to

determine the best model.

Results

Phenotypic data

We had different sample sizes for different treatment combinations (see figure

legends for Figures 3 and 4) due to differential survival of the three karyotype

categories. Three of the 96 combinations had zero flies (ex: αα Kämpinge

females in Kämpinge) but we do not expect that this unduly influenced our results

as we did not investigate any 4 way interactions in any of our statistical models

and thus did not examine these three combinations directly. All morphological

characteristics were highly and significantly correlated with each other

(Supplementary Table S1). To get an overall view of the variation in

morphological traits we conducted a PCA using all four morphological variables

(wing length, leg length, thorax width, and weight). The first principal component

explained 93.5% of the variance and the second principal component explained

6.5% of the variance (Figure 2).

We only examined weight, as this morphological trait is a good proxy of the other

morphological measurements (Pearson correlation coefficients with other traits

varied from 0.78-0.89, Supplemental Table S1) while still being an easily

tractable and understandable measurement. Weight was not strongly correlated

with development time (Pearson's product-moment correlation: 0.10) and

therefore development time was examined separately. F-values from the full

models indicated that effect sizes were much larger for weight than development

time (Table S2). The best model for weight included all main effects, the two-way

interactions: sex*karyotype, population*sex, population*karyotype,

population*substrate, substrate*karyotype, and the three-way interaction:

location*sex*karyotype. Comparing the best model with the next most simplified

model gave a χ^2 value of 43.3 (P = 0.00073). The best model for development

time included the main effects and the two-way interaction karyotype*sex.

Comparing the best model with the next most simplified model gave a χ^2 value of

8.6 (*P* = 0.0135). The best models for both weight and development time

included all main effects and a subset of interactions (Table 2). However, a GxE

interaction (as evidenced by a population x substrate interaction) was only

present for morphological characteristics, but not for development time.

For all measured traits, the Krishnamoorthy and Lee's (2014) modified signed-

likelihood ratio test showed that males had significantly more phenotypic

variation than females (Table 3). This variation could be partially accounted for

by karyotype. Previous work has shown that males are on average bigger than

females and that within male size differences are primarily driven by the αβ

inversion system: αα males are the biggest followed by αβ and ββ (Butlin *et al*,

1982c). This held true in our male data set for all morphological traits (Figure 2)

but the magnitude of these differences depended on both population and

substrate type (Figure 3, Supplementary Figures 2-5). Although the relative sizes

and size differences of the karyotypes differed between combinations of location

and substrate, the pattern of both leg length and wing length, thorax width, and

weight (Figure 3, Supplementary Figures 2-5) in relation to karyotype and sex

was clear and consistent for males, but much less pronounced for females. For

example, while the morphological PCA groups of the different male karyotypes

formed distinct yet overlapping clusters, a clustering of female karyotypes was

absent (Figure 2). Likewise, across treatments (combination of population and

substrate), αα-males had the largest mean size, followed by αβ-males and ββ-

males in all different population x substrate combinations (Figure 3). For females,

less than half of the combinations of population and substrate followed the same

pattern.

Development time did not follow the same pattern as observed for morphology

(Figure 4). While αα-males generally took longer to develop, this was not always

the case for every combination of population and substrate. For example, ββ flies

from Steninge, raised on Kämpinge substrate took longer to develop than any

other flies from Steninge. For females, MS-LRT demonstrated that there was

considerably less variation between karyotypes compared to males and patterns

varied substantially between populations and substrates.

Inversion types

With the exception of Steninge, frequencies of the karyotypes rarely matched

Hardy-Weinberg expectations based on previously estimated allele frequencies

(Figure 5). The frequencies of αα, αβ, and ββ were significantly different across

the Population x Substrate combinations (χ^2 = 71.5, df=30, P < 0.0001).

Generally αβ was the most frequently observed karyotype, but ββ could be just

as frequent or more frequent depending on the substrate. In ¾ of the populations

(all except Steninge) ββ was dominant in at least one substrate and that was

usually the substrate from Mölle.

We used Dirichlet regression to see which parameters best predicted the

karyotype frequencies. With this model we were not able to examine F-values so

we compared all possible models with the 'Anova' function from the R package

car (Fox and Weisberg, 2011). The best model was population + substrate +

population*substrate. Comparing this with the next most simplified model gave a

χ^2 value of 73.99 (P = 0.0002).

Discussion

Our results demonstrate that the environment modulates both the frequencies of

the karyotypes as well as their phenotypic effects in *C. frigida*. The presence of a

significant GxE term for both of these indicates that there is population specific

variation in larval fitness of the different inversion types as well as in the

phenotypic effects of the inversion. The same pool of eggs was used to seed

every substrate treatment per population so the neutral expectation (without

sampling bias) would be similar karyotype frequencies across substrates.

Instead, we found strong differences in karyotype frequencies between

substrates with some trends being consistent across multiple populations (Figure

5). While it is possible that these results are due simply to sampling bias, it is

unlikely that sampling bias would produce consistent trends. Furthermore, this

result confirms previous research showing that the substrate composition has a

strong selective effect on the inversion (Butlin *et al*, 1982b; Day *et al*, 1983;

Edward and Gilburn, 2013). It is likely that population specific variation has

accumulated in the inversion, as not all trends were consistent across

populations. For example, ββ was the most common karyotype on Mölle wrack

for every population except Steninge (Figure 5). Instead, in Steninge, frequencies

were often close to Hardy-Weinberg expectations and little variance was seen

between substrate combinations compared to other populations, possibly

because heterosis was stronger in Steninge than in the other populations.

Our results also indicate that inversion-related size effects in *C. frigida* are

affected by the environment, particularly in males as females show little to no

effect of karyotype on size. Male fly inversion phenotypes from the same

population could vary two-fold for weight when grown on the same substrate (e.g.

Steninge population on Mölle substrate), and three-fold when comparing male

inversion phenotypes across all four growing substrates (e.g. Steninge, Figure 3).

Male size has a high heritability with virtually all the variance in size being

attributable to the chromosomal inversion system (Wilcockson *et al*, 1995). In all

of our treatment combinations, environmental effects never swamped the effect

of the inversion. Male αα were always the biggest, followed by αβ and ββ, which

is consistent with previous work (e.g. Butlin *et al*, 1982a). Population and

substrate changed the magnitude of these differences as well as size averages

but never the pattern of the inversion size effects on males. In contrast to the

strong environment and inversion effects on male size, the inversion exhibited

negligible size effects in females. Specifically, our data show much lower

between inversion variation in size and considerably less variation across

environments for females. A similar difference in effect size between males and

females has also been found in a common garden study on *C. frigida* grown on

artificially mixed substrates containing in excess either the seaweed genus

Laminaria or *Fucus* (Edward and Gilburn, 2013). The study by Edward and

Gilburn (2013) also found that flies grown on an excess of *Laminaria* showed

greater among inversion size differences compared to *Fucus*, suggesting that the

type of seaweed itself, or the associated microbiome of the seaweed, causes

phenotypic differences in size. The wrackbed composition differs markedly along

the populations studied, as a consequence of the declining salinity from the North

Sea to the Baltic Sea (see Figure 1 and 2 in Wellenreuther *et al*, 2017), and have

likely been major selective environmental factors in modulating *C. frigida*

populations. Development time, another trait known to be affected by inversion

type, also showed significantly higher variation across male karyotypes, but

exhibited less extreme environmentally induced effects compared to size, and

consequently no significant GxE effects. These results together with previous

studies (Butlin and Day, 1984a; Day *et al*, 1980) underscore that the inversion

related phenotype effects in *C. frigida* are significantly affected by sex, with male

phenotypes being more labile compared to female phenotypes.

Interactions between genotypes and the environment are commonly observed in

quantitative traits (Gupta and Lewontin, 1982), particularly those associated with

fitness (e.g. Fanara *et al*, 2006), and are thought to provide a potent force for

maintaining genetic variation (Fernández Iriarte and Hasson, 2000; Gillespie and

Turelli, 1989; Hedrick, 1986). The habitat used by *C. frigida* is heterogeneous

and the resulting local population structure is similar to Levene's migration model

of subdivided populations, namely by a single pool of individuals that mate at

random and breed on discrete and ephemeral resources. Environmental

fluctuations in seaweed substrate composition likely favors the persistence of

alternative rearrangements (i.e. the α/β inversion) containing sets of co-adapted

alleles via balancing selection (Carroll *et al*, 2001; Filchak *et al*, 2000; Weinig,

2000), with each set being optimal under different conditions (Dobzhansky,

1970). This hypothesis may account for the so called "supergene" idea

(Schwander *et al*, 2014) such as those determining mating types found in

heterostylous plants and wing mimicry polymorphisms found in *Heliconius*

butterflies (Thompson and Jiggins, 2014). Balancing selection resulting from

seasonal and stochastic temporal and spatial wrackbed changes may thus be a

significant contributor to maintaining the fitness related inversion variation in size.

Other examples for selection on fitness trade-offs come from introduced apple

Malus pumila and native hawthorn *Crataegus* spp. hosts which has led to

divergence in phenology and thermal physiology in the true fruit fly *Rhagoletis*

pomonella (Filchak *et al*, 2000).

Balancing selection has also been implicated in the maintenance of other

inversion polymorphisms, but whether balancing selection can maintain genetic

polymorphisms over long time scales is still debated (Bernatchez, 2016).

Theoretical studies indicate that balancing selection may primarily be a short-

term process because the genetic load that it creates then has a negative

feedback that hinders the long-term maintenance of adaptive alleles (Fijarczyk

and Babik, 2015). Furthermore, rapid environmental changes may cause

dramatic allele frequency shifts that could cause the loss of formerly stable

polymorphisms (Fijarczyk and Babik, 2015). Despite this, recent studies have

provided some evidence that balancing selection can operate even over longer

time scales. Lindtke et al. (2017) investigated the inversion related cryptic colour

polymorphism in *Timema cristinae* walking stick insects using a comprehensive

dataset of natural populations and population genetic analyses. They were able

to demonstrate that inversion colour variation has likely been maintained over

extended time periods by balancing selection, which has previously been thought

to be only possible over short time frames.

In conclusion, our results indicate that the inversion in *C. frigida* likely evolves via

a combination of local mutations, GxE effects, and differential fitness of

karyotypes in heterogeneous environments. Future work should apply high-

resolution genome-wide approaches that target the regions inside the inversion

to identify the true genic targets of balancing varying selection. This approach

would also elucidate which genes inside the inversion are linked to these large

size effects in males, and how the environment has shape the differentiation

between these genes along gradients.

Acknowledgements

MW would like to thank Roger Butlin and Andre Gilburn for helpful advice

regarding the molecular inversion determination and rearing techniques. EB

would like to thank Fabian Roger for helpful advice regarding the figures. Helena

Westerdahl and Jan Åke Nilsson gave some helpful comments on earlier

versions of this work. The work was supported by a grant from the Crafoord

Foundation, from Entomological Foundation in Lund and from the Swedish

Research Vetenskapsrådet Council (2012-3996) to MW. EB was supported by a

Conflict of Interest

450 The authors declare that they have no conflict of interest.

Data Archiving

453 All raw phenotypic data and karyotype frequencies will be deposited on Dryad

454 upon acceptance.

References

Aziz JB (1975). *Investigations into chromosomes 1, 2 and 3 of Coelopa frigida (Fab.)*. Thesis.

Bates D, Mächler M, Bolker B, Walker S (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:14065823*.

Bernatchez L (2016). On the maintenance of genetic variation and adaptation to environmental change: considerations from population genomics in fishes. *J Fish Biol* **89**(6)**:** 2519-2556.

Bland JM, Altman DG (1995). Multiple significance tests: the Bonferroni method. *Bmj* **310**(6973)**:** 170.

Butlin RK (1983). The maintenance of an inversion polymorphism in Coelopa frigida. University of Nottingham.

Butlin RK, Collins P, Skevington S, Day T (1982a). Genetic variation at the alcohol dehydrogenase locus in natural populations of the seaweed fly, *Coelopa frigida*. *Heredity* **48**(1)**:** 45-55.

Butlin RK, Collins PM, Skevington SJ, Day TH (1982b). Genetic-Variation at the Alcohol-Dehydrogenase Locus in Natural-Populations of the Seaweed Fly, Coelopa-Frigida. *Heredity* **48**(Feb)**:** 45-55.

Butlin RK, Day TH (1984a). The effect of larval competition on development time and adult size in the seaweed fly, *Coelopa frigida*. *Oecologia* **63**(1)**:** 122-127.

Butlin RK, Day TH (1984b). The Effect of Larval Competition on Development Time and Adult Size in the Seaweed Fly, Coelopa-Frigida. *Oecologia* **63**(1)**:** 122-127.

Butlin RK, Day TH (1989). Environmental correlates of inversion frequencies in natural populations of seaweed flies (*Coelopa frigida*). *Heredity* **62**(2)**:** 223-232.

Butlin RK, Read IL, Day TH (1982c). The effects of a chromosomal inversion on
adult size and male mating success in the seaweed fly, *Coelopa frigida*. *Heredity*
49(1)**:** 51-62.

Carreira VP, Soto IM, Hasson E, Fanara JJ (2006). Patterns of variation in wing
morphology in the cactophilic Drosophila buzzatii and its sibling D. koepferae. *J*
Evol Biol **19**(4)**:** 1275-1282.

Carroll SP, Dingle H, Famula TR, Fox CW (2001). Genetic architecture of
adaptive differentiation in evolving host races of the soapberry bug, Jadera
haematoloma. *Genetica* **112**(1)**:** 257-272.

Collins PM. (1978). The Genetical Society of Great Britain.

Cullen SJ, Young AM, Day TH (1987). Dietary requirements of seaweed flies
(Coelopa frigida). *Estuarine, Coastal and Shelf Science* **24**(5)**:** 701-710.

Day T, Dobson T, Hillier P, Parkin D, Clarke B (1982a). Associations of enzymic
and chromosomal polymorphisms in the seaweed fly, Coelopa frigida. *Heredity*
48(Pt 1)**:** 35-44.

Day TH, Buckley PA (1980). Alcohol dehydrogenase polymorphism in the
seaweed fly, *Coelopa frigida*. *Biochemical Genetics* **18**(7-8)**:** 727-742.

Day TH, Dawe C, Dobson T, Hillier PC (1983). A chromosomal inversion
polymorphism in Scandinavian populations of the seaweed fly, Coelopa frigida.
Hereditas **99**(1)**:** 135-145.

Day TH, Dobson T, Hillier PC, Parkin DT, Clarke B (1982b). Associations of
enzymic and chromosomal polymorphisms in the seaweed fly, Coelopa frigida.
Heredity **48**(1)**:** 35-44.

Day TH, Dobson T, Hillier PC, Parkin DTa, Clarke B (1980). Different rates of
development associated with the alcohol dehydrogenase locus in the seaweed
fly, Coelopa frigida. *Heredity* **44:** 321-326.

Dobzhansky T (1948). Genetics of natural populations. XVI. Altitudinal and
seasonal changes produced by natural selection in certain populations of
Drosophila pseudoobscura and *Drosophila persimilis*. *Genetics* **33**(2)**:** 158.

Dobzhansky TG (1970). *Genetics of the evolutionary process*. Columbia
University Press: New York.

Edward DA, Gilburn AS (2013). Male-specific genotype by environment
interactions influence viability selection acting on a sexually selected inversion
system in the seaweed fly, *Coelopa frigida*. *Evolution* **67**(1)**:** 295-302.

Endler JA (1977). *Geographic variation, speciation, and clines*. Princeton
University Press.

Falconer D (1990). Selection in different environments: effects on environmental
sensitivity (reaction norm) and on mean performance. *Genetics research* **56**(1)**:**
57-70.

Fanara J, Folguera G, Iriarte PF, Mensch J, Hasson E (2006). Genotype by
environment interactions in viability and developmental time in populations of
cactophilic Drosophila. *JEB* **19**(3)**:** 900-908.

Fernández Iriarte P, Hasson E (2000). The role of the use of different host plants
in the maintenance of the inversion polymorphism in the cactophilic Drosophila
buzzatii. *Evolution* **54**(4)**:** 1295-1302.

Fijarczyk A, Babik W (2015). Detecting balancing selection in genomes: limits
and prospects. *Molecular Ecology* **24**(14)**:** 3529-3545.

Filchak KE, Roethele JB, Feder JL (2000). Natural selection and sympatric
divergence in the apple maggot Rhagoletis pomonella. *Nature* **407**(6805)**:** 739-
742.

Fox J, Weisberg S (2011). *An R Companion to Applied Regression*, 2 edn. SAGE
Publications: Thousand Oaks, California.

Fuller RA (2003). Factors influencing foraging decisions in ruddy turnstones
Arenaria interpres (L.). Durham University.

Gilburn AS, Day TH (1994). Sexual dimorphism, sexual selection and the α β
chromosomal inversion polymorphism in the seaweed fly, *Coelopa frigida*.
Proceeding of the Royal Society of Biological Sciences **257**(1350)**:** 303-309.

Gilburn AS, Foster SP, Day TH (1992). Female mating preference for large size
in *Coelopa frigida* (seaweed fly). *Heredity* **69**(3)**:** 209-216.

Gillespie JH, Turelli M (1989). Genotype-environment interactions and the
maintenance of polygenic variation. *Genetics* **121**(1)**:** 129-138.

Gupta AP, Lewontin R (1982). A study of reaction norms in natural populations of
Drosophila pseudoobscura. *Evolution* **36**(5)**:** 934-948.

Hedrick PW (1986). Genetic polymorphism in heterogeneous environments: a
decade later. *Annual review of ecology and systematics* **17**(1)**:** 535-566.

Hoekstra HE, Coyne JA (2007). The locus of evolution: evo devo and the
genetics of adaptation. *Evolution* **61**(5)**:** 995-1016.

Hoffmann AA, Rieseberg LH (2008). Revisiting the impact of inversions in
evolution: from population genetic markers to drivers of adaptive shifts and
speciation? *Annu Rev Ecol Evol Syst* **39**(1)**:** 21-42.

Kirkpatrick M (2010). How and why chromosome inversions evolve. *PLoS Biol*
8(9)**:** e1000501.

Kirkpatrick M, Barton N (2006). Chromosome inversions, local adaptation and
speciation. *Genetics* **173**(1)**:** 419-434.

Krishnamoorthy K, Lee M (2014). Improved tests for the equality of normal
coefficients of variation. *Computational Statistics* **29**(1-2)**:** 215-232.

Leggett M, Wilcockson R, Day T, Phillips D, Arthur W (1996). The genetic effects
of competition in seaweed flies. *Biol J Linn Soc* **57**(1)**:** 1-11.

Lindtke D, Lucek K, Soria-Carrasco V, Villoutreix R, Farkas TE, Riesch R *et al*
(2017). Long-term balancing selection on chromosomal variants associated with
crypsis in a stick insect. *Mol Ecol* **26:** xxx-xxx.

Lynch M, Walsh B (1998). *Genetics and analysis of quantitative traits*, Vol 1.
Sinauer Sunderland, MA.

Maier MJ (2014). DirichletReg: Dirichlet regression for compositional data in R.
Research Report Series / Department of Statistics and Mathematics **WU Vienna**
University of Economics and Business: 125.

Marwick B, Krishnamoorthy K. (2016).

Mcalpine DK (1991). Review of the Australian Kelp Flies (Diptera, Coelopidae).
Syst Entomol **16**(1)**:** 29-84.

R Development Core Team R (2015). *R: A language and environment for*
statistical computing, Vol 1. R Foundation for Statistical Computing.

Reinhardt JA, Kolaczkowski B, Jones CD, Begun DJ, Kern AD (2014). Parallel
geographic variation in Drosophila melanogaster. *Genetics* **197**(1)**:** 361-373.

Savolainen O, Lascoux M, Merila J (2013). Ecological genomics of local
adaptation. *Nature Review Genetics* **14**(11)**:** 807-820.

Schrider DR, Hahn MW, Begun DJ (2016). Parallel evolution of copy-number
variation across continents in Drosophila melanogaster. *Mol Biol Evo* **33**(5)**:**
1308-1316.

Schwander T, Libbrecht R, Keller L (2014). Supergenes and complex
phenotypes. *Current Biology* **24**(7)**:** 288-294.

Storz JF (2002). Contrasting patterns of divergence in quantitative traits and
neutral DNA markers: analysis of clinal variation. *Mol Ecol* **11:** 2537-2551.

Sturtevant AH (1921). A case of rearrangement of genes in *Drosophila*.
Proceedings of The National Academy of Science of the united States of America
7: 235–237.

Summers RW, Smith S, Nicoll M, Atkinson NK (1990). Tidal and Sexual
Differences in the Diet of Purple Sandpipers Calidris-Maritima in Scotland. *Bird*
Study **37:** 187-194.

Thompson M, Jiggins C (2014). Supergenes and their role in evolution. *Heredity*
113(1)**:** 1-8.

Tigano A, Friesen VL (2016). Genomics of local adaptation with gene flow. *Mol*
Ecol.

Ungerer MC, Halldorsdottir SS, Purugganan MD, Mackay TF (2003). Genotype-
environment interactions at quantitative trait loci affecting inflorescence
development in Arabidopsis thaliana. *Genetics* **165**(1)**:** 353-365.

Via S, Lande S (1985). Genotype–environment interaction and the evolution of
phenotypic plasticity. *Evolution* **39:** 505-522.

Weinig C (2000). Plasticity versus canalization: population differences in the
timing of shade-avoidance responses. *Evolution* **54**(2)**:** 441-451.

Wellenreuther M, Hansson B (2016). Detecting polygenic evolution: problems,
pitfalls, and promises. *Trends Genet* **32**(3)**:** 155–164.

Wellenreuther M, Rosenquist H, Jaksons P, Larson W (2017). Local adaptation
along an environmental cline in a species with an inversion polymorphism. *JEB*.

Wilcockson R, Crean C, Day T (1995). Heritability of a Sexually Selected
Character Expressed in Both Sexes. *Nature* **374**(6518)**:** 158-159.

Yeaman S (2013). Genomic rearrangements and the evolution of clusters of
locally adaptive loci. *PNAS*.

Tables

Table 1. Spatial coordinates (WGS 1984 DD), sea surface salinity (PSU),

inversion frequency, and wrackbed composition of the study sites.

	Coordinates (WGS84, DD)		Sea surface salinity (PSU)[a]		Inversion frequency[b]	Wrackbed composition			
	Latitude	Longitude	Mean annual	Annual SD	Frequency of the α inversion	Percentage *Fucus spp.*	Percentage Kelp[c]	Percentage red algae[d]	Percentage eelgrass[e]
Steninge	56.77	12.63	15.79	16.62	0.49	50	20	30	0
Mölle	56.28	12.49	15.38	16.88	0.42	30	30	30	10
Kämpinge	55.4	12.98	12.82	11.74	0.35	30	0	30	40
Kåseberga	55.38	14.06	9.91	5.4	0.34	30	0	50	20

[a] Data source: Sbrocco & Barber (2013), accessed on 1 November 2016.
[b] Day et al. (1983).
[c] *Saccharina* and *Laminaria spp.*
[d] *Furcellaria lumbricalis*
[e] Zostera marina

Table 2. Best models for adult weight and development time. An X indicates that a main effect or interaction was present in the best model.

	Main Effects				2-way interactions						3-way interactions
	Karyotype	Population	Sex	Substrate	Karyotype:Population	Karyotype:Sex	Karyotype:Substrate	Population:Sex	Population:Substrate	Sex:Substrate	Karyotype:Population:Substrate
Weight	x	x	x	x	x	x	x	x	x		x
Development Time	x	x	x	x		x					

Table 3. Results from Krishnamoorthy and Lee's (2014) modified signed-likelihood ratio test (MS-LRT) looking to see if the coefficient of variation is different between males and females. Bolded p-values are significant at the P< 0.0001 level after a Bonferroni correction.

Trait	MS-LRT Statistic	P-value
Weight	79.22	**0**
Thorax Width	83.79	**0**
Leg Length	61.64	**4.11E-15**
Wing Length	118.05	**0**
Development Time	22.09	**2.61E-06**

Titles and legends to figures

Figure 1: Map of *C. frigida* fly populations used in this study. Base map from d-maps (http://d-maps.com/carte.php?num_car=23102).

Figure 2: Principal Components Analysis (PCA) of four morphological measurements (thorax width, weight, wing length and middle leg tibia length) of adult flies from the reciprocal transplant experiment. Only the first two PCs are shown. Flies are categorized by sex (males-squares, females-circles) and karyotype (black - αα, grey - αβ, and white-ββ).

Figure 3: Boxplot of weight across populations (listed at Top), substrate (listed on x-axis), sex (indicated by top (male) and bottom (female) charts), and karyotype (indicated by color, black - αα, grey - αβ, and white-ββ). Outliers are not shown for clarity. N for each population x sex combination (from left to right for each karyotype and substrate as listed in the figure): Kämpinge Male: 1, 9, 4, 3, 20, 15, 3, 13, 19, 1, 21, 6, Kåseberga Male: 2, 11, 6, 10, 19, 5, 5, 9, 10, 2, 9, 6, Mölle Male: 1, 6, 8, 10, 14, 13, 6, 12, 13, 2, 6, 10, Steninge Male: 0, 3, 2, 7, 17, 7, 5, 15, 5, 7, 16, 14, Kämpinge Female: 0, 5, 4, 1, 12, 11, 0, 13, 18, 3, 17, 9, Kåseberga Female: 2, 3, 6, 7, 24, 11, 4, 12, 18, 5, 13, 5, Mölle Female: 4, 10, 4, 4, 9, 4, 6, 6, 5, 4, 3, 8, Steninge Female: 2, 5, 4, 8, 21, 8, 8, 19, 6, 7, 21, 7.

Figure 4: Boxplot of development time across populations (listed at Top), substrate (listed on x-axis), sex (indicated by top (male) and bottom (female) charts), and karyotype (indicated by color, black - αα, grey - αβ, and white-ββ). Outliers are not shown for clarity. N for each population x sex combination (from left to right for each karyotype and substrate as listed in the figure): Kämpinge Male: 1, 9, 4, 3, 20, 15, 3, 13, 19, 1, 21, 6, Kåseberga Male: 2, 11, 6, 10, 19, 5, 5, 9, 10, 2, 9, 6, Mölle Male: 1, 6, 8, 10, 14, 13, 6, 12, 13, 2, 6, 10, Steninge Male: 0, 3, 2, 7, 17, 7, 5, 15, 5, 7, 16, 14, Kämpinge Female: 0, 5, 4, 1, 12, 11, 0, 13, 18, 3, 17, 9, Kåseberga Female: 2, 3, 6, 7, 24, 11, 4, 12, 18, 5, 13, 5, Mölle Female: 4, 10, 4, 4, 9, 4, 6, 6, 5, 4, 3, 8, Steninge Female: 2, 5, 4, 8, 21, 8, 8, 19, 6, 7, 21, 7.

Figure 5: Box plot of frequencies of different karyotypes (black - αα, grey - αβ, and white-ββ) across populations (listed at the top of each graph) and substrates (listed on the x-axis). Dashed lines indicate HWE expectations (solid - αα, dashed - αβ, and dotted-ββ) based on Day et al (1983) data.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplemental Material

Table S1. Pearson correlations of morphological factors.

	Adult Weight	Thorax Width	Wing Length	Leg Length
Weight	1	0.8306678	0.8850752	0.7786367
Thorax Width	0.8306678	1	0.8452453	0.7628986
Wing length	0.8850752	0.8452453	1	0.865498
Leg Length	0.7786367	0.7628986	0.865498	1

Table S2. ANOVA tables for the full models adult weight (A) and development time (B). Bolded terms are included in the final models.

A.

Term	DF	Sum of Squares	Mean Square	F-Value
Population	**3**	**201.08**	**67.03**	**13.6683**
Substrate	**3**	**327.74**	**109.25**	**22.2781**
Sex	**1**	**1559.27**	**1559.27**	**317.9702**
Karyotype	**2**	**2589.74**	**1294.87**	**264.0532**
Population:Substrate	**9**	**185.65**	**20.63**	**4.2065**
Population:Sex	**3**	**422.75**	**140.92**	**28.7363**
Population:Karyotype	**6**	**228.57**	**38.09**	**7.7683**
Substrate:Sex	3	18.59	6.2	1.264
Substrate:Karyotype	**6**	**116.87**	**19.48**	**3.9719**
Sex:Karyotype	**2**	**1406.79**	**703.4**	**143.4383**
Population:Substrate:Sex	9	64.31	7.15	1.4572
Population:Substrate:Karyotype	**18**	**220.67**	**12.26**	**2.5**
Population:Sex:Karyotype	6	36.49	6.08	1.2401
Substrate:Sex:Karyotype	6	53.67	8.95	1.8242

B.

Term	DF	Sum of Squares	Mean Square	F-Value
Population	**3**	**11.039**	**3.6796**	**3.6796**
Substrate	**3**	**35.277**	**11.759**	**11.759**
Sex	**1**	**20.778**	**20.7776**	**20.7776**
Karyotype	**2**	**6.958**	**3.4792**	**3.4792**
Population:Substrate	9	9.476	1.0528	1.0528
Population:Sex	3	6.46	2.1533	2.1533
Population:Karyotype	6	0.461	0.0769	0.0769
Substrate:Sex	3	0.598	0.1994	0.1994
Substrate:Karyotype	6	1.639	0.2732	0.2732
Sex:Karyotype	**2**	**6.725**	**3.3625**	**3.3625**
Population:Substrate:Sex	9	3.591	0.399	0.399
Population:Substrate:Karyotype	18	4.861	0.2701	0.2701
Population:Sex:Karyotype	6	0.263	0.0438	0.0438
Substrate:Sex:Karyotype	6	0.534	0.0889	0.0889

Figure S2: Boxplot of thorax width across populations (listed at Top), substrate (listed on x-axis), sex (indicated by top (male) and bottom (female) charts), and karyotype (indicated by color, red - αα, blue - αβ, and green-ββ). Outliers are not shown for clarity.

Figure S3: Boxplot of leg length across populations (listed at Top), substrate (listed on x-axis), sex (indicated by top (male) and bottom (female) charts), and karyotype (indicated by color, red - αα, blue - αβ, and green-ββ). Outliers are not shown for clarity.

Figure S4: Boxplot of wing length across populations (listed at Top), substrate (listed on x-axis), sex (indicated by top (male) and bottom (female) charts), and karyotype (indicated by color, red - αα, blue - αβ, and green-ββ). Outliers are not shown for clarity.

Figure S2

Figure S3

Figure S4