Evaluating metabarcoding to analyse diet composition of species foraging in

anthropogenic landscapes using Ion Torrent and Illumina sequencing

Marie-Amélie Forin-Wiart[1,2,3] (MAFW), Maxime Galan[4] (MG), Sylvain Piry[4] (SP), Jean-

François Cosson[4] (JFC), Claire Larose[5] (CL) and Marie-Lazarine Poulle[1,2] (MLP)

[1] Université de Reims Champagne-Ardenne (URCA), SFR Cap-Santé, EA3800-PROTAL,

UFR de Médecine, 1 rue du Maréchal Juin, 51098 Reims Cedex, France

[2] URCA, CERFE, 5 rue de la Héronnière, 08240 Boult-aux-Bois, France

[3] Université de Strasbourg, CNRS, Département Ecologie, Physiologie et Ethologie, IPHC

UMR 7178, F-67087 Strasbourg, France.

[4] CBGP, INRA, CIRAD, IRD, Montpellier SupAgro, Université de Montpellier, 755 avenue

du Campus Agropolis, CS 300 16, 34988 Montferrier-sur-Lez cedex, France

[5] DIANA PetFood Division, Z.A. du Gohélis, 56250, Elven, France

Corresponding author: Marie-Amélie Forin-Wiart, Université de Reims Champagne-Ardenne,

CERFE, 5 rue de la Héronnière, 08240 Boult-aux-Bois, France. email: ma.forinw@hotmail.fr

Running title: HTS platform sensitivity in diet metabarcoding

Abstract

Determining the foraging niche of wild species is a challenging task, recently overcome through DNA metabarcoding. This method is successfully used to identify ingested food items of faecal samples. However, several limitations can affect the output of this method. Here, two benchtop high-throughput sequencing (HTS) platforms were evaluated, the Ion PGM (Life Technologies) and the MiSeq (Illumina). We assessed their ability to identify a large taxonomic array of food items for species having a diet possibly containing human-processed products and prey items. We sequenced faecal DNA samples obtained from domestic cats *Felis silvestris catus*, generalist carnivores presenting numerous specific DNAs in their diet. We assessed platform sensitivity and data reproducibility using both a cat fed on a controlled mixed diet along with a calibration study and free-ranging individuals sampled from a rural area. All the samples were analysed in independent PCR duplicates on each platform. The results of the calibration study showed that the output from MiSeq was more sensitive and reproducible than that from Ion PGM. Detection variability was linked to DNA degradation: 82% of the expected occurrences were detected for intact preys, versus 54% for pet food and 31% for out-of-date raw fish and meat. The Ion PGM detected significantly less human-linked food, birds, field voles, murids and shrews in the field-collected samples than the MiSeq platform. Pooling the replicates from both platforms and filtering HTS data allowed identifying at least one food item in 87.4% of the 326 field-collected samples. A total of 29 prey taxa (90% at species level) and 33 human-related food taxa (76% at genus level) were identified. The findings indicate that using HTS platforms such as MiSeq, which provide reads of sufficiently high quantity and quality, with sufficient numbers of technical replicates, is a robust and non-invasive approach to determine diet composition in animals foraging on a wide range of items in anthropogenic landscapes.

43   Key-words: cyt*b*, HTS data filtering, MiSeq sequencing, non-invasive approach, sequencer

44   bias, variability in DNA degradation

Introduction

With increasing urbanisation and loss of natural habitat, several species have adapted to human-dominated landscapes by shifting their dietary niche to include human-processed food items [1]. For instance, scavengers such as gulls *Larus spp.*, ravens *Corvus spp.*, Eurasian badgers *Meles meles*, opossums *Didelphis virginiana*, racoons *Procyon lotor*, or wild boar *Sus scrofa*, as well as opportunistic generalist carnivores such as coyotes *Canis latrans* [2], or foxes *Vulpes spp.* [3,4], presently have dietary compositions that differ substantially from their natural diet. Several large carnivores such as bears, wolves and hyenas have also altered their diet to include significant parts of human-processed products (see Bateman and colleagues [5] for a review). Understanding to what extent species in close contact with humans may have changed their diet to include human-processed items is critical both from a conservation perspective to determine the likelihood of human-wildlife conflicts, and from an evolutionary perspective to determine species plasticity in foraging niches in a context of global change. Thus, accurately identifying a large diversity of food items in the diet composition of studied species, including those from human-linked food sources is necessary for ecologists.

Foraging ecology studies generally rely on the preservation of hard remains found in the faeces, pellets, or stomach contents of studied species to determine their diet, using visual recognition of morphologically features by macro- or microscopic methods [6]. Whereas useful, these techniques generally present a poor resolution in terms of taxa identification or preclude the identification of food items that do not leave behind hard remains [7,8]. In contrast, DNA metabarcoding has been demonstrated as an accurate method for diet analysis using faecal samples as a source of DNA [9]. As a result, this technique has become a key application in high-throughput sequencing (HTS) technologies in ecological research [10]. Its aim is the

analysis of biological samples (of which there is no prior knowledge) that contain a mixture of

DNA using PCR amplification of a gene marker such as universal DNA barcodes.

Metabarcoding in HTS technologies allows DNA to be read from multiple templates for

hundreds of samples in the same sequencing run. A subunit of a mitochondrial gene – for

example, a 136-bp mini-barcode on the cytochrome *b* gene (cyt*b*) – is enough to distinguish

between closely related rodent species even in degraded or ancient DNA samples [11]. By

comparing obtained sequences to a standard reference library of taxonomically identified

organisms (e.g. Bold or GenBank), the taxa or species present in a sample can be identified

with high confidence. In this way, metabarcoding of faecal samples allows ingested food

items to be identified by characterizing the barcode sequences in the samples [12]. This

technique has been successfully used to analyse the diet of species of birds [13], mammals

[11,14,15], reptiles [16,17] and fish [18].

However promising, metabarcoding for dietary analyses is not free of limitations or bias (see

Table 1 in Galan and colleagues [19] for a review). Contamination from alien DNA during

fieldwork or cross-contamination during lab procedures can occur, resulting in false positive

results [20]. Furthermore, the library preparation protocol, the targeted DNA region, the primer

choice and the sequencing platform can all affect the identification of the composition of

environmental samples [21]. The degree of DNA degradation in an analysed sample also

influences the sensitivity of PCR-based methods [22]. In addition, the sequencing of PCR

products from enriched samples that contain predator DNA could reduce the potential

detection of the prey sequences [15]. As a result of these potential biases, alone or in

combination, the relative abundance of some species in faecal samples can be under- or

overestimated in a DNA analysis, while other species may not be amplified and so remain

absent from the bioinformatics output [23]. Although several studies have described most of

these challenges, to our knowledge the specific use of HTS platforms to analyse samples from

a dietary perspective has never been evaluated.

The main objective of this work was to assess the ability of benchtop HTS platforms in

identifying a large taxonomic array for species having a diet possibly containing human-

processed products. To maximise the variability in DNA samples obtained from the diet and

ensure the presence of human-linked food items, we focused on a high generalist carnivore

species, the domestic cat *Felis silvestris catus*. Cats appear as a particularly interesting model

to test for metabarcoding sensitivity since in addition to human-processed products, free-

ranging individuals are known to feed on a diversity of prey items including rodent species,

birds, reptiles, amphibians and insects [24,25]. Specifically, the sensitivity of the two most

popular sequencers available for metabarcoding analyses, the Life Technologies Ion PGM

(Personal Genome Machine) platform and the Illumina MiSeq platform, in identifying food

items from faecal DNA having undergone different degrees of degradation (fresh prey items

versus processed food versus out-of-date raw fish and meat) was investigated. This was

achieved by collecting faecal samples from (1) a cat fed on a mixed diet of known

composition in an initial calibration study, and (2) free-ranging individuals sampled from a

rural area. Since all samples were analysed in independent PCR duplicates on each HTS

platform, this allowed us to compare the reproducibility of diet results both within the

platform and among platforms.

Materials and methods

Faecal sample collection

Calibration study. A feeding trial was performed with a housebound cat fed a known diet

during 22 consecutive days in May 2012 and 27 consecutive days in November 2012 (see

Table S1 for details). The daily diet of the housebound cat was constant in quantity but

differed in composition from day to day. Daily rations included one or two of the following categories of food item: (1) processed wet or dry pet food (commercial meat-based cat food, Eco+ product line, SCAMARK, France); and (2a) prey items or (2b) out-of-date raw fish or meat. The daily rations of individual prey species, processed pet food, and out-of-date raw fish and meat were weighed and distributed evenly in three to four meals. The faecal samples were then collected from a litter tray that was cleaned out daily, and these were put in labelled plastic bags and stored at -20 °C. See Table S1 for detailed information related to daily ration contents.

Field study. The faecal samples from free-ranging domestic cats were collected in a rural area located in northeastern France (Ardennes region) as part of a joint study (Forin-Wiart et al. unpublished data). Cats were sampled from a monitored free-ranging domestic cat population composed of 162 individuals, 40% of which were free-ranging outdoor cats fed *ad libitum* by their owners, and 60% of which were farm cats fed from time to time with milk, dry pet food or leftovers (see Forin-Wiart and colleagues [26] for the census and cat feeding status definitions). Over 100 fresh faecal samples were collected in June–July 2011, November–December 2011 and March 2012 along an 11-km survey transect delineated in villages and isolated farms and a 15-km transect in two areas surrounding the villages. Additional faecal samples were also collected from the litter trays of free-ranging individuals. Putative cat faecal samples were identified in the field by their shape and size [27] prior to molecular identification based on cat-specific quantitative PCR (qPCR, see below). The 359 putative felid faecal samples were labelled with their location (using a GPS) and stored at -20 °C in plastic bags.

DNA extraction

All the DNA extractions were performed at the SAS SPYGEN (France) laboratory in a room

dedicated to degraded or rare DNA. Total DNA was extracted from approximately 500 mg of

faecal material taken from all parts of the scat, as predator and prey DNA are not evenly

distributed. The QIAamp DNA Stool Mini Kit was used following the manufacturer's

instructions (Qiagen). Ten extraction blanks (containing no faecal sample) were included as a

negative control to test for contamination. DNA extraction success was determined by real-

time PCR (qPCR) amplification with a cat species-specific primer [28] by SAS SPYGEN. This

step was also used to validate the origin of field-collected faecal samples.

DNA metabarcoding

A 136-bp mini-barcode of the cytochrome *b* gene was targeted, as it is effective in identifying

a large range of vertebrates, including ungulates, rodents, birds and fish, from degraded

samples [11,29,30].

Sequencing was successively performed using the Ion Personal Genome Machine (Ion PGM;

Life Technologies) platform and the MiSeq (Illumina) platform to analyse the 400 DNA

extractions obtained from the putative cat faecal samples (41 samples from the calibration

study and 359 from the field study). Each DNA extraction was sequenced four times: twice

independently on each HTS platform. Due to the specificities of each platform, the DNA

library preparation differed slightly (see below).

The Ion PGM libraries were performed twice for each DNA extraction by amplifying the 136-

bp fragment of cytochrome *b* described in Galan and colleagues [11]. We used the fusion

primers L15411F (5'-CCA TCT CAT CCC TGC GTG TCT CCG ACT CAG NNNNNNN

GAY AAA RTY CCV TTY CAY C-3') and H15546R-PGM (5'CCT CTC TAT GGG CAG

TCG GTG AT NNNNNNN AAR TAY CAY TCD GGY TTR AT-3') modified in 5' by the

addition of individual-specific 7-bp MIDs (Multiplex IDentifiers NNNNNNN) and adaptors

required for the emPCR and the Ion PGM sequencing. PCRs were performed following the

procedure detailed in Galan and colleagues [11]. After PCR pooling and size selection by gel

extraction, the amplicon libraries were sequenced by the company Genes Diffusion (Douai,

France) on an Ion PGM system using an Ion 316 Chip Kit version 2 (Ion Torrent, Life

Technologies).

The MiSeq libraries were performed using a two-step PCR protocol (see Illumina Application

Note Part #15044223) combined with the dual-index paired-end sequencing approach

described in Kozich and colleagues [31] due to the significant length of the Illumina adaptors

(see details in Galan and colleagues [32]). During PCR1, the 136-bp fragment of cyt*b* was

amplified twice for each DNA extraction with the primers modified in 5' by the addition of a

partial overhang Illumina adapter: L15411F-MiSeq 5'-TCG TCG GCA GCG TCA GAT GTG

TAT AAG AGA CAG (none or C or TC or ATG) GAY A AA RTY CCV TTY CAY CC-3'

and H15546R-MiSeq 5'-GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA G

(none or C or TG or GCT) AAR TAY CAY TCD GGY TTR AT-3'. The alternative bases

between the partial adaptors and the target-specific primers corresponded to a 0- to 3-bp

'heterogeneity spacer' designed to mitigate the issues caused by low sequence diversity

amplicons in Illumina sequencing [33]. These four versions of each primer were mixed before

PCR1, which was performed in the same conditions as the Ion PGM PCR libraries, i.e.

following the procedure detailed in Galan and colleagues [11].

PCR1 was then used as the DNA template for PCR2, consisting of a limited-cycle

amplification step to add multiplexing indices and Illumina sequencing adapters P5 and P7 at

both ends of each DNA fragment. The primers P5 (5'-AAT GAT ACG GCG ACC ACC

GAG ATC TAC AC NNNNNNNN TCG TCG GCA GCG TC-3') and P7 (5'-CAA GCA

GAA GAC GGC ATA CGA GAT NNNNNNNN GTC TCG TGG GCT CGG-3') were

synthesized with the different index sequences (NNNNNNNN) described in Kozich and

colleagues [31]. PCR2 was carried out in a 10-µL reaction volume using 5 µL of PCR Multiplex

Kit Master Mix (Qiagen) and 0.7 µM of each primer. One microliter of the PCR1 product was

added to each well. PCR2 was started with an initial denaturation step of 95 °C for 15 min,

followed by 5 cycles of denaturation at 94 °C for 40 s, annealing at 45 °C for 45 s and

extension at 72 °C for 60 s, followed by a final extension step at 72 °C for 10 min. After PCR

pooling and size selection by gel extraction, the amplicon library was sequenced at the local

Plateau GPTR-GENOTYPAGE facility (Montpellier, France) on a MiSeq instrument by

paired-end sequencing using a MiSeq 500 Cycle Kit version 2 (Illumina).

Data processing

Bioinformatics and taxonomic identification. The paired-end reads 1 and 2 produced by

MiSeq were assembled with the 'make.contigs' tool available in the mothur program [34].

SESAME barcode software (SEquence Sorter & AMplicon Explorer [35]) was used for both Ion

PGM and MiSeq data to sort the sequences and identify the haplotypes at a taxonomic level

(genus or species) using GenBank and three different similarity thresholds for

BLAST+FASTA assignment (95%, 98% and 100%).

Data filtering for false positive and negative occurrences. In line with Esling and colleagues

[36], a mistagging threshold was calculated to prevent laboratory contamination or sample cross-

contamination. This threshold corresponds to the number of reads for a particular haplotype

below which a positive occurrence could be due to contamination. It is essential to filter out

putative false positive results, especially for food items originating from highly represented

item sequences in the sequencing run. The mistagging threshold was calculated using the

maximal mistagging frequency in a sequencing run, *i.e.* the higher percentage of sequences

assigned to a wrong sample and determined by the known mixed-diet study (e.g. the number

of sequences of a rodent species assigned to a specific sample when this species was not

expected in the targeted sample). The mistagging threshold corresponds to the total number of

reads in a sequencing run for a particular haplotype, multiplied by the maximal mistagging

frequency (e.g. if an Ion PGM run produced 34,000 reads of an *Arvicola terrestris* haplotype

and if the maximal mistagging frequency = 0.05%, then the mistagging threshold = 17 reads

for this *Arvicola terrestris* haplotype).

Another possible source of bias is that DNA from a food item of low biomass or from a

highly degraded sample can be stochastically amplified during the PCR process, resulting in a

proportion of false negative results [37]. To overcome this problem of stochastic amplification, a

specific food item had to occur in at least 50% of the PCR replicates performed for each DNA

extraction of a given sample. Following these stages of data filtering, a table of occurrences of

observed food items was obtained per sequencing run in order to compare the performance of

the two platforms (see below). In a final step, to determine the diet represented by each faecal

sample, the food occurrence data from both HTS platforms was pooled to enhance food item

detection.

Data analysis

Model selection was based on the Information Theoretic approach and was performed with

the Akaike Information Criterion (AIC) following the model selection approach described by

Zuur and colleagues [38]. All statistical analyses were conducted with the statistical software R

3.3 (R Development Core Team 2016 [3939]), and a value of alpha = 0.05 was used for all tests.

Calibration study. The ability of the two HTS platforms to detect consumed items from a diet

of no processed food (prey items only) or of different degrees of processed food (wet and dry

pet food, raw fish or meat) was assessed through descriptive statistics. For each HTS platform

and each food item category (prey, pet food or raw fish and meat), the number of faeces

containing haplotype sequences informative enough to assign a food item (observed

occurrence) was compared to the minimum expected occurrence in order to identify the

sensitivity and bias of each sequencing strategy. The minimum expected occurrence was

determined by considering that the DNA associated with a given food item ingested by the

housebound cat was expected to be detected in the scat emitted at least 12 h after the ingestion

of the food item, according to the transit time previously assessed by Forin-Wiart and

colleagues [26]. The accordance of food item detection (i.e. its occurrence in one, both or none

of the replicates from a sequencing run) for each HTS platform and each successfully

amplified scat was also estimated. Additionally, for mixed meals (i.e. including two food item

categories), the proportion of each food item category by mass ingested by the housebound

cat relative to the total mass of the food ingested was compared to the proportion of each food

item category identified by haplotype information in the sequence reads relative to the total

number of reads of a specific faecal sample. This measure allowed the investigation of

whether the proportion of a given food item category in the diet of cats was reflected in the

HTS sequence counts. These proportions were compared using Fisher's exact tests for count

data.

Field study. Small- and medium-sized rodent, insectivore, wild bird and reptile DNA in field-

collected samples were placed in the 'Prey items' category and its sub-categories (see Table

S2 for details). In contrast, DNA from domestic ungulates (cow, goat, sheep, pig) and farmed

birds (chicken, duck, guinea fowl, Perdicinae *spp.*, turkey), as well as other mammals (e.g.

rabbit *Oryctolagus cuniculus*), farmed fish and game species (*Capreolus capreolus, Cervus*

elaphus and *Sus scrofa*) were placed in the 'Human-linked food' category and its sub-

categories (see Table S2 for details). This latter category was assumed to originate from pet

food, leftovers or hunting scraps. These food item categories were defined to fit those

previously obtained during diet studies.

To identify the sensitivity of each sequencing protocol, the probability of detecting a food

item in field-collected samples, i.e. presence/absence (Table S3), was modelled as a function

of the HTS platform (fPlatform), the food item category (fFoodItem, reptile occurrence was

excluded for modelling purposes) and the interaction between fPlatform and fFoodItem, using

a Generalized Linear Model (GLM) with a binomial link function. The fit of the full model

(lme4 package [4040]) and the reduced model was compared using a deviance test (G^2) as an

alternative to an individual parameter-based approach by considering fSample as a random

effect. Multiple comparisons with interaction terms were then adjusted using Tukey's post-

hoc analyses (lsmeans package [4141]).

Finally, the diet obtained from field-collected faecal samples was assessed through descriptive

statistics.

Results

Sequencing results

The Ion PGM and MiSeq platforms provided 4,042,017 and 4,042,606 raw sequences passing

filter respectively. For the Ion PGM reads, the percentage of bases with a quality score of 30

(Q30: incorrect base call probability of 1 in 1000) or higher was 61.6%; for the MiSeq reads,

the percentage of bases with a quality score of 30 or higher was 96.9%. For the Ion PGM run,

a high proportion of truncated sequences without reverse tags was observed (2,490,973 reads

< 126bp excluding tags and primers) due to a significant decrease in quality at the end of the

sequences. After demultiplexing, 688,530 (using Ion PGM) and 2,509,592 (using MiSeq)

sequences of cyt*b* were assigned to the faecal samples and were identified to the closest

taxonomic rank (genus or species) using the consensus of the three similarity thresholds

(95%, 98% and 100%).

In the calibration study, a high proportion of cat haplotypes were obtained: 41,129 reads

(80%) for Ion PGM and 235,673 reads (98%) for MiSeq at the 95% similarity threshold.

There were also a high proportion of unidentified sequences: 5,069 reads (12%) for Ion PGM

and 1,434 reads (0.6%) for MiSeq at the 95% similarity threshold. In the study of field-

collected data, sequences from predators (cat, but also Canids and Mustelids, see below)

represented 60% of the total read count using Ion PGM, and 88% of the total read count using

MiSeq. For both platforms, a high proportion of cat haplotypes (301,329 reads [51%] for Ion

PGM and 1,690,359 reads [83%] for MiSeq) and unidentified sequences (70,050 reads [12%]

for Ion PGM and 125,086 reads [6%] for MiSeq, with a $\leq$ 95% identity using GenBank

sequences) were obtained. As found in a previous study by Deagle and colleagues [42], some

human DNA sequences were also present, accounting for 0.06% and 0.6% of the sequences

generated by Ion PGM and MiSeq respectively.

DNA metabarcoding on known mixed-diet samples

In the MiSeq run, one out of 1934 sequences of a *Myodes glareolus* haplotype (n = 1,

resulting in a frequency of 0.0517) was detected during the last 21 days of the calibration

study despite the fact the housebound cat was fed only with pet food and raw fish or meat (see

Table S1 for details), so this constituted a mistagging event. Accordingly, the maximal

mistagging frequency in a sequencing run was set at 0.05%.

The DNA of prey items was mostly recovered in duplicates from the MiSeq (7 out of 8

replicated occurrences) and the Ion PGM (3 out of 4 replicated occurrences) runs, i.e. 100%

of prey species expected in a given scat were detected (Table 1 and Table S1). MiSeq was the

only platform to detect the Norway rat, the house mouse and a bird species (Table 1). DNA

belonging to pet food contents was preferentially recovered with the MiSeq platform (85% of

observed occurrences) as compared to the Ion PGM platform (45%) and was largely

recovered by duplicates with the MiSeq platform (71%) (Table 1). The DNA from raw fish

and meat was scarcely detected by either method and was never detected in Ion PGM PCR

replicates (Table 1 and Table S1).

Both platforms were better at detecting DNA from prey items (82% of the expected

occurrences) than DNA from pet food (54%), or from raw fish and meat (31%) (Table 1).

Independently of the food item considered, the percentage of faecal samples in which the

expected DNA of a food item was detected (see Table S1) was higher for the MiSeq run

(25/27, 93%) than for the Ion PGM run (14/27, 52%) (Table 1). Combining the results of the

HTS runs from both platforms, the sequencing of food item DNA in the faeces collected from

the housebound cat succeeded in 66% of the samples (27 out of 41 samples).

Of the six expected prey species, the wood mouse *Apodemus sp.* was not detected, while at

least one of the two HTS platforms sequenced four other rodent species (water vole, field

vole, Norway rat and house mouse) and a bird species (Table 1). DNA from the Norway rat

and field vole, each expected in one scat, as with the other food items, was recovered in two

consecutive faeces. DNA from the house mouse was poorly detected (i.e. only four

informative reads from the MiSeq run), and the wood mouse was not detected at all. A delay

of 12 to 24 hours between prey ingestion and the occurrence of their remains in the scat was

identified for each prey species eaten and detected by both methods.

The proportion of sequences (Ion PGM + MiSeq) associated with the different food item

categories did not match the proportion of the food item consumed ($p < 0.001$; Fig. 1a), even

when the platforms were considered independently (Fig. 1b and 1c). Prey items were

considerably overrepresented: they accounted for 66% of the sequence reads assigned to food

items, whereas they represented only 5% of the biomass ingested by the cat. Within the prey

item category, rodent and bird biomass (90.4% and 9.6% respectively) were representative of

their read proportions (95.7% and 4.3% respectively, p > 0.05). Conversely, human-linked

food items (pet food and raw fish or meat) were considerably underrepresented: pet food

accounted for 33% and raw fish or meat 0.5% of the sequence reads, whereas these

respectively represented 83% and 12% of the biomass ingested.

DNA metabarcoding on field-collected samples

Comparing predator identification using cat-specific PCR amplification and DNA

metabarcoding, 2.4% of faecal samples (10/415) identified as cat faeces using cat-specific

PCR amplification were found to be emitted by another predator species using DNA

metabarcoding; 5% of faecal samples (20/415) not identified as cat faeces using cat-specific

PCR amplification were found to be emitted by a cat using DNA metabarcoding. Of the 359

putative felid faecal samples collected, 90.8% (326/359) were confirmed as originating from

domestic cats using sequencing. The others were determined to be the dejections of Canids

(4.5% from the domestic dog *Canis lupus familiaris* and 3.9% from the red fox,) or Mustelids

(0.6% from the Eurasian badger *Meles meles* and 0.3% from *Martes sp.*). All a priori DNA

metabarcoding identifications of predators were confirmed unambiguously by both HTS

platforms and by the four PCR replicates.

Of the 159 occurrences of prey items detected with Ion PGM, 65% (*n* = 103) were identified

in duplicates compared to 72% (130/180) for MiSeq (Table 2 and Table S2). Of the 115

occurrences of human-linked food items detected with Ion PGM, 39% (*n* = 45) were

identified in duplicates compared to 48% (90/189) for MiSeq (Table 2 and Table S2). The two

platforms jointly confirmed 72.2% (143/198) of the occurrences of prey items, and 61%

(115/189) of the occurrences of human-linked food (pet food and scraps) (Table 2 and Table

S2). In terms of the cat faecal samples, Ion PGM detected at least one food item in 73.6%

(240/326) of the investigated samples, and MiSeq detected at least one food item in 83.1%

(271/326) of cases. Combining the results from both HTS platforms, 87.4% (285/326) of the

investigated samples contained at least one identified food item.

A likelihood ratio test applied to the model selected for food item detection indicated that the

model without random effects was significantly better than the one with random effects (L =

870.09, df = 1, p > 0.05). The top-ranked model included fPlatform and fFoodItem and the

interaction between these two variables (R^2 = 0.17, Table S4). Ion PGM detected significantly

less human-linked food (-2.01 ± 0.31 SE on the log odds ratio scale, p < 0.001), birds (-4.98 ±

1.10 SE, p < 0.001), field voles (-1.50 ± 0.72 SE, p < 0.05), murids (-3.30 ± 1.10 SE, p <

0.01) and shrews (-2.60 ± 1.24 SE, p < 0.05) than MiSeq.

A total of 521 different occurrences of food items were identified and classified as prey items

or human-linked food. A total of 29 different prey taxa were identified, of which 25 were

identified without ambiguity up to the species level (9 rodents, 3 insectivores, 12 birds and 1

reptile), representing 90% of the prey items (287/319, Table S2). One reptile item was

identified as being the slow worm *Anguis fragilis*. Out of 202 different faecal samples, 478

occurrences related to human-linked food items were detected. Game species were also

detected in addition to the expected occurrences of pet food, food leftovers and hunting

remains. A total of 33 different taxa related to human-linked food were identified, of which 17

were identified without ambiguity up to the species level (30%, 155/478) and 23 up to the

genus level (76%, 364/478, Table S2).

Discussion

Effect of food item categories and their DNA integrity

The use of the short cyt*b* mini-barcode provided a good sensitivity of food item detection,

particularly for prey items. Galal-Khallaf and colleagues [18] highlighted the advantage of

amplifying a short mini-barcode when employing a DNA metabarcoding approach on

degraded DNA in highly processed food, based on 454 Next Generation Sequencing (NGS) to

detect fish species in commercial fish feed samples. However, in our study non-reproducible

occurrences between replicates were observed using Ion PGM and MiSeq runs independently

(86% for Ion PGM and 56% for MiSeq runs during the calibration study, and 48% for Ion

PGM and 40% for MiSeq runs during the field study). These findings emphasize the need to

generate sufficient numbers of technical replicates (at least four were required in the present

study) to detect rare or degraded DNA, in order to limit the presence of false negatives as well

as to remove inconsistent results likely to be false positives [37].

Results from the calibration study revealed that DNA related to processed food is rarely

recovered in duplicates and is less detected than the DNA of prey items by both HTS

platforms. This result might be linked to a DNA integrity gradient between prey items, pet

food and raw fish and meat. Prey items were frozen immediately after death during the

calibration study, and in the field study they were probably the result of hunting rather than

scavenging behaviour [25], ensuring maximum DNA integrity before ingestion. In addition, in

this food category, hard parts (e.g. bones versus flesh or other soft parts) limit DNA

degradation during digestion, and so partly solve problems of variable digestibility [7]. In

contrast, the stages of food processing (e.g. cooking, dehydration, heat sterilization, etc.)

involved in producing pet food induce a higher rate of DNA degradation than for prey items,

leading to damaged and degraded DNA [22,43]. However, food processing also suppresses the

activity of micro-organisms such as bacteria, so the integrity of the DNA remains the same

regardless of the duration of food storage before ingestion. In the case of the category of raw

fish and meat, which is not processed, DNA integrity depends on the length of storage and the

activity of micro-organisms. In this study, the use of out-of-date raw fish and meat stored in a

supermarket refrigerator for several days/weeks might explain the low success of DNA

detection related to this food category. The link between food category and DNA integrity

could be confirmed through a calibration study including the evaluation of DNA integrity and

a DNA metabarcoding experiment on various food item categories before and after ingestion.

The effect of the HTS platform

In addition to variability in the DNA detection of food items, variations in the ability of the

different HTS platforms to detect food items were identified. Overall, during the calibration

study, MiSeq was more effective at detecting expected food items and duplicating this (93%

of expected items were detected, of which 71% were duplicate-confirmed) than Ion PGM

(52% of expected items were detected, of which 29% were duplicate-confirmed). A similar

pattern was observed in the analysis of the field-collected faecal samples. Furthermore, MiSeq

was also able to detect more human-linked food, as well as species of bird, field vole, murids

and shrews than Ion PGM in field collected-samples. This outcome could be most easily

explained by the difference in the number of sequence reads assigned to the samples between

the two HTS platforms (i.e. low sequencing depth for the Ion PGM platform), linked with the

better sequencing quality of the MiSeq platform.

Another strong sequencing bias was noticed in results from the Ion PGM platform,

particularly for species haplotypes that included short homopolymer repeats. In our case, a

loss of quality at the end of the cyt*b* mini-barcode, and consequently to the presence of several

short homopolymer stretches in tandem (e.g. the last 22 bases of the mini-barcode from *Felis*

silvestris catus: 5'-CCC TTT AAA TAC CCC TCC CCA T-3') explained a high proportion

of truncated sequences without reverse tags. This bias has been previously observed in other

public datasets generated by this platform and suggests that it is a pervasive problem [44]. This

may also partly explain the bias found between biomass and number of reads by leading to a

lower representation of homopolymer-rich haplotype sequences in the DNA metabarcoding

studies using Ion Torrent technology (e.g. Deagle and colleagues [45]).

Common biases in diet analysis using DNA metabarcoding

The proportion of reads from prey items or human-linked food was not related to the ingested

biomass of these food categories in the housebound cat diet. On a mass-to-mass basis, prey

item DNA was better detected than DNA associated with human-linked food. This result was

not related to a taxonomic bias introduced by primer-template mismatches, but rather related

to food item properties. This discrepancy is likely to be associated with the variable

digestibility of different food items and/or DNA quality (see above). In line with previous

studies [45-47], these results reinforce the preference for a qualitative approach that involves

creating a list of food items consumed and their frequency of occurrence [12,42] rather than

relating reads to ingested biomass, particularly in studies on species with mixed diets. This

issue has been highlighted in several recent DNA-based dietary studies investigating the COI

barcode region [48] or a more conserved mitochondrial region such as the mitochondrial 16S

rRNA gene [49]. Deagle and colleagues [45] also noted a lack of relationship between biomass and

the proportion of reads when comparing the proportion of three fish species detected in seal

scat to the proportion of these species in a constant diet. An alternative would be to use

conserved markers such as those targeting the mitochondrial 12S rRNA or the nuclear 18S

rRNA [50], but these provide limited taxonomic resolution for vertebrates compared to markers

targeting mitochondrial cytochrome *b*. One way to increase pet food detection in dietary

studies would be to target not only vertebrate and invertebrate components of the diet but also

cereals contained in pet food. DNA metabarcoding that targets different genes, including

those of plants, has been successfully used on an omnivorous carnivore, the brown bear *Ursus*

arctos [14]. DNA identified in faeces could also reflect secondary predation cases, e.g. insects

consumed by shrews or seeds ingested by rodents [11,51].

Another bias related to an approach based on amplifying a cyt*b* 136-bp mini-barcode without

a blocking oligonucleotide on the targeted predator species is that it can lead to an over-

representation of the sequences assigned to the predator in the PCR product. As a result, some

samples can exclusively yield predator sequences by inhibiting the amplification of rare DNA

reads (e.g. prey items). This phenomenon is probably exacerbated because predator DNA has

higher integrity than DNA originating from human-linked food. This hypothesis is consistent

with the high proportion of cat haplotypes in the present study (80% for Ion PGM and 98%

for MiSeq when faecal samples were rapidly stored, and 51% for Ion PGM and 83% for

MiSeq in more hazardous sample conservation). Of the field-collected faecal samples, 18%

produced only cat sequences. Similarly, in a study by Shehzad and colleagues [15], 21% of

faecal samples were found to be entirely composed of leopard cat *Prionailurus bengalensis*

DNA when investigating leopard cat diet using metabarcoding without a blocking

oligonucleotide on the targeted predator species. The high proportion of predator sequences in

the PCR product reduces the sensitivity of HTS platforms. This phenomenon contributes to

explaining the relatively low sequencing success rate obtained with Ion PGM, for which the

total number of reads was lower than with MiSeq.

In line with D'Amore and colleagues [21], these results indicate that studies using DNA

metabarcoding for conservation management purposes should evaluate their outputs in

comparison with a calibration study or at least with biological expectations. It should be noted

that our calibration study also indicated underestimation (e.g. small murids) and

overestimation (e.g. DNA from large species [Norway rat] or numerously ingested prey

species [field vole] found in more than one sample) of food items in cat diets; this was linked

to molecular bias (e.g. sequencing bias for particular taxa due to homopolymer stretches or

primer mismatches) or targeted species biology (i.e. pyloric retention in Felids). Several

previous studies also support this pattern [26,52,53].

This study also confirmed that species assignment is subject to errors related to species

sequences from the cyt*b* gene submitted to GenBank (e.g. closely related rodents/birds that

are not distributed within the study range), and thus potentially poorly allocated. DNA

metabarcoding is mainly used to investigate faeces with no previous knowledge about the diet

of the targeted species; a reference database containing voucher specimens from the study

area would be useful to validate the detection of prey species.

DNA metabarcoding improves results in dietary studies

When dealing with conservation or epidemiological concerns, misidentified faecal samples

from non-targeted species within the analysed sample can dramatically affect study

conclusions. In the present study, 9% of the field-collected faecal samples were misidentified

using morphological criterion. Hence, identifying the faeces origin with species-specific

primer pairs and quantitative PCR [15,54] is one alternative. However, the results presented here

and elsewhere [42] have shown that a positive detection in real-time PCR may reflect the

contamination of samples with a small amount of DNA from targeted species. This can be

illustrated through the human contamination usually detected in sequencing runs. In contrast,

a positive species-specific PCR can partly reflect cross-contamination of samples or field

contamination. Combining several replicates of PCR amplification with and without a

blocking oligonucleotide would increase food item detection sensitivity per sample,

particularly for small samples or when targeting the DNA of rare prey species. However, the

number of primer mismatches and the concentration of blocking oligonucleotide can affect

the detection of target species [48].

DNA metabarcoding allowed the identification of 90% of prey items up to species level. It led

to both a high detection rate and the high-resolution identification of closely related prey

species (e.g. *M. arvalis* versus *M. agrestis*, *A. flavicollis* versus *A. sylvaticus*, Table S2). Using

a DNA sequence database and bird-specific primer pairs to analyse faecal samples, Zarzoso-

Lacoste and colleagues [55] highlighted similar efficiency in using molecular ecology for the

identification of bird species in cat and rat diets. Ordinarily, this level of precision is achieved

only in cat diet studies that rely on prey animals brought back to the home (e.g. 92%

identified to at least class level [56]), not with conventional methods analysing scat and stomach

contents (e.g. from 25% [57] to 59% [58]).

The results of this study demonstrate that DNA metabarcoding has the great advantage of

allowing the detection of occurrences of food items that leave undetectable remains in the

field (such as human-related food items) in a non-invasive sampling approach. Even

unprecedented, this method gives insights on the potential origin of ingested food items (e.g.,

pet food or scavenged human-processed food versus scavenged carrion), with a taxonomic

identification of 76% of vertebrate cyt*b* variants to the genus level. This makes this method

useful for evaluating the exploitation of human-linked foods in urban wildlife populations,

above all in endangered species, for which it is preferable to use non-invasive survey

approaches. Nevertheless, the results of the calibration study make it clear that the rate of

false negatives is partly linked to DNA integrity (and so to food item categories). This finding

suggests that even if DNA metabarcoding allows more efficient detection of specific food

items, certain items in the diet may remain highly underestimated (e.g. 46% for pet food and

70% for out-of-date raw fish and meat), which could have a significant impact on

conservation strategies. It should be noted, however, that sequence proportions matched

biomass within a food category (e.g. rodents and birds).

To conclude, the present study joins several metabarcoding studies considering the biases and

limitations of this approach, particularly in terms of using benchtop HTS platforms. In line

with Piñol and colleagues [48], our findings indicate that universal primers bias the detection of

degraded DNA in a heterogeneous mixture. This can drastically affect the identification of the

ingested taxa for species having a diet possibly containing human-linked food items, and thus

the inferences drawn about this diet regarding the impacts of human activities on wildlife

populations and food-web dynamics in anthropogenic landscapes. Nevertheless, generating

sufficient numbers of technical replicates and using a sequencer such as MiSeq to produce a

high quantity and quality of reads appears to be a robust approach for investigating the

exploitation of a large range of food items for populations relying on a mixed-diet through

non-invasive sampling. Indeed, compared to conventional approaches based on morphological

analysis, DNA metabarcoding allows the identification of a large number of food items up to

species level and a wide range of taxa.

Acknowledgements

We are grateful to the people in the study area who allowed us to prospect for and collect

samples, to the cat owners who collected samples for the study, to Matthieu Bastien, Ivan

Puga Gonzales, Nancy Rebout and Claire Saraux for their statistical advice, and to Vincent

Viblanc for its constructive comments. We also sincerely thank Spécialités Pet Food (SPF) for

funding the DNA extraction and the two HTS runs.

Data accessibility

Should the manuscript be accepted, the data supporting the results (raw sequence reads (fastq

format), raw table of abundance and filtered table of occurrence including taxonomic

affiliations) will be published along with the manuscript in a Dryad Digital Repository.

Author contributions

MAFW and MLP contributed to the study design and sample collection. MG conducted the

molecular analysis. MAFW, MG and SP conducted the bioinformatics analysis. MAFW

conducted the data analysis, performed the statistical analysis and prepared the tables and

graphs. MAFW, MLP and MG led the writing of the manuscript. All authors contributed

critically to the drafts and gave final approval for publication.

References

1   McCleery, R. A., Moorman, C. E. & Peterson, M. N. *Urban wildlife conservation:*
theory and practice. (Springer, 2014).
2   Murray, M. *et al.* Greater consumption of protein□poor anthropogenic food by urban
relative to rural coyotes increases diet breadth and potential for human-wildlife
conflict. *Ecography* **38**, 1235-1242 (2015).
3   Robardet, E. *et al.* Infection of foxes by *Echinococcus multilocularis* in urban and
suburban areas of Nancy, France: influence of feeding habits and environment.
Parasite **15**, 77-85 (2008).
4   Newsome, S. D., Ralls, K., Job, C. V. H., Fogel, M. L. & Cypher, B. L. Stable
isotopes evaluate exploitation of anthropogenic foods by the endangered San Joaquin
kit fox (*Vulpes macrotis mutica*). *J. Mammal.* **91**, 1313-1321, doi:10.1644/09-mamm-
a-362.1 (2010).
5   Bateman, P. W., Fleming, P. A. & Le Comber, S. Big city life: carnivores in urban
environments. *J. Zool.* **287**, 1-23, doi:10.1111/j.1469-7998.2011.00887.x (2012).
6   Pompanon, F. *et al.* Who is eating what: Diet assessment using next generation
sequencing. *Mol. Ecol.* **21**, 1931-1950 (2012).
7   Yoccoz, N. G. The future of environmental DNA in ecology. *Mol. Ecol.* **21**, 2031-
2038 (2012).
8   Egeter, B., Bishop, P. J. & Robertson, B. C. Detecting frogs as prey in the diets of
introduced mammals: a comparison between morphological and DNA-based diet
analyses. *Molecular Ecology Resources* **15**, 306-316, doi:10.1111/1755-0998.12309
(2015).
9   Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C. & Willerslev, E. Towards
next-generation biodiversity assessment using DNA metabarcoding. *Mol. Ecol.* **21**,
2045-2050 (2012).
10   Shokralla, S., Spall, J. L., Gibson, J. F. & Hajibabaei, M. Next□generation sequencing
technologies for environmental DNA research. *Mol. Ecol.* **21**, 1794-1805 (2012).
11   Galan, M., Pagès, M. & Cosson, J.-F. Next-generation sequencing for rodent
barcoding: species identification from fresh, degraded and environmental samples.
PLoS ONE **7**, e48374, doi:10.1371/journal.pone.0048374 (2012).
12   Valentini, A., Pompanon, F. & Taberlet, P. DNA barcoding for ecologists. *Trends*
Ecol. Evol. **24**, 110-117, doi:10.1016/j.tree.2008.09.011 (2009).
13   Coghlan, M. L. *et al.* Metabarcoding avian diets at airports: Implications for birdstrike
hazard management planning. *Investigative Genetics* **4** (2013).
14   De Barba, M. *et al.* DNA metabarcoding multiplexing and validation of data accuracy
for diet assessment: Application to omnivorous diet. *Molecular Ecology Resources* **14**,
306-323 (2014).
15   Shehzad, W. *et al.* Carnivore diet analysis based on next-generation sequencing:
Application to the leopard cat (*Prionailurus bengalensis*) in Pakistan. *Mol. Ecol.* **21**,
1951-1965 (2012).
16   Brown, D. S., Ebenezer, K. L. & Symondson, W. O. C. Molecular analysis of the diets
of snakes: Changes in prey exploitation during development of the rare smooth snake
Coronella austriaca. *Mol. Ecol.* **23**, 3734-3743 (2014).
17   Brown, D. S., Jarman, S. N. & Symondson, W. O. C. Pyrosequencing of prey DNA in
reptile faeces: Analysis of earthworm consumption by slow worms. *Molecular*
Ecology Resources **12**, 259-266 (2012).

18 Galal-Khallaf, A., Osman, A. G., Carleos, C. E., Garcia-Vazquez, E. & Borrell, Y. J.
A case study for assessing fish traceability in Egyptian aquafeed formulations using
pyrosequencing and metabarcoding. *Fisheries Research* **174**, 143-150 (2016).
19 Galan, M. *et al.* 16S rRNA amplicon sequencing for epidemiological surveys of
bacteria in wildlife: the importance of cleaning post-sequencing data before estimating
positivity, prevalence and co-infection. *bioRxiv*, 039826 (2016).
20 Ficetola, G. F. *et al.* Replication levels, false presences and the estimation of the
presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources* **15**,
543-556 (2015).
21 D'Amore, R. *et al.* A comprehensive benchmarking study of protocols and sequencing
platforms for 16S rRNA community profiling. *BMC Genomics* **17**, 55 (2016).
22 Sharma, N., Thind, S. & Sharma, D. Effect of meat processing on genomic DNA
quality and specific gene amplification. *Journal of Applied Animal Research* **28**, 69-72
(2005).
23 Coissac, E., Riaz, T. & Puillandre, N. Bioinformatic challenges for DNA
metabarcoding of plants and animals. *Mol. Ecol.* **21**, 1834-1847, doi:10.1111/j.1365-
294X.2012.05550.x (2012).
24 Fitzgerald, B. M. & Turner, D. C. in *The domestic cats: The Biology of its Behaviour*
(eds D.C. Turner & P. Bateson) 151-175 (Cambridge Univ. Press, 2000).
25 Spotte, S. in *Free-ranging cats: Behavior, ecology, management*   (ed Wiley
Blackwell) Ch. 9, 181-213 (John Wiley & Sons, 2014).
26 Forin-Wiart, M.-A., Gotteland, C., Gilot-Fromont, E. & Poulle, M.-L. Assessing the
homogeneity of individual scat detection probability using the bait-marking method on
a monitored free-ranging carnivore population. *European Journal of Wildlife Research*
60, 665-672, doi:10.1007/s10344-014-0833-0 (2014).
27 Corbett, L. K. *Feeding ecology and seasonal organization of wildcats (Felis silvestris)*
and domestic cats (Felis catus) in Scotland., PhD thesis, University of Aberdeen,
Scotland, (1979).
28 Lopez, J. V., Cevario, S. & O'Brien, S. J. Complete Nucleotide Sequences of the
Domestic Cat (Felis catus) Mitochondrial Genome and a Transposed mtDNA Tandem
Repeat (Numt) in the Nuclear Genome. *Genomics* **33**, 229-246,
doi:http://dx.doi.org/10.1006/geno.1996.0188 (1996).
29 Kappel, K., Haase, I., Käppel, C., Sotelo, C. G. & Schröder, U. Species identification
in mixed tuna samples with next-generation sequencing targeting two short
cytochrome b gene fragments. *Food Chem.* **234**, 212-219 (2017).
30 Teletchea, F., Bernillon, J., Duffraisse, M., Laudet, V. & Hänni, C. Molecular
identification of vertebrate species by oligonucleotide microarray in food and forensic
samples. *J. Appl. Ecol.* **45**, 967-975 (2008).
31 Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D.
Development of a dual-index sequencing strategy and curation pipeline for analyzing
amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl. Environ.*
Microbiol. **79**, 5112-5120 (2013).
32 Galan, M. *et al.* Metabarcoding for the parallel identification of several hundred
predators and their prey: Application to bat species diet analysis. *Molecular Ecology*
Resources, 1-16, doi:10.1111/1755-0998.12749 (2017).
33 Fadrosh, D. *et al.* An improved dual-indexing approach for multiplexed 16S rRNA
gene sequencing on the Illumina MiSeq platform. *Microbiome* **2**, 6 (2014).
34 Schloss, P. D. *et al.* Introducing mothur: Open-Source, Platform-Independent,
Community-Supported Software for Describing and Comparing Microbial

Communities. *Appl. Environ. Microbiol.* **75**, 7537-7541, doi:10.1128/aem.01541-09
(2009).

35 Piry, S., Guivier, E., Realini, A. & Martin, J. F. | SE| S| AM| E| Barcode: NGS-
oriented software for amplicon characterization–application to species and
environmental barcoding. *Molecular Ecology Resources* **12**, 1151-1157 (2012).

36 Esling, P., Lejzerowicz, F. & Pawlowski, J. Accurate multiplexing and filtering for
high-throughput amplicon-sequencing. *Nucleic Acids Res.* **43**, 2513-2524 (2015).

37 Ficetola, G. F., Taberlet, P. & Coissac, E. How to limit false positives in
environmental DNA and metabarcoding? *Molecular ecology resources* **16**, 604-607
(2016).

38 Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A. & Smith, G. M. *Mixed effects*
models and extensions in ecology with R. (Springer, 2009).

39 R: A language and environment for statistical computing. (Vienna, Austria, 2016).

40 Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models
using lme4. *arXiv preprint arXiv:1406.5823* (2014).

41 Lenth, R. V. Least-Squares Means: The R Package lsmeans. *Journal of Statistical*
Software **69**, 1-33, doi:10.18637/jss.v069.i01 (2016).

42 Deagle, B. E., Kirkwood, R. & Jarman, S. N. Analysis of Australian fur seal diet by
pyrosequencing prey DNA in faeces. *Mol. Ecol.* **18**, 2022-2038, doi:10.1111/j.1365-
294X.2009.04158.x (2009).

43 De Battisti, C. *et al.* Pyrosequencing as a tool for rapid fish species identification and
commercial fraud detection. *J. Agric. Food Chem.* **62**, 198-205 (2013).

44 Loman, N. J. *et al.* Performance comparison of benchtop high-throughput sequencing
platforms. *Nature Biotechnology* **30**, 434-439 (2012).

45 Deagle, B. E., Thomas, A. C., Shaffer, A. K., Trites, A. W. & Jarman, S. N.
Quantifying sequence proportions in a DNA-based diet study using Ion Torrent
amplicon sequencing: Which counts count? *Molecular Ecology Resources* **13**, 620-
633 (2013).

46 Egge, E. *et al.* 454 pyrosequencing to describe microbial eukaryotic community
composition, diversity and relative abundance: a test for marine haptophytes. *PloS one*
8, e74371 (2013).

47 Pochon, X., Bott, N. J., Smith, K. F. & Wood, S. A. Evaluating detection limits of
next-generation sequencing for the surveillance and monitoring of international marine
pests. *PLoS One* **8**, e73935 (2013).

48 Piñol, J., Mir, G., Gomez-Polo, P. & Agustí, N. Universal and blocking primer
mismatches limit the use of high-throughput DNA sequencing for the quantitative
metabarcoding of arthropods. *Molecular Ecology Resources* **15**, 819-830,
doi:10.1111/1755-0998.12355 (2015).

49 Deagle, B. E., Chiaradia, A., McInnes, J. & Jarman, S. N. Pyrosequencing faecal DNA
to determine diet of little penguins: is what goes in what comes out? *Conserv. Genet.*
11, 2039-2048 (2010).

50 Deagle, B. E., Jarman, S. N., Coissac, E., Pompanon, F. & Taberlet, P. DNA
metabarcoding and the cytochrome c oxidase subunit I marker: not a perfect match.
Biol. Lett. **10**, 20140562 (2014).

51 Sheppard, S. K. *et al.* Detection of secondary predation by PCR analyses of the gut
contents of invertebrate generalist predators. *Mol. Ecol.* **14**, 4461-4468 (2005).

52 Krauze-Gryz, D., Gryz, J. & Goszczyński, J. Predation by domestic cats in rural areas
of central Poland: an assessment based on two methods. *J. Zool.* **288**, 260-266,
doi:10.1111/j.1469-7998.2012.00950.x (2012).

53   Pires, M. M., Widmer, C. E., Silva, C. & Setz, E. Z. F. Differential detectability of
696   rodents and birds in scats of ocelots, *Leopardus pardalis* (Mammalia: Felidae).
697   *Zoologia (Curitiba)* **28**, 280-283 (2011).
698   54   Knapp, J., Umhang, G., Poulle, M.-L. & Millon, L. Development of a real-time PCR
699   for a sensitive one-step coprodiagnosis allowing both the identification of carnivore
700   feces and the detection of Toxocara spp. and Echinococcus multilocularis. *Applied and*
701   *environmental microbiology* **82**, 2950-2958 (2016).
702   55   Zarzoso-Lacoste, D. *et al.* Improving morphological diet studies with molecular
703   ecology: An application for invasive mammal predation on island birds. *Biol. Conserv.*
704   **193**, 134-142, doi:10.1016/j.biocon.2015.11.018 (2016).
705   56   Kauhala, K., Talvitie, K. & Vuorisalo, T. Free-ranging house cats in urban and rural
706   areas in the north: useful rodent killers or harmful bird predators? *Folia Zoologica* **64**
707   (2015).
708   57   Germain, E., Ruette, S. & Poulle, M.-L. Likeness between the food habits of European
709   wildcats, domestic cats and their hybrids in France. *Mammalian Biology* **74**, 412-417
710   (2009).
711   58   Biró, Z., Lanszki, J., Szemethy, L., Heltai, M. & Randi, E. Feeding habits of feral
712   domestic cats (*Felis catus*), wild cats (*Felis silvestris*) and their hybrids: trophic niche
713   overlap among cat groups in Hungary. *Journal of Zoology (London)* **266**, 187-196
714   (2005).

Figures and tables

Table 1 The minimum expected occurrences and the actual observed occurrences of food items in HTS runs of 41 faecal samples collected from a housebound cat. The observed occurrences are shown according to the HTS platform used, the PCR replicates and the pooled occurrences from both platforms.

Food category	Food items	Minimum expected occurrence	Observed occurrence in the Ion PGM run			Observed occurrence in the MiSeq run			Observed occurrences (Ion PGM & MiSeq)
			in one replicate	in duplicates	Total	in one replicate	in duplicates	Total	
Prey items	Water vole *Arvicola terrestris*	3	0	2	2	0	2	2	2
	Field vole *Microtus agrestis*	1	1	1	2	0	1	1	2
	Norway rat *Rattus norvegicus*	1	0	0	0	1	1	2	2
	House mouse *Mus musculus*	2	0	0	0	0	1	1	1
	Wood mouse *Apodemus sp.*	2	0	0	0	0	0	0	0
	Chestnut-eared finch *Taeniopygia guttata*	2	0	0	0	0	2	2	2
	Total prey items	11	1	3	4	1	7	8	9
Pet food*		37	8	1	9	8	10	18	20
Out-of-date raw fish or meat**		13	1	0	1	1	2	4	4
TOTAL		**61**	**10**	**4**	**14**	**10**	**20**	**30**	**33**

* Pet food taxa recovered (Suidae, *Sus sp.*, *Bos sp.*, *Ovis sp.*, Caprie, *Gallus sp.*, *Anser sp.*, *Meleagris gallopavo*, *Numida meleagris*, *Salmo salar*).

** Out-of-date raw meat taxa recovered (*Sus scrofa*, *Ovis aries*, *Cairina moschata*).

Table 2 Observed occurrences of food items in 285 field-collected faecal samples from free-ranging cats. The observed occurrences are shown according to the HTS platform used and the PCR replicates. The final observed occurrence corresponds to the pooled results from the Ion PGM and MiSeq runs. See S1 Table for more details about food sub-categories.

Food category	Food sub-category	Observed occurrence in the Ion PGM run			Observed occurrence in the MiSeq run			Observed occurrence
		in one replicate	in duplicates	Total	in one replicate	in duplicates	Total	
	Small/medium-sized rodents	88	133	221	119	111	230	271
	Insectivores	1	3	4	7	2	9	10
Prey items	Wild birds	3	3	6	22	13	35	37
	Reptiles	0	0	0	1	0	1	1
	Total prey items	92	139	231	149	126	275	319
	Game species	4	1	5	8	6	14	15
Human-linked food	Anthropogenic food	117	53	169	289	151	439	187
	Total human-linked food	121	54	174	297	157	453	202
	TOTAL	**213**	**193**	**405**	**446**	**283**	**728**	**521**

Figure 1 Comparison between the proportion of biomass of the three food categories fed to the housebound cat (black) versus the overall proportion of sequence reads assigned to that food category (light grey) from 28 scat samples for (a) pooled replicates (Ion PGM + MiSeq runs), (b) Ion PGM replicates only, and (c) MiSeq replicates only.