Changes in mRNA abundance drive shuttling of RNA binding proteins, linking
cytoplasmic RNA degradation to transcription

Sarah Gilbertson[1], Joel D. Federspiel[2], Ella Hartenian[1], Ileana M. Cristea[2], and Britt
Glaunsinger[1,3,4*]

[1]Department of Molecular and Cell Biology, University of California, Berkeley CA 94720-3370,
[2]Department of Molecular Biology, Princeton University, Princeton, NJ 08544, [3]Department of
Plant & Microbial Biology, University of California, Berkeley CA 94720-3370, [4]Howard
Hughes Medical Institute.

*Correspondence: Glaunsinger@berkeley.edu

Abstract

Alterations in global mRNA decay broadly impact multiple stages of gene expression, although signals that connect these processes are incompletely defined. Here, we used tandem mass tag labeling coupled with mass spectrometry to reveal that changing the mRNA decay landscape, as frequently occurs during viral infection, results in subcellular redistribution of RNA binding proteins (RBPs) in human cells. Accelerating Xrn1-dependent mRNA decay through expression of a gammaherpesviral endonuclease drove nuclear translocation of many RBPs, including poly(A) tail-associated proteins. Conversely, cells lacking Xrn1 exhibited changes in the localization or abundance of numerous factors linked to mRNA turnover. Using these data, we uncovered a new role for relocalized cytoplasmic poly(A) binding protein in repressing recruitment of TATA binding protein and RNA polymerase II to promoters. Collectively, our results show that changes in cytoplasmic mRNA decay can directly impact protein localization, providing a mechanism to connect seemingly distal stages of gene expression.

Introduction

mRNA decay is a critical stage of gene expression that regulates the abundance and lifespan of cellular mRNAs. Many viruses including alpha and gammaherpesviruses, influenza A virus, and SARS coronavirus accelerate host mRNA degradation through the use of viral proteins that trigger endonucleolytic cleavage of mRNAs in the cytoplasm. Each of these viral proteins bypasses the rate-limiting deadenylation step of the basal decay pathway, resulting in cleaved mRNAs that are rapidly degraded by the major cellular 5'-3' exonuclease Xrn1 (Covarrubias et al., 2011; Gaglia et al., 2012). This process, termed 'host shutoff', allows viruses to rapidly restrict cellular gene expression in order to blunt immune responses and liberate resources for viral replication (Abernathy and Glaunsinger, 2015; Burgess and Mohr, 2015; Gaglia, 2010;

Rivas et al., 2016). Viral endonucleases have also served as tools for deciphering how cells sense

and respond to large changes in mRNA abundance.

While mRNA decay is often considered the terminal stage of gene expression, the rate of

mRNA decay has recently been shown to influence transcription by RNA polymerase II

(RNAPII) in both yeast and mammalian cells (Abernathy et al., 2015; Braun and Young, 2014;

Haimovich et al., 2013; Sun et al., 2013). In yeast, a buffering system exists in which Xrn1 plays

a major role in connecting mRNA synthesis and decay, presumably allowing cells to maintain an

appropriate overall mRNA abundance. Mammalian cells also have a mechanism to sense mRNA

levels, though the pathway appears to operate differently than in yeast. Here, accelerated

cytoplasmic mRNA degradation does not lead to a compensatory increase in mRNA synthesis, as

might be predicted by the homeostatic model, but instead decreases cellular RNAPII promoter

recruitment, thereby amplifying the restrictive gene expression environment (Abernathy et al.,

2015). Significant transcriptional repression as measured by nascent mRNA production was

reported to occur at approximately 9% of host genes, although validation experiments suggested

this number is likely to be an underestimate (Abernathy et al., 2015).

The mRNA decay-transcription feedback pathway is activated in mammalian cells

infected with gammaherpesviruses like Kaposi's sarcoma-associated herpesvirus (KSHV) and

murine gammaherpesvirus 68 (MHV68), as well as upon expression of virally encoded mRNA

endonucleases in uninfected cells. Herpesviral endonucleases, including the muSOX protein of

MHV68, cleave mRNA but do not impact the abundance of noncoding RNAs transcribed by

RNA polymerase I (RNAPI) or III (RNAPIII) (Covarrubias et al., 2011). Correspondingly,

muSOX-induced mRNA decay elicits a significant decrease in RNA polymerase II (RNAPII)

recruitment to cellular promoters (Abernathy et al., 2015). Notably, depletion of Xrn1 from

muSOX-expressing cells prevents the ensuing RNAPII transcriptional repression. This suggests

that the initial mRNA cleavage and translational inactivation are insufficient to restrict RNAPII

recruitment, and that subsequent exonucleolytic degradation of the cleaved mRNA fragments is a

critical signaling step.

Little is currently known about this pathway linking cytoplasmic mRNA decay to

RNAPII activity in mammalian cells, including the nature of the signal that is transmitted

between the two cellular compartments. An attractive hypothesis is that one or more RNA

binding proteins (RBPs) differentially traffics between the cytoplasm and the nucleus when basal

rates of mRNA decay are perturbed, thereby conveying global mRNA abundance information.

Recent analyses indicate that mammalian cells contain hundreds of RBPs that bind

polyadenylated mature mRNAs, and proteins within this group have been shown to regulate all

stages of gene expression (Gerstberger et al., 2014; Mitchell and Parker, 2014; Müller-McNicoll

and Neugebauer, 2013; Singh et al., 2015). Furthermore, RBPs frequently display

nucleocytoplasmic shuttling behavior.

Here, we charted global alterations in protein localization that occur specifically in

response to increased or decreased Xrn1 activity. This revealed a set of mammalian RBPs that

preferentially move from the cytoplasm to the nucleus during accelerated mRNA decay, as well

as components of the 5'-3' decay machinery and other RBPs whose subcellular distribution is

altered in cells lacking Xrn1. Poly(A) tail associated proteins are overrepresented among the

RBPs that accumulate in the nucleus under conditions of global mRNA decay, offering an

explanation for how RNAPII could be selectively sensitive to mRNA abundance. Indeed, we

uncovered a new role for cytoplasmic poly(A) binding protein (PABPC) in mediating mRNA

decay-driven repression of RNAPII promoter recruitment. Furthermore, we show that the

recruitment of TATA binding protein (TBP) to promoters is also impaired in response to PABPC

nuclear translocation, indicating that cytoplasmic mRNA decay impacts early events in

preinitiation complex assembly. Our results reveal how mRNA levels exert significant influence

on RBP localization and suggest that select RBPs transmit mRNA abundance information from

the cytoplasm to the nucleus to broadly influence gene expression, particularly under conditions

of cellular stress.

Results

RNA binding proteins translocate from the cytoplasm to the nucleus in cells undergoing enhanced cytoplasmic mRNA decay

To chart mRNA decay-driven movement of proteins between the cytoplasm and the nucleus, we used a quantitative liquid chromatography/tandem mass spectrometry (LC/MS-MS)-based approach. Specifically, following subcellular fractionation, proteins from nuclear and cytoplasmic fractions were labeled with isobaric tandem mass tags (TMT). TMT labeling enables multiplexing of up to 11 samples per run and was proven to improve the analytical power for quantitation during viral infections (McAlister et al., 2012; Jean Beltran et al., 2016). We used HEK293T cells expressing the MHV68 muSOX endonuclease to create a condition of accelerated, Xrn1-dependent cytoplasmic mRNA decay. We previously demonstrated that muSOX expression in these cells activates the mRNA decay-RNAPII transcription feedback pathway similar to virally infected fibroblasts (Abernathy et al., 2015). Pure populations of cells expressing either WT muSOX or the catalytically dead D219A muSOX point mutant were generated using Thy1.1-based cell sorting. Here, muSOX was fused to the cell surface glycoprotein Thy1.1 with an intervening self-cleaving 2A protease, causing release of Thy1.1 from muSOX for cell surface expression and selection. Three biological replicates of control, WT, and D219A muSOX expressing cells were then separated into nuclear and cytoplasmic fractions, and trypsin-digested proteins from each fraction were differentially TMT labeled prior to LC/MS-MS (**Figure 1A**).

Among the 5,994 total quantifiable nuclear proteins (detected in all replicates), 123 displayed significant nuclear enrichment (adjusted P value of ≤ 0.05) in WT muSOX expressing cells relative to the D219A mutant (**Figure 1B, Table S1-2**). We then removed from further analysis proteins that were simultaneously increased in the cytoplasm in muSOX expressing cells

to remove proteins that increase in overall abundance, as well as proteins displaying significant

differences between the D219A catalytic mutant and the empty vector control. These filtering

steps yielded a final list of 67 proteins that were differentially enriched in the nucleus under

conditions of accelerated mRNA decay (**Figure 1B, Table S3**). Notably, 22 of the 67 proteins

(33%) are annotated as RBPs (Pantherdb) in line with the expectation that mRNA-bound

proteins in particular should be impacted during widespread mRNA degradation. In addition, 31

of the 67 proteins (46%) are listed as localized both to the cytoplasm and nucleus according to

the Database for Annotation, Visualization and Integrated Discovery (DAVID), supporting the

idea that they are shuttling factors. As an independent validation of these results, we evaluated

12 of the top hits by western blotting of fractionated cell lysates in control or muSOX-expressing

cells, 10 of which recapitulated the MS results (**Figure 1-figure supplement 1A**).

Proteins associated with the poly(A) tail display robust mRNA decay-dependent nuclear

translocation

The poly(A) tail is a defining mRNA feature and during basal mRNA decay,

deadenylation is the initiating step that licenses subsequent decapping and exonucleolytic

degradation of an mRNA (Schoenberg and Maquat, 2012). Thus, the binding state of poly(A) tail

associated proteins could theoretically serve as a readout to distinguish the overall abundance of

mRNA over other forms of RNA in the cytoplasm. Notably, nuclear relocalization of PABPC

has been observed during infection with multiple viruses that promote mRNA decay, supporting

the idea that poly(A) tail associated proteins may be particularly sensitive to mRNA abundance

(Harb et al., 2008; Lee, 2009; Park et al., 2014; Piron et al., 1998; Salaun et al., 2010). Indeed, an

overrepresentation analysis using Pantherdb revealed that poly(A) binding proteins, poly(U)

binding proteins, and mRNA 3'UTR binding proteins were significantly overrepresented among

the 67 differentially expressed proteins (**Figure 1C, Figure 1-figure supplement 1B-C**).

Proteins linked to the poly(A) tail consistently arose as robust hits in our MS dataset, including

PABPC proteins 1 and 4 (PABPC1, PABPC4), LA-related protein 4 (LARP4), and

heterogeneous nuclear ribonucleoprotein Q (HNRNPQ) (**Table S3**, **Figure 1D**). We confirmed

that PABPC1 and LARP4 also translocate to the nucleus in NIH3T3 cells infected with WT

MHV68, but not in cells infected with an MHV68 muSOX mutant virus (R443I) with impaired

mRNA cleavage activity (Adler et al., 2000; Richner et al., 2011) (**Figure 1E**). Thus, poly(A)

associated proteins preferentially move from the cytoplasm to the nucleus in response to

muSOX-activated mRNA decay in both transiently transfected and virally infected cells.

Nuclear translocation of RNA binding proteins is dependent on mRNA degradation by

Xrn1

Xrn1 is the major 5'-3' exonuclease in mammalian cells and is responsible for the

degradation of 3' RNA fragments generated upon cleavage by muSOX (Gaglia et al., 2012). In

the absence of Xrn1, muSOX-induced repression of RNAPII promoter occupancy does not

occur, suggesting that Xrn1 activity should be required for release and subsequent nuclear

translocation of RBPs involved in this phenotype (Abernathy et al., 2015). We therefore used

Cas9-based genome editing to generate Xrn1 knockout clones in HEK293T cells and confirmed

that muSOX expression in these cells failed to reduce RNAPII promoter occupancy (**Figure 2-**

figure supplement 1A-B). The Xrn1 knockout cells exhibited a ~2-fold reduction in growth

compared to control Cas9-expressing WT cells (**Figure 2-figure supplement 1C**), in line with

observations in yeast (Larimer and Stevens, 1990). Importantly, this did not lead to broad

changes in gene expression (see below). Given that Xrn1 is a central component of the

mammalian mRNA decay machinery, only low passage versions of these cells were used to

decrease the likelihood of compensatory changes occurring in other decay components. Using

the same TMT-LC/MS-MS strategy described above, we analyzed nuclear and cytoplasmic

fractions from 3 biological replicates of Xrn1 knockout cells expressing muSOX or an empty

vector control (**Figure 1A**). Comparison of these data to the list of proteins from Table S3

indicated that 45 of the 67 hits failed to shuttle in muSOX-expressing Xrn1 knockout cells,

confirming that our workflow identified factors that differentially shuttle in response to mRNA

degradation (**Figure 2A, B**).

Poly(A) tail degradation is normally carried out by deadenylases prior to activation of

Xrn1-mediated decay from the 5' end, but we previously demonstrated that SOX-cleaved

mRNAs are not deadenylated prior to their targeting by Xrn1 (Covarrubias et al., 2011). Indeed,

analysis of endogenous PABPC1 and LARP4 localization by confocal microscopy and western

blot analysis of fractionated cells confirmed that both proteins translocated from the cytoplasm to

the nucleus upon muSOX expression in WT but not Xrn1 knockout cells (**Figure 2C-D, Figure**

2-figure supplement D).

Xrn1 knockout leads to subcellular redistribution of proteins functionally associated with

RNA

Given that increased Xrn1 activity caused nuclear translocation of mRNA-associated

RBPs, we hypothesized that RBPs linked to Xrn1 function might also exhibit altered subcellular

distribution in cells lacking Xrn1. We first looked broadly for proteins with reproducibly altered

abundance in the nucleus or the cytoplasm of Xrn1 knockout cells relative to the vector control

cells. There were 149 and 158 proteins differentially expressed in the absence of Xrn1 in the

nucleus or cytoplasm, respectively (adjusted P value $\leq$ 0.05) (**Figure 3A, Table S4**). Both the

oligosaccharyltransferase (OST) complex and RBPs were significantly overrepresented among

the set of differentially expressed proteins in each compartment (**Figure 3A, Figure 3-figure**

supplement 1A). The significance of the OST enrichment is currently unknown, although the

OST complex has been shown to be critical for infection with flaviviruses, which depend on

Xrn1 for the production of a subgenomic viral noncoding RNA (Chapman et al., 2014; Moon et

al., 2012). However, the RBP enrichment is in line with Xrn1 function, and it is notable that

among the proteins significantly enriched in the nucleus of Xrn1 KO cells were factors that

encompass the first steps of 5'-3' mRNA decay. These included all members of the decapping

complex (DCP1A, DCP1B and DCP2), factors that promote decapping complex formation

(EDC3 and EDC4), and a protein that connects the decapping complex to the deadenylation

machinery (PATL1) (**Figure 3B**).

We next examined whether the absence of Xrn1 also impacted the relative abundance of

its known interaction partners (as listed in the BioGRID database) in the two compartments

(**Figure 3C**). This did not appear to be the case in the TMT data, as the majority of known Xrn1

protein partners were expressed at normal levels in the absence of Xrn1. However, there was a

significant increase in the cytoplasmic levels of UPF1, a mediator of nonsense mediated mRNA

decay (NMD). Similarly, DNASE2, a nuclease which contributes to the degradation of DNA in

dying cells, had increased cytoplasmic abundance. Secernin-2 (SCRN2), a protein involved in

exocytosis, translocated to the nucleus in the absence of Xrn1. Conversely, PABPC4 levels were

decreased in both compartments, and two centrosomal proteins CEP152 and CEP128 were

reduced in the nucleus.

Finally, we considered the possibility that upon loss of Xrn1, cells might upregulate other

components of the mRNA decay machinery. Perhaps surprisingly, out of all 5,994 detected

proteins, only three were significantly upregulated in both the nucleus and the cytoplasm of Xrn1

knockout cells: GW182, Galectin-3, and BAG1. Among these, GW182 stands out because it is a

member of the miRNA-induced silencing complex (miRISC) involved in recruitment of

deadenylases to initiate degradation of target mRNAs (**Figure 3D**). This increase in GW182

abundance, along with the changes to DCP2, DDX6, and PABPC4, were independently validated

by western blot analysis (**Figure 3E**). To determine whether the increases in the whole cell

protein abundance of GW182 and in the nuclear protein abundance in DDX6 and PATL1

occurred at the mRNA level or were a result of translational regulation, we measured steady-

state mRNA expression for each of these factors by RT-qPCR. In each case, the mRNA

abundance was increased in Xrn1 knockout cells compared to WT cells (**Figure 3-figure**

supplement 1B).Importantly, the increases appeared specific to these transcripts and not due to

generalized mRNA abundance changes in the absence of Xrn1, as there was no significant

difference in *gapdh* or *actB* mRNA levels (**Figure 3-figure supplement 1B**).

Collectively, these data suggest that there are not broad increases in cellular proteins in

response to inhibition of 5'-3' mRNA decay. However, there appear to be selective increases in

the whole cell or compartment-specific abundance of select factors associated with mRNA

decay, which likely arises from increases in their mRNA levels in Xrn1 knockout cells.

LARP4 shuttles to the nucleus in a PABPC-dependent manner

Protein relocalization in response to altered cytoplasmic mRNA decay could occur as a

consequence of direct interactions with the nuclear transport machinery that are antagonized by

mRNA, as has been documented for the PABPC nuclear localization signal (NLS) (Kumar et al.,

2011). Alternatively, translocation could occur indirectly via interactions with other proteins that

contain nuclear transport signals. To test for this latter possibility, we first plotted the network of

known interactions among the list of proteins that relocalized in cells undergoing accelerated

mRNA decay using the STRING database (**Figure 4A**). There were significantly more

interactions among this set of proteins than would be predicted for a random group of proteins of

similar size (p = 0.0496), with many of the interactions involving PABPC. This enrichment

suggests that these proteins are biologically related, confirming what was seen in the GO term

analysis. We examined the relocalization mechanism for one of the PABPC interacting proteins,

LARP4 (Yang et al., 2011). We reasoned that if LARP4 relocalization involved direct

interactions with the nuclear import machinery, then it should relocalize in muSOX-expressing

cells in a PABPC independent manner. Conversely, if it was 'escorted' into the nucleus via its

interaction with PABPC, then its relocalization should be blocked by PABPC depletion.

Depletion of PABPC1 has been shown to lead to compensatory induction of PABPC4 , which

can function in a redundant manner (Kumar, 2010). Therefore, we co-depleted both PABPC1

and PABPC4 using siRNAs. Upon co-depletion of the PABPC proteins, LARP4 no longer

accumulated in the nucleus of muSOX-expressing cells (**Figure 4B**). In contrast, siRNA-

mediated depletion of LARP4 had no effect on PABPC1 shuttling in these cells (**Figure 4C**).

These results support a model in which LARP4 is brought into the nucleus in cells undergoing

accelerated mRNA decay through its interaction with PABPC.

PABPC depletion abrogates the muSOX-driven decrease in RNAPII promoter occupancy

Given the nuclear enrichment of many poly(A) and poly(U) associated proteins, we

considered these factors to be strong candidates for involvement in the signaling pathway linking

accelerated mRNA decay to RNAPII transcriptional repression. To determine if they were

required for the mRNA decay-transcription feedback loop, we tested whether depletion of

several of these factors individually altered RNAPII occupancy using chromatin

immunoprecipitation assays (ChIP). To test the role of PABPC we co-depleted both PABPC1

and PABPC4 using siRNAs, then monitored RNAPII occupancy at two cellular promoters

(*gapdh, rplp0*) previously shown to be responsive to mRNA decay-induced transcriptional

repression (Abernathy et al., 2015). In cells depleted of PABPC1 and PABPC4, there was no

longer a reduction in RNAPII occupancy at the *gapdh* and *rplp0* promoters in muSOX

expressing cells relative to vector control cells (**Figure 5A**). In contrast, RNAPII promoter

occupancy remained repressed in muSOX expressing cells upon depletion of LARP4 (**Figure**

5B). In addition to poly(A) tail associated proteins, we tested the effects of depleting three

additional factors that translocated to the nucleus in an mRNA-decay dependent manner: the

poly(U) binding protein MSI1, the CHD3 transcriptional regulator, and one of the top scoring

hits from the MS data, TRIM32 (**Figure 5-figure supplement 1A-C**). RNAPII occupancy

remained reduced in muSOX-expressing cells relative to vector control cells upon depletion of

MSI1, CHD3, and TRIM32 (**Figure 5-figure supplement 1A-C**).

It should be noted that when we measured the effect of depleting the above factors on

RNAPII occupancy in the absence of muSOX, we unexpectedly observed that their knockdown

alone reduced the RNAPII ChIP signal (**Figure 5-figure supplement 2A-E**). However, unlike

the case for PABPC, RNAPII levels were further reduced in muSOX expressing cells after

depletion of LARP4, MSI1, CHD3, and TRIM32 (**Figure 5B, Figure 5-figure supplement 1A-**

C). We hypothesize that knockdown of these factors may lead to broad impacts on cellular

function, in ways that directly or indirectly influence transcription. Therefore, although PABPC

appeared to be selectively involved in suppressing RNAPII occupancy during enhanced mRNA

decay, we sought an alternative strategy to evaluate its connection to this process.

Nuclear accumulation of PABPC1 is sufficient to inhibit RNAPII recruitment to promoters

Endogenous PABPC is subject to translational autoregulation, and our previous data

suggested that the abundance of PABPC in uninfected cells is fine-tuned to match poly(A) tail

availability (Kumar, 2010; Kumar et al., 2011). In this regard, even modest over-expression of

PABPC1 leads to nuclear accumulation of the 'excess' (presumably non-poly(A) bound) protein

in cells lacking muSOX (**Figure 5C**). This feature enabled us to test whether nuclear

accumulation of PABPC1 was sufficient to cause a reduction in RNAPII promoter recruitment in

the absence of muSOX-induced mRNA decay. Indeed, FLAG-PABPC1 transfected cells

displayed a significant decrease in RNAPII occupancy at the *gapdh and rplp0* promoters (**Figure**

5D). These observations suggested that the failure of muSOX to trigger transcriptional repression

in Xrn1 knockout cells might be overcome by driving PABPC into the nucleus via

overexpression. In agreement with this prediction, muSOX-induced transcriptional repression

was restored in Xrn1 knockout cells upon transfection of FLAG-PABPC1, confirming that

nuclear translocation of this RBP plays a central role in connecting cytoplasmic mRNA decay to

RNAPII promoter recruitment (**Figure 5E**).

Nuclear translocation of PABPC selectively impacts early stages of transcription

To more precisely define the stage(s) of transcription impacted by mRNA decay-induced

translocation of PABPC, we began by measuring RNAPII occupancy at both the promoter and

the gene body (exon) of the genes *gapdh*, *actB*, and *tlcd1*. In each of the experiments below, we

evaluated cells transfected with empty vector control, muSOX (to activate cytoplasmic mRNA

decay), or FLAG-PABPC1 (to selectively increase nuclear PABPC levels in the absence of

widespread mRNA decay). Cells expressing muSOX or FLAG-PABPC1 exhibited parallel

phenotypes, in which RNAPII occupancy was reduced at promoters as well as within the gene

body compared to control cells (**Figure 6A**). Western blotting confirmed that the reduced ChIP

signals were not due to a decrease in the overall levels of RNAPII in these cells (**Figure 6-figure**

supplement 1).

The C-terminal domain (CTD) of the RNAPII Rpb1 subunit has unique phosphorylation

patterns associated with each phase of transcription; it initially binds DNA in an

unphosphorylated state, but undergoes progressive serine 5-phosphorylation (Ser5P) during

initiation, then serine 2-phosphorylation (Ser2P) during elongation (Heidemann et al., 2013). To

determine whether mRNA decay-induced PABPC1 translocation impacted RNAPII initiation or

elongation in addition to promoter recruitment, we measured the ratio of total RNAPII to either

Ser5P or Ser2P RNAPII (**Figure 6B, 6C**). In both muSOX and FLAG-PABPC expressing cells,

these ratios were unchanged relative to control cells, suggesting that the primary defect is in

promoter recruitment, and that there are not independent impacts on downstream events. These

data are consistent with previous observations in MHV68-infected cells (Abernathy et al., 2015).

RNAPII promoter recruitment occurs during assembly of the transcription preinitiation

complex (PIC), a multi-step event involving numerous general transcription factors and

transcription associated factors (Roeder, 1996). The initial promoter-defining event in PIC

assembly that occurs prior to RNAPII recruitment is binding of TATA-binding protein (TBP) as

part of the transcription factor TFIID complex, whose recruitment is essential for initiating

transcription (Darzacq et al., 2007; Louder et al., 2016). Notably, TBP ChIP revealed that its

occupancy at the *gapdh*, *actB*, *tlcd1*, and *rplp0* promoters was significantly reduced in cells

expressing either muSOX or FLAG-PABPC1 compared to control cells (**Figure 6D**). Similar to

RNAPII, western blotting confirmed that this reduction in promoter binding was not due to

altered expression of TBP in these cells (**Figure 6-figure supplement 1**). Given that TBP is a

transcription factor required by cellular polymerases other than just RNAPII, we considered the

possibility that all TBP-dependent transcription might be impaired as a consequence of

cytoplasmic mRNA decay. This was not the case however, as 7SK and U6 promoter occupancy

by RNA polymerase III (RNAPIII), which also requires TBP, was unaltered in cells expressing

muSOX or FLAG-PABPC1 compared to control cells (**Figure 6E**). We therefore conclude that

mRNA decay-driven nuclear accumulation of PABPC1 reduces PIC assembly selectively at the

promoters of RNAPII transcribed genes.

Discussion

Cellular mRNA abundance can be dramatically altered in response to a variety of

pathogenic and nonpathogenic stresses including both viral and bacterial infections, and early

apoptosis (Abernathy and Glaunsinger, 2015; Barry et al., 2017; Thomas et al., 2015). In many

of these cases, accelerated cytoplasmic mRNA decay initiates a widespread reduction in

transcript levels, often through the engagement of the major mammalian 5'-3' exonuclease Xrn1

(Covarrubias et al., 2011; Gaglia et al., 2012). In addition to altering the translational landscape,

depletion of cytoplasmic mRNA elicits changes in upstream components of the mammalian gene

expression pathway, including RNAPII transcription, largely by unknown mechanisms

(Abernathy et al., 2015). Here, we tested the hypothesis that cellular RNA binding proteins may

shift their subcellular localization in response to altered mRNA decay, thus conveying mRNA

abundance information between the cytoplasm and the nucleus (**Figure 6F**). Quantitative

proteomics was previously reported to allow the discovery of viral infection-induced protein

translocations (Jean Beltran et al., 2017). Indeed, our unbiased TMT-based proteomics approach

revealed that among the total cellular protein pool, an RBP-enriched protein subset concentrates

in the nucleus specifically in response to increased mRNA decay in an Xrn1 dependent manner.

RBPs have critical roles in all stages of gene expression (Müller-McNicoll and Neugebauer,

2013), and our data further emphasize their multifunctional capacity.

We also found that RBPs are enriched in the set of proteins with altered nuclear or

cytoplasmic localization in Xrn1 knockout cells. Interestingly, factors involved in decapping, the

event that directly precedes Xrn1 attack during basal mRNA decay, were selectively increased in

the nuclei of cells lacking Xrn1. Furthermore, we detected increased levels of the NMD factor

UPF1 in the cytoplasm and overall elevated levels of GW182 in these cells. One speculative

possibility is that these changes occur in response to cellular 'reprogramming' of the mRNA

decay network, for example shifting emphasis towards 3' end targeting mechanisms to

compensate for the absence of the primary 5' end decay mechanism. This scenario might explain

the increase in GW182 levels, as it recruits the cellular deadenylase complexes PAN2-PAN3 and

CCR4-NOT to mRNA targets to promote mRNA decay by Xrn1 (Braun et al., 2011; Chekulaeva

et al., 2011; Fabian et al., 2011; Huntzinger and Izaurralde, 2011). In its absence, the increased

GW182 levels may accelerate deadenylation to instead promote 3'-5' decay. Alternatively, the

RBP nuclear and/or cytoplasmic enrichment in Xrn1 knockout cells may reflect changes that

occur when the cytoplasmic mRNA decay rate is reduced. The fact that we did not observe

significant redistribution of the majority of Xrn1 interacting proteins argues against a model in

which physical association with Xrn1 helps control decay factor protein localization.

Aside from RBPs, there was a clear overrepresentation of the OST complex, which

catalyzes co-translational N-glycosylation, among the set of differentially expressed proteins in

Xrn1 knockout cells. Although OST does not have established links to RNA decay or Xrn1, it

has been shown to be a critical component of the replication cycle of flaviviruses such as Dengue

and Zika (Marceau et al., 2016; Puschnik et al., 2017). Furthermore, these arthropod-borne

flaviviruses inhibit Xrn1 activity through a subgenomic viral noncoding RNA that contains an

Xrn1 blocking sequence (Chapman et al., 2014; Moon et al., 2012). In this context, it will be

exciting to explore possible links between these two processes, especially given that small

molecule OST inhibitors are now being tested for their pan-flaviviral inhibition (Puschnik et al.,

2017).

Among the set of proteins that translocated in cells undergoing accelerated Xrn1-

dependent mRNA decay, there was a striking enrichment in factors that bind the 3'end of

mRNAs. This supports the hypothesis that this class of RBPs would be significantly impacted by

the mRNA abundance and availability. PABPC nuclear translocation in particular has been well

documented in the context of infection with viruses that drive mRNA decay (Bablanian et al.,

1991; Borah et al., 2012; Harb et al., 2008; Lee, 2009; Park et al., 2014; Piron et al., 1998;

Salaun et al., 2010), and our unbiased proteomics approach establishes it as one of the most

robustly relocalized RBPs under these conditions. Several features of PABPC render it an ideal

indicator of mRNA abundance. First, its association with poly(A) tails implies that depletion of

mRNAs but no other type of abundant non-polyadenylated RNAs should selectively alter the

level of PABPC in the RNA bound versus unbound state. Second, nuclear import of PABPC is

antagonized by cytoplasmic mRNA abundance. We previously reported that PABPC harbors

noncanonical NLSs within its RNA recognition motifs (RRMs); upon poly(A) binding, these

elements are masked and the protein is thus retained in the cytosol (Kumar et al., 2011).

However, release of PABPC from mRNA exposes the NLSs, enabling its interaction with

importin α and its subsequent nuclear import. The observation that PABPC localization is

directly influenced by mRNA abundance suggest that cells must carefully calibrate the ratio of

PABPC to mRNA. Indeed, PABPC protein binds an autoregulatory A-rich sequence in the

5'UTR of its own mRNA to disrupt 40S ribosomal scanning and reduce its translation (Neto et

al., 1995; Wu and Bag, 1998).

When bound to poly(A) tails in the cytoplasm, PABPC contributes to mRNA stability

and facilitates protein-protein interactions for efficient translation by the ribosome (Burgess and

Gray, 2010; Fatscher et al., 2014). However, when concentrated in the nucleus, PABPC

functions instead to restrict gene expression. One previously established mechanism by which

gene expression is inhibited involves disruption of mRNA processing, where PABPC drives

hyperadenylation of nascent mRNAs (Kumar, 2010). In this study, we reveal that nuclear

accumulation of PABPC phenotypically mimics muSOX-dependent repression of RNAPII

promoter binding; it appears necessary and sufficient to repress RNAPII promoter recruitment as

a consequence of accelerated mRNA decay. Both muSOX and FLAG-PABPC1 expression target

early stages of PIC assembly, as TBP and RNAPII occupancy are reduced at promoters.

Interestingly, the *S. cerevisiae* nuclear poly(A) binding protein Nab 2 has been shown to

potentiate RNAPIII activity by directly binding RNAPIII and stabilizing TFIIIB with promoter

DNA (Reuter et al., 2015), providing a precedent for PABPs influencing transcription. However,

although TBP is required for the activity of other polymerases including RNAPIII, we found that

the impact of mRNA decay-induced PABPC translocation appears specific to RNAPII

responsive promoters. Furthermore, while RNAPII levels are reduced at both promoters and in

the gene body, the residual promoter-bound population of RNAPII does not appear to have

additional defects in promoter escape or elongation, as measured by polymerase CTD

phosphorylation patterns. Collectively, these observations suggest that altered PABPC

trafficking primarily impacts the very earliest stages of PIC assembly. Determining which factors

govern the specificity for RNAPII responsive promoters during accelerated mRNA decay and

their connection to nuclear PABPC remain important challenges for the future.

Although we did not detect a role for LARP4, MSI1, CHD3, or TRIM32 in muSOX-

induced transcriptional repression, these findings are complicated by the observation that their

depletion alone impaired RNAPII recruitment. In our hands, this phenotype is common to the

depletion of a number of different RBPs (though not all), suggesting that their absence may

cause secondary effects on gene expression. This also underscores the importance of using

alternative assays to evaluate their contributions, as we did for PABPC. Interestingly, some of

these proteins are likely to engage in distinct gene regulatory functions in the nucleus that could

also be impacted by their altered nuclear-cytoplasmic trafficking. For example, a nuclear role for

MSI1 has recently been uncovered during mouse spermatogenesis, when it translocates from the

cytoplasm to the nucleus (Sutherland et al., 2015). In the cytoplasm, MSI1 negatively regulates

the translation of its target RNAs by competing with eukaryotic initiation factor eIF4G for

binding to PABPC (Kawahara et al., 2008). However, upon spermatocyte differentiation, MSI1

relocalizes to the nucleus through direct interaction with importin-5 (IPO5), where it

concentrates at the silent XY chromatin domain. This not only releases its repression on

translation, but also alters its repertoire of RNA targets in the nucleus. LARP4 also binds

PABPC, but unlike MSI1, this interaction promotes mRNA poly(A) tail lengthening and

stabilization in the cytoplasm (Yang et al., 2011). Our findings suggest that nuclear accumulation

of LARP4 is also dependent on its interaction with PABPC. LARP4 protein levels are controlled

post-transcriptionally via an instability determinant within its coding sequence, suggesting that

akin to PABPC, its protein abundance is tightly regulated (Mattijssen et al., 2017). The functions

of LARP4 in the nucleus, as well as other RBPs identified in this work, are currently unknown.

Exploring these roles and how they become manipulated during times of cellular stress are areas

ripe for future studies.

Finally, it is notable that connections between Xrn1-driven mRNA decay and RNAPII

transcription have also been made in yeast, providing further evidence that these seemingly

divergent stages of the gene expression cascade are intimately linked (Haimovich et al., 2013;

Sun et al., 2013). However, one key difference between this pathway in yeast and mammalian

cells is that in yeast it appears to operate as a compensatory mechanism to maintain optimal

mRNA abundance: reduced mRNA decay results in reduced transcription, and vice versa

(Haimovich et al., 2013; Sun et al., 2013). This potentially represents an evolutionary divergence

in which a unicellular eukaryote 'buffers' its overall gene expression for continued maintenance

of the organism. In multicellular eukaryotes like mammals, global mRNA decay (which is

induced by numerous pathogens) may instead serve as a stress signal, and the ensuing response is

thus geared towards shutdown of major cellular programs.

Acknowledgements

This work was funded by NIH grant CA136367 to B.G., who is also an investigator of

the Howard Hughes Medical Institute, and by NIH GM114141 to I.M.C. We thank all members

of the Glaunsinger and Coscoy labs for helpful comments and suggestions.

Author Contributions

Conceptualization, S.G., B.G.; Methodology, J.D.F., I.M.C., S.G., B.G.; Validation, S.G., E.H.;

Formal Analysis, S.G., J.D.F., I.M.C., B.G.; Investigation, S.G., J.D.F., E.H.; Writing – Original

Draft, S.G.; Writing – Review & Editing, S.G., B.G., J.D.F., E.H., I.M.C.; Supervision, B.G.,

I.M.C.

Declaration of Interests

The authors declare no competing interests.

References

454
455 Abernathy, and Glaunsinger (2015). Emerging roles for RNA degradation in viral replication and
456 antiviral defense. Virology 479, 600–608.

457
458 Abernathy, Gilbertson, Alla, and Glaunsinger (2015). Viral Nucleases Induce an mRNA
459 Degradation-Transcription Feedback Loop in Mammalian Cells. Cell Host & Microbe 18, 243–
460 253.

Adler, Messerle, Wagner, and Koszinowski (2000). Cloning and Mutagenesis of the Murine Gammaherpesvirus 68 Genome as an Infectious Bacterial Artificial Chromosome. Journal of Virology 74, 6964–6974.

Bablanian, R., Goswami, S.K., Esteban, M., Banerjee, A.K., and Merrick, W.C. (1991). Mechanism of selective translation of vaccinia virus mRNAs: differential role of poly(A) and initiation factors in the translation of viral and cellular mRNAs. J. Virol. *65*, 4449–60.

Barry, Ingolia, and Vance (2017). Global analysis of gene expression reveals mRNA superinduction is required for the inducible immune response to a bacterial pathogen. ELife 6, e22707.

Borah, Nichols, Hassman, Kedes, and Steitz (2012). Tracking expression and subcellular localization of RNA and protein species using high-throughput single cell imaging flow cytometry. RNA 18, 1573–1579.

Braun, and Young (2014). Coupling mRNA Synthesis and Decay. Molecular and Cellular Biology 34, 4078–4087.

Braun, Huntzinger, Fauser, and Izaurralde (2011). GW182 proteins directly recruit cytoplasmic deadenylase complexes to miRNA targets. Molecular Cell 44, 120–33.

Burgess, H.M., and Gray, N.K. (2010). mRNA-specific regulation of translation by poly(A)-binding proteins. Biochem. Soc. Trans. *38*, 1517–22.

Burgess, and Mohr (2015). Cellular 5′-3′ mRNA Exonuclease Xrn1 Controls Double-Stranded RNA Accumulation and Anti-Viral Responses. Cell Host & Microbe 17, 332–344.

Chapman, Costantino, Rabe, Moon, Wilusz, Nix, and Kieft (2014). The Structural Basis of Pathogenic Subgenomic Flavivirus RNA (sfRNA) Production. Science 344, 307–310.

Chekulaeva, M., Mathys, H., Zipprich, J.T., Attig, J., Colic, M., Parker, R., and Filipowicz, W. (2011). miRNA repression involves GW182-mediated recruitment of CCR4-NOT through conserved W-containing motifs. Nat. Struct. Mol. Biol. *18*, 1218–26.

Covarrubias, Gaglia, Kumar, Wong, Jackson, and Glaunsinger (2011). Coordinated Destruction of Cellular Messages in Translation Complexes by the Gammaherpesvirus Host Shutoff Factor and the Mammalian Exonuclease Xrn1. PLoS Pathogens 7, e1002339.

Covarrubias, S., Richner, J., Clyde, K., Lee, Y., and Glaunsinger, B. (2009). Host Shutoff Is a Conserved Phenotype of Gammaherpesvirus Infection and Is Orchestrated Exclusively from the Cytoplasm. Journal of Virology 83, 9554–9566.

502 Darzacq, X., Shav-Tal, Y., Turris, V. de, Brody, Y., Shenoy, S.M., Phair, R.D., and Singer, R.H.
503 (2007). In vivo dynamics of RNA polymerase II transcription. Nat. Struct. Mol. Biol. *14*, 796–
504 806.
505
Fabian, Cieplak, Frank, Morita, Green, kumar, Nagar, Yamamoto, Raught, Duchaine, et al.
(2011). miRNA-mediated deadenylation is orchestrated by GW182 through two conserved
motifs that interact with CCR4–NOT. Nature Structural and Molecular Biology nsmb.2149.

Fatscher, T., Boehm, V., Weiche, B., and Gehring, N.H. (2014). The interaction of cytoplasmic
poly(A)-binding protein with eukaryotic initiation factor 4G suppresses nonsense-mediated
mRNA decay. RNA *20*, 1579–92.

512
513 Federspiel, J.D., & Cristea, I.M. (2018). Considerations for identifying endogenous protein
514 complexes from tissue via immunoaffinity purification and quantitative mass spectrometry.
515 Methods in Molecular Biology. (Under Review)
516
517 Gaglia, and Glaunsinger (2010). Viruses and the cellular RNA decay machinery. Wiley
518 Interdisciplinary Reviews: RNA 1, 47–59.
519
520 Gaglia, Covarrubias, Wong, and Glaunsinger (2012). A Common Strategy for Host RNA
521 Degradation by Divergent Viruses. Journal of Virology 86, 9527–9530.
522
523 Gerstberger, Hafner, and Tuschl (2014). A census of human RNA-binding proteins. Nature
524 Reviews Genetics 15, 829–845.
525
526 Haimovich, Medina, Causse, Garber, Millán-Zambrano, Barkai, Chávez, Pérez-Ortín, Darzacq,
527 and Choder (2013). Gene Expression Is Circular: Factors for mRNA Degradation Also Foster
528 mRNA Synthesis. Cell 153, 1000–1011.
529
530 Harb, Becker, Vitour, Baron, Vende, Brown, Bolte, Arold, and Poncet (2008). Nuclear
531 Localization of Cytoplasmic Poly(A)-Binding Protein upon Rotavirus Infection Involves the
532 Interaction of NSP3 with eIF4G and RoXaN. Journal of Virology 82, 11283–11293.

533 Heidemann, M., Hintermair, C., Voß, K., and Eick, D. (2013). Dynamic phosphorylation patterns
534 of RNA polymerase II CTD during transcription. Biochim. Biophys. Acta *1829*, 55–62.
535
536 Huntzinger, and Izaurralde (2011). Gene silencing by microRNAs: contributions of translational
537 repression and mRNA decay. Nature Reviews. Genetics 12, 99–110.
538
539 Kawahara, Imai, Imataka, Tsujimoto, Matsumoto, and Okano (2008). Neural RNA-binding
540 protein Musashi1 inhibits translation initiation by competing with eIF4G for PABP. The Journal
541 of Cell Biology 181, 639–653.
542
543 Kumar, and Glaunsinger (2010). Nuclear Import of Cytoplasmic Poly(A) Binding Protein

Restricts Gene Expression via Hyperadenylation and Nuclear Retention of mRNA. Molecular and Cellular Biology 30, 4996–5008.

Kumar, Shum, and Glaunsinger (2011). Importin α-Mediated Nuclear Import of Cytoplasmic Poly(A) Binding Protein Occurs as a Direct Consequence of Cytoplasmic mRNA Depletion. Molecular and Cellular Biology 31, 3113–3125.

Larimer, F., and Stevens, A. (1990). Disruption of the gene XRN1, coding for a 5'----3' exoribonuclease, restricts yeast cell growth. Gene *95*, 85–90.

Lee, and Glaunsinger (2009). Aberrant Herpesvirus-Induced Polyadenylation Correlates With Cellular Messenger RNA Destruction. PLoS Biology 7, e1000107.

Louder, R.K., He, Y., López-Blanco, J.R.R., Fang, J., Chacón, P., and Nogales, E. (2016). Structure of promoter-bound TFIID and model of human pre-initiation complex assembly. Nature *531*, 604–9.

Marceau, Puschnik, Majzoub, Ooi, Brewer, Fuchs, Swaminathan, Mata, Elias, Sarnow, et al. (2016). Genetic dissection of Flaviviridae host factors through genome-scale CRISPR screens. Nature 535, 159–63.

Mattijssen, S., Arimbasseri, A., Iben, J., Gaidamakov, S., Lee, J., Hafner, M., and Maraia, R. (2017). LARP4 mRNA codon-tRNA match contributes to LARP4 activity for ribosomal protein mRNA poly(A) tail length protection. ELife 6, e28889.

McAlister, Huttlin, Haas, Ting, Jedrychowski, Rogers, Kuhn, Pike, Grothe, Blethrow, et al. (2012). Increasing the Multiplexing Capacity of TMTs Using Reporter Ion Isotopologues with Isobaric Masses. Analytical Chemistry 84, 7469–7478.

Mitchell, S.F., and Parker, R. (2014). Principles and properties of eukaryotic mRNPs. Mol. Cell *54*, 547–58.

Moon, Anderson, Kumagai, Wilusz, Akira, Khromykh, and Wilusz (2012). A noncoding RNA produced by arthropod-borne flaviviruses inhibits the cellular exoribonuclease XRN1 and alters host mRNA stability. RNA 18, 2029–2040.

Müller-McNicoll, and Neugebauer (2013). How cells get the message: dynamic assembly and function of mRNA–protein complexes. Nature Reviews Genetics 14, nrg3434.

Neto, O.P., Standart, N., and Sa, C. (1995). Autoregulation of poly (A)-binding protein synthesis in vitro. Nucleic Acids Research *23*, 2198–2205.

Park, El-Guindy, Heston, Lin, Yu, Nagy, Borah, Delecluse, Steitz, and Miller (2014). Nuclear Translocation and Regulation of Intranuclear Distribution of Cytoplasmic Poly(A)-Binding Protein Are Distinct Processes Mediated by Two Epstein Barr Virus Proteins. PLoS ONE 9,

e92593.

Piron, Vende, Cohen, and Poncet (1998). Rotavirus RNA☐binding protein NSP3 interacts with eIF4GI and evicts the poly(A) binding protein from eIF4F. The EMBO Journal 17, 5811–5821.

Jean Beltran, P.M., Mathias, R.A., and Cristea, I.M. (2016). A Portrait of the Human Organelle Proteome In Space and Time during Cytomegalovirus Infection. Cell Syst *3*, 361–373.e6.

Jean Beltran, P.M., Federspiel, J.D., Sheng, X., and Cristea, I.M. (2017). Proteomics and integrative omic approaches for understanding host-pathogen interactions and infectious diseases. Mol. Syst. Biol. *13*, 922.

Puschnik, Marceau, Ooi, Majzoub, Rinis, Contessa, and Carette (2017). A Small-Molecule Oligosaccharyltransferase Inhibitor with Pan-flaviviral Activity. Cell Reports 21, 3032–3039.

Reuter, Meinel, and Sträßer (2015). The poly(A)-binding protein Nab2 functions in RNA polymerase III transcription. Genes & Development 29, 1565–1575.

Richner, J., Clyde, K., Pezda, A., Cheng, B., Wang, T., Kumar, G., Covarrubias, S., Coscoy, L., and Glaunsinger, B. (2011). Global mRNA Degradation during Lytic Gammaherpesvirus Infection Contributes to Establishment of Viral Latency. Plos Pathog *7*, e1002150.

Rivas, H.G., Schmaling, S.K., and Gaglia, M.M. (2016). Shutoff of Host Gene Expression in Influenza A Virus and Herpesviruses: Similar Mechanisms and Common Themes. Viruses *8*, 102.

Roeder, R.G. (1996). The role of general initiation factors in transcription by RNA polymerase II. Trends Biochem. Sci. 21, 327–35.

Salaun, MacDonald, Larralde, Howard, Lochtie, Burgess, Brook, Malik, Gray, and Graham (2010). Poly(A)-Binding Protein 1 Partially Relocalizes to the Nucleus during Herpes Simplex Virus Type 1 Infection in an ICP27-Independent Manner and Does Not Inhibit Virus Replication. Journal of Virology 84, 8539–8548.

Sauls, Greco, Wang, Zou, Villasmil, Qian, Cristea, and Conlon (2018). Initiating Events in Direct Cardiomyocyte Reprogramming. Cell Reports 22, 1913–1922.

Schoenberg, and Maquat (2012). Regulation of cytoplasmic mRNA decay. Nature Reviews Genetics 13, 246–259.

Sanjana, N., Shalem, O., and Zhang, F. (2014). Improved vectors and genome-wide libraries for CRISPR screening. Nat Methods 11, nmeth.3047.

Shalem, O., Sanjana, N., Hartenian, E., Shi, X., Scott, D., Mikkelsen, T., Heckl, D., Ebert, B., Root, D., Doench, J., et al. (2014). Genome-Scale CRISPR-Cas9 Knockout Screening in Human

Cells. Science 343, 84–87.

Singh, Pratt, Yeo, and Moore (2015). The Clothes Make the mRNA: Past and Present Trends in
mRNP Fashion. Annual Review of Biochemistry 84, 325–354.

Sun, Schwalb, Pirkl, Maier, Schenk, Failmezger, Tresch, and Cramer (2013). Global Analysis of
Eukaryotic mRNA Degradation Reveals Xrn1-Dependent Buffering of Transcript Levels.
Molecular Cell 52, 52–62.

Sutherland, Sobinoff, Fraser, Redgrove, Davidson, Siddall, Koopman, Hime, and McLaughlin
(2015). RNA binding protein Musashi-1 directly targets Msi2 and Erh during early testis germ
cell development and interacts with IPO5 upon translocation to the nucleus. The FASEB Journal
29, 2759–2768.

Thomas, Liu, Whangbo, McCrossan, Sanborn, Basar, Walch, and Lieberman (2015). Apoptosis
Triggers Specific, Rapid, and Global mRNA Decay with 3' Uridylated Intermediates Degraded
by DIS3L2. Cell Reports 11, 1079–89.

Vizcaíno, J.A., Deutsch, E.W., Wang, R., Csordas, A., Reisinger, F., Ríos, D., Dianes, J.A.A.,
Sun, Z., Farrah, T., Bandeira, N., et al. (2014). ProteomeXchange provides globally coordinated
proteomics data submission and dissemination. Nat. Biotechnol. *32*, 223–6

Wu, and Bag (1998). Negative Control of the Poly(A)-binding Protein mRNA Translation Is
Mediated by the Adenine-rich Region of Its 5′-Untranslated Region. Journal of Biological
Chemistry 273, 34535–34542.

Yang, Gaidamakov, Xie, Lee, Martino, Kozlov, Crawford, Russo, Conte, Gehring, et al. (2011).
La-Related Protein 4 Binds Poly(A), Interacts with the Poly(A)-Binding Protein MLLE Domain
via a Variant PAM2w Motif, and Can Promote mRNA Stability. Molecular and Cellular Biology
31, 542–556.

Figure Legends

Figure 1. RNA binding proteins are translocated from the cytoplasm to the nucleus in cells

undergoing enhanced cytoplasmic mRNA decay. (A) Diagram depicting the experimental

setup. (B) Venn diagram of nuclear proteins that are specifically and significantly (p < 0.05)

enriched in muSOX-expressing cells compared to D219A-expressing cells that also show either

no change or a decrease in cytoplasmic abundance. (C) Gene ontology molecular function

overrepresentation analysis by Pantherdb, graphed according to their P value. (D) Graphs

showing the nuclear and cytoplasmic distribution of poly(A) binding proteins from the TMT-MS

data. Graphs display the mean with SEM of 3 biological replicates. (E) Western blot of nuclear,

cytoplasmic, and whole cell fractions of NIH3T3 fibroblasts mock infected or infected with WT

or R443I MHV68 for 24h. GAPDH and histone H3 serve as fractionation and loading controls.

Shown is a representative example of 3 biological replicates.

Figure 2. Nuclear translocation of RNA binding proteins is dependent on mRNA

degradation by Xrn1. (A) Pie chart showing the percent of shuttling proteins that fail to

translocate in Xrn1 KO cells. (B) Heat map depicting the average nuclear abundance in WT or

Xrn1 KO HEK293T cells of the 67 significantly shifted proteins in samples expressing muSOX

or D219A, relative to the empty vector control. (C) Western blots of nuclear, cytoplasmic, and

whole cell fractions of WT (left panel) or Xrn1 KO (right panel) HEK293T cells transfected with

the indicated plasmid. GAPDH and histone H3 serve as fractionation and loading controls. (D)

Confocal microscopy and quantification of WT or Xrn1 KO HEK293T cells transfected with

GFP or GFP-muSOX, showing signals for DAPI stained nuclei (blue), PABPC (red), GFP

(green), and the merged images (overlay). Arrow heads point to representative GFP-muSOX

expressing cells. The number of cells displaying either cytoplasmic or nuclear PABPC

localization by immunofluorescence (IFA) was quantified for WT or Xrn1 KO cells expressing

GFP-muSOX. A total of 75 GFP-muSOX WT cells and 80 GFP-muSOX Xrn1 KO cells were

counted. The graph displays individual biological replicates as dots, with the mean and SEM.

Figure 3. Xrn1 knockout leads to subcellular redistribution of proteins functionally associated with RNA. (A) The number of proteins that are differentially expressed in Xrn1 knockout (KO) cells from the nucleus (149) and the cytoplasm (158). Gene ontology molecular function overrepresentation analysis by Pantherdb is shown for each compartment, graphed according to their false discovery rate (FDR). (B) Graphs showing the nuclear and cytoplasmic distribution of decapping-related proteins from the TMT-LC/MS-MS data. Graphs display the mean with SEM of 3 biological replicates. (C) Heatmap depicting the Log_2 abundance ratio in Xrn1 KO HEK293T cells compared to WT HEK293T cells of proteins identified as Xrn1 interactors using the BioGRID database. Proteins with a significant difference in abundance between WT and Xrn1 KO are listed in red. (D) Graph of nuclear and cytoplasmic distribution of GW182 from the TMT-LC/MS-MS data. Graph displays the mean with SEM of 3 biological replicates. (E) Western blot of nuclear, cytoplasmic, and whole cell fractions of WT and Xrn1 KO HEK293T cells. GAPDH and histone H3 serve as fractionation and loading controls.

Figure 4. LARP4 translocates to the nucleus in a PABPC-dependent manner. (A) STRING network of reported protein-protein interactions between the 67 proteins that shuttle in muSOX-expressing cells. Medium and high confidence interactions are shown with thin and thick connector lines, respectively. (B, C) Western blots of nuclear and cytoplasmic fractions of vector- or muSOX-transfected HEK293T cells treated with the indicated siRNA. GAPDH and histone H3 serve as fractionation and loading controls.

Figure 5. PABPC depletion prevents muSOX-induced repression of RNAPII recruitment. (A, B) HEK293T cells treated with siRNAs targeting PABPC1 and 4 (A), LARP4 (B), or non-targeting scramble siRNAs were subsequently transfected with either empty vector or muSOX, then subjected to chromatin immunoprecipitation (ChIP) using antibodies to RNAPII or IgG.

Western blots showing protein levels of PABPC1, PABPC4, and Larp4 after siRNA depletion

are shown in the lower panels, along with a GAPDH loading control. (C) Western blots of

nuclear and cytoplasmic fractions of HEK293T cells transfected with an empty vector or a

plasmid containing FLAG-PABPC1. GAPDH and histone H3 serve as fractionation and loading

controls. (D) HEK293T cells transfected with either empty vector or FLAG-PABPC1 were

subjected to ChIP using antibodies to RNAPII or IgG. (E) WT or Xrn1 KO HEK293T cells

transfected with either empty vector or FLAG-PABPC1 alone or together with muSOX were

subjected to ChIP using antibodies to RNAPII or IgG. Purified chromatin in each of the above

experiments was quantified by qPCR. Western blots showing the levels of Xrn1 in WT or

Xrn1KO HEK293Ts are shown, along with a GAPDH loading control. All graphs display

individual biological replicates as dots, with the mean and SEM. Statistical significance was

determined using Student's t test *p <0.05 **p <0.005 ***p <0.0005.

Figure 6. Nuclear translocation of PABPC selectively impacts early stages of transcription.

(A) HEK293T cells transfected with empty vector, muSOX, or FLAG-PABPC1 were subjected

to ChIP using antibodies to RNAPII or IgG at the indicated gene matched promoters and exons.

(B) ChIP using antibodies to serine 5-phosphorylated (Ser5P) RNAPII or IgG at gene promoters.

The level of Ser5P RNAPII was determined by dividing the Ser5P values over the total RNAPII

values within the same region of the gene in HEK293T cells transfected with empty vector,

muSOX, or FLAG-PABPC1. (C) ChIP was performed as described in (B), but using antibodies

to serine 2-phosphorylated (Ser2P) RNAPII or IgG at gene exons. The level of Ser2P RNAPII

was determined by dividing the Ser2P values over the total RNAPII values within the same

region of the gene. (D) ChIP was performed as described in (B), but using antibodies to TATA-

binding protein (TBP) or IgG at gene promoters. (E) ChIP was performed as described in (B),

but using antibodies to the POLR3A subunit of RNAPIII or IgG. In each experiment, chromatin

was quantified by qPCR and all graphs display individual biological replicates as dots, with the

mean and SEM. Statistical significance was determined using Student's t test $*p < 0.05$ $**p$

<0.005 $***p < 0.0005$. (F) Model summarizing the impact of SOX and Xrn1-driven mRNA

degradation on RBP trafficking and RNAPII transcription. See text for details.

Materials and Methods

Key Resources Table										
Reagent type (species)	Designation	Source or referenc	Identifiers	Additional information						
strain, strain backgroun	MHV68	PMID: 10888635	NCBI_refseq ID:NC_0018	Koszinowski Lab						
strain, strain backgroun	R443I MHV68	PMID: 21811408		Glaunsinger Lab						
cell line (Homo sapiens	HEK293T	American Type Cu	Cat# CRL-11268; RRID: CVCL_1926							
cell line (Mus musculus	NIH3T3	American Type Cu	Cat# CRL-1658; RRID: CVCL_0594							
transfected construct (	Xrn1 knockout (KO) cel	This paper		HEK293T clone stably expressing Cas9 and Xrn1 single-guide RNA.						
transfected construct (	Cas9-expressing WT ce	This paper		HEK293T clone stably expressing Cas9 alone with no guide RNA.						
recombinant DNA reag	GFP-muSOX (plasmid)	This paper		Progenitors: pcDNA3-HA-muSOX (Covarrubius et al., 2009); Gateway vect						
recombinant DNA reag	Thy1.1-muSOX (plasmi	This paper		Progenitors: GFP-muSOX						
recombinant DNA reag	Thy1.1-muSOX D219A (	This paper		Progenitors: Thy1.1-muSOX						
recombinant DNA reag	Thy1.1-GFP (plasmid)	This paper		Progenitors: Thy1.1-muSOX						
recombinant DNA reag	pCDEF3-FLAG-PABPC1	PMID: 20823266		Glaunsinger Lab						
recombinant DNA reag	lentiCas9-Blast (lentiv	Addgene; PMID: 2	52962							
recombinant DNA reag	lentiGuide-Puro (lenti	Addgene; PMID: 2	52963							
antibody	Mouse monoclonal ant	Santa Cruz Biotech	Clone 10 E 10; SC32318	IFA (1:25)						
antibody	Rabbit polyclonal anti-	Thermofisher	PA5-58727	IFA (1:200); Western (1:1000)						
antibody	Rabbit polyclonal anti-	Cell Signaling Tech	4992S	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Bethyl	A301-466A-M	Western (1:1000)						
antibody	Mouse monoclonal ant	Abcam	Clone 6C5; ab8245	Western (1:5000)						
antibody	Rabbit monoclonal ant	Cell Signaling Tech	Clone D1H2; 4499S	Western (1:2000)						
antibody	Rabbit polyclonal anti-	Abcam	ab124789	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Bethyl	A303-996A-T	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Abcam	ab52865	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Abcam	ab71415	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Novus Biologicals	NBP15707220	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Cell Signaling Tech	4241T	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Life Technologies	PA543172	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Life Technologies	PA535053	Western (1:1000)						
antibody	Rabbit polyclonal anti-	This paper		Western (1:1000)						
antibody	Rabbit polyclonal anti-	Abcam	ab10530	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Abcam	ab131223	Western (1:1000)						
antibody	Mouse monoclonal ant	Abcam	Clone 4B6; ab70522	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Bethyl	A300-460	Western (1:1000)						
antibody	Rabbit polyclonal anti-	Bethyl	A302-597A-M	Western (1:1000)						
antibody	Mouse monoclonal ant	BioLegend	8WG16	Western (1:2000); ChIP (10 mg)						
antibody	Rabbit polyclonal anti-	Abcam	ab28175	Western (1:2000); ChIP (10 mg)						
antibody	Rabbit polyclonal anti-	Abcam	ab5131	ChIP (10mg)						
antibody	Rabbit polyclonal anti-	Abcam	ab5095	ChIP (10mg)						
antibody	Rabbit polyclonal anti-	Abcam	ab96328	ChIP (10mg)						
sequence-based reage	siPABPC1	PMID: 20823266		Custom siRNA; see Table S5 for sequence						
sequence-based reage	siPABPC4	PMID: 20823266		Custom siRNA; see Table S5 for sequence						
sequence-based reage	siLARP4	Dharmacon	M-016523-00-0020							
sequence-based reage	siCHD3	Dharmacon	M-023015-01-0020							
sequence-based reage	siMSI1	Dharmacon	M-011338-01-0010							

Plasmids

Primers used for cloning are listed in Table S5. MHV68 muSOX was cloned into the

Gateway entry vector pDON207 (Invitrogen), and then transferred into the Gateway-compatible

peGFP-C1 destination vector to generate GFP-muSOX. Thy1.1-muSOX was generated by

Infusion cloning (Clontech) of Thy1.1 (CD90.1) followed by a self-cleaving 2A peptide from

foot-and-mouth disease virus in place of GFP into the Nhe1 and SacII restriction enzyme sites of

GFP-muSOX. The D219A muSOX mutant was made using Quikchange site-directed

mutagenesis (Agilent). Thy1.1-GFP was created with Infusion cloning to insert GFP back into

the vector with the BamHI and EcoRI restriction enzyme sites to replace muSOX with GFP.

pCDEF3-Flag-PABPC1 was described previously (Kumar, 2010). The Cas9 (lentiCas9-Blast)

and sgRNA (lentiGuide-Puro) viral vectors were made as previously described (Sanjana et al.,

2014; Shalem et al., 2014). The Xrn1 sgRNA was chosen using the Broad sgRNA design website

(Doench et al., 2014).

Cells and transfections

NIH3T3 cells and HEK293T cells, both from ATCC and obtained through the UC

Berkeley Tissue Culture Facility, were maintained in DMEM (Invitrogen) supplemented with

10% fetal bovine serum. Cell lines were authenticated by STR analysis, and determined to be

free of mycoplasma by PCR screening. DNA transfections were carried out in HEK293T cells at

70% confluency in 15cm plates with 25μg DNA using PolyJet (SignaGen) for 24h. For small

interfering RNA (siRNA) transfections, HEK293T cells were transfected twice over 48h with

100μM siRNA using Lipofectomine RNAiMAX (Thermo Fisher), whereupon the cells were

transfected with the indicated DNA plasmid for an additional 24h. Non-targeting scramble

siRNAs, LARP4, MSI1, CHD3, and TRIM32 siRNAs were obtained from Dharmacon

(Scramble: D-001206-13-50, LARP4: M-016523-00-0020, MSI1: M-011338-01-0010, CHD3:

M-023015-01-0020, TRIM32: M-006950-01-0010). PABPC1 and PABPC4 siRNAs have been

previously described and are listed in Table S5. (Kumar, 2010; Lee, 2009).

The Xrn1 knockout clone and control Cas9-expressing cells were made by transducing

HEK293T cells as previously described (Sanjana et al., 2014; Shalem et al., 2014). Briefly, lenti-

Cas9-blast lentivirus was spinfected onto a monolayer of HEK293T cells, which were then

incubated with 20μg/ml blasticidin to remove non-transduced cells. These Cas9-expressing

HEK293T cells were then spinfected with lentivirus made from lentiGuide-Puro containing the

Xrn1 sgRNA sequence and selected with 1μg/ml puromycin. The pool of Xrn1 knockout cells

was then single-cell cloned in 96-well plates and individual clones were screened by western blot

to determine knockout efficiency.

Pure populations of cells expressing muSOX were generated using the Miltenyi Biotec

MACS cell separation system. HEK293T cells were transfected with either Thy1.1-GFP,

Thy1.1-muSOX, or Thy1.1-muSOX D219A for 24h, whereupon cells were washed twice with

PBS and cell pellets were resuspended in 95μl auto-MACS rinsing buffer supplemented with

0.5% FBS and incubated with 3μl anti-CD90.1 microbeads on ice for 10-15 min, and mixed by

flicking the tube every 5 min. Cells were then magnetically separated according to the

manufacturer's instructions. Thy1.1 positive cells were used in all downstream experiments

unless otherwise stated.

Viruses and infections

The MHV68 bacterial artificial chromosome (BAC), and the construction of the R443I

muSOX mutant were previously described (Adler et al., 2000; Richner et al., 2011). MHV68 was

produced by transfecting NIH3T3 cells in 6-well plates with 2.5μg BAC DNA using Mirus

TransIT-X2 (Mirus Bio) for 24h, whereupon the cells were split into a 10cm dish and harvested

after 5-7 days, once all the cells were green and dead. Virus was amplified in NIH 3T12 cells and

titered by plaque assay. Cells were infected with MHV68 at an MOI of 5 for 24h.

Immunofluorescence assays

Cells were plated on coverslips coated with 100ug/mL poly-L-lysine and transfected at

70% confluency with either GFP or GFP-muSOX for 24h. Transfected cells were fixed in 4%

formaldehyde, permeabilized with ice-cold methanol, and incubated with blocking buffer [1%

Triton X-100, 0.5% Tween-20, 3% Bovine Serum Albumin] prior to incubation with mouse

monoclonal PABPC diluted 1:25 (Santa Cruz Biotechnologies, 10E10) or rabbit polyclonal

LARP4 diluted 1:200 (Thermo Fisher) in blocking buffer at 4°C overnight, followed by

incubation with Alexa Fluor 594-conjugated goat anti-mouse, or anti-rabbit secondary antibody

(Thermo Fisher, 1:1000) and DAPI (Pierce, 1:1000). Coverslips were mounted on slides using

Vectashield hard-set mounting medium (VectorLabs) and imaged by confocal microscopy on a

Zeiss LSM 710 AxioObserver microscope.

Fractionation

HEK293T cells were fractionated using the REAP method (Nabbi and Riabowol, 2015).

Briefly, cells were washed twice with ice-cold PBS and the cell pellet was lysed in 0.1% NP-40

PBS lysis buffer. The nuclei were then isolated by differential centrifugation at 10,000 x g for 10

sec and the supernatant retained as the cytoplasmic fraction. For western blotting, the nuclei were

sonicated in 0.1% NP-40 PBS lysis buffer.

Cell lysis and protein digestion

WT Cas9-HEK293T cells were transfected with Thy1.1-GFP, Thy1.1-muSOX, or

Thy1.1-muSOX D219A. Xrn1 knockout HEK293T cells were transfected with Thy1.1-GFP or

Thy1.1-muSOX for 24h, followed by Thy1.1 separation. Separated cells were then fractionated

as described above, and nuclear pellets were snap-frozen in liquid nitrogen. Cytoplasmic

fractions were concentrated using an Amicon ultra filtration unit with a molecular weight cutoff

of 3kDa (Millipore) and exchanged into a 50mM NH_4HCO_3, 2% Deoxycholate buffer and then

snap frozen in liquid nitrogen. The nuclear pellets were lysed in 200µL of 100mM Tris-HCl, pH

8.0, 4% SDS, 1mM EDTA preheated to 70°C. Cytoplasmic fractions were thawed and adjusted

to 1% SDS with a 10% SDS solution. Complete lysis of samples was achieved via five

successive rounds of heating at 95°C for 3 min followed by sonication for 10 sec in a cup horn

sonicator set on 1 sec pulses at medium output. Protein amounts were assessed by BCA protein

assay (Pierce) and 50 µg of protein from each sample was simultaneously reduced and alkylated

with 20mM tris(2-carboxyethyl)phosphine (Pierce) and chloroacetamide respectively for 20 min

at 70°C. Protein samples were then cleaned up by methanol-chloroform precipitation (Wessel et

al., 1984; Federspiel and Cristea, 2018). LC-MS grade methanol, chloroform, and water (at a

4:1:3 ratio) were added to the sample with vortexing following each addition. The samples were

spun at $2,000 \times g$ for 5 min at room temperature and the top phase was removed. Three volumes

of cold methanol were then added and the samples were spun at $9,000 \times g$ for 2 min at 4°C. All

liquid was removed and the protein pellets were washed with five volumes of cold methanol and

then spun at $9,000 \times g$ for 2 min at 4°C. All liquid was removed again and the dried protein

pellets were resuspended in 50mM HEPES pH 8.5 at a 0.5µg/µL concentration. Trypsin (Pierce)

was added at a 1:50 trypsin:protein ratio and the samples were incubated at 37°C overnight.

TMT labeling

Digested samples were concentrated by speed vac to one half the original volume prior to

labeling and adjusted to 20% acetonitrile (ACN). All three biological replicates were labeled

concurrently with a 10-plex TMT kit (Thermo Fisher Scientific) as in (Sauls et al., 2018). The

TMT reagents (0.8mg per channel) were dissolved in 42µL of anhydrous ACN and 14µL of this

was added to each sample following the scheme in Figure 1A and allowed to react at RT for 1h.

The labeling was quenched by the addition of hydroxylamine to a final 0.5% (v/v) concentration

followed by incubation at RT for 15 min. Labeled peptides were pooled at equal peptide amounts

thereby generating three 10-plex experiments, each of which was an individual biological

replicate. An initial test mix for each replicate was analyzed, and the apparent peptide ratios were

determined. Mixing ratios were adjusted using the information from the test mix to correct for

sample losses and generate mixes with equal peptide amounts per channel.

Peptide fractionation

Pooled peptides were acidified and fractionated by 2D StageTip (Sauls et al., 2018).

Peptides were first desalted via C18 StageTips to remove unreacted TMT reagent by washing the

bound peptides with 5% ACN, 0.5% formic acid (FA) and then eluting the peptides in 70%

ACN, 0.5% FA. The eluted peptides were then bound to SCX StageTips and eluted in four

fractions with sequential elution (100µL) as follows: (1) 0.05 M ammonium formate/20% ACN,

(2) 0.05 M ammonium acetate/20% ACN, (3) 0.05 M ammonium bicarbonate/20% ACN, and (4)

0.1% ammonium hydroxide/20% ACN. Each of these fractions were diluted 1:1 with 1%

trifluoroacetic acid and further fractionated by SDB-RPS StageTips with sequential elution

(50µL) into three fractions as follows: (1) 0.2 M ammonium formate/0.5% FA/60% ACN, (2)

0.2 M ammonium acetate/0.5% FA/60% ACN, (3) 5% ammonium hydroxide/80% ACN. The

resulting 12 fractions for each 10-plex experiment were dried *in vacuo* and resuspended in 5µL

of 1% FA, 1% ACN in water.

LC-MS/MS analysis

Peptides (2µL) were analyzed by LC-MS/MS using a Dionex Ultimate 3000 UPLC

coupled online to an EASYSpray ion source and Q Exactive HF. Peptides were separated on an

EASYSpray C18 column (75µm x 50cm) heated to 50°C using a linear gradient of 5% ACN to

42% ACN in 0.1% FA over 150min at a flow rate of 250nL/min and ionized at 1.7kv. MS/MS

analysis was performed as follows: an MS1 scan was performed from 400 to 1800 m/z at

120,000 resolution with an automatic gain control (AGC) setting of 3e6 and a maximum

injection time (MIT) of 30ms recorded in profile. The top 18 precursors were then selected for

fragmentation and MS2 scans were acquired at a resolution of 60,000 with an AGC setting of

2e5, a MIT of 105ms, an isolation window of 0.8 m/z, a fixed first mass of 100 m/z, normalized

collision energy of 34, intensity threshold of 1e5, peptide match set to preferred, and a dynamic

exclusion of 45s recorded in profile.

Informatic analysis of TMT data

MS/MS data were analyzed by Proteome Discoverer (Thermo Fisher Scientific,

v2.2.0.388). The nuclear channels (126-128C) and cytoplasmic channels (129N-131) were

analyzed in separate Proteome Discoverer studies to not bias the quantitation due to the expected

protein expression differences between these two compartments. The Spectrum Files RC node

was utilized to perform post-acquisition mass recalibration and the recalibrated spectra were

passed to Sequest HT where two successive rounds of searching were employed against a

Uniprot human database appended with common contaminants (2016-04, 22,349 sequences).

Both search rounds required 5ppm accuracy on the precursor and 0.02Da accuracy on the

fragments and included static carbamidomethyl modifications to cysteine, static TMT additions

to peptide N-termini and lysine residues, dynamic oxidation of methionine, dynamic deamidation

of asparagine, and dynamic methionine loss and acetylation of protein n-termini. The first

Sequest HT search was for fully tryptic peptides only and any unmatched spectra were sent to a

second Sequest HT search, which allowed semi-tryptic peptide matches. All matched spectra

were scored by Percolator and reporter ion signal-to-noise (S/N) values were extracted (The et

al., 2016). The resulting peptide spectrum matches were parsimoniously assembled into a set of

identified peptide and protein identifications with a false discovery rate of less than 1% for both

the peptide and protein level and at least 2 unique peptides identified per protein. TMT reporter

ion quantification was performed for unique and razor peptides with an average S/N of at least

10 and a precursor co-isolation threshold of less than 30% which did not contain a variable

modification. Reporter ion values were normalized to the total detected signal in each channel

and protein abundances were calculated as the sum of all normalized reporter ion values for each

channel in each protein. Missing values were input using the low abundance resampling

algorithm. The reporter ion values for the empty vector WT samples (channels 126 and 129N)

were set as 100 and the other channels were scaled to this value. Statistically differential proteins

were assessed via a background based ANOVA analysis implemented in Proteome Discoverer.

Proteins and associated TMT reporter ion abundances and adjusted p-values from the ANOVA

analysis were exported to Excel for further analysis. The mass spectrometry proteomics data

reported in this paper have been deposited at the ProteomeXchange Consortium via the PRIDE

partner repository (Vizcaíno et al., 2014). The PRIDE accession number is PXD009487.

Gene Ontology analysis and informatic software used

Differential proteins (adjusted p-value ≤ 0.05) were analyzed via over representation

analysis (www.pantherdb.org) for associated gene ontology enrichments (Mi et al., 2016).

Example proteins of different classes, along with all heatmaps, were graphed in GraphPad Prism

v7.

Western blotting

Nuclear, cytoplasmic, and whole cell lysates were quantified by Bradford assay and

resolved by SDS-PAGE and western blotted with antibodies against PABPC (Cell Signaling,

1:1000), PABPC4 (Bethyl, 1:1000), LARP4 (Thermo Fisher, 1:1000), Gapdh (Abcam, 1:3000),

Histone H3 (Cell Signaling, 1:2000), LYRIC (Abcam, 1:1000), RRBP1 (Bethyl, 1:1000), MSI1

(Abcam, 1:1000), Lin28b (Abcam, 1:1000), CHD3 (Cell Signaling, 1:1000), RPP20 (Novus,

1:1000), THOC6 (Life Technologies, 1:1000), PNN (Life Technologies, 1:1000), EXO4 (rabbit

polyclonal produced using recombinant EXO4 with an MBP tag, 1:1000), NPM (Abcam,

1:1000), GW182 (Abcam, 1:1000), DDX6 (Bethyl, 1:1000), DCP2 (Bethyl, 1:1000), TRIM32

(Abcam, 1:1000), RNAPII Rpb1 (BioLegend, 1:2000), TBP (Abcam, 1:2000).

Chromatin immunoprecipitation (ChIP)

ChIP was performed on 15 cm plates of HEK293T cells transfected twice 4 h apart with

the indicated plasmid DNA. 24 h after the first transfection, cells were crosslinked in 1%

formaldehyde in PBS for 10 min at room temperature, quenched in 0.125 M glycine, and washed

twice with ice-cold PBS. Crosslinked cell pellets were mixed with 1 ml ice-cold ChIP lysis

buffer (5mM PIPES pH 8.0, 85mM KCl, 0.5% NP-40) and incubated on ice for 10 min,

whereupon the lysate was dounce homogenized to release nuclei and spun at 1.5 x g for 5 min at

4°C. Nuclei were then resuspended in 500 µl of nuclei lysis buffer (50mM Tris-HCl pH 8.0,

0.3% SDS, 10mM EDTA) and rotated for 10 min at 4°C followed by sonication using a QSonica

Ultrasonicator with a cup horn set to 75 amps for 20 min total (5 min on, 5 min off). Chromatin

was spun at 16,000 x *g* for 10 min at 4°C and the pellet was discarded. 100µl of chromatin was

diluted 1:5 in ChIP dilution buffer (16.7 mM Tris-HCl pH 8.0, 1.1% Triton X-100, 1.2 mM

EDTA, 167 mM NaCl) and incubated with 10 µg mouse monoclonal anti-RNAPII (BioLegend,

8WG16), rabbit IgG (Fisher Scientific), rabbit polyclonal anti-RNAPII phospho S5 (Abcam

ab5131), rabbit polyclonal anti-RNAPII phospho S2 (Abcam ab5095), rabbit polyclonal anti-

TBP (Abcam ab28175), or rabbit polyclonal anti-POLR3A (Abcam ab96328) overnight,

whereupon samples were rotated with 20 µl protein G dynabeads (with mouse antibodies), or

20µl mixed protein G and A dynabeads (with rabbit antibodies) (Thermofisher) for 2 h at 4°C.

Beads were washed with low salt immune complex (20 mM Tris pH 8.0, 1% Triton-x-100, 2

mM EDTA, 150 mM NaCl, 0.1% SDS), high salt immune complex (20 mM Tris pH 8.0, 1%

Triton-x-100, 2 mM EDTA, 500 mM NaCl, 0.1% SDS), lithium chloride immune complex

(10mM Tris pH 8.0, 0.25 M LiCl, 1% NP-40, 1% Deoxycholic acid, 1 mM EDTA), and Tris-

EDTA for 5 min each at 4°C with rotation. DNA was eluted from the beads using 100 µl of

elution buffer (150 mM NaCl, 50 µg/ml proteinase K) and incubated at 50°C for 2h, then 65°C

overnight. DNA was purified using a Zymo Oligo Clean & Concentrator kit. Purified DNA was

quantified by qPCR using iTaq Universal SYBR Mastermix (BioRad) with the indicated primers

(Table S5). Each sample was normalized to its own input.

Replicates

In this study, individual biological replicates are experiments performed separately on

biologically distinct samples representing identical conditions and/or time points. For cell

culture-based assays, this means that the cells are maintained in different flasks. Technical

replicates are experiments performed on the same biological sample multiple times. See **Figure**

Legends for the number of experimental replicates performed for each experiment. No outliers

were encountered in this study. Criteria for the inclusion of data was based on the performance

of positive and negative controls within each experiment.

Supplementary Information

Figure 1-figure supplement 1. RNA binding proteins are translocated from the cytoplasm

to the nucleus in cells undergoing enhanced mRNA decay. (A) Western blots of nuclear,

cytoplasmic, and whole cell fractions of HEK293T cells transfected with either empty vector

control, WT muSOX, or D219A muSOX. Shown are two independent biological replicates.

GAPDH and histone H3 serve as fractionation and loading controls. (B, C) Graphs showing

poly(U) (B) and 3' UTR (C) binding protein distributions in the nucleus and cytoplasm from the

TMT-LC/MS-MS data. Graphs display the mean with SEM of 3 biological replicates.

Figure 2-figure supplement 2. Nuclear translocation of RNA binding proteins is dependent

on mRNA degradation by Xrn1. (A) Western blots showing the levels of Xrn1 or the GAPDH

loading control in WT or Xrn1 KO HEK293T cells. (B) WT or Xrn1 KO HEK293T cells

transfected with either empty vector or muSOX were subjected to ChIP using antibodies to

RNAPII or IgG. Purified chromatin was quantified by qPCR. Graphs display individual

biological replicates as dots, with the mean and SEM. Statistical significance was determined

using Student's t test *$p < 0.05$ **$p < 0.005$ ***$p < 0.0005$. (C) Growth curve of WT or Xrn1 KO

HEK293T cells. Statistical significance was determined using Student's t test *$p < 0.05$ **p

< 0.005 ***$p < 0.0005$. (D) Confocal microscopy of WT or Xrn1 KO HEK293T cells transfected

with GFP or GFP-muSOX, showing signals for DAPI stained nuclei (blue), LARP4 (red), GFP

(green), and the merged images (overlay). Arrow heads point to representative GFP-muSOX

expressing cells.

Figure 3-figure supplement 3. Xrn1 knockout leads to subcellular redistribution of proteins

functionally associated with RNA. (A) Graphs showing the distribution of proteins of the

oligosaccharyltransferase (OST) complex in the nucleus and cytoplasm from the TMT-LC/MS-

MS data. Graphs display the mean with SEM of 3 biological replicates. (B) mRNA levels from

WT and Xrn1 KO HEK293T cells were measured by RT-qPCR. Graphs display individual

biological replicates as dots, with the mean and SEM. Statistical significance was determined

using Student's t test *p <0.05 **p <0.005 ***p <0.0005.

Figure 5-figure supplement 1. PABPC is required for the connection between cytoplasmic

mRNA decay and RNAPII promoter occupancy. (A,-C) HEK293T cells were treated with

siRNAs targeting CHD3 (A), MSI1 (B), or TRIM32 (C), or non-targeting scramble siRNAs, then

transfected with either empty vector or muSOX and subjected to chromatin immunoprecipitation

(ChIP) using antibodies to RNAPII or IgG. Purified chromatin was quantified by qPCR. Western

blots showing the levels of CHD3 and MSI1 after siRNA depletion, along with a GAPDH or

histone H3 loading control are shown below. Graphs display individual biological replicates as

dots, with the mean and SEM. Statistical significance was determined using Student's t test *p

<0.05 **p <0.005 ***p <0.0005.

Figure 5-figure supplement 2. Effects of depleting PABPC, LARP4, CHD3, MSI1 and

TRIM32 on RNAPII promoter occupancy. (A-E) HEK293T cells were treated with siRNAs

targeting PABPC1 and PABPC4 (A), LARP4 (B), CHD3 (C), MSI1 (D), or TRIM32 (E), and

subjected to ChIP using antibodies to RNAPII or IgG. Purified chromatin was quantified by

qPCR. All graphs display individual biological replicates as dots, with the mean and SEM.

Statistical significance was determined using Student's t test *p <0.05 **p <0.005 ***p <0.0005.

Figure 6-figure supplement 1. RNAPII Rpb1 and TBP protein levels are unchanged in cells

expressing muSOX or FLAG-PABPC1. Western blot of cells expressing empty vector,

muSOX, or FLAG-PABPC1, with VINCULIN serving as a loading control.

Table S1. Nuclear proteome. Identified and quantified proteins in the nuclear fraction with

associated TMT abundance values.

Table S2. Cytoplasmic proteome. Identified and quantified proteins in the cytoplasmic fraction

with associated TMT abundance values.

Table S3. Proteins enriched in the nucleus in muSOX expressing cells. Proteins significantly

(p < 0.05) enriched in the nucleus in muSOX expressing cells compared to D219A expressing

cells that also show either no change or a decrease in cytoplasmic abundance. Listed from left to

right: Accession number, gene symbol, nuclear abundance ratio, muSOX compared to D219A

muSOX, subcellular localization, and gene ontology (GO) Molecular Function. The Database for

Annotation, Visualization, and Integrated Discovery (DAVID v6.8) was used for subcellular

localization and GO molecular function information.

Table S4. Proteins differentially expressed in Xrn1 knockout cells. Proteins that are

differentially expressed in Xrn1 knockout cells in the nucleus (left) and the cytoplasm (right).

Listed are the protein accession numbers and the corresponding gene symbol.

Table S5. Primer and custom siRNA sequences.

Figure 1

Figure 1-figure supplement 1

A.

B. Poly(U) binding proteins

MSI1

MSI2

● Nucleus
■ Cytoplasm

C. mRNA 3'UTR binding proteins

PUM2

IGF2BP2

ELAVL2

● Nucleus
■ Cytoplasm

Figure 2

A. Xrn1 requirement for shuttling

33% independent

67% dependent

B.

Nuclear abundance / vector

C.

WT

Xrn1 KO

D.

DAPI   α PABPC1   GFP   Overlay

IFA Quantification

GFP-muSOX
WT
Xrn1 KO

Figure 2-figure supplement 1

Figure 3

A. Differential nuclear and cytoplasmic abundance in Xrn1 KO

C. Expression of Xrn1 interactors

Figure 3-figure supplement 1

A.   OST complex

B.   Gene Expression

Figure 4

A. STRING interaction networks among shuttling proteins

B.

C.

Figure 5

A. PABPC knockdown RNAPII ChIP

B. LARP4 knockdown RNAPII ChIP

C.

D. RNAPII ChIP

E. Xrn1 KO RNAPII ChIP

Figure 5-supplement 1

A. CHD3 knockdown RNAPII ChIP

B. MSI1 knockdown RNAPII ChIP

C. TRIM32 knockdown RNAPII ChIP

Figure 5-figure supplement 2

A. PABPC knockdown RNAPII ChIP

B. LARP4 knockdown RNAPII ChIP

C. CHD3 knockdown RNAPII ChIP

D. MSI1 knockdown RNAPII ChIP

E. TRIM32 knockdown RNAPII ChIP

Figure 6

A. RNAPII ChIP

B. ChIP: Ser5P / RNAPII

C. ChIP: Ser2P / RNAPII

D. TBP ChIP

E. RNAPIII ChIP

F.

Figure 6-figure supplement 1