1   *Lactobacillus acidophilus* **disrupts collaborative multispecies bile acid metabolism**
2

3   Sydney Dautel[1,2], Nymul Khan[1], Kristopher R. Brandvold[1], Colin J. Brislawn[1], Janine
4   Hutchison[3], Karl K. Weitz[1], Heino M. Heyman[1], Hyun-Seob Song[1], Zehra Esra Ilhan[4], Eric A.
5   Hill[1], Joshua R. Hansen[1], Xueyun Zheng[1], Erin S. Baker[1], John R. Cort[1], Young-Mo Kim[1],
6   Nancy G. Isern[5], John K. DiBaise[6], Rosa Krajmalnik-Brown[4], Janet K. Jansson[1], Aaron T.
7   Wright[1,7], Thomas O. Metz[1], Hans C. Bernstein[1,7,8*]
8
Affiliations:
[1]Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA, USA;
[2]School of Medicine, University of Washington, Seattle, WA, USA; [3]Chemical and Biological
Signature Science, Pacific Northwest National Laboratory, Richland, WA, USA; [4]Biodesign
Swette Center for Environmental Biotechnology and Biodesign Center for Fundamental and
Applied Microbiomics, Arizona State University, Tempe, AZ, USA; [5]Environmental Molecular
Sciences Laboratory, Pacific Northwest National Laboratory, Richland, WA, USA; [6]Division of
Gastroenterology, Mayo Clinic, Scottsdale, AZ, USA; [7]The Gene and Linda Voiland School of
Chemical Engineering and Bioengineering, Washington State University, Pullman, WA, USA;
[8]Norwegian College of Fishery Science, UiT - The Arctic University of Norway, 9019, Tromsø,
Norway
20
21   ***Correspondence**: Hans C. Bernstein, 902 Battelle Boulevard, P.O. Box 999, MSIN J4-18,
22   Richland WA, 99352; Phone: 509-371-6989, Email: Hans.Bernstein@pnnl.gov; Email:
23   Hans.C.Bernstein@uit.no

24
Keywords: probiotic, bile acid, microbiome, consortia, multi-omics, chemical probes

ABSTRACT

Bile acids are metabolic links between hosts and their gut microbiomes, yet little is known about

the roles they play in microbe-to-microbe interactions. Here we present a study designed to

investigate the effect that a common probiotic, *Lactobacillus acidophilus*, has on microbial

interactions that lead to formation of secondary bile acids. A model microbial consortium was

built from three human gut isolates, *Clostridium scindens, Collinsella aerofaciens,* and *Blautia*

obeum, and cultured under different bile acid and probiotic treatments. A multi-omics platform

that included mass spectrometry-based metabolomics and activity-based proteomic probes was

used to produce two major results. The first, was that an uncommon secondary bile acid –

ursocholate – was produced by a multi-species chemical synthesis pathway. This result

highlights a new microbe-to-microbe interaction mediated by bile acids. The second finding was

that the probiotic strain, *L. acidophilus,* quenched the observed interactions and effectively

halted consortial synthesis of ursocholate. Little is known about the role that ursocholate plays in

human health and development. However, we did discover that a decrease in ursocholate

abundance corresponded with successful weight loss in patients after gastric bypass surgery

versus those who did not lose weight after surgery. Hence, this study uncovered basic knowledge

that may aid future designs of custom probiotic therapies to combat obesity.

INTRODUCTION

The human gastrointestinal (GI) tract is a complex ecosystem that functions in symbiosis with oral and intestinal microbiomes[1,2]. It has long been recognized that the composition of the gut microbiome has a significant effect on host digestion but more recent research has implicated the microbiome in human health and disease states that include cardiovascular disease risk[3], neurological function[4], and autoimmunity[5]. Rapid gains in knowledge of host-microbiome associations will undoubtedly lead to new practical applications[6]. Of these, the use of probiotics to modulate both the function and composition of gut microbiomes is especially promising[7,8]. Probiotic supplementation likely reduces the risk of developing antibiotic associated diarrhea[9] and necrotizing enterocolitis in infants[10]. However, the therapeutic opportunities for probiotics are advancing to more precisely target specific processes carried out by the gut microbiome to impart health benefits beyond enhanced digestion[11-13]. Probiotic therapies are being explored relieve symptoms of autism[14], depression[4], autoimmune diseases[15,16], and irritable bowel syndrome[17] among many other conditions with positive – albeit conflicting – results. The efficacy of probiotic treatments is variable. Differing results can obviously arise from inconsistent study design – e.g., probiotic strain, dose – trial size, but they are also indicative of a large scientific knowledge gap and incomplete understanding of the mechanisms by which probiotics impact the GI-tract microbial ecosystem.

There are many hypotheses about the modes of action by which the gut microbiome and probiotic microbes impact human health. One proposed model is through modulation of the host immune system[18], which has been concluded from studies that showed probiotic treatments affecting host immune function in humans facing pathogenic challenges[19-21] or with autoimmune disorder[15,16,22]. Another hypothesis is that probiotics increase microbial competition within the

intestinal ecosystem, thereby making it more difficult for pathogenic bacteria to survive[23,24].

Studies have also speculated that observed therapeutic action from probiotics results from their

effect on the intestinal physiology by modulating endothelial junctions[25] and the mucosal

lining[12,25] through a variety of proposed metabolic pathways. Another possible mode of action

for probiotics is through bile acids. It has long been known that bile acids are important linkers

between host and gut microbes. Intestinal bacteria produce secondary bile acids by

deconjugation, reduction, oxidation, and epimerization of their host's primary bile acids. Many

probiotics can alter bile acid pools in humans[26,27].

The primary bile acids (cholic acid and chenodeoxycholic acid) are synthesized in human

hepatocytes and conjugated to the amino acids glycine and taurine to increase solubility[27,28].

They are then released into the duodenum and moved through the small intestine to assist

emulsification of dietary lipophilic substances[29]. Approximately 95% of the bile acids are then

passively reabsorbed in transit through the small intestine, resulting in approximately 5% (400-

800 mg) passed on to the colon[27]. In the colon, these bile acids are rapidly deconjugated by the

microbiome and reduced, oxidized and epimerized to a variety of secondary bile acids[27]. These

secondary bile acids are known to have diverse effects on human health ranging from direct

cytotoxicity[30], to altered probability of cancer[31], to hormonal function as cell messengers[29,32-34].

More recently, this list of known host-related effects has grown to include modulating the

composition and function of the gut microbiome[35,36] – e.g., by disassembling lipid membranes.

The intestinal microbiome represents a major modifier of the human bile acid pool. This

is evinced by the fact that bile in the gall bladder is comprised of 70% primary bile acids but

only 4% primary bile acids in the feces[27]. Not only does the microbiome determine bile acid

composition, but bile acids also direct microbial communities[37]. For example, studies have

shown that high fat diets and diets high in resistant starch[38] have an effect on both the bile acid

pool[39] and the microbiome[40]. It has also been shown in rats that oral administration of certain

bile acids can shift the microbiome composition[35,41]. Collectively, these studies suggest that there

is a complex interplay between microbial species and bile acids that is not fully understood.

The relationships between the gut microbiome, probiotics and the bile acid pool are

particularly relevant due to the current epidemic of obesity and the comorbidities associated with

high adiposity (high cholesterol, high blood pressure, diabetes)[42]. Due to high rates of obesity,

bariatric surgery has become a more common procedure with the number and types of surgeries

increasing with time[43,44]. It is now well established that bariatric surgery results in reduced

weight and a reduction in many comorbidities of obesity[45]. Several different advantages (weight

loss, reduction of comorbidity) and disadvantages (surgical complications, malnutrition, weight

regain, re-surgery, infection, etc.)[45,46] have been have been identified for different bariatric

procedures. Yet, there remains an incomplete understanding of the exact mechanisms of many of

these outcomes, making it difficult to predict which patients will benefit most from these

procedures. Elucidation of the root causes will require consideration of the impacts that the bile

acid pool, microbiome composition and probiotic administration can have[47,48]. Deeper

mechanistic insight could not only result in increased surgery success, potentially by pre-emptive

modulation of the microbiome by bile acid pool[49,50], but could also result in less surgeries

necessitated if some of the positive results can indeed be realized via targeted use of probiotics.

Here we present a study that was designed to investigate the community dynamics of

microbes commonly found in the gut and the impact that both the addition of bile acids and a

probiotic has on interspecies interactions. An *in vitro* model was built as a three-species bacterial

consortium, *Clostridium scindens, Collinsella aerofaciens,* and *Blautia obeum,* each of which

occurs naturally in the human gut. This consortium was then treated with the probiotic strain

Lactobacillus acidophilus to make an altered four-member consortium. Multi-omics assays were

used to elucidate the interspecies microbial interactions with and without bile acid treatments

(cholic acid and deoxycholic acid). We found that a secondary bile acid, ursocholic acid (7-

epicholic acid), was produced from cholic acid through a multispecies chemical synthesis route

mediated by an interaction between *B. obeum* and *C. aerofaciens*. This process was quenched by

the addition of *L. acidophilus*, which disrupted the coordination between *B. obeum* and *C.*

aerofaciens and shut down ursocholic acid production. These results were then contextualized by

performing targeted metabolite measurements in fecal samples from a human clinical study[44] that

investigated secondary bile acid abundances as outcomes of gastric bypass surgery. The

abundance of ursocholic acid corresponded with successful post-operative weight loss,

highlighting that it may be important to investigate the implications of both patient- and microbe-

derived metabolites to help gain a predictive understanding of a patient's response to bariatric

surgery. New knowledge in this area will yield opportunities to design custom probiotic

therapies. More broadly, this study supports an emerging theme in microbiome sciences that

microbial interactions are context dependent and the presence or absence of select species and/or

metabolites can have a strong effect on the overall function.

RESULTS

A model microbial consortium responds to probiotic and bile acid treatments. We

constructed an *in vitro* microbial consortium from human gut bacterial isolates. A 3-member

consortium (3MC) composed of *Clostridium scindens* ATCC 35704, *Collinsella aerofaciens*

ATCC 25986, and *Blautia obeum* ATCC 29174 was compared to a 4-member consortium (3MC

+ L) that included an added a probiotic strain, *Lactobacillus acidophilus* ATCC 4356. The

species were chosen for this model based on their capacity to perform one-of-three unique

microbial transformations on human bile acids (Table S1): 7α-dehydroxylation, hydroxysteroid

dehydrogenatation, or bile salt hydrolsis[27,37,51,52].

Anaerobic culturing treatments included supplementation with cholic acid and/or

deoxycholic acid and were compared to axenic controls. We found that each member of the

consortia grew together under anaerobic culturing conditions but that each consortium (3MC

versus 3MC + L) had very different growth and extracellular metabolite profiles (Fig. 1). Despite

the anticipated anti-microbial effect of bile acids[26,53,54], all cultures in media containing added

cholic acid and deoxycholic acid (0.1 mM) showed faster specific growth rates compared to their

corresponding treatments without bile acids (Fig. 1a and b). The axenic controls confirmed that

L. acidophilus had the fastest specific growth rate, followed by *B. obeum*, *C. aerofaciens* and *C.*

scindens. The probiotic, *L. acidophilus*, was the least affected by the addition of the bile acids to

the growth media, with only a 17% increase in the specific growth rate as compared to 170%,

76% and 167% increases for *C. scindens*, *C. aerofaciens* and *B. obeum,* respectively. An adonis

test was used to show that the species composition of each consortium – i.e., presence/absence of

L. acidophilus – was the strongest determiner of variance (R2 = 0.49, p < 0.001) in the

extracellular metabolome. This was in contrast to the effect of treatments that tested for changes

in the global metabolome based on bile acid or no bile acid inputs, which were not a statistically

significant source of variance (Fig. 1C). Hence, the probiotic *L. acidophilus* was a major

modifier of the extracellular chemical environment.

Figure 1. Probiotic influence on consortial growth and the extracellular chemical environment. (A) Growth of the 3-member consortium (3MC) in bile acid supplemented media as compared to (B) the 4-member consortium (3MC + L) that differed by the addition of probiotic, *L. acidophilus*. The y-axis of A and B represent means from triplicate measurements of absolute species abundance defined as the total optical density (OD_{630nm}) at each time point multiplied by each respective measurement of relative abundance obtained from qPCR; error bars represent ± 1 standard deviation. (C) A principle component analysis on the extracellular metabolome ordinated by the Euclidean distance between the metabolic profiles from each treatment. The colored ellipses represent 95% confidence limits assuming a multi-variate t-distribution.

L. acidophilus quenches secondary bile acid production. Ursocholic acid was produced in the

3MC (Fig. 2A), but not by any one species grown under axenic conditions. Hence, ursocholic

acid was produced by a multispecies synthesis route that required at least two species from our

model consortium. Addition of the probiotic, *L. acidophilus* quenched the production of

ursocholic acid to negligible levels as compared to those measured in the 3MC (Fig. 2b). In

addition to ursocholic acid production, *L. acidophilus* also attenuated growth of *B. obeum* and *C.*

aerofaciens as observed in the 3MC + L species-specific growth dynamics, which were in stark

contrast to the 3MC (Fig.2c). Within the 3MC + L, *L. acidophilus* became a dominant member

of the community, but in the absence of the probiotic, the 3MC was dominated by *B. obeum* with

C. scindens showing the lowest relative abundance in both consortia.

The dynamic profiles of bile acids and bacterial species were correlated for each

treatment group, 3MC and 3MC + L (Fig. 2d), respectively. Pearson's correlations were

determined between species and for species to bile acids but not between bile acids. The results

show that ursocholic acid shared strong positive correlations (Pearson's coefficient; $r > 0.85$)

with the abundances of *C. aerofaciens* and *B. obeum* in the 3MC. As expected, cholic acid

showed strong negative correlations with *C. aerofaciens* and *B. obeum* within the 3MC,

establishing that it was the most likely substrate for ursocholic acid synthesis. Cholic acid, was

not correlated with *C. scindens*, implying that this species may not have been involved in

production of secondary bile acids directly from cholic acid. The lack of *C. scindens'*

participation in secondary bile acid synthesis in the 3MC was also evinced by its strong negative

correlation with deoxycholic acid, a known 7α-dehydroxylation product of *C. scindens* that

utilizes cholic acid as the substrate[27].

L. acidophilus membership

changed the correlations between species

and bile acids. Most notably, ursocholic

acid was not correlated with any species

in the 3MC + L under the probiotic

treatment. *L. acidophilus* did not

correlate with any of the bile acids and

shared strong negative correlations with

C. aerofaciens and *B. obeum*, indicating

that competition and/or antagonism are

the likely mechanisms by which the

multi-species ursocholic acid synthesis

was quenched.

Multispecies synthesis of ursocholic

acid. Based on the metabolomics results,

we hypothesized a multi-species

cooperative synthesis of ursocholic acid

that involved both *B.*

obeum and *C.*

aerofaciens. The initial

inference was derived by

comparing the metabolite

Figure 2. Secondary bile acid – ursocholic acid – was produced by consortia but not axenic cultures; *L. acidophilus* disrupted the consortial synthesis. (A) Comparative abundances of bile acids measured after 24 h incubations (3MC and 3MC + L) and axenic controls. Time course measurements of (B) bile acid abundances and (C) species abundance as shown by the inverse qPCR cycle thresholds (CT). Comparisons are shown between the 3MC and 3MC + L (with *L. acidophilus*) treated with the 0.1 mM bile acid mixture. Each data point shown in panels A-C represent the mean from a minimum of three biological replicates ± 1 standard deviation. (D) Pearson's correlations between bacterial species and bile acid abundances; thicker lines correspond to greater correlation coefficients (cut-off below 0.85); red and grey colors correspond to positive and negative correlations, respectively.

profiles between the axenic and consortial treatments (Fig. 2a). The next piece of evidence was

obtained from correlations between species and bile acids in the context of *a priori* knowledge of

the metabolic reactions that were the basis for choosing each species in the model consortium

(Fig. 2d). A possible mechanism for this could start with *B. obeum* conversion of cholic acid into

a transient 7-keto intermediate, such as 7-oxodeoxycholic acid. A ketone intermediate was not

identified by our bile-acid-targeting LC-MS metabolomics approach but that does not exclude

the possibility of its existence. The next step could be achieved by the known genome encoded

functions of *C. aerofaciens*, which contains *hdhB* (GenBank accession ZP_01773061)[55]. This

gene encodes for a 7β-hydroxysteroid dehydrogenase (7β-HSDH), known to catalyze a reaction

that takes a 7-oxodeoxycholic acid to ursocholic acid[56]. However, the *B. obeum* ATCC 29174

genome does not contain an oxidoreductase that is clearly annotated to catalyze our hypothesized

reaction in this first step. *B. obeum* does contain a *baiA* gene (RUMOBE_03494) that encodes a

putative 7α-dehydroxylase, but genes encoding for a 7α-HSDH have yet to be identified. Yet our

experiments clearly showed synthesis of ursocholic acid and we confirmed a required

participation of *B. obeum*; hence, we concluded that a non-specific or previously unidentified

oxidoreductase catalyzed this first step.

To test our hypothesis that *B. obeum* expresses enzymes that react with cholic acid (other

than the known BaiA protien), we synthesized and employed a custom cholic acid photo affinity

probe (CAP; cholic acid probe). This probe is a cholic acid derivative designed to bind and

enrich proteins that can then be identified by MS-based proteomics (Fig. 3a). The results were a

suite of proteins from each species in the consortium that were significantly enriched by the CAP

($>10^2$ fold-change; p-value < 0.001) (Table S2). Surprisingly, the *B. obeum* BaiA protein

(*RUMOBE_03694*) was not enriched by the CAP, indicating the possibility of an incorrect

annotation from homology of UniProt-KB A5ZWVO or lack of expression in these consortial

conditions. We identified a suite of *B. obeum* proteins that were significantly enriched by the

CAP and specific to the treatments corresponding to ursocholic acid synthesis and the *B. obeum*

axenic treatment. Of these, we focused on a pair of oxidoreductases as candidates for the

hypothesized two-step, two-species reaction that lead to ursocholic acid in our consortium. While

they did not share strong homology with known 7α-HSDH proteins, they were annotated as

enzymes that may catalyze the hypothesized alcohol-ketone inter-conversions (Fig. 3b). These

CAP-binding proteins were annotated based on homology as an alcohol dehydrogenase

(A5ZM66; *RUMOBE_00083*) and an alcohol-aldehyde dehydrogenase (A5ZNA3;

RUMOBE_00470).

246
Figure 3. **The multispecies synthesis hypothesis and cholic acid binding oxidoreductases.** (A) The cholic acid photoaffinity
probe (CAP) was synthesized and used to enrich proteins for mass spectrometry-based proteomics. (B) Of the proteins enriched
by CAP (> 100 fold-change and p < 0.001), we identified two oxidoreductases annotated alcohol or alcohol-ketone
dehydrogenases (ADH and ALDH, respectively). These proteins are B. obeum candidates for the hypothesized two-step, two-
species mechanism (C) leading to the synthesis of ursocholic acid. The abbreviations used to describe experimental treatments
are as follows: Bo (*B. obeum*); Ca (*C. aerofaciens*); and Cs (*C. scindens*).

Implications toward obesity and

bariatric surgery. The proposed

mechanism for physiological effect for

Roux-en-Y gastric bypass surgery

(RYGB) are BRAVE "Bile flow

alteration, Reduction of gastric size

Anatomical gut re-arrangement, Vagal

manipulation, Enteric gut hormone

modulation[57]. Hence, the GI tract of

patients that have undergone bariatric

surgery is a potential model system to

study bile acid metabolism/alterations

Figure 4. Normalized abundance of ursocholic acid in feces of patients who had undergone gastric bypass surgery – Roux-en-Y gastric bypass (RYGB) – as compared to normal weight and obese controls. Unpaired two-tailed t-tests were used to infer a statistical difference between the means of the obese controls – a group that contained pre-operative patients – and those that experience successful RYGB ($p < 0.005$). Different ursocholic acid abundances were also observed – albeit statistically less conclusive – between successful and unsuccessful RYGB patients ($p < 0.076$).

given that bile acid profiles change by increasing abundance of secondary bile acids[49]. To

investigate the clinical relevance of ursocholic acid, we leveraged access to a cohort of patients

that had undergone gastric bypass surgery. Targeted measurements of ursocholic acid were

performed and compared in fecal samples collected from 24 patients that underwent RYGB

surgery: 10 patients with normal weight and 14 morbidly obese controls, which included those

scheduled for surgery[44]. We found that the abundance of ursocholic acid corresponded with

obesity and the success of gastric bypass surgical procedures (Fig. 4). Success was defined when

patients exhibited at least 50% excess weight loss and less than 20% regain. Ursocholic acid

levels were significantly higher in the morbidly obese controls (pre-surgery) compared to

patients that had experienced successful gastric bypass surgery. There was no statistically

significant difference between the fecal abundance of ursocholic acid in the obese controls and

the unsuccessful surgical patients.

<u>DISCUSSION</u>

The human gut microbiome is integral to human health and development[58,59]. However, the

function and composition of the human GI-tract ecosystem is complex[60], which often makes

detailed studies of individual processes difficult. The use of model systems is a valuable

approach to dissect complex biological functions. In particular, simplified model consortia

gaining interest in microbiome research because they enable experimentalists to manage

complexity by controlling multiple system components under defined treatments. The utility of

simplified consortia, or bench-top microbiomes, has been demonstrated for a variety of human-

associated communities[61,62] and other complex microbial ecosystems related to plants[63],

sediment/biofilms[64-66] and marine habitats[67,68]. Here, we developed a simple model microbial

consortium that was specifically designed to investigate how the addition of a common probiotic

(*L. acidophilus*) effects microbial interactions mediated through bile acid transformations. This

study was not designed to directly inform microbial physiologies that should occur in the human

GI-tract *in situ*. Rather, it was designed and successfully implemented for controlling the

membership of microbial species and presence/absence of bile acids that are associated with

human digestive systems. Our model bile acid consortium helped produce two major results. The

first was that ursocholic acid was synthesized by the coordinated metabolism of a consortium

and not by any single species included within this study. The second important finding was that

probiotic, *L. acidophilus* quenched the observed multi-species interactions that resulted in

secondary bile acid synthesis.

Ursocholic acid is the 7 beta-hydroxyepimer of cholic acid. It is rarely mentioned in the

scientific literature and has been termed an "unusual secondary bile acid" as compared to more

commonly studied metabolites such as deoxycholic and ursodeoxycholic acid[69]. Previous studies

have investigated ursocholic acid as a potential therapeutic to modulate the host's synthesis of

primary bile acids[70], or to improve the liver function of patients with primary biliary cirrhosis[71]

and reduction of bile cholesterol saturation[72]. In these previous studies, ursocholic acid was

typically tested along with ursodeoxycholic acid and found to be notably less effective as a

treatment for biliary cirrhosis[71]. Despite these therapeutic studies, little is known about the role

that ursocholic acid plays in modulating human-microbe and/or microbe-microbe interactions.

The data derived from our clinical study showed that ursocholic acid is in fact present

and abundant within the human GI-tract and its relative abundances change when drastic changes

to microbiome occur (pre- and post- bariatric surgery)[44]. Our limited understanding of the role

that ursocholic acid plays in human health and specifically the host-microbiome interactions that

lead to its production represent a major knowledge gap. This is punctuated by the fact that our

current study – and a previous study[73] – have observed microbial synthesis of ursocholic acid

and/or 7-oxodeoxycholic acid but not conclusively identified the genes and 7α-HSDH proteins

responsible.

Cataloguing the bacterial genes from the "sterolbiome" is an active area of research[74] that

can yield new biological insight and help to improve human health by understanding how

microbes modify chemical environments within the human GI tract. In this study, we

hypothesized a multispecies chemical synthesis route in which *B. obeum* converts cholic acid

into an intermediate ketone via a 7α-HSDH-like reaction which is then proceeded by the known

7β-HSDH reaction catalyzed by *C. aerofaciens*. Intraspecies 7-epimerization has been

demonstrated in *Clostridium limosum*[75] and *Clostridium absonum*[76], which express both the

required enzymes, 7α-HSDH and 7β-HSDH. However, genes encoding for 7α-HSDH, have yet

to be identified in *B. obeum* and other bacteria such as an *Eggerthella* sp. known to express this

protien[73]. Yet, we conclusively found that a collaborative reaction between *B. obeum* and *C.*

aerofaciens does occur, which highlights an increase in our understanding of bile acid

metabolism of bacteria.

We hypothesized a 7-keto intermediate that was transferred between *B. obeum* and *C.*

aerofaciens in the observed multispecies chemical synthesis route. We did not identify an

intermediate such as 7-oxodeoxycholic acid in the supernatant of the samples and therefore

cannot categorically confirm its existence. However, 7α-HSDH mediated synthesis of 7-

oxodeoxycholic acid has been previously observed in similar multi-step bile acid transformation

processes[56]. It is possible that *C. aerofaciens* has a high affinity uptake mechanism for the

hypothesized 7-keto intermediate such that extracellular concentrations were below the detection

limits of our LC- and GC-MS metabolite identification methods. Another point of uncertainty is

whether the CAP-enriched alcohol dehydrogenase (A5ZM66; *RUMOBE_00083*) and alcohol-

aldehyde dehydrogenase (A5ZNA3; *RUMOBE_00470*) were responsible for the hypothesized

reactions. Certainly, other proteins were enriched by CAP (Table S2), yet these were the only *B.*

obeum proteins annotated with an enzymatic function capable of alcohol-aldehyde

interconversion. We cannot rule out the possibility that enriched proteins of unknown function

participated in the observed reaction. However, there is some precedent for associating

secondary bile acid synthesis genes with alcohol dehydrogenases. BaiA proteins from *C.*

scindens, encoding for 3α-HSDH proteins, have previously been shown to align well to short

chain alcohol dehydrogenases in *Eubacterium* sp. Strain VPI 12708 and to alcohol/polyol

dehydrogenase genes[77,78]. We chose to use *C. scindens* in the 3MC because it has genes that

encode for Bai proteins and express HSDH proteins. In fact, the *C. scindens* reference proteome

(VPI 12708) does contain a putative 7α-HSDH (UniProtKB – Q03906); however, *C. scindens*

abundances and correlation-based inferences from this study did not provide evidence of *C.*

scindens' participation in the transformation of cholic acid to secondary acids. Our conclusion

was that *C. scindens* played a minor role in the system and was largely outcompeted by other

members.

Towards precision probiotics to complement bariatric surgery outcomes. The effect of

probiotics on the human gut microbiome and the bile acid pools are particularly relevant due to

the current epidemic of obesity and the comorbidities associated with high adiposity (high

cholesterol, high blood pressure, diabetes)[42]. Hence, bariatric surgery is becoming more common

as a treatment strategy. Yet knowledge gaps still exist with respect to how metabolites and

microbes could play a role in successful outcomes. Here, we show that the abundance of

ursocholic acid corresponds with the efficacy of gastric bypass surgery. This new knowledge

about the role of specific probiotic strains and/or their metabolic products, are therefore leading

towards promising novel treatments for patients undergoing bariatric surgery. It has already been

shown that post-operative administration of *Lactobacillus* sp. improves weight loss and vitamin

B absorption in RYGB patients[48]. It is possible that the cessation of ursocholic acid production or

increased abundances of *Lactobacillus* sp. could result in better control over weight loss. We also

note that there is some precedent derived from mouse models for the idea that probiotics or

introduction of non-adapted microbial taxa can modulate a hosts' microbiome[7] and microbe-

associated bile acid pool[79].

Conclusion. This current study was a fundamental investigation of microbial interactions

and the role that a probiotic bacterium plays in modulating the synthesis of secondary bile acids.

It was not intended to inform clinical practice. However, the results and conclusions presented

establish an important idea related to broader microbiome sciences; microbial interactions are

context dependent[64,65] and the presence or absence of select species and/or metabolites can have

a strong effect on the overall community-level function.

METHODS

Bacterial strains and cultivation. *Clostridium scindens* ATCC 35704, *Colllineslla aerofaciens*

ATCC 25986, *Blautia obeum* ATCC 29174 (formerly *Ruminococcus obeum*) and *Lactobacillus*

acidophilus ATCC 4356 were grown under axenic conditions and in consortia on Lactobacilli

MRS Broth (BD Difco, Houston, TX, USA). The primary treatment the addition of bile salts,

equivalent mixtures of cholic and deoxycholic acid (Sigma-Aldrich 48305, St. Louis, MO,

USA), supplemented to 0.1 mM. The secondary treatment was the presence and absence of

probiotic, *Lactobacillus acidophilus* ATCC 4356 rendering either a 3-member (*L. acidophilus*

negative) or 4-member (*L. acidophilus* positive) consortia. Anaerobic growth conditions were

prepared by boiling the media and subsequently sparging with an 80% N_2, 10% H_2, 10% CO_2 gas

mixture and transferring to 30 ml sealed Balch tubes under the same gas headspace prior to

autoclaving. Each culture was inoculated to a starting $OD_{630nm} = 0.077 \pm 0.033$ by each axenic

cell suspension resulting in 1 serial passage of log phase cells; consortia were inoculated with an

equivalent volume ratio mL. The optical density (OD_{630nm}) was measured over a 24 h period

using a Spectronic 20D+ spectrophotometer (ThermoSpectronic, Madison, WI, USA); each time

point was sampled in triplicate via destructive sampling.

PCR quantification of gene target. Bacterial DNA was extracted from consortia and axenic

culture using the MoBio (Carlsbad, CA, USA) PowerSoil DNA Isolation Kit following the

manufacturer's protocol. A total volume of 5 µL of undiluted consortia DNA or standard curve

DNA was analyzed in triplicate with an Applied Biosystems 7500 fast instrument (Foster City,

CA). Samples were analyzed in triplicate with the primers shown in Table S3 targeting *rpoB*.

PCR reactions were run using the FAST cycling conditions: initial denaturation was done for 20

seconds at 95 °C followed by 40 cycles of denaturation (95 °C for 3 seconds), annealing (60 °C

for 30 seconds). The output from the real-time PCR assays were C_T values that represent the

PCR cycle at which the amplification crosses a given threshold (0.1). All C_T values in Table S4;

data are plotted and analyzed as inverse C_T representing the relevant abundance[66,80] assuming

equal between species in the consortia.

Bile acid identification and quantification. Stock solutions of the cholic and deoxycholic acids

(Steraloids Inc., Newport, RI, USA) were made in methanol (1 mg mL^{-1}) and were then pooled

together and diluted in a series in 0.1% formic acid to generate a 7 pt of the calibration curve

(0.0062, 0.025, 0.050, 0.1, 0.5 and 1 µg mL^{-1}). An internal standard (23-nor-5β-cholanic acid-3α,

12α-diol) was added to the filtered media (0.2 µm) collected from each microbial growth sample.

Cold methanol (-20˚C) was added at a ratio of 1:4 (filtrate:MeOH). The samples were mixed via

vortexing, chilled at -20˚C for 30 minutes and separated via centrifugation (1725 rpm, 10

minutes). The supernatant was removed and dried and then re-suspended in 0.1% formic acid in

deionized water solution. These samples were analyzed on a Waters nano-Acquity UPLC system

(Milford, MA, USA) configured for direct 5 µL sample injections onto an in-house packed fused

silica column (360 µm o.d. x 150µm i.d. x 30 cm long; Polymicro Technologies Inc., Phoenix,

AZ, USA) containing Waters HSS T3 media (1.8 µm particle size). A flow of 600 nL min^{-1} was

maintained using mobile phases consisting of (A) 0.1% formic acid in water and (B) 0.12 %

formic acid and 5 mM ammonium acetate in methanol with the following gradient profile (min,

%B): 0, 1; 5, 1; 10, 65; 59, 99; 60, 1. Total run time including column re-equilibration was 75

min. Mass spectrometry (MS) analysis was performed using an Agilent model 6490 triple

quadrupole mass spectrometer (Agilent Technologies, Santa Clara, CA, USA) outfitted with a

custom nano-electrospray ionization interface built using 150 um o.d. x 20 um i.d. chemically

etched fused silica [81]. The hexabore ion transfer tube temperature and spray voltage were held at

200˚C and -4.0 kV, respectively. Data were acquired in negative ion mode for 75 min from

sample injection using a dwell time of 200 µs, fragmentation of 380 volts, and collision energy

of 10 volts. Selected reaction monitoring (SRM) transitions were acquired as shown in

supplementary Table S5. Ursocholic acid was identified as an unknown in the initial LC-MS

trials. After fractionation and purification, we isolated the unknown and verified that it was

ursocholic acid via NMR and ion mobility mass spectrometry analyses. Authentic ursocholic

acid was purchased from Toronto Research Chemicals (N. York, Ontario, Canada).

Comprehensive details of these procedures are provided in the supplementary materials.

Untargeted Metabolomics. The spent media was dried, chemically derivated and analyzed by

GC-MS as previously reported[82]. GC-MS raw data files collected by GC-MS were processed

using the Metabolite Detector software, version 2.5 beta[83]. Agilent .D files were converted to

netCDF format using Agilent Chemstation (Agilent, Santa Clara, CA, USA) and then converted

to binary files using Metabolite Detector. Samples were aligned chromatographically across all

analyses after deconvolution. Metabolites were identified by matching experimental spectra to a

Pacific Northwest National Laboratory (PNNL) augmented version of FiehnLib[84]. This library

has spectra and validated retention indices for over 850 metabolites. In order to minimize errors

in deconvolution and identification, all metabolite identifications were manually validated after

automated data-processing.

Bile acid photoaffinity probes and proteomics. Custom photoaffinity probes were synthesized

as derivatives of cholic and deoxycholic acid for this study as described in detail within the

supplementary materials. Bacterial lysate samples were normalized to 500 µL 1.8 mg/mL

proteome in PBS buffer. Cholic acid photoaffinity probe (CAP) or an equal volume of DMSO

control was incubated with proteome for 60 min at 37 °C. Final DMSO concentration was 1%.

Samples were exposed to UV light (wavelength: 365nm; 115V, 15W) using a Fisher UVP95

lamp (Fisher Scientific, Hampton, NH, USA) for 7 minutes on ice. Subsequent to UV irradiation,

the samples were subjected to click chemistry, with final concentrations of reagents being:

biotin-azide (60 µM) in DMSO, sodium ascorbate (10 mM), THPTA (4 mM), and $CuSO_4$ (8

mM). Each reagent was added individually in that sequence, vortexed, centrifuged, and

incubated at room temperature in the dark for 90 min. 800 µL of pre-chilled MeOH was then

added to each sample and incubated at -80 °C freezer for 30 min to induce protein precipitation.

Samples were centrifuged at 14,000 x g at 4°C for 5 min. The supernatant was discarded, and the

pellet was allowed to air-dry for 5 min. Samples were reconstituted and sonicated in 520 µl SDS

(1.2%) in PBS; followed by incubation at 95 °C for 2 min. Samples were centrifuged at 14,000 x

g for 4 min at room temperature. Protein concentrations were determined via BCA assay and

samples were normalized to a volume of 500 µL at 1.2 mg/mL. Trypsin digestion was performed

on protein bound to 100 µL Streptavidin-agarose beads. Peptides were reconstituted by adding

40 µl of 25 mM NH_4HCO_3 and heating the samples at 37°C for 5 min. Samples were transferred

to ultracentrifuge tubes and were centrifuged at 100,000 x g to remove debris. 25 µL was added

to glass vials for storage at -20 °C until analysis. All proteomics samples prepared for LC-MS

were analyzed using a Velos Orbitrap MS as previously described [85,86].

Clinical data and experimentation. Fecal samples were collected from patients at the Mayo

Clinic, Scottsdale, AZ, USA. The metabolomics assays were performed at the Pacific Northwest

National Laboratory by the methods described above. The experimental design has been

previously described[44] and was approved by The Institutional Review Boards of Mayo Clinic

and Arizona State University (IRB# 10-008725).

Statistics. Analysis and graphing was performed in R[87] making use of the 'vegan'[88] and

'igraph'[89] packages, along with many packages in the Tidyverse[90]. The adonis test (permutation

MANOVA) was used to partition a matrix of Euclidean distances between global metabolite

samples based on bile acid and probiotic treatments. An unpaired two-tailed Student's t-test was

used to compare fecal ursocholic acid measurements between treatment groups. Linear models

were fit to the log of the exponential growth phase of microbial consortia, and for each

regression, the goodness of fit and the probability of observing a similar slope if the true slope

coefficient was zero was reported. Global proteomics was used to identify proteins enriched by

the CAP probes, by selecting proteins with a fold-change increase of >100 and a t-test p-value of

< 0.001. The methods for all hypothesis testing and descriptive statistical procedures are

included in the R markdown supplied for this study.

Data Repositories and Reproducibility. The raw data sets for this study along with the R

scripts used for analysis and graphing are available from the Open Science Framework (OSF)

under the name "Bile Acids Consortia" at https://osf.io/5meyd/. The mass spectrometry

proteomics data have been deposited to ProteomeXchange with the dataset identifier

PXD008617.

ACKNOWLEDGMENT

This research was supported by the Microbiomes in Transition (M*in*T) Initiative, a Laboratory

Directed Research and Development (LDRD) Program of Pacific Northwest National Laboratory

(PNNL). Measurements of bile acids in samples from the human clinical trial were supported by

the PNNL LDRD Program via the Signature Discovery Initiative. Research reported in this

publication was supported by the National Institute of Diabetes and Digestive and Kidney

Diseases of the National Institutes of Health under Award Number R01DK090379. The content

is solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health. Metabolomics measurements and chemical characterization of

ursocholic acid were performed in the Environmental Molecular Sciences Laboratory (user

project 49356), a national scientific user facility sponsored by the Department of Energy Office

of Biological and Environmental Research, located at PNNL. PNNL is operated for the DOE by

Battelle under contract no. DE-AC05-76RLO-1830. Specific acknowledgments are given to

Heather Brewer, Athena Schepmoes, Rose Perry and Erika Zink for their technical contributions.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

REFERENCES CITED

Carballa, M., Regueiro, L. & Lema, J. M. Microbial management of anaerobic digestion: exploiting the microbiome-functionality nexus. *Curr Opin Biotech* **33**, 103-111 (2015).

Simon, G. L. & Gorbach, S. L. The human intestinal microflora. *Digest Dis Sci* **31**, 147-162 (1986).

Brown, J. M. & Hazen, S. L. Microbial modulation of cardiovascular disease. *Nat Rev Microbiol* (2018).

Foster, J. A. & Neufeld, K.-A. M. Gut–brain axis: how the microbiome influences anxiety and depression. *Trends Neurosci* **36**, 305-312 (2013).

Bach, J.-F. The hygiene hypothesis in autoimmunity: the role of pathogens and commensals. *Nat Rev Immunol* (2017).

Schmidt, T. S., Raes, J. & Bork, P. The human gut microbiome: from association to modulation. *Cell* **172**, 1198-1215 (2018).

Martin, F. P. J. *et al.* Probiotic modulation of symbiotic gut microbial–host metabolic interactions in a humanized microbiome mouse model. *Mol Sys Biol* **4**, 157 (2008).

Kang, D.-W. *et al.* Microbiota Transfer Therapy alters gut ecosystem and improves gastrointestinal and autism symptoms: an open-label study. *Microbiome* **5**, 10 (2017).

McFarland, L. V. Meta-analysis of probiotics for the prevention of antibiotic associated diarrhea and the treatment of Clostridium difficile disease. *Am J Gastroenterol* **101**, 812-822 (2006).

AlFaleh, K. & Anabrees, J. Probiotics for prevention of necrotizing enterocolitis in preterm infants. *Evid-Based Child Health: A Cochrane Rev J* **9**, 584-671 (2014).

Baquero, F. & Nombela, C. The microbiome as a human organ. *Clin Microbiol Inf* **18**, 2-4 (2012).

Bron, P. A., van Baarlen, P. & Kleerebezem, M. Emerging molecular insights into the interaction between probiotics and the host intestinal mucosa. *Nat Rev Microbiol* **10**, 66-78 (2012).

Hooper, L. V., Littman, D. R. & Macpherson, A. J. Interactions between the microbiota and the immune system. *Science* **336**, 1268-1273 (2012).

Gilbert, J. A., Krajmalnik-Brown, R., Porazinska, D. L., Weiss, S. J. & Knight, R. Toward effective probiotics for autism and other neurodevelopmental disorders. *Cell* **155**, 1446-1448 (2013).

Pandey, G., Pandey, A. K., Pandey, S. & Pandey, B. Microbiota in Immune Pathogenesis and the Prospects for Pre and Probiotic Dietetics in Psoriasis. *Biomed Res J*, 220.

Vaghef-Mehrabany, E. *et al.* Probiotic supplementation improves inflammatory status in patients with rheumatoid arthritis. *Nutrition* **30**, 430-435 (2014).

Zhang, Y. *et al.* Effects of probiotic type, dose and treatment duration on irritable bowel syndrome diagnosed by Rome III criteria: a meta-analysis. *BMC Gastroenterol* **16**, 62 (2016).

Rooks, M. G. & Garrett, W. S. Gut microbiota, metabolites and host immunity. *Nat Rev Immunol* **16**, 341-352 (2016).

Bäumler, A. J. & Sperandio, V. Interactions between the microbiota and pathogenic bacteria in the gut. *Nature* **535**, 85-93 (2016).

Haywood, B. A. *et al.* Probiotic supplementation reduces the duration and incidence of infections but not severity in elite rugby union players. *J Sci Med Sport* **17**, 356-360 (2014).

21 King, S., Glanville, J., Sanders, M. E., Fitzgerald, A. & Varley, D. Effectiveness of probiotics on the
duration of illness in healthy children and adults who develop common acute respiratory
infectious conditions: a systematic review and meta-analysis. *Brit J Nutr* **112**, 41-54 (2014).
22 Lomax, A. & Calder, P. Probiotics, immune function, infection and inflammation: a review of the
evidence from studies conducted in humans. *Curr Pharm Design* **15**, 1428-1518 (2009).
23 Fooks, L. & Gibson, G. Probiotics as modulators of the gut flora. *Brit J Nutr* **88**, s39-s49 (2002).
24 Hemarajata, P. & Versalovic, J. Effects of probiotics on gut microbiota: mechanisms of intestinal
immunomodulation and neuromodulation. *Ther Adv Gastroenter*, 1756283X12459294 (2012).
25 Ohland, C. L. & MacNaughton, W. K. Probiotic bacteria and intestinal epithelial barrier function.
Am J Physiol-Gastr L **298**, G807-G819 (2010).
26 Begley, M., Gahan, C. G. & Hill, C. The interaction between bacteria and bile. *FEMS Microbiol Rev*
29, 625-651 (2005).
27 Ridlon, J. M., Kang, D.-J. & Hylemon, P. B. Bile salt biotransformations by human intestinal
bacteria. *J Lipid Res* **47**, 241-259 (2006).
28 Monte, M. J., Marin, J., Antelo, A. & Vazquez-Tato, J. Bile acids: chemistry, physiology, and
pathophysiology. *World J Gastroenterol* **15**, 804-816 (2009).
29 Kuipers, F., Bloks, V. W. & Groen, A. K. Beyond intestinal soap [mdash] bile acids in metabolic
control. *Nat Rev Endocrinol* **10**, 488-498 (2014).
30 Attili, A., Angelico, M., Cantafora, A., Alvaro, D. & Capocaccia, L. Bile acid-induced liver toxicity:
relation to the hydrophobic-hydrophilic balance of bile acids. *Med Hypotheses* **19**, 57-69 (1986).
31 Louis, P., Hold, G. L. & Flint, H. J. The gut microbiota, bacterial metabolites and colorectal cancer.
Nat Rev Microbiol **12**, 661-672 (2014).
32 Degirolamo, C., Rainaldi, S., Bovenga, F., Murzilli, S. & Moschetta, A. Microbiota modification
with probiotics induces hepatic bile acid synthesis via downregulation of the Fxr-Fgf15 axis in
mice. *Cell Rep* **7**, 12-18 (2014).
33 Kawamata, Y. *et al.* AG protein-coupled receptor responsive to bile acids. *J Biol Chem* **278**, 9435-
9440 (2003).
34 Thomas, C. *et al.* TGR5-mediated bile acid sensing controls glucose homeostasis. *Cell Metabol*
10, 167-177 (2009).
35 Islam, K. S. *et al.* Bile acid is a host factor that regulates the composition of the cecal microbiota
in rats. *Gastroenterol* **141**, 1773-1781 (2011).
36 Kurdi, P., Kawanishi, K., Mizutani, K. & Yokota, A. Mechanism of growth inhibition by free bile
acids in lactobacilli and bifidobacteria. *J Bacteriol* **188**, 1979-1986 (2006).
37 Ridlon, J. M., Kang, D. J., Hylemon, P. B. & Bajaj, J. S. Bile acids and the gut microbiome. *Curr
Opin Gastroenerol* **30**, 332 (2014).
38 Maier, T. V. *et al.* Impact of Dietary Resistant Starch on the Human Gut Microbiome,
Metaproteome, and Metabolome. *mBio* **8**, e01343-01317 (2017).
39 Sagar, N. M., Cree, I. A., Covington, J. A. & Arasaradnam, R. P. The interplay of the gut
microbiome, bile acids, and volatile organic compounds. *Gastroent Res Pract* **2015** (2015).
40 Daniel, H. *et al.* High-fat diet alters gut microbiota physiology in mice. *ISME J* **8**, 295-308 (2014).
41 Nie, Y.-f., Hu, J. & Yan, X.-h. Cross-talk between bile acids and intestinal microbiota in host
metabolism and health. *J Zhejiang University* **16**, 436 (2015).
42 Swinburn, B. A. *et al.* The global obesity pandemic: shaped by global drivers and local
environments. *Lancet* **378**, 804-814 (2011).
43 Buchwald, H. & Oien, D. M. Metabolic/bariatric surgery worldwide 2008. *Obes Surg* **19**, 1605-
1611 (2009).
44 Ilhan, Z. E. *et al.* Distinctive microbiomes and metabolites linked with weight loss after gastric
bypass, but not gastric banding. *ISME J* (2017).

45   Cottam, D. R., Atkinson, J., Anderson, A., Grace, B. & Fisher, B. A case-controlled matched-pair
cohort study of laparoscopic Roux-en-Y gastric bypass and Lap-Band® patients in a single US
center with three-year follow-up. *Obes Surg* **16**, 534-540 (2006).
46   Buchwald, H. *et al.* Bariatric surgery: a systematic review and meta-analysis. *Jama* **292**, 1724-
1737 (2004).
47   Sweeney, T. E. & Morton, J. M. The human gut microbiome: a review of the effect of obesity and
surgically induced weight loss. *JAMA Surg* **148**, 563-569 (2013).
48   Woodard, G. A. *et al.* Probiotics improve outcomes after Roux-en-Y gastric bypass surgery: a
prospective randomized trial. *J Gastrointest Surg* **13**, 1198-1204 (2009).
49   Albaugh, V. L. *et al.* Early increases in bile acids post Roux-en-Y gastric bypass are driven by
insulin-sensitizing, secondary bile acids. *J Clin Endocrinol Metab* **100**, E1225-E1233 (2015).
50   Ryan, K. K. *et al.* FXR is a molecular target for the effects of vertical sleeve gastrectomy. *Nature*
509, 183 (2014).
51   Cayuela, C. & Juste, C. Isolates from Normal Human Intestinal Flora but not Lactic Acid Bacteria
Exhibit 7a-and 7b-Hydroxysteroid Dehydrogenase Activities. *Microb Ecol Health D* **16** (2004).
52   Hirano, S. & Masuda, N. Characterization of NADP-dependent 7 beta-hydroxysteroid
dehydrogenases from Peptostreptococcus productus and Eubacterium aerofaciens. *Appl Environ*
Microbiol **43**, 1057-1063 (1982).
53   Floch, M. H., Binder, H. J., Filburn, B. & Gershengoren, W. The effect of bile acids on intestinal
microflora. *The American journal of clinical nutrition* **25**, 1418-1426 (1972).
54   Binder, H., Filburn, B. & Floch, M. Bile acid inhibition of intestinal anaerobic organisms. *The*
American journal of clinical nutrition **28**, 119-125 (1975).
55   Liu, L., Aigner, A. & Schmid, R. D. Identification, cloning, heterologous expression, and
characterization of a NADPH-dependent 7β-hydroxysteroid dehydrogenase from Collinsella
aerofaciens. *Appl Microbiol Biotechnol* **90**, 127-135 (2011).
56   Ferrandi, E. E. *et al.* In search of sustainable chemical processes: cloning, recombinant
expression, and functional characterization of the 7α-and 7β-hydroxysteroid dehydrogenases
from Clostridium absonum. *Appl Environ Microbiol* **95**, 1221-1233 (2012).
57   Li, J. V. *et al.* Metabolic surgery profoundly influences gut microbial–host metabolic cross-talk.
Gut **60**, 1214-1223 (2011).
58   Yatsunenko, T. *et al.* Human gut microbiome viewed across age and geography. *Nature* **486**, 222
(2012).
59   Huttenhower, C. *et al.* Structure, function and diversity of the healthy human microbiome.
Nature **486**, 207 (2012).
60   Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K. & Knight, R. Diversity, stability and
resilience of the human gut microbiota. *Nature* **489**, 220 (2012).
61   James, G. A. *et al.* Microsensor and transcriptomic signatures of oxygen depletion in biofilms
associated with chronic wounds. *Wound Rep Regen* **24**, 373-383 (2016).
62   Shu, M., Wong, L., Miller, J. & Sissons, C. Development of multi-species consortia biofilms of oral
bacteria as an enamel and root caries model system. *Arch Oral Biol* **45**, 27-40 (2000).
63   Niu, B., Paulson, J. N., Zheng, X. & Kolter, R. Simplified and representative bacterial community
of maize roots. *Proc Nat Acada Sci* **114**, E2450-E2459 (2017).
64   Khan, N. *et al.* Phenotypic responses to interspecies competition and commensalism in a
naturally-derived microbial co-culture. *Sci Rep* **8**, 297 (2018).
65   Bernstein, H. C. *et al.* Indirect interspecies regulation: transcriptional and physiological
responses of a cyanobacterium to heterotrophic partnership. *MSystems* **2**, e00181-00116
(2017).

66   Cole, J. K. *et al.* Phototrophic biofilm assembly in microbial-mat-derived unicyanobacterial
642   consortia: model systems for the study of autotroph-heterotroph interactions. *Front Microbiol* **5**,
643   109 (2014).

644   67   Beliaev, A. S. *et al.* Inference of interactions in cyanobacterial–heterotrophic co-cultures via
645   transcriptome sequencing. *ISME J* **8**, 2243 (2014).

646   68   Boetius, A. *et al.* A marine microbial consortium apparently mediating anaerobic oxidation of
647   methane. *Nature* **407**, 623 (2000).

648   69   Nakashima, T. *et al.* A paucity of unusual trihydroxy bile acids in the urine of patients with
649   severe liver diseases. *Hepatology* **29**, 1518-1522 (1999).

650   70   Tint, G. S. *et al.* Metabolism of ursocholic acid in humans: Conversion of ursocholic acid to
651   deoxycholic acid. *Hepatology* **15**, 645-650 (1992).

652   71   Batta, A. K., Salen, G. & Abroon, J. Ursocholic acid, a hydrophilic bile acid, fails to improve liver
653   function parameters in primary biliary cirrhosis: comparison with ursodeoxycholic acid. *Am J*
654   *Gastroenterol* **92** (1997).

655   72   Loria, P. *et al.* Effect of ursocholic acid on bile lipid secretion and composition. *Gastroenterol* **90**,
656   865-874 (1986).

657   73   Mythen, S. M., Devendran, S., Méndez-García, C., Cann, I. & Ridlon, J. M. Targeted synthesis and
658   characterization of a gene-cluster encoding NAD (P) H-dependent 3α-, 3β-, and 12α-
659   hydroxysteroid dehydrogenases from Eggerthella CAG: 298, a gut metagenomic sequence. *Appl*
660   *Environ Microbiol*, AEM. 02475-02417 (2018).

661   74   Ridlon, J. M. & Bajaj, J. S. The human gut sterolbiome: bile acid-microbiome endocrine aspects
662   and therapeutics. *Acta Pharm Sin B* **5**, 99-105 (2015).

663   75   Sutherland, J. D. & Macdonald, I. A. The metabolism of primary, 7-oxo, and 7 beta-hydroxy bile
664   acids by Clostridium absonum. *J Lipid Res* **23**, 726-732 (1982).

665   76   Macdonald, I. A. & Roach, P. D. Bile salt induction of 7α-and 7β-hydroxysteroid dehydrogenases
666   in Clostridium absonum. *Biochimica et Biophysica Acta (BBA)-Lipids and Lipid Metabolism* **665**,
667   262-269 (1981).

668   77   Mallonee, D. H., Lijewski, M. A. & Hylemon, P. B. Expression in Escherichia coli and
669   characterization of a bile acid-inducible 3α-hydroxysteroid dehydrogenase from Eubacterium sp.
670   strain VPI 12708. *Curr Microbiol* **30**, 259-263 (1995).

671   78   White, W., Franklund, C., Coleman, J. & Hylemon, P. Evidence for a multigene family involved in
672   bile acid 7-dehydroxylation in Eubacterium sp. strain VPI 12708. *J Bacteriol* **170**, 4555-4561
673   (1988).

674   79   Martin, F. P. J. *et al.* A top-down systems biology view of microbiome-mammalian metabolic
675   interactions in a mouse model. *Mol Sys Biol* **3**, 112 (2007).

676   80   Habibi, A. *et al.* The effects of vitrification on gene expression in mature mouse oocytes by
677   nested quantitative PCR. *Journal of assisted reproduction and genetics* **27**, 599-604 (2010).

678   81   Kelly, R. T. *et al.* Chemically etched open tubular and monolithic emitters for nanoelectrospray
679   ionization mass spectrometry. *Anal Chem* **78**, 7796-7801 (2006).

680   82   Kim, Y.-M. *et al.* Diel metabolomics analysis of a hot spring chlorophototrophic microbial mat
681   leads to new hypotheses of community member metabolisms. *Front Microbiol* **6**, 209 (2015).

682   83   Hiller, K. *et al.* MetaboliteDetector: comprehensive analysis tool for targeted and nontargeted
683   GC/MS based metabolome analysis. *Anal Chem* **81**, 3429-3439 (2009).

684   84   Kind, T. *et al.* FiehnLib: mass spectral and retention index libraries for metabolomics based on
685   quadrupole and time-of-flight gas chromatography/mass spectrometry. *Anal Chem* **81**, 10038-
686   10048 (2009).

687   85   Bernstein, H. C. *et al.* Multi-omic dynamics associate oxygenic photosynthesis with nitrogenase-
688   mediated H2 production in Cyanothece sp. ATCC 51142. *Sci Rep* **5** (2015).

86   Sadler, N. C. *et al.* Dinitrogenase-Driven Photobiological Hydrogen Production Combats
Oxidative Stress in Cyanothece sp. Strain ATCC 51142. *Appl Environ Microbiol* **82**, 7227-7235
(2016).
87   Team, R. C. R: A language and environment for statistical computing. (2013).
88   Oksanen, J. *et al.* The vegan package. *Community ecology package* **10**, 631-637 (2007).
89   Csardi, G. & Nepusz, T. The igraph software package for complex network research. *InterJournal,*
Complex Systems **1695**, 1-9 (2006).
90   Wickham, H. Tidyverse: Easily install and load'tidyverse'packages. *R package version* **1** (2017).