1 **Environmental sources of bacteria and genetic variation in behavior influence**
2 **host-associated microbiota**
3

Alexandra A. Mushegian*[1,3], Roberto Arbore*[1,4], Jean-Claude Walser[1,2], Dieter Ebert[1]

*These authors contributed equally to this work

[1]Zoological Institute, Universität Basel, Switzerland

[2]Genetic Diversity Centre, ETH Zürich, Switzerland

[3]To whom correspondence should be addressed: amushegian@gmail.com. Current

affiliation: American Society for Biochemistry and Molecular Biology, USA

[4]Current affiliation: École normale supérieure de Lyon, France

Keywords: microbiome, Daphnia, habitat selection, genotype by environment

interactions

Summary statement: Genetic differences in Daphnia behavior contribute to the

amount of environmental bacteria present in their microbiome

Abstract

        In many organisms, host-associated microbial communities are acquired horizontally after birth. This process is believed to be shaped by a combination of environmental and host genetic factors. We examined whether genetic variation in animal behavior could affect the composition of the animal's microbiota in different environments. The freshwater crustacean *Daphnia magna* is primarily planktonic, but exhibits variation in the degree to which it browses in benthic sediments. We performed an experiment with clonal lines of *D. magna* showing different levels of sediment-browsing intensity exposed to either bacteria-rich or bacteria-poor sediment or whose access to sediments was prevented. We find that the bacterial composition of the environment and genotype-specific browsing intensity together influence the diversity and composition of the *Daphnia*-associated bacterial community. Exposure to more diverse bacteria did not lead to a more diverse microbiome, but greater abundances of environment-specific bacteria were found associated with host genotypes that exhibited greater browsing behavior. Our results indicate that individual behavior can mediate genotype-by-environment interaction effects on microbiome composition.

Introduction

        Every multicellular organism is colonized by a community of microorganisms: its microbiota (McFall-Ngai et al. 2013). The host provides a habitat for a complex and dynamic consortium of microorganisms, many of which have fundamental influences on the host's well-being (Zilber-Rosenberg & Rosenberg 2008). A central concern in both infectious disease epidemiology and in studies of host-associated microbial community ecology is the transmission of microbes between host individuals and between hosts and the environment. Many bacterial assemblages are transmitted from host mother to offspring (Funkhouser & Bordenstein 2013; Salem et al. 2015), but the diversity of microbiota typically changes over time depending on the microbes available in the environment (Rajilic-Stojanovic et al. 2009; Shin et al. 2015). In some cases, environmentally acquired microbes are even essential for the completion of postembryonic development (e.g.

Cheesman et al. 2011; Diaz Heijtz et al. 2011). Thus microbes from the environment

can be co-opted as part of the microbiota, or can affect host health during a transient

occupation (Voss et al. 2015).

Environmental effects on microbiota community structure have been

extensively documented (Fan et al. 2013; Seedorf et al. 2014) and studies on model

organisms have started to shed light on the relative importance of environmental

and host genetic factors in determining microbiota composition (Campbell et al.

2012; Spor et al. 2011). Recently, the focus has been moving towards a better

understanding of the mechanisms of bacterial aqusition from the environment. Host

genetics have been shown to play a role in the establishment of microbial

associations through microbial recognition, immune selection, and determination of

the biochemical niche (Spor et al. 2011). Importantly, these processes select

microbes after the host has come in contact with bacterial communities in the

environment. The initial encounter may be a key phase of the host's colonization by

microbes. If host genetics influence interaction with the environment, for example

through the expression of behavioral variation, it may influence the initial

encounters with environmental bacteria and thus affect the composition of the host

microbiota.

Many animals utilize different habitats according to behavioral strategies

collectively termed habitat selection. If habitats differ in their microbial

communities, host behavior influencing habitat choice and the microbiome may

influence each other. Hosts may have evolved strategies to ensure or avoid

encounter with beneficial and pathogenic microorganisms. Avoidance behaviors of

harmful bacteria are well documented, and behavior is considered one of the first

lines of defense against infectious disease. For example, the nematode

Caenorhabditis elegans actively avoids pathogenic bacteria and the genetic

determinants of this behavior have been worked out (Meisel & Kim 2014). The

opposite case, where a host's behavior is involved in the acquisition of beneficial

bacteria from the environment, has received less attention, despite speculation

about the role of human behaviors such as outdoor play in preventing autoimmune

diseases (Rook 2013). The overall effects of host habitat choice behavior on

microbiome composition have not, to our knowledge, been explored in any system.

An analysis of natural genetic variation in behavioral traits potentially influencing

microbiota acquisition is therefore timely (Ezenwa et al. 2012). If variation in

behavior affects the composition of the host's microbial community, then behavior

could underlie some genotype-environment interaction effects on microbiota. The

goal of this study was to examine the effect of genetic variation in host behavior on

microbiota composition in different environments using the freshwater planktonic

crustacean *Daphnia magna*.

Recently, it has been shown that *D. magna* microbiota play a major role in

host fitness (Sison-Mangus et al. 2015), that both host clonal line and environmental

factors are determinants of microbiota community structure (Sullam et al. 2017)

and that genotype-specific microbiomes can mediate daphnids' adaptive traits

(Macke et al. 2017). However, little is known about the mechanisms by which the

host acquires microbiota from the environment. A specific behavior, termed

sediment browsing, mediates the interaction between *D. magna* and bottom

sediments of ponds and lakes (Horton et al. 1979; Decaestecker et al. 2002). During

browsing, the animals swim along the sediment surface, stirring up particles, and

then ingest the particles by filter feeding. Besides representing valuable food

reservoirs, sediments are likely important environmental sources of bacteria.

Therefore, the physical contact with the sediments resulting from browsing might

present both disease risks and benefits from increased contact with bacteria.

Previous work found evidence of genetic variation for the levels of browsing activity

in *D. magna* (Arbore et. al. 2016).

We performed a laboratory experiment in which we analyzed the browsing

behavior and microbiota of 12 genetically distinct *D. magna* clones allowed to

browse in sediment. The animals were exposed to three different treatment

conditions, where they had access to either previously autoclaved (i) or untreated

("natural" and therefore microbe-rich) sediments (ii), or where their access to

natural sediment was prevented (iii) (Figure 1A). We hypothesized that *D. magna*

clones exhibiting more intense browsing behavior would have more diverse

microbiota in conditions where they had access to bacteria-rich sediment. In this

experiment, we made no assumptions about whether bacteria found in the

sediments were beneficial, harmful, or neutral for the host, nor whether they

colonized *Daphnia* stably or transiently; therefore, the patterns observed here could

be applicable to studies of disease, microbiota, or general environmental microbial

community dynamics. Our analysis illustrates how a behavioral trait can mediate

the interplay between genetic and environmental variation in the establishment of

host-microbe associations.

Methods

Overview of the experiment

In this study, we combined the analysis of animals with constitutive (genetic)

differences in browsing behavior with manipulations of the environment that

affected animals' access to the sediments. Animals were either exposed to natural

sediment, to autoclaved sediment, or to natural sediment blocked by a permeable

net barrier (Figure 1A). In order to analyze both the browsing behavior of the

animals and their microbiota, we placed two animals in each jar; of these pairs of

animals, after 6 days of exposure to the different treatments, one animal was used to

assay browsing behavior while the other was used for microbiota analyses.

Experimental animals

D. magna reproduces by cyclical parthenogenesis. Clonal populations can be

generated and propagated in the laboratory through asexual reproduction. Here, we

refer to such genetically identical individuals as "replicates" or "animals" while we

refer to different genetic lines as "clones." In this study we used 12 *D. magna* clones

from our stock collection, originating from different populations (Table 1). The

animals were propagated from stock cultures maintained in the laboratory in

standardized conditions and without any effort to modify their microbiota. The

browsing behavior of these clones has been assessed before (Arbore et al. 2016) and

was shown to differ among genotypes.

All animals used in this study were females. Prior to the experiment, every

clone was propagated in individual replicates for three generations in order to

minimize variation due to maternal effects. These animals were kept individually in

100-ml glass jars filled with 80 ml of ADaM (*Daphnia* medium (Kluttgen et al. 1994))

randomly distributed within trays in incubators with a 16:8 light/dark cycle and

constant temperature of 20 °C. To establish every generation, the animals were

isolated at 4 days old and fed daily with chemostat-grown green algae *Scenedesmus*

sp: 1×10^6 algae cells/animal until day 5, 2×10^6 until day 8, 2.5×10^6 until day 10, 3

$\times 10^6$ until day 12, and 5×10^6 onwards. The animals were transferred to fresh

medium when they were 12 days old and thereafter every day.

For the experiment, we used animals from the 4th generation of each of the

12 clones. These animals were kept in groups of 8 siblings belonging to one clutch of

one mother. At 4 days old (± 1 day), 6 animals from every clutch were randomly

assigned in pairs to individual jars divided into the three different treatments (split

brood design); each such jar containing a pair of animals was an experimental

replicate. In total, we included in the experiment 540 animals (270 pairs)

corresponding to 15 pairs of clone BE-OHZ-T10, 18 pairs of clones DE-K35-Mu10

and NO-V-7, 21 pairs of clone F2-918, and 24 pairs of each of the remaining clones.

Variation in replicate numbers resulted from differences in availability of female

offspring at the time that the treatments were established.

Experimental design

The experiment was conducted in cylindrical glass jars (height = 20 cm;

diameter = 6.5 cm) (Fig. 1A) kept in cardboard boxes on shelves in a climate room

(16:8 light/dark cycle at 20 °C), loosely covered with transparent plastic film and

top-illuminated with neon lights. In this way, light only entered the jars from the

top. All the experimental jars were first filled with 400 ml of medium. 15 ml of a

suspension of loess (fine silt) was then carefully deposited on the bottom using a

serological pipette. The loess was previously collected from a soil stock, suspended

in water, passed through a 200 μm filter and washed to remove very fine particles.

After two days of sedimentation, the loess formed a 1 cm layer at the bottom of the

jar. Then, an acrylic tube (height = 21 cm; diameter = 5 cm) was inserted into the

jars and kept in position with a plastic ring fixed to the opening of the jar, so that its

lower end was positioned close to the sediment surface. In one treatment (NET), the

acrylic tube was closed with a 500 μm net at the lower end (suspended 5 mm above

the sediment surface) preventing animals from direct contact with the sediment

(Fig.1A left). In the other two treatments (AUT and SED), the acrylic tubes had no

net so that the animals had free access to the sediment (Fig.1A right). In the AUT

treatment, the loess was previously autoclaved while in the SED and the NET

treatment the loess was left untreated ("natural"). (After autoclaving, AUT sediment

was handled in the same way as natural sediment, i.e. exposed to nonsterile media

and laboratory environment.) After inserting the tubes, the jars were left

undisturbed for two days before the animals were introduced in order to allow the

sediment to settle. Immediately before the experiment, the sediments of three jars

of each of the SED and the AUT treatment were sampled and frozen at -20 °C; these

sampled jars were not used further.

Two animals from the same clutch were carefully introduced into the inner

tube of each jar. The 264 jars, each containing one pair of animals, were evenly

distributed among the treatments and their positions in the incubator room were

randomized. The animals remained in the experimental jars for 6 days. During this

time, the animals were carefully fed twice daily with 2.5 x 10^6 algal cells. At day 6, all

animals were collected and one member of every pair was assigned to the

behavioral assay (see below) and the other was frozen for later DNA extraction. 32

pairs of animals were lost or damaged during the experiment and were excluded

from further analyses. At the end of the experiment, 3 sediment samples from the

NET treatment and 3 sediment samples from the AUT treatment were collected and

frozen at -20°C.

Behavioral analysis

The animals for the behavioral assay were transferred individually from the

sediment jars to 100-ml glass jars filled with medium and kept in an incubator and

fed daily with 5 x 10^6 algal cells. The behavior assay was conducted over two days

when the animals were 12 to 14 days old with all replicates for the different clone
by treatment combinations evenly distributed across time. The behavior assay was
performed as described in Arbore et al. (2016). Briefly, we measured the traces left
by individual replicate animals on a smooth surface layer of sediment (loess) at the
bottom of tall cylindrical glass jars (20 cm tall, 6.5 cm diameter; Fig. 1B) during 30
minutes. The sediment surface was photographed before animals were released
(time 0), using a ring light to ensure uniform illumination. The jar was then
transferred into a cardboard tube and illuminated from the top with a neon light
and one animal was introduced in each jar. After exactly 30 minutes, the animal was
removed and the sediment surface was again photographed (time 1), in the same
position and under the same light conditions. Using the software ImageJ
(http://rsb.info.nih. gov/ij/), the pictures were converted to grey scale and a central
circular area was cropped to exclude shadows from the edge of the jar (Fig. 1B).
Pictures were processed such that the browsing traces of the animals on the
sediment surface resulted black areas against a white background. Then the number
of black pixels was quantified. Pictures taken at time 0 were used to correct the
values calculated for the browsing traces when irregularities on the sediment
surface were detected (i.e. in cases the picture of time 0 was not entirely white). The
pixel values were then log-transformed ($[\log_{10}(X+1000)]$; 1000 corresponds
approximately to the number of pixels of one individual browsing trace). During the
assay, four animals were accidentally damaged while handling and were excluded
from the analyses. The body lengths of the animals used for behavior analysis were
measured after the behavioral assay.
The adjusted intra-class correlation coefficient for the browsing behavior
(equivalent to broad sense heritability) was calculated with a linear mixed effect
(LMM) model, with treatment as a fixed effect and clone as a random effect (R
software package rptR developmental version; (Nakagawa & Schielzeth 2010)).
Confidence intervals and statistical significance were calculated using parametric
bootstrapping with 5000 iterations and a randomization procedure with 5000
permutations.
233

DNA extraction, library preparation and sequencing

The animals assigned to the microbiota analysis were transferred

individually from the sediment treatment jars to 40 mL of autoclaved ADaM for

about 2 hours to dilute carryover of unattached bacteria. Then, the animals were

transferred into 1.5 ml Eppendorf tubes, the ADaM was removed and the tubes were

stored at -20 °C until DNA extraction.

DNA was extracted from single animals using a cetyltrimethylammonium

bromide (CTAB)-based protocol. The animals were ground with a sterile pestle in

1.5 ml Eppendorf tubes in a 10 mg/ml lysozyme solution and mixed at 850 rpm and

37 °C for 45 minutes. Then, a 20 mg/ml solution of proteinase K was added and the

tubes mixed at 850 rpm and 55 °C for 1 hour. After an RNase treatment (20 mg/ml)

at room temperature for 10 minutes, a preheated 2X solution of CTAB was added

and the tubes mixed at 300 rpm and 65 °C for 1 hour. After two rounds of

chloroform isoamyl alcohol (CIA) purification (1 volume CIA; 8 minutes

centrifugation at 12,0000 rpm and 15 °C), a solution of sodium acetate 3M pH 5.2

and isopropanol were added to the DNA solution and the tubes were stored

overnight at -20 °C. The following day, DNA was purified by two rounds of 70 %

ethanol precipitation and suspended in water. The extractions were then incubated

at 4 °C overnight and then stored at -20 °C.

All DNA extractions were conducted over a period of 6 days with samples

from the different clone by treatment combinations randomly distributed between

the days and one reagent-only negative control extraction included every day. DNA

from the sediment samples and from one negative control was extracted on a

different day using a commercial kit (PowerSoil® DNA Extraction kit; MO BIO

Laboratories, cat. 12888-100).

We sequenced amplicons of the V3-V4 variable region of the bacterial 16S

rRNA gene using the Illumina MiSeq platform. Amplicons were generated using

NEBNext High Fidelity PCR Master Mix (New England Biolabs catalog#M0541L) for

27 cycles in 25 µl reactions containing 3% DMSO. The primers used were 341F (5'-

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGA-3') and 785R (5'-

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGCAGA-3') with Illumina adapter

sequences and 0-3 bp random frameshifts. PCR product was purified with Ampure

beads at a 0.6x ratio of beads to PCR product, amplified for 9 cycles with Nextera XT

v2 indexing primers, and purified again. Libraries were quantified with Qubit and

quantitative PCR, normalized, and pooled, followed by additional bead purification

to remove remaining short fragments before sequencing on the Illumina MiSeq

(reagent kit v3, 300 bp paired-end reads). The same library pool was used for two

MiSeq runs; after checking that there was no statistical difference in community

composition between the runs (Adonis analysis of Bray-Curtis dissimilarity between

samples, p=0.394), the data from the two runs were merged.

Sequence quality control

Raw reads were quality controlled with FastQC (Babraham Institute, UK).

Paired reads were merged (FLASH v1.2.9), primers trimmed (Cutadapt v1.5), and

quality filtered (PRINSEQ-lite v0.20.4). OTU clustering including abundance sorting

and chimera removal was performed using the UPARSE workflow (Edgar 2013).

Only those OTUs represented by 5 or more reads in the run were included.

Taxonomic assignment was performed using UTAX against the GreenGenes v13/5

database. We analyzed samples with more than 5000 total reads. This left 214

samples; numbers of replicates for each combination of variables are reported in

Table 1.

Since individual *Daphnia* contain low bacterial biomass, we considered the

issue of reagent contamination with bacterial DNA (Salter et al. 2014). Samples were

processed in haphazard order, so erroneous sequences originating from reagent

contamination were expected to be distributed randomly and not confounded with

any treatment or genotype. For our research question, we were interested in

patterns of diversity and changes in composition in response to experimental

factors rather than in the presence or absence of any particular strain. For all

analyses, we first tested for processing batch effects and stratified the main analysis

by batch if they were significant.

For statistical analyses in which host clone was a fixed effect, we excluded

clone NO-V-7, since it did not have at least 3 replicates in each treatment; we

included this clone in analyses where clone was treated as a random effect. We

examined the effects of experimental factors on both overall diversity and the

community composition of each animal's microbiota using standard ecological

diversity indices and ordination methods. To evaluate the effect of animal behavior

on microbiota, we used as proxies for individual behavior either the mean browsing

intensity index of the clone or the browsing intensity of the individual co-housed

with the sequenced individual in the same jar ("jar-mate").

Analyses were carried out in R (3.4.3), using the packages phyloseq (1.22.3),

vegan (2.4.6), plyr (1.8.4), dplyr (0.7.4), DESeq2 (1.18.1), nlme (3.1.131), lme4

(1.1.15), metacoder (0.2.0.9012), and ggplot2 (2.2.1).

Results

Browsing intensity, animal microbiota, and sediment bacteria

Consistently with previous studies (Arbore et al. 2016), browsing behavior

intensity varied among _Daphnia_ clones (Fig. 2). Clone and treatment, but not their

interaction, had a significant effect on browsing behavior (analysis of variance:

clone F=12.717, df=11, p<0.0001; treatment F=4.100, df=2, p=0.018;

clone*treatment F=1.274, df=22, p=0.193). The average browsing intensity of

animals from the NET treatment was lower than that of animals in the SED and AUT

treatments (Fig. S1). The total phenotypic variance for browsing behavior explained

by clone, after controlling for the treatment effect, corresponded to 36.5% (95% CI

= [13.2, 55.3%], p = 0.0002). Clone but not treatment had a significant effect on body

size, so we assume that access to (and type of) sediment did not substantially affect

nutrition and growth over the timeframe of the experiment (analysis of variance:

clone F=8.08, df=11, p < 0.001; treatment F=2.01, df=2, p=0.137; clone*treatment

F=1.43, df=22, p=0.103). Individual body size was uncorrelated with behavior

(analysis of variance: F=0.346, df=1, p=0.56; Fig. S2).

A total of 370 OTUs were found among the animal samples; of these, 318

were found in less than 10% of samples. (See Fig. S3A-C for taxonomic heat trees of

OTUs with presence/absence information) (Foster et al. 2016). Consistently with

multiple previous studies of *Daphnia* microbiota (Peerakietkhajorn et al. 2015; Qi et

al. 2009; Callens et al. 2015; Eckert & Pernthaler 2014), the most abundant bacterial

species was a single strain (OTU_1) of *Limnohabitans* sp (Betaproteobacteria,

Comamonadaceae), with a mean relative abundance across all clones of 0.39 (s.e.m.

0.02). Interestingly, a second strain of *Limnohabitans* (OTU_2) was a dominant

strain only in the three clones originating from clones bred in the laboratory as part

of a genetic breeding design (QTL panel; 0.32 mean relative abundance among

individuals of clones IXF1, F2-82, F2-918; 0.0016 mean relative abundance in

remaining clones). As expected, the sediment originating from the SED treatment

had much higher bacterial species richness than that originating from the AUT

treatment (Fig. S4).

Effects of treatment and clone on alpha diversity

Both *Daphnia* clone and treatment, but not their interaction, had significant

effect on the Shannon and inverse Simpson alpha diversity indices (Table 2). For

further analyses, we focused on the Shannon index, because it takes into account not

only species richness but also evenness (with additional species given more weight

as they become more abundant). The Shannon index displayed no significant effect

of processing batch (df=5, F=1.42, p=0.22). Shannon diversity estimates for the 12

clones arranged in order of increasing average browsing intensity and the three

groups (AUT, NET and SED) are shown in Fig. 3 (species richness and inverse

Simpson index are shown in Fig. S5A-B). Unexpectedly, the highest average alpha

diversity in most clones (9/12) was observed in the AUT treatment group, despite

their exposure to less-diverse sediment than the SED group. Therefore, diversity of

animal microbiota does not directly reflect diversity of bacteria in the environment.

To specifically investigate the effect of direct access to the same bacteria-rich

sediment, we compared the NET and SED treatment groups' diversity as a function

of clonal average behavior in each group (Fig. 4A). The difference in mean Shannon

diversity between SED and NET animals was greatest at the highest average clonal

level of browsing intensity (Fig. 4B; linear regression p=0.055). A similar tendency

could be seen when the browsing intensity of each individual's jar-mate was used as

the proxy for individual behavior (Fig. S6). Shannon diversity significantly depended

on the interaction between treatment and clonal average browsing intensity in a

linear mixed-effects model with clone included as a random effect (Table 3); the

same was true when treatment-specific clonal average behavior was used as the

behavior proxy, but not when individual jar-mate behavior was used (Table S1).

Community composition and acquisition of bacteria from sediment

365   To examine shifts in bacterial community composition in response to

environmental treatments, we Hellinger-transformed the bacterial abundances by

taking the square root of the relative abundance of each taxon in each sample to

reduce the influence of rare taxa, and then calculated pairwise Bray-Curtis distances

between samples. The average distance to the centroid (dispersion) was lower in

the NET group than in the AUT and the SED groups (Fig. 5), suggesting that access to

sediment increases variability of microbiota regardless of the composition of the

sediment. To see whether the different sediments resulted in systematically

different microbiota composition, we excluded the NET group and performed

principal coordinates analysis (PCoA) (Fig.6). Adonis analysis stratified by

processing batch showed that both treatment and clone had a significant effect

(treatment: R^2=0.05, p=0.001; clone: R^2=0.16, p=0.001), but not their interactions

(treatment:clone p=0.63). However, clones also showed significant differences in

dispersion (p<0.001).

379   Having confirmed that differences in the sediment environment resulted in

differences in animal microbiota composition, we next explored the extent to which

environment-specific bacteria contributed to these differences. We used DESeq2 to

determine which bacteria were significantly more present in natural sediment than

autoclaved sediment (n=3 each). 115 OTUs were calculated to be significantly

differentially present between the two sediment types; of these, 48 had at a

log2-fold increase of at least 8 in natural sediment compared to autoclaved

sediment. We refer to these as natural-sediment-derived taxa. The 8-fold threshold

was chosen based on inspecting the data; similar results were seen when sediment-

derived bacteria were defined by a log2-fold change of 5 or 10; see Fig. S7. Only one

of these OTUs was found in a majority of animals, and the median number of

animals in which a given OTU was found was 6.5. We therefore concluded that

animals likely acquired environmental bacteria randomly rather than selectively

from the environment. Accordingly, we looked at the total relative abundance of all

natural sediment–derived bacteria in each individual.

The relative abundance of natural-sediment-derived bacteria was generally

low in both the AUT and NET treatment groups, and increased with browsing

intensity in the SED treatment group (Fig. 7), with a significant interaction effect

between treatment and clonal average browsing intensity (Table 4). Treatment-

specific clonal average behavior showed the same significant interaction effect with

treatment, but the interaction effect was not significant when jar-mate behavior was

used as the behavior proxy. Among the set of clones examined here, an appreciably

high relative abundance of sediment-derived bacteria was detectable mainly in

clones with a browsing intensity index higher than mean 4.4 (clones IXF1, NO-V-7,

DE-KA-F28, CZ-N2-6, CZ-N1-1, F2-918). The mean relative abundance of sediment-

derived bacteria in the SED treatment in the pooled animals from these clones was

0.14 (s.e.m. 0.026), whereas it was 0.048 (s.e.m. 0.0097) in the lower-browsing

clones. Across all clones in the AUT and NET treatment groups, the average relative

abundance of sediment-derived bacteria was 0.051, nearly identical to that of the

low-browsing clones in the SED conditions.

Discussion

Our results have several implications for studies of animal-associated

microbiota in diverse environmental settings. First, we confirm the intuition that

environmental sources of bacteria affect the diversity of animal microbiota, but not

because more diverse environments always create more diverse microbiota; rather,

the animals we exposed to the less species-rich autoclaved sediments had higher

overall diversity in their microbiota than those exposed to untreated, bacterial-

species-rich sediment. We hypothesize that this might be due to competitive

interactions between *Daphnia* microbiota and the particular microbes found in

these sediments. The untreated sediments may contain bacteria that can

outcompete multiple strains of "native," preexisting *Daphnia* microbiota. If this were

the case, then browsing in sediment could have multiple opposing effects on overall

microbiota diversity: on the one hand, it would bring daphnids into contact with

more diverse bacteria, but on the other hand those bacteria could reduce existing

microbiota diversity. In the NET treatment, animals might be exposed to some

sediment-derived bacteria in the water column but lack access to the full diversity of

bacteria in the sediment. We also saw that having access to either sediment

increased the variability of community composition as measured by multivariate

dispersion. These results suggest that having access to multiple habitats with

different bacterial communities can affect the diversity and composition of an

animal's microbiota. Therefore, fine-scale heterogeneity in a host's habitat might be

a relevant aspect to take into account when examining effects of environment on

animal microbiota. This is especially important when considering ecological

immunology, because disease-causing bacteria in the environment may cause short-

term risk but also long-term fitness benefit via processes like immune priming

(Kaltenpoth & Engl 2014; Olszak et al. 2012).

436   Our data further suggest that the diversity of *Daphnia*-associated microbiota

in a particular environment may to some extent be mediated by genotype-specific

sediment browsing intensity. This was apparent as the net barrier made the greatest

difference in microbial alpha diversity in high-browsing host clones. However, this

effect may be partially obscured by several factors: the hypothesized competitive

exclusion effects we allude to above, and also non-behavior-related host genotype

effects on microbiota diversity. While host genotype had an effect on microbial

diversity, the highest- and lowest-browsing clones in our study had similar

microbial alpha diversity overall. It was only in evaluating the difference between

presence and absence of the barrier that an effect of browsing on diversity could be

seen. We conclude that the effect of environmental bacteria on host-associated

microbiota is not additive. The clearest effect of environmental bacteria on host-

associated microbiota was not on alpha diversity, but relative abundances of certain
taxa.

Clones with low average browsing intensity had no greater amount of
sediment-specific bacteria than animals exposed to autoclaved sediment or
prevented from browsing, whereas those with high browsing intensity could reach
over 60% of reads from environment-derived bacteria in some individuals. While
many studies of animal microbiota rightly concern themselves with distinguishing
between truly "host-associated" microbiota versus "transient environmental"
microbiota, these results raise the possibility that the amount of environmental
microbes found in association with an animal could itself be a host-genotype-
specific feature of the microbiome.

A key question for understanding symbiosis that future studies should
address is whether browsing behavior affects community composition by simple
exposure to more colonizing bacteria, or by more frequent replenishment of
bacterial taxa that would not otherwise persist in association with the host (Blum et
al. 2013). For example, browsing frequently enough may replenish bacteria that
would otherwise be lost when the animal molts. In this study, we made no
assumptions about the type of the interactions between the sediment-associated
bacteria and the *Daphnia*, but still were able to see that presence of sediment-
associated bacteria affected the bacteria that *Daphnia* were carrying.

It would also be interesting to investigate whether carriage of bacteria on
Daphnia from the sediment into the water column affects bacterial dynamics in the
larger environment; previous studies have shown that movement of *Daphnia*
between benthic and limnetic environments represents a mechanism of bacterial
dispersal in the environment (Grossart et al. 2010). Studies using classification
methods more sensitive than 16S-based taxonomy may be necessary to
unambiguously distinguish and assign sources to different bacterial strains.
475

Conclusion

We show that at least some characteristics of host-associated microbial
community composition result from genotype-by-microhabitat interactions,

specifically ones resulting from genotype-specific variation in behavior. Behavior

may thus be considered a genetic factor that shapes microbial exposure in a given

environment. Overall, these results provide further evidence that environment,

behavior, genetics, disease risk, and microbial community composition are

interrelated in potentially complex ways. Our observations indicate a need for more

integrative eco-immunology studies, in which the interfaces between behavioral

ecology, microbial community ecology and evolution of immune function are

explored. Studies can take advantage of the experimental tractability of the

Daphnia-microbiota system to further investigate these relationships in mechanistic

detail.

Acknowledgments

We thank Daniel Lüscher for designing the experimental jars, Jürgen Hottinger for

laboratory support, and the Botanical Institute of the University of Basel for

providing sediment. Illumina sequencing was performed at the Genetic Diversity

Centre of the ETH Zürich; thank you to Aria Minder and Silvia Kobel for advice on

library preparation.

Funding

This work was funded by an ERC Advanced Grant (268596-MicrobiotaEvolution).

Author contributions

AAM and RA conceived the study and performed the experiment with input from

DE. RA performed behavioral assays and analysis. AAM and RA prepared sequencing

libraries. JCW performed quality control and OTU clustering of sequence data. AAM

analyzed sequencing data. AAM and RA wrote the paper. DE and JCW revised the

paper.

Data deposition

Sequence data will be deposited in the European Nucleotide Archive of the EBI

upon publication. Data tables and code used for analysis can be found on Github at

References

Arbore, R. et al., 2016. Ecological genetics of sediment browsing behaviour in a

planktonic crustacean. *Journal of Evolutionary Biology*, 29(10), pp.1999–2009.

Blum, J.E. et al., 2013. Frequent Replenishment Sustains the Beneficial Microbiome

of Drosophila melanogaster. *mBio*, 4(6), pp.1–8.

Callens, M. et al., 2015. Food availability affects the strength of mutualistic host–

microbiota interactions in Daphnia magna. *The ISME Journal*, 10, pp.911–920.

Campbell, J.H. et al., 2012. Host genetic and environmental effects on mouse

intestinal microbiota. *The ISME Journal*, 6(11), pp.2033–2044.

Cheesman, S.E. et al., 2011. Epithelial cell proliferation in the developing zebrafish

intestine is regulated by the Wnt pathway and microbial signaling via Myd88.

Proceedings of the National Academy of Sciences of the United States of America,

108 Suppl, pp.4570–7.

Decaestecker, E., De Meester, L. & Ebert, D., 2002. In deep trouble: Habitat selection

constrained by multiple enemies in zooplankton. *Proceedings of the National*

Academy of Sciences, 99(8), pp.5481–5485.

Diaz Heijtz, R. et al., 2011. Normal gut microbiota modulates brain development and

behavior. *Pnas*, 108(7), pp.3047–3052.

Eckert, E.M. & Pernthaler, J., 2014. Bacterial epibionts of Daphnia: a potential route

for the transfer of dissolved organic carbon in freshwater food webs. *The ISME*

Journal, 8(9), pp.1808–1819.

Edgar, R.C., 2013. UPARSE: highly accurate OTU sequences from microbial amplicon

reads. *Nature Methods*, 10(10), pp.996–8.

Ezenwa, V.O. et al., 2012. Animal Behavior and the Microbiome. *Science*, 338(6104),

pp.198–199.

Fan, L. et al., 2013. Marine microbial symbiosis heats up: the phylogenetic and

functional response of a sponge holobiont to thermal stress. *The ISME journal*,

7(5), pp.991–1002.

Foster, Z.S.L., Sharpton, T. & Grunwald, N.J., 2016. MetacodeR: An R package for
manipulation and heat tree visualization of community taxonomic data from
metabarcoding. *bioRxiv*, p.71019.

Funkhouser, L.J. & Bordenstein, S.R., 2013. Mom knows best: the universality of
maternal microbial transmission. *PLoS biology*, 11(8), p.e1001631.

Grossart, H.-P. et al., 2010. Bacteria dispersal by hitchhiking on zooplankton.
Proceedings of the National Academy of Sciences of the United States of America,
107(26), pp.11959–11964.

Horton, P. et al., 1979. Browsing and grazing by cladoceran filter feeders. *Canadian*
Journal of Zoology-Revue Canadienne De Zoologie, 57(206–212).

Kaltenpoth, M. & Engl, T., 2014. Defensive microbial symbionts in Hymenoptera.
Functional Ecology, 28(2), pp.315–327.

Kluttgen, B. et al., 1994. ADaM, an artificial freshwater for the culture of
zooplankton. *Water Research*, 28(3), pp.743–746.

Macke, E. et al., 2017. Host-genotype dependent gut microbiota drives zooplankton
tolerance to toxic cyanobacteria. *Nature Communications*, 8(1).

McFall-Ngai, M. et al., 2013. Animals in a bacterial world, a new imperative for the
life sciences. *Proceedings of the National Academy of Sciences of the United*
States of America, 110(9), pp.3229–3236.

Meisel, J.D. & Kim, D.H., 2014. Behavioral avoidance of pathogenic bacteria by
Caenorhabditis elegans. *Trends in Immunology*, 35(10), pp.465–470.

Nakagawa, S. & Schielzeth, H., 2010. Repeatability for Gaussian and non-Gaussian
data: A practical guide for biologists. *Biological Reviews*, 85(4), pp.935–956.

Olszak, T. et al., 2012. Microbial exposure during early life has persistent effects on
natural killer T cell function. *Science (New York, N.Y.)*, 336(6080), pp.489–93.

Peerakietkhajorn, S. et al., 2015. Betaproteobacteria Limnohabitans strains increase
fecundity in the crustacean Daphnia magna: Symbiotic relationship between
major bacterioplankton and zooplankton in freshwater ecosystem.
Environmental Microbiology, 18, pp.2366–2374.

Qi, W. et al., 2009. Comparative metagenomics of Daphnia symbionts. *BMC Genomics*,
10(1), p.172.

Rajilic-Stojanovic, M. et al., 2009. Development and application of the human
intestinal tract chip, a phylogenetic microarray: Analysis of universally
conserved phylotypes in the abundant microbiota of young and elderly adults.
Environmental Microbiology, 11(7), pp.1736–1751.

Rook, G.A., 2013. Regulation of the immune system by biodiversity from the natural
environment: an ecosystem service essential to health. *Proc Natl Acad Sci USA*,
110(46), pp.18360–18367.

Salem, H. et al., 2015. An out-of-body experience : the extracellular dimension for
the transmission of mutualistic bacteria in insects. *Proceedings of the Royal*
Society B, 282(1804), pp.1–10.

Salter, S.J. et al., 2014. Reagent and laboratory contamination can critically impact
sequence-based microbiome analyses. *BMC Biology*, 12(1), p.87.

Seedorf, H. et al., 2014. Bacteria from diverse habitats colonize and compete in the
mouse gut. *Cell*, 159(2), pp.253–266.

Shin, H. et al., 2015. The first microbial environment of infants born by C-section: the
operating room microbes. *Microbiome*, 3(1), p.59.

Sison-Mangus, M.P., Mushegian, A.A. & Ebert, D., 2015. Water fleas require
microbiota for survival, growth and reproduction. *The ISME journal*, 9, pp.59–
67.

Spor, A., Koren, O. & Ley, R., 2011. Unravelling the effects of the environment and
host genotype on the gut microbiome. *Nature Publishing Group*, 9(4), pp.279–
290.

Sullam, K.E. et al., 2017. The Combined Effect of Temperature and Host Clonal Line
on the Microbiota of a Planktonic Crustacean. *Microbial Ecology*.

Voss, J.D. et al., 2015. Pawnobiome: manipulation of the hologenome within one host
generation and beyond. *Frontiers in Microbiology*, 6(August), pp.1–5.

Zilber-Rosenberg, I. & Rosenberg, E., 2008. Role of microorganisms in the evolution
of animals and plants: the hologenome theory of evolution. *FEMS Microbiology*
Reviews, 32(5), pp.723–735.

 Figure legends

Figure 1: Experimental set-up and browsing behavior assay. A: the jars used in

the experiment had a bottom layer of fine loess and contained two animals each; the

animals were prevented from browsing on untreated sediments by a net placed 5

mm above the sediment surface (NET, right) or were allowed to browse on

autoclaved sediments (AUT) or untreated sediments (SED) (left). B: traces left by

one animal browsing on a sediment surface for 30 minutes and the same picture

after processing for quantification of the browsing behavior.

Figure 2: Browsing intensity of 12 *D. magna* clones (mean and SE). Browsing

intensity was defined as the Log_{10} of the area of the browsing traces left by

individual replicate animals browsing on a sediment surface for 30 minutes (see Fig.

1).

 Figure 3: Microbiota diversity (Shannon index) of *Daphnia* **clones under three**

different treatment conditions (AUT, NET and SED). AUT: Exposure to

autoclaved sediment; NET: prevented exposure to untreated sediment; SED:

exposure to untreated sediments. Error bars represent standard error of the mean.

Clones are arranged left-right by increasing average clone browsing intensity.

Figure 4: Average browsing intensity and average microbiota diversity in the

NET and SED treatments. A: average clone browsing intensity and average clone

microbiota diversity in the NET and SED treatments. Average browsing intensities

were calculated based on samples whose jar-mates passed the sequence quality

control (N=214, Table 1). B: average clone browsing intensity and the difference

between average Shannon diversity in the SED treatment and average diversity in

the NET treatment. Here, average browsing intensities were calculated based on the

complete set of samples (N=228, i.e. all assayed jar-mates). Error bars represent

standard error of the mean.

Figure 5: Within-group dispersion of community similarity. The median

distance to the centroid is lower in the NET treatment group than in the others

(Permutation test of multivariate dispersion p<0.0001, 999 permutations), meaning

that NET communities are less variable than AUT or SED microbiotas.

Figure 6: Similarity of bacterial community composition in the AUT and SED

treatments. A: first and second axis of a principal coordinates analysis (PCoA) of

bacterial community composition based on Hellinger-transformed Bray-Curtis

dissimilarities. B: first and second axis of a principal coordinates analysis (PCoA) of

bacterial community composition by *Daphnia* clone.

Figure 7: Analysis of sediment-derived bacteria. Proportion of sediment-derived
bacteria in the microbiota of animals from AUT, NET and SED treatments. Sediment-
derived bacteria were identified by comparing autoclaved and untreated sediment
samples (log2-fold increase of at least 8 in natural sediment compared to autoclaved
sediment). Error bars correspond to standard errors of the mean. Clones are
arranged left-right by increasing average clone browsing intensity.

Table 1. Names, number of individual replicates included in the microbiota analyses in the three treatments (AUT, NET and SED) and origin information of the 12 *Daphnia magna* clones used in this study. AUT: Exposure to autoclaved sediment; NET: prevented exposure to untreated sediment; SED: exposure to untreated sediments.

Clone ID	N (AUT)	N (NET)	N (SED)	Country	Latitude, N	Longitude E/W	Source	Description
BE-OHZ-T10	4	5	3	Belgium	50º50'00"N	4º39'00"E		
CZ-N1-1	8	8	8	Czech Rep.	48°46'31.14"N	16°43'24.70"E		
CZ-N2-6	6	7	6	Czech Rep.	48°46'31.14"N	16°43'24.70"E		
DE-K35-Mu10	4	3	4	Germany	48°12'23.93"N	11°42'34.98"E	*D. magna* Diversity panel	A geographically diverse collection of clones maintained asexually in the laboratory since 2012
DE-KA-F28	8	7	7	Germany	50°56'02"N	6°55'41"E		
ES-DO1-1	7	6	7	Spain	36°58'42.1"N	6°28'39.5"W		
TR-EG-1	5	7	8	Turkey	39°49' 25"N	32°49' 50"E		
BE-WE-G59	7	8	7	Belgium	51°04'04"N	3°46'25"E		
No-V-7	4	3	2	Norway	67°41'13.06"N	12°40'19.09E		

Clone ID				Description			Source	Description
IXF1	5	8	7	F1 clone			*D. magna* QTL panel	An intercross F2 recombinant panel maintained asexually in the laboratory since 2006/2007
F2-82	7	7	4	F2 clone				
F2-918	5	6	6	F2 clone				

Table 2. Results of analyses of variance of different alpha diversity indices. All treatments are included; clone NO-V-7 is excluded.

	Richness	Shannon	Inverse Simpson
Clone	F=0.40, df=10, **p=.944**	F=2.43, df=10, **P=.00997 ***	F=1.98, df=10, **p=.0383 ***
Treatment	F=4.91, df=2, **p=.00842 ***	F=12.21, df=2, **p<.0001 ***	F=13.25, df=2, **p<.0001 ***
Clone:Treatment	F=0.75, df=20, **p=.770**	F=0.78, df=20, **P=.734**	F=0.65, df=20, **P=.8124**

Table 3. Effect on Shannon index. NET and SED treatments only, all clones included. Linear mixed-effects model with treatment, clonal average browsing intensity and clonal average size as fixed effects and clone as random effect.

	numDF	denDf	F-value	p-value
(Intercept)	1	129	270.12	<.0001
Treatment	**1**	**129**	**12.26**	**0.0006 ***
Clone average behavior	1	9	0.275	0.613
Clone average size	1	9	2.568	0.144
Treatment:Clone average behavior	**1**	**129**	**4.677**	**0.0324 ***
Treatment:Clone average size	1	129	0.0980	0.755

Table 4. Effect on relative abundance of sediment-derived bacteria. All treatments and all animals included. Linear mixed-effects model with treatment and clonal average browsing intensity as fixed effects and clone as random effect.

	numDF	denDf	F-value	p-value
(Intercept)	1	198	53.999	<.0001
Treatment	**2**	**198**	**6.500**	**0.0018***
Clone average behavior	1	10	0.490	0.4998
Treatment:Clone average behavior	**2**	**198**	**4.185**	**0.0166***