Viral Fitness Across a Continuum from Lysis to Latency

Joshua S. Weitz,[1, 2, *] Guanlin Li,[2] Hayriye Gulbudak,[3] Michael H. Cortez,[4] and Rachel J. Whitaker[5, 6]

[1] *School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA, USA*
[2] *School of Physics, Georgia Institute of Technology, Atlanta, GA, USA*
[3] *Department of Mathematics, University of Louisiana at Lafayette, Lafayette, LA, USA*
[4] *Department of Mathematics and Statistics, Ecology Center, Utah State University Logan, UT, USA*
[5] *Department of Microbiology, University of Illinois Urbana-Champaign, Champaign, IL, USA*
[6] *Carl R. Woese Institute for Genomic Biology,*
University of Illinois Urbana-Champaign, Champaign, IL, USA
(Dated: June 25, 2018)

The prevailing paradigm in ecological studies of viruses and their microbial hosts is that the reproductive success of viruses depends on the proliferation of the "predator", i.e., the virus particle. Yet, viruses are obligate intracellular parasites, and the virus genome – the actual unit of selection – can persist and proliferate from one cell generation to the next without lysis or the production of new virus particles. Here, we propose a theoretical framework to quantify the 'fitness' of viruses using an epidemiological cell-centric metric that focuses on the proliferation of viral genomes inside cells instead of virus particles outside cells. This cell-centric metric enables direct comparison of viral strategies characterized by obligate killing of hosts (e.g., via lysis), persistence of viral genomes inside hosts (e.g., via lysogeny), and strategies along a continuum between these extremes (e.g., via chronic infections). As a result, we can identify environmental drivers, life history traits, and key feedbacks that govern variation in viral propagation in nonlinear population models. For example, we identify threshold conditions given relatively low densities of susceptible cells and relatively high growth rates of infected cells in which lysogenic and other chronic strategies have higher potential viral reproduction than lytic strategies. Altogether, the theoretical framework helps unify the study of eco-evolutionary drivers of viral strategies in natural environments.

I. INTRODUCTION

Viral infections begin with the physical interaction between a virus particle (the "virion") and the host cell. Infection dynamics within the cell often culminate in lysis, i.e., the active disruption of the integrity of the cell surface, leading to the death of the host cell and the release of virus particles [1, 2]. At population scales, virus-induced lysis can be a significant driver of microbial mortality, whether in the oceans, lakes, soil, extreme environments, or in plant and animal microbiomes [3–9]. As a result, studies of the ecological influence of viruses of microorganisms have, for the most part, emphasized the impact of the lytic mode of infection. However, the spread of viruses through microbial populations need not involve the immediate lysis of the infected cell.

Indeed, many viruses have alternative strategies. Temperate phage – like phage λ – can integrate their genomes with that of their bacterial hosts, such that the integrated viral DNA, i.e., the prophage, is replicated along with the infected cell, i.e., the lysogen [10]. Chronic viruses, like the filamentous phage M13, infect cells and persist episomally [11, 12], whereby the genome is replicated and then packaged into particles which are released extracellularly without necessarily inducing cell death [13, 14]. An analogous mode of chronic infection has been observed in archaeal virus-host systems [15]. These examples raise a critical question (see [16–18]): are temperate or chronic modes prevalent or rare in nature?

More than a decade ago, studies of marine, hydrothermal, and soil environments suggested that lysogeny could be more prevalent than assumed based on culture-based analysis of virus-microbe interactions [19–22]. This evidence has been augmented by recent studies identifying viral dark matter - including integrated and extrachromosal viral sequences - in microbial genomes [23–26]. Yet, despite increasing evidence of the relevance of persistent infections *in situ* the ecological study of phage has not integrated a common metric to compare the context-dependent fitness of lytic, temperate, and other chronic viral strategies.

A landmark theoretical study provides a setting off point for investigating the potential benefits of non-lytic strategies [27]. This study proposed that temperate phage could persist over the long term if prophage integration directly enhanced host fitness or enhanced resistance to infections by other lytic phage ("superinfection immunity"). The same study predicted that oscillations in population abundances could provide an ecological "niche" for temperate phage. In essence, if bacterial densities were too low to support the spread of lytic phage, then temperate phage already integrated into lysogens could persist until "conditions become favorable for the bacteria to proliferate" [27]. Yet this finding does not exclude the possibility that lytic strategies could outcompete temperate strategies – even if lysis at low densities leads to population collapse.

More recently, efforts to understand why viruses should be temperate have drawn upon the mathematical theory of portfolio balancing [28]. According to portfolio balancing theory, the temperate strategy enables viruses to expand rapidly during stable periods for hosts (via lysis) and mitigate risks of population collapse, particu-

*Corresponding author: jsweitz@gatech.edu; URL: http://ecotheory.biology.gatech.edu

larly during unfavorable periods for hosts (via lysogeny). Such arguments rely on generalized estimates of long-term growth rates without invoking the nonlinear feedback mechanisms underlying virus-microbe interactions. Moreover, a focus on *long-term* estimates of growth does not directly address whether killing a microbial host cell is the advantageous strategy for a virus at a given *moment* in time. As noted by [28], ecological models that incorporate feedback mechanisms of virus-host interactions are required to understand the viability of realized viral strategies.

Viruses have evolved many mechanisms to propagate with microbial hosts. Here, we use the word "strategies" to denote the *type* of mechanism underlying viral propagation. Comparing the relative fitness of viral strategies requires some means to quantify reproduction and survival across an entire viral life cycle, even if the molecular details, host strain, or virus strain differ. Here we propose to utilize an epidemiological framework to quantify viral fitness in which viral proliferation amongst microbial hosts is measured in terms of *infected cells* instead of virus particles. We show how doing so can help predict and explain a continuum of infection strategies observed in different environmental contexts.

II. ON HORIZONTAL AND VERTICAL TRANSMISSION

Viruses are obligate intracellular parasites. As such, virus-microbe dynamics can be re-cast in terms of the spread of an infectious disease through a microbial population. The risk for the spread of an infectious disease can be quantified in terms of the basic reproduction number, $\mathcal{R}_0$, "arguably the most important quantity in infectious disease epidemiology" [29]. In mathematical epidemiology, $\mathcal{R}_0$ is defined as the average number of new infected individuals caused by a single (typical) infected individual in an otherwise susceptible population [30]. Measuring $\mathcal{R}_0$ is the *de facto* standard for assessing pathogen invasion, *e.g.*, when $\mathcal{R}_0 > 1$ then a pathogen is expected to increase its relative abundance in a population [31, 32].

Thus far, estimates of $\mathcal{R}_0$ have had limited application to *in situ* studies of viruses of single-celled microorganisms, in part because counts of the number of *virus particles* have been used as a proxy for eco-evolutionary success (e.g., [33]). Yet, there are precedents for doing so. A 'phage reproductive number' has been proposed to quantify the proliferation of lytic viruses [34–36] and the ratio of new phage particles produced to those introduced has been used to identify invasision and equilibrium conditions for lytic viruses [37, 38]. However, the production of new virus particles does not, in and of itself, constitute a new infection. In addition, particle production is not the only way for viruses of microbes to proliferate at the population scale.

The virocell paradigm offers a path forward towards a unified notion of viral fitness [39, 40]. The paradigm centers on the idea that the "real living [viral] organism" [39] is an infected cell actively producing new virions, i.e., the "virocell". In contrast, conventional definitions of a virus refer to the physical properties of the virus particle, e.g., nucleic acids surrounded by a protein coat. As a consequence, it would seem logical to surmise that the viability of a viral strategy should be measured in terms of the number of new virocells produced.

Here we reconcile the virocell and conventional paradigms by adapting definitions of $\mathcal{R}_0$ to the ecological study of viruses of microorganisms. Specifically, we propose the following definition:

> $\mathcal{R}_0$: *the average number of new infected cells produced by a single (typical) infected cell and its progeny virions in an otherwise susceptible population.*

This definition counts viral reproduction in terms of infected cells, *i.e.*, with progeny viral genomes in them, rather than in terms of virus particles. For reasons that we will make clear in subsequent sections, this definition of $\mathcal{R}_0$ includes a critical asymmetry: we use infected cell, and not virion, production to measure viral spread, irrespective of whether we are evaluating purely lytic, latent, or chronic viral strategies. Indeed, the basic reproduction number has been used to compare the fitness of variants of temperate phage λ within eco-evolutionary studies [41, 42] (see Discussion for more details).

Characterizing the dynamics of virus genomes inside cells and virus particles outside of cells also enables comparisons amongst viruses with different life cycles. In particular, this definition accounts for infections caused by "vertical" transmission (i.e., from mother to daughter cell), those caused by "horizontal" transmission (i.e. from an infected cell to another susceptible cell in the population), and those caused by a combination of both routes (e.g., as in chronic viruses). Next, we explain how to calculate $\mathcal{R}_0$ and conditions for invasion within nonlinear models of virus and microbial population dynamics.

III. OBLIGATELY LYTIC VIRAL STRATEGIES – A BASELINE FOR COMPARISON

We begin our examination of obligately lytic strategies given a virion-centric perspective. Obligately lytic viruses infect and lyse their microbial hosts, thereby modifying the population densities of viruses and cells. Virus-host interactions can be represented via the following nonlinear differential equations (see Figure 1 for this and other model schematics):

$$\frac{dS}{dt} = \overbrace{bS\left(1 - S/K\right)}^{\text{logistic growth}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{dS}^{\text{cell washout}}$$

$$\frac{dV}{dt} = \overbrace{\beta\phi SV}^{\text{lysis}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{mV}^{\text{viral decay}}$$

(1)

FIG. 1: Population models for three types of viral strategies: obligately lytic, chronic viruses, and latent viruses. Boxes denote susceptible S, actively infected I, latently infected L, and virus particles, V. The top and bottom panels are modified from [38].

This system of equations represents changes in the density of virus particles, V, and susceptible microbial cells, S, using a resource-implicit model of bacterial growth. Model variants include terms representing nutrient uptake, fixed delays between infection and lysis, and other forms of cell mortality (e.g., due to grazing by eukaryotic predators) [38, 43, 44]. Given the model in Eq. (1), we are interested in determining the likelihood that a virus will spread when introduced to a susceptible cellular population.

The linearized virus population dynamics near the virus-free steady state are:

$$\frac{dV}{dt} = \left(\beta\phi S^* - \phi S^* - m\right)V \qquad (2)$$

where the steady-state density is $S^* = K(1 - d/b)$. This equation represents the potential exponential growth or decay of viruses. The growth constant is the term in the parentheses, $\beta\phi S^* - \phi S^* - m$. When this constant is greater than 0 then virus particles should increase in number, whereas when this constant is less than 0 then virus particles should decrease in number. In other words, viruses should spread when $\beta\phi S^* > \phi S^* + m$ or, alternatively, when $\mathcal{R}_{hor} > 1$ where

$$\mathcal{R}_{hor} \equiv \beta\left(\frac{\phi S^*}{\phi S^* + m}\right) \qquad (3)$$

is the (exclusively) horizontal contributions to the basic reproduction number. This inequality can be understood in two ways (see Figure 2).

First, consider a single virion. Virions successfully adsorb to susceptible hosts at a rate ϕS^*. In contrast, virions decay at a rate m. When two independent, random processes take place concurrently, the probability of one event - in this case adsorption - taking place before the other - in this case decay - is the ratio of one process relative to the sum of the rates of all processes. The factor $\phi S^*/(\phi S^* + m)$ denotes the probability that a virion is adsorbed before it decays. The present model assumes that adsorption implies successful infection and lysis. Hence, this probability must be multiplied by the burst size β, i.e., the number of new virions released, to yield the average number of new infectious virions produced by a single virion in a susceptible host population. This product is equal to the basic reproduction number, $\mathcal{R}_{hor}$. When $\mathcal{R}_{hor}$ exceeds 1 then a single virion produces, on average, more than one virion, of which each in turn produces, on average, more than one virion and so on. This process leads to exponential proliferation of virus particles, at least initially. As is evident, the spread of an obligately lytic virus depends on its life history traits and the ecological conditions (see Figure 3).

Second, we can revisit this same calculation beginning with an assumption that there is a single infected cell in an otherwise susceptible population. In that event, the infected cell produces β virions, of which only a fraction $\phi S^*/(\phi S^* + m)$ are adsorbed before they decay. The product represents the number of newly infected cells produced by a single infected cell in an otherwise susceptible population. The product is the same, but in this alternative approach we have counted proliferation in terms of a viral life cycle that starts and ends inside cells, requiring that contributions "complete the cycle". Figure 3 shows how viral proliferation varies with life history traits (in this case, the burst size) and the ecological context (in this case, the initial cell density). As is apparent, there is a threshold between regimes of viral extinction and proliferation corresponding to the transition of $\mathcal{R}_0$ from below to above one.

Thus far, we have not considered the explicit population dynamics of infected cells. In the Appendix we show that including an explicitly modeled infectious cell state leads to the same qualitative result. The only change is that the horizontal spread includes another factor: the probability that an infected cell releases virions before it dies or is washed out of the system by some other means. If η is the reciprocal of the average latent period and d' is the loss rate of infected cells, then only a fraction $\eta/(\eta + d')$ of infected cells will release virions before being washed out of the system. As such, the corrected basic reproduction number for obligately lytic viruses is:

$$\mathcal{R}_{hor} = \beta\left(\frac{\phi S^*}{\phi S^* + m}\right)\left(\frac{\eta}{\eta + d'}\right) \qquad (4)$$

Although both interpretations - the virion-centric and the cell-centric - lead to equivalent estimates of $\mathcal{R}_0$ for

FIG. 2: Schematic of cell-centric counting of the reproduction of obligately lytic viruses. Here, the mother virus generates dozens, if not hundreds of virions, most of these decay or are otherwise removed from the environment. Only three virions infect cells, these are progeny viruses, aka new mothers. Hence, the horizontal $\mathcal{R}_0$ of this virus is 3.

FIG. 3: Virus reproduction as a function of burst size and susceptible cell density. The contours denote the $\log_{10}$ of $\mathcal{R}_0$, as measured using Eq. (3), given variation in burst size, β, and susceptible cell density, S. Viruses invade when $\mathcal{R}_{hor} > 1$ or, equivalently when $\log_{10} \mathcal{R}_{hor} > 0$. Contours denote combinations of (β, S^*) of equivalent $\mathcal{R}_{hor}$. Additional parameters that affect viral reproduction are $\phi = 6.7 \times 10^{-10}$ ml/hr and $m = 1/24$ hr^{-1}.

obligately lytic viruses, we will use the cell-centric definition to unify comparisons across a spectrum of viral strategies.

IV. LATENT VIRAL STRATEGIES

In this section we consider the dynamics of latent viral strategies, such as temperate phage, in which proliferation may be either horizontal or vertical (but not both simultaneously). We model the dynamics of latent viruses using the following set of nonlinear differential equations:

$$\frac{dS}{dt} = \overbrace{bS\left(1 - N/K\right)}^{\text{logistic growth}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{dS}^{\text{cell death}}$$

$$\frac{dL}{dt} = \overbrace{qb'L\left(1 - N/K\right)}^{\text{lysogen growth}} + \overbrace{\phi SV}^{\text{infection}} - \overbrace{p\eta L}^{\text{lysis}} - \overbrace{d'L}^{\text{cell death}} \quad (5)$$

$$\frac{dV}{dt} = \overbrace{\beta p\eta L}^{\text{lysis}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{mV}^{\text{viral decay}}$$

This system of equations represents changes in the density of virus particles, V, lysogens, L, and susceptible microbial cells, S, in which the total density of cells is denoted as $N = S + L$. In this formulation, the relative rate of lysogenic growth and cellular lysis is controlled by the scaling factors q and p. When $q = 1$ and $p = 0$ then all infections are strictly latent and only lead to lysogenic growth. In contrast, when $q = 0$ and $p = 1$ then all infections are strictly lytic and only lead to cellular lysis. This is a variant of a nutrient-explicit formulation considered as part of an analysis of the tradeoffs underlying lysis and lysogeny for marine viruses [45]. Note that this model includes only a single infected state for cells; analysis of a related model, including detailed processes of integration and induction, will be the subject of follow-up work.

Using Eq. (5), we first consider the case $p = 0$ and $q = 1$ to focus on the vertical pathway. In the vertical pathway, virus genomes exclusively integrate with host cell genomes which can then be passed on to daughter cells. We use the cell-centric interpretation as before, and consider infection dynamics given a single lysogen in an otherwise susceptible population with no virus particles:

$$\frac{dL}{dt} = \left(b'\left(1 - \frac{S^*}{K}\right) - d'\right)L \quad (6)$$

This exponential growth equation predicts that lysogens will spread in abundance as long as $\left(b'\left(1 - \frac{S^*}{K}\right) - d'\right) > 0$. We can rewrite this condition for proliferation as:

$$\mathcal{R}_{ver} = \frac{b'\left(1 - \frac{S^*}{K}\right)}{d'} > 1. \quad (7)$$

Here, the subscript denotes the fact that $\mathcal{R}_0$ is entirely derived from *vertical* transmission of viral genomes among lysogens.

The basic reproduction number also has a mechanistic interpretation. The term $b'(1 - S^*/K)$ represents the birth rate of lysogens, which decreases with increasing number of cells - whether susceptibles or lysogens. Given that d' is the death rate of lysogens, the term $1/d'$ denotes the average lifespan of an individual lysogen. Therefore, this reproduction number is equal to the average number of newly infectious cells produced in the lifetime of the original infection (see Figure 4). If this number is greater than one, then a single lysogen will beget more than one

FIG. 4: Schematic of cell-centric counting of the reproduction of latent viruses, e.g., temperate phage. The mother virus could, in principle, lyse the cell or integrate its genome with that of the host. Here, the mother virus integrates and forms a lysogen. The lysogen divides three times before it is removed, thereby producing three daughter cells with virus genomes. The vertical $\mathcal{R}_0$ of this virus is 3.

FIG. 5: Basic reproduction number of temperate viruses as a function of susceptible cell density. The increasing (red) line denotes the horizontal $\mathcal{R}_0$ if temperate phage infect then always lyse cells. The decreasing (blue) line denotes the vertical $\mathcal{R}_0$ if temperate viruses always integrate with their hosts. Relevant parameters are $\beta = 50$, $\phi = 6.7 \times 10^{-10}$ ml/hr, $K = 7.5 \times 10^7$ ml^{-1}, and $b' = 0.32$, 0.54 and 1 hr^{-1} as well as $d' = 0.75$, 0.44, and 0.24 hr^{-1} for the three lysogeny curves from bottom to top respectively.

lysogen, on average, and those lysogens will do the same, and so on.

Measuring reproduction in this way also enables a mechanistic interpretation to the value of vertical transmission without invoking environmental fluctuations or other long-term measures of fitness. As is evident, lysogens reproduce more frequently when they are subject to less competition with hosts, i.e., when S^* is small relative to K. Given the value of S^* in the particular ecological model of Eq. (5), the basic reproduction number can be written as $\mathcal{R}_{ver} = (b'/d') \big/ (b/d)$. Hence, if lysogens

have more advantageous life history traits than do susceptible cells (as measured by a higher birth to death rate ratio, i.e., b'/d' larger than b/d), then viruses can spread exclusively via vertical transmission. This benefit of lysogeny applies in the immediate term and constitutes direct support for how a lysogen that benefits its host can also benefit the virus. However, if lysogeny comes with a cost (i.e., b'/d' lower than b/d), then vertical transmission alone will not be enough for $\mathcal{R}_{ver} > 1$. Note that $\mathcal{R}_{ver}$ is a monotonically decreasing function of S^*, such that increased abundances – all things being equal – diminishes the advantage for vertical transmission.

To consider horizontal transmission, consider the case where $p = 1$ and $q = 0$. In that case, analysis of the full model in Eq. (5) reduces to that of the obligately lytic virus already presented in Eq. (A5). This raises the question: does a strictly lytic or strictly lysogenic strategy have a higher basic reproduction number? Recall that the horizontal $\mathcal{R}_0$ is an *increasing* function of susceptible cell density, i.e., when there are more hosts then the value of horizontal transmission increases. The value of $\mathcal{R}_{hor}$ and $\mathcal{R}_{ver}$ cross at a critical value, S_c, which satisfies

$$\frac{b'\left(1 - \frac{S_c}{K}\right)}{d'} = \frac{\beta \phi S_c}{\phi S_c + m} \tag{8}$$

For $S > S_c$, then $p = 1$ and $q = 0$ has the higher basic reproduction number, (i.e., horizontal transmission is favored) whereas for $S < S_c$, then $p = 0$ and $q = 1$ has the higher basic reproduction number, (i.e., vertical transmission is favored). Extending prior analysis, we identify threshold conditions separating out when lysis should be favored at high density vs. when lysogeny should be favored at low density (see Figure 5). The use of a cell-centric metric makes it evident that vertical transmission can be evolutionarily advantageous given low densities of permissive hosts without invoking group selection or long-term fitness (see [46]).

V. CHRONIC VIRAL STRATEGIES

Next, we consider the dynamics of "chronic" virus strategies, or what have been termed "persister" or "producer" strains in other contexts. The dynamics of viruses, V, chronically infected cells, I, and susceptible microbial cells, S, can be represented via the following nonlinear differential equations:

$$\frac{\mathrm{d}S}{\mathrm{d}t} = \overbrace{bS\left(1 - N/K\right)}^{\text{logistic growth}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{dS}^{\text{cell death}}$$

$$\frac{\mathrm{d}I}{\mathrm{d}t} = \overbrace{b'I\left(1 - N/K\right)}^{\text{logistic growth}} + \overbrace{\phi SV}^{\text{infection}} - \overbrace{d'I}^{\text{cell death}}$$

$$\frac{\mathrm{d}V}{\mathrm{d}t} = \overbrace{\alpha I}^{\text{virion production}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{mV}^{\text{viral decay}}$$

(9)

in which the total number of cells is denoted as $N = S + I$. Although it can be remapped to the latency model, this

system of equations represents distinct mechanistic processes, including establishment of a chronically infected cell and release of virions from chronically infected cells without lysis at a per-capita rate α. As such, we expect that both vertical and horizontal transmission can take place concurrently.

As before, consider a newly infected cell in an environment in which all other cells are susceptible and there are no additional virus particles. The chronic cell will remain viable for an average duration of $1/d'$. In that time, the chronic cell will produce new virions at a rate α, of which only $\phi S^*/(\phi S^* + m)$ will survive to enter another cell. This is the horizontal component of reproduction for the chronic cell. Concurrently in that time, the chronic cell will divide at a rate $b'(1 - N/K)$. This is the vertical component of reproduction for the chronic cell. Hence a chronic virus will spread at the population scale, on average, as long as

$$\mathcal{R}_{chron} \equiv \overbrace{\frac{\alpha}{d'}\left(\frac{\phi S^*}{\phi S^* + m}\right)}^{\text{horizontal}} + \overbrace{\frac{b'(1 - S^*/K)}{d'}}^{\text{vertical}} > 1. \quad (10)$$

This decomposition of reproduction into horizontal and vertical components (see Figure 6) enables simple and interpretable calculations (see Appendix for the next-generation matrix method and derivation).

This analysis shows how the spread of chronic viruses depends on both infected cell traits and virion-associated traits. As a consequence, it would suggest that chronic viruses should evolve adaptations to improve the sum of horizontal and vertical reproduction. Without trade-offs, this would lead to chronic viruses with arbitrarily high virion release rates and arbitrarily low cell death rates. Yet, there will likely be trade-offs. For example, increasing the virion production rate, α, may improve horizontal reproduction, but if doing so increases cell death, d', then the overall change in $\mathcal{R}_{chron}$ may be negative. As a result, it is possible that chronic viruses could have the largest reproduction number in an intermediate density regime (see example in Figure 7). Understanding the pleiotropic effects of changes to chronic virus genotypes may provide one route to characterizing the evolution of viral strategies in which both horizontal and vertical transmission rates operate concurrently [47].

VI. DISCUSSION

We have proposed a unified theoretical framework to measure the spread of viral strategies across a continuum from lysis to chronic to latent infections. By defining viral reproduction in terms of infected cells, we are able to directly compare the spread of obligately lytic viruses, latent viruses, and chronic viruses in the context of nonlinear population models (see Figure 1). The invasibility of a newly introduced virus is measured in terms of the basic reproduction number, specifically adapted

FIG. 6: Schematic of cell-centric counting of the basic reproduction number of chronic viruses. In this example, the mother virus divides once, yielding a vertical $\mathcal{R}_0$ of 1. The mother virus chronically produces multiple virions of which 2 infect new cells, yielding a horizontal $\mathcal{R}_0$ of 2. The total is 3, i.e., the sum of horizontal and vertical components.

FIG. 7: Viral strategies with the highest $\mathcal{R}_0$ vary with susceptible host density, including exclusively vertical (bold blue, left), mixed (bold green, middle), and horizontal (bold red, right) modes of transmission. Relevant parameters are (i) for obligately lytic viruses (red): $\beta = 100$, $\phi = 6.7 \times 10^{-10}$ ml/hr, and $m = 0.13$ hr^{-1}; (ii) for chronic viruses (green): $b' = 0.68$ hr^{-1}, $d' = 0.63$ hr^{-1}, $\alpha = 20$ hr^{-1}, $\phi = 3.4 \times 10^{-10}$ ml/hr, and $m = 0.04$ hr^{-1}; (iii) for temperate viruses, given vertical transmission (blue) $b' = 0.54$ hr^{-1}, $d' = 0.44$ hr^{-1}, where $K = 7, 5 \times 10^7$ ml^{-1} in all three scenarios given variation in S^*.

to the life cycle of viral infections – in which new cellular infections can arise through horizontal and vertical transmission.

At its core, the theoretical framework re-envisions life history theory for viruses that infect microorganisms. In our calculations, a focal virus genome inside a cell can be thought of as a "mother virus". These mother viruses may lyse cells and produce "juvenile" offspring, i.e., virus particles. When a virion successfully infects a susceptible host this new infection becomes, once again, a

mother virus. This is an example of horizontal transmission. For latent and chronic viruses, the viral genome inside an infected cell may be passed on to both cells upon division. This division is equivalent to direct reproduction of a mother virus, bypassing the juvenile state. This is an example of vertical transmission. Combinations of these two scenarios (as shown in Figures 2, 4, and 6) are precisely those that emerge in applying next-generation matrix theory for calculating the basic reproduction number of viral strategies (see the Appendix).

The critical invasion fitness of a virus strategy, as calculated in terms of R_0, depends on life history traits as well as susceptible cell density. Obligately lytic viruses have increasing values of R_0 in populations with larger numbers of susceptible hosts. This trend is consistent with experimental findings that fitness of virulent phage λcI857 declines with decreasing susceptible cell density [41]. In contrast, latent viruses have decreasing values of R_0 in populations with larger numbers of susceptible hosts, arising from increased niche competition between a rare lysogenized cell and resident cells. Chronic viruses benefit from both transmission modes, although there can be trade-offs involving allocation toward horizontal and vertical transmission. As a consequence, latent, chronic, and lytic strategies can have higher potential reproductive success at low, intermediate, and high susceptible cell densities, respectively (see Figure 7). This result may help inform ongoing debates regarding the environmental conditions favoring lysogeny in marine systems [16–18, 48]. In our models, strictly vertically transmitted viruses may have a R_{ver} above 1 if the ratio of infected cell growth and death rates exceed that of susceptible hosts. This suggests a rationale for the evolution of viral traits that directly benefit host competitive fitness, e.g., toxin production and antibiotic resistance.

The present approach to measuring viral fitness focuses on a particular short-term invasion scenario: in which either a single virus particle or a single infected cell is added to an otherwise susceptible population. The framework is more general in the sense that it could apply to partially susceptible populations. However, it is important to note that comprehensive understanding of viral strategies requires analysis of long-term dynamics. This is particularly relevant given that evolutionary dynamics need not lead to the maximization of $\mathcal{R}_0$ (reviewed in [49]). Indeed, long-term fates are influenced by the Malthusian growth rate of viruses which we denote as r. For phage, population growth rates have been used as proxies for invasibility in multiple contexts [38, 50]. The basic reproduction number $\mathcal{R}_0$ and r are related, but they are not equivalent [51] – $\mathcal{R}_0$ measures the speed of viral proliferation in generations (i.e., at the individual level) whereas r measures the speed of viral proliferation in time (i.e., at the population level) [52]. The threshold condition $\mathcal{R}_0 > 1$ indicates whether the population growth rate r is positive, but does not predict changes in fitness given viral-host feedback. For example, virus proliferation depletes susceptible hosts, thereby decreas-

ing the 'effective' viral fitness – an outcome concordant with prior findings from mathematical models and eco-evolutionary dynamics of phage λ and *E. coli* [41].

Systematic analysis of the evolution of viral traits spanning lysis, chronic, and latent strategies in an ecological context is likely to draw upon a substantial body of work on the evolution of virulence (e.g., [53–60]), and in particular on the evolution of transmission mode [61]. For example, there can be tension and even conflicts at different scales of selection between viral genotypes that are effective at spreading within hosts but relatively ineffective at spreading between hosts [62–64]. In leveraging the insights of prior work, it is important to recall that virus-host dynamics unfold as part of complex ecosystems, whether in animal-associated or environmental microbiomes. As such, drivers of viral fitness will include spatial effects [65–68], temporal variation [69, 70], interactions with other strains [71–73], as well as feedback with other components of multi-trophic systems [74–76].

In moving forward, one immediate opportunity is to assess how viruses of microbes evolve virulence levels, or even strategy types, when co-infecting the same microbial population. For example, the cell-centric approach suggests new mechanisms of coexistence among viruses, e.g., viral-induced lysis may reduce niche competition between cells, increasing the benefits of vertical transmission, and enable invasion by latent/chronic viruses [77]. In addition, competition by multiple viruses within the same host cell could lead to emergent new strategies, e.g., as seen in a prisoner's dilemma in an RNA virus [78]. Finally, analyzing the evolution of temperate phage in the present theoretical framework may also shed light on plastic strategies in which infection outcome depends on the multiplicity of infection [42, 79–83]. How viruses sense cellular state, and perhaps even modify the state of cells through the release of small molecules [84, 85], remains an open question.

Altogether, the theory presented here provides an additional imperative to develop new measurement approaches to assess the entangled fates of viruses and cells. Measurements of the fitness of viruses with latent and chronic strategies should prioritize estimates of the life history traits of infected cells. Screening for viral genomes and their expression inside cells – whether integrated or persisting episomally – may reveal benefits of viral strategies that have thus far remained hidden when utilizing lysis-based assays or virion counts. By combining new measurements and theory, we hope that the present framework provides new opportunities to explore how viruses transform the fates of populations, communities, and ecosystems.

Acknowledgements

JSW thanks Jay Lennon, Maureen O'Malley, and the participants of a 2017 CIFAR-GBMF workshop on 'A Continuum of Persistence' for discussions critical to the development of ideas in this paper. The authors thank Sam Brown, Jonathan Dushoff, Santi-

ago Elena, Matthew Sullivan, and Steven Wilhelm for comments and feedback on the manuscript. This work was supported by a grant from the Simons Foundation (SCOPE Award ID 329108, J.S.W.), an Allen Distinguished Investigator Grant (to R.J.W.), and a NSF Dimensions of Biodiversity grant 1342876 (to J.S.W. and R.J.W.).

1 Calendar, R. L. (ed.) *The Bacteriophages* (Oxford University Press USA, 2005).

2 Dimmock, N. J., Easton, A. J. & Leppard, K. N. *Introduction to Modern Virology* (Blackwell Publishing, Malden, MA, 2007), 5th edn.

3 Wommack, K. & Colwell, R. Virioplankton: Viruses in aquatic ecosystems. *Microbiology and Molecular Biology Reviews* **64**, 69–114 (2000).

4 Weinbauer, M. Ecology of prokaryotic viruses. *FEMS Microbiology Reviews* **28**, 127–181 (2004).

5 Suttle, C. A. Marine viruses - major players in the global ecosystem. *Nature Reviews Microbiologygy* **5**, 801–812 (2007).

6 Rohwer, F. & Thurber, R. Viruses manipulate the marine environment. *Nature* **459**, 207–212 (2009).

7 Weitz, J. S. & Wilhelm, S. W. Ocean viruses and their effects on microbial communities and biogeochemical cycles. *F1000 Biology Reports* **4**, 17 (2012).

8 Breitbart, M. Marine viruses: truth or dare. *Annual Review of Marine Science* **4**, 425–48 (2012).

9 Brum, J. R. & Sullivan, M. B. Rising to the challenge: accelerated pace of discovery transforms marine virology. *Nat Rev Micro* **13**, 147–159 (2015).

10 Ptashne, M. *A Genetic Switch: Phage Lambda Revisited* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 2004), 3rd edn.

11 Salivar, W. O., Tzagoloff, H. & Pratt, D. Some physical-chemical and biological properties of the rod-shaped coliphage m13. *Virology* **24**, 359 – 371 (1964).

12 Jaenisch, R., Jacob, E. & Hofschneider, P. H. Replication of the small coliphage m13: evidence for long-living m13 specific messenger rna. *Nature* (1970).

13 Smeal, S. W., Schmitt, M. A., Pereira, R. R., Prasad, A. & Fisk, J. D. Simulation of the m13 life cycle i: Assembly of a genetically-structured deterministic chemical kinetic simulation. *Virology* **500**, 259 – 274 (2017).

14 Smeal, S. W., Schmitt, M. A., Pereira, R. R., Prasad, A. & Fisk, J. D. Simulation of the m13 life cycle ii: Investigation of the control mechanisms of m13 infection and establishment of the carrier state. *Virology* **500**, 275 – 284 (2017).

15 Snyder, J. C., Bolduc, B. & Young, M. J. 40 years of archaeal virology: Expanding viral diversity. *Virology* **479-480**, 369 – 378 (2015). 60th Anniversary Issue.

16 Knowles, B. *et al.* Lytic to temperate switching of viral communities. *Nature* **531**, 466–470 (2016).

17 Weitz, J. S., Beckett, S. J., Brum, J. R., Cael, B. B. & Dushoff, J. Lysis, lysogeny and virus–microbe ratios. *Nature* **549**, E1–E3 (2017).

18 Knowles, B. & Rohwer, F. Knowles & rohwer reply. *Nature* **549**, E3–E4 (2017).

19 Williamson, S. J., Houchin, L. A., McDaniel, L. & Paul, J. H. Seasonal variation in lysogeny as depicted by prophage induction in tampa bay, florida. *Applied and Environmental Microbiology* **68**, 4307–4314 (2002). http://aem.asm.org/content/68/9/4307.full.pdf+html.

20 Williamson, K. E., Radosevich, M., Smith, D. W. & Wommack, K. E. Incidence of lysogeny within temperate and extreme soil environments. *Environmental Microbiology* **9**, 2563–2574 (2007).

21 Williamson, S. J. *et al.* The sorcerer ii global ocean sampling expedition: metagenomic characterization of viruses within aquatic microbial samples. *PloS one* **3**, e1456 (2008).

22 Paul, J. H. Prophages in marine bacteria: dangerous molecular time bombs or the key to survival in the seas? *The Isme Journal* **2**, 579–589 (2008).

23 Roux, S., Enault, F., Hurwitz, B. & Sullivan, M. Virsorter: mining viral signal from microbial genomic data. *PeerJ* **3**, e985 (2015).

24 Howard-Varona, C., Hargreaves, K. R., Abedon, S. T. & Sullivan, M. B. Lysogeny in nature: mechanisms, impact and ecology of temperate phages. *ISME J* **11**, 1511–1520 (2017).

25 Luo, E., Aylward, F. O., Mende, D. R. & DeLong, E. F. Bacteriophage distributions and temporal variability in the ocean's interior. *mBio* **8** (2017). http://mbio.asm.org/content/8/6/e01903-17.full.pdf+html.

26 Kim, M.-S. & Bae, J.-W. Lysogeny is prevalent and widely distributed in the murine gut microbiota. *The ISME Journal* (2018).

27 Stewart, F. M. & Levin, B. R. The population biology of bacterial viruses: why be temperate. *Theoretical Population Biology* **26**, 93–117 (1984).

28 Maslov, S. & Sneppen, K. Well-temperate phage: optimal bet-hedging against local environmental collapses. *Scientific Reports* **5**, 10523 (2015).

29 Diekmann, O., Heesterbeek, H. & Britton, T. *Mathematical Tools for Understanding Infectious Disease Dynamics* (Princeton University Press, Princeton, NJ, 2012).

30 Keeling, M. J. & Rohani, P. *Modelling Infectious Diseases* (Princeton University Press, Princeton, 2007).

31 Diekmann, O., Heesterbeek, J. & Metz, J. On the definition and the computation of the basic reproduction ratio r_0 in models for infectious diseases in heterogeneous populations. *J. Math. Biol.* **28**, 365–382 (1990).

32 Diekmann, O., Heesterbeek, J. A. P. & Roberts, M. G. The construction of next-generation matrices for compartmental epidemic models. *Journal of the Royal Society Interface* **7**, 873–885 (2010).

33 Güemes, A. G. C. *et al.* Viruses as winners in the game of life. *Annual Review of Virology* **3**, 197–214 (2016). PMID: 27741409, https://doi.org/10.1146/annurev-virology-100114-054952.

34 Jones, D. & Smith, H. Bacteriophage and bacteria in a flow reactor. *Bull. Math. Biol.* **73**, 2357–2383 (2011).

35 Smith, H. L. & Thieme, H. R. Persistence of bacteria and phages in a chemostat. *J. Math. Biol.* **64**, 951–979 (2012).

36 Krysiak-Baltyn, K., Martin, G. J. O., Stickland, A. D., Scales, P. J. & Gras, S. L. Computational models of populations of bacteria and lytic phage. *Critical Reviews in Microbiology* **42**, 942–968 (2016). PMID: 26828960, https://doi.org/10.3109/1040841X.2015.1114466.

37 Beretta, E. & Kuang, Y. Modeling and analysis of a marine bacteriophage infection. *Mathematical Biosciences* **149**, 57–76 (1998).

38 Weitz, J. S. *Quantitative Viral Ecology: Dynamics of Viruses and Their Microbial Hosts* (Princeton University Press, Princeton, NJ, 2015).

39 Forterre, P. Manipulation of cellular syntheses and the nature of viruses: The virocell concept. *Comptes Rendus Chimie* **14**, 392 – 399 (2011). De la chimie de synthse la biologie de synthse.

40 Forterre, P. The virocell concept and environmental microbiology. *ISME J* **7**, 233–236 (2013).

41 Berngruber, T. W., Froissart, R., Choisy, M. & Gandon, S. Evolution of virulence in emerging epidemics. *PLoS pathogens* **9**, e1003209 (2013).

42 Gandon, S. Why be temperate: lessons from bacteriophage λ. *Trends in microbiology* **24**, 356–365 (2016).

43 Campbell, A. Conditions for the existence of bacteriophage.

Evolution **15**, 153–165 (1961).

Levin, B. R., Stewart, F. M. & Chao, L. Resource-limited growth, competition, and predation: a model and experimental studies with bacteria and bacteriophage. *American Naturalist* **111**, 3–24 (1977).

Record, N. R., Talmy, D. & Vage, S. Quantifying tradeoffs for marine viruses. *Frontiers in Marine Science* **3**, 251 (2016).

Anderson, R. M. & May, R. M. The population dynamics of microparasites and their invertebrate hosts. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **291**, 451–524 (1981).

Goldhill, D. H. & Turner, P. E. The evolution of life history trade-offs in viruses. *Current opinion in virology* **8**, 79–84 (2014).

Thingstad, T. F. & Bratbak, G. Microbial oceanography: Viral strategies at sea. *Nature* **531**, 454–455 (2016).

Lion, S. & Metz, J. A. Beyond r0 maximisation: On pathogen evolution and environmental dimensions. *Trends in Ecology and Evolution* **33**, 458 – 473 (2018).

Jensen, M. A., Faruque, S. M., Mekalanos, J. J. & Levin, B. R. Modeling the role of bacteriophage in the control of cholera outbreaks. *Proceedings of the National Academy of Sciences, USA* **103**, 4652–4657 (2006).

Wallinga, J. & Lipsitch, M. How generation intervals shape the relationship between growth rates and reproductive numbers. *Proceedings of the Royal Society B: Biological Sciences* **274**, 599–604 (2007).

Park, S. W., Champredon, D., Weitz, J. & Dushoff, J. Exploring how generation intervals link strength and speed of epidemics. *bioRxiv* (2018). https://www.biorxiv.org/content/early/2018/05/02/312397.full.pdf

Ewald, P. W. *Evolution of Infectious Disease* (Oxford University Press, Oxford, UK, 1994).

Lenski, R. E. & May, R. M. The evolution of virulence in parasites and pathogens: reconciliation between two competing hypotheses. *Journal of theoretical biology* **169**, 253–265 (1994).

Lipsitch, M., Siller, S. & Nowak, M. A. The evolution of virulence in pathogens with vertical and horizontal transmission. *Evolution* **50**, 1729–1741 (1996).

Frank, S. A. Models of parasite virulence. *The Quarterly review of biology* **71**, 37–78 (1996).

Turner, P. E., Cooper, V. S. & Lenski, R. E. Tradeoff between horizontal and vertical modes of transmission in bacterial plasmids. *Evolution* **52**, 315–329 (1998).

Brown, S. P., Hochberg, M. E. & Grenfell, B. T. Does multiple infection select for raised virulence? *Trends in microbiology* **10**, 401–405 (2002).

Ebert, D. & Bull, J. J. Challenging the trade-off model for the evolution of virulence: is virulence management feasible? *Trends in microbiology* **11**, 15–20 (2003).

Alizon, S., Hurford, A., Mideo, N. & Van Baalen, M. Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future. *Journal of evolutionary biology* **22**, 245–259 (2009).

Antonovics, J. *et al.* The evolution of transmission mode. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **372**, 20160083 (2017).

Gilchrist, M. A. & Coombs, D. Evolution of virulence: interdependence, constraints, and selection using nested models. *Theoretical Population Biology* **69**, 145–153 (2006).

Coombs, D., Gilchrist, M. A. & Ball, C. L. Evaluating the importance of within-and between-host selection pressures on the evolution of chronic pathogens. *Theoretical population biology* **72**, 576–591 (2007).

Mideo, N., Alizon, S. & Day, T. Linking within-and between-host dynamics in the evolutionary epidemiology of infectious diseases. *Trends in Ecology & Evolution* **23**, 511–517 (2008).

Kerr, B., Neuhauser, C., Bohannan, B. J. M. & Dean, A. M. Local migration promotes competitive restraint in a host–pathogen "tragedy of the commons". *Nature* **442**, 75–78 (2006).

Heilmann, S., Sneppen, K. & Krishna, S. Coexistence of phage and bacteria on the boundary of self-organized refuges. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 12828–12833 (2012).

Ashby, B., Gupta, S. & Buckling, A. Spatial structure mitigates fitness costs in host-parasite coevolution. *The American Naturalist* **183**, E64–E74 (2014). PMID: 24561607, https://doi.org/10.1086/674826.

Berngruber, T. W., Lion, S. & Gandon, S. Spatial structure, transmission modes and the evolution of viral exploitation strategies. *PLOS Pathogens* **11**, 1–13 (2015).

Brum, J., Hurwitz, B., Schofield, O., Ducklow, H. & Sullivan, M. Seasonal time bombs: dominant temperate viruses affect southern ocean microbial dynamics. *ISME J* **10**, 437–449 (2016).

Aylward, F. O. *et al.* Diel cycling and long-term persistence of viruses in the ocean's euphotic zone. *Proceedings of the National Academy of Sciences* **114**, 11446–11451 (2017). http://www.pnas.org/content/114/43/11446.full.pdf.

Weitz, J. S. *et al.* Phage-bacteria infection networks. *Trends in Microbiology* **21**, 82–91 (2013).

Roux, S. *et al.* Ecogenomics and biogeochemical impacts of uncultivated globally abundant ocean viruses. *Nature* **537**, 689–693 (2016). http://biorxiv.org/content/early/2016/05/12/053090.full.pdf.

Paez-Espino, D. *et al.* Uncovering Earth's virome. *Nature* **536**, 425–430 (2016).

Thingstad, T. F. Elements of a theory for the mechanisms controlling abundance, diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems. *Limnology and Oceanography* **45**, 1320–1328 (2000).

Winter, C., Bouvier, T., Weinbauer, M. G. & Thingstad, T. F. Trade-offs between competition and defense specialists among unicellular planktonic organisms: the "killing the winner" hypothesis revisited. *Microbiology and Molecular Biology Reviews* **74**, 42–57 (2010).

Weitz, J. S. *et al.* A multitrophic model to quantify the effects of marine viruses on microbial food webs and ecosystem processes. *ISME J* **9**, 1352–1364 (2015).

Gulbudak, H. & Weitz, J. S. Heterogeneous virus strategies promote coexistence in virus-microbe systems. *preprint* (2018).

Turner, P. E. & Chao, L. Prisoner's dilemma in an RNA virus. *Nature* **398**, 441–443 (1999).

Kourilsky, P. Lysogenization by bacteriophage-lambda 1. Multiple infection and lysogenic response. *Molecular and General Genetics* **122**, 183–195 (1973).

St. Pierre, F. & Endy, D. Determination of cell fate selection during phage lambda infection. *Proc. Natl. Acad. Sci. USA* **105**, 20705–20710 (2008).

Weitz, J. S., Mileyko, Y., Joh, R. I. & Voit, E. O. Collective decision making in bacterial viruses. *Biophysical Journal* **95**, 2673–2680 (2008).

Avlund, M., Dodd, I. B., Semsey, S., Sneppen, K. & Krishna, S. Why do phage play dice? *Journal of virology* **83**, 11416–11420 (2009).

Zeng, L. *et al.* Decision making at a subcellular level determines the outcome of bacteriophage infection. *Cell* **141**, 682–691 (2010).

Erez, Z. *et al.* Communication between viruses guides lysis–lysogeny decisions. *Nature* **541**, 488–493 (2017).

Schatz, D. *et al.* Communication via extracellular vesicles enhances viral infection of a cosmopolitan alga. *Nature Microbiology* **2**, 1485–1492 (2017).

Appendix A: Next-generation matrix approach for calculating viral invasion

The next-generation matrix (NGM) approach can be used to calculate $\mathcal{R}_0$ in mathematical models of interactions between cells

and viruses. We follow the convention of Dieckmann and colleagues in analyzing the subset of the model including infected subclasses [32]. In the case of viruses of microbes, we denote those infected subclasses to include any population type that has an infectious viral genome, i.e., both infected cells and virus particles.

1. Obligately lytic interactions

The main text considers a model of interactions between obligately lytic viruses and cellular hosts. Here, we modify this model to consider the dynamics of susceptible cells, infected cells, and virus particles:

$$\frac{dS}{dt} = \overbrace{bS\left(1 - N/K\right)}^{\text{logistic growth}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{dS}^{\text{cell death}}$$

$$\frac{dI}{dt} = \overbrace{\phi SV}^{\text{infection}} - \overbrace{\eta I}^{\text{lysis}} - \overbrace{d'I}^{\text{cell death}} \tag{A1}$$

$$\frac{dV}{dt} = \overbrace{\beta \eta I}^{\text{lysis}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{mV}^{\text{viral decay}}$$

We linearize the dynamics around the virus-free equilibrium, $(S^*, 0, 0)$ where $S^* = K(1 - d/b)$, and focus on the infected subsystem of $X(t) = [I(t)\ V(t)]^\intercal$. The linearized infected subsystem dynamics can be written as $\dot{X} = (T + \Sigma)X$ where

$$T = \left[\begin{array}{cc} 0 & \phi S^* \\ 0 & 0 \end{array} \right] \tag{A2}$$

denote transmissions events (i.e., corresponding to epidemiological births) and

$$\Sigma = \left[\begin{array}{cc} -\eta - d' & 0 \\ \beta \eta & -\phi S^* - m \end{array} \right] \tag{A3}$$

denote transition events (i.e., corresponding to changes in the state of viral genomes, including loss of infections). Via the NGM theory, the basic reproduction number $\mathcal{R}_0$ corresponds to the largest eigenvalue of the matrix $-T\Sigma^{-1}$. The i, j matrix elements of Σ^{-1} correspond to the expected duration in state i of a viral genome that begins in state j. For this model,

$$-\Sigma^{-1} = \left[\begin{array}{cc} \frac{1}{\eta + d'} & 0 \\ \frac{\beta}{\phi S^* + m}\left(\frac{\eta}{\eta + d'}\right) & \frac{1}{\phi S^* + m} \end{array} \right] \tag{A4}$$

As a consequence, the basic reproduction number is:

$$\mathcal{R}_{hor} = \frac{\beta \phi S^*}{\phi S^* + m}\left(\frac{\eta}{\eta + d'}\right) \tag{A5}$$

2. Latent strategies

Consider the dynamics of susceptible cells, infected cells (which denote lysogens), and virus particles:

$$\frac{dS}{dt} = \overbrace{bS\left(1 - N/K\right)}^{\text{logistic growth}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{dS}^{\text{cell death}}$$

$$\frac{dL}{dt} = \overbrace{qb'I\left(1 - N/K\right)}^{\text{lysogen growth}} + \overbrace{\phi SV}^{\text{infection}} - \overbrace{p\eta L}^{\text{lysis}} - \overbrace{d'L}^{\text{cell death}} \tag{A6}$$

$$\frac{dV}{dt} = \overbrace{\beta p\eta L}^{\text{lysis}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{mV}^{\text{viral decay}}$$

where $N = S + I$ is the total cell population density. As before, we linearize the dynamics around the virus-free equilibrium, $(S^*, 0, 0)$

where $S^* = K(1 - d/r)$, and focus on the infected subsystem of $X(t) = [L(t)\ V(t)]^\intercal$. The linearized infected subsystem dynamics can be written as $\dot{X} = (T + \Sigma)X$ where

$$T = \left[\begin{array}{cc} qb'(1 - S^*/K) & \phi S^* \\ 0 & 0 \end{array} \right] \tag{A7}$$

denote transmission events and

$$\Sigma = \left[\begin{array}{cc} -p\eta - d' & 0 \\ \beta p\eta & -\phi S^* - m \end{array} \right] \tag{A8}$$

denote transition events. For this model,

$$-\Sigma^{-1} = \left[\begin{array}{cc} \frac{1}{d' + p\eta} & 0 \\ \left(\frac{\beta p\eta}{p\eta + d'}\right)\frac{1}{\phi S^* + m} & \frac{1}{\phi S^* + m} \end{array} \right] \tag{A9}$$

As a consequence, the basic reproduction number is:

$$\mathcal{R}_0 = \frac{qb'(1 - S^*/K)}{d' + p\eta} + \frac{\beta \phi S^*}{\phi S^* + m}\left(\frac{p\eta}{p\eta + d'}\right) \tag{A10}$$

3. Chronic strategies

Consider the model of interactions between chronic viruses and microbial hosts:

$$\frac{dS}{dt} = \overbrace{bS\left(1 - N/K\right)}^{\text{logistic growth}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{dS}^{\text{cell death}}$$

$$\frac{dI}{dt} = \overbrace{b'I\left(1 - N/K\right)}^{\text{logistic growth}} + \overbrace{\phi SV}^{\text{infection}} - \overbrace{d'I}^{\text{cell death}} \tag{A11}$$

$$\frac{dV}{dt} = \overbrace{\alpha I}^{\text{virion production}} - \overbrace{\phi SV}^{\text{infection}} - \overbrace{mV}^{\text{viral decay}}$$

where $N = S + I$ is the total population density of cells. We recognize that the previous temperate model and the current chronic strategy model can be re-scaled to be equivalent. Nonetheless, the trade-off mechanisms are different and we note that in chronic infections, release of new virions need not be associated with lysis.

As before, we linearize the dynamics around the virus-free equilibrium, $(S^*, 0, 0)$ where $S^* = K(1 - d/b)$, and focus on the infected subsystem of $X(t) = [I(t)\ V(t)]^\intercal$. The linearized infected subsystem dynamics can be written as $\dot{X} = (T + \Sigma)X$ where

$$T = \left[\begin{array}{cc} b'(1 - S^*/K) & \phi S^* \\ 0 & 0 \end{array} \right] \tag{A12}$$

denote transmission events and

$$\Sigma = \left[\begin{array}{cc} -d' & 0 \\ \alpha & -\phi S^* - m \end{array} \right] \tag{A13}$$

denote transition events. For this model,

$$-\Sigma^{-1} = \left[\begin{array}{cc} \frac{1}{d'} & 0 \\ \left(\frac{\alpha}{d'}\right)\frac{1}{\phi S^* + m} & \frac{1}{\phi S^* + m} \end{array} \right] \tag{A14}$$

As a consequence, the basic reproduction number is:

$$\mathcal{R}_0 = \frac{b'(1 - S^*/K)}{d'} + \frac{\phi S^*}{\phi S^* + m}\left(\frac{\alpha}{d'}\right) \tag{A15}$$