Classification: BIOLOGICAL SCIENCES

Title: Autism sensory dysfunction in an evolutionarily
conserved system

Authors

Greta Vilidaite[a]*†, Anthony M. Norcia[b]*†, Ryan J. H. West[c], Christopher J. H.
Elliott[c], Francesca Pei[d], Alex R. Wade[a,c], & Daniel H. Baker[a]

Author affiliations

[a] Department of Psychology, University of York, York, YO10 5DD, UK.

[b] Department of Psychology, Stanford University, CA 94305, USA.

[c] Department of Biology, University of York, York, YO10 5DD, UK.

[d] Department of Psychiatry, Stanford University, CA 94305, USA.

† Authors contributed equally

Corresponding authors

* Correspondence and requests for materials should be addressed to
gv529@york.ac.uk or amnorcia@stanford.edu

Key words

Autism, animal model, sustained, *Drosophila*, sensory processing, visual
system

Abstract

Individuals with Autism Spectrum Disorder (ASD) report a host of sensory
symptoms, which suggests a fundamental, genetic (Miles, 2011) neural
signaling deficit in autistic brains (Rubenstein & Merzenich, 2003). However,
neither animal models nor previous theories explaining sensory symptoms
have been able to predict neurophysiological data in autistic humans. Here we
show a strikingly similar trajectory of visual development in a genetic
Drosophila Nhe3 model of autism and in autistic human participants. We
report a dissociation between first- and second-order electrophysiological
visual responses to steady-state stimulation in adults with ASD as well as a
large sample of neurotypical individuals with high numbers of autistic traits.
We also report a strikingly similar impairment in the adult fruit fly model of
ASD. We explain this as a selective signaling abnormality in the transient
response mechanisms in the visual system. In contrast to adults, autistic
children show decreases in both first- and second-order responses, which are
closely matched by the fruit fly model, suggesting a compensatory change in
processing occurs over the course of development. Our results provide the
first animal model of autism comprising a developmental sensory pathway
phenotype.

Significance statement

Autism Spectrum Disorder exhibits strong and widespread sensory symptoms
that have not yet been explained by previous research. Here we have
developed a novel *Drosophila* model of sensory deficits in autism that is highly
predictive of electrophysiological visual data in both autistic adults and
children. Both the animal model, and human data from three samples (total
N=154), point towards a deficit of a fundamental signaling mechanism in early
parts of the sensory system. This deficit shows signs of change during
development indicating a possible partial rescue of function at later stages of
life. Our findings can explain previous inconsistencies in research into visual
perception in autism. The *Drosophila* model can be used in future biomarker
and treatment development.

Introduction

Autistic individuals report a host of sensory symptoms including unusual sensory interests, overstimulation and hyper- and hyposensitivity to intense stimuli such as bright lights or loud noises (*1*, *2*). Several theories have been proposed to account for sensory abnormalities in autism – one of the most common proposing an imbalance in excitation/inhibition (E/I) (*3*). The E/I imbalance theory of autism has gained mixed but largely supportive evidence from neurochemical, neurobiological and genetic investigations (*4*, *5*), implicating GABA-ergic mechanism deficits (*6*). Alternatively, a vision-specific theory has linked visual processing abnormalities in ASD to a selective deficit in the magnocellular pathway (*7*, *8*). Such signaling-level impairments indicate a genetic cause of ASD sensory symptoms (*9*). However, neural measurements that can convincingly delineate these competing explanations are yet to be reported.

One attempt in mice to model a related developmental condition, Rett syndrome, found similar cross-species decreases in visual neural responses and poor visual acuity (*10*). However, Rett syndrome lacks the pervasive sensory symptoms characteristic of autism, thus it is difficult to link cellular and genetic abnormalities to ASD sensory deficits (*11*). As an alternative to rodents, *Drosophila* have provided successful models of human neurological disorders such as Parkinson's (*12*) and Alzheimer's (*13*). Fruit flies share 75% of human disease causing genes (*14*) and have a visual system exhibiting similar nonlinear neural properties, including a color- and luminance-selective module as well as a motion-selective module (*15*). The neural dynamics of these modules closely resemble those of the transient and sustained neural populations in humans (*16–18*). This provides an excellent framework for modeling low-level changes in sensory neuronal signaling (*18*) which may lie behind sensory abnormalities in autism.

In this study we aimed to develop an animal model of sensory abnormalities in human ASD and to investigate the mechanism underlying these impairments. We measured visual neural responses both in autistic humans and in *Drosophila* with a genetic mutation linked to autism. In humans, loss-of-function mutations in the gene *SLC9A9* have been linked to ASD (*19*). Here we used a *Drosophila* orthologue of *SLC9A9* – *Nhe3*. A homozygous P-element insertion loss-of-function mutants (*Nhe3*^{KG08307}) and *Nhe3* hemizygotes *(Nhe3*^{KG08307}/Df(2L)BSC187) were used to inhibit *Nhe3* function in fly. To assess the functionality of sustained and transient pathways in these species, we measured two steady-state electrophysiological responses: first harmonic (stimulation) and second harmonic frequencies, in both humans and fruit flies. The use of two *Nhe3* mutations in different genetic backgrounds

ruled out the possibility of other mutations influencing the flies' visual
responses. The dynamics of the sustained pathway manifest in the
electrophysiological stimulation frequency (first harmonic) response whereas
the dynamics of the transient pathway can be observed in the second
harmonic response (*17*, *20*). Furthermore, to investigate the progression of
ASD sensory abnormalities over the course of development, we also
measured visual responses at two stages of fruit fly maturation and acquired
similar responses from autistic children. Finally, as the ASD phenotype is
complex and non-binary, we validated our sensory model with a large sample
of neurotypical participants with high and low numbers of autistic traits.

Results

Increased sustained/transient response ratio in *Nhe3* fruit flies. Using a
steady-state visual evoked potential (ssVEP) paradigm (*12*) (see *Fig. 1*) we
measured fruit fly visual responses to flickering stimuli via an electrode on the
fly's eye. Wild type, eye-color matched flies (a cross between isogenic and
Canton-S) were used as controls (+). Flies from each genotype (n=12) were
tested at three days (when the flies are young) and at 14 days post eclosion
(older). First harmonic (12Hz) and second harmonic (24Hz) response
amplitudes were derived by fast Fourier transform (see *Methods*). Although
the first harmonic responses of mutant and wild-type flies were the same, the
second harmonic response was significantly reduced in the *Nhe3* mutants
(*Fig. 2a, 2b*).

Figure 1. Human and *Drosophila* steady-state electrophysiology methods.

To quantify this functional dissociation between the two response components we calculated a normalized ratio between first (1F) and second (2F) harmonics ($\frac{1F-2F}{1F+2F}$) and averaged over the highest contrast conditions (where the response rises above the noise floor, see *Methods*). This allowed us to measure the differences between sustained and transient responses whilst normalizing for overall responsiveness of the visual pathway. The ratio was

significantly higher in both mutant strains than in the controls (ANOVA, $F_{2,33}$ =
20.53, P < 0.0001, both paired contrasts P < 0.001; *Fig. 2c*). These data
suggest an impairment in the post-receptoral neural pathways (structures
downstream of the photoreceptors) of the older mutant flies.

Figure 2. Older ASD-mimic flies and autistic humans have visual deficits in the
transient pathway.

Interestingly, unlike the older flies, the young 3 day old flies showed a reduced
response at both frequencies (see *Fig. 3a,b*) relative to controls. Importantly,

there was no effect of genotype on the ratio between harmonics ($F_{2,33} = 1.38$,
$P = 0.27$; *Fig. 3c*). These results suggest a deficit in the sustained visual
module of young mutant flies, which most likely causes a cascading
impairment in the transient module. These differences between visual
responses at two stages of life suggest a change in visual processing over the
course of development.

Figure 3. Young ASD-mimic flies and autistic children have visual deficits in
the sustained pathway.

High autistic trait population show similar ssVEPs to *Nhe3* flies. To
assess the relevance of the *Nhe3* model to the human ASD phenotype we
used a comparable and equally sensitive steady-state EEG paradigm in
human participants. One hundred neurotypical participants with putative
autistic traits measured using the Autism Spectrum Quotient (AQ)
questionnaire (*21*) were tested with the ssVEP paradigm. Visual responses
were recorded from an occipital electrode (Oz, located at the back of the head
over the visual cortex) to grating stimuli flickered at 7Hz. Seven contrast
conditions (each repeated eight times) were presented in a randomized order.
First and second harmonic ssVEP responses were again derived via Fourier
analysis. The evoked response data were averaged separately over

participants split by their median AQ score: high (n = 53, AQ mean = 11.53,
SD = 3.68) and low (n = 47, AQ mean = 5.32, SD = 1.81) AQ (high AQ
implying many autistic traits). The first harmonic response was similar in both
groups whereas the second harmonic was notably reduced in the high AQ
group (*Fig. 2d, 2e*). A two-way ANOVA showed the interaction between group
and frequency to be significant ($F_{1,98}$ = 6.17, P = 0.015). The high AQ group
also had a significantly higher frequency ratio than the low AQ group (t_{98} =
2.86, P < 0.01, *Fig. 2f*). Moreover, a regression analysis showed that AQ
scores correlated with the frequency ratio, with high AQ scores being
predictive of higher ratios (R = 0.26 $F_{1,98}$ = 6.87, P = 0.01; see *Fig. 4*). This
result shows a relationship between the amplitude of the second harmonic
response and the severity of the subclinical ASD phenotype, however, this
effect cannot be directly generalized to clinical autism as the AQ is not
diagnostic of full-blown ASD.

Figure 4. Positive relationship between the number of autistic traits and
first/second harmonic ratio.

Autistic individuals show the same pattern of responses as *Nhe3* flies.
We assess this difference between harmonics in clinical ASD by testing 12
typical-IQ autistic adults (diagnosis confirmed with the Autism Diagnostic
Observation Schedule, Second Edition (ADOS-2), Lord et al., 2000) and 12
age- and gender-matched controls using the same human ssVEP paradigm.
The pattern of data again mimicked that of the previous adult datasets: there
was a significant interaction between group and frequency ($F_{1,22}$ = 5.85, P =

0.02; *Fig. 2g, 2h*). The ratio between harmonics was again significantly larger
in the ASD group than in the control group (t_{22} = 2.13, P = 0.04; *Fig. 2i*).

Young *Nhe3* fly responses predict autistic children's responses.
Considering the striking similarity between the adult human datasets and the
adult fruit fly model, it is reasonable to ask if similarities also exist between
younger humans and young Drosophila ASD models. Specifically, our fly
model predicts that the visual system of autistic children should show reduced
responses in both the first and second harmonics. To examine this, we
recorded from 13 autistic children (5 – 13 years old) and 17 neurotypical age-
and gender-ratio-matched controls using an ssVEP contrast-sweep paradigm.
Artifact rejection was employed to control for movement and blinking in both
groups. The stimulus in each sweep trial increased continuously in contrast
from 0% to 50% in logarithmic steps. Data were binned into 9 contrast levels
before being Fourier transformed to compute response amplitudes.

As predicted by the fly data, the ASD group showed impaired processing in
the sustained response (t_{28} = 2.07, P = 0.04; *Fig. 3d, 3e*). A two-way ANOVA
also revealed a significant group effect over both frequencies ($F_{1,28}$ = 4.23, P
= 0.049). Unlike adults, children exhibited no difference in frequency ratios
between the groups (t = 1.41, P = 0.17; *Fig. 3f*). Children showed impaired
processing in the sustained pathway as predicted by the fruit fly model.
However, the smaller effect of amplitude reduction observed in the fruit fly
second harmonic responses was present, but not statistically reliable in the
children (t_{28} = 1.26, P = 0.219). This may be due to the more variable nature
of human EEG data.

Discussion

We found a selective deficit in second order responses in autistic adults,
individuals with high levels of autistic traits and *Nhe3* fruit flies suggesting that
this impairment is specific to the autistic phenotype. Autistic children and
young *Nhe3* flies showed a more global deficit in visual processing at early
stages of life. The *Nhe3* fruit fly model of autism was predictive of these ASD
visual response deficits both in children and in adults, suggesting a
fundamental and pervasive change in visual impairment during development.
High AQ individuals showed consistent visual responses compared to
participants with diagnosed ASD suggesting common visual response
properties between samples. This was unsurprising as previous research has
found that AQ scores in the general population are highly correlated (R =
0.77) with sensory processing difficulties, as measured by the Glasgow

The intact first harmonic response in adult flies and humans indicates normal
functioning of mechanisms which give rise to the sustained response.
Conversely, the reduced second harmonic response suggests a deficit in the
transient dynamics of the visual system. In fly, the first harmonic has been
associated with sustained photoreceptor polarization and the second
harmonic with second-order lamina cells (*17*). In human, an association has
been made between simple cell and sustained responses to pattern onset and
complex cells and transient responses at both stimulus onset and offset (*24*).
Although simple cells exhibit some transient response properties as well (*24*,
25), the intact first harmonic suggests that the impairment is specific to human
complex cells that only generate even-order response components. This
early, cell-type-specific deficit may explain previous findings of atypical neural
dynamics of spatial frequency processing in ASD in the face of normal
sensitivity thresholds (*26*, *27*).

The differential impairment of sustained and transient modules observed in
our *Nhe3* model mimics the pathway abnormality in autistic adults. *Nhe3*
affects the exchange of sodium and hydrogen ions in cell membranes directly
affecting neural signaling (*19*, *28*). Differential expression of *Nhe3* and other
genes in ASD, which has been observed in other parts of the brain (*19*, *29*)
may extend to differential expression in color and motion modules in the
Drosophila visual system. As *Nhe3* (*SLC9A9* in humans) is only a single gene
in a multifaceted genetic etiology of autism, it is likely that the expression of
several genes in human autism affects simple and complex cell dynamics,
producing similar effects at the neural population level. Furthermore, such
abnormality in gene expression in other parts of autistic brains, as well as
environmental influences and gene-environment interactions, may give rise to
a wide range of cognitive and social impairments in childhood and adulthood.

Our data indicate little or no over-responsivity in the visual responses
predicted by the E/I theory, consistent with some previous studies (*30*, *31*).
However, it is possible that E/I imbalance in autism, stemming from GABA-
ergic mechanism deficits, affects different neuron types or processing
pathways in distinct ways and to different extents. Regardless, cell-type level
processing abnormalities may explain previous inconsistencies in sensory
deficits in ASD in which pathways were not differentiated (*32*). Furthermore,
the current results can provide an amended explanation to the magnocellular
(M pathway) dysfunction hypothesis. As it is difficult to isolate the M pathway

by changing stimulus properties (*20*), the paradigms previously used to
investigate magnocellular dysfunction in ASD may have been selectively
activating complex cells rather than the M pathway in particular (*33, 34*).

Developmentally, the observed change in the nature of the abnormality in
both species with increasing age is in accordance with previous findings
showing reduction or complete rescue of neuroanatomical abnormalities
present in early ASD childhood over the course of maturation (*35*). Previous
longitudinal research has also shown that symptom severity in individuals
diagnosed with ASD in childhood decreases over time (*36, 37*). McGovern &
Sigman (*37*) found that 48 adolescents, who were diagnosed with ASD as
children, showed marked improvement in social interaction,
repetitive/stereotyped behaviors and other symptoms, with two no longer
meeting criteria for ASD under ADI-R criteria, and four under ADOS criteria.
This might be explained by a change in neural processing during
development, which would likely affect both behavioral and sensory
outcomes.

One possible mechanism that would explain the present results is that the
neural signaling abnormalities (such as ion balance in the case of *Nhe3*),
change over time. In flies, reduced *Nhe3* expression may reduce the rate at
which sodium ions and protons are exchanged across the cell membrane. At
least in Mosquito, this exchanger is found in the gut, and Malpighian tubules
(the fly equivalent of the kidney) (*38*). Failure to properly regulate ionic
balance in young adult flies might affect the sodium concentration, or proton
levels in the body and brain, and affect the speed an intensity of action
potentials. Later in life, the normal balance may be restored. A similar
reduction in efficacy of *SLC9A9,* linked to ASD, may also be present and
explain the homology. In this respect, we note that another transporter, the
potassium/chloride exchanger, has been linked to epilepsy in young people:
with age the kcc/KCC2 eventually achieves a normal ionic balance and proper
inhibitory GABA signaling (*39*).

The *Nhe3* model may serve for further research into the development of ASD
in young brains as well as early biomarkers and treatments. Consistency
between the fly and human datasets at both ages indicates a breakdown of a
fundamental sensory mechanism comprising two components that have been
conserved over 500 million years of evolution. The conservation of the
phenotype and mechanisms from fly to human opens up the option to utilize
the unrivaled genetic tractability of the fly to dissect the molecular
mechanisms underpinning the disorder.

Methods

Drosophila stocks

Two *Drosophila melanogaster* genotypes were used as ASD models. The *Nhe3* loss-of-function P-element insertion (*Nhe3*[KG08307] homozygotes) mutation was homozygous P{SUPor-P}Nhe3[KG08307] (Bloomington Drosophila Stock Center (BDSC) 14715). The deficiency was Df(2L)BSC187 (BDSC 9672). To avoid second site mutations in the P-element stock, we used the hemizygote *Nhe3*[KG08307] /Df(2L)BSC187 as a second experimental genotype.

For our control cross we mated the lab stock of *Canton-S (CS)* flies with those with isogenic chromosomes 2C and 3J (*40*). All tested flies had dark red eyes. All genotypes were raised in glass bottles on yeast-cornmeal-agar-sucrose medium (10g agar, 39g cornmeal, 37g yeast, 93.75g sucrose per litre). They were kept at 25°C on a 12 hour light-dark cycle. Male flies were collected on CO_2 the day after eclosion and placed on Carpenter (1950) (*41*) medium in the same environmental conditions for either 3 days or 14 days. Flies were tested approximately between the 4th and 9th hour of the daylight cycle.

Drosophila electroretinography

Steady-state visual evoked potentials (SSVEPs) were obtained from the fruit flies (*12*, *17*). Flies were recorded in pairs in a dark room. They were placed in small pipette tips and secured in place with nail varnish. One glass saline-filled electrode was placed inside the proboscis of the fly and another on the surface of the eye. A blue (467nm wavelength) LED light (Prizmatix FC5-LED) with a Gaussian spectral profile (FWHM 34nm) was placed in front of the flies together with a diffuser screen and used for temporal contrast stimulation. Flies were dark adapted for at least two minutes and then tested for signal quality with six light flashes. Steady-state stimulation lasted 12 min and comprised seven contrast levels (0 – 69% in linear steps) each with five repetitions. The frequency of the light flicker was 12Hz. Each trial (contrast level repetition) was 11 s. The order of the contrast conditions was randomized. The stimulation and the recording from the fly was controlled by in-house MATLAB scripts (scripts can be found in https://github.com/wadelab/flyCode).

Adult EEG

One-hundred neurotypical adult participants (32 males, mean age 21.87, range 18 – 49, no reported diagnosis of ASD, reportedly normal or corrected

to normal vision) took part in the autism spectrum quotient (AQ) measurement
study. The AQ is an instrument used for quantifying autistic traits in the
neurotypical population and has been shown to have high face validity and
reliability in these populations (*21*). The AQ consists of 50 Likert scale items
and the scores in this study were calculated according to Baron-Cohen et al.
(*21*). Each participant completed the AQ questionnaire on a computer in the
laboratory.
367
For the autistic adult ssVEP study, 12 typical-IQ autistic participants and 12
gender- and age- matched controls (11 males, mean age 23.53, range 18 –
39, reportedly normal or corrected to normal vision) took part. ASD diagnosis
was confirmed with the Autism Diagnostic Observation Schedule, second
edition (ADOS-2). Although IQ was not explicitly measured in this study, all
adults had normal speech and a high level of independence (the majority
were university students). The absence of ASD diagnosis in the neurotypical
participants was also confirmed with ADOS-2 (none of the control participants
met criteria for ASD). All participants in the study gave informed consent and
were debriefed on the purpose of the study after the experiment. The
experiments were approved by the Department of Psychology Ethics
Committee at the University of York.
380
Steady-state VEPs were recorded using an ANT Neuro system with a 64-
channel Waveguard cap. EEG data were acquired at 1kHz and were recorded
using ASALab, with stimuli presented using MATLAB. The timing of the
recording and the stimulation was synchronized using 8-bit low-latency digital
triggers. All sessions were performed in a darkened room, testing lasted 45-
60min with approximately 20min set up time.
387
Stimuli were presented on a ViewPixx display (VPixx Technologies Inc.,
Quebec, Canada) with a mean luminance of 51cd/m^2 and a refresh rate of
120Hz. Stimuli were 0.5 cycle/deg sine-wave gratings enveloped by a raised
cosine envelope. Gratings subtended 3 degrees of visual angle and were tiled
in a 17x9 grid. The participants fixated on a circle in the middle of the screen
and performed a fixation task (two-interval-forced-choice contrast
discrimination) to maintain attention. All participants were able to perform the
task at above chance levels. There were seven contrast conditions for the
flickering gratings (0%, and 2 - 64% in logarithmic steps, where $C_\%$ =
$100(L_{max}-L_{min})/(L_{max}+L_{min})$, L is luminance) and eight repetitions. Stimuli
flickered on/off sinusoidally at 7Hz. Trials were presented in random order in
four testing blocks with short breaks in between. Each trial was 11 seconds
long and contained gratings of a random spatial orientation to avoid
orientation adaptation effects. These trials were intermixed with orthogonal
masking trials that are not presented as part of this study. Data were taken
from the occipital electrode Oz.

Child EEG

Thirteen children with a diagnosis of ASD and 20 neurotypical controls matched on gender ratio (10 and 12 males respectively) and average age (mean age 9.31 and 8.94 respectively, range 5 – 13) completed the study. Three of the neurotypical children were tested but excluded due to having autistic siblings (17 participants were included). All children were in mainstream local schools (if they were old enough) and did not have other (or any – in the case of the neurotypical group) reported history of serious medical, psychiatric, or neurological conditions.

Steady state EEG data were acquired with a 128-channel HydroCell Geodesic Sensor Net (Electrical Geodesics Inc.). Data were digitized at 432Hz and band-pass filtered from 0.3Hz to 50Hz and were recorded using NetStation 4.3 Software. Highly noisy data were excluded by removing repetitions with amplitudes that were four standard deviations away from the group mean (for each contrast level and harmonic individually). There were 10 repetitions in total, however, two autistic and one neurotypical child only completed 8 repetitions.

Increasing contrast sweep ssVEPs were used. Stimuli for this experiment were presented on an HP1320 CRT monitor with 800x600 pixel resolution, 72Hz refresh rate and mean luminance of 50cd/m^2. Stimuli were random binary noise patterns of two luminance levels that increased in contrast in 9 logarithmic steps (0% – 50%) of 1 second each. Each trial contained a prelude at the initial value of the sweep and a postlude at the final sweep value, lasting 12 seconds in total. Stimuli flickered at 5.12Hz. Data from the middle 9 seconds during the sweep were binned according to contrast steps. Methodological differences between the adult and child datasets were due to different conventions being used by the two laboratories in which data were collected.

Data analysis

A Fast Fourier transform (in MATLAB) was used to retrieve steady-state response amplitudes at the stimulation frequency (12Hz for fruit flies, 7Hz for adult participants and 5.12Hz for children) and at the second harmonic (24Hz, 14Hz and 10.24Hz respectively). Fourier transforms were applied to 10 s of each trial (first 1s discarded; total trial length was 11s) for the fruit fly and the adult participant datasets and to 1 second binned data for the children's dataset. Contrast response functions were obtained by coherently averaging the amplitudes over repetitions for each contrast level within a participant. Group/genotype scalar means over response amplitude (discarding phase angle) were then calculated for each contrast across participants/flies.

Two-way (harmonic x group) ANOVAs were performed on amplitudes at the
highest contrast level to investigate the interactions and group effects in all
human datasets where only two groups were compared. To identify at which
harmonic the autistic children showed a decreased response, two
independent samples t-tests were also conducted. One-way ANOVAs with
simple planned contrasts were conducted to assess the genotype differences
in fruit fly first and second harmonic responses separately as that aided the
interpretability of the results between the three genotypes.
456
To investigate the dissociation between first and second harmonic responses
a scaled ratio $\frac{1F-2F}{1F+2F}$ (where 1F is the first and 2F is the second harmonic) was
calculated for each participant/fly and each contrast condition. To increase the
power of statistical analyses and to decrease the type I error rate, the ratios
were then averaged over the contrast conditions that had first harmonic
amplitudes significantly above the baseline response (0% contrast condition).
For fruit flies this was six conditions (11.5 – 69%), for adult participants this
was four conditions (8 – 64%) and for children this was five conditions (8.5 –
50%). This procedure resulted in a single frequency-ratio index for each
participant/fly. One-way ANOVAs with simple planned contrasts (comparing
mutant genotypes with the control genotype) were conducted on the fly
frequency ratios for each age separately. Independent t-tests were used to
compare frequency ratios in all human datasets between groups. Additionally,
a linear regression was conducted on the adult AQ measurement dataset to
assess the predictive power of AQ scores on the ratios between frequencies.
All statistical tests were two-tailed.

Acknowledgements

This research was supported by the Experimental Psychology Society Study Visit Grant awarded to G.V.

We would like to thank E. Cole for second-coding ADOS2 assessments; N. Gutmanis, S. Harris, R.E. Kitching, H.A. Melton, R. Norman, C.J.T. Scott, C. Simpson, A.K. Smith, A. Stockton and K. Wailes-Newson for contributing to EEG data collection. A.M.N. and F.P. were supported by a grant from the Simons Foundation Autism Research Initiative. We would also like to thank the Experimental Psychology Society for awarding G.V. the Grant for Study Visits for travel to Stanford University. Lastly, we would like to thank all our participants and their families for their contribution.

References

1. Jones RSP, Quigney C, Huws JC (2003) First-hand accounts of sensory perceptual experiences in autism: a qualitative analysis. *Journal of Intellectual & Developmental Disability* 28(2):112-121.

2. Ben-Sasson A et al. (2008) A Meta-Analysis of Sensory Modulation Symptoms in Individuals with Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders* 39:1-11.

3. Rubenstein JLR, Merzenich MM (2003) Model of autism: increased ratio of excitation/inhibition in key neural systems. *Genes, Brain and Behavior* 2(5):255-267.

4. Nelson SB, Valakh V (2015) Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders. *Neuron* 87(4):684-698.

5. Hahn N et al. (2013) Monogenic heritable autism gene neuroligin impacts Drosophila social behaviour. *Behavioural Brain Research* 252:450-457.

6. Gao R, Penzes P (2015) Common Mechanisms of Excitatory and Inhibitory Imbalance in Schizophrenia and Autism Spectrum Disorders. *Current Molecular Medicine* 15(2):146-167.

7. Plaisted K, Swettenham J, Rees L (1999) Children with Autism Show Local Precedence in a Divided Attention Task and Global Precedence in a Selective Attention Task. *Journal of Child Psychology and Psychiatry* 40:733-742.

8. Milne E et al. (2002) High motion coherence thresholds in children with autism. *Journal of Child Psychology and Psychiatry* 43(2):255-263.

9. Miles JH (2011) Autism spectrum disorders—A genetics review. *Genetics in Medicine* 13(4):278-294.

10. LeBlanc JJ et al. (2015) Visual evoked potentials detect cortical processing deficits in Rett syndrome. *Annals of Neurology* 78:775-786.

11. Burd L, Gascon GG (1988) Rett Syndrome: Review and Discussion of Current Diagnostic Criteria. *Journal of Child Neurology* 3:263-268.

12. West RJH, Furmstron R, Williams CAC, Elliott CJH (2015) Neurophysiology of Drosophila Models of Parkinson's Disease. *Parkinson's Disease* 2015:1-11.

13. Moloney A, Sattelle DB, Lomas DA, Crowther DC (2010) Alzheimer's disease: insights from Drosophila melanogaster models. *Trends in Biochemical Sciences* 35(4):228-235.

14. Reiter LT, Potocki L, Chien S, Gribskov M, Bier M (2001) A Systematic Analysis of Human Disease-Associated Gene Sequences In Drosophila melanogaster. *Genome Research* 11:1114-1125.

15. Fischbach K-F, Dittrich APM (1989) The optic lobe of Drosophila
melanogaster. I. A Golgi analysis of wild-type structure. *Cell and Tissue*
Research 258.

16. Behnia R, Desplan C (2015) Visual circuits in flies: beginning to see the
whole picture. *Current Opinion in Neurobiology* 34:125-132.

17. Afsari F et al. (2014) Abnormal visual gain control in a Parkinson's
disease model. *Human Molecular Genetics* 23(17):4465-4478.

18. Clark DA et al. (2014) Flies and humans share a motion estimation
strategy that exploits natural scene statistics. *Nature*
Neuroscience 17(2):296-303.

19. Kondapalli K, Hack A, Schushan M, Ben-Tal N, Rao R (2013) Functional
Evaluation of Autism Associated Mutations in NHE9. *Nature*
Communications 4(2510):1-12.

20. Skottun BC, Skoyles JR (2007) Some remarks on the use of visually
evoked potentials to measure magnocellular activity. *Clinical*
Neurophysiology 118:1903-1905.

21. Baron-Cohen S, Wheelwright, S, Skinner R, Martin J, Clubley E (2001)
The Autism-Spectrum Quotient (AQ): evidence from Asperger
Syndrome/high-functioning autism, males and females, scientists and
mathematicians. *Journal of Autism and Developmental Disorders* 31:5-17.

22. Lord C et al. (2000) The autism diagnostic observation schedule-generic:
a standard measure of social and communication deficits associated with
the spectrum of autism. *J Autism Dev Disord* 30:205-223.

23. Robertson AE, Simmons DR (2013) The Relationship between Sensory
Sensitivity and Autistic Traits in the General Population. *J Autism Dev*
Disord 43:775–784.

24. McKeefry DJ, Russell MHA, Murray IJ, Kulikowski JJ (1996) Amplitude
and phase variations of harmonic components in human achromatic and
chromatic visual evoked potentials. *Visual Neuroscience* 13:639-653.

25. McLelland D, Baker PM, Ahmed B, Bair W (2010) Neuronal Responses
during and after the Presentation of Static Visual Stimuli in Macaque
Primary Visual Cortex. *Journal of Neuroscience* 30(38):12619-12631.

26. Pei F, Baldassi S, Norcia AM (2014) Electrophysiological measures of
low-level vision reveal spatial processing deficits and hemispheric
asymmetry in autism spectrum disorder. *Journal of Vision* 14(11):3-3.

27. Jemel B, Mimeault D, Saint-Amour D, Mottron L (2012) VEP contrast
sensitivity responses reveal reduced functional segregation of visual
processing channels in autism. *International Journal of*
Psychophysiology 10(6):85:298.

28. Schwede M, Garbett K, Mirnics K, Geschwind DH, Morrow EM (2013)
Genes for endosomal NHE6 and NHE9 are misregulated in autism
brains. *Molecular Psychiatry* 19:277-279.

29. Voineagu I et al. (2011) Transcriptomic analysis of autistic brain reveals
convergent molecular pathology. *Nature* 474(7351):380-384.

30. Said CP, Egan RD, Minshew NJ, Behrmann M, Heeger DJ (2013) Normal
binocular rivalry in autism: Implications for the excitation/inhibition
imbalance hypothesis. *Vision Research* 77:59-66.

31. Dickinson A, Jones M, Milne E (2016) Measuring neural excitation and
inhibition in autism: Different approaches, different findings and different
interpretations. *Brain Research* 1648:277-289.

32. Simmons DR et al. (2009) Vision in autism spectrum disorders. *Vision*
Research 49:2705-2739.

33. Greenaway R, Davis G, Plaisted-Grant K (2013) Marked selective
impairment in autism on an index of magnocellular
function. *Neuropsychologia* 51:592-600.

34. Sutherland A, Crewther DP (2010) Magnocellular visual evoked potential
delay with high autism spectrum quotient yields a neural mechanism for
altered perception. *Brain* 133:2089-2097.

35. Courchesne E, Redcay E, Kennedy DP (2004) The autistic brain: birth
through adulthood. *Current Opinion in Neurology* 17:489-496.

36. Seltzer MM et al. (2003) The Symptoms of Autism Spectrum Disorders in
Adolescence and Adulthood. *J Autism Dev Disord* 33(6):565-581.

37. Mcgovern CW, Sigman M (2005) Continuity and change from early
childhood to adolescence in autism. *J Child Psychol Psychiatry*
46(4):401–408.

38. Pullikuth AK (2006) Molecular characterization of sodium/proton
exchanger 3 (NHE3) from the yellow fever vector, Aedes aegypti. *Journal*
of Experimental Biology 209:3529-3544.

39. Ben-Ari Y (2014) The GABA excitatory/inhibitory developmental
sequence: A personal journey. *Neuroscience* 279:187-219.

40. Sharma P et al. (2005) Isogenic autisomes to be applied in optimal
screening for novel mutants with viable phenotypes in Drosophila
melanogaster. *Journal of Neurogenetics* 19:57-85.

41. Carpenter JM (1950) A new semisynthetic food medium for
Drosophila. *Dros Inf Serv* 24:96-97.

 Figure legends

Figure 1. Human and *Drosophila* steady-state electrophysiology
methods. Panel A illustrates an example of electrophysiological responses (in
the adult AQ dataset) as a function of time (red trace) in response to stimulus
flicker (black trace). Two peaks can be observed after each increase in
stimulus contrast giving rise to the first (1F) and second (2F) responses,
which are evident in the frequency domain after Fourier transform (lower
panel A). Adult participants (for both the autistic adult dataset and the
neurotypical AQ dataset) were presented with a grid of sinusoidal gratings
flickering at 7Hz whilst SSVEPs were recorded using a 64-channel EEG cap
(C). Children were tested with a binary noise stimulus flickering at 5.12Hz and
a 128-channel EEG cap (D). SSVEPs were measured from the occipital
electrode Oz (indicated by green dots in e, f, g) where the highest first
harmonic amplitude was centered, indicated by the heatmaps for the adult AQ
dataset (E), autistic adult dataset (F) and child dataset (G). Fruit fly
electrophysiological data were acquired as shown (B) using square wave
stimulus flicker; see Methods for more details.

Figure 2. Older ASD-mimic flies and autistic humans have visual deficits
in the transient pathway. Contrast response functions for adult *Nhe3* mutant
flies (*Nhe3^{KG08307}* homozygotes, red squares and *Nhe3^{KG08307} /Df(2L)BSC187*,
purple diamonds) were similar at the first harmonic (a one-way ANOVA
showed no effect of group $F_{2,33} = 0.05$, $P = 0.95$, panel A) but responses were
reduced for P/P (simple contrast, P=0.025) and P/Df mutants compared to
controls at the second harmonic (simple contrast $P = 0.001$; ANOVA group
effect $F_{2,33} = 6.71$, $P < 0.01$; panel B). Ratios between frequencies ($\frac{1F-2F}{1F+2F}$)
were significantly higher for P/P ($P < 0.001$) and for P/Df ($P < 0.0001$) than for
the control genotype (C). First harmonic responses were also similar for the
high AQ and low AQ groups (panel D) and for autistic and neurotypical adults
(panel G). However, second harmonic responses were reduced for both
adults with high AQ (panel E) and autistic adults compared to controls (panels
H). The ratio between harmonics was also higher in both experimental groups
compared to controls (panels F and I, $P = 0.005$ and $P = 0.04$, respectively).
Curved lines are hyperbolic function fits to the data. Frequency ratios are
baselined in respect to the mean over groups of each comparison for display
purposes. Error bars in all panels represent ±SEM.
636

Figure 3. Young ASD-mimic flies and autistic children have visual
deficits in the sustained pathway. Young fruit flies showed reduced
responses at the first harmonic ($F_{2,33} = 3.73$, $P = 0.035$; panel A) with P/P and
P/Df flies showing a significant difference from control flies (respectively, P =

0.016 and P = 0.040). There was also a significant effect of genotype at the
second harmonic ($F_{2,33}$ = 3.39, P = 0.046, panel B). P/Df flies showed a
significant difference from control flies (P = 0.018), however, P/P showed a
non-significant difference from controls (P = 0.064). The flies had normal
frequency ratios (panel C). Autistic children also showed reduced first
harmonic (t_{28} = 2.065, P = 0.048; panel D) but not second harmonic
responses (t_{28} = 1.26, P = 0.22; panel E) and had frequency ratios similar to
that of control children (t_{28} = 1.21, P = 0.24; panel F). Curved lines are
hyperbolic function fits to the data. Frequency ratios are baselined in respect
to the mean over groups of each comparison for display purposes. Error bars
in all panels represent ±SEM.

Figure 4. Positive relationship between the number of autistic traits and
first/second harmonic ratio. Scatterplot showing a significant positive
relationship between AQ scores and frequency ratios in the 100 neurotypical
adult dataset indicating a gradual increase of impairment with the number of
reported autistic traits. The black line indicates the regression line of best fit.
Shaded grey areas show histograms of AQ scores and frequency ratios. Blue-
red color transition indicates number of AQ traits with participants split by
median into low and high AQ groups as presented in *Fig. 2*.