1 # Long-read sequence capture of the hemoglobin gene clusters
2 # across species

4 Siv Nam Khang Hoff*[a], Helle T. Baalsrud*[a‡], Ave Tooming-Klunderud[a], Morten
5 Skage[a], Todd Richmond[b], Gregor Obernosterer[c], Reza Shirzadi[d], Ole Kristian
6 Tørresen[a], Kjetill S. Jakobsen[a] and Sissel Jentoft[a‡]
7
8 [a] Centre for Ecological and Evolutionary Synthesis, Department of Biosciences,
9 University of Oslo, Oslo, Norway
10 [b] Roche NimbleGen Inc., Madison, WI, USA
11 [c] Roche Diagnostics, Mannheim, Germany
12 [d] Roche Diagnostics, Oslo, Norway
13
14 * these authors contributed equally to this work
15
16 ‡ Correspondence to: h.t.baalsrud@ibv.uio.no, sissel.jentoft@ibv.uio.no

Abstract

Combining high-throughput sequencing with targeted sequence capture has become an attractive tool to study specific genomic regions of interest. Most studies have so far focused on the exome using short-read technology. These approaches are not designed to capture intergenic regions needed to reconstruct genomic organization, including regulatory regions and gene synteny. Here, we demonstrate the power of combining targeted sequence capture with long-read sequencing technology for comparative genomic analyses of the hemoglobin (Hb) gene clusters across eight species separated by up to 70 million years. Guided by the reference genome assembly of the Atlantic cod (*Gadus morhua*) together with genome information from draft assemblies of selected codfishes, we designed probes covering the two Hb gene clusters. Use of custom-made barcodes combined with PacBio RSII sequencing led to highly continuous assemblies of the LA (~100kb) and MN (~200kb) clusters, which include syntenic regions of coding and intergenic sequences. Our results revealed an overall conserved genetic organization and synteny of the *Hb* genes within this lineage, yet with several, lineage-specific gene duplications. Moreover, for some of the species examined, we identified amino acid substitutions at two sites in the *Hbb1* gene as well as length polymorphisms in its regulatory region, which has previously been linked to temperature adaptation in Atlantic cod populations. This study highlights the use of targeted long-read capture as a versatile approach for comparative genomic studies by generation of a cross-species genomic resource elucidating the evolutionary history of the *Hb* gene family across the highly divergent group of codfishes.

Key words: Targeted sequence capture, comparative genomics, Gadiformes, PacBio sequencing, teleosts

Introduction

The rapid advancement of high-throughput sequencing has over the last decade revolutionized genomic research with the increasing numbers of whole genome resources available for multiple vertebrate species, including the diverse group of teleost fishes (Volff 2004; Ellegren 2014; Goodwin *et al.* 2016; Malmstrøm *et al.* 2016; 2017). However, whole genome sequencing (WGS) and generation of high quality genome assemblies are still considered costly and time-consuming (Jones & Good 2015). For investigations concerning specific genomic regions there is no need for complete genome information, which has spurred the development of reduction complexity approaches such as targeted sequence capture (Teer *et al.* 2010; Grover *et al.* 2012; Samorodnitsky *et al.* 2015). The basic idea of targeted sequence capture involves design of specific probes covering the particular genomic area of interest generating an enriched coverage of the targeted sequences (Turner *et al.* 2009; Grover *et al.* 2012). Most studies using targeted sequence capture have to a large extent been directed towards the exome, often supported by the existence of a reference genome (Broeckx *et al.* 2014; Yoshihara *et al.* 2016), or transcriptome assemblies (Syring *et al.* 2016). Recent reports have, however, been focusing on off-target sequences in noncoding regions (Guo *et al.* 2012; Syring *et al.* 2016; Yoshihara *et al.* 2016; Morin *et al.* 2016), as they may contain crucial regulatory elements varying in sequence and length between populations or species and could be of functional and evolutionary importance (Woolfe *et al.* 2004).

To date, comparative studies using sequence capture have been mainly based on short-read sequencing technology and probe design targeting genic regions (George *et al.* 2011; Samorodnitsky *et al.* 2015; Bragg *et al.* 2015; Li *et al.* 2015). Consequently, construction of continuous sequences enabling resolution of gene organization across species has not yet been looked into. Comparative genetic studies of gene organization or synteny requires longer, more continuous stretches of DNA containing more than one gene (Huddleston *et al.* 2014). By its ability to span long stretches of repeats, long-read sequencing technology has been successfully applied to improve genome assembly statistics and generation of highly continuous genome assemblies for a growing number of species (English *et al.* 2012; Kim *et al.* 2014; Tørresen *et al.* 2017; Kolmogorov *et al.* 2018; Tørresen *et al.* 2018). Incorporation of long PacBio reads have for instance resulted in a significantly improved version of the Atlantic cod (*Gadus morhua*) genome assembly, i.e. a 50-fold increase in sequence continuity and a 15-fold reduction in the proportion of gap (Tørresen *et al.* 2017). Furthermore, recent studies report the combination of capture and long read

sequencing as highly efficient in enriching and assembly of full length complex genes as well as detailed characterization of chromosomal structural variations (Wang *et al.* 2015; Witek *et al.* 2016; Giolai *et al.* 2017). Correspondingly, utilizing long-read sequencing technology in combination with targeted capture could yield longer continuous assemblies of specific genomic regions of interest, allowing in-depth comparative genetic studies including synteny analyses, in species where reference genomes are not available.

In fishes, the hemoglobin (*Hb*) gene family, encoding the protein subunits Hba and Hbb, has shown to be of importance for ecological adaptation, as environmental factors such as temperature directly influences the ability of Hb to bind O_2 at respiratory surfaces and its subsequent release to tissues (Wells 2005). In a recent report, a characterization of the *Hb* gene repertoire by comparative draft genome analysis uncovered a remarkably high *Hb* gene copy variation within the codfishes (Baalsrud *et al.* 2017). Based on the gene copy number a negative correlation between the number of *Hb* genes and depth of which the species occur was observed, as well as signs of diversifying selection on the gene paralogues suggesting that the variable environment in epipelagic waters have facilitated a larger more diverse *Hb* gene repertoire (Baalsrud *et al.* 2017). However, the rather fragmented draft genomes did not allow for reconstruction of the gene synteny and a deeper understanding of the evolution of the Hb-clusters.

Moreover, two tightly linked polymorphisms at amino acid positions 55 and 62 of the Hbb1-globin suggested associated with thermal adaptation are demonstrated in Atlantic cod populations. These polymorphisms exhibit a latitudinal cline in allele frequency in populations inhabiting varying temperature and oxygen regimes for Atlantic cod in the North Atlantic and Baltic Sea (Andersen *et al.* 2009). Populations found in the southern regions display the Hbb1-1 variant (Met55Lys62), whereas more northern populations largely display the Hbb1-2 variant (Val55Ala62) (Andersen *et al.* 2009). The Hbb1-1 variant has been shown to be insensitive to temperature whereas Hbb1-2 is temperature dependent with a higher O_2 affinity than Hbb1-1 at colder temperatures (Andersen *et al.* 2009), however, this has been questioned by Barlow *et al.* 2017. Additionally, an indel polymorphism within the promoter of the *Hbb1* gene has been reported to be in linkage disequilibrium with the above-mentioned polymorphisms (Star *et al.* 2011). Examination of multiple Atlantic cod populations uncovered that a longer promoter variant is associated with *Hbb1-2* and found to up-regulate its gene expression at higher temperatures, i.e. aiding in the maintenance of total oxygen-carrying capacity (Star *et al.* 2011).

In teleosts, the *Hb* genes are found to reside at two distinct genomic regions, the MN and LA cluster. Earlier reports have shown that there is a high evolutionary turnover of *Hb* genes across teleosts, with lineage-specific duplications and losses, which is in stark contrast to genes flanking the *Hb* genes, where the synteny is highly conserved (Quinn *et al.* 2010; Opazo *et al.* 2012; Feng *et al.* 2014). In this study, the overall goal was to elucidate the evolutionary past of the *Hb* clusters – including *Hb* genes, flanking genes and intergenic sequences – within the phylogenetically diverse group of codfishes (Gadiformes) by taking advantage of long read sequencing technology combined with targeted sequence capture. Eight codfish species were carefully selected on the basis of both phylogenetic and habitat divergence, implying that they are exposed to a variety of environmental factors as well as displaying distinct life-history traits. A highly continuous genome assembly of Atlantic cod (Tørresen *et al.* 2017) as well as low coverage draft genome assemblies of all eight species (Malmstrøm *et al.* 2017) were used in the design of the probes covering the genomic regions of interest. To enable targeted sequence capture for PacBio RSII sequencing, we modified the standard protocol for sequence capture offered by NimbleGen, i.e. the SeqCap EZ (Roche NimbleGen), as well as generating custom-made barcodes. This combined approach resulted in successful capturing and assembling of the two *Hb* gene clusters across the codfishes examined. The generation of highly continuous assemblies – for most of the species – enabled reconstruction of micro-synteny revealing lineage-specific gene duplications and identification of a relatively large and inter-species variable indel located in the promoter region between the *Hbb1* and *Hba1* genes.

Our study demonstrates that combining sequence capture technology with long-read sequencing is a highly efficient and versatile method to investigate specific genomic regions of interest – with respect to micro-synteny, regulatory regions and genetic organization – across distantly related species where genome sequences are lacking.

Results

Capture and *de novo* assembly of the target regions

The probe design (workflow schematically shown in Figure 1) resulted in a total of 7057 capture probes based on the target region in Atlantic cod, covering 337 kbp of sequence. 26774 probes were designed for the additional codfishes, covering a total of 1.82 Mbp of target sequence. The target region and the *Hb* gene clusters were successfully captured and enriched for eight codfishes; Atlantic cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus)*, silvery pout (*Gadiculus argenteus*), cusk (*Brosme brosme*), burbot (*Lota lota*), European hake (*Merluccius merluccius*), marbled moray cod (*Muraenolepus marmoratus*), and roughhead grenadier (*Macrourus berglax*), with number of reads spanning from 35573 to 73005 (Table 1). The average read length was 3032 bp, varying from 2836 bp in European hake to 3265 bp in burbot, resulting in the capture of an average of 16.71 Mbp per species (Table 1). By mapping reads back to the capture target region we found that the average mapping depth was variable across the target region for all species (Figure 2 and 3). Because of the skewed distribution of mapping depth, we also calculated median depth, which was, as expected, the highest for Atlantic cod at 242x (Table S1). The median mapping depth was consistently high for most of the other species as well, with the lowest for roughhead grenadier (12x). Both median and average depths for the MN region were persistently higher than for the LA region for all species, with the exception of silvery pout (Table S1). Furthermore, positions with high degree of mapping corresponded to the location of the genes used in the design of the capture probes across all species (Figure 2 and Figure 3). The percentage of reads mapping to the target region ranged from 25-43%, however, the percentage of the target region covered by reads ranged from 53-100% with five species having more than 90% of the target region covered by reads (Figure 4c and Table S1).

To address factors influencing capture success we compared various capture statistics to overall genomic divergence between the Atlantic cod genome and independent WGS data for each species from (Malmstrøm *et al.* 2017) (Table S1). We found a strong negative correlation between genomic divergence to Atlantic cod and median mapping depth against the target region (r=-0.90, Figure 4a), percent of reads mapped to the target region (r=-0.90, Figure 4b), and percentage of reads mapped to the target region (r=-0.84, Figure 4c).

We constructed *de novo* assemblies with quite consistent assembly statistics across species. Contig N50 ranged from 8055 bp in burbot to 6523 bp in European hake and the total number of contigs varied from 205 in burbot to 455 in marbled

moray cod. However, there was some variation in the size of the largest contig,

which ranged from 79 kbp in Atlantic cod to 30 kbp in marbled moray cod (Table 1).

To evaluate whether the assemblies represent the actual target regions we mapped

the *de novo* assemblies for each species to the target region in Atlantic cod, for which

the capture design is largely based upon (Figure 2 and 3). As expected, the

assemblies corresponded to the regions with high coverage of reads, i.e. the areas of

the target region containing genes included in the probe design.

Synteny of the *Hb* gene regions

Our capture design combined with long-read PacBio sequencing allowed us to

reconstruct micro-synteny of the MN and LA regions for Atlantic cod, haddock,

silvery pout, cusk, burbot, European hake, marbled moray cod and roughhead

grenadier (Figure 5). From the *de novo* assemblies, we were able to identify the

majority of the *Hb* genes and all of the flanking genes, which show that our capture

design was successful. However, the degree of continuity varied in the different

assemblies. In Atlantic cod, haddock, silvery pout, cusk, burbot and European hake

we could infer micro-synteny revealing that *Hb* and their flanking genes organization

largely followed what has previously been reported for Atlantic cod (Figure 5) (Star

et al. 2011). We found *Hbb4* only to be present in Atlantic cod (Figure 5b), which is in

line with Baalsrud *et al.* 2017. Furthermore, the *de novo* assemblies confirmed a linage-

specific duplication of *Hbb2* in the roughhead grenadier (Baalsrud *et al.* 2017).

Additionally, we identified a complete *Hba4*-like gene in the assembly of the marbled

moray cod, not earlier identified in this species. However, the *Hba4*-like gene in

marbled moray cod is likely a pseudogene due to a frameshift mutation causing

multiple stop codons. Furthermore, we were able to identify most of the *Hb* genes

reported in the recent study by Baalsrud *et al.* 2017, however, a few are missing from

our dataset (Figure 5a and b). Pairwise sequence alignment of these paralogous *Hb*

genes from Baalsrud *et al.* 2017 revealed sequence identities up to 98 % (Table S2).

Target region in the haddock and Atlantic cod genome assemblies

As a proof of concept, we reconstructed synteny of the target region in the most

recent genome assemblies of Atlantic cod (Tørresen *et al.* 2017; gadMor2) and

haddock (Tørresen *et al.* 2018; melAeg). In Atlantic cod, the MN region is located on

linkage group 2 (Figure 5a) and LA on linkage group 18 (Figure 5b), in haddock MN

is located on scaffold MeA_20160214_scaffold_771 (Figure 5a) and LA on scaffold

MeA_20160214_scaffold_1676 (Figure 5b). The overall synteny in Atlantic cod was

congruent with Wetten *et al.* 2010 except for the relative direction of the genes *foxj1a*

and *rhbdf1*. Furthermore, the organization of *Hb*s and their flanking genes in the

genome assembly of haddock is conserved compared to Atlantic cod with the
exception of *Hbb4* in the MN region, which is absent in haddock (Figure 5).

Repetitive sequences in the in the *Hb* gene regions

Quantifying the amount of repetitive sequences in the target region(s) was only
possible for Atlantic cod (gadMor2) and haddock (melAeg), for which high-quality
genome assemblies exist. The amount of repetitive sequences in the target region
differed between the MN cluster and the LA cluster in Atlantic cod. The MN region
(214 kb) contained a total of 10.7% repeated sequences, including 1.0% retro-
elements, 1.3% transposons, 5.8% simple repeats, and 2.6% of various low complexity
and unclassified repeated sequences (Table S3). In comparison, in the smaller LA
region (123 kb) the proportion of repeated sequences was twice as high (20.3%),
which comprised of 2.8% retro-elements, 2.4% transposons, 13.8% simple repeats,
and 1.3% of various low complexity and unclassified repeated sequences.
Furthermore, the orthologous target regions in haddock followed the same pattern.
The MN region contained 16.3 % repeated sequences, in contrast to 19.8 % found in
the LA region (Table S3).

Insertions and deletions in the promoter region of *Hba1 – Hbb1*

The previously shown 73 bp indel in the bi-directional promoter region of *Hba1* and
Hbb1 – discerning the cold-adapted migratory Northeast Artic (NEA) cod from the
more temperate-adapted southern Norwegian coastal (NC) cod (Star *et al.* 2011) –
was confirmed by the improved version of the NEA cod assembly (gadMor2). The
continuity of our capture assemblies (Figure 5) enabled location of the orthologous
captured regions in haddock, silvery pout and cusk. In each of the species an indel of
variable length were identified (Figure 6). Compared to the long promoter variant –
found to be linked with the *Hbb1-2* in Atlantic cod – the indel is shorter in the other
species by 11 bp in haddock, 22 bp in silvery pout and 56 bp in cusk (Figure 6).
Although the indels are varying in length, the conserved flanking sequences in the
alignment clearly show that they represent orthologous regions. Moreover, we found
the amino acid positions at 55 and 62 of the *Hbb1* gene to vary between species;
Haddock has Val55-Lys62, silvery pout has Met55-Gln62, while cusk has Met55-
Lys62 similarly to NEA cod (Figure 6). Additionally, we investigated amino acid
positions 55 and 62 in the *Hbb1* gene across a number additional codfish species for
which we have available gene sequences from Baalsrud *et al.* 2017, revealing these
sites to be variable across this lineage (Table S4). Ancestral reconstruction of *Hbb1*
demonstrated that the ancestral state in position 55 was Met in codfishes, and in

position 62 was Lys in all codfishes except *Bregmaceros cantori* (Supplementary
Figures S1 and S2).

Discussion

Capture of *Hb* gene clusters with 70 million years divergence time reveal conserved synteny and lineage-specific *Hb* duplications

We here demonstrate a successful in-solution targeted sequence capture and assembling of coding and noncoding sequences of the *Hb* clusters from codfish species separated by up to 70 million years (My) of evolution. Two features make our approach unique from earlier studies. First, the target regions consisted of both coding and noncoding genomic sequences. Second, we designed custom-made probes in order to utilize the long-read PacBio sequencing platform. In contrast to previous targeted capture sequencing studies based on short-read sequencing technologies (George *et al.* 2011; Mascher *et al.* 2013) our approach enabled the generation of highly continuous assemblies of the *Hb* clusters across distantly related codfishes.

The organization and orientation of the *Hb* flanking genes that we identified were conserved across all species (Figure 5a and b). However, in concordance with earlier studies of the *Hb* region, we found significant variation in copy numbers of the *Hb* genes, with linage specific duplications and losses (Star *et al.* 2011; Opazo *et al.* 2012; Feng *et al.* 2014; Baalsrud *et al.* 2017). We only found *Hbb4* in Atlantic cod, supporting earlier studies showing that *Hbb4* is the result of a recent duplication in this species (Borza *et al.* 2009; Baalsrud *et al.* 2017). Interestingly, the presence of two copies of *Hbb2* on the same contig in the roughhead grenadier *de novo* assembly confirmed a lineage-specific gene duplication of *Hbb2* in this species. This duplication was postulated in recent study of *Hbs* in codfishes (Baalsrud *et al.* 2017), but due to the lack of synteny in the draft genome assemblies, it was not possible to determine with certainty. Additionally, a copy of the *Hba4* was found in the *de novo* assembly of the marbled moray cod not found in the previous study by Baalsrud *et al.* (2017). The presence of a frame-shifting mutation that is causing multiple stop codons indicated that this *Hba4* gene is most likely a pseudogene. *Hba4* is also a pseudogene in the closely related species *Mora moro, Trachyrincus scabrus, T. murrayi* and *Melanonus zugmayeri* (Baalsrud *et al.* 2017). Although we identified most of the *Hb* genes from Baalsrud *et al.* (2017), a few were absent from this dataset (Figure 5a and b), which we suspect may be due to collapse of paralogous *Hb* genes, as they may have as high as 98% sequence identity (Table S2).

Length variation in the bi-directional *Hba1-Hbb1* promoter within the codfishes

The discovery of a promoter of variable length between *Hba1* and *Hbb1* in different species (Figure 6) was concordant with earlier findings of length variation in the homologous region in different populations of Atlantic cod (Star *et al.* 2011). The migratory NEA cod population has been shown to harbor the 73 bp longer variant at a higher frequency compared to coastal cod populations (see Figure 6 and Star *et al.* 2011). Interestingly, we found relatively long promoters with high sequence similarity to the NEA cod indel in haddock and silvery pout. In contrast, cusk displayed a relatively short promoter, however, still 17 bp longer than in NC cod (Figure 6). Furthermore, we found the amino acid positions 55 and 62 in *Hbb1*, known to be polymorphic in Atlantic cod, to be variable across all codfishes included in this study (Figure 6). Investigations of the same positions in a number additional codfishes for which we have available gene sequences (Baalsrud *et al.* 2017), revealed that these positions are highly variable across this linage (Table S4). Notably, the most likely ancestral state of codfish *Hbb1* is Met55Lys62 (Supplementary Figures S1 and S2). Cusk and the coastal/southern Atlantic cod thus both display the ancestral state as well as a short promoter, although the cusk promoter was 17 bp longer (Figure 6). Collectively, these results suggest two different scenarios for promoter length evolution. Scenario 1: The short promoter represents the ancestral state of the Gadidae-family (including cusk and Atlantic cod; see Malmstrøm *et al.* 2016) and that silvery pout and some populations of Atlantic cod have evolved a longer promoter. Scenario 2: The long promoter is the ancestral state with independent deletions of variable lengths in cusk, silvery pout, haddock and coastal/southern Atlantic cod (*Hbb1-1*). To disentangle this, we would need to obtain promoter sequences from additional gadiform species. Regardless, the short-long promoter polymorphism has been maintained throughout speciation events based on the presence of both variants in Atlantic cod. Moreover, in both scenarios, cusk and coastal/southern Atlantic cod (*Hbb1-1*) have maintained the ancestral Met55Lys62, while silvery pout, haddock and NEA cod (*Hbb1-2*) have acquired substitutions at these positions due to similar selection pressures or genetic drift. In this regard, it could be mentioned that the NEA cod, haddock and silvery pout display migratory behavior (e.g. diurnally feeding movements as well as seasonal spawning migrations) compared to the more stationary cusk and coastal cod (Eschemeyer & Fricke 2017) which could mean that they have a higher O_2 demand and are exposed to greater temperature variation, which in turn has selected for a temperature-dependent long promoter. Furthermore, given that promoter length and positions 55/62 at *Hbb1* are important genetic

components of temperature adaptation in Atlantic cod populations (Star *et al.* 2011),
they most likely play a role in temperature adaptation in the other codfishes.

Assembly success affected by probe design and repeat content

In some species, nearly the complete target region is assembled in large contigs
containing multiple genes including cusk, whereas in other species such as the more
distantly related roughhead grenadier, the cluster is more fragmented (Figure 5). In
all species, the areas of the target regions that harbor genes of which probes are
designed for, as well as any areas containing repeated sequences, have very high
depths in comparison to the areas of intergenic sequences (Figure 2 and 3). This
poses a challenge for the assembly software, which is based in the assumption of
uniform depth over the sequencing data (Miller *et al.* 2010).
Overall, the MN cluster seems to be more successfully assembled than the LA
cluster, which is more fragmented (Figure 5). Differences in assembly completeness
between the two regions might be a result of several factors. Firstly, the MN region
has more flanking genes in closer proximity to the *Hb* region, which results in a
higher density of probes. Secondly, the overall repeat content of the LA region is one
order of magnitude larger than in the MN region, largely due to the larger
proportion of simple repeats. Repeat content is a major interference in capture
experiments because unwanted repetitive DNA may be enriched for, especially if
there are repeated sequences included in the probe design. Furthermore, if probes
were not completely covered by target DNA they get single-stranded sticky ends that
can hybridize to repetitive or other non-target DNA (Newman & Austin 2016).
Lastly, unless there were some longer reads that bridged such areas, this would in
turn have led to gaps in the downstream *de novo* assemblies. Following that the
assembly success was possibly a result of read length, we reason that a future
increase of the average read length from 3 kbp to 5-10 kbp, would be sufficient to
substantially increase the completeness of the assemblies. Due to the current circular
consensus (CCS) PacBio sequencing technology, however, which is a trade-off
between accuracy and length of reads, longer reads with sufficient accuracy are not
feasible.

Long-read sequencing capture across species harbors new potential for
comparative genomic studies

The number of reads mapping to the target region was in the range of 23-43%, which
may be seemingly low compared with other capture studies. For instance, a whole
exome capture study on humans reported 56.1% of reads mapped to the target
region (Guo *et al.* 2012) and a similar study in rats reported to have 78.3% of reads on

target (Yoshihara *et al.* 2016). In contrast to our study however, these capture

experiments enriched either the exome or ultra-conserved elements within a single

species, allowing for more efficient capture of conserved sequences. We were

however, able to cover up to 98% of the target region with a sequencing depth of >10

reads across species (Table S1) which is similar to what mentioned experiments

within human and rat exomes reported (Guo *et al.* 2012; Yoshihara *et al.* 2016) and the

main difference is the higher percentage of non-target sequences in our study.

We were able to capture complete genes for species with 70 My divergence

time from the Atlantic cod (Figure 5). As expected, we found that capture success

declines with increased sequence divergence between the reference genome of which

we chiefly based our capture probes and the genomes of the included codfishes

(Figure 4). It has been reported that orthologous exons were successfully captured in

highly divergent frog species (with 200 My of separation), nevertheless the capture

success greatly decreased with increased evolutionary distance (Hedtke *et al.* 2013).

Similarly, it has been demonstrated that it is possible to capture >97% of orthologous

sequences in four species of primates that diverged from humans 40 My ago, using

probes entirely based on the human exome (George *et al.* 2011). Further, exomes were

effectively captured from skink species that diverged up to 80 My from the reference,

yet reporting a substantial decline in capture efficiency for sequences >10 % different

from the reference species (Bragg *et al.* 2015). Our study stands out from previous

capture experiments because intergenic, noncoding sequences in addition to genes

were captured across distantly related species. Efficient capture of intergenic

sequences requires less divergence time, as these regions usually evolve faster than

genes (Koonin & Wolf 2010). Thus, the most distantly related species from Atlantic

cod for which we captured both coding and noncoding sequences was burbot, which

diverged from Atlantic cod 46 My (Figure 5). We argue, in line with (Schott *et al.*

2017), that sequence divergence may be a more exact predictor of capture success

than evolutionary distance, as the sequence capture process is mainly influenced by

the difference between the probe sequence and the target sequence. European hake,

marbled moray cod and roughhead grenadier all diverged from cod about 70 My

ago, however, the European hake *Hb* regions was more successfully captured and

assembled (Table 1; Figure 2). This could be due to European hake having a lower

genome-wide divergence to Atlantic cod than marbled moray cod and roughhead

grenadier (809k vs 879k and 907k SNPs; Table S1).

Finally, it should be mentioned that cusk – which diverged from Atlantic cod

39 My ago – was added to the experimental design after the probes were generated.

Thus, the successful capture of cusk was therefore solely based on cross-species
target enrichment, demonstrating the power of heterologous probe targeting.

Concluding remarks and future perspectives

Here, we have successfully demonstrated that combining targeted sequence capture
with long-read sequencing technology is as an efficient approach to obtain high
quality sequence data of a specific genomic region, including both coding and
noncoding sequences, across evolutionary distant species. We show that genome-
wide divergence is of importance for capture success across species. Furthermore, the
use of long-read sequencing augmented the *de novo* assembly of regions containing
repeated sequences that would otherwise fragment assemblies based on short-read
sequencing. This is crucial for capturing complete intergenic sequences that may be
highly divergent compared to genic regions even among fairly closely related
species. Given the rapid development in sequencing technologies future methods
will enable read-through of repeated regions and thus further increase the
completeness of assemblies. Moreover, a less stringent hybridization protocol should
make it possible to capture sequences across even deeper evolutionary time. In sum,
our approach has generated a cross-species genomic resource across distantly related
codfishes and show the potential of enhancing comparative genomic studies of
continuous genic and intergenic regions between any eukaryotic species-group
where genomic resources are scarce.

Material and methods

Defining target region and probe design

The probe design was chiefly based on the high-quality genome of Atlantic cod, known as gadMor2 (Tørresen *et al.* 2017). In addition, species-specific probes were designed based on low-coverage assembled genomes (Malmstrøm *et al.* 2016) for ten selected species representing six families in the Gadiformes order. These species were Atlantic cod (*Gadus morhua*), Alaskan Pollock (*Gadus chalcogrammus*), polar cod (*Boreogadus saida*), haddock (*Melanogrammus aeglefinus*), Silvery pout (*Gadiculus argenteus*), burbot (*Lota lota*), European hake (*Merluccius merluccius*), roughhead grenadier (*Macrourus berglax*), roughsnout grenadier (*Trachyrincus scabrus*) and marbled moray cod (*Muraenolepus marmoratus*).

To retrieve relevant sequence data for the probe design, the MN and LA *Hb* regions were extracted from gadMor2 (Figure 1). These sequences, hereby known as the target region, were then used as queries in BLAST (Altschul *et al.* 1990) searches with an E-value threshold of <0.1 against the genome assembly data of all ten species.

In total, 5604 sequences from the chosen species were supplied to NimbleGen probe design. Protein coding genes from the ENSEMBL database were used to define the regions to be tiled in the probe design (Table S5) within the target region of the Atlantic cod, and the unitigs for each of the additional codfishes.

NimbleGen SeqCap EZ capture probes were designed by NimbleGen (Roche, Madison, USA) using a proprietary design algorithm. NimbleGen offers an in-solution sequence capture protocol, which includes custom made probes. Uniquely, the capture probes from NimbleGen are tiled to overlap the target area. 50 – 100 bp (average 75 bp) probes where designed tiled over the target region (subset of gadMor2) resulting in each base, on average, being covered by two probes. Additionally, raw reads from Illumina sequencing from (Malmstrøm *et al.* 2016; 2017) were used for each species to estimate repetitive sequences in each of the species genomes, aiming to discard probes containing any repeats. A more detailed description of the probe design is provided in Supplementary Materials and Methods and Table S6.

Sample collection and DNA extraction

Our goal working with animals is always to limit any harmful effects of our research on populations and individuals. Whenever possible we try to avoid animals being euthanized to serve our scientific purpose alone by collaborating with commercial fisheries or museums. The tissue samples used in this study are either from

commercially fished individuals intended for human consumption or museum
specimen. The commercially caught fish were immediately stunned, by bleeding
following standard procedures by a local fisherman. There is no specific legislation
applicable to this manner of sampling in Norway, however it is in accordance with
the guidelines set by the 'Norwegian consensus platform for replacement, reduction
and refinement of animal experiments' (www.norecopa.no).
DNA was extracted from tissue samples using High Salt DNA Extraction
method by Phill Watts (https://www.liverpool.ac.uk/~kempsj/IsolationofDNA.pdf ,
last day accessed: December 2017). The concentration and purity of the DNA samples
were quantified using NanoDrop (Thermo Fisher Scientific, Waltham, MA, USA) and
a Qubit fluorometer (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA). Due
to poor DNA quality, three species included in the probe design; Alaskan Pollock,
polar cod and roughsnout grenadier were excluded from further analysis. In total,
eight species were sequenced; seven of these species were included in the probe
design and one closely related species (cusk), which serves as a cross species capture
experiment without species-specific probes.

Capture, library preparation and sequencing

The sequencing libraries were prepared following a modified Pacific Biosciences
SeqCap EZ protocol. As multiplexing of the samples before capture was required,
barcodes were designed at the Norwegian Sequencing Centre
(http://www.sequencing.uio.no) using guidelines from Pacific Biosciences, for more
information see Supplementary Materials and methods, Table S7 and Figure S3.
Genomic DNA was sheared to 5 kb fragments using MegaRuptor (Diagenode,
Seraing (Ougrée), Belgium). Due to poorer DNA quality, fragmenting was not done
for European hake. For this sample together with fragmented DNA from roughhead
grenadier, short fragments were removed using BluePippin (Sage Science, Beverly,
MA, USA) before library preparation. Illumina libraries were prepared using KAPA
Hyper Prep kit (Kapa Biosystems, Wilmington, MA, USA) and barcoded using
different Illumina barcodes. PacBio barcodes were implemented during pre-capture
amplification of libraries. After amplification, fragment length distribution was
evaluated using Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA) and
samples were pooled in equimolar ratio. During hybridization, SeqCap EZ
Developer Reagent (universal repeat blocker for use on vertebrate genomes) and
oligos corresponding to Illumina and PacBio barcodes were used for blocking.
Captured gDNA was amplified to ensure that sufficient amount of DNA was
available for PacBio library preparation. Size selection of the libraries was performed

using BluePippin. Final libraries were quality checked using Bioanalyzer and Qubit

fluorometer (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA) and

sequenced on RS II instrument (PacBio, Menlo Park, CA, USA) using P6-C4

chemistry with 360 minutes movie time. In total, 9 SMRT cells were used for

sequencing.

De novo assemblies

Reads were filtered and de-multiplexed using the 'RS_reads of insert.1' pipeline on

SMRT Portal (SMRT Analysis version smrtanalysis_2.3.0.140936.p2.144836). Each set

of reads corresponding to a given species was crossed-checked with their respective

six-nucleotide Illumina adapter. Reads containing an incorrect Illumina adapter were

removed. Adapter sequences were then trimmed using the application Prinseq-lite

v0.20.4 (Schmieder & Edwards 2011). The trimmed reads were assembled *de novo*

using Canu v1.4 +155 changes (r8150 c0a988b6a106c27c6f993dfe586d2336282336a6)

(Berlin *et al.* 2015). The Canu software is optimized for assembling single molecule

high noise sequence data. We specified genome size as the size of the target region

(300 kbp). Additionally, we ran PBJelly (English *et al.* 2012) on the Canu *de novo*

assemblies, using the raw reads to possible bridge gaps between scaffolds, settings

given in Supplementary Materials and Methods.

We assessed the assemblies by running Assemblathon 2 (Bradnam *et al.* 2013),

which reports assembly metrics such as the longest contig, the number of contigs,

and the N50 value. *De novo* assemblies of the MN and LA regions of Atlantic cod and

haddock were aligned and compared to their reference genomes, gadMor2 and

melAeg respectively, using BLAST and BWA v0.7.10 (Li & Durbin 2009) to determine

syntenic similarities and assembly completeness.

Estimating capture success

PacBio reads for all the species were mapped back to the Atlantic cod genome

assembly (gadMor2) in order to determine sequence capture success and target

mapping depths. Mapping was done using BWA-MEM v0.7.10 (Li & Durbin 2009).

Target-area read depth for all the species based on mapping against gadMor2, were

calculated using Samtools v1.3.1 (Li *et al.* 2009). We calculated both average and

median mapping depth against the target region as a whole and for the MN and LA

region separately. We also calculated percentage of reads that mapped to the target

region, and the percentage of the target regions covered by reads to a minimum

depth of 10x. To compare assemblies to the target region we additionally mapped the

assemblies to the target region. In order to verify the sequence capture process,

sequence data for Atlantic cod and haddock were mapped back to their reference

genomes using BWA-mem v0.7.10 (Li & Durbin 2009). The results were visualized
using Integrative Genome Viewer (Robinson *et al.* 2011).
To obtain an independent measure of divergence between species in the
capture experiment we calculated genome wide level of divergence of each species to
the reference genome of Atlantic cod using low-coverage whole-genome sequence
data from (Malmstrøm *et al.* 2016; 2017). We mapped raw reads to Atlantic cod using
BWA-MEM (Li & Durbin 2009) and called SNPs using the Freebayes variant caller
(Garrison & Marth 2012). Some species are more closely related to Atlantic cod than
others, which could introduce a bias in mapping. To avoid this, we only looked at
genomic regions where all species mapped. The number of SNPs was then used as an
estimate of genome-wide divergence of each species to Atlantic cod. We also mapped
a low-coverage genome of Atlantic cod to the Atlantic cod reference genome as a
control.
In pursuance of factors explaining capture success we tested for correlations
and plotted the relationship between the genome wide level of divergence and the
following variables; median mapping depth against the target region (for total, LA
and MN, respectively); percentage of reads that mapped to the target region; and the
percentage of the target region covered by reads. All tests and plots were done using
R version 3.2.5 (R Core Team 2016).
Assembly continuity is very often hampered by the presence of repeats, which
create gaps. We therefore quantified repeat-content in the target region extracted
from gadMor2 and orthologous regions in haddock using Repeatmasker Open 3.0
(Smit A *et al.* 2010) for the MN region and the LA region separately.

543 Identifying gene location and synteny

In order to identify the genes of interest and their location in the assembly we used
local sequence alignment algorithm BLAST v2.4.0 (Altschul *et al.* 1990) with protein
sequences of the genes of interest (Table S5) as queries. tblastn was used with an e-
value of 0.1. Investigation of *Hbb1-Hba1* promoter region was done for four species,
Atlantic cod, haddock, silvery cod and cusk. Sequences were aligned with ClustalW
default settings using MEGA7 (Kumar *et al.* 2016). Ancestral sequence reconstruction
was carried out for *Hbb-1* gene sequences from 24 species of codfishes from (Baalsrud
et al. 2017) using a maximum likelihood method implemented in MEGA7 (Kumar *et*
al. 2016).
Additionally, we estimated sequence identity using EMBOSS Needle (Rice *et*
al. 2000) with default settings, between *Hbb* gene sequences from (Baalsrud *et al.*

2017) that were missing and present in the *de novo* assemblies to evaluate similarity

(Table S2).

Acknowledgements

We would like to thank Marianne H. Selander Hansen and Alexander J. Nederbragt

for help with the initial design of this project. All computational work was

performed on the Abel Supercomputing Cluster (Norwegian Metacenter for High-

Performance Computing (NOTUR) and the University of Oslo), operated by the

Research Computing Services group at USIT, the University of Oslo IT Department.

Sequencing library creation and high-throughput sequencing were carried out at the

Norwegian Sequencing Centre (NSC), University of Oslo, Norway. This work was

funded by a grant awarded to K.S.J. from the Research Council of Norway (RCN

grant 222378).

Author contributions

H.T.B. and S.J. initially conceived and designed the study, with input from S.N.K.H,

A.T.-K., M.S., G.O., R.S., and K.S.J. Tissue samples were provided by S.J. and H.T.B.

Probe design was carried out by T.R. with assistance from S.N.K.H and H.T.B. DNA

extraction and sequence library preparation was performed by S.N.K.H and A.T.-K,

respectively. Sequence capture was carried out by S.N.K.H, A.T.-K., M.S. and G.O.

Filtering, mapping of sequences and *de novo* assemblies was done by S.N.K.H.,

assisted by O.K.T and H.T.B. Annotation of genes, synteny analyses, statistical

analyses and construction of all figures and tables was done by S.N.K.H and H.T.B.

The manuscript was written by S.N.K.H and H.T.B. with input from S.J. and K.S.J.

Competing interests

579   The authors declare that they have no competing interests.

Data and materials availability

581   All reads and assemblies (unitigs) reported on here, and the target region, subset of

582   gadMor2, as well as the relevant sequence data for the probe design from the chosen

583   species supplied to NimbleGen and the probe sequences have been deposited at

584   figshare under doi/xxx.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology*, **215**, 403–410.

Andersen O, Wetten OF, De Rosa MC *et al.* (2009) Haemoglobin polymorphisms affect the oxygen-binding properties in Atlantic cod populations. *Proceedings of the Royal Society B: Biological Sciences*, **276**, 833–841.

Baalsrud HT, Voje KL, Tørresen OK *et al.* (2017) Evolution of Hemoglobin Genes in Codfishes Influenced by Ocean Depth. *Scientific Reports*, **7**, 1–10.

Barlow SL, Metcalfe J, Righton DA, Berenbrink M (2017) Life on the edge: O2 binding in Atlantic cod red blood cells near their southern distribution limit is not sensitive to temperature or haemoglobin genotype. *The Journal of Experimental Biology*, **220**, 414–424.

Berlin K, Koren S, Chin C-S *et al.* (2015) Assembling large genomes with single-molecule sequencing and locality-sensitive hashing. *Nature Biotechnology*, **33**, 623–630.

Borza T, Stone C, Gamperl AK, Bowman S (2009) Atlantic cod (*Gadus morhua*) hemoglobin genes: multiplicity and polymorphism. *BMC Genetics*, **10**, 51–14.

Bradnam KR, Fass JN, Alexandrov A *et al.* (2013) Assemblathon 2: evaluating de novo methods of genome assembly in three vertebrate species. *Giga Science,* **2**, 10.

Bragg JG, Potter S, Bi K, Moritz C (2015) Exon capture phylogenomics: efficacy across scales of divergence. *Molecular Ecology Resources*, **16**, 1059–1068.

Broeckx BJG, Coopman F, Verhoeven GEC *et al.* (2014) Development and performance of a targeted whole exome sequencing enrichment kit for the dog (Canis Familiaris Build 3.1). *Scientific Reports*, **4**, 1522–4.

Ellegren H (2014) Genome sequencing and population genomics in non-model organisms. *Trends in Ecology & Evolution*, **29**, 51–63.

English AC, Richards S, Han Y *et al.* (2012a) Mind the Gap: Upgrading Genomes with Pacific Biosciences RS Long-Read Sequencing Technology (Z Liu, Ed,). *PLoS ONE*, **7**, e47768–12.

Eschemeyer WN, Fricke R. Catalog of fishes. http://research.calacademy.org/researchichthyology/catalog/fishcatmain.asp.2017. (Last accessed December 2017)

Feng J, Liu S, Wang X *et al.* (2014) Channel catfish hemoglobin genes: Identification, phylogenetic and syntenic analysis, and specific induction in response to heat stress. *Comparative Biochemistry and Physiology - Part D: Genomics and Proteomics*, **9**, 11–22.

621 Garrison E, Marth G (2012) Haplotype-based variant detection from short-read
622  sequencing. *bioRxiv*, 1207.3907.

623 George RD, McVicker G, Diederich R *et al.* (2011) Trans genomic capture and
624  sequencing of primate exomes reveals new targets of positive selection. *Genome*
625  *Research*, **21**, 1686–1694.

626 Giolai M, Paajanen P, Verweij W *et al.* (2017) Comparative analysis of targeted long
627  read sequencing approaches for characterization of a plant's immune receptor
628  repertoire. *BMC Genomics*, **18**, 521.

629 Goodwin S, McPherson JD, McCombie WR (2016) Coming of age: ten years of next-
630  generation sequencing technologies. *Nature Reviews Genetics*, **17**, 333–351.

631 Grover CE, Salmon A, Wendel JF (2012) Targeted sequence capture as a powerful
632  tool for evolutionary analysis1. *American Journal of Botany*, **99**, 312–319.

633 Guo Y, Long J, He J *et al.* (2012) Exome sequencing generates high quality data in
634 non-target regions. *BMC Genomics*, **13**, 194.

635 Hedtke SM, Morgan MJ, Cannatella DC, Hillis DM (2013) Targeted Enrichment:
636  Maximizing Orthologous Gene Comparisons across Deep Evolutionary Time (U
637  Joger, Ed,). *PLoS ONE*, **8**, e67908–10.

638 Huddleston J, Ranade S, Malig M *et al.* (2014) Reconstructing complex regions of
639  genomes using long-read sequencing technology. *Genome Research*, **24**, 688–696.

640 Jones MR, Good JM (2015) Targeted capture in evolutionary and ecological
641  genomics. *Molecular Ecology*, **25**, 185–202.

642 Kim KE, Peluso P, Babayan P *et al.* (2014) Long-read, whole-genome shotgun
643  sequence data for five model organisms. *Scientific Data*, **1**, 140045–10.

644 Kolmogorov M, Yuan J, Lin Y, Pevzner P (2018) Assembly of Long Error-Prone
645  Reads Using Repeat Graphs. *bioRxiv*, 247148.

646 Koonin EV, Wolf YI (2010) Constraints and plasticity in genome and molecular-
647  phenome evolution. *Nature Reviews Genetics* , **11**, 487–498.

648 Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics
649  Analysis Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, **33**,
650  1870–1874.

651 Li C, Corrigan S, Yang L *et al.* (2015) DNA capture reveals transoceanic gene flow in
652  endangered river sharks. *Proceedings of the National Academy of Sciences*, **112**,
653  13302–13307.

654 Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler
655  transform. *Bioinformatics*, **25**, 1754–1760.

656 Li H, Handsaker B, Wysoker A *et al.* (2009) The Sequence Alignment/Map format and
657  SAMtools. *Bioinformatics*, **25**, 2078–2079.

Malmstrøm M, Matschiner M, Tørresen OK *et al.* (2016) Evolution of the immune
system influences speciation rates in teleost fishes. *Nature Genetics*, **48**, 1204–1210.
Malmstrøm M, Matschiner M, Tørresen OK, Jakobsen KS, Jentoft S (2017) Whole
genome sequencing data and de novo draft assemblies for 66 teleost species.
Scientific Data, **4**, 160132–13.
Mascher M, Muehlbauer GJ, Rokhsar DS *et al.* (2013) Anchoring and ordering NGS
contig assemblies by population sequencing (POPSEQ). *The Plant Journal,* **76**, 718–
727.
Miller JR, Koren S, Sutton G (2010) Assembly algorithms for next-generation
sequencing data. *Genomics*, **95**, 315–327.
Morin A, Kwan T, Ge B *et al.* (2016) Immunoseq: the identification of functionally
relevant variants through targeted capture and sequencing of active regulatory
regions in human immune cells. *BMC Medical Genomics*, **9**, 1–12.
Newman CE, Austin CC (2016) Sequence capture and next-generation sequencing of
ultraconserved elements in a large-genome salamander. *Molecular Ecology*, **25**,
6162–6174.
Opazo JC, Butts GT, Nery MF, Storz JF, Hoffmann FG (2012) Whole-Genome
Duplication and the Functional Diversification of Teleost Fish Hemoglobins.
Molecular Biology and Evolution, **30**, 140–153.
Quinn NL, Boroevich KA, Lubieniecki KP *et al.* (2010) Genomic organization and
evolution of the Atlantic salmon hemoglobin repertoire. *BMC Genomics*, **11**, 539–
17.
Rice P, Longden I, Bleasby A (2000) EMBOSS: The European Molecular Biology Open
Software Suite. *Trends in Genetics*, **16**, 276–277.
Robinson JT, Thorvaldsdóttir H, Winckler W *et al.* (2011) Integrative genomics
viewer. *Nature Biotechnology*, **29**, 24–26.
Samorodnitsky E, Datta J, Jewell BM *et al.* (2015) Comparison of Custom Capture for
Targeted Next-Generation DNA Sequencing. *The Journal of Molecular Diagnostics*,
17, 64–75.
Schmieder R, Edwards R (2011) Quality control and preprocessing of metagenomic
datasets. *Bioinformatics*, **27**, 863–864.
Schott RK, Panesar B, Card DC *et al.* (2017) Targeted Capture of Complete Coding
Regions across Divergent Species. *Genome Biology and Evolution*, **9**, 398–414.
Smit A, Hubley R, Green P. RepeatMasker Open 3.0. http://www.repeatmasker.org.
2010. (Last accessed December 2017)
Star B, Nederbragt AJ, Jentoft S *et al.* (2011) The genome sequence of Atlantic cod
reveals a unique immune system. *Nature*, **477**, 207–210.

Syring JV, Tennessen JA, Jennings TN *et al.* (2016) Targeted Capture Sequencing in
Whitebark Pine Reveals Range-Wide Demographic and Adaptive Patterns
Despite Challenges of a Large, Repetitive Genome. *Frontiers in Plant Science*, **7**,
403–15.

Team RC. R Core Team. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL
http://www. R-project. org/.(3.3. 1) Software Vienna, Austria: R Foundation for
Statistical Computing; 2013.

Teer JK, Bonnycastle LL, Chines PS *et al.* (2010) Systematic comparison of three
genomic enrichment methods for massively parallel DNA sequencing. *Genome*
Research, **20**, 1420–1431.

Turner EH, Ng SB, Nickerson DA, Shendure J (2009) Methods for Genomic
Partitioning. *Annual Review of Genomics and Human Genetics*, **10**, 263–284.

Tørresen OK, Brieuc MSO, Solbakken MH *et al.* (2018) Genomic architecture of
haddock (*Melanogrammus aeglefinus*) shows expansions of innate immune genes
and short tandem repeats. *BMC Genomics*, **19**, 51.

Tørresen OK, Star B, Jentoft S *et al.* (2017) An improved genome assembly uncovers
prolific tandem repeats in Atlantic cod. *BMC Genomics*, **18**, 95.

Volff J-N (2004) Genome evolution and biodiversity in teleost fish. *Heredity*, **94**, 280–
294.

Wang M, Beck CR, English AC *et al.* (2015) PacBio-LITS: a large-insert targeted
sequencing method for characterization of human disease-associated
chromosomal structural variations. *BMC Genomics*, **16**, 214.

Wells RMG (2005) Blood-Gas Transport and Hemoglobin Function in Polar Fishes:
Does Low Temperature Explain Physiological Characters? In: *Physiology of Polar*
Fishes Fish Physiology. pp. 281–316. Academic Press.

Wetten OF, Nederbragt AJ, Wilson RC *et al.* (2010) Genomic organization and gene
expression of the multiple globins in Atlantic cod: conservation of globin-
flanking genes in chordates infers the origin of the vertebrate globin clusters.
BMC Evolutionary Biology, **10**, 315.

Witek K, Jupe F, Witek AI *et al.* (2016) Accelerated cloning of a potato late blight–
resistance gene using RenSeq and SMRT sequencing. *Nature Biotechnology*, **34**,
656–660.

Woolfe A, Goodson M, Goode DK *et al.* (2004) Highly Conserved Non-Coding
Sequences Are Associated with Vertebrate Development (Sean Eddy, Ed,). *PLoS*
Biology, **3**, e7–15.

Yoshihara M, Saito D, Sato T *et al.* (2016) Design and application of a target capture
sequencing of exons and conserved non- coding sequences for the rat. *BMC*
Genomics, **17**, 1–11.

734
735

Figure legends

Figure 1: Flowchart of sequence capture approach. a) Sequence data from the
Atlantic cod genome (Tørresen *et al.* 2017; gadMor2) combined with gene sequences
of target genes and sequences from low coverage genomes of the additional
codfishes are combined to generate probes. b) Isolated DNA is multiplexed with
Illumina and PacBio barcodes. c) Raw reads for each species are used to score all
probes, ensuring that no repeated sequences are present. DNA Probes are used in
solution on isolated DNA for all of the included species, hybridizing to the target
sequences. Target sequences are then captures and sequences on the PacBio RSII
sequencing platform. d) Downstream bioinformatics includes de-multiplexing of
reads and trimming, making the reads ready for downstream analysis such as
mapping and *de novo* assembly.

Figure 2: Mapping of reads and assemblies against the MN target region. Each
panel shows the reads and *de novo* assembly mapped against the MN target region in
grey and orange, respectively, for species a.) Atlantic cod, b) haddock, c) silvery
pout, d) cusk, e) burbot, f) European hake, g) marbled moray cod and h) roughhead
grenadier. The positions of genes in the target region are indicated at the top.

Figure 3: Mapping of reads and assemblies against the LA target region. Each
panel shows the reads and *de novo* assembly mapped against the LA target region in
grey and orange, respectively, for species a) Atlantic cod, b) haddock, c) silvery pout,
d) cusk, e) burbot, f) European hake, g) marbled moray cod and h) roughhead
grenadier. The positions of genes in the target region are indicated at the top.

Figure 4: The relationship between capture success and genomic divergence to
Atlantic cod. Linear regression of the relationship between the genomic divergence
to Atlantic cod (SNPs x 10^5) and a) median mapping depth for the MN region (blue),
LA region (red) and the combined target region (black); b) the percentage of reads
mapping to the target region; c) the percentage of the target region covered by reads
to a minimum depth of 10x. For each regression the correlation coefficient, r, is
shown along with a p-value. Each data point is labeled by species according to this
code: Ac=Atlantic cod, H=haddock, Sp=silvery pout, C=cusk, B=burbot, Eh=European
hake, Mm=marbled moray cod and Rg=roughhead grenadier.
770

Figure 5: Synteny of the Hb gene clusters. Genomic synteny of the hemoglobin gene
clusters shown at the top for the genomes of Atlantic cod (Tørresen *et al.* 2017;
gadMor2) and haddock (Tørresen *et al.* 2018; melAeg). Below, the genomic synteny
inferred from the *de novo* assemblies for all of the species included in the capture
experiment. Stippled lines indicate assembly gaps – here we assume that the
orientation of genes corresponds to the genomes of Atlantic cod and haddock. Gray
boxes indicate genes that have been identified in Baalsrud *et al.* (2017) but are absent
in the *de novo* assemblies. a) Synteny across the MN region b) Synteny across the LA
region.
780

Figure 6: Polymorphisms in the bi-directional promoter between *Hba1* and *Hbb1*
for five species in the Gadidae family.
A schematic representation of *Hba1* and *Hbb1* with the promoter region between
them. The region contains an indel polymorphism of variable length across the five
species, as indicated by gaps. For each species/variant the alignment is shown along
with amino acid substitutions at positions 55 and 62 in the translated part of the *Hbb1*
gene.

Figures

795
796   Figure 3
797

800
801 Figure 5
802

803
804   Figure 6
805

Tables

Table 1. Number of reads and bases captured and sequenced for each species, and number of utgs, largest utg and N50 in the assemblies.

Species	Latin name	Number of reads	Number of bases	Number of utgs	Largest utg (bp)	N50 (bp)
Atlantic cod	*Gadus morhua*	73005	217252583	278	79 020	7 728
Haddock	*Melanogrammus aeglefinus*	35573	107839552	227	52 433	7 227
Silvery pout	*Gadiculus argenteus*	69775	212519845	410	35 801	7 098
Cusk	*Brosme brosme*	55348	175883008	394	64 145	7 322
Burbot	*Lota lota*	56155	165360828	205	70 602	8 055
European hake	*Merluccius merluccius*	65661	180558336	311	31 558	6 523
Marbled moray cod	*Muraenolepus marmoratus*	52076	148100933	455	30 019	6 632
Roughhead grenadier	*Macrourus berglax*	46195	129085001	325	35 216	7122