Long-read sequence capture elucidates the evolution of the hemoglobin gene clusters in codfishes

Siv Nam Khang Hoff*[a], Helle T. Baalsrud*[a‡], Ave Tooming-Klunderud[a], Morten Skage[a], Todd Richmond[b], Gregor Obernosterer[c], Reza Shirzadi[d], Ole Kristian Tørresen[a], Kjetill S. Jakobsen[a] and Sissel Jentoft[a‡]

[a] Centre for Ecological and Evolutionary Synthesis, Department of Biosciences, University of Oslo, Oslo, Norway
[b] Roche NimbleGen Inc., Madison, WI, USA
[c] Roche Diagnostics, Mannheim, Germany
[d] Roche Diagnostics, Oslo, Norway

* these authors contributed equally to this work

‡ Correspondence to: h.t.baalsrud@ibv.uio.no, sissel.jentoft@ibv.uio.no

Abstract

Combining high-throughput sequencing with targeted sequence capture has become an attractive tool to study specific genomic regions of interest. Most studies have so far focused on the exome using short-read technology. This approach does not capture intergenic regions needed to reconstruct genomic organization, including regulatory regions and gene synteny. In this study, we demonstrate the power of combining targeted sequence capture with long-read sequencing technology, leading to the successful sequencing and assembling of the two hemoglobin (Hb) gene clusters LA (~100kb) and MN (~200kb) across eight species of codfishes that are separated by up to 70 million years of evolution. The highly continuous assemblies – capturing both intergenic and coding sequences – revealed overall conserved genetic organization and synteny of the *Hb* genes within this lineage, yet with several, lineage-specific gene duplications. Moreover, for some of the species examined we identified amino acid substitutions at two sites in the *Hbb1* gene as well as length polymorphisms in its regulatory region, which has previously been linked to temperature adaptation in Atlantic cod populations. Taken together, our study highlights the efficiency of targeted long-read capture for comparative genomic studies by shedding light on the evolutionary history of the *Hb* gene family across the highly divergent group of codfishes.

Article summary

Hemoglobins (Hbs) are key respiratory proteins in most vertebrates. In fishes, Hbs are shown to be of great importance for ecological adaptation, as environmental factors including temperature, directly influences the solubility of O_2 in surrounding waters as well as the ability of Hb to bind O_2 at respiratory surfaces.

We here combine targeted sequence capture and long-read sequencing to reconstruct and resolve the organization of *Hb* genes and their flanking genes in a selection of codfishes, inhabiting different environmental conditions. Our results shed light on the evolutionary history of *Hb* genes across species separated up to 70 million years of evolution, revealing genetic variations possibly linked to thermal adaptation.

Introduction

The rapid advancement of high-throughput sequencing has over the last decade
revolutionized genomic research with the increasing numbers of whole genome
resources available for multiple vertebrate species, including the diverse group of
teleost fishes (Volff 2005; Ellegren 2014; Goodwin *et al.* 2016; Malmstrøm *et al.* 2016;
2017). However, whole genome sequencing (WGS) and generation of high quality
genome assemblies are still considered costly and time-consuming (Jones and Good
2015). For investigations concerning specific genomic regions there is no need for
complete genome information, which has spurred the development of reduction
complexity approaches such as targeted sequence capture (Teer *et al.* 2010; Grover *et*
al. 2012; Samorodnitsky *et al.* 2015). The basic idea of targeted sequence capture
involves design of specific probes covering the particular genomic area of interest
generating an enriched coverage of the targeted sequences (Turner *et al.* 2009; Grover
et al. 2012). Most studies using targeted sequence capture have to a large extent been
directed towards the exome, often supported by the existence of a reference genome
(Ng *et al.* 2009; Broeckx *et al.* 2014; Yoshihara *et al.* 2016), or transcriptome assemblies
(Syring *et al.* 2016). Recent reports have, however, been focusing on off-target
sequences in noncoding regions (Guo *et al.* 2012; Samuels *et al.* 2013; Syring *et al.*
2016; Yoshihara *et al.* 2016; Morin *et al.* 2016), as they may contain crucial regulatory
elements varying in sequence and length between populations or species, and could
be of functional and evolutionary importance (Woolfe *et al.* 2004; Patrushev and
Kovalenko 2014).

To our knowledge, sequence capture studies have so far been based on short-
read sequencing technology (George *et al.* 2011; Samorodnitsky *et al.* 2015; Li *et al.*
2015; Bragg *et al.* 2016), and construction of continuous sequences enabling resolution
of gene organization have therefore not been possible. Comparative genetic studies
of gene organization or synteny requires longer, more continuous stretches of DNA
containing more than one gene (Huddleston *et al.* 2014). By its ability to span long
stretches of repeats, long-read sequencing technology has been successfully applied
to improve genome assembly statistics and generation of highly continuous genome
assemblies for a growing number of species (English *et al.* 2012; Kim *et al.* 2014;
Bickhart *et al.* 2017; Korlach *et al.* 2017; Tørresen *et al.* 2017a; Tørresen *et al.* 2017b). For
example, incorporation of long PacBio reads resulted in a significantly improved
version of the Atlantic cod (*Gadus morhua*) genome assembly, i.e. a 50-fold increase in
sequence continuity and a 15-fold reduction in the proportion of gaps (Tørresen *et al.*
2017b). Correspondingly, utilizing long-read sequencing technology in combination

with targeted capture could yield longer continuous assemblies of specific genomic

regions of interest, allowing in-depth comparative genetic studies, even in species

where a reference genome is not available.

In fishes, the hemoglobin (*Hb*) gene family, encoding the protein subunits Hba

and Hbb, has shown to be of importance for ecological adaptation, as environmental

factors such as temperature directly influences the ability of Hb to bind O_2 at

respiratory surfaces and its subsequent release to tissues (Wells 2005). In a recent

report, a full characterization of the *Hb* gene repertoire – using comparative

genomics analysis – uncovered a remarkably high *Hb* gene copy variation within the

codfishes (Baalsrud *et al.* 2017). Notably, a negative correlation between the number

of *Hb* genes and depth of which the species occur was observed, suggesting that the

more variable environment in epipelagic waters have facilitated a larger and more

diverse *Hb* gene repertoire, which was supported by evidence of diversifying

selection (Baalsrud *et al.* 2017). Furthermore, in Atlantic cod, two tightly linked

polymorphisms at amino acid positions 55 and 62 of the Hbb1-globin – suggested to

be associated with thermal adaptation – exhibit a latitudinal cline in allele frequency

in populations inhabiting varying temperature and oxygen regimes in the North

Atlantic and Baltic Sea (Andersen *et al.* 2009). Populations found in the southern

regions display the Hbb1-1 variant (Met55Lys62), whereas more northern

populations largely display the Hbb1-2 variant (Val55Ala62) (Andersen *et al.* 2009).

The Hbb1-1 variant has been shown to be insensitive to temperature whereas Hbb1-2

is temperature dependent with a higher O_2 affinity than Hbb1-1 at colder

temperatures (Karpov and Novikov 1980; Andersen *et al.* 2009), however, this has

been questioned by (Barlow *et al.* 2017). Additionally, an indel polymorphism within

the promoter of the *Hbb1* gene has been reported to be in linkage disequilibrium with

the above-mentioned polymorphisms (Star *et al.* 2011). Examination of multiple

Atlantic cod populations uncovered that a longer promoter variant is associated with

Hbb1-2 and found to up-regulate its gene expression at higher temperatures, i.e.

aiding in the maintenance of total oxygen-carrying capacity (Star *et al.* 2011).

In teleosts, the *Hb* genes are found to reside at two distinct genomic regions –

the MN and LA cluster. Earlier reports have shown that there is a high evolutionary

turnover of *Hb* genes across teleosts, with lineage-specific duplications and losses,

which is in stark contrast to genes flanking the *Hb* genes, where the synteny is highly

conserved (Quinn *et al.* 2010; Opazo *et al.* 2013; Feng *et al.* 2014). In this study, the

overall goal was to elucidate the evolutionary past of the *Hb* clusters – including *Hb*

genes, flanking genes and intergenic sequences – within the phylogenetically diverse

group of codfishes (Gadiformes) by taking the advantage of long read sequencing

technology combined with targeted sequence capture. Eight codfish species were
carefully selected on the basis of both phylogenetic and habitat divergence, implying
that they are exposed to a variety of environmental factors as well as displaying
distinct life-history traits. A highly continuous genome assembly of Atlantic cod
(Tørresen *et al.* 2017b) as well as low coverage draft genome assemblies of all eight
species (Malmstrøm *et al.* 2017) were used in the design of the probes covering the
genomic regions of interest. To enable targeted sequence capture for PacBio RSII
sequencing, we modified the standard protocol for sequence capture offered by
NimbleGen, i.e. the SeqCap EZ (Roche NimbleGen), as well as generating custom-
made barcodes. This combined approach resulted in successful capturing and
assembling of the two *Hb* gene clusters across the codfishes examined. The
generation of highly continuous assemblies – for most of the species – enabled
reconstruction of micro-synteny revealing lineage-specific gene duplications and
identification of a relatively large and inter-species variable indel located in the
promoter region between the *Hbb1* and *Hba1* genes.
Our study demonstrates that this approach, combining sequence capture
technology with long-read sequencing is a highly efficient and versatile method to
investigate specific genomic regions of interest – with respect to micro-synteny,
regulatory regions and genetic organization – across distantly related species where
genome sequences are lacking.

Results

Capture and *de novo* assembly of the target regions

The capture probe design (workflow schematically shown in Figure 1) resulted in a
total of 7057 probes based on the target region in Atlantic cod, covering 337 kbp of
sequence. 26774 probes were designed for the additional codfishes, covering in total
an area of 1.82 Mbp of target sequence. The target region and the *Hb* gene clusters
were successfully captured and enriched for eight codfishes; Atlantic cod (*Gadus*
morhua), haddock (*Melanogrammus aeglefinus)*, silvery pout (*Gadiculus argenteus*), cusk
(*Brosme brosme*), burbot (*Lota lota*), European hake (*Merluccius merluccius*), marbled
moray cod (*Muraenolepus marmoratus*), and roughhead grenadier (*Macrourus berglax*),
with number of reads spanning from 35573 to 73005 (Table 1). The average read
length was 3032 bp, varying from 2836 bp in European hake to 3265 bp in burbot,
resulting in the capture of an average of 16.71 Mbp per species (Table 1). By mapping
reads back to the capture target region we found that the average mapping depth
was variable across the target region for all species (Figure 2 and 3). Because of the
skewed distribution of mapping depth, we also calculated median depth, which was,
as expected, the highest for Atlantic cod at 242x (Table S1). The median mapping
depth was consistently high for most of the other species as well, with the lowest for
roughhead grenadier (12x). Both median and average depths for the MN region are
persistently higher than for the LA region for all species, with the exception of silvery
pout (Table S1). Furthermore, positions with high degree of mapping corresponded
to the location of the genes used in the design of the capture probes across all species
(Figure 2 and Figure 3). The percentage of reads mapping to the target region ranged
from 25-43%, however, the percentage of the target region covered by reads ranged
from 53-100% with five species having more than 90% of the target region covered by
reads (Figure 4c and Table S1).

To address factors influencing capture success we compared various capture
statistics to overall genomic divergence between the Atlantic cod genome and
independent WGS data for each species from (Malmstrøm *et al.* 2017) (Table S1). We
found a strong negative correlation between genomic divergence to Atlantic cod and
median mapping depth against the target region (r=-0.90, Figure 4a), percent of reads
mapped to the target region (r=-0.90, Figure 4b), and percentage of reads mapped to
the target region (r=-0.84, Figure 4c).

We constructed *de novo* assemblies with quite consistent assembly statistics
across species. Contig N50 ranged from 8055 bp in burbot to 6523 bp in European
hake and the total number of contigs varied from 205 in burbot to 455 in marbled

moray cod. However, there was some variation in the size of the largest contig,
which ranged from 79 kbp in Atlantic cod to 30 kbp in marbled moray cod (Table 1).
To evaluate whether the assemblies represent the actual target regions we mapped
the *de novo* assemblies for each species to the target region in Atlantic cod, for which
the capture design is largely based upon (Figure 2 and 3). As expected, the
assemblies corresponded to the regions with high coverage of reads, i.e. the areas of
the target region containing genes included in the probe design.

Synteny of the *Hb* gene regions

Our capture design combined with long-read PacBio sequencing allowed us to
reconstruct micro-synteny of the MN and LA regions for Atlantic cod, haddock,
silvery pout, cusk, burbot, European hake, marbled moray cod and roughhead
grenadier (Figure 5). From the *de novo* assemblies, we were able to identify the
majority of the *Hb* genes and all of the flanking genes, which show that our capture
design was successful. However, the degree of continuity varied in the different
assemblies. In Atlantic cod, haddock, silvery pout, cusk, burbot and European hake
we could infer micro-synteny revealing that *Hb* and their flanking genes organization
largely followed what has previously been reported for Atlantic cod (Figure 5) (Star
et al. 2011). We found *Hbb4* only to be present in Atlantic cod (Figure 5b), which is in
line with (Baalsrud *et al.* 2017). Furthermore, the *de novo* assemblies confirmed a
linage-specific duplication of *Hbb2* in the roughhead grenadier (Baalsrud *et al.* 2017).
Additionally, we identified a complete *Hba4*-like gene in the assembly of the marbled
moray cod, not earlier identified in this species. However, the *Hba4*-like gene in
marbled moray cod is likely a pseudogene due to a frameshift mutation causing
multiple stop codons. Furthermore, we were able to identify most of the *Hb* genes
reported in the recent study of (Baalsrud *et al.* 2017), however, a few are missing from
our dataset (Figure 5a and b). Pairwise sequence alignment of these paralogous *Hb*
genes from (Baalsrud *et al.* 2017) revealed sequence identities up to 98 % (Table S2).

Target region in the haddock and Atlantic cod genome assemblies

As a proof of concept, we reconstructed synteny of the target region in the most
recent genome assemblies of Atlantic cod (gadMor2 (Tørresen *et al.* 2017b)) and
haddock (melAeg (Tørresen *et al.* 2017a)). In Atlantic cod, the MN region is located
on linkage group 2 (Figure 5a) and LA on linkage group 18 (Figure 5b), in haddock
MN is located on scaffold MeA_20160214_scaffold_771 (Figure 5a) and LA on
scaffold MeA_20160214_scaffold_1676 (Figure 5b). The overall synteny in Atlantic
cod was congruent with (Wetten *et al.* 2010) except for the relative direction of the
genes *foxj1a* and *rhbdf1*. Furthermore, the organization of *Hb*s and their flanking

genes in the genome assembly of haddock is conserved compared to Atlantic cod
with the exception of *Hbb4* in the MN region, which is absent in haddock (Figure 5).

Repetitive sequences in the in the *Hb* gene regions

Quantifying the amount of repetitive sequences in the target region(s) was only
possible for Atlantic cod (gadMor2) and haddock (melAeg), for which high-quality
genome assemblies exist. The amount of repetitive sequences in the target region
differed between the MN cluster and the LA cluster in Atlantic cod. The MN region
(214 kb) contained a total of 10.7% repeated sequences, including 1.0% retro-
elements, 1.3% transposons, 5.8% simple repeats, and 2.6% of various low complexity
and unclassified repeated sequences (Table S3). In comparison, in the smaller LA
region (123 kb) the proportion of repeated sequences was twice as high (20.3%),
which comprised of 2.8% retro-elements, 2.4% transposons, 13.8% simple repeats,
and 1.3% of various low complexity and unclassified repeated sequences.
Furthermore, the orthologous target regions in haddock followed the same pattern.
The MN region contained 16.3 % repeated sequences, in contrast to 19.8 % found in
the LA region (Table S3).

Insertions and deletions in the promoter region of *Hba1 – Hbb1*

The previously shown 73 bp indel in the bi-directional promoter region of *Hba1* and
Hbb1 – discerning the cold-adapted migratory Northeast Artic cod (NEAC) from the
more temperate-adapted southern Norwegian coastal cod (NCC) (Star *et al.* 2011) –
was confirmed by the improved version of the NEAC assembly (gadMor2). The
continuity of our capture assemblies (Figure 5) enabled location of the orthologous
captured regions in haddock, silvery pout and cusk. In each of the species an indel of
variable length were identified (Figure 6). Compared to the long promoter variant –
found to be linked with the *Hbb1-2* in Atlantic cod – the indel is shorter in the other
species by 11 bp in haddock, 22 bp in silvery pout and 56 bp in cusk (Figure 6).
Although the indels are varying in length, the conserved flanking sequences in the
alignment clearly show that they represent orthologous regions. Moreover, we found
the amino acid positions at 55 and 62 of the *Hbb1* gene to vary between species;
Haddock has Val55-Lys62, silvery pout has Met55-Gln62, while cusk has Met55-
Lys62 similarly to NEAC (Figure 6). Additionally, we investigated amino acid
positions 55 and 62 in the *Hbb1* gene across a number additional codfish species for
which we have available gene sequences from (Baalsrud *et al.* 2017), revealing these
sites to be variable across this lineage (Table S4). Ancestral reconstruction of *Hbb1*
demonstrated that the ancestral state in position 55 was Met in codfishes, and in

position 62 was Lys in all codfishes except *Bregmaceros cantori* (Supplementary
Figures S1 and S2).

Discussion

Capture of *Hb* gene clusters with 70 million years divergence time reveal conserved synteny and lineage-specific *Hb* duplications

We here demonstrate a successful in-solution targeted sequence capture and assembling of coding and noncoding sequences of the *Hb* clusters from codfish species separated by up to 70 million years (My) of evolution. Two features make our approach unique from earlier studies. First, the target regions consisted of both coding and noncoding genomic sequences. Second, we designed capture of large fragments – combined with development of custom-made probes – in order to utilize the long-read PacBio sequencing platform. This is in contrast to current targeted capture sequencing protocols that are based on short-read sequencing technologies (George *et al.* 2011; Mascher *et al.* 2013).

The organization and orientation of the *Hb* flanking genes that we identified were conserved across all species (Figure 5a and b). However, in concordance with earlier studies of the *Hb* region, we found significant variation in copy numbers of the *Hb* genes, with linage specific duplications and losses (Star *et al.* 2011; Opazo *et al.* 2013; Feng *et al.* 2014; Baalsrud *et al.* 2017). We only found *Hbb4* in Atlantic cod, supporting earlier studies showing that *Hbb4* is the result of a recent duplication in this species (Borza *et al.* 2009; Baalsrud *et al.* 2017). Interestingly, the presence of two copies of *Hbb2* on the same contig in the roughhead grenadier *de novo* assembly confirmed a lineage specific gene duplication of *Hbb2*, which was found in a recent study of *Hb*s in codfishes (Baalsrud *et al.* 2017). Additionally, a copy of the *Hba4* was found in the *de novo* assembly of the marbled moray cod not found in (Baalsrud *et al.* 2017). The presence of a frame-shifting mutation that is causing multiple stop codons indicated that this *Hba4* gene is most likely a pseudogene. *Hba4* is also a pseudogene in the closely related species *Mora moro*, *Trachyrincus scabrus*, *T. murrayi* and *Melanonus zugmayeri* (Baalsrud *et al.* 2017). Although we identified most of the *Hb* genes from (Baalsrud *et al.* 2017), a few were absent from this dataset (Figure 5a and b), which we suspect may be due to collapse of paralogous *Hb* genes, as they may have as high as 98% sequence identity (Table S2).

Length variation in the bi-directional *Hba1-Hbb1* promoter within the codfishes

The discovery of a promoter of variable length between *Hba1* and *Hbb1* in different species (Figure 6) was concordant with earlier findings of length variation in the homologous region in different populations of Atlantic cod (Star *et al.* 2011). The

migratory NEAC population has been shown to harbor the 73 bp longer variant at a
higher frequency compared to coastal cod populations (see Figure 6 and (Star *et al.*
2011)). Interestingly, we found relatively long promoters with high sequence
similarity to the NEAC indel in haddock and silvery pout. In contrast, cusk
displayed a relatively short promoter, however, still 17 bp longer than in NCC
(Figure 6). Furthermore, we found the amino acid positions 55 and 62 in *Hbb1*,
known to be polymorphic in Atlantic cod, to be variable across all codfishes included
in this study (Figure 6). Investigations of the same positions in a number additional
codfishes for which we have available gene sequences (Baalsrud *et al.* 2017), revealed
that these positions are highly variable across this linage (Table S2). Notably, the
most likely ancestral state of codfish *Hbb1* is Met55Lys62 (Supplementary Figures S1
and S2). Cusk and the coastal/southern Atlantic cod thus both display the ancestral
state as well as a short promoter, although the cusk promoter was 17 bp longer
(Figure 6). Collectively, these results suggest two different scenarios for promoter
length evolution. Scenario 1: The short promoter represents the ancestral state of the
Gadidae-family (including cusk and Atlantic cod; see (Malmstrøm *et al.* 2016)) and
that silvery pout and some populations of Atlantic cod have evolved a longer
promoter. Scenario 2: The long promoter is the ancestral state with independent
deletions of variable lengths in cusk, silvery pout, haddock and costal/southern
Atlantic cod (*Hbb1-1*). To disentangle this, we would need to obtain promoter
sequences from additional gadiform species. Regardless, the short-long promoter
polymorphism has been maintained throughout speciation events based on the
presence of both variants in Atlantic cod. Moreover, in both scenarios, cusk and
Atlantic cod (*Hbb1-1*) have maintained the ancestral Met55Lys62, while silvery pout,
haddock and Atlantic cod (*Hbb1-2*) have acquired substitutions at these positions due
to similar selection pressures or genetic drift. In this regard, it could be mentioned
that the NEAC, haddock and silvery pout display migratory behavior (e.g. diurnally
feeding movements as well as seasonal spawning migrations) compared to the more
stationary cusk and coastal cod (Eschemeyer and Fricke 2017) which could mean that
they have a higher O_2 demand and are exposed to greater temperature variation,
which in turn has selected for a temperature-dependent long promoter. Furthermore,
given that promoter length and positions 55/62 at *Hbb1* are important genetic
components of temperature adaptation in Atlantic cod populations (Star *et al.* 2011),
they most likely play a role in temperature adaptation in the other codfishes.

Assembly success affected by probe design and repeat content

In some species, nearly the complete target region is assembled in large contigs
containing multiple genes including cusk, whereas in other species such as the more

distantly related roughhead grenadier, the cluster is more fragmented (Figure 5). In all species, the areas of the target regions that harbor genes of which probes are designed for, as well as any areas containing repeated sequences, have very high depths in comparison to the areas of intergenic sequences (Figure 2 and 3). This poses a challenge for the assembly software, which is based in the assumption of uniform depth over the sequencing data (Miller *et al.* 2010).

Overall, the MN cluster seems to be more successfully assembled than the LA cluster, which is more fragmented (Figure 5). Differences in assembly completeness between the two regions might be a result of several factors. Firstly, the MN region has more flanking genes in closer proximity to the *Hb* region, which results in a higher density of probes. Secondly, the overall repeat content of the LA region is one order of magnitude larger than in the MN region, largely due to the larger proportion of simple repeats. Repeat content is a major interference in capture experiments because unwanted repetitive DNA may be enriched for, especially if there are repeated sequences included in the probe design. Furthermore, if probes were not completely covered by target DNA they get single-stranded sticky ends that can hybridize to repetitive or other non-target DNA (Newman and Austin 2016). Lastly, unless there were some longer reads that bridged such areas, this would in turn have led to gaps in the downstream *de novo* assemblies. Following that the assembly success was possibly a result of read length, we reason that a future increase of the average read length from 3 kbp to 5-10 kbp, would be sufficient to substantially increase the completeness of the assemblies. Due to the current circular consensus (CCS) PacBio sequencing technology, however, which is a trade-off between accuracy and length of reads, longer reads with sufficient accuracy are not feasible.

Long-read sequencing capture across species harbors new potential for comparative genomic studies

The number of reads mapping to the target region was in the range of 23-43%, which may seem low compared with other capture studies. For instance, a whole exome capture study on humans reported 56.1% of reads mapped to the target region (Guo *et al.* 2012) and a similar study in rats reported to have 78.3% of reads on target (Yoshihara *et al.* 2016). In contrast to our study however, these capture experiments enriched either the exome or ultra-conserved elements within a single species. Furthermore, we were able to cover up to 98% of the target region with >10 reads across species (Table S1) which is similar to what mentioned experiments within human and rat exomes reported (Guo *et al.* 2012; Yoshihara *et al.* 2016) and the main difference is the higher percentage of non-target sequences in our study.

We were able to capture complete genes for species with 70 My divergence time from the Atlantic cod (Figure 5). As expected, we found that capture success declines with increased sequence divergence between the reference genome of which we chiefly based our capture probes and the genomes of the included codfishes (Figure 4). It has been reported that orthologous exons were successfully captured in highly divergent frog species (with 200 My of separation), nevertheless the capture success greatly decreased with increased evolutionary distance (Hedtke *et al.* 2013). Similarly, it has been demonstrated that it is possible to capture >97% of orthologous sequences in four species of primates that diverged from humans 40 My ago, using probes entirely based on the human exome (George *et al.* 2011). Further, exomes were effectively captured from skink species that diverged up to 80 My from the reference, yet reporting a substantial decline in capture efficiency for sequences >10 % different from the reference species (Bragg *et al.* 2016). Our study stands out from previous capture experiments because intergenic, noncoding sequences in addition to genes were captured. Efficient capture of intergenic sequences requires less divergence time, as these regions usually evolve faster than genes (Koonin and Wolf 2010). Thus, the most distantly related species from Atlantic cod for which we captured both coding and noncoding sequences was burbot, which diverged from Atlantic cod 46 My (Figure 5). We argue, in line with (Schott *et al.* 2017), that sequence divergence may be a more exact predictor of capture success than evolutionary distance, as the sequence capture process is mainly influenced by the difference between the probe sequence and the target sequence. European hake, marbled moray cod and roughhead grenadier all diverged from cod about 70 My ago, however, the European hake *Hb* regions was more successfully captured and assembled (Table 1; Figure 2). This could be due to European hake having a lower genome-wide divergence to Atlantic cod than marbled moray cod and roughhead grenadier (809k vs 879k and 907k SNPs; Table S1).

Finally, it should be mentioned that cusk – which diverged from Atlantic cod 39 My ago – was added to the experimental design after the species-specific probes were generated. Thus, the successful capture of cusk was therefore solely based on cross-species target enrichment, and could most likely been further improved if species-specific probes for this species have been included.

Concluding remarks and future perspectives

Here, we have successfully demonstrated that combining targeted sequence capture with long-read sequencing technology is as an efficient approach to obtain high quality sequence data of a specific genomic region, including both coding and

noncoding sequences, across evolutionary distant species. We show that genome-
wide divergence is of importance for capture success across species. Furthermore, the
use of long-read sequencing augmented the *de novo* assembly of regions containing
repeated sequences that would otherwise fragment assemblies based on short-read
sequencing. This is crucial for capturing complete intergenic sequences that may be
highly divergent compared to genic regions even among fairly closely related
species. Given the rapid development in sequencing technologies future methods
will enable read-through of repeated regions and thus further increase the
completeness of assemblies. Moreover, a less stringent hybridization protocol should
make it possible to capture sequences across even deeper evolutionary time. In sum,
our approach has the potential of enhancing comparative genomic studies of
continuous genic and intergenic regions between any eukaryotic species-group
where genomic resources are scarce.

Material and methods

Defining target region and probe design

The probe design was chiefly based on the high-quality genome of Atlantic cod, known as gadMor2 (Tørresen *et al.* 2017b). In addition, species-specific probes were designed based on low-coverage assembled genomes (Malmstrøm *et al.* 2016) for ten selected species representing six families in the Gadiformes order. These species were Atlantic cod (*Gadus morhua*), Alaskan Pollock (*Gadus chalcogrammus*), polar cod (*Boreogadus saida*), haddock (*Melanogrammus aeglefinus*), Silvery pout (*Gadiculus argenteus*), burbot (*Lota lota*), European hake (*Merluccius merluccius*), roughhead grenadier (*Macrourus berglax*), roughsnout grenadier (*Trachyrincus scabrus*) and marbled moray cod (*Muraenolepus marmoratus*).

To retrieve relevant sequence data for the probe design, the MN and LA *Hb* regions were extracted from gadMor2 (Figure 1). These sequences, hereby known as the target region, were then used queries in BLAST (Altschul *et al.* 1990) searches with an E-value threshold of <0.1 against the genome assembly data of all ten species.

In total, 5604 sequences from the chosen species were supplied to NimbleGen probe design. Protein coding genes from the ENSEMBL database were used to define the regions to be tiled in the probe design (Table S5) within the target region of the Atlantic cod, and the unitigs for each of the additional codfishes.

NimbleGen SeqCap EZ capture probes were designed by NimbleGen (Roche, Madison, USA) using a proprietary design algorithm. NimbleGen offers an in-solution sequence capture protocol, which includes custom made probes. Uniquely, the capture probes from NimbleGen are tiled to overlap the target area. 50 – 100 bp (average 75 bp) probes where designed tiled over the target region (subset of gadMor2) resulting in each base, on average, being covered by two probes. Additionally, raw reads from Illumina sequencing from (Malmstrøm *et al.* 2017) were used for each species to estimate repetitive sequences in each of the species' genomes, aiming to discard probes containing any repeats.

Sample collection and DNA extraction

Our goal working with animals is always to limit any harmful effects of our research on populations and individuals. Whenever possible we try to avoid animals being euthanized to serve our scientific purpose alone by collaborating with commercial fisheries or museums. The tissue samples used in this study are either from commercially fished individuals intended for human consumption or museum specimen. The commercially caught fish were immediately stunned, by bleeding

following standard procedures by a local fisherman. There is no specific legislation
applicable to this manner of sampling in Norway, however it is in accordance with
the guidelines set by the 'Norwegian consensus platform for replacement, reduction
and refinement of animal experiments' (www.norecopa.no).
DNA was extracted from tissue samples using High Salt DNA Extraction
method by Phill Watts (https://www.liverpool.ac.uk/~kempsj/IsolationofDNA.pdf ,
last day accessed: December 2017). The concentration and purity of the DNA samples
were quantified using NanoDrop (Thermo Scientific, Thermo Fisher Scientific,
Waltham, MA, USA) and a Qubit fluorometer (Invitrogen, Thermo Fisher Scientific,
Waltham, MA, USA). Due to poor DNA quality, three species included in the probe
design; Alaskan Pollock, polar cod and roughsnout grenadier were excluded from
further analysis. In total, eight species were sequenced; seven of these species were
included in the probe design and one closely related species (cusk, *Brosme brosme*),
which serves as a cross species capture experiment without species-specific probes.

Capture, library preparation and sequencing

The sequencing libraries were prepared following a modified Pacific Biosciences
SeqCap EZ protocol. As multiplexing of the samples before capture was required,
barcodes were designed at the Norwegian Sequencing Centre
(http://www.sequencing.uio.no) using guidelines from Pacific Biosciences
(Supplementary Materials and methods). Genomic DNA was sheared to 5 kb
fragments using MegaRuptor (Diagenode, Seraing (Ougrée), Belgium). Due to poorer
DNA quality, fragmenting was not done for European hake. For this sample together
with fragmented DNA from roughhead grenadier, short fragments were removed
using BluePippin (Sage Science, Beverly, MA, USA) before library preparation.
Illumina libraries were prepared using KAPA Hyper Prep kit (Kapa Biosystems,
Wilmington, MA, USA) and barcoded using different Illumina barcodes. PacBio
barcodes were implemented during pre-capture amplification of libraries. After
amplification, fragment length distribution was evaluated using Bioanalyzer (Agilent
Technologies, Santa Clara, CA, USA) and samples were pooled in equimolar ratio.
During hybridization, SeqCap EZ Developer Reagent (universal repeat blocker for
use on vertebrate genomes) and oligos corresponding to Illumina and PacBio
barcodes were used for blocking. Captured gDNA was amplified to ensure that
sufficient amount of DNA was available for PacBio library preparation. Size selection
of the libraries was performed using BluePippin. Final libraries were quality checked
using Bioanalyzer and Qubit fluorometer (Invitrogen, Thermo Fisher Scientific,
Waltham, MA, USA) and sequenced on RS II instrument (PacBio, Menlo Park, CA,

USA) using P6-C4 chemistry with 360 minutes movie time. In total, 9 SMRT cells
were used for sequencing.

De novo assemblies

Reads were filtered and de-multiplexed using the 'RS_reads of insert.1' pipeline on
SMRT Portal (SMRT Analysis version smrtanalysis_2.3.0.140936.p2.144836). Each set
of reads corresponding to a given species was crossed-checked with their respective
six-nucleotide Illumina adapter. Reads containing an incorrect Illumina adapter were
removed. Adapter sequences were then trimmed using the application Prinseq-lite
v0.20.4 (Schmieder and Edwards 2011). The trimmed reads were assembled *de novo*
using Canu v1.4 +155 changes (r8150 c0a988b6a106c27c6f993dfe586d2336282336a6)
(Berlin *et al.* 2015). The Canu software is optimized for assembling single molecule
high noise sequence data. We specified genome size as the size of the target region
(300 kbp). Additionally, we ran PBJelly (English *et al.* 2012) on the Canu *de novo*
assemblies, using the raw reads to possible bridge gaps between scaffolds, settings
given in Supplementary Materials and Methods.
We assessed the assemblies by running Assemblathon 2 (Bradnam *et al.* 2013),
which reports assembly metrics such as the longest contig, the number of contigs,
and the N50 value. *De novo* assemblies of the MN and LA regions of Atlantic cod and
haddock were aligned and compared to their reference genomes, gadMor2 and
melAeg respectively, using BLAST and BWA v0.7.10 (Li and Durbin 2009) to
determine syntenic similarities and assembly completeness.

Estimating capture success

PacBio reads for all the species were mapped back to the Atlantic cod genome
assembly (gadMor2) in order to determine sequence capture success and target
mapping depths. Mapping was done using BWA-MEM v0.7.10 (Li and Durbin 2009).
Target-area read depth for all the species based on mapping against gadMor2, were
calculated using Samtools v1.3.1 (Li *et al.* 2009). We calculated both average and
median mapping depth against the target region as a whole and for the MN and LA
region separately. We also calculated percentage of reads that mapped to the target
region, and the percentage of the target regions covered by reads to a minimum
depth of 10x. To compare assemblies to the target region we additionally mapped the
assemblies to the target region. In order to verify the sequence capture process,
sequence data for Atlantic cod and haddock were mapped back to their reference
genomes using BWA-mem v0.7.10 (Li and Durbin 2009). The results were visualized
using Integrative Genome Viewer (Robinson *et al.* 2011).

To obtain an independent measure of divergence between species in the
capture experiment we calculated genome wide level of divergence of each species to
the reference genome of Atlantic cod using low-coverage whole-genome sequence
data from (Malmstrøm *et al.* 2017). We mapped raw reads to Atlantic cod using
BWA-MEM (Li and Durbin 2009) and called SNPs using the Freebayes variant caller
(Garrison and Marth 2017). Some species are more closely related to Atlantic cod
than others, which could introduce a bias in mapping. To avoid this, we only looked
at genomic regions where all species mapped. The number of SNPs was then used as
an estimate of genome-wide divergence of each species to Atlantic cod. We also
mapped a low-coverage genome of Atlantic cod to the Atlantic cod reference genome
as a control.
In pursuance of factors explaining capture success we tested for correlations
and plotted the relationship between the genome wide level of divergence and the
following variables; median mapping depth against the target region (for total, LA
and MN, respectively); percentage of reads that mapped to the target region; and the
percentage of the target region covered by reads. All tests and plots were done using
R version 3.2.5(Team 2013).
Assembly continuity is very often hampered by the presence of repeats, which
create gaps. We therefore quantified repeat-content in the target region extracted
from gadMor2 and orthologous regions in haddock using Repeatmasker Open 3.0
(Smit *et al.* 2010) for the MN region and the LA region separately.

Identifying gene location and synteny

In order to identify the genes of interest and their location in the assembly we used
local sequence alignment algorithm BLAST v2.4.0 (Altschul *et al.* 1990) with protein
sequences of the genes of interest (Table S5) as queries. tblastn was used with an e-
value of 0.1. Investigation of *Hbb1-Hba1* promoter region was done for four species,
Atlantic cod, haddock, silvery cod and cusk. Sequences were aligned with ClustalW
default settings using MEGA7 (Kumar *et al.* 2016). Ancestral sequence reconstruction
was carried out for *Hbb-1* gene sequences from 24 species of codfishes from (Baalsrud
et al. 2017) using a maximum likelihood method implemented in MEGA7 (Kumar *et
al.* 2016).
Additionally, we estimated sequence identity using EMBOSS Needle (Rice *et
al.* 2000) with default settings, between *Hbb* gene sequences from (Baalsrud *et al.*
2017) that where missing and present in the *de novo* assemblies to evaluate similarity
(Table S2).

Acknowledgements

We would like to thank Marianne H. Selander Hansen and Alexander J. Nederbragt for help with the initial design of this project. All computational work was performed on the Abel Supercomputing Cluster (Norwegian Metacenter for High-Performance Computing (NOTUR) and the University of Oslo), operated by the Research Computing Services group at USIT, the University of Oslo IT Department. Sequencing library creation and high-throughput sequencing were carried out at the Norwegian Sequencing Centre (NSC), University of Oslo, Norway. This work was funded by a grant awarded to K.S.J. from the Research Council of Norway (RCN grant 222378).

Author contributions

H.T.B. and S.J. initially conceived and designed the study, with input from S.N.K.H, A.T.-K., M.S., G.O., R.S., and K.S.J. Tissue samples were provided by S.J. and H.T.B. Probe design was carried out by T.R. with assistance from S.N.K.H and H.T.B. DNA extraction and sequence library preparation was performed by S.N.K.H and A.T.-K, respectively. Sequence capture was carried out by S.N.K.H, A.T.-K., M.S. and G.O. Filtering, mapping of sequences and *de novo* assemblies was done by S.N.K.H., assisted by O.K.T and H.T.B. Annotation of genes, synteny analyses, statistical analyses and construction of all figures and tables was done by S.N.K.H and H.T.B. The manuscript was written by S.N.K.H and H.T.B. with input from S.J. and K.S.J.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All reads and assemblies (unitigs) reported on here, and the target region, subset of gadMor2 have been deposited at figshare under doi/xxx.

References

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman, 1990 Basic local alignment search tool. J. Mol. Biol. 215: 403–410.

Andersen, Ø., O. F. Wetten, M. C. De Rosa, C. Andre, C. Carelli Alinovi *et al.*, 2009 Haemoglobin polymorphisms affect the oxygen-binding properties in Atlantic cod populations. Proc. Biol. Sci. 276: 833–841.

Baalsrud, H. T., K. L. Voje, O. K. Tørresen, M. H. Solbakken, M. Matschiner *et al.*, 2017 Evolution of Hemoglobin Genes in Codfishes Influenced by Ocean Depth. Sci Rep 7: 168–10.

Barlow, S. L., J. Metcalfe, D. A. Righton, and M. Berenbrink, 2017 Life on the edge: O 2binding in Atlantic cod red blood cells near their southern distribution limit is not sensitive to temperature or haemoglobin genotype. Journal of Experimental Biology 220: 414–424.

Berlin, K., S. Koren, C.-S. Chin, J. P. Drake, J. M. Landolin *et al.*, 2015 Assembling large genomes with single-molecule sequencing and locality-sensitive hashing. Nature Biotechnology 33: 623–630.

Bickhart, D. M., B. D. Rosen, S. Koren, B. L. Sayre, A. R. Hastie *et al.*, 2017 Single-molecule sequencing and chromatin conformation capture enable de novo reference assembly of the domestic goat genome. Nat Genet 431: 931–11.

Borza, T., C. Stone, A. K. Gamperl, and S. Bowman, 2009 Atlantic cod (*Gadus morhua*) hemoglobin genes: multiplicity and polymorphism. BMC Genetics 10: 51.

Bradnam, K. R., J. N. Fass, A. Alexandrov, P. Baranay, M. Bechner *et al.*, 2013 Assemblathon 2: evaluating de novo methods of genome assembly in three vertebrate species. Giga Sci 2: 10.

Bragg, J. G., S. Potter, K. Bi, and C. Moritz, 2016 Exon capture phylogenomics: efficacy across scales of divergence. Molecular Ecology Resources 16: 1059–1068.

Broeckx, B. J. G., F. Coopman, G. E. C. Verhoeven, V. Bavegems, S. De Keulenaer *et al.*, 2014 Development and performance of a targeted whole exome sequencing enrichment kit for the dog (Canis Familiaris Build 3.1). Sci Rep 4: 1522–4.

Ellegren, H., 2014 Genome sequencing and population genomics in non-model organisms. Trends in Ecology & Evolution 29: 51–63.

English, A. C., S. Richards, Y. Han, M. Wang, V. Vee *et al.*, 2012 Mind the Gap: Upgrading Genomes with Pacific Biosciences RS Long-Read Sequencing

Technology (Z. Liu, Ed.). PLoS ONE 7: e47768.

Eschemeyer, W. N., and R. Fricke, 2017 Catalog of fishes.
http://research.calacademy.org/researchichthyology/catalog/fishcatmain.asp.

Feng, J., S. Liu, X. Wang, R. Wang, J. Zhang *et al.*, 2014 Channel catfish hemoglobin
genes: Identification, phylogenetic and syntenic analysis, and specific induction
in response to heat stress. Comparative Biochemistry and Physiology Part D:
Genomics and Proteomics 9: 11–22.

Garrison, E., and G. Marth, 2017 Haplotype-based variant detection from short-read
sequencing. 1–9.

George, R. D., G. McVicker, R. Diederich, S. B. Ng, A. P. MacKenzie *et al.*, 2011 Trans
genomic capture and sequencing of primate exomes reveals new targets of
positive selection. Genome Research 21: 1686–1694.

Goodwin, S., J. D. McPherson, and W. R. McCombie, 2016 Coming of age: ten years
of next-generation sequencing technologies. Nat Rev Genet 17: 333–351.

Grover, C. E., A. Salmon, and J. F. Wendel, 2012 Targeted sequence capture as a
powerful tool for evolutionary analysis. American Journal of Botany 99: 312–319.

Guo, Y., J. Long, J. He, C.-I. Li, Q. Cai *et al.*, 2012 Exome sequencing generates high
quality data in non-target regions. BMC Genomics 13:.

Hedtke, S. M., M. J. Morgan, D. C. Cannatella, and D. M. Hillis, 2013 Targeted
Enrichment: Maximizing Orthologous Gene Comparisons across Deep
Evolutionary Time (U. Joger, Ed.). PLoS ONE 8: e67908.

Huddleston, J., S. Ranade, M. Malig, F. Antonacci, M. Chaisson *et al.*, 2014
Reconstructing complex regions of genomes using long-read sequencing
technology. Genome Research 24: 688–696.

Jones, M. R., and J. M. Good, 2015 Targeted capture in evolutionary and ecological
genomics. Molecular Ecology 25: 185–202.

Karpov, A. K., and G. G. Novikov, 1980 Hemoglobin alloforms in cod, *Gadhus morhua*
(Gadiformes, Gadidae), their functional characteristics and occurrence in
populations. Journal of Ichthyology 20: 45–50.

Kim, K. E., P. Peluso, P. Babayan, P. J. Yeadon, C. Yu *et al.*, 2014 Long-read, whole-
genome shotgun sequence data for five model organisms. Sci. Data 1: 140045–10.

Koonin, E. V., and Y. I. Wolf, 2010 Constraints and plasticity in genome and
molecular-phenome evolution. Nat Rev Genet 11: 487–498.

Korlach, J., G. Gedman, S. B. Kingan, C.-S. Chin, J. T. Howard *et al.*, 2017 Giga Sci 6:
1–16.

Kumar, S., G. Stecher, and K. Tamura, 2016 MEGA7: Molecular Evolutionary
Genetics Analysis Version 7.0 for Bigger Datasets. Molecular Biology and
Evolution 33: 1870–1874.

Li, C., S. Corrigan, L. Yang, N. Straube, M. Harris *et al.*, 2015 DNA capture reveals
transoceanic gene flow in endangered river sharks. Proc. Natl. Acad. Sci. U.S.A.
112: 13302–13307.

Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-
Wheeler transform. Bioinformatics 25: 1754–1760.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.

Malmstrøm, M., M. Matschiner, O. K. Tørresen, K. S. Jakobsen, and S. Jentoft, 2017
Whole genome sequencing data and de novo draft assemblies for 66 teleost
species. Sci. Data 4: 1–13.

Malmstrøm, M., M. Matschiner, O. K. Tørresen, B. Star, L. G. Snipen *et al.*, 2016
Evolution of the immune system influences speciation rates in teleost fishes. Nat
Genet 48: 1204–1210.

Mascher, M., G. J. Muehlbauer, D. S. Rokhsar, J. Chapman, J. Schmutz *et al.*, 2013
Anchoring and ordering NGS contig assemblies by population sequencing
(POPSEQ). Plant J 76: 718–727.

Miller, J. R., S. Koren, and G. Sutton, 2010 Assembly algorithms for next-generation
sequencing data. Genomics 95: 315–327.

Morin, A., T. Kwan, B. Ge, L. Letourneau, M. Ban *et al.*, 2016 Immunoseq: the
identification of functionally relevant variants through targeted capture and
sequencing of active regulatory regions in human immune cells. BMC Med
Genomics 9: 7–12.

Newman, C. E., and C. C. Austin, 2016 Sequence capture and next-generation
sequencing of ultraconserved elements in a large-genome salamander. Molecular
Ecology 25: 6162–6174.

Ng, S. B., E. H. Turner, P. D. Robertson, S. D. Flygare, A. W. Bigham *et al.*, 2009
Targeted capture and massively parallel sequencing of 12 human exomes. Nature
461: 272–276.

Opazo, J. C., G. T. Butts, M. F. Nery, J. F. Storz, and F. G. Hoffmann, 2013 Whole-

genome duplication and the functional diversification of teleost fish
hemoglobins. Molecular Biology and Evolution 30: 140–153.

Patrushev, L. I., and T. F. Kovalenko, 2014 Functions of noncoding sequences in
mammalian genomes. Biochemistry (Mosc) 79: 1442–1469.

Quinn, N. L., K. A. Boroevich, K. P. Lubieniecki, W. Chow, E. A. Davidson *et al.*, 2010
Genomic organization and evolution of the Atlantic salmon hemoglobin
repertoire. BMC Genomics 11: 539.

Rice, P., I. Longden, and A. Bleasby, 2000 EMBOSS: the European Molecular Biology
Open Software Suite. Trends in Genetics 16: 276–277.

Robinson, J. T., H. Thorvaldsdottir, W. Winckler, M. Guttman, E. S. Lander *et al.*, 2011
Integrative genomics viewer. Nature Biotechnology 29: 24–26.

Samorodnitsky, E., J. Datta, B. M. Jewell, R. Hagopian, J. Miya *et al.*, 2015 Comparison
of custom capture for targeted next-generation DNA sequencing. J Mol Diagn 17:
64–75.

Samuels, D. C., L. Han, J. Li, S. Quanghu, T. A. Clark *et al.*, 2013 Finding the lost
treasures in exome sequencing data. Trends in Genetics 29: 593–599.

Schmieder, R., and R. Edwards, 2011 Quality control and preprocessing of
metagenomic datasets. Bioinformatics 27: 863–864.

Schott, R. K., B. Panesar, D. C. Card, M. Preston, T. A. Castoe *et al.*, 2017 Targeted
Capture of Complete Coding Regions across Divergent Species. Genome Biology
and Evolution 9: 398–414.

Smit, A., R. Hubley, and P. Green, 2010 RepeatMasker Open 3.0.
http://www.repeatmasker.org.

Star, B., A. J. Nederbragt, S. Jentoft, U. Grimholt, M. Malmstrøm *et al.*, 2011 The
genome sequence of Atlantic cod reveals a unique immune system. Nature 477:
207–210.

Syring, J. V., J. A. Tennessen, T. N. Jennings, J. Wegrzyn, C. Scelfo-Dalbey *et al.*, 2016
Targeted Capture Sequencing in Whitebark Pine Reveals Range-Wide
Demographic and Adaptive Patterns Despite Challenges of a Large, Repetitive
Genome. Front Plant Sci 7: 484.

Team, R. C., 2013 R Core Team. R: A language and environment for statistical
computing. R Foundation for Statistical Computing, Vienna, Austria: ISBN 3-
900051-07-0, URL http://www. R-project. org/.(3.3. 1) Software Vienna, Austria: R
Foundation for Statistical Computing.

Teer, J. K., L. L. Bonnycastle, P. S. Chines, N. F. Hansen, N. Aoyama *et al.*, 2010
713  Systematic comparison of three genomic enrichment methods for massively
714  parallel DNA sequencing. Genome Research 20: 1420–1431.

715 Turner, E. H., S. B. Ng, D. A. Nickerson, and J. Shendure, 2009 Methods for Genomic
716  Partitioning. Annu. Rev. Genom. Human Genet. 10: 263–284.

717 Tørresen, O. K., M. S. O. Brieuc, M. H. Solbakken, E. Sørhus, A. J. Nederbragt *et al.*,
718  2017a Genomic architecture of codfishes featured by expansions of innate
719  immune genes and short tandem repeats. bioRxiv 1–42.

720 Tørresen, O. K., B. Star, S. Jentoft, W. B. Reinar, H. Grove *et al.*, 2017b An improved
721  genome assembly uncovers prolific tandem repeats in Atlantic cod. BMC
722  Genomics 18: 311–23.

723 Volff, J.-N., 2005 Genome evolution and biodiversity in teleost fish. Heredity 94: 280–
724  294.

725 Wells, R. M. G., 2005 Blood-Gas Transport and Hemoglobin Function in Polar Fishes:
726  Does Low Temperature Explain Physiological Characters?, pp. 281–316 in *Fish
727  Physiology*, Fish Physiology, Elsevier.

728 Wetten, O. F., A. J. Nederbragt, R. C. Wilson, K. S. Jakobsen, R. B. Edvardsen *et al.*,
729  2010 Genomic organization and gene expression of the multiple globins in
730  Atlantic cod: conservation of globin-flanking genes in chordates infers the origin
731  of the vertebrate globin clusters. BMC Evolutionary Biology 10: 315.

732 Woolfe, A., M. Goodson, D. K. Goode, P. Snell, G. K. McEwen *et al.*, 2004 Highly
733  Conserved Non-Coding Sequences Are Associated with Vertebrate Development
734  (Sean Eddy, Ed.). PLoS Biol 3: e7.

735 Yoshihara, M., D. Saito, T. Sato, O. Ohara, T. Kuramoto *et al.*, 2016 Design and
736  application of a target capture sequencing of exons and conserved non-coding
737  sequences for the rat. BMC Genomics 17: 1522–11.

Figure legends

Figure 1: Flowchart of sequence capture approach. a) Sequence data from the
Atlantic cod genome (gadMor2 (Tørresen *et al.* 2017b)) combined with gene
sequences of target genes and sequences from low coverage genomes of the
additional codfishes are combined to generate probes. b) Isolated DNA is
multiplexed with Illumina and PacBio barcodes. c) Raw reads for each species are
used to score all probes, ensuring that no repeated sequences are present. DNA
Probes are used in solution on isolated DNA for all of the included species,
hybridizing to the target sequences. Target sequences are then captures and
sequences on the PacBio RSII sequencing platform. d) Downstream bioinformatics
includes de-multiplexing of reads and trimming, making the reads ready for
downstream analysis such as mapping and *de novo* assembly.
749
750

Figure 2: Mapping of reads and assemblies against the MN target region. Each
751 panel shows the reads and *de novo* assembly mapped against the MN target region in
752 grey and orange, respectively, for species a.) Atlantic cod, b) haddock, c) silvery
753 pout, d) cusk, e) burbot, f) European hake, g) marbled moray cod and h) roughhead
754 grenadier. The positions of genes in the target region are indicated at the top.
755
756

Figure 3: Mapping of reads and assemblies against the LA target region. Each
757 panel shows the reads and *de novo* assembly mapped against the LA target region in
758 grey and orange, respectively, for species a) Atlantic cod, b) haddock, c) silvery pout,
759 d) cusk, e) burbot, f) European hake, g) marbled moray cod and h) roughhead
760 grenadier. The positions of genes in the target region are indicated at the top.
761
762

**Figure 4: The relationship between capture success and genomic divergence to
Atlantic cod**. Linear regression of the relationship between the genomic divergence
to Atlantic cod (SNPs x 10^5) and a) median mapping depth for the MN region (blue),
LA region (red) and the combined target region (black); b) the percentage of reads
mapping to the target region; c) the percentage of the target region covered by reads
to a minimum depth of 10x. For each regression the correlation coefficient, r, is
shown along with a p-value. Each data point is labeled by species according to this
code: Ac=Atlantic cod, H=haddock, Sp=silvery pout, C=cusk, B=burbot, Eh=European
hake, Mm=marbled moray cod and Rg=roughhead grenadier.
771
772

Figure 5: Synteny of the Hb gene clusters. Genomic synteny of the hemoglobin gene
clusters shown at the top for the genomes of Atlantic cod (gadMor2 (Tørresen *et al.*
2017b)) and haddock (MelAeg (Tørresen *et al.* 2017a)). Below, the genomic synteny
inferred from the *de novo* assemblies for all of the species included in the capture
experiment. Stippled lines indicate assembly gaps – here we assume that the
orientation of genes corresponds to the genomes of Atlantic cod and haddock. Gray
boxes indicate genes that have been identified in (Baalsrud *et al.* 2017), but are absent

in the *de novo* assemblies. a) Synteny across the MN region b) Synteny across the LA
region.
782
Figure 6: Polymorphisms in the bi-directional promoter between *Hba1* and *Hbb1*
for five species in the Gadidae family.
A schematic representation of *Hba1* and *Hbb1* with the promoter region between
them. The region contains an indel polymorphism of variable length across the five
species, as indicated by gaps. For each species/variant the alignment is shown along
with amino acid substitutions at positions 55 and 62 in the translated part of the *Hbb1*
gene.

Supporting Information Legends

Table S1: For each species, the average and median depth of reads mapped against the target region (for MN, LA and total), the genomic divergence to Atlantic cod (number of SNPs), percentage of nucleotides mapped to the target and the percentage of the target region with more than 10x coverage.

Table S2: Estimated sequence identity using EMBOSS Needle (Rice *et al.* 2000) with default settings, between paralogous Hbb gene sequences from (Baalsrud *et al.* 2017). Genes highlighted in bold are missing from the assemblies in figure 5.

Table S3: Amino acids at positions 55 and 62 in Hbb1 in various codfishes taken from (Baalsrud *et al.* 2017).

Table S4: Amount of repeated sequences in the target region of the Atlantic cod (gadMor2 (Tørresen *et al.* 2017b)) and haddock (melAeg (Tørresen *et al.* 2017a)) given in percentage.

Table S5: Genes provided Nimblegen for the probe design, and used to identify genes in *de novo* assemblies. For each gene, the gene name is given with its ENSEMBL name and ENSEMBL identifier.

Figure S1: Ancestral reconstruction of amino acids at position 55 in the *Hbb-1* gene in Gadiformes. Phylogenetic trees and ancestral reconstruction was carried out in MEGA 7.0.

Figure S2: Ancestral reconstruction of amino acids at position 62 in the *Hbb-1* gene in Gadiformes. Phylogenetic trees and ancestral reconstruction was carried out in MEGA 7.0.

818   **Figures**

819
820   Figure 1
821

822
823   Figure 2
824

825
826   Figure 3
827

Figure 4

830
831 Figure 5
832

833
834   Figure 6
835

Tables

Table 1. Number of reads and bases captured and sequenced for each species, and number of utgs, largest utg and N50 in the assemblies.

Species	Latin name	Number of reads	Number of bases	Number of utgs	Largest utg (bp)	N50 (bp)
Atlantic cod	*Gadus morhua*	73005	217252583	278	79 020	7 728
Haddock	*Melanogrammus aeglefinus*	35573	107839552	227	52 433	7 227
Silvery pout	*Gadiculus argenteus*	69775	212519845	410	35 801	7 098
Cusk	*Brosme brosme*	55348	175883008	394	64 145	7 322
Burbot	*Lota lota*	56155	165360828	205	70 602	8 055
European hake	*Merluccius merluccius*	65661	180558336	311	31 558	6 523
Marbled moray cod	*Muraenolepus marmoratus*	52076	148100933	455	30 019	6 632
Roughhead grenadier	*Macrourus berglax*	46195	129085001	325	35 216	7122