Title

A critical assessment of single-cell transcriptomes sampled following patch-clamp electrophysiology

Authors

Shreejoy J. Tripathy[1,2,*], Lilah Toker[1,2], Claire Bomkamp[2], B. Ogan Mancarci[1,2], Manuel Belmadani[1,2], Paul Pavlidis[1,2*]

[1] Michael Smith Laboratories, University of British Columbia, Vancouver BC, Canada

[2] Department of Psychiatry, University of British Columbia, Vancouver BC, Canada

[*] Corresponding Author
Email: stripathy@msl.ubc.ca
Email: paul@msl.ubc.ca

Abstract

Patch-seq, combining patch-clamp electrophysiology with single-cell RNA-sequencing (scRNAseq), enables unprecedented single-cell access to a neuron's transcriptomic, electrophysiological, and morphological features. Here, we present a systematic review and re-analysis of scRNAseq profiles from 4 recent patch-seq datasets, benchmarking these against analogous profiles from cellular-dissociation based scRNAseq. We found an increased likelihood for off-target cell-type mRNA contamination in patch-seq, likely due to the passage of the patch-pipette through the processes of adjacent cells. We also observed that patch-seq samples varied considerably in the amount of mRNA that could be extracted from each cell, strongly biasing the numbers of detectable genes. We present a straightforward marker gene-based approach for controlling for these artifacts and show that our method improves the correspondence between gene expression and electrophysiological features. Our analysis suggests that these technical confounds likely limit the interpretability of patch-seq based single-cell transcriptomes. However, we provide concrete recommendations for quality control steps that can be performed prior to costly RNA-sequencing to optimize the yield of high quality samples.

Introduction

Linking gene expression to a neuron's electrical and morphological features has long been a goal of cellular neuroscience. To this end, one strategy is to use the same patch-clamp electrode for electrophysiological characterization for mRNA sampling, for example, by aspirating the cell's cytosol into the patch-pipette (Eberwine et al., 1992; Sucher and Deitcher, 1995; Toledo-Rodriguez et al., 2004; Toledo-Rodriguez and Markram, 2014; Kodama et al., 2012; Rossier et al., 2014). The aspirated mRNA transcripts can then be detected and quantified using RT-PCR (Eberwine et al., 1992; Sucher and Deitcher, 1995; Cauli et al., 1997; Toledo-Rodriguez et al., 2004; Kodama et al., 2012; Rossier et al., 2014) or other methods (Subkhankulova et al., 2010), allowing the quantification of multiple genes or transcripts.

Recently, a number of groups have published protocols for patch-seq that extend previous RT-
PCR-based methods by quantifying patch-pipette sampled cellular mRNA transcripts using
next-generation RNA-sequencing (Cadwell et al., 2015; Fuzik et al., 2016; Földy et al., 2016;
Bardy et al., 2016; Cadwell et al., 2017b, 2017a). These protocols make use of recent technical
improvements in single-cell RNA-sequencing (scRNAseq) that enable gene expression
quantification from very low starting volumes of mRNA (Poulin et al., 2016; Tasic et al., 2017),
such as those present in a single-cell or single-nucleus.
56
Patch-seq mRNA sample collection differs from standard single-cell or single-nucleus RNAseq,
in two major ways (Cadwell et al., 2017b, 2017a). First, as opposed to relying on dissociating
cells into suspension, the micropipette used for electrical recording is used for mRNA extraction
via aspiration. While guiding the patch pipette to (or from) the soma of a cell of interest, the
pipette often must travel through the processes of other cells, presenting an opportunity for
contamination. Second, the effectiveness of cell content aspiration is difficult to control, so the
amount of mRNA extracted may tend to vary from cell to cell.
64
Here, our goal was to investigate the quality of scRNAseq data profiled using patch-seq. Our
strategy was to compare patch-seq derived scRNAseq data with analogous data sampled using
cellular-dissociation based methods, from which multiple large and high-quality single-cell
transcriptomic datasets are available (Tasic et al., 2016; Zeisel et al., 2015). Our findings
suggest that sampling cellular mRNA using a patch-pipette induces technical artifacts that tend
not to be present to the same degree in cellular-dissociation based scRNAseq data. Based on
our findings, we provide approaches for detecting these technical issues and discuss strategies
for generating high-quality patch-seq datasets in the future.
73
74 **Methods**

75 *Dataset overview*
76
We made use of 4 previously published patch-seq datasets (Cadwell, Földy, Fuzik, Bardy)
(Bardy et al., 2016; Cadwell et al., 2015; Földy et al., 2016; Fuzik et al., 2016) , reflecting, to our
knowledge, all of the published patch-seq datasets as of January 2018. We compared these to
2 cellular dissociation-based single-cell RNAseq datasets (Tasic, Zeisel) (Tasic et al., 2016;
Zeisel et al., 2015). We downloaded single-cell transcriptomic data from each study from
accessions provided in Table 1 and Supplementary Table 1 or by contacting the authors directly.
We obtained patch-seq-based electrophysiological data for the Cadwell and Fuzik datasets from
the authors. For all patch-seq datasets, electrophysiological data were provided as a
spreadsheet containing a set of summarized electrophysiological features per cell (e.g., input
resistance, resting membrane potential, etc.). Electrophysiological data from the Allen Institute
Cell Types database (celltypes.brain-map.org) were obtained and processed as described
previously (Tripathy et al., 2017).
89
90 *Transcriptome data pre-processing*
91
We reprocessed transcriptomic data for the Cadwell, Földy, and Tasic datasets directly from
Gene Expression Omnibus (GEO) or Array Express. Data from GEO was downloaded using
fastq-dump version 2.8.2 from the Sequence Read Archive Toolkit. Technical reads such as
barcodes and primers were filtered out during extraction. Adapter sequences were clipped from
the raw reads. The list of option used is as follows: '--gzip --skip-technical --readids --dumpbase
--split-files --clip'. Data from ArrayExpress was downloaded and used directly as prepared by
the European Bioinformatics Institute.

99
The reference mouse transcriptome was produced using the 'rsem-prepare-reference' script
provided by the RSEM RNA-Seq transcript quantifier (Li and Dewey, 2011). The assembly
version used was Ensembl GRCm38, packaged by Illumina for the iGenomes collection.
Alignment was performed using STAR (Dobin et al., 2013) version 2.4.0h, provided as the
aligner to RSEM v1.2.31. Default parameters were used (with the exception of parallel
processing and logging related options). Transcript definitions used to detect ERCC spike-ins
were obtained from the ERCC92 version fasta and GTF files. Spike-ins were concatenated to
the GRCm38 assembly before applying rsem-prepare-reference, and independently to create a
standalone ERCC assembly. Both the concatenated and standalone spike-ins assemblies
showed highly comparable proportions of spike-in expression. For the Fuzik and Zeisel
datasets, we made use of the quantified summarized unique molecule counts (UMIs) made
available at GEO. For the Bardy dataset, we used the summarized count matrices directly
provided by the authors.
113
114 *Mapping of mouse patch-seq cell types onto taxonomies derived from dissociated cells*
115
Using descriptions for cellular identities provided in the original patch-seq publications, we
manually mapped each of the cell types represented across the three mouse patch-seq
datasets onto transcriptomically-defined cellular clusters reported in the two dissociated cell
datasets (shown in Supplementary Table 2). For example, given that the elongated
neurogliaform cells and single bouquet cells characterized in Cadwell are both cortical layer 1
cells, we manually mapped these to the layer 1 cells defined in Tasic as Ndnf cells. Similarly, we
mapped the hippocampal regular-spiking interneurons characterized in Foldy to the Sncg cluster
from Tasic (personal communication with Csaba Földy). To align cell subtype clusters between
Tasic and Zeisel, we used mappings provided by MetaNeighbor (Crow et al., 2018) (shown in
Supplementary Table 2). The mappings between broad cell types in Tasic with Zeisel are
provided in Supplementary Table 3. As with our previous work mapping cells and cell types
across datasets (Mancarci et al., 2017; Tripathy et al., 2017), we note that these cross-dataset
mappings are approximate and ideally would be guided by the use methods for unambiguously
aligning cell types across experiments (e.g., transgenic mouse lines with specific cell types
labeled by fluorescent proteins).
131
132 *Identification of cell type-specific marker genes*
133
For this study, we defined two classes of marker genes, termed "on" and "off" markers. The first
class, "on" markers, are genes that are highly and ubiquitously expressed in the cell type of
interest with enriched expression relative to other cell types. The second class, "off" markers,
are expected to be expressed at low levels in a given patch-seq cell type. These are genes that
are specifically expressed in a single cell type (e.g., microglia) and, if expressed, are an
indicator of possible cellular contamination. To identify marker genes, we employed two recent
surveys of mouse cortical diversity from Tasic et al. and Zeisel et al. (Tasic et al., 2016; Zeisel et
al., 2015).
142
To identify "on" marker genes, we initially used the Tasic dataset, and selected genes whose
average expression in the chosen cell type was >10 times relative all other cell types in the
dataset, with an average expression in the cell type of >100 TPM. From this initial gene list, we
next filtered these genes to only include those that were expressed >10 TPM/cell in >75% of all
cells of that type in Tasic, and >1 UMI/cell in >50% of all cells of that type in Zeisel. Using the
Tasic nomenclature, we defined "on" markers for Ndnf, Sncg, Pvalb, and Pyramidal cell types.

149
To identify "off" marker genes for broad cell types (shown in Supplementary Table 3), as an
initial listing we used the set of cell type-specific marker genes for broad cell classes in the
mouse cortex, defined in our previous work using the NeuroExpresso database (Tasic et al.,
2016). Specifically, we used the set of cortical markers derived from single-cell RNA-seq for
astrocytes, endothelial cells, microglia, oligodendrocytes, oligodendrocyte precursor cells, and
pyramidal cells. From this list, we first filtered out lowly expressed genes that were expressed
<10 TPM/cell in >50% of all cells of that type in Tasic, and <1 UMI/cell in >50% of all cells of
that type in Zeisel. Next, we filtered genes too broadly expressed in our patch-seq cell types of
interest by assessing the expression of these genes in the Ndnf, Sncg, Pvalb, and Pyramidal
cell types, removing genes that were expressed at a level greater than >10 TPM/cell in >33% of
all cells of that type in Tasic, and >2 UMI/cell in >33% of all cells of that type in Zeisel.
161
When defining on and off marker genes for inhibitory cell subtypes (e.g., the Ndnf cell type), we
did not compare these cells to other GABAergic cells. For example, when defining "on" markers
for Ndnf cells, we did not compare these cells' expression to Pvalb or Sst cells. We note that
this choice limits our ability to identify inhibitory-to-inhibitory cell contamination, for example, an
Ndnf cell contaminated by Sst-cell specific markers. To define an initial set of "off" markers for
GABAergic inhibitory cells, we first obtained a list of genes based on Tasic where in GABAergic
cells had average expression >10 times all other non-GABAergic cells in the dataset and with
an average expression of at least 100 TPM.
170
171   The final list of filtered mouse cell type specific marker genes used in this study are provided in
172   Supplementary Table 4.
173
174   To obtain a list of human cell type specific marker genes for use for the Bardy dataset, we made
175   use of classic cell-type specific markers for astrocytes and microglia, based on human purified
176   cell types shown in Figure 4A of reference (Zhang et al., 2016).
177
178   *Summarizing cell type-specific marker expression*
179
When directly comparing expression values from patch-seq data to dissociated cell data, we
compared the Cadwell and Földy datasets to Tasic, as these all were quantified using TPM and
employed Smart-seq-based methods. Similarly, we compared Fuzik dataset to Zeisel, as these
both used C1-STRT and were quantified using unique molecule identifiers (UMIs), normalized
as UMI counts per million. We summarized a single-cell sample's expression of multiple cell
type-specific markers using the sum of the $\log_2$ normalized expression values. Given a patch-
seq sample of cell type identity A (e.g., a pyramidal cell) and wanting to quantify its normalized
expression of "off" markers for cell type B (e.g., microglial markers), we used the dissociated cell
data to estimate the median expression of cell type B's "markers in cells of type A (e.g., median
expression level of microglial markers in pyramidal cells) and the median expression of cell type
B's markers in cells of type B (e.g., median expression level of microglial markers in microglia
cells). Specifically, we normalized expression to a value of approximately 0 to 1, as follows:
192
193   *(PatchSeqCellTypeA*_markersB – median(*DissocCellTypeA*_markersB))/
194   (median(*DissocCellTypeB*_markersB) - median(*DissocCellTypeA*_markersB))
195
196   where we set all negative values to 0. Next, to obtain a single contamination index per single
197   cell, we summed all contamination scores for all broad cell types, excluding the patch-seq cell's
198   assigned broad cell type.

199
Lastly, to obtain a scalar quality score for transcriptomic data from patch-seq samples (e.g., for
analysis of electrophysiological data), we used the Spearman correlation of each patch-seq
sample's expression of "on" and "off" marker genes to the average expression profile of
dissociated cells of the same cell type (shown in Supplement Figure 3). For example, for an
Ndnf patch-seq sample from Cadwell, we first calculated the average expression profile of Ndnf
cells from Tasic across the set of all "on" and "off" marker genes (i.e., Ndnf markers, pyramidal
cell markers, astrocyte markers, etc.), and then calculated the correlation between the patch-
seq cell's marker expression to the mean dissociated cell expression profile. Since these
correlations could potentially be negative, we set quality scores to a minimum of 0.1. A
convenient feature of this quality score is that it yields low correlations for samples with
relatively high contamination as well as those where contamination is largely undetected but
expression of endogenous "on" markers is also low (Supplement Figure 3).
212
213 *Analysis of factors influencing the numbers of genes detected per cell*
214
We analyzed how the following factors influenced the numbers of genes detected per cell:
library size, defined as the total numbers of reads sequenced per cell; spike-in ratio, defined as
the number of reads mapping to ERCC spike-ins divided by total sequenced reads; the
unmapped ratio, defined as the ratio of reads not mapping to the exonic reference divided by all
non-ERCC sequenced reads; and cellular contamination indices, as defined in the previous
section. For the Cadwell, Tasic, and ERCC-containing subsets of the Földy, and Bardy
datasets, we fit a linear model (implemented using the 'lm' function in R) for numbers of
detected genes per each cell as follows:
223
224 num_genes ~ log10(library_size) + spike-in_ratio + unmapped_ratio + contam_index
225
226 where each term above was first scaled to z-scores, yielding standardized beta coefficients.
227
228 *Combined analysis of transcriptomic and electrophysiological features*
229
We analyzed correlations between transcriptomic and electrophysiological features using an
approach similar to our previous work (Tripathy et al., 2017). For each patch-seq dataset, we
first filtered for genes whose average expression was > 30^{th} percentile relative to all genes in
the dataset. We analyzed electrophysiological features overlapping with our previous analysis,
specifically, input resistance (Rin), resting membrane potential (Vrest , action potential threshold
(APthr) , action potential amplitude (APamp) , action potential half-width (APhw), membrane
time constant (Tau), after-hyperpolarization amplitude (AHPamp), rheobase (Rheo), maximum
firing rate (FRmax), and capacitance (Cm). We calculated Pearson correlations between the set
of electrophysiology features and gene expression values, both without weighting cells by their
overall quality scores (based on correlation of markers to dissociated cell samples), and after
weighting cells using their quality scores.
241
We performed an analogous analysis for comparison of pooled-cell correlations based on the
AIBS/Tasic dataset, where we computationally merged different groups of cells characterized
using dissociated cell scRNAseq (based on Tasic et al, (Tasic et al., 2016)) with cells
characterized using patch-clamp electrophysiology (Teeter et al., 2018) based on the overlap of
same mouse transgenic lines and coarse cortical layers (i.e., upper vs lower mouse visual
cortex). For example, we merged 14 QC-passing scRNAseq samples from the Sst-IRES-cre
mouse line from visual cortex dissections specific to lower layers with 89 patch-clamp samples

from the same mouse line from cortical layers 4 through 6b. After merging single-cells into cell
types, we averaged expression and electrophysiological values; since cell types tended to be
represented by differing numbers of cells, in our gene-electrophysiology correlation analyses we
weighted cell types based on the numbers of cells available using the square root of the
harmonic mean of the number of cells characterized by electrophysiology and
electrophysiology.
255
256 *Statistical information*
257 We used the R weights toolbox (v0.85) to calculate weighted Pearson correlations and raw p-
258 values. We used the Benjamini-Hochberg False Discovery Rate (FDR) to account for analysis of
259 multiple correlations.
260
261 *Computer code and data availability*
262
263 All computational code and associated data has been made accessible at
264 https://github.com/PavlidisLab/patchSeqQC and code for the RNAseq pipeline is accessible at
265 https://github.com/PavlidisLab/rnaseq-pipeline.
266
267
268 **Results**
269
270 To quantitatively assess the influence of patch-seq specific technical confounds, we performed
271 a re-analysis of four recently published patch-seq datasets. We focused our analyses on three
272 datasets obtained from mouse acute brain slices (Cadwell et al., 2015; Földy et al., 2016; Fuzik
273 et al., 2016) and contrast these against one dataset obtained from human stem-cell derived
274 neurons and astrocytes in culture (Bardy et al., 2016) (Table 1).
275

Dataset	Description	Preparation	RNA amplification	Number of cells	Accession
Cadwell (Cadwell et al., 2015)	Cortical layer 1 interneurons	Acute mouse slices	Smart-seq2	58	E-MTAB-4092
Fuzik (Fuzik et al., 2016)	Cortical layer 1/2 interneurons and pyramidal cells	Acute mouse slices	STRT-C1 (with unique molecule identifiers)	80	GSE70844
Földy (Földy et al., 2016)	Hippocampal CA1 and Subiculum pyramidal cells and regular- and fast-spiking interneurons	Acute mouse slices	SMARTer	93	GSE75386
Bardy (Bardy et al., 2016)	Stem-cell derived neurons and astrocytes	Differentiated human cells in culture	SMARTer	56	NA*

276 *Table 1: Description of patch-seq datasets re-analyzed in this study. *Expression data obtained by contacting the authors*
277 *directly.*

278 *Expression of off-target cell type marker genes in patch-seq samples*
279

We first assessed if patch-seq based single-cell transcriptomes might have been contaminated
by mRNA from other cells adjacent to the patched cell (Figure 1A, B), termed off-target cell-type
contamination (Okaty et al., 2011). For example, is there paradoxical expression of genes
specific to microglia in the scRNAseq profile of a recorded pyramidal cell? To address this
question, we made use of the fact that the broad identities of the recorded cells can be
ascertained from morphological and electrophysiological features without relying on the
transcriptomic data (see Methods). Furthermore, we used multiple mouse forebrain scRNAseq
datasets collected from dissociated cells to define lists of marker genes specific to various
cortical and hippocampal cell types (Supplementary Table 4) (Mancarci et al., 2017; Tasic et al.,
2016; Zeisel et al., 2015).
290
We detected that some of the single cell samples from the three mouse datasets collected from
acute brain slices expressed markers for multiple distinct cell types (Figure 1, Supplement
Figure 1). For example, some of the cortical layer 1 elongated neurogliaform cells (eNGCs)
characterized in the Cadwell dataset appeared to also express multiple marker genes specific to
pyramidal cells (Figure 1C), such as *Slc17a7*, the vesicular glutamatergic transporter VGLUT1.
Similarly, many of the cells identified as hippocampal regular spiking GABAergic interneurons in
the Földy dataset also expressed microglial and pyramidal cell markers (Figure 1H).
298
We sought to quantify the extent of off-target cell type contamination in the mouse patch-seq
samples. We directly compared the patch-seq-based expression profiles to cellular dissociation-
based transcriptomes from two recent surveys of mouse cortical diversity from Tasic et al. and
Zeisel et al. (Tasic et al., 2016; Zeisel et al., 2015). After matching cell type identities across
studies (shown in Supplementary Table 2), we found that compared to dissociated cells, patch-
seq-based samples expressed markers for multiple cell types at considerably higher levels
(Figure 1C, H, J; Supplement Figure 2A, B). We defined a simple contamination index,
providing a scalar value for greater than expected off-target cell type marker expression across
multiple classes of broad cell types, by comparison to analogous cells from the dissociated-cell
reference (see Methods). Importantly, patch-seq-based samples with larger contamination
indices also expressed markers of their own cell type at lower levels (Supplement Figure 3).
We note that we saw less off-target cell type marker expression in the Fuzik dataset relative to
the Cadwell and Földy datasets (Supplement Figure 2), suggesting either less contamination
in these cells or that the lower gene detection rate in this dataset (Figure 3B) obscures our
ability to use expression profiles to identify cellular contamination.
314

We next assessed the degree of off-target cell type contamination in the Bardy patch-seq
dataset of human stem-cell derived neurons and astrocytes obtained from cultured cells (Bardy
et al., 2016). Since the cells in this dataset were cultured relatively sparsely, allowing the
processes of each cultured cell to be easily visualized (personal communication with Cedric
Bardy), we wondered if this dataset would show less off-target cell type marker expression

compared to the three mouse acute brain slice datasets. Indeed, when assessing astrocyte
marker expression in the population of electrophysiologically-mature neurons (with markers
based on purified human cells (Zhang et al., 2016)), we found these neurons showed some, but
overall very little, expression of astrocyte markers relative to the mature astrocytes also profiled
in this dataset (Figure 2A, B). In addition, both neurons and astrocytes showed almost no
expression of microglia markers (Figure 2A), perhaps unsurprisingly, since microglial cells are
not present in these cultures (Bardy et al., 2016). This example provides suggestive evidence
that the density of processes of adjacent cells might contribute to off-target mRNA
contamination.
341

342
Figure 2. Expression of cell type-specific marker genes in patch-seq samples obtained from human astrocytes and neurons
differentiated in culture from the Bardy dataset. A) Gene expression profiles for differentiated astrocytes (green) and
electrophysiologically-mature neurons (red) for astrocyte and microglial-specific (grey) marker genes. Each column
reflects a single-cell sample. Two astrocyte cells were removed because they expressed fewer than 3 astrocyte
markers. B) Summed astrocyte marker expression for astrocyte and mature neuron single-cells, for the same cells
shown in part A.

Technical factors strongly influence the numbers of genes detected per cell
350
Next, we wondered if there are identifiable technical factors that can help explain the large
ranges in the numbers of genes detected per cell in each dataset, from 6000-13000 genes/cell
in Cadwell to 800-7000 genes/cell in Fuzik (Figure 3B). Because patch-seq mRNA collection
requires the experimenter to manually aspirate cellular mRNA into the patch-pipette, we
reasoned that mRNA harvesting would be difficult to consistently control from cell to cell, leading
there to be different amounts of extracted mRNA per cell. To estimate how much cellular mRNA
was extracted per cell, we made use of ERCC spike-ins (Tasic et al., 2017), which are synthetic
control mRNAs that are added to single-cell samples prior to library preparation and sequencing
(Figure 3A). Specifically, since the same amount of ERCC spike-in mRNAs are added to each
sample, we can use the ratio of spike-in reads to the total count of sequenced reads to estimate
the relative amount of extracted mRNA per cell (Lun et al., 2017; Vallejos et al., 2017). Here,
every cell in the Cadwell and Tasic datasets and a subset of cells in the Földy and Bardy
datasets contained ERCC spike-ins.
364
We used a multivariate regression approach to ask how various technical factors contribute to
the numbers of genes detected per cell in the Cadwell, Földy, and Bardy patch-seq datasets
and the Ndnf cell subset of the Tasic dissociated-cell dataset (Figure 3C; the Fuzik dataset did

not include spike-ins). Library size (the number of sequenced reads per cell) was positively
correlated with detected gene counts in the Tasic and Cadwell datasets (Figure 3C, D).
Similarly, cells with a larger ratio of spike-in reads to total sequenced reads (i.e., with lower
initial amounts of cellular mRNA; Figure 3A), had lower numbers of detected genes across all of
the datasets (Figure 3D), pointing to the importance of mRNA extraction efficiency. In addition,
we saw considerably greater ranges in the spike-in ratio in the patch-seq datasets relative to the
Tasic dataset (Cadwell: 3-17%, Bardy: 3-37%, Tasic: .4-4%).
375
Next, we reasoned that though many mRNA transcripts might be extracted from a cell, not all of
these would be sufficiently high quality to map to the reference (e.g., they might reflect
degraded mRNAs (Cadwell et al., 2017b, 2017a), other contaminants, etc.). To account for this
possibility, we calculated the ratio of unmapped to mapped reads, after excluding reads
mapping to spike-ins. Cells with very large ratios of unmapped to mapped reads had fewer
genes detected (Figure 3C). This technical factor was especially important in the Földy and
Bardy datasets, with some cells in the Földy dataset having fewer than 10% of reads mapped to
the transcriptome (Figure 3G). Lastly, we further wondered if cells showing greater amounts of
off-target cell type contamination would also have a greater number of detected genes. We
found that cells with greater contamination indices from the Cadwell and Földy datasets (i.e., the
acute slice-based patch-seq datasets) had more genes detected, consistent with previous
reports (Ilicic et al., 2016; Vallejos et al., 2017). In total, these simple technical factors explain
between 50-85% of the cell-to-cell variance in the detected gene counts per patch-seq datasets
(Figure 3I).
390

391
Figure 3. Patch-seq experimental confounds affect the numbers of genes detected per cell. A) Schematic illustrating
how spike-in mRNAs can be used to estimate how much mRNA was extracted per cell. B) Violin plots showing
numbers of protein-coding genes detected per cell across patch-seq datasets or the Ndnf subset of the Tasic
dissociated-cell dataset. C) Technical factors associated with numbers of genes detected per cell across datasets
(dataset color shown in B). Bars show standardized beta model coefficients with y-axis in units of standard
deviations, allowing comparison of effects across factors and across datasets. Error bars indicate coefficient standard
deviations. Positive (negative) coefficients indicate factor is correlated with increased (decreased) gene counts.
Regression models calculated using only cells containing mRNA spike-ins. D-H) Examples of univariate relationships
between technical factors and detected gene count per cell (dots) across patch-seq datasets. Grey line shows best fit
line. D) Library size (count of sequenced reads per cell). F) Spike-ins as a fraction of all sequenced reads per cell.
Samples with lower cellular mRNA content (indicated by higher spike-in ratios) have lower gene counts. G)
Unmapped ratio, calculated as the ratio of exonic reads to all other reads (excluding spike-ins). H) Cellular
contamination index, quantified by summing normalized contamination values across tested cell types (arbitrary
units). F) Overall percent variance explained by each dataset-specific statistical model shown in E.

Accounting for technical factors improves the correspondence with electrophysiological features

Lastly, we performed an integrated analysis of gene expression and electrophysiological features for the 3 mouse-based patch-seq datasets, reasoning that more lower quality patch-seq samples would be less informative of relationships between cellular electrophysiology and gene expression (Tripathy et al., 2017). We first calculated a quality score for each patch-seq sample, based on the similarity of its marker expression to dissociated cells of its same type (see Methods; Supplement Figure 3). After statistically down-weighting lower quality cells, we observed a modest improvement in the correspondence between gene expression and electrophysiology, as evidenced by an increase in the number of genes significantly correlated with electrophysiological features (FDR < 0.1, Figure 4A, B). In addition, after correction, we found more genes overlapping with those identified in our previous gene-electrophysiology correlation analysis based on pooled cell types (Tripathy et al., 2017) (Figure 4C). While the biological implications of these correlations require further investigation, this analysis suggests that controlling for these technical factors can help improve the interpretability of patch-seq data.

Figure 4. Adjusting for patch-seq experimental confounds improves the correspondence with electrophysiological measures. A) Comparison of gene expression (Nek7; x-axis) with electrophysiological features (action potential half-width; AP_{hw}; y-axis). Left panel shows single-cell samples (circles) from the Földy dataset. Middle panel shows same data as left, but size of circles proportional to each sample's quality score, defined as the similarity of marker

Discussion

The patch-seq technique reflects a considerable leap in our ability to interrogate a neuron
across multiple features of its activity. However, across our analyses of multiple patch-seq
datasets, we noticed several technical issues that appeared to be shared across experiments.
First, in the three mouse datasets collected from acute brain slices (Cadwell et al., 2015; Földy
et al., 2016; Fuzik et al., 2016), we observed that many single cell samples appeared to strongly
express marker genes from off-target cell types. We interpret this as mRNA contamination from
cells adjacent to the recorded cell, but note that there are alternative explanations. Second, we
observed that mRNA extraction efficiency differs between sampled cells, leading to varying
numbers of genes detected even among cells of the same broad type. These technical artifacts
can be mitigated in part through post hoc analyses, such as our attempt to weight single-cells by
the similarity of their marker gene expression to analogous dissociated cells of the same broad
cell type.

To detect off-target cell type contamination, our main approach was to compare patch-seq
based single-cell transcriptomes to dissociated-cell based reference scRNAseq data from
similar cell types. We used these reference data to identify cell type-specific marker genes as
well as to determine approximately how much off-target marker expression would be expected
in each cell type. We note that there are obvious methodological differences between
dissociated-cell scRNAseq and patch-seq (Cadwell et al., 2017b, 2017a), such as the strain
induced by dissociating cells (Wu et al., 2017) or that patch-seq might be more likely to sample
transcripts from distal cellular processes. Thus we cannot conclusively rule out that some of the
off-target cell type marker expression might reflect a true biological signal, as opposed to mRNA
contamination from adjacent cells. However, we note that the use of marker genes to identify
suspected off-target contamination is a routine quality control step in cell type-specific gene
expression analyses (Mancarci et al., 2017; Okaty et al., 2011), including recent methods for
identifying suspected "doublets" or multi-cell contamination in droplet-based scRNAseq (Zeisel
et al., 2018).

We speculate that the sources of off-target contamination are the processes of cells adjacent to
the patch-pipette. For example, while there are relatively few cell bodies in layer 1 of the
neocortex, there are processes of other cell types like pyramidal cells, and it is well established
that these processes contain mRNA transcripts (Glock et al., 2017). In addition, we noticed that
we routinely observed expression of microglial markers in the mouse patch-seq samples. This is
interesting because the presence of even 1 mM ATP in the patch-pipette is sufficient to induce
rapid chemotaxis of microglial processes towards the pipette (Madry et al., 2018). Patch-clamp
intracellular solutions usually use 2 or 4 mM ATP (Tebaykin et al., 2017), including those of the
patch-seq datasets here (Bardy et al., 2016; Cadwell et al., 2015; Földy et al., 2016; Fuzik et al.,
2016). At present, it is unclear whether this suspected off-target contamination might occur
while the pipette is actively manipulated under positive pressure towards the recorded cell.

Alternatively, such contamination might take place following mRNA extraction during the
retraction of the pipette from the neuropil and recording chamber. Assuming that neuropil is the
major source of off-target contamination, this suggests that there may be advantages to
performing patch-seq on sparsely cultured or acutely dissociated cells (Bardy et al., 2016;
Kodama et al., 2012; Schulz et al., 2006).

Our analyses identified several technical factors that influence the numbers of genes detected
per cell. First, to obtain a sufficient number of detected genes, it is essential to extract a large
amount of mRNA from the targeted cell. However, this itself is not sufficient, as other factors,
such as mRNA degradation can lead the extracted transcripts being too low quality to map to
the genomic reference (Cadwell et al., 2017b, 2017a). Second, given sufficient extraction of
non-degraded transcripts, because of the extremely high sensitivity of modern ultra-low mRNA
capture kits (Poulin et al., 2016; Tasic et al., 2017), any off-target cell-type contamination will
inflate the numbers of genes detected per cell. This suggests that the detected gene count,
often used as a proxy for the quality of scRNAseq data, should not be the only quality control
metric for single-cell transcriptomes sampled using patch-seq.

The effect of these technical confounds on downstream analyses of patch-seq data is likely
context specific. For example, the presence of a small degree of off-target contamination is
likely to be of little consequence if the patch-seq data is used as a "Rosetta stone", to help
connect cellular classifications based on different methodologies, such as transcriptomically-
defined cell clusters with electrophysiological clusters (Fuzik et al., 2016; Tasic, 2018).
However, accurately quantifying single-cell transcriptomes is likely to be much more important
when using these data to investigate how transcriptomic heterogeneity gives rise to subtle cell
to cell variability in physiological features (Cadwell et al., 2015; Schulz et al., 2006; Tripathy et
al., 2017).

Our analyses point to quality control steps that can improve the yield of high-quality patch-seq
samples. An advantage of patch-seq over traditional dissociated-cell based scRNA-seq is that a
cell's electrophysiological and morphological features are often sufficient to determine its broad
cell type (Cadwell et al., 2015; Földy et al., 2016; Fuzik et al., 2016). We argue that knowing a
cell's broad type can help quality control its sampled transcriptome: the cell should express
marker genes of its own type, including highly expressed markers as well as more lowly
expressed markers, such as some transcription factors and long non-coding RNAs (Mancarci et
al., 2017). In addition, the cell should not express marker genes specific to other cell types. This
quality control step can be performed following RNAseq, as we pursue here. However, this
quality control could also be performed after library preparation and amplification but prior to
costly sequencing, for example, using qPCR to detect the expression of a small number of
expected and unexpected marker genes (Bardy et al., 2016).

To summarize, though patch-seq provides a powerful method for multi-modal neuronal
characterization (Bardy et al., 2016; Cadwell et al., 2017b; Földy et al., 2016; Fuzik et al., 2016),
it is susceptible to a number of methodology-specific technical artifacts, such as an increased
likelihood of mRNA contamination from adjacent cells. These artifacts strongly bias traditional
scRNAseq quality metrics such as the numbers of genes detected per cell. By leveraging high-
quality reference atlases of single-cell transcriptomic diversity (Tasic et al., 2016; Zeisel et al.,
2015), we argue that inspection of cell type-specific marker expression should be an essential
patch-seq quality control step prior to downstream analyses.

Acknowledgements
529

530 We thank Cathryn Cadwell, Janos Fuzik, Csaba Földy, and Cedric Bardy for sharing data and
531 Jim Berg, Dmitry Kobak, and Philipp Berens for helpful discussions. This work was supported by
532 Kids Brain Health Network, a Canadian Institute for Health Research Post-doctoral Fellowship,
533 and NIH grant MH111099.

534
535 **Author contributions**

536
537 SJT and PP conceived the project. SJT implemented the methodology and generated the
538 results with assistance from LT, OBM, CB and MB. All authors contributed to interpreting the
539 results. SJT and PP wrote the paper with assistance from all authors.

540
541 **Competing interests**

542
543 The authors declare no competing financial interests.

544
545
546 **References**

547
Bardy, C., van den Hurk, M., Kakaradov, B., Erwin, J. A., Jaeger, B. N., Hernandez, R. V., et al.
(2016). Predicting the functional states of human iPSC-derived neurons with single-cell
RNA-seq and electrophysiology. *Mol. Psychiatry*. doi:10.1038/mp.2016.158.

Cadwell, C. R., Palasantza, A., Jiang, X., Berens, P., Deng, Q., Yilmaz, M., et al. (2015).
Electrophysiological, transcriptomic and morphologic profiling of single neurons using
Patch-seq. *Nat. Biotechnol.* doi:10.1038/nbt.3445.

Cadwell, C. R., Sandberg, R., Jiang, X., and Tolias, A. S. (2017a). Q&A: using Patch-seq to
profile single cells. *BMC Biol.* 15, 58. doi:10.1186/s12915-017-0396-0.

Cadwell, C. R., Scala, F., Li, S., Livrizzi, G., Shen, S., Sandberg, R., et al. (2017b). Multimodal
profiling of single-cell morphology, electrophysiology, and gene expression using Patch-
seq. *Nat. Protoc.* 12, nprot.2017.120. doi:10.1038/nprot.2017.120.

Cauli, B., Audinat, E., Lambolez, B., Angulo, M. C., Ropert, N., Tsuzuki, K., et al. (1997).
Molecular and Physiological Diversity of Cortical Nonpyramidal Cells. *J. Neurosci.* 17,
3894–3906.

Crow, M., Paul, A., Ballouz, S., Huang, Z. J., and Gillis, J. (2018). Characterizing the
replicability of cell types defined by single cell RNA-sequencing data using
MetaNeighbor. *Nat. Commun.* 9, 884. doi:10.1038/s41467-018-03282-0.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., et al. (2013). STAR:
ultrafast universal RNA-seq aligner. *Bioinforma. Oxf. Engl.* 29, 15–21.
doi:10.1093/bioinformatics/bts635.

Eberwine, J., Yeh, H., Miyashiro, K., Cao, Y., Nair, S., Finnell, R., et al. (1992). Analysis of
gene expression in single live neurons. *Proc. Natl. Acad. Sci.* 89, 3010–3014.
doi:10.1073/pnas.89.7.3010.

Földy, C., Darmanis, S., Aoto, J., Malenka, R. C., Quake, S. R., and Südhof, T. C. (2016).
Single-cell RNAseq reveals cell adhesion molecule profiles in electrophysiologically
defined neurons. *Proc. Natl. Acad. Sci.* 113, E5222–E5231.
doi:10.1073/pnas.1610155113.

Fuzik, J., Zeisel, A., Máté, Z., Calvigioni, D., Yanagawa, Y., Szabó, G., et al. (2016). Integration
of electrophysiological recordings with single-cell RNA-seq data identifies neuronal
subtypes. *Nat. Biotechnol.* 34, 175–183. doi:10.1038/nbt.3443.

Glock, C., Heumüller, M., and Schuman, E. M. (2017). mRNA transport & local translation in
neurons. *Curr. Opin. Neurobiol.* 45, 169–177. doi:10.1016/j.conb.2017.05.005.

Ilicic, T., Kim, J. K., Kolodziejczyk, A. A., Bagger, F. O., McCarthy, D. J., Marioni, J. C., et al.
(2016). Classification of low quality cells from single-cell RNA-seq data. *Genome Biol.*
17, 29. doi:10.1186/s13059-016-0888-1.

Kodama, T., Guerrero, S., Shin, M., Moghadam, S., Faulstich, M., and Lac, S. du (2012).
Neuronal Classification and Marker Gene Identification via Single-Cell Expression
Profiling of Brainstem Vestibular Neurons Subserving Cerebellar Learning. *J. Neurosci.*
32, 7819–7831. doi:10.1523/JNEUROSCI.0543-12.2012.

Li, B., and Dewey, C. N. (2011). RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* 12, 323. doi:10.1186/1471-
2105-12-323.

Lun, A. T. L., Calero-Nieto, F. J., Haim-Vilmovsky, L., Göttgens, B., and Marioni, J. C. (2017).
Assessing the reliability of spike-in normalization for analyses of single-cell RNA
sequencing data. *Genome Res.* doi:10.1101/gr.222877.117.

Madry, C., Kyrargyri, V., Arancibia-Cárcamo, I. L., Jolivet, R., Kohsaka, S., Bryan, R. M., et al.
(2018). Microglial Ramification, Surveillance, and Interleukin-1β Release Are Regulated
by the Two-Pore Domain K+ Channel THIK-1. *Neuron* 97, 299-312.e6.
doi:10.1016/j.neuron.2017.12.002.

Mancarci, B. O., Toker, L., Tripathy, S. J., Li, B., Rocco, B., Sibille, E., et al. (2017). Cross-
Laboratory Analysis of Brain Cell Type Transcriptomes with Applications to
Interpretation of Bulk Tissue Data. *eNeuro*, ENEURO.0212-17.2017.
doi:10.1523/ENEURO.0212-17.2017.

Okaty, B. W., Sugino, K., and Nelson, S. B. (2011). A Quantitative Comparison of Cell-Type-
Specific Microarray Gene Expression Profiling Methods in the Mouse Brain. *PLoS ONE*
6, e16493. doi:10.1371/journal.pone.0016493.

Poulin, J.-F., Tasic, B., Hjerling-Leffler, J., Trimarchi, J. M., and Awatramani, R. (2016).
Disentangling neural cell diversity using single-cell transcriptomics. *Nat. Neurosci.* 19,
1131–1141. doi:10.1038/nn.4366.

Rossier, J., Bernard, A., Cabungcal, J.-H., Perrenoud, Q., Savoye, A., Gallopin, T., et al. (2014). Cortical fast-spiking parvalbumin interneurons enwrapped in the perineuronal net express the metallopeptidases Adamts8, Adamts15 and Neprilysin. *Mol. Psychiatry*. doi:10.1038/mp.2014.162.

Schulz, D. J., Goaillard, J.-M., and Marder, E. (2006). Variable channel expression in identified single and electrically coupled neurons in different animals. *Nat. Neurosci.* 9, 356–362. doi:10.1038/nn1639.

Subkhankulova, T., Yano, K., Robinson, H. P. C., and Livesey, F. J. (2010). Grouping and classifying electrophysiologically-defined classes of neocortical neurons by single cell, whole-genome expression profiling. *Front. Mol. Neurosci.* 3, 10. doi:10.3389/fnmol.2010.00010.

Sucher, N. J., and Deitcher, D. L. (1995). PCR and patch-clamp analysis of single neurons. *Neuron* 14, 1095–1100. doi:10.1016/0896-6273(95)90257-0.

Tasic, B. (2018). Single cell transcriptomics in neuroscience: cell classification and beyond. *Curr. Opin. Neurobiol.* 50, 242–249. doi:10.1016/j.conb.2018.04.021.

Tasic, B., Levi, B. P., and Menon, V. (2017). "Single-Cell Transcriptomic Characterization of Vertebrate Brain Composition, Development, and Function," in *Decoding Neural Circuit Structure and Function* (Springer, Cham), 437–468. doi:10.1007/978-3-319-57363-2_18.

Tasic, B., Menon, V., Nguyen, T. N., Kim, T. K., Jarsky, T., Yao, Z., et al. (2016). Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat. Neurosci.* 19, 335–346. doi:10.1038/nn.4216.

Tebaykin, D., Tripathy, S. J., Binnion, N., Li, B., Gerkin, R. C., and Pavlidis, P. (2017). Modeling sources of interlaboratory variability in electrophysiological properties of mammalian neurons. *J. Neurophysiol.* 119, 1329–1339. doi:10.1152/jn.00604.2017.

Teeter, C., Iyer, R., Menon, V., Gouwens, N., Feng, D., Berg, J., et al. (2018). Generalized leaky integrate-and-fire models classify multiple neuron types. *Nat. Commun.* 9, 709. doi:10.1038/s41467-017-02717-4.

Toledo-Rodriguez, M., Blumenfeld, B., Wu, C., Luo, J., Attali, B., Goodman, P., et al. (2004). Correlation maps allow neuronal electrical properties to be predicted from single-cell gene expression profiles in rat neocortex. *Cereb. Cortex N. Y. N 1991* 14, 1310–1327. doi:10.1093/cercor/bhh092.

Toledo-Rodriguez, M., and Markram, H. (2014). Single-cell RT-PCR, a technique to decipher the electrical, anatomical, and genetic determinants of neuronal diversity. *Methods Mol. Biol. Clifton NJ* 1183, 143–158. doi:10.1007/978-1-4939-1096-0_8.

Tripathy, S. J., Toker, L., Li, B., Crichlow, C.-L., Tebaykin, D., Mancarci, B. O., et al. (2017). Transcriptomic correlates of neuron electrophysiological diversity. *PLOS Comput. Biol.* 13, e1005814. doi:10.1371/journal.pcbi.1005814.

Vallejos, C. A., Risso, D., Scialdone, A., Dudoit, S., and Marioni, J. C. (2017). Normalizing
single-cell RNA sequencing data: challenges and opportunities. *Nat. Methods*.
doi:10.1038/nmeth.4292.

Wu, Y. E., Pan, L., Zuo, Y., Li, X., and Hong, W. (2017). Detecting Activated Cell Populations
Using Single-Cell RNA-Seq. *Neuron* 96, 313-329.e6. doi:10.1016/j.neuron.2017.09.026.

Zeisel, A., Hochgerner, H., Lonnerberg, P., Johnsson, A., Memic, F., Zwan, J. van der, et al.
(2018). Molecular architecture of the mouse nervous system. *bioRxiv*, 294918.
doi:10.1101/294918.

Zeisel, A., Muñoz-Manchado, A. B., Codeluppi, S., Lönnerberg, P., Manno, G. L., Juréus, A., et
al. (2015). Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-
seq. *Science* 347, 1138–1142. doi:10.1126/science.aaa1934.

Zhang, Y., Sloan, S. A., Clarke, L. E., Caneda, C., Plaza, C. A., Blumenthal, P. D., et al. (2016).
Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals
Transcriptional and Functional Differences with Mouse. *Neuron* 89, 37–53.
doi:10.1016/j.neuron.2015.11.013.

660
661

Supplemental Figures

Supplement Figure 1. Expression of cell type-specific marker genes in patch-seq samples from Fuzik. A) Gene expression profiles for sampled pyramidal cells for various cell type-specific markers. B) Summed expression of cell type-specific marker genes for Pyramidal cell (x-axis) and Astrocyte (y-axis) markers. Dots reflect cortical Pyramidal cell (turquoise) and Astrocyte (green) single cells collected in the Zeisel dataset, based on dissociated scRNAseq. Dashed lines reflect 95% intervals of marker expression for each cell type. C) Same as B, but showing summed marker expression for Pyramidal cells shown in A based on patch-seq data. Arrow denotes the same single-cell highlighted in A. D,E) Same as B and C, but showing comparison of microglial marker expression.

 Supplementary Tables

Dataset	Experiment type	Preparation	Description	Accession	Number of cells
Tasic (Tasic et al., 2016)	dissociated cell scRNAseq	Dissociated cells	Visual cortex neurons and glia	GSE71585	1366
Zeisel (Zeisel et al., 2015)	dissociated cell scRNAseq	Dissociated cells	Somatosensory cortex and hippocampus neurons and glia	GSE60361	3005
Allen Institute Cell Types (Teeter et al., 2018)	patch-clamp electrophysiology	Acute mouse slices	Visual cortex neurons	celltypes.brain-map.org	952

700 *Supplementary Table 1: Description of dissociated-cell scRNAseq datasets and patch-clamp*
701 *electrophysiological datasets used. For RNA amplification, the Tasic scRNAseq dataset*
702 *employed SMARTer (i.e., Smart-seq based, consistent with the Cadwell, Foldy, and Bardy*
703 *datasets) whereas the Zeisel dataset employed C1-STRT (consistent with the Fuzik dataset).*

Patch-seq dataset	Cell type (patch-seq)	Matched cell type (dissociated cell; Tasic)	Matched cell type (dissociated cell; Zeisel)
Cadwell	Cortex Layer 1 elongated neuragliaform cell (eNGC)	Ndnf cluster (Ndnf Car4, Ndnf Cxcl14)	Int12, Int15
Cadwell	Cortex Layer 1 single bouquet cells (SBC)	Ndnf cluster (Ndnf Car4, Ndnf Cxcl14)	Int12, Int15
Földy	Hippocampus regular-spiking (RS) interneurons	Sncg	Int 5
Földy	Hippocampus CA1 and Subiculum Pyramidal cells	Pyramidal cluster	Pyramidal cluster (excluding CA1PyrInt)
Földy	Hippocampus fast-spiking (FS) interneurons	Pvalb cluster (Pvalb Gpx3, Pvalb Wt1, Pvalb Tacr3,	Int 3

		Pvalb Tpbg, Pvalb Cpne5, Pvalb Rspo2, Pvalb Obox3)	
Fuzik	Cortex Layer 1 and 2 interneurons	Ndnf cluster (Ndnf Car4, Ndnf Cxcl14)	Int12, Int15
Fuzik	Cortex Pyramidal cells	Pyramidal cluster	Pyramidal cluster (excluding CA1PyrInt)

Supplementary Table 2: Matching of patch-seq cell types to dissociated cell reference atlases.

Broad cell type	Tasic subtypes	Zeisel subtypes
Astrocyte	Astro Gja1	Astro2, Astro1
Endothelial	Endo Myl9, Endo Tbc1d4	Vsmc
Inhibitory	Vip Chat, Vip Parm1, Vip Mybpc1, Vip Gpc3, Pvalb Gpx3, Ndnf Cxcl14, Vip Sncg, Ndnf Car4, Sst Myh8, Sst Th, Sst Chodl, Sst Tacstd2, Sst Cdk6, Pvalb Wt1, Sncg, Sst Cbln4, Pvalb Tacr3, Igtp, Smad3, Pvalb Tpbg, Pvalb Cpne5, Pvalb Rspo2, Pvalb Obox3	Int10, Int6, Int9, Int2, Int4, Int1, Int3, Int13, Int16, Int14, Int11, Int5, Int7, Int8, Int12, Int15
Microglia	Micro Ctss	Mgl1, Mgl2
Oligodendrocyte	Oligo Opalin, Oligo 96_Rik	Oligo1, Oligo3, Oligo4, Oligo2, Oligo6, Oligo5
OPC	OPC Pdgfra	*
Pyramidal	L2/3 Ptgs2, L2 Ngb, L4 Ctxn3, L4 Scnn1a, L5a Batf3, L5a Pde1c, L6a Mgp, L6b Serpinb11, L6b Rgs12, L5a Hsd11b1, L4 Arf5, L5a Tcerg1l, L6a Sla, L6a Syt17, L6a Car12, L5b Cdh13, L5 Ucma, L5b Tph2, L5 Chrna6	S1PyrL4, ClauPyr, S1PyrL5, S1PyrL23, S1PyrDL, S1PyrL5a, SubPyr, CA1Pyr1, S1PyrL6b, S1PyrL6, CA1Pyr2, CA2Pyr2

*Supplementary Table 3. Mapping of broad cell types between Tasic and Zeisel dissociated cell reference datasets. * denotes oligodendrocyte precursor cell type not being explicitly labelled in Zeisel.*

Supplementary Table 4: List of cell type-specific markers based on re-analysis of published dissociated cell-based scRNAseq experiments from mouse brain.