1 **Title**
2 Protein interaction energy landscapes are shaped by functional and also non-functional
3 partners
4
5
6 **Short Title**
7 The interaction propensity of the whole surface of proteins is conserved during evolution
8
9 **Authors**

10 Hugo Schweke[a], Marie-Hélène Mucchielli[ab], Sophie Sacquin-Mora[c], Wanying Bei[a], Anne
11 Lopes[a]
12
13
14
Authors affiliations
[a] Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud,
Université Paris-Saclay, 91198, Gif-sur-Yvette cedex, France
[b] Sorbonne Universités, UPMC Univ Paris 06, UFR927, F-75005 Paris, France.
[c] Laboratoire de Biochimie Théorique, UPR 9080 CNRS Institut de Biologie Physico-
Chimique, Paris, France
21
22
23 **Corresponding author**
24 Anne Lopes, Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-
25 Sud, Université Paris-Saclay, 1 avenue de la Terrasse, 91198 Gif-sur-Yvette, France.
26 Tel: +33 (0)1 69 15 35 60
27 email: anne.lopes@u-psud.fr

28
29
30
31 The authors have declared that no competing interests exist.

Abstract

In the crowded cell, a strong selective pressure operates on the proteome to limit the competition between functional and non-functional protein-protein interactions. Understanding how this competition constrains the behavior of proteins with respect to their partners or random encounters is very difficult to address experimentally. Here, we developed an original theoretical framework in order to investigate the propensity of protein surfaces to interact with functional and arbitrary partners and ask whether their interaction propensity is conserved during evolution. Therefore, we performed 5476 cross-docking simulations to systematically characterize the energy landscapes of 74 proteins interacting with different sets of homologs, corresponding to their functional partner's family or arbitrary protein families. Our framework relies on an original representation of interaction energy landscapes with two-dimensional energy maps that reflect the propensity of a protein surface to interact. To address the evolution of interaction energy landscapes, we systematically compared the energy maps resulting from the docking of a protein with several homologous partners. Strikingly, we show that the interaction propensity of not only binding sites but also of the rest of protein surfaces is conserved for homologous partners, and this feature holds for both functional and arbitrary partners. While most studies aiming at depicting protein-protein interactions focus on native binding sites of proteins, our analysis framework enables in an efficient and automated way, the physical characterization of not only known binding sites, but also of the rest of the protein surfaces, and provides a wealth of valuable information to understand mechanisms driving and regulating protein-protein interactions. It enables to address the energy behavior of a protein in interaction with hundreds of selected partners, providing a functional and systemic point of view of protein interactions, and opening the way for further developments to study the behavior of proteins in a specific environment.

Author Summary

In the crowded cell, the competition between functional and non-functional interactions is severe. Understanding how a protein binds the right piece in the right way in this complex jigsaw puzzle is crucial and very difficult to address experimentally. To interrogate how this competition constrains the behavior of proteins with respect to their partners or random encounters, we (i) performed thousands of cross-docking simulations to systematically characterize the interaction energy landscapes of functional and non-functional protein pairs and (ii) developed an original theoretical framework based on two-dimensional energy maps that reflect the propensity of a protein surface to interact. Strikingly, we show that the interaction propensity of not only binding sites but also of the rest of protein surfaces is conserved for homologous partners be they functional or not. We show that exploring non-functional interactions (i.e. non-functional assemblies and interactions with non-functional partners) is a viable route to investigate the mechanisms underlying protein-protein interactions. Precisely, our 2D energy maps based strategy enables it in an efficient and automated way. Moreover, our theoretical framework opens the way for the developments of a variety of applications covering functional characterization, binding site prediction, or characterization of protein behaviors in a specific environment.

Introduction

Biomolecular interactions are central for many physiological processes and are of utmost importance for the functioning of the cell. Particularly protein-protein interactions have attracted a wealth of studies these last decades [1–5]. The concentration of proteins in a cell has been estimated to be approximately 2-4 million proteins per cubic micron [6]. In such a highly crowded environment, proteins constantly encounter each other and numerous non-specific interactions are likely to occur [7,8]. For example, in the cytosol of *S. cerevisiae* a protein can encounter no less than 2000 different proteins [9]. In this complex jigsaw puzzle, each protein has evolved to bind the right piece in the right way (positive design) and to prevent misassembly and non-functional interactions (negative design) [10–14]).

Consequently, positive design constrains the physico-chemical properties and the evolution of protein-protein interfaces. Indeed, a strong selection pressure operates on binding sites to maintain the functional assembly. For example, homologs sharing at least 30% sequence identity almost invariably interact in the same way [15]. Conversely, negative design prevents proteins to be trapped in the numerous competing non-functional interactions inherent to the crowded environment of the cell. Particularly, the misinteraction avoidance shapes the evolution and physico-chemical properties of abundant proteins, resulting in a slower evolution and less sticky surfaces than what is observed for less abundant ones [16–21]. The whole surface of abundant proteins is thus constrained, preventing them to engage deleterious non-specific interactions that could be of dramatic impact for the cell at high concentration [20]. Recently, it has been shown in *E. coli* that the net charge as well as the charge distribution on protein surfaces affect the diffusion coefficients of proteins in the cytoplasm [22]. Positively charged proteins move up to 100 times more slowly as they get caught in non-

specific interactions with ribosomes which are negatively charged and therefore, shape the

composition of the cytoplasmic proteome [22].

All these studies show that both positive and negative design effectively operate on the whole

protein surface. Binding sites are constrained to maintain functional assemblies (i.e.

functional binding modes and functional partners) while the rest of the surface is constrained

to avoid non-functional assemblies. Consequently, these constraints should shape the energy

landscapes of functional but also non-functional interactions so that non-functional

interactions do not prevail over functional ones. This should have consequences (i) on the

evolution of the propensity of a protein to interact with its environment (including functional

and non-functional partners) and (ii) on the evolution of the interaction propensity of the

whole surface of proteins, non-interacting surfaces being in constant competition with

functional binding sites. We can hypothesize that the interaction propensity of the whole

surface of proteins is constrained during evolution in order to (i) ensure that proteins correctly

bind functional partners, and (ii) limit non-functional assemblies as well as interactions with

non-functional partners.

In this work, we focus on protein surfaces as a proxy for functional and non-functional

protein-protein interactions. We investigate their interaction energy landscapes with native

and non-native partners and ask whether their interaction propensity is conserved during

evolution. With this aim in mind, we performed large-scale docking simulations to

characterize interactions involving either native and/or native-related (i.e. partners of their

homologs) partners or arbitrary partners. Docking simulations enable the characterization of

all possible interactions involving either functional or arbitrary partners, and thus to simulate

the interaction of arbitrary partners which is very difficult to address with experimental

approaches. Docking algorithms are now fast enough for large-scale applications and allow

for the characterization of interaction energy landscapes for thousand of protein couples.

Typically, a docking simulation takes from a few minutes to a couple of hours on modern

processors [23–25], opening the way for extensive cross-docking experiments [26–29].

Protein docking enables the exploration of the interaction propensity of the whole protein

surface by simulating alternative binding modes. Here, we performed a cross-docking

experiment involving 74 selected proteins docked with their native-related partners and their

corresponding homologs, as well as arbitrary partners and their corresponding homologs. We

represented the interaction energy landscape resulting from each docking calculation with a

two dimensional (2D) energy map in order to (i) characterize the propensity of all surface

regions of a protein to interact with a given partner (either native-related or not) and (ii) easily

compare the energy maps resulting from the docking of a same protein with different

homologous partners, thus addressing the evolution of the propensity of the whole protein

surface to interact with homology-related partners either native or arbitrary.

Results

The interaction propensity of a protein to interact either with native-related or arbitrary partners is conserved during evolution

We ask whether the interaction propensity of a protein surface is conserved for homologous native-related partners, and whether this remains true for homologous arbitrary partners. For a protein A, we refer as native-related partners its native partner (when its three dimensional (3D) structure is available) and native partners of proteins that are homologous to the protein A. Arbitrary pairs refer to pairs of proteins for which no interaction between them or their respective homologs has been experimentally characterized in the Protein Data Bank [30]. To test the aforementioned hypothesis, we built a database comprising 74 protein structures divided into 12 families of homologs (S1 Table and *Materials and Methods*). Each family displays different degrees of structural variability and sequence divergence in order to see the impact of these properties on the conservation of the interaction propensity inside a protein family. Each family has at least a native-related partner family (S1 Fig). Docking calculations were performed with the ATTRACT software [25]. ATTRACT enables a homogeneous and exhaustive conformational sampling and is well suited to investigate the propensity of the whole surface of a protein to interact with a given ligand. Our procedure is asymmetrical since we aim at characterizing the interaction propensity of a protein (namely the receptor) with a subset of proteins (namely the ligands). Therefore, a given receptor is docked with a subset of ligands (here the 74 proteins of the dataset) (Fig 1A and *Materials and Methods*). For each docking calculation, we produced a 2D energy map, which provides the distribution of interaction energies of all docking solutions over the whole receptor surface (Fig 1B and *Materials and Methods*, Fig 2A-C). The resulting energy map reflects the propensity of the whole surface of the receptor to interact with the docked ligand. One should notice that

energy maps computed for two unrelated receptors are not comparable since their surfaces are

not comparable. Therefore, the procedure is ligand-centered and allows only the comparison

of energy maps produced by different ligands docked with the same receptor. The comparison

of two energy maps enables the evaluation of the similarity of the interaction propensity of the

receptor with the two corresponding ligands. In order to investigate the interaction propensity

of all proteins of the dataset, each protein plays alternately the role of receptor and ligand.

Consequently, the procedure presented in Fig 1 is repeated for the whole dataset where each

protein plays the role of the receptor and is docked with the 74 proteins that play the role of

ligands.

Fig 1. Experimental Protocol. (*A*) A receptor protein is docked with all proteins of the

dataset (namely the ligands) resulting in 74 docking calculations. (*B*) For each docking

calculation, an energy map is computed as well as its corresponding five-color and one-color

energy maps, with the procedure described in Fig 2 and *Materials and Methods*. (*C*) An

energy map distance (EMD) matrix is computed, representing the pairwise distances between

the 74 energy maps resulting from the docking of all ligands with this receptor. Each cell (*i,j*)

of the matrix represents the Manhattan distance between the two energy maps resulting from

the docking of ligands *i* and *j* with the receptor. A small distance indicates that the ligands *i*

and *j* produce similar energy maps when docked with this receptor. In other words, it reflects

that the interaction propensity of this receptor is similar for these two ligands. To prevent any

bias from the choice of the receptor, the whole procedure is repeated for each receptor of the

database, leading to 74 EMD matrices.

Fig 2. 2D asymmetrical representation of docking energy landscapes and resulting

energy maps. (*A*) Three-dimensional (3D) representation of the ligand docking poses around

the receptor. Each dot corresponds to the center of mass (CM) of a ligand docking pose. It is

colored according to its docking energy score. (*B*) Representation of the CM of the ligand

docking poses after an equal-area 2D sinusoidal projection. CMs are colored according to the

same scale as in A. (*C*) Continuous energy map (see *Materials and Methods* for more details).

(*D*) Five-color map. The energy map is discretized into five energy classes (*E*) One-color

maps. Top to bottom: red, orange, green, dark green and blue maps highlight respectively hot,

warm, lukewarm, cool and cold regions.

Fig 3A represents the energy maps computed for the receptor 2AYN_A, the human ubiquitin

carboxyl-terminal hydrolase 14 (family UCH) docked with (i) its native partner (1XD3_B,

ubiquitin-related family), a homolog of its partner (1NDD_B) and (ii) two arbitrary

homologous ligands (1YVB_A and 1NQD_B from the papain-like family). For all four

ligands, either native-related or arbitrary partners, docking calculations lead to an

accumulation of low-energy solutions (hot regions in red) around the two experimentally

known binding sites of the receptor. The first one corresponds to the interaction site with the

native partner, ubiquitin (pdb id 2ayo). The second one corresponds to its homodimerisation

site (pdb id 2ayn). This indicates that native-related but also arbitrary partners tend to bind

onto the native binding sites of native partners as observed in earlier studies [29,31]. The

same tendency is observed for all 74 ligands in the database (Fig 3B). Their 20 best docking

poses systematically tend to accumulate in the vicinity of the two native interaction sites.

Whereas the low-energy solutions for most ligands accumulate around the same interaction

sites (i.e. the native binding sites), we observe that, globally, 2-D energy maps (i) seem to be

more similar between ligands of a same family than between ligands belonging to different

families (Fig 3A). The two energy maps obtained with the ligands of the native-related

partners family both reveal two sharp hot regions around the native sites and a subset of well-

defined cold regions (i.e. blue regions corresponding to high energy solutions) placed in the

same area in the map's upper-right quadrant. In contrast, the energy maps obtained for the two

ligands of the papain-like family display a large hot region around the two native binding sites

of the receptor, extending to the upper-left and bottom-right regions of the map, suggesting a

large promiscuous binding region for these ligands.

Fig 3. Subset of energy maps and of ligand docking poses for receptor 2AYN_A. (*A*)

Examples of maps for the receptor 2AYN_A (ubiquitin carboxyl-terminal hydrolase (UCH)

family) docked with the ligands 1XD3_B (native partner), 1NDD_B (homolog of the native

partner), 1YVB_A and 2NQD_B (false partners). The star indicates the localization of the

experimentally determined interaction site of the ubiquitin, the circle-cross indicates the

homodimerization site of 2AYN_A. (*B*) Centers of mass (CM) of the 20 best docking poses

obtained for each of the 74 ligands of the database docked with the receptors 2AYN_A.

Receptor protein is represented in cartoon (black), its native ligand and its homodimere are

represented in cartoon with transparency (red and black respectively). CMs of the ligands

belonging to the ubiquitin-related family are colored in red, CMs of the proteins belonging to

the papain-like family are colored in blue.

We ask whether the observation made for the receptor 2AYN_A, that energy maps produced

with homologous ligands are more similar than those produced with unrelated ligands could

be generalized to all proteins of the dataset. Therefore, we systematically compared the

energy maps computed for a single receptor docked successively with the 74 ligands of the

dataset by calculating of the Manhattan distance between each pair of maps (Fig 1C and

Materials and Methods). The resulting distances are stored in an energy map distance (EMD)

matrix, where each entry (i,j) corresponds to the distance $d_{i,j}$ between the energy maps of

ligands i and j docked with the receptor of interest (Fig 1C and *Materials and Methods*).

Consequently, a small distance $d_{i,j}$ between ligands i and j docked with the receptor k, reflects

that their energy maps are similar. In other words, the interaction propensity of the surface of

the receptor k is similar for both ligands i and j. The procedure is repeated for each receptor of

the dataset resulting in 74 EMD matrices. In order to quantify the extent to which the

interaction propensity of the receptor is conserved for homologous ligands, we investigate

whether distances calculated between homologous ligand pairs (be they native-related to the

receptor or not) are smaller than distances calculated between random pairs. Fig 4 represents

the boxplots of energy map distances calculated between random ligand pairs or between

homologous ligand pairs docked with their native-related receptors or with the other receptors

of the dataset. Homologous ligands docked either with their native-related or arbitrary

receptors display significantly lower energy map distances than random ligand pairs

(Wilcoxon test $p = 0$). This indicates that energy maps produced by homologous ligands

docked with a given receptor are more similar than those produced with non-homologous

ligands. Interestingly, this observation holds whether the receptor-ligand pair is a native pair

or not. This suggests that the interaction propensity of a receptor is conserved for homologous

ligands be they native-related or not.

Fig 4. Boxplots of energy map pairwise distances between homologous ligand pairs from

native-related partner families, homologous ligand pairs from arbitrary partner families

and random ligand pairs. For each receptor, we computed (i) the average of energy map

distances of pair of homologous ligands belonging to its native-related partner family(ies), (ii)

the average of energy map distances of pair of homologous ligands belonging to its non-

native-related partner families, and (iii) the average of energy map distances of random pairs.

P-values are calculated with an unilateral Wilcoxon test.

Energy maps are specific to protein families

The results presented above prompt us to assess the extent to which the interaction propensity

of a receptor is specific to the ligand families. In other words, we quantify the extent to which

energy maps are specific to ligand families. If so, we should be able to retrieve ligand

homology relationships solely with the comparison of their corresponding 2D energy maps.

Therefore, we tested our ability to predict the homologs of a given ligand based only on the

comparison of its energy maps with those of the other ligands. In order to prevent any bias

from the choice of the receptor, the 74 EMD matrices are averaged in an averaged distances

matrix (ADM) (see *Materials and Methods*). Each entry *(i,j)* of the ADM corresponds to the

averaged distance between two sets of 74 energy maps produced by two ligands *i* and *j*. A

low distance indicates that the two ligands display similar energy maps whatever the receptor

is. We computed a receiver operating characteristic (ROC) curve from the ADM (see

Materials and Methods) which evaluates our capacity to discriminate the homologs of a given

ligand from non-homologous ligands by comparing their respective energy maps computed

with all 74 receptors of the dataset. The true positive set consists in the homologous protein

pairs while the true negative set consists in any homology-unrelated protein pair. The

resulting Area Under the Curve (AUC) is equal to 0.79 (Fig 5). We evaluated the robustness

of the ligand's homologs prediction depending on the size of the receptor subset with a

bootstrap procedure by randomly removing receptor subsets of different sizes (from 1 to 73

receptors). The resulting AUCs range from 0.769 to 0.79, and show that from a subset size of

five receptors, the resulting prediction accuracy no longer significantly varies (risk of wrongly

rejecting the equality of two variances (F-test) >5%), and is thus robust to the nature of the

receptor subset (S2 Fig). Finally, we evaluated the robustness of the predictions according to

the number of grid cells composing the energy maps. Therefore, we repeated the procedure

using energy maps with resolutions ranging from 144x72 to 48x24 cells. S2 Table presents

the AUCs calculated with different grid resolutions. The resulting AUCs range from 0.78 to

0.8 showing that the grid resolution has a weak influence on the map comparison. All

together, these results indicate that homology relationships between protein ligands can be

detected solely on the basis of the comparison of their energy maps. In other words, the

energy maps calculated for a given receptor docked with a set of ligands belonging to a same

family are specific to these families. Interestingly, this observation holds for families

displaying important sequence variations (S1 Table). For example, the AUC computed for the

UCH and ubiquitin-related families are 0.98 and 0.88 respectively despite the fact that the

average sequence identity of these families does not exceed 45% (S3 Fig and S1 Table). This

indicates that energy maps are similar even for homologous ligands displaying large sequence

variations.

Fig 5. Receiver operating characteristic (ROC) curve and its Area Under the Curve

(AUC). ROC are calculated on the averaged distance matrix (ADM) including either all pairs

(blue) or only arbitrary pairs (red) (see *Materials and Methods* for more details).

We then specifically investigate the similarity of the energy maps produced by ligands

belonging to a same family in order to see whether some ligands behave energetically

differently from their family members. On the 74 ligands, only five (2L7R_A, 4BNR_A,

1BZX_A, 1QA9_A, 1YAL_B) display energy maps that are significantly different from those

of their related homologs (Z-tests *p-values* for the comparison of the averaged distance of

each ligand with their homologs versus the averaged distance of all ligands with their

homologous ligands ≤ 5%). In order to identify the factors leading to differences between

energy maps involving homologous ligands, we computed the pairwise sequence identity and

the root mean square deviation (RMSD) between the members of each family. Interestingly,

none of these criteria can explain the energy map differences observed within families (Fisher

test p of the linear model estimated on all protein families >0.1) (see Fig 6B-C for the

ubiquitin-related family, S4-S14B-C Fig for the other families, and S3 Table for details). Fig

6A represents a subsection of the ADM for the ubiquitin-related family (i.e. the energy map

distances computed between all the members of the ubiquitin-related family and averaged

over the 74 receptors). Low distances reflect pairs of ligands with similar energy behaviors

(i.e. producing similar energy maps when interacting with a same receptor) while high

distances reveal pairs of ligands with distant energy behaviors. 2L7R_A distinguishes itself

from the rest of the family, displaying high-energy map distances with all of its homologs.

RMSD and sequence identity contribute modestly to the energy map distances observed in Fig

6A (Spearman correlation test $p^{RMSD} = 0.01$ and $p^{seq} = 0.02$ (S3 Table, Fig 6B-C)). Fig 6D

shows a projection of the contribution from the electrostatic term in the energy function of

ATTRACT on the surface of the seven ubiquitin-related family members (for more details,

see S15 Fig and *Materials and Methods*). Fig 6E represents the electrostatic maps distances

computed between all members of the family. 2L7R_A stands clearly out, displaying a

negative electrostatic potential over the whole surface while its homologs harbor a remarkable

fifty-fifty electrostatic distribution (Fig 6D). The negatively charged surface of 2L7R_A is

explained by the absence of the numerous lysines that are present in the others members of

the family (referred by black stars, Fig 6D). Lysines are known to be essential for ubiquitin

function by enabling the formation of polyubiquitin chains on target proteins. Among the

seven lysines of the ubiquitin, K63 polyubiquitin chains are known to act in non-proteolytic

events while K48, K11, and the four other lysines polyubiquitin chains are presumed to be

involved into addressing proteins to the proteasome [32]. 2L7R_A is a soluble UBL domain

resulting from the cleavage of the fusion protein FAU [33]. Its function is unrelated to

proteasomal degradation, which might explain the lack of lysines on its surface and the

differences observed in its energy maps. Interestingly, the differences observed for the energy

maps of 1YAL_B (Papain-like family) (S4 Fig) and 4BNR_A (eukaryotic proteases family)

(S5 Fig) regarding their related homologs can be explained by the fact that they both display a

highly charged surface. These two proteins are thermo-stable [34,35], which is not the case

for their related homologs, and probably explains the differences observed in their relative

energy maps. The V-set domain family is split into two major subgroups according to their

averaged energy map distances (S6A Fig). The first group corresponds to CD2 proteins

(1QA9_A and its unbound form 1HNF_A) and differs significantly from the second group (Z-

test $p = 0.03$ and $p = 0.05$ respectively). The second group corresponds to CD58 (1QA9_B

and its unbound form 1CCZ_A) and CD48 proteins (2PTT_A). Interestingly, CD2 is known

to interact with its homologs (namely CD58 and CD48) through an interface with a striking

electrostatic complementarity [36]. The two subgroups have thus evolved distinct and specific

binding sites to interact together. We can hypothesize that they have different interaction

propensities resulting in the differences observed between their corresponding energy maps.

These five cases illustrate the capacity of our theoretical framework to reveal functional or

biophysical specificities of homologous proteins that could not be revealed by classical

descriptors such as RMSD or sequence identity.

Fig 6. Ubiquitin-related family. (*A*) Energy map distances matrix. It corresponds to the

subsection of the ADM for the ubiquitin-related family (for the construction of the ADM, see

Materials and Methods). Each entry *(i,j)* represents the pairwise energy map distance of the

ligand pair (*i,j*) averaged over the 74 receptors of the dataset. (*B*) Pairwise sequence identity

matrix between all members of the family. (*C*) Pairwise root mean square deviation (RMSD)

matrix between all members of the family. (*D*) Electrostatic maps and cartoon representations

of the seven members of the family. An electrostatic map represents the distribution of the

electrostatic potential on the surface of a protein (for more details, see S15 Fig and *Materials*

and Methods). On the electrostatic maps, lysines positions are indicated by stars. Cartoon

structures are colored according to the distribution of their electrostatic potential. (*E*)

Electrostatic map distances matrix. Each entry *(i,j)* of the matrix represents the Manhattan

distance between the electrostatic maps of the proteins *(i,j)*.

The AUC of 0.79 calculated previously with energy maps produced by the docking of either

native-related or arbitrary pairs indicates that energy maps are specific to ligand families. To

see whether this observation is not mainly due to the native-related pairs, we repeated the

previous test while removing that time all energy maps computed with native-related pairs

and calculated the resulting ADM. We then measured our ability to retrieve the homologs of

each ligand by calculating the ROC curve as previously. The resulting AUC is still equal to

0.79, revealing that our ability to identify a ligand's homologs is independent from the fact

that the corresponding energy maps were computed with native-related or arbitrary pairs (Fig

5). This shows that the energy maps are specific to protein families whether the docked pairs

are native-related or not. Consequently, the propensity of the whole protein surface to interact

with a given ligand is conserved and specific to the ligand family whether the ligand is native-

related or not. This striking result may reflect both positive and negative design operating on

protein surfaces to maintain functional interactions and to limit random interactions that are

inherent to a crowded environment.

The interaction propensity of all surface regions of a receptor is evolutionary conserved for homologous ligands

To see whether some regions contribute more to the specificity of the maps produced by homologous ligands, we next dissected the effective contribution of the surface regions of the receptor defined according to their docking energy value, in the identification of ligand's homologs. We discretized the energy values of each energy map into five categories, leading to a palette of five energy classes (or colors) (see Fig 2D and *Materials and Methods*). These five-color maps highlight low-energy regions (i.e. hot regions in red), intermediate-energy regions (i.e. warm, lukewarm and cool regions in orange, light-green and dark-green respectively) and high-energy regions (i.e. cold regions in blue). We first checked that the discretization of the energy maps does not affect our ability to identify the homologs of each of the 74 ligands from the comparison of their five-colors maps. The resulting AUC is 0.77 (Table 1)**,** showing that the discretization step does not lead to an important loss of information.

Table 1. AUC obtained with different types of energy maps.

type of map	continuous energy maps	five-colors energy maps	red energy maps	orange energy maps	light green energy maps	dark green energy maps	blue energy maps
AUC	0.79	0.77	0.73	0.76	0.76	0.76	0.79

The AUC are calculated from the ADM with the continuous energy maps (Fig 2C), the five-color energy maps (Fig 2D) and the one-color energy maps (Fig 2E) (see *Materials and Methods* for more details).

Then, we evaluated the contribution of each of the five energy classes separately in the

ligand's homologs identification by testing our ability to retrieve the homologs of the 74

ligands from their one-color energy maps (either red, orange, yellow, green or blue) (see

Materials and Methods). Table 1 shows the resulting AUCs. Interestingly, the information

provided by each energy class taken separately is sufficient for discriminating the homologs

of a given ligand from the rest of the dataset (Table 1). The resulting AUCs range from 0.76

to 0.79 for the orange, light green, dark green, and blue classes and are comparable to those

obtained with all classes taken together (0.77). This shows that (i) warm, lukewarm, cool, and

cold regions alone are sufficient to retrieve homology relationships between ligands and (ii)

the localization on the receptor surface of a given energy class is specific to the ligand

families. Hot regions are less discriminative and lead to an AUC of 0.73. In order to see how

regions corresponding to a specific energy class are distributed over a receptor surface, we

summed its 74 corresponding one-color maps into a stacked map (S16 Fig – see *Materials*

and Methods for more details). For each color, the resulting stacked map reflects the tendency

of a map cell to belong to the corresponding energy class. Fig 7 shows an example of the five

stacked maps (i.e. for cold, cool, lukewarm, warm and hot regions) computed for the receptor

1P9D_U. Intermediates regions (i.e. warm, lukewarm and cool regions) are widespread on the

stacked map while cold and hot regions are localized on few small spots (three and one

respectively) no matter the nature of the ligand. S17 Fig shows for the receptor 1P9D_U the

12 blue and red stacked maps computed for each ligand family separately. We can see that

some cold spots are specific to ligand families and that their area distribution is specific to

families while all 12 ligand families display the same hot spot in the map's upper-right

quadrant. These observations can be generalized to each receptor. On average, intermediate

regions are widespread on the stacked maps and cover respectively 744, 1164 and 631 cells

for cool, lukewarm and warm regions, while cold and hot regions cover no more than

respectively 104 and 110 cells respectively (S18 Fig). Interestingly, hot regions are more

colocalized than cold ones and are restricted to 2 distinct spots on average per stacked map,

while cold regions are spread on 3.7 spots on average (t-Test $p = 7.42e\text{-}13$). These results

show that ligands belonging to different families tend to dock preferentially on the same

regions and thus lead to similar hot region distributions on the receptor surface. This

observation recalls those made by *Fernandez-Recio et al.* [31], who showed that docking

random proteins against a single receptor leads to an accumulation of low-energy solutions

around the native interaction site and who suggested that different ligands will bind

preferentially on the same localization.

Fig 7. Stacked maps of 1P9D_U after the filtering of cells with too low intensity and

areas of too small size. The protocol to generate stacked maps is presented in S16 Fig. (*A*)

Blue stacked map (i.e. stacked cold regions). (*B*) Dark green stacked map (i.e. stacked cool

regions). (*C*) Light green stacked map (i.e. stacked lukewarm regions). (*D*) Orange stacked

map (i.e. stacked warm regions). (*E*) Red stacked map (i.e. stacked hot regions). One should

notice that stacked maps of two different colors can overlap because a cell can be associated

to different energy classes depending on the docked ligands. S17 Fig presents blue and red

stacked maps of 1P9D_U computed for each ligand family.

We can hypothesize that hot regions present universal structural and biochemical features that

make them more prone to interact with other proteins. To test this hypothesis, we computed

for each protein of the dataset, the 2D projection of three protein surface descriptors (see

Materials and Methods and S15 Fig): the Kyte-Doolittle (KD) hydrophobicity [37], the

circular variance (CV) [38] and the stickiness [20]. The CV measures the density of protein

around an atom and is a useful descriptor to reflect the local geometry of a surface region. CV

values are comprised between 0 and 1. Low values reflect protruding residues and high values

indicate residues located in cavities. Stickiness reflects the propensity of amino acids to be

involved in protein-protein interfaces [20]. It has been calculated as the log ratio of the

residues frequencies on protein surfaces versus their frequencies in protein-protein interfaces.

For each receptor, we calculated the correlation between the docking energy and the

stickiness, hydrophobicity or CV over all cells of the corresponding 2D maps. We found a

significant anti-correlation between the docking energy and these three descriptors

(correlation test p between docking energies and respectively stickiness, hydrophobicity and

CV < 2.2e-16, see S4 Table)). Fig 8 represents the boxplots of the stickiness, hydrophobicity

and CV of each energy class (see S15 Fig and *Materials and Methods* section for more

details). We observe a clear effect of these factors on the docking energy: cold regions (i.e.

blue class) are the less sticky, the less hydrophobic and the most protruding while hot ones

(i.e. red class) are the most sticky, the most hydrophobic and the most planar (Tukey HSD test

[39], p of the differences observed between each energy classes < 2.2e-16). One should notice

that stickiness has been defined from a statistical analysis performed on experimentally

characterized protein interfaces and therefore between presumed native partners. The fact that

docking energies (physics-based) calculated either between native-related or arbitrary partners

is anti-correlated with stickiness (statistics-based) defined from native interfaces, strengthens

strongly the concept of stickiness as the propensity of interacting promiscuously and provides

physics-based pieces of evidence for sticky regions as a proxy for promiscuous interactions.

We show that not only the area distribution on a receptor surface of hot regions but also those

of intermediate and cold regions are similar for homologous ligands and are specific to ligand

families (AUC ranging from 0.73 to 0.79) whether the ligands are native-related or not. This

tendency is even stronger for intermediate and cold regions. Interestingly, the information

contained in the cold regions that cover on average no more than 5.0% of the energy maps is

sufficient to identify homology relationships between ligands.

Fig 8. Boxplots of three descriptors of the protein surface. (*A*) the stickiness values, (*B*) the

Kyte-Doolittle hydrophobicity and (*C*) the CV values, depending on the energy class. The

stickiness, hydrophobicity and CV values are calculated for each protein following the

protocol described in *Materials and Methods*. For each of these criteria, *p-values* between the

median values of two "successive" energy classes were computed using the Tukey HSD

statistical test [39].

Discussion

In this study, we address the impact of both positive and negative design on thousands of

interaction energy landscapes by the mean of a synthetic and efficient representation of the

docking energy landscapes: two-dimensional energy maps that reflect the interaction

propensity of the whole surface of a protein (namely the receptor) with a given partner

(namely the ligand). We show that all regions of the energy maps, including cold,

intermediate and hot regions are similar for homologous ligands and are specific to ligand

families whether the ligands are native-related or arbitrary. This reveals that the interaction

propensity of the whole surface of proteins is constrained by functional and non-functional

interactions, reflecting both positive and negative design operating on the whole surface of

proteins, thus shaping the interaction energy landscapes of functional partners and random

encounters. These observations were made on a dataset of 74 protein structures belonging to

12 families of structural homologs. 54 out of the 74 proteins of the dataset have at least one

known partner in the dataset. For the 20 remaining proteins, we were not able to find

evidences that they indeed interact with a protein of the dataset. However, we showed that the

interaction propensity of a receptor is conserved for homologous ligands independently from

the fact that these ligands correspond to native partners or not. Indeed, we showed that ligand

homology relationships could be retrieved from their energy maps whether the maps were

computed with native-related pairs or not (the corresponding AUCs calculated with and

without native pairs both equal to 0.79).

While most studies that aim at depicting protein-protein interactions focus on native binding

sites of proteins [12,40–44], we bring a new perspective on protein-protein interactions by

providing a systematic and physical characterization of all regions of the surface of a protein

in interaction with a given ligand (i.e. cold, intermediate and hot regions). Here, we address

the energy behavior of not only known binding sites, but also of the rest of the protein surface,

which plays an important role in protein interactions by constantly competing with the native

binding site. We show that the interaction propensity of the rest of the surface is not

homogeneous and displays regions with different binding energies that are specific to ligand

families. This may reflect the negative design operating on these regions to limit non-

functional interactions [12,14,45]. We can hypothesize that non-interacting regions participate

to favor functional assemblies (i.e. functional assembly modes with functional partners) over

non-functional ones and are thus evolutionary constrained by non-functional assemblies. The

fact that cold regions seem to be more specific to ligand families than hot ones may be

explained by the fact that they are on average more protuberant and more charged. They thus

display more variability than hot ones. Indeed, there is more variability in being positively or

negatively charged and protuberant (with an important range of protuberant shapes) than in

being neutral and flat. S19 Fig presents the electrostatic potential distribution of all energy

classes. Cold regions display a larger variability of electrostatic potential (F-test, $p < 2.2e\text{-}16$)

than hot regions that are mainly hydrophobic thus displaying neutral charge distributions in

average. Consequently, a same hot region may be attractive for a large set of ligands while a

cold region may be unfavorable to specific set of ligands, depending on their charges, shapes

and other biophysical properties.

On the other hand, we show that hot regions are very localized (4.9% of the cells of an energy

map) and tend to be similar no matter the ligand. Similarly to protein interfaces that have been

extensively characterized in previous studies [2,40-43], hot regions are likely to display

universal properties of binding, i.e. they are more hydrophobic and more planar, and thus

more "sticky" than the other regions. They may provide a non-specific binding patch that is

suitable for many ligands. However, we can hypothesize that native partners have evolved to

optimize their interfaces (positive design) so that native interactions prevail over non-native

competing ones. Indeed, we have previously shown that the docking of native partners lead to

more favorable binding energies than the docking of non-native partners when the ligand is

constrained to dock around the receptor's native binding site [28,46]. All these results suggest

a new physical model of protein surfaces where protein surface regions, in the crowded

cellular environment, serve as a proxy for regulating the competition between functional and

non-functional interactions. In this model, intermediate and cold regions play an important

role by preventing non-functional assemblies and by guiding the interaction process towards

functional ones and hot regions may select the functional assembly among the competing ones

through optimized interfaces with the native partner.

In this work, we used and extended the application of the 2D energy map representation

developed in [31] to develop an original theoretical framework that enables the efficient,

automated and integrative analysis of different protein surface features. 2D maps provide the

area distribution of a given feature on the whole protein surface and their discretization

enables the study of a given surface property (e.g. protuberance, planarity, stickiness,

positively charged regions, or cold and hot regions for example). They are easy to manipulate

and their straightforward comparison enables (i) the study of relationships between different

surface properties through the comparison of their area distributions on a protein surface and

(ii) the highlight of the evolutionary constraints exerted on a given feature by comparing its

area distribution on the surfaces of homologous proteins. Particularly, this enables the

identification and characterization of hot regions on a protein surface which can be either

specific or conserved for all ligands and opens up new possibilities for the development of

novel methods for protein binding sites prediction and their classification as functional or

promiscuous in the continuity of previous developments based on arbitrary docking

[28,29,31,46].

Our framework provides a proxy for further protein functional characterization as shown with

the five proteins discussed in the *Results* section *Energy maps are specific to protein families*.

The comparison of their respective energy maps enables us to reveal biophysical and

functional properties that could not be revealed with classical monomeric descriptors such as

RMSD or sequence identity. Indeed, our framework can reflect the energy behavior of a

protein interacting with a subset of selected partners either functional or arbitrary, thus

revealing functional and systemic properties of proteins. This work goes beyond the classical

use of binary docking to provide a systemic point of view of protein interactions, for example

by exploring the propensity of a protein to interact with hundreds of selected ligands, and thus

addressing the behavior of a protein in a specific cellular environment. Particularly, exploring

the dark interactome (i.e. non-functional assemblies and interactions with non-functional

partners) can provide a wealth of valuable information to understand mechanisms driving and

regulating protein-protein interactions. Precisely, our 2D energy maps based strategy enables

its exploration in an efficient and automated way.

Materials and Methods

Protein dataset

The dataset comprises 74 protein structures divided into 12 families of structural homologs (see S1 Table for a detailed list of each family). Each family is related to at least one other family (its native-related partners family) through a pair of interacting proteins for which the 3D structure of the complex is characterized experimentally (except the V set domain family: the two native partners are homologous and belong to the same family) (S1 Fig). Each family is composed of a monomer selected from the protein-protein docking benchmark 5.0 [47] in its bound and unbound forms, which is called the master protein. Each master protein has a native partner (for which the 3D structure of the corresponding complex has been characterized experimentally) in the database, which is the master protein for another family, except the V set domain family, which is a self-interacting family. When available, we completed families with interologs (i.e. pairs of proteins which have interacting homologs in an other organism) selected in the INTEREVOL database [48] according to the following criteria: (i) experimental structure resolution better than 3.25 Å, (ii) minimum alignment coverage of 75% with the rest of the family members and (iii) minimum sequence identity of 30% with at least one member of the family. Since we were limited by the number of available interologs, we completed families with unbound monomers homologous to the master following the same criteria and by searching for their partners in the following protein-protein interactions databases [49–54]. We consider that all members of a family correspond to native-related partners of all members of their native-related partner family. To address the impact of conformational changes of a protein on its interaction energy maps, we added different NMR conformers. We show that energy maps involving pairs of conformers are significantly more similar than those obtained for other pairs of homologous ligands

(unilateral Wilcoxon test, $p < 2.2e\text{-}16$) showing that the conformational changes in a protein

(lower than 3Å) have a low impact on the resulting energy maps (S20 Fig).

Docking experiment and construction of energy maps

A complete cross-docking experiment was realized with the ATTRACT software [25] on the

74 proteins of the dataset, leading to 5476 (74 x 74) docking calculations (Fig 1A).

ATTRACT uses a coarse-grain reduced protein representation and a simplified energy

function comprising a pseudo Lennard-Jones term and an electrostatic term. The calculations

took approximately 20000 hours on a 2.7GHz processor. Prior to docking calculations, all

PDB structures were prepared with the DOCKPREP software [55].

During a docking calculation, the ligand L_i explores exhaustively the surface of the receptor

R_k (whose position is fixed during the procedure), sampling and scoring thousands of different

ligand docking poses (between 10000 and 50000 depending on the sizes of the proteins) (Fig

2A). For each protein couple R_k-L_i, a 2D energy map is computed which shows the

distribution of the energies of all docking solutions over the receptor surface. To compute

these maps, for all docking poses, the spherical coordinates (ϕ, θ) (with respect to the

receptor center of mass (CM)) of the ligand CM are represented onto a 2D map in an equal-

area 2D sinusoidal projection (Fig 2B) (see [31] for more details). Each couple of coordinates

(ϕ, θ) is associated with the energy of the corresponding docking conformation (Fig 2B). A

continuous energy map is then derived from the discrete one, where the map is divided into a

grid of 36 x 72 cells. Each cell represents the same surface and, depending on the size of the

receptor, can span from 2.5 $Å^2$ to 13$Å^2$. For each cell, all solutions with an energy score below

2.7 kcal/mol^{-1} from the lowest solution of the cell are retained, according to the conformations

filtering protocol implemented in [28]. The average of the retained energy scores is then

assigned to the cell. If there is no docking solution in a cell, a score of 0 is assigned to it.

Finally, the energies of the cells are smoothed, by averaging the energy values of each cell

and of the eight surrounding neighbors (Fig 2C).

For each map, the energy values are discretized into five energy classes of same range leading

to a discrete five-colors energy map (Fig 2D). The range is calculated for each energy map

and spans from the minimum to the maximum scores of the map cells. The range of the

energy classes of the map R_k-L_i is equal to (maxE – minE)/5, where maxE and minE

correspond to the maximal and minimal energy values in the R_k-L_i map. Each five-colors

energy map is then split into five one-color maps, each one representing an energy class of the

map (Fig 2E). The continuous, five-colors and one-color energy maps are calculated for the

5476 energy maps.

Comparison of energy maps and identification of ligand's homologs

Since, we cannot compare energy maps computed for two unrelated receptors, the procedure

is ligand-centered and only compares energy maps produced with different ligands docked

with the same receptor. The referential (i.e. the receptor) is thus the same (in other words all

grid cells are comparable) for all the energy maps that are compared. For each receptor R_k, we

computed a 74x74 energy map distance (EMD) matrix where each entry (*i,j*) corresponds to

the pairwise distance between the energy maps R_k-L_i and R_k-L_j resulting from the docking of

the ligands L_i and L_j on the receptor R_k (Fig 1). The pairwise distance $d_{Man}(R_k$-L_i, R_k-$L_j)$

between the energy maps is calculated with a Manhattan distance according to equation (1)

$$d_{Man}\left(R_k L_i, R_k L_j\right) = \sum_{n=1}^{36} \sum_{m=1}^{72} |a_{nm} - b_{nm}| \quad (1)$$

where a_{nm} and b_{nm} are the cells of row index n and column index m of the energy maps R_k-L_i

and R_k-L_j respectively. Low distances reflect pairs of ligands that induce similar energy maps

when they are docked on the same receptor. The procedure presented in Fig 1 is repeated for

each receptor of the database resulting in 74 EMD matrices. The 74 EMD matrices are

averaged into an averaged distances matrix (ADM). Each entry (i,j) of the ADM reflects the

similarity of the R_k-L_i and R_k-L_j energy maps averaged over all the receptors R_k in the dataset.

In order to estimate the extent to which family members display similar energy maps when

they are docked with the same receptor, we tested our ability to correctly identify the

homologs of the 74 ligands from the only comparison of its energy maps with those of the

other ligands. Because, energy maps are receptor-centered, we cannot compare the energy

maps computed for two unrelated receptors. The procedure consists in the comparison of

energy maps produced with different ligands docked with a same receptor. Two ligands (i,j)

are predicted as homologs according to their corresponding distance (i,j) in the ADM. Values

close to zero should reflect homologous ligand pairs, while values close to one should reflect

unrelated ligand pairs. A Receiver Operating Characteristic (ROC) curve and its Area Under

the Curve (AUC) are computed from the ADM. True positives (TP) are all the homologous

ligand pairs and predicted as such, true negatives (TN) are all the unrelated ligand pairs and

predicted as such. False positives (FP) are unrelated ligand pairs but incorrectly predicted as

homologous pairs. False negatives (FN) are homologous ligand pairs but incorrectly predicted

as unrelated pairs. ROC curves and AUC values were calculated with the R package pROC

[56]. The ligand's homologs identification was also realized using the five-color energy maps

or the one-color energy maps taken separately. The five energy class regions display very

different sizes, with median ranging from 63 and 66 cells for the blue and red regions to 633

cells for the yellow one. To prevent any bias due to the size of the different classes, we

normalized the Manhattan distance by the size of the regions compared in the map. The rest of

the procedure is the same than those used for continuous energy maps (Fig 1).

To visualize the area distribution of the regions of a given energy class for all ligands on the

receptor surface, the 74 corresponding one-color maps are summed into a stacked map where

each cell's intensity varies from 0 to 74 (S16 Fig). To remove background-image from these

maps, i.e. cells with low intensity (intensity < 17) and the areas of small size (< 4 cells), we

used a Dirichlet process mixture model simulation for image segmentation (R package

dpmixsim) [57].

2D projection of monomeric descriptors of protein surfaces

We computed KD hydrophobicity [37], stickiness [20], CV [38] maps of each protein of the

dataset, in order to compare their topology with the energy maps. Prior to all, proteins

belonging to the same families were structurally aligned with TM-align [58] in order to place

them in the same reference frame, making their maps comparable. Particles were generated

around the protein surface with a slightly modified Shrake-Rupley algorithm [59]. The density

of spheres is fixed at 1Å^2, representing several thousands particles per protein. Each particle is

located at 5Å from the surface of the protein. The CV, stickiness and KD hydrophobicity

values of the closest atom of the protein are attributed to each particle. We also generated

electrostatic maps reflecting the distribution of the contribution of the coulombic term as

encoded in the ATTRACT force field on a protein surface. The procedure is slightly different:

each particle *i* has a +1 positive charge, and receives the coulombic value Q_i (see equation

(2)).

$$Q_i = \sum_{j=1}^{n} q_i q_j / \mathcal{E} r_{ij} \quad (2)$$

with n the number of pseudo-atom in the protein, q_i the charge of the particle, q_j the charge of

the pseudo-atom j, r_{ij} the distance between the particle i and the pseudo-atom j, and ε a

distant-dependent dielectric constant ($\varepsilon = 15 r_{ij}$). CV was calculated following the protocol

described in [38] on the all-atom structures. Stickiness, electrostatics and hydrophobicity were

calculated on ATTRACT coarse-grain models. Pseudo-atom charges are defined according to

the ATTRACT force field [25]. After attributing a value to each particle, the position of their

spherical coordinates is represented in a 2-D sinusoidal projection, following the same

protocol as described in Fig 2 and *Materials and Methods* section *Docking experiment and*

construction of energy maps. The map is then smoothed following the protocol in Fig 2.

Acknowledgments

We thank F. Fraternali, R. Guerois, E. Laine, and M. Montes for their constructive comments on the manuscript.

References

[1].Garzón JI, Deng L, Murray D, Shapira S, Petrey D, Honig B. A computational interactome and functional annotation for the human proteome. ELife Sciences 2016;5:e18715. doi:10.7554/eLife.18715.

[2] Janin J, Bahadur RP, Chakrabarti P. Protein-protein interaction and quaternary structure. Q Rev Biophys 2008;41:133–80. doi:10.1017/S0033583508004708.

[3] Nobeli I, Favia AD, Thornton JM. Protein promiscuity and its implications for biotechnology. Nature Biotechnology 2009;27:157–67. doi:10.1038/nbt1519.

[4] Nooren IMA, Thornton JM. Diversity of protein–protein interactions. The EMBO Journal 2003;22:3486–92. doi:10.1093/emboj/cdg359.

[5] Robinson CV, Sali A, Baumeister W. The molecular sociology of the cell. Nature 2007;450:973–82. doi:10.1038/nature06523.

[6] Milo R. What is the total number of protein molecules per cell volume? A call to rethink some published values. Bioessays 2013;35:1050–5. doi:10.1002/bies.201300066.

[7] McGuffee SR, Elcock AH. Diffusion, crowding & protein stability in a dynamic molecular model of the bacterial cytoplasm. PLoS Comput Biol 2010;6:e1000694. doi:10.1371/journal.pcbi.1000694.

[8] Yu I, Mori T, Ando T, Harada R, Jung J, Sugita Y, et al. Biomolecular interactions modulate macromolecular structure and dynamics in atomistic model of a bacterial cytoplasm. Elife 2016;5. doi:10.7554/eLife.19274.

[9] Levy ED, Kowarzyk J, Michnick SW. High-resolution mapping of protein concentration reveals principles of proteome architecture and adaptation. Cell Rep 2014;7:1333–40. doi:10.1016/j.celrep.2014.04.009.

[10] Richardson JS, Richardson DC. Natural beta-sheet proteins use negative design to avoid edge-to-edge aggregation. Proc Natl Acad Sci USA 2002;99:2754–9. doi:10.1073/pnas.052706099.

[11] Deeds EJ, Ashenberg O, Gerardin J, Shakhnovich EI. Robust protein protein interactions in crowded cellular environments. Proc Natl Acad Sci USA 2007;104:14952–7. doi:10.1073/pnas.0702766104.

[12] Pechmann S, Levy ED, Tartaglia GG, Vendruscolo M. Physicochemical principles that regulate the competition between functional and dysfunctional association of proteins. Proc Natl Acad Sci USA 2009;106:10159–64. doi:10.1073/pnas.0812414106.

[13] Karanicolas J, Corn JE, Chen I, Joachimiak LA, Dym O, Peck SH, et al. A de novo protein binding pair by computational design and directed evolution. Mol Cell 2011;42:250–60. doi:10.1016/j.molcel.2011.03.010.

[14] Garcia-Seisdedos H, Empereur-Mot C, Elad N, Levy ED. Proteins evolve on the edge of supramolecular self-assembly. Nature 2017;548:244–7. doi:10.1038/nature23320.

[15] Aloy P, Ceulemans H, Stark A, Russell RB. The relationship between sequence and interaction divergence in proteins. J Mol Biol 2003;332:989–98.

[16] Pál C, Papp B, Hurst LD. Highly expressed genes in yeast evolve slowly. Genetics 2001;158:927–31.

[17] Drummond DA, Wilke CO. Mistranslation-induced protein misfolding as a dominant constraint on coding-sequence evolution. Cell 2008;134:341–52. doi:10.1016/j.cell.2008.05.042.

[18] Zhang J, Maslov S, Shakhnovich EI. Constraints imposed by non-functional protein–protein interactions on gene expression and proteome size. Molecular Systems Biology 2008;4:210.

[19] Heo M, Maslov S, Shakhnovich E. Topology of protein interaction network shapes protein abundances and strengths of their functional and nonspecific interactions. Proc Natl Acad Sci USA 2011;108:4258–63. doi:10.1073/pnas.1009392108.

[20] Levy ED, De S, Teichmann SA. Cellular crowding imposes global constraints on the chemistry and evolution of proteomes. Proc Natl Acad Sci USA 2012;109:20461–6. doi:10.1073/pnas.1209312109.

[21] Yang J-R, Liao B-Y, Zhuang S-M, Zhang J. Protein misinteraction avoidance causes highly expressed proteins to evolve slowly. Proc Natl Acad Sci USA 2012;109:E831-840. doi:10.1073/pnas.1117408109.

[22] Schavemaker PE, Śmigiel WM, Poolman B. Ribosome surface properties may impose limits on the nature of the cytoplasmic proteome. Elife 2017;6. doi:10.7554/eLife.30084.

[23] Ritchie DW, Venkatraman V. Ultra-fast FFT protein docking on graphics processors. Bioinformatics 2010;26:2398–405. doi:10.1093/bioinformatics/btq444.

[24] Pierce BG, Hourai Y, Weng Z. Accelerating protein docking in ZDOCK using an advanced 3D convolution library. PLoS ONE 2011;6:e24657. doi:10.1371/journal.pone.0024657.

[25] de Vries S, Zacharias M. Flexible docking and refinement with a coarse-grained protein model using ATTRACT. Proteins 2013;81:2167–74. doi:10.1002/prot.24400.

[26] Wass MN, Fuentes G, Pons C, Pazos F, Valencia A. Towards the prediction of protein interaction partners using physical docking. Mol Syst Biol 2011;7:469. doi:10.1038/msb.2011.3.

[27] Ohue M, Matsuzaki Y, Shimoda T, Ishida T, Akiyama Y. Highly precise protein-protein interaction prediction based on consensus between template-based and de novo docking methods. BMC proceedings, vol. 7, BioMed Central; 2013, p. S6.

[28] Lopes A, Sacquin-Mora S, Dimitrova V, Laine E, Ponty Y, Carbone A. Protein-protein interactions in a crowded environment: an analysis via cross-docking simulations and evolutionary information. PLoS Comput Biol 2013;9:e1003369. doi:10.1371/journal.pcbi.1003369.

[29] Vamparys L, Laurent B, Carbone A, Sacquin-Mora S. Great interactions: How binding incorrect partners can teach us about protein recognition and function. Proteins 2016;84:1408–21. doi:10.1002/prot.25086.

[30] Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, et al. The Protein Data Bank. Nucleic Acids Res 2000;28:235–42.

[31] Fernandez-Recio J, Totrov M, Abagyan R. Identification of protein–protein interaction sites from docking energy landscapes. Journal of Molecular Biology 2004;335:843–865.

[32] Xu P, Duong DM, Seyfried NT, Cheng D, Xie Y, Robert J, et al. Quantitative Proteomics Reveals the Function of Unconventional Ubiquitin Chains in Proteasomal Degradation. Cell 2009;137:133–45. doi:10.1016/j.cell.2009.01.041.

[33] Welchman RL, Gordon C, Mayer RJ. Ubiquitin and ubiquitin-like proteins as multifunctional signals. Nat Rev Mol Cell Biol 2005;6:599–609. doi:10.1038/nrm1700.

[34] Molnár T, Vörös J, Szeder B, Takáts K, Kardos J, Katona G, et al. Comparison of complexes formed by a crustacean and a vertebrate trypsin with bovine pancreatic trypsin inhibitor - the key to achieving extreme stability? FEBS J 2013;280:5750–63. doi:10.1111/febs.12491.

[35] Sumner IG, Harris GW, Taylor MA, Pickersgill RW, Owen AJ, Goodenough PW. Factors effecting the thermostability of cysteine proteinases from Carica papaya. Eur J Biochem 1993;214:129–34.

[36] Wang JH, Smolyar A, Tan K, Liu JH, Kim M, Sun ZY, et al. Structure of a heterophilic adhesion complex between the human CD2 and CD58 (LFA-3) counterreceptors. Cell 1999;97:791–803.

[37] Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a protein. J Mol Biol 1982;157:105–32.

[38] Mezei M. A new method for mapping macromolecular topography. Journal of Molecular Graphics and Modelling 2003;21:463–72. doi:10.1016/S1093-3263(02)00203-6.

[39] Tukey JW. Comparing Individual Means in the Analysis of Variance. Biometrics 1949;5:99–114. doi:10.2307/3001913.

[40] Lo Conte L, Chothia C, Janin J. The atomic structure of protein-protein recognition sites. J Mol Biol 1999;285:2177–98.

[41] Chakrabarti P, Janin J. Dissecting protein–protein recognition sites. Proteins: Structure, Function, and Bioinformatics 2002;47:334–343.

[42] Li X, Keskin O, Ma B, Nussinov R, Liang J. Protein–protein interactions: hot spots and structurally conserved residues often locate in complemented pockets that pre-organized in the unbound states: implications for docking. Journal of Molecular Biology 2004;344:781–795.

[43] Keskin O, Ma B, Nussinov R. Hot regions in protein--protein interactions: the organization and contribution of structurally conserved hot spot residues. J Mol Biol 2005;345:1281–94. doi:10.1016/j.jmb.2004.10.077.

[44] Andreani J, Faure G, Guerois R. Versatility and invariance in the evolution of homologous heteromeric interfaces. PLoS Comput Biol 2012;8:e1002677. doi:10.1371/journal.pcbi.1002677.

[45] Kastritis PL, Rodrigues JPGLM, Folkers GE, Boelens R, Bonvin AMJJ. Proteins feel more than they see: fine-tuning of binding affinity by properties of the non-interacting surface. J Mol Biol 2014;426:2632–52. doi:10.1016/j.jmb.2014.04.017.

[46] Sacquin-Mora S, Carbone A, Lavery R. Identification of protein interaction partners and protein–protein interaction sites. Journal of Molecular Biology 2008;382:1276–1289.

[47] Vreven T, Moal IH, Vangone A, Pierce BG, Kastritis PL, Torchala M, et al. Updates to the integrated protein–protein interaction benchmarks: docking benchmark version 5 and affinity benchmark version 2. Journal of Molecular Biology 2015;427:3031–3041.

[48] Faure G, Andreani J, Guerois R. InterEvol database: exploring the structure and evolution of protein complex interfaces. Nucleic Acids Res 2012;40:D847-856. doi:10.1093/nar/gkr845.

[49] Salwinski L, Miller CS, Smith AJ, Pettit FK, Bowie JU, Eisenberg D. The Database of Interacting Proteins: 2004 update. Nucleic Acids Res 2004;32:D449–51. doi:10.1093/nar/gkh086.

[50] Alfarano C, Andrade CE, Anthony K, Bahroos N, Bajec M, Bantoft K, et al. The Biomolecular Interaction Network Database and related tools 2005 update. Nucleic Acids Res 2005;33:D418–24. doi:10.1093/nar/gki051.

[51] Güldener U, Münsterkötter M, Oesterheld M, Pagel P, Ruepp A, Mewes H-W, et al. MPact: the MIPS protein interaction resource on yeast. Nucleic Acids Res 2006;34:D436–41. doi:10.1093/nar/gkj003.

[52] Stark C, Breitkreutz B-J, Reguly T, Boucher L, Breitkreutz A, Tyers M. BioGRID: a general repository for interaction datasets. Nucleic Acids Res 2006;34:D535–9. doi:10.1093/nar/gkj109.

[53] Kerrien S, Alam-Faruque Y, Aranda B, Bancarz I, Bridge A, Derow C, et al. IntAct—open source resource for molecular interaction data. Nucleic Acids Res 2007;35:D561–5. doi:10.1093/nar/gkl958.

[54] Alonso-López D, Gutiérrez MA, Lopes KP, Prieto C, Santamaría R, De Las Rivas J. APID interactomes: providing proteome-based interactomes with controlled quality for multiple species and derived networks. Nucleic Acids Res 2016;44:W529–35. doi:10.1093/nar/gkw363.

[55] Lang PT, Brozell SR, Mukherjee S, Pettersen EF, Meng EC, Thomas V, et al. DOCK 6: combining techniques to model RNA-small molecule complexes. RNA 2009;15:1219–30. doi:10.1261/rna.1563609.

[56] Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J-C, et al. pROC: an open-source package for R and S+ to analyze and compare ROC curves. BMC Bioinformatics 2011;12:77. doi:10.1186/1471-2105-12-77.

[57] Ferreira da Silva AR. A Dirichlet process mixture model for brain MRI tissue classification. Med Image Anal 2007;11:169–82. doi:10.1016/j.media.2006.12.002.

[58] Zhang Y, Skolnick J. TM-align: a protein structure alignment algorithm based on the TM-score. Nucleic Acids Res 2005;33:2302–9. doi:10.1093/nar/gki524.

[59] Saladin A, Fiorucci S, Poulain P, Prévost C, Zacharias M. PTools: an opensource molecular docking library. BMC Struct Biol 2009;9:27. doi:10.1186/1472-6807-9-27.

[60] Fox NK, Brenner SE, Chandonia J-M. SCOPe: Structural Classification of Proteins--extended, integrating SCOP and ASTRAL data and classification of new structures. Nucleic Acids Res 2014;42:D304-309. doi:10.1093/nar/gkt1240.

[61] Snedecor GW, Cochran WC. Statistical methods. Iowa state university press, Ames. Iowa N Vadivukkarasi et Al 1989.

Supporting information

S1 Table. List of proteins of the dataset and their structural families. Proteins are referred by their PDB identifiers, followed by their chain identifier. The NMR conformers are referred with their conformation identifier. The conformational state of the structures are indicated in brackets ((b) for bound conformation, (u) for unbound conformation). Structural families are named according to the SCOPe database [60] at the family level. Averaged sequence identity and RMSD are given for each family.

S2 Table. AUC according to the grid resolution used for the energy maps. A linear model was constructed from the dataset constituted of all the intra-family ligand pairs (202 protein pairs). This model allows the estimation of the linear correlation between the three descriptors and the pairwise ADM distance. The model takes into account the individual contribution of each descriptor as well as their crossed contributions with each other. The p-value of each individual contribution calculated over the 202 pairs is estimated with a Fisher test and are given in the table line "all proteins". We then individually looked each family to see whether the contribution of the descriptors is dependent from the family. Inside each family, the number of protein pairs is too small to estimate a linear model. Consequently, we used a Spearman correlation coefficient test to estimate the p-value of each contribution.

S3 Table. Estimation of the effective contribution of sequence identity, RMSD and electrostatic distance in the pairwise ADM distances for each ligand pair belonging to a same family. The correlation is computed between each cell of the 74 energy maps of each of the 74 receptors and the corresponding cell in receptor's maps of stickiness, hydrophobicity and CV.

S4 Table. Correlation between energy scores and stickiness, hydrophobicity and circular variance (CV). The grid resolution corresponds to the number of cells composing the energy maps. The AUC is calculated following the same protocol used in the main text (see *Materials and Methods*)

S1 Fig. Interactions between structural families of the dataset. Interactions are symbolized by links between families. An interaction is established between two families when, there is at least one PDB reporting a structure of complex involving members of the two families [30]. Consequently, all members of a family do not necessarily have its native partner in its native-related partner family. The V set domains family is a special case of self-interacting family, where members form dimers of structural homologs.

S2 Fig. AUC values calculated on random subsets of receptor of different sizes. The AUC is computed following the protocol described in Fig. 1 with random subsets composed from 1 to 73 receptors. Receptors of each subset are randomly chosen among the 74 receptors of the dataset. For each subset size, the procedure is repeated 100 times. Red vertical lines indicate the standard deviation of the AUC for each subset size. Above a subset size of five receptors, the AUC does not significantly fluctuate (risk of wrongly rejecting the equality of two variances (F-test) >5% [61]).

S3 Fig. Receiver operating characteristic (ROC) curve and Area Under this Curve (AUC) calculated for each family.

S4 Fig. Papain-like family. (*A*) Energy map distances matrix. It corresponds to the subsection of the ADM for the papain-like family (for the construction of the ADM, see

Materials and Methods). Each entry *(i,j)* represents the pairwise energy map distance of the

ligand pair *(i,j)* averaged over the 74 receptors of the dataset (for more details, see *Materials*

and Methods). (*B*) Pairwise sequence identity matrix between all members of the family. (*C*)

Pairwise root mean square deviation (RMSD) matrix between all members of the family. (*D*)

Electrostatic maps and cartoon representations of the seven members of the family. An

electrostatic map represents the distribution of the electrostatic potential on the surface of a

protein (see Fig. S15 and *Materials and Methods*). Cartoon structures are colored according to

the distribution of their electrostatic potential. (*E*) Electrostatic map distances matrix. Each

entry *(i,j)* of the matrix represents the Manhattan distance between the electrostatic maps of

the proteins *(i,j)*.

S5 Fig. Eukaryotic-proteases family. (*A*) Energy map distances matrix. It corresponds to the

subsection of the ADM for the Eukaryotic proteases family (for the construction of the ADM,

see *Materials and Methods*). Each entry *(i,j)* represents the pairwise energy map distance of

the ligand pair *(i,j)* averaged over the 74 receptors of the dataset (for more details, see

Materials and Methods). (*B*) Pairwise sequence identity matrix between all members of the

family. (*C*) Pairwise root mean square deviation (RMSD) matrix between all members of the

family. (*D*) Electrostatic maps and cartoon representations of the seven members of the

family. An electrostatic map represents the distribution of the electrostatic potential on the

surface of a protein (for more details, see Fig. S15 and *Materials and Methods*). Cartoon

structures are colored according to the distribution of their electrostatic potential. (*E*)

Electrostatic map distances matrix. Each entry *(i,j)* of the matrix represents the Manhattan

distance between the electrostatic maps of the proteins *(i,j)*.

S6 Fig. V set domains family. (*A*) Energy map distances matrix. It corresponds to the

subsection of the ADM for the V set domain family (for the construction of the ADM, see

Materials and Methods). Each entry *(i,j)* represents the pairwise energy map distance of the

ligand pair *(i,j)* averaged over the 74 receptors of the dataset (for more details, see *Materials*

and Methods). (*B*) Pairwise sequence identity matrix between all members of the family. (*C*)

Pairwise root mean square deviation (RMSD) matrix between all members of the family. (*D*)

Electrostatic maps and cartoon representations of the six members of the family. An

electrostatic map represents the distribution of the electrostatic potential on the surface of a

protein (for more details, see Fig. S15 and *Materials and Methods*). Cartoon structures are

colored according to the distribution of their electrostatic potential. (*E*) Electrostatic map

distances matrix. Each entry *(i,j)* of the matrix represents the Manhattan distance between the

electrostatic maps of the proteins *(i,j)*.

S7 Fig. UCH-L family. (*A*) Energy map distances matrix. It corresponds to the subsection of

the ADM for the UCH-L family (for the construction of the ADM, see *Materials and*

Methods). Each entry *(i,j)* represents the pairwise energy map distance of the ligand pair *(i,j)*

averaged over the 74 receptors of the dataset (for more details, see *Materials and Methods*).

(*B*) Pairwise sequence identity matrix between all members of the family. (*C*) Pairwise root

mean square deviation (RMSD) matrix between all members of the family. (*D*) Electrostatic

maps and cartoon representations of the seven members of the family. An electrostatic map

represents the distribution of the electrostatic potential on the surface of a protein (for more

details, see Fig. S15 and *Materials and Methods*). Cartoon structures are colored according to

the distribution of their electrostatic potential. (*E*) Electrostatic map distances matrix. Each

entry *(i,j)* of the matrix represents the Manhattan distance between the electrostatic maps of

the proteins *(i,j)*.

S8 Fig. UCH family. (*A*) Energy map distances matrix. It corresponds to the subsection of the

ADM for the UCH family (for the construction of the ADM, see *Materials and Methods*).

Each entry *(i,j)* represents the pairwise energy map distance of the ligand pair *(i,j)* averaged

over the 74 receptors of the dataset (for more details, see *Materials and Methods*). (*B*)

Pairwise sequence identity matrix between all members of the family. (*C*) Pairwise root mean

square deviation (RMSD) matrix between all members of the family. (*D*) Electrostatic maps

and cartoon representations of the seven members of the family. An electrostatic map

represents the distribution of the electrostatic potential on the surface of a protein (for more

details, see Fig. S15 and *Materials and Methods*). Cartoon structures are colored according to

the distribution of their electrostatic potential. (*E*) Electrostatic map distances matrix. Each

entry *(i,j)* of the matrix represents the Manhattan distance between the electrostatic maps of

the proteins *(i,j)*.

S9 Fig. Ubiquitin activating enzymes family. (*A*) Energy map distances matrix. It

corresponds to the subsection of the ADM for the Ubiquitin activating enzymes family (for

the construction of the ADM, see *Materials and Methods*). Each entry *(i,j)* represents the

pairwise energy map distance of the ligand pair *(i,j)* averaged over the 74 receptors of the

dataset (for more details, see *Materials and Methods*). (*B*) Pairwise sequence identity matrix

between all members of the family. (*C*) Pairwise root mean square deviation (RMSD) matrix

between all members of the family. (*D*) Electrostatic maps and cartoon representations of the

seven members of the family. An electrostatic map represents the distribution of the

electrostatic potential on the surface of a protein (for more details, see Fig. S15 and *Materials*

and Methods). Cartoon structures are colored according to the distribution of their

electrostatic potential. (*E*) Electrostatic map distances matrix. Each entry *(i,j)* of the matrix

represents the Manhattan distance between the electrostatic maps of the proteins *(i,j)*.

S10 Fig. UBC-related family. (*A*) Energy map distances matrix. It corresponds to the

subsection of the ADM for the UBC-related family (for the construction of the ADM, see

Materials and Methods). Each entry *(i,j)* represents the pairwise energy map distance of the

ligand pair *(i,j)* averaged over the 74 receptors of the dataset (for more details, see *Materials*

and Methods). (*B*) Pairwise sequence identity matrix between all members of the family. (*C*)

Pairwise root mean square deviation (RMSD) matrix between all members of the family. (*D*)

Electrostatic maps and cartoon representations of the seven members of the family. An

electrostatic map represents the distribution of the electrostatic potential on the surface of a

protein (for more details, see Fig. S15 and *Materials and Methods*). Cartoon structures are

colored according to the distribution of their electrostatic potential. (*E*) Electrostatic map

distances matrix. Each entry *(i,j)* of the matrix represents the Manhattan distance between the

electrostatic maps of the proteins *(i,j)*.

S11 Fig. Kunitz (STI) inhibitors family. (*A*) Energy map distances matrix. It corresponds to

the subsection of the ADM for the Kunitz (STI) inhibitors family (for the construction of the

ADM, see *Materials and Methods*). Each entry *(i,j)* represents the pairwise energy map

distance of the ligand pair *(i,j)* averaged over the 74 receptors of the dataset (for more details,

see *Materials and Methods*). (*B*) Pairwise sequence identity matrix between all members of

the family. (*C*) Pairwise root mean square deviation (RMSD) matrix between all members of

the family. (*D*) Electrostatic maps and cartoon representations of the seven members of the

family. An electrostatic map represents the distribution of the electrostatic potential on the

surface of a protein (for more details, see Fig. S15 and *Materials and Methods*). Cartoon

structures are colored according to the distribution of their electrostatic potential. (*E*)

Electrostatic map distances matrix. Each entry *(i,j)* of the matrix represents the Manhattan

distance between the electrostatic maps of the proteins *(i,j)*.

S12 Fig. Retrovirus capsid proteins family. (*A*) Energy map distances matrix. It

corresponds to the subsection of the ADM for the retrovirus capsid proteins family (for the

construction of the ADM, see *Materials and Methods*). Each entry *(i,j)* represents the pairwise

energy map distance of the ligand pair *(i,j)* averaged over the 74 receptors of the dataset (for

more details, see *Materials and Methods*). (*B*) Pairwise sequence identity matrix between all

members of the family. (*C*) Pairwise root mean square deviation (RMSD) matrix between all

members of the family. (*D*) Electrostatic maps and cartoon representations of the seven

members of the family. An electrostatic map represents the distribution of the electrostatic

potential on the surface of a protein (for more details, see Fig. S15 and *Materials and*

Methods). Cartoon structures are colored according to the distribution of their electrostatic

potential. (*E*) Electrostatic map distances matrix. Each entry *(i,j)* of the matrix represents the

Manhattan distance between the electrostatic maps of the proteins *(i,j)*.

S13 Fig. Cystatins family. (*A*) Energy map distances matrix. It corresponds to the subsection

of the ADM for the cystatins family (for the construction of the ADM, see *Materials and*

Methods). Each entry *(i,j)* represents the pairwise energy map distance of the ligand pair *(i,j)*

averaged over the 74 receptors of the dataset (for more details, see *Materials and Methods*).

(*B*) Pairwise sequence identity matrix between all members of the family. (*C*) Pairwise root

mean square deviation (RMSD) matrix between all members of the family. (*D*) Electrostatic

maps and cartoon representations of the seven members of the family. An electrostatic map

represents the distribution of the electrostatic potential on the surface of a protein (for more

details, see Fig. S15 and *Materials and Methods*). Cartoon structures are colored according to

the distribution of their electrostatic potential. (*E*) Electrostatic map distances matrix. Each

entry *(i,j)* of the matrix represents the Manhattan distance between the electrostatic maps of

the proteins *(i,j)*.

S14 Fig. Cyclophilins family. (*A*) Energy map distances matrix. It corresponds to the

subsection of the ADM for the cyclophilins family (for the construction of the ADM, see

893 *Materials and Methods*). Each entry *(i,j)* represents the pairwise energy map distance of the

894 ligand pair *(i,j)* averaged over the 74 receptors of the dataset (for more details, see *Materials*

895 *and Methods*). (*B*) Pairwise sequence identity matrix between all members of the family. (*C*)

896 Pairwise root mean square deviation (RMSD) matrix between all members of the family. (*D*)

897 Electrostatic maps and cartoon representations of the seven members of the family. An

898 electrostatic map represents the distribution of the electrostatic potential on the surface of a

899 protein (for more details, see Fig. S15 and *Materials and Methods*). Cartoon structures are

900 colored according to the distribution of their electrostatic potential. (*E*) Electrostatic map

901 distances matrix. Each entry *(i,j)* of the matrix represents the Manhattan distance between the

902 electrostatic maps of the proteins *(i,j)*.

904 **S15 Fig. Generation of electrostatics, stickiness, hydrophobicity and circular variance**

905 **(CV) maps.** Here is presented an example of generation of the stickiness map for the structure

906 1AVW_A. (*A*) Generation of particles with a slightly modified Shrake-Rupley algorithm [59]

907 around the protein surface, leads to a homogenous shell of particles with a 1Å^2 density. Each

908 sphere is located at 5Å from the surface of the protein. The stickiness value of the closest

909 atom of the protein is attributed to each particle. In this example, spheres are colored

910 according to the stickiness of the protein surface. The procedure is similar for hydrophobicity

911 and CV. (*B*) The spherical coordinates of each sphere is represented on a 2-D map with an

912 equal-area sinusoidal projection, following the same protocol as described in Fig. 2 and

913 *Materials and Methods*. Each resulting dot is colored according to the same scale of (*A*). (*C*)

914 The map is smoothed following the protocol in Fig. 2 and *Materials and Methods*. The scale

915 is the same as in (*A*).

S16 Fig. Generation of stacked maps of a receptor. (*A*) Calculation of the 74 one-color

maps (red ones in the example) of receptor #1. A value of one is associated to colored cells

while zero is assigned to white cells. (*B*) Sum of the 74 one-color maps into a stacked map.

Cell's intensity varies from 0 to 74 and corresponds to the number of time the cell is colored

over the 74 ligands. (*C*) Filtering of the cells of low cell intensity (intensity < 17) and areas of

too small size (< 4 cells) with a Dirichlet process mixture model simulation for image

segmentation [57]. The procedure is repeated for each color stacked map.

S17 Fig. Blue and red stacked maps of 1P9D_U computed for each ligand family. (*A-L*)

We compute the one-color stacked map of each family as the sum of the one-color maps

resulting from the docking of each ligand of a same family with 1P9D_U.

S18 Fig. Boxplots of the size (in number of cells) of each energy class for all stacked

maps. One should notice that the sum of the sizes of the 5 energy classes is superior to 1548

cells, which is the total size of a map, because a same cell of a stacked map can be assigned to

several energy classes (Fig 8).

S19 Fig. Boxplots of the electrostatic potential of the protein surfaces depending on the

energy class. The electrostatic potential is calculated for each protein following the protocol

described in *Materials and Methods*. *p-values* between the variances of two "successive"

energy classes were computed using the F-test.

S20 Fig. Boxplots of energy map pairwise distances between ligand pairs of conformers

and pairs of homologous ligands (i.e. non-conformers pairs). For each receptor, we

computed (i) the average of energy map distances of pairs of conformers, (ii) the average of

energy map distances of pairs of homologous ligands. P-values are calculated with an

unilateral Wilcoxon test.

Fig 1.

Fig 2.

Fig 3.

Fig 4.

Fig 5.

Fig 6.

Fig 7.

Fig 8.