1

Confounding factors in algal phosphorus limitation experiments

Algal phosphorus limitation

Whitney S. Beck[1,2] *, Ed K. Hall[2, 3, 4]

[1]Department of Biology, Colorado State University, Fort Collins, Colorado, United States of
America

[2]Graduate Degree Program in Ecology, Colorado State University, Fort Collins, Colorado,
United States of America

[3]Department of Ecosystem Science and Sustainability, Colorado State University, Fort Collins,
Colorado, United States of America

[4]Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, Colorado,
United States of America

*Corresponding author

Email: whitney.beck@colostate.edu

Abstract

Assessing algal nutrient limitation is critical for understanding the interaction of primary production and nutrient cycling in streams, and nutrient diffusing substrate (NDS) experiments are often used to determine limiting nutrients such as nitrogen (N) and phosphorus (P). Unexpectedly, many experiments have also shown decreased algal biomass on NDS P treatments compared to controls. To address whether inhibition of algal growth results from direct P toxicity, NDS preparation artifacts, or environmental covariates, we first quantified the frequency of nutrient inhibition in published experiments. We also conducted a meta-analysis to determine whether heterotrophic microbial competition or selective grazing could explain decreases in algal biomass with P additions. We then deployed field experiments to determine whether P-inhibition of algal growth could be explained by P toxicity, differences in phosphate cation (K vs. Na), differences in phosphate form (monobasic vs. dibasic), or production of H_2O_2 during NDS preparation. We found significant inhibition of algal growth in 12.9% of published NDS P experiments as compared to 4.7% and 3.6% of N and NP experiments. The meta-analysis did not show enhanced heterotrophy on NDS P treatments or selective grazing of P-rich algae. Our field experiments did not show inhibition of autotrophic growth with P additions, but we found significantly lower gross primary productivity (GPP) and biomass-specific GPP of benthic algae on monobasic phosphate salts as compared to dibasic phosphate salts, likely because of reduced pH levels. Additionally, we note that past field experiments and meta-analyses support the plausibility of direct P toxicity or phosphate form (monobasic vs. dibasic) leading to inhibition of algal growth, particularly when other resources such as N or light are limiting. Given that multiple mechanisms may be acting simultaneously, we recommend practical, cost-

44   effective steps to minimize the potential for P-inhibition of algal growth as an artifact of NDS

45   experimental design.

Introduction

Benthic algal production provides an important energy source to higher trophic levels [1], and

in low productivity streams, growth of macroinvertebrate and fish grazers may be limited by the

availability of algal food resources [2]. Freshwater algal growth is often limited by the

availability of nitrogen (N), phosphorus (P), or both nutrients [3, 4], but human activities are

increasing N and P inputs to streams via sources such as wastewater treatment effluent,

agricultural runoff, and atmospheric N deposition [5]. These excess nutrients may result in

harmful levels of algal biomass that degrade ecological habitat [5], stream aesthetics [6], and

drinking water quality [5]. Identifying nutrients that limit algal productivity in individual stream

reaches can inform stream management plans that promote human and ecosystem health.

For over thirty years, nutrient diffusing substrate (NDS) experiments have been used to

determine nutrient limitation of benthic algal communities [7, 8]. Nutrient diffusing substrate

experiments are constructed by filling a small vessel (e.g., a plastic vial or clay pot) with agar

and a nutrient solute of choice (e.g., KH_2PO_4), and comparing growth with a paired control

vessel containing only agar [9]. Differences in algal responses can then be compared across the

NDS treatments. Contrary to the long-held paradigm that P frequently limits algal productivity in

freshwater ecosystems [10], one of the first published NDS experiments observed that treatments

with 0.5 M P had lower algal biomass than treatments with 0.05 M P [7]. Many studies have

since found that algal biomass can be inhibited by P as compared to controls (e.g., [11, 12]), and

it is unclear how often or why this phenomenon occurs. Whereas P may not always enhance

growth due to limitation by N or other resources (light, Fe, etc.), it is surprising that increasing

levels of P in NDS can result in decreased algal biomass relative to the control treatment. These

results suggest that artifacts associated with NDS experiments may be leading to the

underreporting or misrepresentation of P-limitation in freshwater ecosystems.

Several hypotheses have been introduced to explain why addition of P in NDS would result in

a decrease of algal biomass. As a macronutrient, P is required for algal growth and maintenance

[13], but high P concentrations may result in direct physiological toxicity and this has been

hypothesized as a reason for observed P-inhibition [7]. While mechanisms for this toxicity in

algae are unclear, excessive P concentrations in growth media of terrestrial plants can negatively

affect the availability, uptake, and metabolic processing [14] of Fe [15], K [15], and Z [16],

leading to deficiency of these essential nutrients and slowing or inhibiting plant growth. Other

hypothesized mechanisms for P-inhibition of algal growth have focused on artifacts related to

preparation of the NDS, including: 1) phosphate cation type (K vs. Na) and toxicity, 2)

phosphate form (monobasic vs. dibasic), and 3) H_2O_2 production from phosphate reacting with

agar during autoclaving. A limited number of laboratory studies have tested whether high

concentrations of the phosphate salt cations (K^+, Na^+, H^+) may inhibit algal growth. These

studies have shown that K phosphates and KCl are toxic to algae at lower concentrations than Na

phosphates and NaCl [17, 18]. The phosphate form may also influence algae, as monobasic

forms (KH_2PO_4 and NaH_2PO_4) tend to have lower NDS effect sizes than dibasic forms (K_2HPO_4

and Na_2HPO_4 [19]). This suggests algae are either inhibited by acidic pH levels (induced by

monobasic forms) or are experiencing cation limitation (alleviated by dibasic forms). Lower pH

may influence algae directly by changing concentrations of H^+ around the cell or indirectly

through the effects of pH on metal toxicity or nutrient availability (e.g., via slowed nitrification

rates or binding of P by Al [20]). Finally, the preparation of the NDS media may affect how the

P treatments influence algal growth. Autoclaving phosphate and agar together produces H_2O_2, a

toxin that may inhibit microbial growth [21]. It has been suggested that the common NDS

construction method of combining the two compounds on a hotplate could also produce the same

result [9] thus leading to inhibition of algal growth in treatments that contain P; however, to our

knowledge this has never been directly tested.

Beyond these direct artifacts of NDS preparation, there may also be a series of indirect effects

of adding phosphate to NDS that could inhibit algal growth. For example, P could

disproportionately stimulate heterotrophic microbes [22] and increase competition between

autotrophs and heterotrophs for other limiting nutrients, ultimately suppressing algal growth. It is

also possible that P amendments may induce additional top-down pressure if insect grazers

selectively graze P-rich algal biofilms [23], resulting in lower algal biomass on NDS P

treatments as compared to controls [11]. Selective foraging has been supported by a theoretical

analysis [24], and laboratory experiments provide some additional support that grazers can

engage in P-specific foraging [23, 25].

Determining why P-inhibition of algal growth has been observed in NDS experiments is not

only an important methodological question but one with important implications for an ecological

understanding of lotic ecosystems. Each of the previously-described mechanisms may potentially

affect the response of algae to P treatments in NDS experiments, but there is no single study that

simultaneously evaluates each mechanism. We used quantitative analyses of published data and

our own field experiments to investigate how frequently and why NDS P treatments inhibit algal

growth. First, we surveyed the literature to determine how frequently significant inhibition of

algal growth was reported for P, N, and NP treatments. We also completed a meta-analysis to

determine if there was consistent evidence for heterotrophic microbial competition or top-down

grazing control that could be leading to P-inhibition of algal growth across multiple study

systems. Finally, we completed field experiments to directly address the effects of NDS

preparation on several common response variables used to evaluate algal growth: chlorophyll *a*

(a measure of algal biomass), ash-free dry mass (AFDM, a measure of total biofilm organic

matter), a calculated index of autotrophy (AI), gross primary productivity (GPP), and biomass-

specific GPP (GPP/chlorophyll *a*).

Methods

Quantitative Review

To explicitly quantify how often nutrient treatments inhibit algal growth in NDS experiments,

we used the database assembled by Beck et al. [19]. Briefly, this database includes 649 NDS

experiments from 1985-2015 that used algal biomass (chlorophyll *a*) as a response variable. The

database was previously used to determine overall effect sizes of P, N , and NP additions and to

quantify the influence of over thirty experimental, environmental, and geographic covariates.

However, in this study our goal was to determine how many individual experiments detected

significantly lower algal biomass levels on P treatments (n=534), N treatments (n=553), and NP

treatments (n=591) as compared to controls (α=0.05). Although we could have used a meta-

analysis approach, we used separate two-tailed t-tests for each NDS experiment to better

represented how investigators analyzed data in individual studies.

To determine whether P treatments significantly influenced the proportion of autotrophy in

microbial communities and whether grazers selected for algal biomass on NDS P treatments, we

used meta-analysis models (see below). To assess whether P additions changed algal-

heterotrophic interactions, we identified any experiments from the previously compiled database

[19] that also reported AFDM as a response variable, as a proxy for total biofilm organic matter.

For this study, we extracted AFDM data using Webplot Digitizer version 3.12 [26], and

calculated an autotrophic index (AI) for each experiment and treatment as follows:

$$\text{AI} = \frac{\text{AFDM}}{\text{chlorophyll } a}$$ (1)

We interpret lower values of the AI as a higher proportion of autotrophy in the microbial

community [22]. For NDS P treatments we calculated AI log response ratios (LRRs, a measure

of effect size) as:

$$\text{LRR} = \ln \frac{Y_1}{Y_2} = \ln(Y_1) - \ln(Y_2)$$ (2)

Where Y_1 is the mean AI from the P treatments and Y_2 is the mean AI from the control

treatments in a given experiment [27]. LRRs greater than zero indicate there is a positive P

treatment effect, but LRRs less than zero indicate an inhibitory effect of the P treatment as

compared to controls. We also calculated the variance of effect sizes using:

$$\text{LRR\_Var} = \frac{s_1^2}{n_1 Y_1^2} + \frac{s_2^2}{n_2 Y_2^2}$$ (3)

Where s_1 and s_2 are the standard deviations of the P and control treatments, and n_1 and n_2 are the

number of P and control replicates [27]. To determine whether P additions influence grazer

selection of algal biofilms, we identified NDS experiments that also incorporated grazer

exclusion treatments. We calculated LRR and LRR_Var metrics for P treatments compared to

controls in grazed and ungrazed plots (Equations 2-3).

We used the metafor package in R to build meta-analysis models based on experimental effect

sizes and variances [28]. Meta-analysis models account for variability within and among

experiments by weighting effect sizes by their variances [28]. In all models, we used "site" as a

random effect, to account for the correlated effects of experiments deployed in the same stream

reach [29]. Models were used to determine how P additions influenced AI effect size, and to

determine how grazers influenced P treatment effects on algal biomass. For the grazer models,

we included grazer treatment as a covariate in the meta-analysis models [28].

Experiments

To investigate how NDS preparation methods influence P-inhibition of algal growth, we

prepared and deployed a series of complementary NDS experiments in a sub-alpine stream. To

address the influence of cation type, monobasic vs. dibasic phosphate form, and the potential for

H_2O_2 formation, the first two experiments involved crossing four different phosphate chemicals

(KH_2PO_4, K_2HPO_4, NaH_2PO_4, or Na_2HPO_4 all at 0.1 M concentrations) with two laboratory

heating methods (boiling agar and phosphate together vs. separately) for a total of eight

preparation treatments (Fig 1). In a third experiment, we again crossed the four phosphate

compounds with two laboratory heating methods but to assess direct toxicity of excess P, we also

used two different concentrations of P (0.05 M and 0.5 M) for a total of 16 preparation

treatments. Concentrations were chosen based on the most commonly used NDS concentrations.

Furthermore, Fairchild et al. [7] previously observed a difference in algal biomass responses to

0.05 M and 0.5 M of P.

We constructed NDS according to standard methods in the literature [9]. Briefly, we boiled 2%

agar with deionized water, poured the solutions into 30 mL vials (Item #66159, U.S. Plastic

Corps, Lima OH), and capped the cooled agar with fritted glass discs (5.7 cm^2, Item #C4505, EA

Consumables, Pennsauken, NJ). A plain agar solution was used for the control, and the specified

phosphate salt was added to the agar according to the experimental design described above (Fig

1). For the "heated together" treatment, we boiled the agar and the phosphate salt together. For

the "heated separately" treatment, we added the phosphate salt once the agar had cooled to

handling temperature (55-65° C) and mixed it thoroughly with a magnetic stir plate before

solidification.

While preparing the NDS treatments we also measured pH, a putative mechanism for how

phosphate form mediates P-inhibition of algal growth. We tested the effects of phosphate

compound and heat treatment on agar and water pH in the lab. To do this we used litmus pH test

strips to measure the pH of cooling agar for four vials from each phosphate treatment and the

control. To measure effects of phosphate compound and heat treatment on water pH, we

constructed two replicate NDS representing the four phosphate compounds crossed with two heat

treatments, as well as a control. These NDS were placed in separate plastic Ziploc® bags with 1

L of distilled water. We used a multimeter with a pH probe (Thermo Fisher Scientific Orion

Star™ A329, Waltham, MA) to measure the initial pH and remeasured the pH after 24 hours.

We deployed all NDS experiments at Little Beaver Creek, a low-order, open-canopy stream in

the mountains of the Roosevelt National Forest in Colorado (40.625° N, -105.527° W).

Experiments 1 and 2 were deployed in summer 2016, while experiment 3 was deployed in

summer 2017 (S1 Table). Previous NDS experiments in summer 2015 showed that Little Beaver

was co-limited by N and P, but primarily N-limited, with P treatments causing inhibition of algal

biomass relative to the control (Beck, unpublished). We randomized 6 replicates of each

treatment and attached 6-8 individual NDS vials to plastic L-shaped bars (Item #45031, U.S.

Plastic Corp, Lima, OH) that were anchored into the streambed using metal stakes. Upon

deployment, collection, or both, we measured in-stream pH, conductivity, and temperature using

a multimeter and probes (Thermo Fisher Scientific Orion Star™ A329, Waltham, MA); collected

duplicate filtered (0.45 µm Type A/E filters, Pall Corporation, Port Washington, NY) water

samples in 60 mL nalgene™ bottles for nutrient analysis; measured canopy cover using a

densiometer (Forestry Suppliers, Jackson, MS); and measured flow using a Marsh McBirney

meter (Hach, Loveland, CO). We used a MiniWater® 20 meter (Schiltknecht, Switzerland) to

measure 2.5 cm-scale current velocity above three evenly-spaced vials on each L-bar. We

measured NO_3^- using the Cd reduction method [30] and orthophosphate using the ascorbic acid

method [31] on an ALPKEM® Flow Solution IV autoanalyzer (O.I. Analytical, College Station,

TX).

We analyzed primary production rates on NDS discs after each in-stream experiment [9].

Briefly, upon collecting the discs, we immediately placed them in 50 mL centrifuge tubes with

unfiltered river water, capping the tubes underwater to exclude any air bubbles. Tubes were

stored on ice during transport back to the laboratory (less than two hours), where we used filtered

(0.45 µm Type A/E filters, Pall Corporation, Port Washington, NY) river water to run light and

dark incubations with the NDS discs. We measured the initial temperature and dissolved oxygen

(DO) values of the water before light and dark incubations using a ProDO meter (YSI, Yellow

Springs, OH). We also included four "blank" tubes with water only, to control for background

changes in DO because of temperature changes or exposure to atmospheric oxygen during the

measurement process. During light treatments, tubes were incubated in sunlight for two hours in

the afternoon, then we measured the ending DO concentration in each tube and ending

temperature in several representative tubes. For dark incubations, we replaced the filtered water

and incubated tubes in the dark for two hours, after which we measured the final DO

concentrations in each tube and ending temperature in several representative tubes.

We calculated net primary productivity (NPP) as the increase in oxygen during the light

incubations, correcting for the change in the blank stream water tubes. We calculated respiration

as the decrease in oxygen during the dark incubations, correcting for the change in the blank

stream water tubes. Gross primary productivity was calculated as follows, then all variables were

standardized by disc area and time:

$\text{GPP} = \text{NPP} + |\text{Respiration}|$ (4)

After the incubations, we immediately placed the discs in black film canisters and extracted

chlorophyll *a* for 12-24 hours using buffered 90% ethanol. We measured chlorophyll *a* with an

acidification correction [32] using an Aquafluor® fluorometer (Turner Designs, San Jose, CA).

We calculated biomass-specific GPP (GPP/ chlorophyll *a*) as an additional response metric for

each disc.

For experiments 1 and 2, we saved all liquid slurry from the chlorophyll *a* extractions and

filtered the liquid through a pre-combusted filter (500° C for one hour, 0.45 µm Type A/E, Pall

Corporation, Port Washington, NY). We then used both the filter and glass disc associated with

each NDS to measure AFDM [22]. For experiment 3, we allowed the chlorophyll *a* extraction

slurry to evaporate in a weigh boat under a fume hood and used all remaining material and the

glass disc to measure AFDM. We dried the samples for 48 hours at 50° C, pre-weighed their

masses, and combusted them at 500° C for one hour in a muffle furnace. We then rehydrated the

discs with deionized water, dried them for another 48 hours at 50° C, and weighed the final

masses. This procedure was to account for any water that might have been lost from clay

particles in the muffle furnace. The difference in weights was calculated as AFDM. We also used

the chlorophyll *a* and AFDM measurements to calculate AI as described in Equation 1.

Statistical Analyses for Experiments

We completed all statistical analyses in R version 3.5.0 (R Core Team 2018). To test the effect

of treatment classes on the pH of the NDS agar and water incubated with the NDS, we used two-

way ANOVAs with phosphate form (monobasic vs. dibasic) and heating method as factors. We

compared means using post-hoc Tukey's HSD tests ($\alpha=0.05$).

Because the field experiments comprised an incomplete factorial design (i.e., crossed

treatments and a separate control, Fig 1), we used three separate approaches to analyze the

results. First, to determine whether P preparation treatments significantly stimulated or inhibited

response variables relative to controls, we used one-way ANOVAs with post-hoc Dunnett tests

to compare each of the eight P preparation treatments (Fig 1) with the controls. The Dunnett test

is a multiple comparison procedure that compares individual treatment means to control means

while maintaining a family-wise error rate that is below α ($\alpha = 0.05$ in this study). We used only

data from experiments 1 and 2 and included experiment as a fixed effect block to control for

experiment-specific artifacts. Second, to determine whether the treatment classes (cation type,

phosphate form, and heat, Fig 1) influenced NDS P effect sizes, we used data from experiments

1 and 2 to run separate one-way ANOVAs for each treatment class and response variable (algal

biomass, AFDM, AI, GPP, and biomass-specific GPP). We included experiment as a fixed effect

block, and the model response variables were P treatment values of each response variable

divided by their respective experimental control means [9]. Third, to determine whether

treatment classes interacted with P concentration to influence NDS P effect sizes, we used data

from experiment 3 to run separate two-way ANOVAs crossing each treatment class with

concentration.

Results

Quantitative Review

In our analysis of 649 of experiments from the literature we found that algal biomass was more

commonly inhibited by NDS P treatments than either N or NP treatments. Phosphorus additions

produced a significant negative effect in 12.9% of experiments. However, N and NP additions

produced a significant negative effect in only 4.7% and 3.6% of experiments, respectively (both

within the commonly assumed type I error rate of 5%).

We next looked for published evidence of hypothesized biological mechanisms that would

explain inhibition of algal growth by P. To address the potential for heterotrophic suppression of

autotrophs in microbial communities on NDS P treatments, we identified 45 experiments from

11 studies where an AI effect size could be calculated (S1 References). However, the meta-

analysis of AI effect sizes showed neither a positive nor negative response to P treatments (Fig

2). We found even fewer examples of studies that could be used to examine nutrient additions in

conjunction with grazer exclusions. Only five experiments from three studies reported the effects

of grazer exclusions on P treatments in NDS experiments (S2 References). In these studies, algal

biomass effect sizes on the P treatments were not significantly influenced by the presence or

absence of grazers (Fig 2, $Q_M = 2.594$, p=0.107).

Experiments

To address the remaining hypothesized mechanisms for inhibition of algal growth by P

amendments we conducted a series of field experiments in a small sub-alpine stream. We found

no significant differences between any of the eight P preparation treatments (Fig 1) and control

treatments (all p >0.05) for any of the algal response variable raw values, indicating neither

inhibition nor stimulation of algal growth was induced by the addition of P relative to the control

in our experiments.

When we grouped the eight P preparation treatments by three different treatment classes

(cation type, phosphate form, and heating method, Fig 1) and tested treatment class effects on

response variable effect sizes (S2 Table), we found a significant effect of phosphate form

(monobasic or dibasic) on GPP ($F_{1,88}$=5.057, p=0.027) and biomass-specific GPP ($F_{1,87}$=5.578,

p=0.020). Dibasic treatments produced higher rates for both measures of primary production

(Figs 3-4). Furthermore, phosphate form significantly altered the pH of the agar ($F_{1,28}$=1408.333,

p<0.001) and the water ($F_{1,12}$=503.244, p<0.001) in the laboratory experiments. Monobasic

chemicals significantly lowered pH means (agar pH=4.81, water pH=5.45) and dibasic chemicals

significantly raised pH means (agar pH=8.88, water pH=8.28) relative to controls (agar pH=7.25,

water pH=6.35). We did not find an effect of the phosphate cation salt or heating treatment on

any algal response variable effect sizes (p>0.05) in the field experiments (Figs 3-7, S2 Table).

We also did not find any significant main effects of phosphate concentration or interactions

between concentration and other treatment classes on response variable effect sizes in the field

experiments (S3 Table).

Discussion

Our experiments and quantitative literature analyses did not identify a clear mechanism to

explain why 12.9% of past NDS experiments reported a significant negative effect of P

treatments on algal growth, a number more than twice as high as the type I error rate (5%) and

higher compared to what was observed on N and NP treatments (both <5%). In our experiments

we observed slight but non-significant P stimulation rather than inhibition, consistent with the

evidence for P-limitation in many freshwater streams [4]. However, past laboratory experiments

[17, 18] and a previous meta-analysis [19] support the plausibility of direct P toxicity, cation

toxicity, or phosphate form as mechanisms that may inhibit algal growth. Given that multiple

mechanisms may be acting simultaneously, and some mechanisms are most likely ecosystem

dependent, we discuss the implications of each of our findings and recommend a series of

practical steps for future NDS experiments to reduce the likelihood of P-inhibition of algal

growth resulting from artifacts of experimental design or experimental preparation.

P toxicity

While our experiments did not show evidence of P toxicity as a mechanism for P-inhibition of

algal growth, previous research supports the possibility of P toxicity occurring across a range of

stream ecosystems. In contrast with previous research [7], in our experiments we saw no

significant difference in algal biomass between the low (0.05 M) P concentrations and high (0.5

M) P concentrations. Phosphorus toxicity is dependent upon biological and environmental

context (see below), and the P concentrations we used in this experiment may have been too low

to induce toxicity. There is evidence that NDS experiments in other systems may commonly

exceed the concentrations required to detect toxicity, as a previous meta-analysis of 534 NDS

experiments showed that higher P concentrations in NDS treatments significantly decreased P

effect sizes [19]. The physiological mechanisms that cause P toxicity in algae are not well

defined, but some insight can be gained from the terrestrial plant literature.

Terrestrial plant studies have demonstrated that excess P within a cell can induce Fe [15], K

[15], or Zn [16] deficiencies. This leads to leaf necrosis and discoloration [33], reduced growth

rates [15], and plant death [34, 35]. These plant symptoms have occurred even when studies

maintained optimal levels of other nutrients and pH [15], and when multiple phosphate forms

have been tested in the same study [36]. Algae [37], like terrestrial plants [38], can take up

excess P for storage (i.e. "luxury P uptake"), a strategy to deal with heterogenous nutrient

supplies common in stream ecosystems. The mechanisms and potential consequences of luxury P

consumption for algae are less clear, but it is plausible that P could accumulate to toxic levels

within cells and induce other elemental deficiencies as has been shown for terrestrial plants.

However, toxicity from luxury P consumption would only occur if NDS nutrients diffuse at high

enough rates that excess P could accumulate in the water-cell boundary layer or biofilm.

Previous studies have measured NDS nutrient diffusion rates in beakers of distilled water to

estimate stream water diffusion rates. For instance, a study showed that plastic vial NDS (5.1

cm^2 area) constructed with 50 mM KH_2PO_4 can release 0.321 mM PO_4^{3-}-P·hr^{-1} at day 0, but that

diffusion rate declines to 0.001 mM PO_4^{3-}-P·hr^{-1} by day 14 [39]. Clay pot NDS (86.8 cm^2 area)

constructed with 50 mM KH_2PO_4 can release 0.113 mM PO_4^{3-}-P ·hr^{-1} P at day 0, with a diffusion

rate that declines to 0.011 mM PO_4^{3-}-P ·hr^{-1} by day 14 [39]. Because diffusion rates decline in a

log-linear fashion over time, algal populations may initially experience concentrations of P from

NDS that are sufficiently high to induce toxicity and inhibition of growth (e.g., 0.019 mM PO_4^{3-}

P [18]). To optimize NDS experiments, future studies could complete pilot experiments that

empirically measure diffusion rates to determine appropriate P starting concentrations and

experimental lengths (described in detail by Costello et al. [40]). Ultimately, studies should be

long enough to surpass the initial high pulses of P released from NDS when direct toxicity may

occur, but short enough to maintain a measurable nutrient flux that is significantly enriched from

that of the control treatment.

There is also evidence that the availability of other resources can influence P toxicity.

Nitrogen- or light-limitation could potentially induce toxicity at lower concentrations of P, if

luxury P [37] accumulates without being used for growth due to limitation by other resources.

Terrestrial plant studies have shown a positive relationship between P toxicity concentrations and

N:P and K:P resource ratios [41], likely because high growth rates supported by N and K

availability can reduce tissue P concentrations [36]. Furthermore, P-inhibition of algal growth

tends to be stronger in shaded areas [19] where light may be limiting and where NO_3^- reduction

could be limited by the availability of NADPH from photosynthesis [42]. Thus, thresholds for P

toxicity appear to be closely linked with the probability of secondary limitation by another

resource, which would in part explain the inconsistency of reported P-inhibition in NDS studies

as environmental conditions change among different ecosystems.

Cation and Phosphate Form

Few NDS studies have controlled for the effects of phosphate cation or phosphate form

(monobasic vs. dibasic). However, experimental evidence suggests that toxicity thresholds differ

based on the cation in the phosphate salt [18]. In general, it appears K leads to toxicity at lower

phosphate salt concentrations than Na. Growth of the freshwater chrysophyte *Dinobryon sociale*

was maintained in the laboratory at P concentrations of 0.032 mM PO_4^{3-}-P for NaH_2PO_4 but

declined when P concentrations were raised from 0.005 mM PO_4^{3-}-P to 0.019 mM PO_4^{3-}-P for

KH$_2$PO$_4$ [18]. Similarly, a laboratory study on a cyanobacteria, *Microcystis spp.*, showed lower

toxicity thresholds for KCl as compared to NaCl [43], further supporting the potential for cation

toxicity with the same cation but for different salts. However, across hundreds of published field

studies, P effect sizes for algal biomass were higher for K phosphates as compared to Na

phosphates [19]. Taken together these results suggest that K-toxicity of algal biomass can be

induced under laboratory conditions, but K concentrations in NDS phosphate salts may not be

high enough to induce toxicity in field experiments.

Although the cation in the phosphate salt does not appear to strongly influence the effect of

NDS P treatments on algal growth, it is likely that the phosphate form (monobasic vs. dibasic)

used influences experimental outcomes by modifying pH at the surface of the NDS. In our

experimental stream, the pH varied between 7.79 and 8.10, which differed substantially from the

agar amended with monobasic phosphates (pH 4.81 $\pm$ 0.09) but was more comparable to the agar

amended with dibasic phosphates (pH 8.88 $\pm$ 0.06). These differences in pH between the P

treatments may have contributed to the difference in GPP and biomass-specific GPP we saw

between these treatments, i.e., increased productivity on dibasic treatments as compared to the

monobasic treatments. These experimental results are consistent with a previous meta-analysis

which showed P and NP effect sizes for algal biomass were significantly higher on dibasic NDS

treatments relative to those effects on monobasic treatments [19]. Our meta-analysis in this study

showed that P-inhibition was reported more commonly than N-inhibition of algal growth, which

could be driven by P compounds containing easily disassociated H$^+$ ions while N compounds

often do not. To avoid the artifact of pH on NDS P treatments we recommend that NDS

experiments mix monobasic and dibasic phosphates to reflect the background pH of study

streams as best as possible. This simple step would avoid the confounding influence of alteration

of pH in the P treatment of NDS experiments.

Heating Method

We also investigated the potential for H_2O_2 production during the preparation of NDS P

treatments to create an artifact in NDS experimental results, as has been hypothesized [44]. We

found that heating phosphate and agar together vs. separately on a hotplate did not produce

significant differences in algal response metrics. We did not directly test whether H_2O_2 was

produced in our experiments as was found by a laboratory study that autoclaved phosphate and

agar together [21], but either H_2O_2 production requires the combination of heat and pressure

(from autoclaves) and was not produced in this experiment, or the concentrations of H_2O_2 in our

experiments were not high enough to inhibit algal growth. A simple solution to avoid the

potential inhibiting effect of H_2O_2 would be to heat P and agar separately (described by Tank et

al. [9]) to avoid the possibility of H_2O_2 interference with algal growth. This approach would

require little extra effort in NDS experiment preparation and completely remove this potentially

confounding artifact.

Microbial Competition

Indirect mechanisms have also been proposed to explain why P additions commonly inhibit

algal growth. Heterotrophic and autotrophic microbial communities interact in complex ways

that may change along nutrient gradients. At low concentrations of P, heterotrophic microbes are

expected to be competitively dominant because of their strong affinity for P and high surface

area relative to volume [45]. However, heterotrophs may also regenerate nutrients that can

stimulate autotrophic production and autotrophs produce organic C that fuels heterotrophy,

leading to a coupling of the two communities [46, 47]. When nutrients are added to the system

these community dynamics are altered, affecting the biomass and diversity of both heterotrophs

and autotrophs within the biofilm [48]. A study of Idaho streams found a strongly stimulated AI

(i.e., a higher proportion of heterotrophs) when C was added, and a weakly stimulated AI when P

was added [22]. Furthermore, a study of Texas streams showed a decoupling of autotrophic and

heterotrophic production when nutrients were added [46]. Our experiments and meta-analysis

produced no evidence that heterotrophic-autotrophic interactions were influenced by P additions

(i.e., no change in AI), but future studies should consider alternative response metrics for

measuring heterotrophic microbial biomass if heterotrophic estimates are required to answer

study-specific research questions.

Grazer Selection

In addition to heterotrophic microbes affecting algal growth through ecological interactions,

we hypothesized that P effect sizes on grazer exclusion treatments may be larger than on grazed

treatments because some grazers have been shown to selectively consume P-enriched resources

[25]. Our meta-analysis did not produce evidence to suggest that P effect sizes differed between

grazed and ungrazed plots. However, because of the difficulty of conducting grazer exclusion

NDS studies, few studies to date have investigated resource selectivity in grazers under field

conditions. We recommend that future experiments consider the interactions between nutrient-

specific foraging and algal nutrient limitation to determine whether grazing leads to apparent P-

inhibition of algal growth. One option is to construct electrical exclusion treatments which can

prevent grazing from macroinvertebrates and vertebrates across a wide range of body sizes [49–

51].

Additional Considerations

There are several potential reasons why our experiments did not show significant P treatment effects that have previously been described in the literature. First, deploying only six replicates per treatment produced low statistical power, given the large number of treatments and the small effect sizes of P additions in our NDS experiment. Furthermore, phosphate and cation inhibition of algae have been demonstrated using controlled laboratory experiments that can achieve a wider and more precise concentration gradient as compared to field experiments. We previously observed P-inhibition of algal biomass at the same stream reach used in this study. However, NDS diffusion rates [52] and environmental characteristics in the field are clearly variable over space and time, which may have obscured our ability to connect field-scale results with proposed mechanisms that are largely based on laboratory studies and meta-analyses of field studies.

Conclusion

Phosphorus additions have significantly inhibited algal biomass in 12.9% of past NDS studies, and investigators have hypothesized that this may be an artifact of NDS preparation or an indirect effect of increased heterotrophic microbial competition or top-down grazer control. We found that phosphate form (monobasic vs. dibasic) likely influences algal growth on NDS P treatments by mediating biofilm pH levels, and acidic monobasic treatments may inhibit algal growth. Furthermore, the literature supports direct P toxicity as a mechanism for P-inhibition of algal growth [17–19], particularly when other resources such as light or N are limiting [37]. We did not find support for phosphate salt cation toxicity occurring under field conditions, nor did we find evidence that laboratory heating method influenced algal responses to P. Based on our

analyses, it is also unlikely that P stimulates heterotrophic microbes relative to autotrophic

microbes or that P stimulates grazing rates.

Considering that multiple mechanisms may be operating simultaneously to inhibit algal growth

on NDS P treatments, we recommend several low effort, cost-effective steps for the NDS

preparation process that could reduce the potential for P-inhibition in future experiments. First,

future experiments could measure background stream nutrient concentrations to determine the

most appropriate P treatment concentrations for NDS construction [53], with the goal of avoiding

the potential for levels of P that are directly toxic to algae. In addition, measuring NDS diffusion

rates [39, 40, 54] under conditions that mimic natural systems would allow investigators to

further determine appropriate concentrations and experimental lengths for NDS studies.

Experiments should be long enough to surpass the potential for initial P toxicity when diffusion

concentrations are at their highest but short enough to maintain stimulatory P diffusion from

NDS. Because monobasic and dibasic phosphates influence NDS pH levels, we encourage

investigators to mix the two phosphate forms to reflect the background pH in experimental

streams to the extent possible. Finally, while we did not find evidence that laboratory

construction methods inhibited algal growth, it seems prudent (and logistically simple) to use the

separate agar and phosphate heating methods outlined by Tank et al. [9] to avoid the potential for

H_2O_2 production that inhibits microbial growth [21]. Avoiding confounding factors in NDS

experiments will ensure that studies are not underestimating P-limitation of primary producers in

aquatic ecosystems, improving our understanding of how resources and environmental

conditions interact to affect algal growth and stream ecosystems as a whole.

Acknowledgements

This manuscript benefited from reviews by Keeley MacNeill and Steven Francoeur. We also

thank Mitchell Ralson for help in the field and laboratory.

References

1. Lamberti GA. The role of periphyton in benthic food webs. In: Stevenson RJ, Bothwell ML,

Lowe RL, editors. Algal ecology. San Diego: Academic Press; 1996. pp. 533-573.

2. Lewis WM, McCutchan JH. 2010. Ecological responses to nutrients in streams and rivers of

the Colorado mountains and foothills. Freshw Biol. 2010;55: 1973-1983.

3. Francoeur SN. Meta-analysis of lotic nutrient amendment experiments: detecting and

quantifying subtle responses. J North Am Benthol Soc. 2001;20: 358-368.

4. Elser JJ, Bracken MES, Cleland EE, Gruner DS, Harpole WS, Hillebrand H, et al. Global

analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and

terrestrial ecosystems. Ecol Lett. 2007;10: 1135-1142.

5. Carpenter SR, Caraco NF, Correll DL, Howarth RW, Sharpley AN, Smith VH. Nonpoint

pollution of surface waters with phosphorus and nitrogen. Ecol Appl. 1998;8: 559–568.

6. Suplee MW, Watson V, Teply M, McKee H. How green is too green? Public opinion of what

constitutes undesirable algae levels in streams. J Am Water Resour Assoc. 2009;45: 123–140.

7. Fairchild GW, Lowe RL, Richardson WB. Algal periphyton growth on nutrient-diffusing

substrates: an in situ bioassay. Ecology. 1985;66: 465-472.

8. Pringle CM. Effects of water and substratum nutrient supplies on lotic periphyton growth: an

integrated bioassay. Can J Fish Aquat Sci. 1987;44: 619-629.

9. Tank JL, Reisinger AJ, Rosi EJ. Nutrient limitation and uptake. In: Hauer FR, Lamberti GA,

editors. Methods in stream ecology 3rd edition. Amsterdam: Elsevier; 2017. pp. 141-171.

10. Schindler DW. Evolution of phosphorus limitation in lakes. Science 1977;195: 260–262.

11. Bernhardt ES, Likens GE. Controls on periphyton biomass in heterotrophic streams. Freshw

Biol. 2004;49: 14–27.

12. Sanderson BL, Coe HJ, Tran CD, Macneale KH, Harstad DL, Goodwin AB. Nutrient

limitation of periphyton in Idaho streams: results from nutrient diffusing substrate experiments. J

North Am Benthol Soc. 2009;28: 832-845.

13. Borchardt MA. Nutrients. In: Stevenson RJ, Bothwell ML, Lowe RL, editors. Algal ecology.

San Diego: Academic Press; 1996. pp. 183-227.

14. Jones Jr JB. Phosphorus toxicity in tomato plants: when and how does it occur? Commun

Soil Sci Plant Anal. 1998;29: 1779-1784.

15. Christie EK, Moorby J. Physiological responses of semiarid grasses. I. The influence of

phosphorus supply on growth and phosphorus absorption. Aust J Agric Res. 1975;26: 423-436.

16. Loneragan JF, Grunes DL, Welch RM, Aduayi EA, Tengah A, Lazar A, Cary EE.

Phosphorus accumulation and toxicity in leaves in relation to zinc supply. Soil Sci Soc Am J.

1982;46: 345-352.

17. Chu SP. The influence of the mineral composition of the medium on the growth of

planktonic algae: part II. The influence of the concentration of inorganic nitrogen and phosphate

phosphorus. J Ecol. 1943;31: 109-148.

18. Lehman JT. Ecological and nutritional studies on *Dinobryon Ehrenb.*: Seasonal periodicity

and the phosphate toxicity problem. Limnol Oceanogr. 1976;21: 646-658.

19. Beck WS, Rugenski AT, Poff NL. Influence of experimental, environmental, and geographic

factors on nutrient-diffusing substrate experiments in running waters. Freshw Biol. 2017;62:

1667–1680.

20. Planas D. Acidification effects. In: Stevenson RJ, Bothwell ML, Lowe RL, editors. Algal

ecology. San Diego: Academic Press; 1996. pp. 497-530.

21. Tanaka T, Kawasaki K, Daimon S, Kitagawa W, Yamamoto K, Tamaki H, et al. A hidden

pitfall in the preparation of agar media undermines microorganism cultivability. Appl Environ

Microbiol. 2014;80: 7659-7666.

22. Bechtold HA, Marcarelli AM, Baxter CV, Inouye RS. Effects of N, P, and organic carbon on

stream biofilm nutrient limitation and uptake in a semi-arid watershed. Limnol Oceanogr.

2012;57: 1544–1554.

23. Hood JM, McNeely C, Finlay JC, Sterner RW. Selective feeding determines patterns of

nutrient release by stream invertebrates. Freshw Sci. 2014;33: 1093-1107.

24. Neeson TM, Salomon M, Coll M. Nutrient-specific foraging leads to Allee effects and

dynamic functional responses. Oikos. 2013;122: 265-273.

25. Mooney RJ, Strauss EA, Haro RJ. Nutrient-specific foraging by *Glossosoma intermedium*

larvae leads to conspecific case grazing. Freshw Sci. 2016;35: 873-881.

26. Rohatgi A. WebPlotDigitizer Version 3.9. Austin; 2015. Retrieved from

http://arohatgi.info/WebPlotDigitizer

27. Koricheva J, Gurevitch J, Mengersen K, editors. Handbook of meta-analysis in ecology and

evolution. Princeton: Princeton University Press; 2013.

28. Viechtbauer W. Conducting meta-analyses in R with the metafor package. J Stat Softw.

2010;36: 1-48.

29. Gelman A, Hill J. Data analysis using regression and multilevel/hierarchical models. 1st ed.

Cambridge: Cambridge University Press; 2007.

30. U.S. EPA. Method 353.2: determination of nitrate-nitrite nitrogen by automated colorimetry.

1993.

31. Murphy J, Riley JP. A modified single solution method for the determination of phosphate in

natural waters. Anal Chim Acta. 1962;27: 31-36.

32. U.S. EPA. Method 445.0: in vitro determination of chlorophyll *a* and pheophytin *a* in marine

and freshwater algae by fluorescence. 1997.

33. Musick HB. Phosphorus toxicity in seedlings of *Larrea divaricata* grown in solution culture.

Bot Gaz. 1978;139: 108-111.

34. Heddle EM, Specht RL. Dark Island Heath (Ninety-Mile Plain, South Australia). VIII. The

effect of fertilizers on composition and growth, 1950-1972. Aust J Bot. 1972;23: 151-164.

35. Groves RH, Keraitis K. Survival and growth of seedlings of three sclerophyll species at high

levels of phosphorus and nitrogen. Australian Journal of Botany. 1976;24: 681-690.

36. Rossiter RC. Phosphorus toxicity in subterranean clover and oats grown on Muchea sand,

and the modifying effects of lime and nitrate-nitrogen. Aust J Agric Res. 1952; 3: 227-243.

37. Stevenson RJ, Stoermer EF. Luxury consumption of phosphorus by five *Cladophora*

epiphytes in Lake Huron. Trans Am Microsc Soc. 1982;101: 151-161.

38. Chapin FS, Vitousek PM, Van Cleve K. The nature of nutrient limitation in plant

communities. Am Nat. 1986;127: 48-58.

39. Capps KA, Booth MT, Collins SM, Davison MA, Moslemi JM, El-Sabaawi RW, et al.

Nutrient diffusing substrata: a field comparison of commonly used methods to assess nutrient

limitation. J North Am Benthol Soc. 2011;30 :522–532.

40. Costello DM, Rosi-Marshall EJ, Shaw LE, Grace MR, Kelly JJ. A novel method to assess

effects of chemical stressors on natural biofilm structure and function. Freshw Biol. 2016;61:

2129-2140.

41. Grundon NJ. Mineral nutrition of some Queensland Heath plants. Journal of Ecology.

1972;60: 171-181.

42. Grant BR, Turner IM. Light-stimulated nitrate and nitrite assimilation in several species of

algae. Comp Biochem Physiol. 1969;29: 995–1004.

43. Parker DL, Kumar HD, Rai LC, Singh JB. Potassium salts inhibit growth of the

cyanobacteria *Microcystis spp*. in pond water and defined media: implications for control of

microcystin-producing aquatic blooms. Appl Environ Microbiol. 1997;63: 2324-2329.

44. Reisinger AJ, Tank JL, Dee MM. Regional and seasonal variation in nutrient limitation of

river biofilms. Freshw Sci. 2016;35: 474–89.

45. Brown EJ, Button DK, Lang DS. Competition between heterotrophic and autotrophic

microplankton for dissolved nutrients. Microb Ecol. 1981;7: 199–206.

46. Scott JT, Back JA, Taylor JM, King RS. Does nutrient enrichment decouple algal–bacterial

production in periphyton? J North Am Benthol Soc. 2008;27: 332-344.

47. Hoagland KD, Rosowski JR, Gretz MR, Roemer SC. Diatom extracellular polymeric

substances: function, fine structure, chemistry, and physiology. J Phycol. 1993;29: 537-566.

48. Pepe-Ranney C, Hall EK. The effect of carbon subsidies on marine planktonic niche

partitioning and recruitment during biofilm assembly. Front Microbiol. 2015; 6: 703.

49. Lourenço-Amorim C, Neres-Lima V, Moulton TP, Sasada-Sato CY, Oliveira-Cunha P,

Zandonà E. Control of periphyton standing crop in an Atlantic Forest stream: the relative roles of

nutrients, grazers and predators. Freshw Biol. 2014;59: 2365-2373.

50. Pringle CM, Blake GA. Quantitative effects of Atyid shrimp (Decapoda: *Atyidae*) on the

depositional environment in a tropical stream: use of electricity for experimental exclusion. Can

J Fish Aquat Sci. 1994;51: 1443-1450.

51. Moulton TP, de Souza ML, Silveira RML, Krsulović FAM. Effects of Ephemeropterans and

shrimps on periphyton and sediments in a coastal stream (Atlantic forest, Rio de Janeiro, Brazil).

J North Am Benthol Soc. 2004;23: 868-881.

52. Corkum LD. Patterns of nutrient release from nutrient diffusing substrates in flowing water.

Hydrobiologia. 1996;333: 37-43.

53. Keck F, Lepori F. Can we predict nutrient limitation in streams and rivers? Freshw Biol.

2012;57: 1410-1421.

54. Rugenski AT, Marcarelli AM, Bechtold HA, Inouye RS. Effects of temperature and

concentration on nutrient release rates from nutrient diffusing substrates. J North Am Benthol

Soc. 2008;27: 52–7.

Figures

Fig 1. Field experiment preparation methods and treatment classes. Four different

phosphate chemicals crossed with laboratory heating methodology were deployed in two NDS

experiments in 2016, for a total of eight preparation treatments. The eight treatments were

grouped by three treatment classes including cation, phosphate form, and heating method. The

same preparation treatments were crossed with two different phosphate concentrations in a 2017

experiment, for a total of sixteen preparation treatments. The sixteen treatments were grouped by

four treatment classes including cation, phosphate form, heating method, and concentration.

Heated Together = phosphate and agar boiled together, Heated Separately = agar boiled and

phosphate added at pouring temperature of 55-65° C.

Fig 2. Meta-analysis model results. (A) Results of a meta-analysis model testing the

significance of NDS P treatment autotrophic index (AI) effect sizes (n=45 experiments), where

lower values indicated a higher proportion of autotrophy in microbial communities. (B) Results

of a meta-analysis model testing the effect of grazing on NDS P treatment algal biomass

(chlorophyll *a*) effect sizes (n=5 experiments per grazing treatment). Squares are log response

ratio (LRR, see equation 2) mean estimates surrounded by 95% confidence interval bars.

Fig 3. **Effect of experimental preparation treatments on gross primary productivity**

(GPP). Gross primary productivity means ± 1 standard error from nutrient diffusing substrate

(NDS) field experiments. (A) Results from experiments 1-2 (total n=103), where treatments

consisted of crossing four chemicals (K_2HPO_4, KH_2PO_4, Na_2HPO_4, and NaH_2PO_4) and two heat

treatments (agar and phosphate heated together vs. heated separately, denoted as "HT" and

"HS"). An agar-only control treatment was included in each experiment. (B) Results from

experiment 3 (total n=104), where treatments included the same factors as experiments 1-2,

except low (0.05 M) and high (0.5 M) phosphate concentrations were included as an additional

factor. Gross primary productivity values by concentration are presented by averaging over all

other factors.

Fig 4. Effect of experimental preparation treatments on biomass-specific gross primary productivity (GPP). Biomass-specific GPP means ± 1 standard error from NDS field experiments (see Figure 2 caption). (A) For experiments 1-2, total n=102. (B) For experiment 3, total n=103.

Fig 5. Effect of experimental preparation treatments on chlorophyll *a*. Chlorophyll *a*

means ± 1 standard error from NDS field experiments (see Figure 2 caption). (A) For

experiments 1-2, total n=107. (B) For experiment 3, total n=105.

Fig 6. Effect of experimental preparation treatments on ash-free dry mass (AFDM). Ash-free dry mass means ± 1 standard error from NDS field experiments (see Figure 2 caption). (A) For experiments 1-2, total n=93. (B) For experiment 3, total n=104.

Fig 7. Effect of experimental preparation treatments on an autotrophic index (AI).

Autotrophic index means ± 1 standard error from NDS field experiments (see Figure 2 caption).

(A) For experiments 1-2, total n=92. (B) For experiment 3, total n=104.

Supporting Information Captions

S1 Table. Environmental variables measured during nutrient diffusing substrate experiments at Little Beaver Creek, CO. Multiple measurements are recorded as mean ± standard deviation. Several nutrient measurements were below detection. pH was not measured during experiment 3 because of an instrument malfunction.

S2 Table. Results of ANOVAs testing the effect of three treatment classes on P effect sizes. Four different phosphate chemicals (KH_2PO_4, K_2HPO_4, NaH_2PO_4, Na_2HPO_4) crossed with laboratory heating methodology were deployed in two NDS experiments in 2016. Treatment classes in the statistical analyses included phosphate cation, phosphate form, and heating method. ANOVA results testing the effect of each treatment class on each response variable's P treatment effect size are presented, with experiment included as a fixed effect block in all models. P<0.05 is indicated as "*".

S3 Table. Results of ANOVAs testing the effect of four treatment classes on P effect sizes. Four different phosphate chemicals (KH_2PO_4, K_2HPO_4, NaH_2PO_4, Na_2HPO_4) crossed with laboratory heating methodology and two phosphate concentrations were deployed in an NDS experiment in 2017. Treatment classes in the statistical analyses included phosphate cation, phosphate form, and heating method, along with concentration as a potential modifier. Two-way ANOVA results are presented where each treatment class was crossed with concentration, to determine effects on each response variable's P treatment effect size. P<0.05 is indicated as "*".

S1 References. Studies used in autotrophic index meta-analysis.

S2 References. Studies used in grazer exclusion meta-analysis.