2
3
4 TNF-α induces reactivation of human cytomegalovirus independently of myeloid cell

5 differentiation following post-transcriptional establishment of latency

7 Running title: Differentiation-independent reactivation of HCMV

8
9

Eleonora Forte[a], Suchitra Swaminathan[b], Mark W. Schroeder[a,] Jeong Yeon Kim[a], Scott S.

Terhune[c,d], and Mary Hummel[a#]

[a]Comprehensive Transplant Center, Department of Surgery, Northwestern University Feinberg

School of Medicine, Chicago, IL

[b]Division of Rheumatology, Department of Medicine, Northwestern University Feinberg School

of Medicine, Chicago, IL

[c]Department of Microbiology and Immunology, Medical College of Wisconsin, Milwaukee, WI

[d]Biotechnology and Bioengineering Center, Medical College of Wisconsin, Milwaukee, WI

Abstract word count: 243

Text word count: 5080

#Address correspondence to Mary Hummel, m-hummel@northwestern.edu

ABSTRACT: We used the Kasumi-3 model to study HCMV latency and reactivation in

myeloid progenitor cells. Kasumi-3 cells were infected with HCMV strain TB40/E*wt*-GFP, flow

sorted for GFP+ cells, and cultured for various times to monitor establishment of latency, as

judged by repression of viral gene expression (RNA/DNA ratio) and loss of virus production.

We found that, in the vast majority of cells, latency was established post-transcriptionally in the

GFP+ infected cells: transcription was initially turned on, and then turned off. We also found

that some of the GFP- cells were infected, suggesting that latency might be established in these

cells at the outset of infection. We were not able to test this hypothesis because some GFP- cells

expressed lytic genes, and thus, it was not possible to separate them from GFP- quiescent

cells. In addition, we found that the pattern of expression of lytic genes that have been

associated with latency, including UL138, US28, and RNA2.7, was the same as that of other

lytic genes, indicating that there was no preferential expression of these genes once latency is

established. We confirmed previous studies showing that TNF-α induced reactivation of

infectious virus, and by analyzing expression of the progenitor cell marker CD34 as well as

myeloid cell differentiation markers in IE+ cells after treatment with TNF-α, we showed that

TNF-α induced transcriptional reactivation of IE gene expression independently of

differentiation. TNF-α-mediated reactivation in Kasumi-3 cells was correlated with activation of

NF-κB, KAP-1 and ATM.

IMPORTANCE: HCMV is an important human pathogen that establishes lifelong latent

infection in myeloid progenitor cells, and reactivates frequently to cause significant disease in

immunocompromised people. Our observation that viral gene expression is first turned on, and

then turned off to establish latency suggests that there is a host defense, which may be myeloid-

specific, responsible for transcriptional silencing of viral gene expression. Our observation that

TNF-α induces reactivation independently of differentiation provides insight into molecular

mechanisms that control reactivation.

INTRODUCTION

HCMV is a ubiquitous human pathogen of the beta herpesvirus family. Infection is

transmitted through contact with body fluids, including saliva, urine, blood, and genital

secretions. Hematogenous spread of the virus leads to a systemic infection, with multiple cell

types infected in many organs. In immunocompetent hosts, primary infection is typically sub-

clinical, and is resolved by activation of innate and adaptive immunity. However, latently

infected cells, which carry viral DNA, but do not produce infectious virus, are able to escape

immune surveillance, and can persist for the life of the host.

Under the appropriate conditions, the virus in these cells wakes up from its dormant state

to reactivate the infectious cycle. In immunocompetent hosts, viral replication is controlled by

the immune response, but reactivation of latent virus can be a significant infectious complication

in immunocompromised hosts, such as recipients of solid organ or bone marrow transplants.

Reactivation of CMV in solid organ transplant recipients is associated with increased risk of

CMV disease, acute and chronic allograft rejection, infection with other opportunistic pathogens,

graft failure, and death (1). Risk factors for reactivation include CMV serostatus of the donor

(D) and recipient (R), with the highest risk in the D+/R- combination, specific

immunosuppression protocols, and inflammatory conditions associated with high TNF-α

secretion, including allograft rejection and sepsis (1).

The cell type harboring latent CMV DNA in solid organs has not been clearly identified.

However, it is known that latent virus is transmitted through blood transfusion, and monocytes in

the blood and hematopoietic progenitor cells (HPCs) in the bone marrow are sites of latency in

vivo (2-10). Experimental models have shown that these cells are less permissive to lytic

replication, and that they support a latent infection (11-16). Cell type-specific establishment of

latency is thought to be due to a combination of host and viral factors. Infection activates a host

intrinsic immune response, which recognizes viral DNA invading the nucleus and silences viral

gene expression at the outset of infection through heterochromatinization of viral genomes (13,

17-26). Factors present in the viral particle, including the tegument protein pp71, enter the cell

upon infection, and counteract this host defense response to activate viral gene expression. In

cells that support latency, pp71 is sequestered in the cytoplasm and is therefore unable to

perform this function (26-28). Differentiation of myeloid cells to dendritic cells increases

permissiveness of these cells to infection and also induces reactivation of both naturally and

experimentally latently infected cells (12, 13, 15, 20, 29-32). Thus, the generally accepted

paradigm is that myeloid progenitor cells are not permissive to infection due to

heterochromatinization and transcriptional repression of invading viral genomes. The process of

myeloid cell differentiation is thought to change the balance of repressive and activating cellular

factors that control both viral and cellular transcription, resulting in changes to the viral

epigenome that activate viral gene expression (17, 33, 34).

However, some recent observations suggest that the current paradigm may be in need of

revision. First, several investigators have demonstrated that, although myeloid progenitors may

be less permissive than other cell types, some expression of viral lytic genes is detectable at early

times post-infection in various models of latency, including primary CD34+ HPCs, Kasumi-3

cells, a CD34+ myeloblastic cell line derived from a patient with acute myeloid leukemia (AML)

(35), which is a tractable model for CMV latency and reactivation (27, 36, 37), and primary

CD14+ peripheral blood mononuclear cells (11, 12, 36-43). Expression of these genes is lost as

latency is established. These observations suggest that latency is not necessarily established at

the outset of infection, but may also be established following activation of viral transcription.

Second, differentiation of myeloid cells into dendritic cells *in vivo* occurs in the context

of infection and inflammation (44), and all of the physiologically relevant differentiation factors

that have been used to induce reactivation of CMV in primary hematopoietic cells, including

GM-CSF (45), TNF-α, IL-6, and LPS, are mediators of inflammation. Thus, it is difficult to

distinguish the roles of inflammation versus differentiation in these models. In contrast, Kasumi-

3 cells, like other transformed cell lines, are refractory to normal physiological differentiation

cues, and therefore can be used to dissect the contributions of these two pathways. Here, we

have used the Kasumi-3 model to further investigate these issues.

RESULTS

Latency is established after activation of transcription. The time at which latency is

established following infection with HCMV has not been clearly determined. Some previous

studies have shown that primary CD34+ myeloid progenitor cells do not express lytic viral genes

upon infection, but rather, that viral genomes in these cells are inactivated upon entry and remain

quiescent until they receive signals that induce them to differentiate (26, 46). However, other

studies have shown that both primary CD34+ cells and Kasumi-3 cells express an SV40

promoter-driven GFP reporter, as well as viral genes at early times post-infection (11, 36-41),

and that latency can be established in the GFP+ population following cell sorting (11, 36, 38, 40,

41, 47). To investigate the question of timing further, we infected Kasumi-3 cells with the

TB40/E*wt*-GFP strain of HCMV (48), and analyzed the cells by FACS after 24 hr (Fig. 1A).

GFP expression was observed in 20-50% percent of cells (Fig. 1B), as previously reported (36).

Following inactivation of input virus, we purified the GFP+ population by cell sorting and

analyzed viral RNA expression, DNA copy number, and production of infectious virus over the

course of infection. Latency has been operationally defined as the lack of virus production in

cells that carry viral DNA with the capacity to reactivate (49).   Production of infectious virus

was detectable at 4 dpi and peaked at 8 dpi, but was no longer detectable at 11dpi (Fig. 1C).

A second characteristic of latency is transcriptional repression of genes involved in lytic

replication. We therefore analyzed expression of viral genes representative of the immediate

early (IE-2, UL122 and IE-1, UL123), early (viral DNA polymerase UL54), and late (UL32)

phases of the viral life cycle. Expression of these genes peaked at 4 dpi and then fell throughout

the course of infection. Viral DNA copy number also increased following infection, peaked 4-8

dpi, and then decreased significantly (Fig. 1D and Fig. S1). Because both viral DNA and RNA

copy number dropped over the course of infection, analysis of RNA expression alone does not

indicate repression of viral transcription. We therefore calculated the ratio of viral RNA to DNA.

Relative to 1 dpi, this ratio fell for all genes from 4 to 14 dpi (Fig. 1D and E). TNF-α induced

reactivation of infectious virus in these cells (see below). It is important to note that, because the

cells were sorted for GFP expression to obtain a pure population of infected cells in which viral

genomes were transcriptionally active, these results show that latency can be established by

shutting off gene expression after it has been turned on. We have introduced the term post-

transcriptional latency to describe this state.

139
Some GFP- cells contain viral DNA and express lytic transcripts. Although our

studies indicate that latency can be established post-transcriptionally, we wondered whether there

might be a second population of cells in which latency is established at the outset of infection.

These cells would carry viral DNA, but would not express GFP or viral genes. To test this

hypothesis, we analyzed both the GFP-positive and GFP-negative populations for the presence of

viral DNA at 1 dpi. To preclude cross-contamination of these fractions, the gates for sorting were

set far apart. qPCR analysis revealed that, although the vast majority of the DNA was present in

the GFP+ cells, some DNA was detectable in the GFP- population (Fig. 2C). We then analyzed

the RNA from these cells for expression of immediate early, early, and late genes (Fig. 2D).

Expression of immediate early (IE-2, UL122) RNA was detectable in the GFP- population, albeit

at a lower level than the GFP+ cells. When we calculated the ratio of UL122 RNA to DNA, we

found that there was no difference between the GFP+ and GFP- cells, indicating that

transcription of the IE genes was equally active in these two populations at this time (Fig. 2E).

However, when we analyzed the RNA/DNA ratios of early (UL54) and late (UL32) genes, we

found that this ratio was significantly lower in the GFP- cells (Fig. 2E), indicating that these

classes of genes were transcriptionally repressed in the GFP- cells. We then analyzed the GFP-

cells for changes in GFP expression, viral RNA expression and DNA copy number over time.

We found that some GFP- cells become GFP+ by 4 dpi (Fig. 2B). Expression of all classes of

viral RNA and DNA copy increased in these cells from 1 to 4 dpi, and then declined after 7 dpi

(Fig. S2).

In order to determine whether there was a subset of cells in the GFP- population

containing quiescent viral DNA, we re-sorted these cells at 4 dpi into GFP-positive and -negative

cells for analysis of viral DNA and RNA (Fig. 2A and B). The results show that viral DNA was

detectable in the GFP- population at 4 dpi (Fig. 2C, GFP-D4). Analysis of RNA expression

showed that some expression of lytic genes was still detectable in the GFP- population at 4 dpi

(Fig. 2D,GFP-D4). We conclude that *i)* the GFP reporter is regulated as an early gene in

TB40/E*wt*-GFP infected Kasumi-3 cells, as previously noted in fibroblasts (50); *ii)* there is

heterogeneity in the kinetics of viral gene expression after infection of Kasumi-3 cells, with

activation of early and late gene expression delayed in some cells; *iii)* it is not possible to use

sorting for GFP expression to determine whether a small minority of Kasumi-3 cells infected

with TB40/E*wt*-GFP contain viral genomes, but remain quiescent.

171
US28, UL138, and 2.7 kb RNAs are not differentially expressed in latently infected

Kasumi-3 cells. A number of studies have analyzed the viral transcriptome in latency, and some

of these have indicated that some lytic genes are also specifically expressed in latency, including

US28, UL138, and a non-coding 2.7 kb RNA (RNA2.7) (12, 14, 37, 46, 47, 49, 51-53). We

analyzed expression of these genes in GFP+ cells over the course of infection in Kasumi-3 cells.

RNA2.7 is expressed at very high levels during lytic infection of fibroblasts (54), and at early

times after infection of Kasumi-3 cells (Fig. 3A). UL138 and US28 RNAs are also relatively

abundant transcripts during lytic infection. To analyze repression of these genes relative to a

lytic gene, we analyzed UL32, a gene that is expressed at levels similar to UL138 and US28 at 4

dpi. Our results show that expression of UL138, US28, and RNA2.7 followed the same pattern

as that observed for UL32, with a peak of expression at 4 dpi, followed by a significant decrease

at 7 dpi (Fig. 3A). When we analyzed the RNA/DNA ratio, we found that expression of these

genes was repressed over the course of infection, with kinetics similar to that of UL32 (Fig. 3C).

The RNA/DNA ratio of RNA2.7 was significantly higher than other genes at 4 dpi, consistent

with its high level of expression during lytic infection, but there was no difference between

UL32 and UL138 or US28 at any time point, and there were no differences in the RNA/DNA

ratios between UL32 and any of the other genes analyzed after 4 dpi (Fig. 3C). Thus, there was

no preferential expression of UL138, US28, or RNA2.7 over the lytic gene UL32 when latency

was established.

TNF-α induces reactivation of HCMV in Kasumi-3 cells independently of myeloid

cell differentiation. Previous studies showed that TNF-α induced reactivation of HCMV in

latently infected Kasumi-3 cells (36). We confirmed this result using slightly different

conditions. Kasumi-3 cells were infected with TB40/E*wt*-GFP virus at an MOI of 1-2, flow-

sorted for GFP+ cells and cultured for 14 days to establish latency. The cells were then divided

into different pools and incubated for a further 3 days in the presence of 5 ng/ml TNF-α or left

untreated. Relative to untreated controls, TNF-α induced statistically significant increases in

viral DNA copy, RNA expression and the RNA/DNA ratio, as well as production of infectious

virus (Fig. 4A and additional RNA analyses of early and late genes in Fig. S3). In addition, we

analyzed the percentage of reactivating cells by flow cytometry after staining with an IE1/2-

specific antibody. Expression of IE proteins was essentially undetectable in latently infected,

untreated Kasumi-3 cells (Fig. 4B, upper right panel). Treatment with TNF-α reproducibly

induced expression of IE1/2 proteins in 2-3% of the cells (representative results shown in Fig. 4B,

lower left panel).

Differentiation of myeloid progenitor cells to a dendritic cell phenotype has long been

associated with transcriptional activation of HCMV IE gene expression and reactivation of

infectious virus (13, 20, 34, 40). Transformed cell lines are generally resistant to differentiation,

and therefore it seemed unlikely that differentiation would play a role in this model. However,

the fact that reactivation occurred in only a small percentage of latently infected Kasumi-3 cells

left open the possibility that reactivation occurred in a small number of cells with the capacity to

differentiate. To investigate this question, we used immunophenotyping to characterize the

differentiation state of Kasumi-3 cells carrying reactivating virus. Latently infected TNF-

α treated or untreated cells were stained for IE1/2 antibody in combination with the

hematopoietic progenitor marker CD34, and for markers of various stages of myeloid

differentiation, including CD64, a marker of granulo-monocytic lineage commitment (55), CD14

(monocytes), CD15 (granulocytes), and CD1c and CD11c, which identify major subpopulations

of human myeloid dendritic cells (56). Uninfected Kasumi-3 cells are homogeneous with respect

to CD34 expression, and do not express other markers of myeloid cell differentiation (Fig. 4B,

upper left and Fig. S4). Expression of CD34 was lost in a small percentage of cells after

establishment of latency, but latent cells did not differentiate (Fig. S4B). All the latently infected

cells were negative for expression of IE proteins (Fig. 4B, upper right panel). 2-3% of the cells

became IE+ after treatment with TNF-α, and these cells retained expression of CD34 (Fig. 4B,

lower left panel). We then gated on the IE+/CD34+ population to analyze expression of

differentiation markers in reactivating cells. ~95% of the IE+/CD34+ cells were negative for

CD64- and for markers of mature monocytes and dendritic cells (Fig. 4C). Thus, our data shows

that differentiation was not required for transcriptional reactivation of IE gene expression in this

model.

TNF-α induces activation of NF-κB and a DNA damage response in Kasumi-3 cells.

Binding of TNF-α to the TNFR1 receptor activates signaling cascades, which lead to activation

of transcription factors that control activation of the HCMV Major Immediate Early Promoter

(MIEP), including canonical NF-κB (p65/p50) (57-59). Because the IE genes are

transcriptionally silent in latently infected cells and they are required for activation of lytic

replication, activation of the MIEP is thought to be a requisite first step in reactivation of the

virus. Activation of NF-κB is controlled at multiple levels, including cellular localization and

post-translational modification at multiple sites (60). TNF/TNFR1-mediated activation of IKK

leads to degradation of the inhibitory IκB subunit, which permits translocation of the active

p65/p50 complex from the cytoplasm into the nucleus. In addition, IKK mediates

phosphorylation of p65 S536 (61, 62), which leads to enhanced transactivation through increased

CBP/p300 binding and acetylation of p65 K310 (63). We therefore analyzed total levels of p65,

p-p65 S536, and the ratio of p-p65 S536/total p65 in both uninfected and latently infected

Kasumi-3 cells treated with or without TNF-α (Fig. 5A). Our results show that all of these

increased following treatment with TNF-α in latently infected cells.

Previous studies showed that TNF-α induces activation of ATM (64), and that activation

of ATM is sufficient to induce reactivation of HCMV in primary hematopoietic progenitor cells

through phosphorylation of KAP-1 (S824) bound to latent viral genomes (40). We therefore

analyzed total and phospho-ATM (S1981), total and phospho-KAP-1 (S824) and the ratio of

phospho/total proteins. In addition, we analyzed phosphorylation of H2AX (γ-H2AX), a variant

histone that is recruited to sites of DNA damage (65). Our results show that all of these markers

of DNA damage are increased by treatment of Kasumi-3 cells with TNF-α (Fig. 5B-D).

DISCUSSION

Mechanisms of establishment of latency. HCMV establishes latency in myeloid

progenitor cells through heterochromatinization and repression of viral gene expression, but the

timing and molecular mechanisms that control this process are not well understood. While some

studies have shown that viral gene expression is silenced immediately after infection (26, 46),

others have shown that viral gene expression is activated in various models at early times post-

infection, and that latency is subsequently established in this population of cells (11, 36, 38, 40,

41, 47). Our studies are in agreement with this latter observation. We found that ~90% of

infected Kasumi-3 cells become GFP+ at 1 dpi. From 4-8 dpi, expression of all classes of viral

genes was activated in this population, viral DNA was amplified, and infectious virus was

produced. At later times, viral RNA expression and DNA copy number, and the RNA/DNA

ratio decreased. Some lytically infected cells die after infection, and the drop in RNA/DNA may

be due in part to loss of these cells. However, we observed a significant decrease in RNA/DNA

between live GFP+ cells isolated at 1 dpi, prior to robust viral replication, and 17 dpi, when the

cells maintain viral DNA, but expression of viral genes is nearly undetectable. This difference is

due at least in part to transcriptional repression of viral genes during the transition from a lytic to

latent infection.

Thus, we find that, although activation of the MIEP is lower in myeloid progenitor cells

than permissive fibroblasts (26), Kasumi-3 cells support the full cycle of lytic replication. In

contrast to fibroblasts, where the infection is propagated throughout the culture to kill all the

cells, our studies suggest that some infected Kasumi-3 cells are able to shut down viral

replication and survive. Our studies are in agreement with previous studies of Kasumi-3 cells,

which showed increased expression of lytic RNAs and amplification of viral DNA relative to

early times post-infection, and, in some cases, production of infectious virus, followed by

transcriptional repression (36, 37, 66). Although Kasumi-3 cells may be more permissive than

primary CD34+ HPCs, previous studies have also shown activation of viral gene expression in

HPCs prior to establishment of latency (11, 38, 41, 47).

Recent studies indicate that post-transcriptional establishment of latency may be a more

general feature of herpesvirus latency (67-70). Previous studies have shown that activation of

the immediate early ICP0 promoter was detectable in some, but not all neurons of mice latently

infected with HSV (68). Activation of the late gC promoter was also detectable in the acute

stage of infection, but not in latently infected mice. This observation suggests that, although

some viral genes may be expressed prior to establishment of latency, there is a point of no return,

and full replication of the virus is not compatible with survival. Interestingly, these authors also

demonstrated activation of the HCMV MIEP prior to establishment of latency in neurons

infected with HSV.

We considered the possibility that there may be two routes to establishment of latency:

repression prior to activation of gene expression as well as post-transcriptional repression. Our

results show that ~10% of the infected cells are GFP- at 1 dpi. We found that there is

heterogeneity in the kinetics of viral gene expression, and that some GFP- cells become GFP+,

activate the early and late phases of viral gene expression, and amplify viral DNA with delayed

kinetics. Previous studies showed that initiation of HCMV gene expression requires that the cells

be in the G0 or G1 phase of the cell cycle (71), and heterogeneity in cell cycle phase may

account for some of the observed variation in the kinetics of gene expression. The majority of

the cells that were GFP- at 1 dpi remain GFP- at 4 dpi. But, because some GFP- cells expressed

lytic viral transcripts, we were unable to determine whether there were other cells that retain viral

genomes, but do not express viral RNAs. However, since 90% of the infected cells are GFP+ at

1 dpi, these cells would constitute a very small minority of the infected cells.

Our findings have two important implications. First, the observation that viral gene is

expression is first turned on, and then turned off, suggests that there is a mechanism for shutting

off viral genomes that are actively engaged in transcription. This may be due to expression of

viral genes that facilitate establishment of latency (37, 41, 46-49, 66, 72, 73). However, these

genes are also expressed in models of lytic infection, and could not act directly to repress viral

gene expression, since they are not localized to the nucleus. An attractive hypothesis that would

account for the highly restricted tropism for HCMV latency is that there is a host defense

response that shuts off viral transcription specifically in myeloid lineage cells.

Second, shedding of HCMV is often observed in healthy, immunocompetent individuals,

suggesting that the virus reactivates with high frequency (74). The finding that HCMV

establishes latency in cells following activation of transcription raises the possibility that these

genomes may retain a memory of prior activation, and may therefore be poised to reactivate

readily under the appropriate conditions. Interestingly, previous studies have shown that HCMV

chromatin is specifically marked by H3 lysine 4 methylation following replication of viral DNA

in fibroblasts (75). It is tempting to speculate that there may be cell type-specific differences in

histone modifications bound to post-replicative viral DNA that affect the potential to establish

latency and to reactivate.

Viral transcription in latency. Previous studies showed that some viral genes that are

expressed during lytic infection, particularly UL138 and US28, facilitate establishment of

latency in myeloid progenitor cells (37, 41, 46-49, 66, 72, 73). However, analysis of expression

of these genes in latently infected cells has been contradictory and confusing. Some studies

have shown that some genes are selectively expressed in latency (12, 14, 37, 46, 47, 51-53). In

contrast, studies using unbiased transcriptome analyses to detect viral transcripts in both

experimental and natural latency, have identified many genes expressed under conditions of

latency, and none of these show specific expression of some RNAs in the absence of all other

lytic genes (11, 39, 42, 43, 76). This raises the possibility that detection of these transcripts is

due to the presence of a small population of lytically infected cells. With the exception of the

noncoding miR-UL148D microRNA (37), most previous studies have not included analyses of

the RNA/DNA ratio to demonstrate differences in transcriptional activity between lytic genes

and latency-associated lytic genes. With the exception of LUNA (14), there has been no analysis

of histones bound to latency-associated genes in latently infected cells to support the idea that

they are selectively active. We have investigated expression of US28, UL138, as well as the non-

coding RNA2.7 in the Kasumi-3 model. We find that the pattern of expression of these genes

parallels that of other lytic genes, including UL122, UL123, UL54, and UL32, and analysis of

the RNA/DNA ratio shows that transcription of all of these genes is repressed in latency. We

suggest that this type of analysis should be included in future studies to demonstrate latency-

specific expression of a gene. Collectively, our studies and others are consistent with the idea

that these genes facilitate establishment of latency through changes in the relative abundance of

viral transcripts, but that they are shut off when latency is established. This situation would be

analogous to that of EBV latency, where the EBNA genes are expressed in naive B cells, but are

shut off in memory B cells, which are the long term site of latency (77) and HSV, where ICP0 is

initially expressed to turn on LAT expression and facilitate heterochromatinization of viral

genomes (67).

What is the role of myeloid cell differentiation in reactivation of HCMV? Previous

studies using primary CD34-derived or monocyte-derived dendritic cells (MoDCs) showed that

LPS-induced reactivation of latent HCMV required differentiation to a dendritic cell phenotype

(9, 13, 20, 30, 32, 52, 78). Reactivation in immature dendritic cells in response to LPS occurred

through induction of IL-6 and activation of ERKs and mitogen and stress-activated kinases

(MSK) (52, 79). However, the factors that were used to differentiate dendritic cells in these

models are also mediators of inflammation, and it is therefore difficult to distinguish the roles of

a general inflammatory response, which could occur in many cell types, from a dendritic cell-

specific mechanism of reactivation. In contrast, Kasumi-3 cells, like many transformed cell lines,

are refractory to the normal cues that induce differentiation, and are therefore more suitable for

studies seeking to dissect the roles of these different processes. Our studies show that

differentiation is not required for reactivation in response to TNF-α.

Previous studies showed that, in latently infected primary CD34+ hematopoietic

progenitor cells, viral genomes are bound to KAP-1, and that KAP-1 is required to establish

latency through heterochromatinization and transcriptional silencing (40). Reactivation could be

induced by treatment of the cells with chloroquine, an agent that induces phosphorylation of

KAP-1 through activation of ATM, or by knock-down of KAP-1, and this effect was potentiated

by co-treatment with TNF-α. These cells retained expression of CD34, suggesting that

differentiation was not required for reactivation in this model, although expression of dendritic

cell markers was not analyzed (40).

ATM is the master regulator of the response to double-stranded DNA breaks (DSB) (80).

ATM is recruited to damage sites by the Mre11-Rad50-Nbs1 (MRN) complex, where it

phosphorylates a variety of substrates that facilitate DNA repair, including KAP-1. KAP-1 is a

multifunctional protein whose activity is regulated through post-translational modifications (81).

The sumoylated form of KAP-1 acts as a repressor of transcription through recruitment of factors

that mediate H3K9 de-acetylation/methylation. ATM-mediated phosphorylation of KAP-1 leads

to changes in its interaction partners that result in de-condensation of chromatin to allow repair

of damaged sites, and this is especially important in repair of breaks that occur in regions of

heterochromatin (81-83). The results of Rauwel, et al. (40) therefore suggest that reactivation of

HCMV can be induced through chromatin remodeling induced by activation of a DNA damage

response independently of differentiation to dendritic cells. Co-treatment with TNF-α

potentiated reactivation in response to chloroquine, suggesting that inflammatory mediators that

activate NF-κB can act synergistically with chromatin remodeling to induce reactivation of

HCMV. Our studies showed that TNF-α was sufficient to induce reactivation in Kasumi-3 cells

without inducing differentiation, and that this correlated with activation of NF-κB, as well as

ATM and KAP-1. Further studies will be required to fully understand the roles of differentiation,

inflammation, and chromatin remodeling in reactivation of HCMV. However, the results of our

studies, as well as those of others, suggest that the mechanisms of reactivation of HCMV are not

necessarily limited to a dendritic cell specific pathway, but may be induced by more general

pathways of inflammation and cellular damage.

MATERIALS AND METHODS

Cells and reagents: Kasumi-3 and MRC-5 cells were obtained from the ATCC. Human

recombinant TNF-α was obtained from PreproTech (#300-01A). Fluorescent conjugated

monoclonal antibodies used in this study are listed in Table 1. TB40/E*wt*-GFP (48) viral stocks

were produced and titered as described in supplementary methods.

Infection of Kasumi-3 cells: Kasumi-3 cells were infected with HCMV strain TB40/E*wt*-GFP

(48) (MOI 1-2) using centrifugal enhancement of infection, as described in supplementary

methods (84). GFP+ and GFP- cells were sorted on a FACSAria-6 laser cell sorter (BD

Biosciences). Infected cells were cultured for 13 days to establish latency, with media changes

every 2-3 days. For reactivation studies, cells were treated with 5ng/ml TNF-α at 14 dpi for 3

days at 37°C. A $TCID_{50}$ assay was used to determine the infectious titer of infected cell

supernatants.

RNA and DNA extraction: DNA was isolated with the Arcturus pico Pure DNA extraction kit

(Thermo Fisher Scientific) as recommended by the manufacturer. RNA was purified with the

Direct-zol RNA mini prep kit (Zymo Research) with a DNAse digestion step following

manufacturer's instructions.

Real time PCR analysis. For all analyses, mock-infected cells were analyzed in parallel with

infected cells. These samples were uniformly positive for cellular genes, but negative for viral

genes. TaqMan assays specific for viral genes were custom-designed by Life Technologies

(Table 2). Relative viral DNA and RNA quantity was analyzed by the 2-($\Delta\Delta$Ct) method, using

pre-designed assays (Life Technologies) for RNaseP (catalog number 4403326) or GAPDH,

respectively, as the normalization controls. This method was validated by determining that the

efficiencies of the target and reference genes were approximately equal (85) as described in

supplementary methods (Fig. S5 and S6). Previous studies have shown that GAPDH is an

appropriate normalization control for RNA analysis of CMV-infected cells (86), and we verified

that GAPDH Ct values are stable following infection of Kasumi-3 cells (Fig. S7). TaqMan gene

expression assays purchased from Life Technologies were used for analysis of cellular RNAs

(Hs00174128_m1 for TNF-α and Hs99999905_m1 for GAPDH).

Immunophenotyping. Cells were stained with antibodies specific for CD34, CD64, CD14,

CD15, CD11c, CD1c and IE 1/2 (CH160) (Table 1) as described in supplementary methods.

Data on live cells was acquired on 6-laser BD LSRFortessa SORP flow cytometer and analyzed

using FlowJo software (version 9).

Cytokine stimulation and intracellular phospho-protein analysis: Kasumi-3 cells were

treated or untreated with 5 ng/ml TNF for 30' at 37^{o}C, fixed, permeabilized and washed

according to previously described protocols (87, 88). Cells were stained with antibodies specific

for p65 (NF-κB), p-p65 (pNF-κB), ATM, p-ATM, KAP-1, p-KAP-1 or γ-H2AX (Table 1).

Antibodies were validated using published protocols (89) as described in supplementary methods

and Fig. S8. Data were acquired on BD LSRFortessa SORP flow cytometer and analyzed using

FlowJo (version 9). The fraction of responding cells for each population was determined as

described in supplementary methods as previously described (88, 89).

ACKNOWLEDGEMENTS. This work was supported by grant P01AI112522 to Michael

Abecassis and R01 AI083281 to Scott Terhune from the National Institute of Allergy and

Infectious Diseases and by a Cancer Center Support Grant (NCI CA060553) to the

Northwestern University Flow Cytometry Core Facility. Cell sorting was performed on a BD

FACSAria SORP system, purchased through the support of NIH 1S10OD011996-01. The

authors thank Katie Cataldo for preparation of virus stocks, Paul Mehl and Carolina Ostiguin

for cell sorting, Dr. Robert Kalejta for the suggestion to analyze RNA/DNA ratios during

establishment of latency, and Dr. Michael Abecassis for critical discussions and support. The

authors have no conflicts of interest to disclose.

434
435
436

REFERENCES

1.   **Eid AJ, Razonable RR.** 2010. New developments in the management of

cytomegalovirus infection after solid organ transplantation. Drugs **70:**965-981.

2.   **Gilbert GL, Hayes K, Hudson IL, James J.** 1989. Prevention of transfusion-acquired

cytomegalovirus infection in infants by blood filtration to remove leucocytes. Neonatal

Cytomegalovirus Infection Study Group. Lancet **1:**1228-1231.

3.   **Hahn G, Jores R, Mocarski ES.** 1998. Cytomegalovirus remains latent in a common

precursor of dendritic and myeloid cells. Proc Natl Acad Sci U S A **95:**3937-3942.

4.   **Mendelson M, Monard S, Sissons P, Sinclair J.** 1996. Detection of endogenous human

cytomegalovirus in CD34+ bone marrow progenitors. J Gen Virol **77 (Pt 12):**3099-3102.

5.   **Sindre H, Tjoonnfjord GE, Rollag H, Ranneberg-Nilsen T, Veiby OP, Beck S, Degre**

M, Hestdal K. 1996. Human cytomegalovirus suppression of and latency in early

hematopoietic progenitor cells. Blood **88:**4526-4533.

6.   **Zhuravskaya T, Maciejewski JP, Netski DM, Bruening E, Mackintosh FR, St Jeor S.**

1997. Spread of human cytomegalovirus (HCMV) after infection of human

hematopoietic progenitor cells: model of HCMV latency. Blood **90:**2482-2491.

7.   **Taylor-Wiedeman J, Sissons JG, Borysiewicz LK, Sinclair JH.** 1991. Monocytes are a

major site of persistence of human cytomegalovirus in peripheral blood mononuclear

cells. Journal of General Virology **72:**2059-2064.

8.   **Bolovan-Fritts CA, Mocarski ES, Wiedeman JA.** 1999. Peripheral blood CD14+ cells

from healthy subjects carry a circular conformation of latent cytomegalovirus genome.

Blood **93:**394-398.

9.   **Minton EJ, Tysoe C, Sinclair JH, Sissons JG.** 1994. Human cytomegalovirus infection

of the monocyte/macrophage lineage in bone marrow. Journal of Virology **68:**4017-4021.

10.   **Slobedman B, Mocarski ES.** 1999. Quantitative analysis of latent human

462   cytomegalovirus. J Virol **73:**4806-4812.

463   11.   **Goodrum FD, Jordan CT, High K, Shenk T.** 2002. Human cytomegalovirus gene

464   expression during infection of primary hematopoietic progenitor cells: a model for

465   latency. Proc Natl Acad Sci U S A **99:**16255-16260.

466   12.   **Hargett D, Shenk TE.** 2010. Experimental human cytomegalovirus latency in CD14+

467   monocytes. Proc Natl Acad Sci U S A **107:**20039-20044.

468   13.   **Reeves MB, Lehner PJ, Sissons JG, Sinclair JH.** 2005. An in vitro model for the

469   regulation of human cytomegalovirus latency and reactivation in dendritic cells by

470   chromatin remodelling. J Gen Virol **86:**2949-2954.

471   14.   **Reeves MB, Sinclair JH.** 2010. Analysis of latent viral gene expression in natural and

472   experimental latency models of human cytomegalovirus and its correlation with histone

473   modifications at a latent promoter. J Gen Virol **91:**599-604.

474   15.   **Hertel L, Lacaille VG, Strobl H, Mellins ED, Mocarski ES.** 2003. Susceptibility of

475   immature and mature Langerhans cell-type dendritic cells to infection and

476   immunomodulation by human cytomegalovirus. J Virol **77:**7563-7574.

477   16.   **Rice GP, Schrier RD, Oldstone MB.** 1984. Cytomegalovirus infects human

478   lymphocytes and monocytes: virus expression is restricted to immediate-early gene

479   products. Proc Natl Acad Sci U S A **81:**6134-6138.

480   17.   **Kalejta RF.** 2013. Pre-Immediate Early Tegument Protein Functions, p 141-151. *In*

481   Reddehase MJ (ed), Cytomegaloviruses: From Molecular Pathogenesis to Intervention,

482   vol 1. Caister Academic Press, Norfolk, UK.

18. **Saffert RT, Kalejta RF.** 2006. Inactivating a cellular intrinsic immune defense mediated

by Daxx is the mechanism through which the human cytomegalovirus pp71 protein

stimulates viral immediate-early gene expression. J Virol **80:**3863-3871.

19. **Saffert RT, Kalejta RF.** 2007. Human cytomegalovirus gene expression is silenced by

Daxx-mediated intrinsic immune defense in model latent infections established in vitro. J

Virol **81:**9109-9120.

20. **Reeves MB, MacAry PA, Lehner PJ, Sissons JG, Sinclair JH.** 2005. Latency,

chromatin remodeling, and reactivation of human cytomegalovirus in the dendritic cells

of healthy carriers. Proceedings of the National Academy of Sciences of the United States

of America **102:**4140-4145.

21. **Cuevas-Bennett C, Shenk T.** 2008. Dynamic histone H3 acetylation and methylation at

human cytomegalovirus promoters during replication in fibroblasts. J Virol **82:**9525-9536.

22. **Groves IJ, Reeves MB, Sinclair JH.** 2009. Lytic infection of permissive cells with

human cytomegalovirus is regulated by an intrinsic 'pre-immediate-early' repression of

viral gene expression mediated by histone post-translational modification. J Gen Virol

90:2364-2374.

23. **Nitzsche A, Paulus C, Nevels M.** 2008. Temporal dynamics of cytomegalovirus

chromatin assembly in productively infected human cells. J Virol **82:**11167-11180.

24. **Woodhall DL, Groves IJ, Reeves MB, Wilkinson G, Sinclair JH.** 2006. Human Daxx-

mediated repression of human cytomegalovirus gene expression correlates with a

repressive chromatin structure around the major immediate early promoter. J Biol Chem

281:37652-37660.

25. **Zalckvar E, Paulus C, Tillo D, Asbach-Nitzsche A, Lubling Y, Winterling C,**

Strieder N, Mucke K, Goodrum F, Segal E, Nevels M. 2013. Nucleosome maps of the

human cytomegalovirus genome reveal a temporal switch in chromatin organization

linked to a major IE protein. Proc Natl Acad Sci U S A **110:**13126-13131.

26. **Saffert RT, Penkert RR, Kalejta RF.** 2010. Cellular and viral control over the initial

events of human cytomegalovirus experimental latency in CD34+ cells. J Virol **84:**5594-

5604.

27. **Albright ER, Kalejta RF.** 2013. Myeloblastic Cell Lines Mimic Some but Not All

Aspects of Human Cytomegalovirus Experimental Latency Defined in Primary CD34+

Cell Populations. J Virol **87:**9802-9812.

28. **Penkert RR, Kalejta RF.** 2010. Nuclear localization of tegument-delivered pp71 in

human cytomegalovirus-infected cells is facilitated by one or more factors present in

terminally differentiated fibroblasts. J Virol **84:**9853-9863.

29. **Coronel R, Takayama S, Juwono T, Hertel L.** 2015. Dynamics of Human

Cytomegalovirus Infection in CD34+ Hematopoietic Cells and Derived Langerhans-Type

Dendritic Cells. J Virol **89:**5615-5632.

30. **Huang MM, Kew VG, Jestice K, Wills MR, Reeves MB.** 2012. Efficient human

cytomegalovirus reactivation is maturation dependent in the Langerhans dendritic cell

lineage and can be studied using a CD14+ experimental latency model. J Virol **86:**8507-

8515.

31. **Riegler S, Hebart H, Einsele H, Brossart P, Jahn G, Sinzger C.** 2000. Monocyte-

derived dendritic cells are permissive to the complete replicative cycle of human

cytomegalovirus. J Gen Virol **81:**393-399.

32.   **Taylor-Wiedeman J, Sissons P, Sinclair J.** 1994. Induction of endogenous human

cytomegalovirus gene expression after differentiation of monocytes from healthy carriers.

J Virol **68:**1597-1604.

33.   **Reeves M, Sinclair J.** 2013. Epigenetic Regulation of Human Cytomegalovirus Gene

Expression: Impact on Latency and Reactivation, p 330-346. *In* Reddehase MJ (ed),

Cytomegaloviruses: From Molecular Pathogenesis to Intervention, vol 1. Caister

Academic Press, Norfolk, UK.

34.   **Sinclair J, Reeves M.** 2014. The intimate relationship between human cytomegalovirus

and the dendritic cell lineage. Front Microbiol **5:**389.

35.   **Asou H, Suzukawa K, Kita K, Nakase K, Ueda H, Morishita K, Kamada N.** 1996.

Establishment of an undifferentiated leukemia cell line (Kasumi-3) with t(3;7)(q27;q22)

and activation of the EVI1 gene. Jpn J Cancer Res **87:**269-274.

36.   **O'Connor CM, Murphy EA.** 2012. A myeloid progenitor cell line capable of supporting

human cytomegalovirus latency and reactivation, resulting in infectious progeny. J Virol

86:9854-9865.

37.   **Pan C, Zhu D, Wang Y, Li L, Li D, Liu F, Zhang CY, Zen K.** 2016. Human

Cytomegalovirus miR-UL148D Facilitates Latent Viral Infection by Targeting Host Cell

Immediate Early Response Gene 5. PLoS Pathog **12:**e1006007.

38.   **Goodrum F, Jordan CT, Terhune SS, High K, Shenk T.** 2004. Differential outcomes

of human cytomegalovirus infection in primitive hematopoietic cell subpopulations.

Blood **104:**687-695.

39. **Rossetto CC, Tarrant-Elorza M, Pari GS.** 2013. Cis and trans acting factors involved

in human cytomegalovirus experimental and natural latent infection of CD14 (+)

monocytes and CD34 (+) cells. PLoS Pathog **9:**e1003366.

40. **Rauwel B, Jang SM, Cassano M, Kapopoulou A, Barde I, Trono D.** 2015. Release of

human cytomegalovirus from latency by a KAP1/TRIM28 phosphorylation switch. Elife

4.

41. **Zhu D, Pan C, Sheng J, Liang H, Bian Z, Liu Y, Trang P, Wu J, Liu F, Zhang CY,**

Zen K. 2018. Human cytomegalovirus reprogrammes haematopoietic progenitor cells

into immunosuppressive monocytes to achieve latency. Nat Microbiol **3:**503-513.

42. **Shnayder M, Nachshon A, Krishna B, Poole E, Boshkov A, Binyamin A, Maza I,**

Sinclair J, Schwartz M, Stern-Ginossar N. 2018. Defining the Transcriptional

Landscape during Cytomegalovirus Latency with Single-Cell RNA Sequencing. MBio **9**.

43. **Cheung AK, Abendroth A, Cunningham AL, Slobedman B.** 2006. Viral gene

expression during the establishment of human cytomegalovirus latent infection in

myeloid progenitor cells. Blood **108:**3691-3699.

44. **Banchereau J, Steinman RM.** 1998. Dendritic cells and the control of immunity. Nature

392:245-252.

45. **Becher B, Tugues S, Greter M.** 2016. GM-CSF: From Growth Factor to Central

Mediator of Tissue Inflammation. Immunity **45:**963-973.

46. **Lee SH, Albright ER, Lee JH, Jacobs D, Kalejta RF.** 2015. Cellular defense against

latent colonization foiled by human cytomegalovirus UL138 protein. Sci Adv

1:e1501164.

47.   **Goodrum F, Reeves M, Sinclair J, High K, Shenk T.** 2007. Human cytomegalovirus

sequences expressed in latently infected individuals promote a latent infection in vitro.

Blood **110:**937-945.

48.   **Umashankar M, Petrucelli A, Cicchini L, Caposio P, Kreklywich CN, Rak M,**

Bughio F, Goldman DC, Hamlin KL, Nelson JA, Fleming WH, Streblow DN,

Goodrum F. 2011. A novel human cytomegalovirus locus modulates cell type-specific

outcomes of infection. PLoS Pathog **7:**e1002444.

49.   **Goodrum F.** 2016. Human Cytomegalovirus Latency: Approaching the Gordian Knot.

Annu Rev Virol **3:**333-357.

50.   **Qin Q, Penkert RR, Kalejta RF.** 2013. Heterologous viral promoters incorporated into

the human cytomegalovirus genome are silenced during experimental latency. J Virol

87:9886-9894.

51.   **Beisser PS, Laurent L, Virelizier JL, Michelson S.** 2001. Human cytomegalovirus

chemokine receptor gene US28 is transcribed in latently infected THP-1 monocytes. J

Virol **75:**5949-5957.

52.   **Reeves MB, Compton T.** 2011. Inhibition of inflammatory interleukin-6 activity via

extracellular signal-regulated kinase-mitogen-activated protein kinase signaling

antagonizes human cytomegalovirus reactivation from dendritic cells. J Virol **85:**12750-

12758.

53.   **Keyes LR, Hargett D, Soland M, Bego MG, Rossetto CC, Almeida-Porada G, St**

Jeor S. 2012. HCMV protein LUNA is required for viral reactivation from latently

infected primary CD14(+) cells. PLoS One **7:**e52827.

54. **Gatherer D, Seirafian S, Cunningham C, Holton M, Dargan DJ, Baluchova K,**

Hector RD, Galbraith J, Herzyk P, Wilkinson GW, Davison AJ. 2011. High-

resolution human cytomegalovirus transcriptome. Proc Natl Acad Sci U S A **108:**19755-

19760.

55. **Olweus J, Lund-Johansen F, Terstappen LW.** 1995. CD64/Fc gamma RI is a granulo-

monocytic lineage marker on CD34+ hematopoietic progenitor cells. Blood **85:**2402-

2413.

56. **Collin M, McGovern N, Haniffa M.** 2013. Human dendritic cell subsets. Immunology

140:22-30.

57. **Bhatt D, Ghosh S.** 2014. Regulation of the NF-kappaB-Mediated Transcription of

Inflammatory Genes. Front Immunol **5:**71.

58. **Oeckinghaus A, Hayden MS, Ghosh S.** 2011. Crosstalk in NF-kappaB signaling

pathways. Nat Immunol **12:**695-708.

59. **Meier JL, Stinski MF.** 2013. Major Immediate-Early Enhancer and Its Gene Products, p

152-173. *In* Reddehase MJ (ed), Cytomegaloviruses: From Molecular Pathogenesis to

Intervention, vol 1. Caister Academic Press, Norfolk, UK.

60. **Chaturvedi MM, Sung B, Yadav VR, Kannappan R, Aggarwal BB.** 2011. NF-

kappaB addiction and its role in cancer: 'one size does not fit all'. Oncogene **30:**1615-

1630.

61. **Buss H, Dorrie A, Schmitz ML, Hoffmann E, Resch K, Kracht M.** 2004. Constitutive

and interleukin-1-inducible phosphorylation of p65 NF-{kappa}B at serine 536 is

mediated by multiple protein kinases including I{kappa}B kinase (IKK)-{alpha},

IKK{beta}, IKK{epsilon}, TRAF family member-associated (TANK)-binding kinase 1

(TBK1), and an unknown kinase and couples p65 to TATA-binding protein-associated

factor II31-mediated interleukin-8 transcription. J Biol Chem **279:**55633-55643.

62.   **Sakurai H, Chiba H, Miyoshi H, Sugita T, Toriumi W.** 1999. IkappaB kinases

phosphorylate NF-kappaB p65 subunit on serine 536 in the transactivation domain. J Biol

Chem **274:**30353-30356.

63.   **Chen LF, Williams SA, Mu Y, Nakano H, Duerr JM, Buckbinder L, Greene WC.**

2005. NF-kappaB RelA phosphorylation regulates RelA acetylation. Mol Cell Biol

25:7966-7975.

64.   **Choudhary S, Rosenblatt KP, Fang L, Tian B, Wu ZH, Brasier AR.** 2011. High

throughput short interfering RNA (siRNA) screening of the human kinome identifies

novel kinases controlling the canonical nuclear factor-kappaB (NF-kappaB) activation

pathway. J Biol Chem **286:**37187-37195.

65.   **Lukas J, Lukas C, Bartek J.** 2011. More than just a focus: The chromatin response to

DNA damage and its role in genome integrity maintenance. Nat Cell Biol **13:**1161-1169.

66.   **Humby MS, O'Connor CM.** 2015. Human Cytomegalovirus US28 Is Important for

Latent Infection of Hematopoietic Progenitor Cells. J Virol **90:**2959-2970.

67.   **Raja P, Lee JS, Pan D, Pesola JM, Coen DM, Knipe DM.** 2016. A Herpesviral Lytic

Protein Regulates the Structure of Latent Viral Chromatin. MBio **7**.

68.   **Proenca JT, Coleman HM, Connor V, Winton DJ, Efstathiou S.** 2008. A historical

analysis of herpes simplex virus promoter activation in vivo reveals distinct populations

of latently infected neurones. J Gen Virol **89:**2965-2974.

69.   **Proenca JT, Coleman HM, Nicoll MP, Connor V, Preston CM, Arthur J, Efstathiou**

S. 2011. An investigation of herpes simplex virus promoter activity compatible with

latency establishment reveals VP16-independent activation of immediate-early promoters

in sensory neurones. J Gen Virol **92:**2575-2585.

70.   **Proenca JT, Nelson D, Nicoll MP, Connor V, Efstathiou S.** 2016. Analyses of herpes

simplex virus type 1 latency and reactivation at the single cell level using fluorescent

reporter mice. J Gen Virol **97:**767-777.

71.   **Spector DH.** 2013. Exploitation of Host Cell Cycle Regulatory Pathways by HCMV, p

247-263. *In* Reddehase MJ (ed), Cytomegaloviruses: From Molecular Pathogenesis to

Intervention, vol 1. Caister Academic Press, Norfolk, UK.

72.   **Lee SH, Caviness K, Albright ER, Lee JH, Gelbmann CB, Rak M, Goodrum F,**

Kalejta RF. 2016. Long and Short Isoforms of the Human Cytomegalovirus UL138

Protein Silence IE Transcription and Promote Latency. J Virol **90:**9483-9494.

73.   **Petrucelli A, Rak M, Grainger L, Goodrum F.** 2009. Characterization of a novel Golgi

apparatus-localized latency determinant encoded by human cytomegalovirus. J Virol

83:5615-5629.

74.   **Wylie KM, Mihindukulasuriya KA, Zhou Y, Sodergren E, Storch GA, Weinstock**

GM. 2014. Metagenomic analysis of double-stranded DNA viruses in healthy adults.

BMC Biol **12:**71.

75.   **Nitzsche A, Steinhausser C, Mucke K, Paulus C, Nevels M.** 2012. Histone H3 lysine 4

methylation marks postreplicative human cytomegalovirus chromatin. J Virol **86:**9817-

9827.

76.   **Cheng S, Caviness K, Buehler J, Smithey M, Nikolich-Zugich J, Goodrum F.** 2017.

Transcriptome-wide characterization of human cytomegalovirus in natural infection and

experimental latency. Proc Natl Acad Sci U S A **114:**E10586-E10595.

77.   **Thorley-Lawson DA, Hawkins JB, Tracy SI, Shapiro M.** 2013. The pathogenesis of

Epstein-Barr virus persistent infection. Curr Opin Virol **3:**227-232.

78.   **Kew VG, Wills MR, Reeves MB.** 2017. LPS promotes a monocyte phenotype

permissive for human cytomegalovirus immediate-early gene expression upon infection

but not reactivation from latency. Sci Rep **7:**810.

79.   **Kew VG, Yuan J, Meier J, Reeves MB.** 2014. Mitogen and stress activated kinases act

co-operatively with CREB during the induction of human cytomegalovirus immediate-

early gene expression from latency. PLoS Pathog **10:**e1004195.

80.   **Stracker TH, Roig I, Knobel PA, Marjanovic M.** 2013. The ATM signaling network in

development and disease. Front Genet **4:**37.

81.   **Iyengar S, Farnham PJ.** 2011. KAP1 protein: an enigmatic master regulator of the

genome. J Biol Chem **286:**26267-26276.

82.   **Ziv Y, Bielopolski D, Galanty Y, Lukas C, Taya Y, Schultz DC, Lukas J, Bekker-**

Jensen S, Bartek J, Shiloh Y. 2006. Chromatin relaxation in response to DNA double-

strand breaks is modulated by a novel ATM- and KAP-1 dependent pathway. Nat Cell

Biol **8:**870-876.

83.   **Goodarzi AA, Noon AT, Deckbar D, Ziv Y, Shiloh Y, Lobrich M, Jeggo PA.** 2008.

ATM signaling facilitates repair of DNA double-strand breaks associated with

heterochromatin. Mol Cell **31:**167-177.

84.   **Umashankar M, Goodrum F.** 2014. Hematopoietic long-term culture (hLTC) for

human cytomegalovirus latency and reactivation. Methods Mol Biol **1119:**99-112.

85.   **Livak KJ, Schmittgen TD.** 2001. Analysis of relative gene expression data using real-

684   time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods (Duluth) **25:**402-

685   408.

686   86.   **Watson S, Mercier S, Bye C, Wilkinson J, Cunningham AL, Harman AN.** 2007.

687   Determination of suitable housekeeping genes for normalisation of quantitative real time

688   PCR analysis of cells infected with human immunodeficiency virus and herpes viruses.

689   Virol J **4:**130.

690   87.   **Chow S, Minden MD, Hedley DW.** 2006. Constitutive phosphorylation of the S6

691   ribosomal protein via mTOR and ERK signaling in the peripheral blasts of acute

692   leukemia patients. Exp Hematol **34:**1183-1191.

693   88.   **Marvin J, Swaminathan S, Kraker G, Chadburn A, Jacobberger J, Goolsby C.** 2011.

694   Normal bone marrow signal-transduction profiles: a requisite for enhanced detection of

695   signaling dysregulations in AML. Blood **117:**e120-130.

696   89.   **Krutzik PO, Nolan GP.** 2003. Intracellular phospho-protein staining techniques for flow

697   cytometry: monitoring single cell signaling events. Cytometry A **55:**61-70.

 FIGURE LEGENDS

Fig. 1. HCMV latency is established after activation of transcription at 14 days post

infection. (A) Schematic outlining the infection model used for studies of latency and

reactivation in Kasumi-3 cells. GFP+ infected cells were purified by flow cytometry at 1 dpi. On

day 14, latently infected cells were treated with TNF-α for 3 days to induce reactivation. (B)

Representative FACS analysis of GFP expression in Kasumi-3 infected cells at 1 dpi compared

to uninfected cells. (C) Release of viral particles into the media was measured by a $TCID_{50}$ assay

on MRC-5 cells after 2 weeks. (D) UL122 mRNA expression and DNA amount were analyzed

by at the indicated times post infection and expressed relative to D1 after normalization to

GAPDH or RNAseP. (E) RNA/DNA ratios of UL123, UL54, and UL32 over the course of

infection. For Panels C to E, statistical significance was calculated by a one-way ANOVA with a

Dunnett's multiple comparison test (n=4). The error bars represent standard error of the mean

(SEM) and the asterisks indicate p-values (* $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$; **** $P \leq 0.0001$)

calculated by the comparison to the peak (D4 for RNA, D8 for DNA and virus, and D1 for the

RNA/DNA ratio).

Fig. 2. Analysis of GFP- infected cells at 1 and 4 dpi. (A) Schematic outlining the sorting

strategy to purify GFP- cells at D1 and D4 post infection. (B) Representative FACS analysis of

GFP sorting at D1 and D4 post infection. (C) DNA amount, (D) mRNA expression, (E)

RNA/DNA ratio of UL122, UL54, UL32, US28, UL138, and RNA2.7 analyzed at the indicated

times post infection and expressed relative to D1. RNA expression values were normalized to

GAPDH and the DNA amount was normalized to RNAseP. The statistical significance was

calculated by 2-way ANOVA with a Tukey's multiple comparison test (n=3). The error bars

represent the SEM and the asterisks indicate p-values (* $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$;

**** $P \leq 0.0001$) calculated by comparison of the different genes at each time point.

Fig. 3. Expression of latency associated RNAs. mRNA expression (A), DNA amount (B) and

RNA/DNA ratio (C) of UL32, RNA2.7, UL138 and US28 analyzed at the indicated times post

infection and expressed relative to D1. RNA expression values were normalized to GAPDH and

the DNA amount was normalized to RNAseP. The statistical significance was calculated by 2-

way ANOVA with a Tukey's multiple comparison test (n=3). The error bars represent the SEM

and the asterisks indicate p-values (* $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$; **** $P \leq 0.0001$)

calculated by comparison of the different genes at each time point.

732
733
Fig 4. TNF-α induces reactivation independently of differentiation. (A) Latently infected

cells (14 dpi) were treated or not with TNF-α for 3 days. On day 17, DNA and RNA from

infected cells was analyzed for DNA copy number and expression of viral genes as decribed in

Materials and Methods. The release of viral particles in the supernatant was measured by a

$TCID_{50}$ assay. The statistical significance was calculated by unpaired *t-test* (n=4). The error bars

represent the SEM and the asterisks indicate p-values (* $P<0.05$; ** $P< 0.01$). (B) Representative

FACS analysis of uninfected, latently infected cells and TNF-α treated cells for the expression of

the hematopoietic progenitor marker CD34 and the viral immediate early proteins IE1/2.

Reactivating cells were immunophenotyped by gating on the CD34+/IE1/2+ population and

analyzing expression of markers of myeloid differentiation (CD64, CD14, CD15, CD11c and

CD1c). (C) Percentage of undifferentiated cells within the IE1/2 expressing population in the

TNF-α treated cells (CD14-, CD64-, CD15-, CD11c-and CD1c-). The error bars represent the

SEM calculated from 3 experiments.

Fig 5. TNF-α mediated reactivation is correlated with activation of NFκB and ATM

signaling.

Latently infected cells (14 dpi) were treated for 3 days with or without TNF-α 5ng/ml. (A to D)

Flow cytometry analysis were performed to assess the activation of NFκB (p65) and DDR

(ATM, KAP1 and γH2AX). The activation of p65, ATM and KAP1 and are expressed as the

Ratio of Phosphoprotein and total protein compared to untreated cells (p-p65-S536/p65, pATM-

S1981/ATM, and p-Kap-S824/Kap1). γH2AX is expressed as ratio of MFI compared to

untreated cells. The error bars represent the SEM calculated from 3 independent experiments (*

P$\leq$ 0.05; ** P$\leq$ 0.01; *** P$\leq$ 0.001).

TABLE 1. Antibodies used in this study.

Marker	Clone	Supplier
CD34	581	BD Biosciences
CD45	HI30	BD Biosciences
CD1c	F10/21A3	BD Biosciences
CD11c	B-ly6	BD Biosciences
CD14	MOP9	BD Biosciences
CD15	HI98	BD Biosciences
CD64	10.1	BD Biosciences
IE1/2	CH160	Virusys
NF-kB p65	L8F6	Cell Signaling Technology #9460
phospho-NF-kB p65 (Ser536)	93H1	Cell Signaling Technology
ATM	D2E2	Cell Signaling Technology #2873
phospho-ATM (S1981)	D25E5	Cell Signaling Technology #13050
KAP-1	EPR5216	Abcam #109287
phospho-KAP-1 (S824)	EPR5248	Abcam #133440
γ-H2AX (S139)	EP854(2)Y	Abcam #195189

765 TABLE 2. TaqMan assays for analysis of viral DNA and RNA

Assay name	Forward primer sequence (5'-3')	Reverse primer sequence (5'-3')	Reporter sequence (5'-3')
UL122	GGCTCACCTCGTCAATCTTGA	CAACGAGAAGGTACGCAATATCATG	CCCCTTCTGCACACCC
UL123	GCACCCGACAAAACTCACTTAAGA	TGACGCTTGTATGATGACCATGT	ACGGGTCCATCTCTC
UL54	GAAACATAGCCGCCACAGAAC	ACGAATAGTGTTGCCGTGTCA	AACGCCGCTATCATCT
UL32	GGTGAAACGCGGATCTTGAC	GCTCACGGAGACCAGAGG	CTGTCGGAATCCTCG
UL138	CTTCCTCCCAACGGACCAA	CGGACAGACGATACCGTTTCT	ACGGCCCGATGAGATC
US28	CCGATCATCCTCAACGTAGAACT	CGGTAGTAGCAGTAGCTGATGAC	CATGCTCGGTGCTTTC
2.7kb RNA	TCACCGTTTTCTCTCTTCTCTCTCT	TCGAAGAATGAAAGACGACGATGATT	CCGCTCCTGCCACCCG

SUPPLEMENTARY FIGURES

SUPPLEMENTARY FIGURES

Fig. S1. Analysis of viral RNA and DNA over the course of infection in Kasumi-3 cells.

RNAs and DNAs from representative immediate early, early and late genes were analyzed by

RT-qPCR and qPCR, respectively, at various times post-infection as described in Fig. 1 and

Materials and Methods.

Fig S2. Kinetic analysis of HCMV DNA and RNA expression in GFP- cells isolated at 1 dpi.

(A-C) Relative mRNA expression and DNA amount of UL122, UL54 and UL32 analyzed by

RT-qPCR and qPCR, respectively, at the indicated times post infection and expressed relative to

D1. Statistical significance was calculated by a one-way ANOVA with a Dunnett's multiple

comparison test (n=3). The error bars represent the SEM and the asterisks indicate p-values (* $P \leq$

0.05; ** $P \leq 0.01$; *** $P \leq 0.001$; **** $P \leq 0.0001$*) calculated by comparison to the peak at D4.

Fig. S3. TNF induces expression of HCMV early and late genes. RNAs from the experiments

shown in Fig. 6 were analyzed for relative expression of the early gene UL54 and the late gene

UL32.

Fig. S4. Phenotyping of uninfected and latently infected Kasumi-3 cells. Representative

FACS analysis of uninfected (A) and latently infected (B) cells for the expression of the

hematopoietic progenitor marker CD34 and markers of myeloid differentiation (CD64, CD14,

CD15, CD11c and CD1c).

Fig. S5. Analysis of the efficiencies of amplification of viral genes versus RNaseP. Viral

genes and the cellular gene RNaseP were amplified in samples prepared from serial dilutions of

DNA isolated from lytically infected MRC-5 fibroblasts. The ΔCt values ($C_T$viral gene –

C_TRNaseP) for each dilution were calculated and plotted against the log ng DNA.

Fig. S6. Analysis of the efficiencies of amplification of viral RNAs versus GAPDH. Viral

RNAs and cellular GAPDH RNA were amplified in samples prepared from serial dilutions of

cDNA prepared from RNA isolated from lytically infected MRC-5 fibroblasts. The ΔCt values

($C_T$viral gene – C_TGAPDH) for each dilution were calculated and plotted against the log ng

cDNA.

Fig. S7. Validation of GAPDH as a normalization control in HCMV-infected Kasumi-3

cells. Data shows average Ct values ± SD for GAPDH at various times after infection. N=4.

Fig. S8. Antibody staining validation. (A) Representative flow cytometric analysis of HeLa

cells, untreated (red) or treated with hTNF-α (20ng/ml) and Calyculin A (100nM) for 15 mins

(blue), using Phospho-NF-*k*B p65 (Ser563) rabbit mAb and total NF-*k*B p65.

(B and C) Representative flow cytometric analysis of HCT116 treated with 200nM NCS, using

phospho-ATM (S1981), phospho-KAP1 (S824) mAb, ATM and total KAP1 mAb (blue)

compared to untreated control cells (red).

Fig. 1. HCMV latency is established after activation of transcription at 14 days post infection. (A) Schematic outlining the infection model used for studies of latency and reactivation in Kasumi-3 cells. GFP+ infected cells were purified by flow cytometry at 1 dpi. On day 14, latently infected cells were treated with TNF-α for 3 days to induce reactivation. (B) Representative FACS analysis of GFP expression in Kasumi-3 infected cells at 1 dpi compared to uninfected cells. (C) Release of viral particles into the media was measured by a TCID50 assay on MRC-5 cells after 2 weeks. (D) UL122 mRNA expression and DNA amount were analyzed by at the indicated times post infection and expressed relative to D1 after normalization to GAPDH or RNAseP. (E) RNA/DNA ratios of UL123, UL54, and UL32 over the course of infection. For Panels C to E, statistical significance was calculated by a one-way ANOVA with a Dunnett's multiple comparison test (n=4). The error bars represent standard error of the mean (SEM) and the asterisks indicate p-values (* P≤ 0.05; ** P≤ 0.01; *** P≤ 0.001; **** P ≤ 0.0001) calculated by the comparison to the peak (D4 for RNA, D8 for DNA and virus, and D1 for the RNA/DNA

Fig. 2. Analysis of the GFP- infected cells at 1 and 4 days post infection.
(A) Schematic outlining the sorting strategy to purify GFP- cells at D1 and D4 post infection.
(B) Representative FACS analysis of GFP sorting at D1 and D4 post infection. DNA amount (C), mRNA expression (D), and RNA/DNA ratio (E) of UL122, UL54, UL32, US28, UL138, and RNA2.7 analyzed at the indicated times post infection and expressed relative to D1. RNA expression values were normalized to GAPDH and the DNA amount was normalized to RNAseP. The statistical significance was calculated by 2-way ANOVA with a Tukey's multiple comparison test (n=3). The error bars represent the SEM and the asterisks indicate p-values (* P≤ 0.05; ** P≤ 0.01; *** P≤ 0.001; **** P ≤ 0.0001) calculated by comparison of the different genes at each time point.

Fig. 3. Expression of latency associated RNAs. mRNA expression (A), DNA amount (B) and RNA/DNA ratio (C) of UL32, RNA2.7, UL138 and US28 analyzed at the indicated times post infection and expressed relative to D1. RNA expression values were normalized to GAPDH and the DNA amount was normalized to RNAseP. The statistical significance was calculated by 2-way ANOVA with a Tukey's multiple comparison test (n=3). The error bars represent the SEM and the asterisks indicate p-values (* P≤ 0.05; ** P≤ 0.01; *** P≤ 0.001; **** P ≤ 0.0001) calculated by comparison of the different genes at each time point.

Fig 4. TNF-α induces reactivation independently of differentiation. (A) Latently infected cells (14 dpi) were treated or not with TNF-α for 3 days. On day 17, DNA and RNA from infected cells was analyzed for DNA copy number and expression of viral genes as decribed in Materials and Methods. The release of viral particles in the supernatant was measured by a TCID50 assay. The statistical significance was calculated by unpaired t-test (n=4). The error bars represent the SEM and the asterisks indicate p-values (* P< 0.05; ** P< 0.01). (B) Representative FACS analysis of uninfected, latently infected cells and TNF-α treated cells for the expression of the hematopoietic progenitor marker CD34 and the viral immediate early proteins IE1/2. Reactivating cells were immunophenotyped by gating on the CD34+/IE1/2+ population and analyzing expression of markers of myeloid differentiation (CD64, CD14, CD15, CD11c and CD1c). (C) Percentage of undifferentiated cells within the IE1/2 expressing population in the TNF-α treated cells (CD14-, CD64-, CD15-, CD11c-and CD1c-). The error bars represent the SEM calculated from 3 experiments.

Fig 5. TNF-α mediated reactivation is correlated with activation of NFκB and ATM signaling.
Latently infected cells (14 dpi) were treated for 3 days with or without TNF-α 5ng/ml. (A to D) Flow
cytometry analysis were performed to assess the activation of NFκB (p65) and DDR (ATM, KAP1 and
γH2AX). The activation of p65, ATM and KAP1 and are expressed as the Ratio of Phosphoprotein and
total protein compared to untreated cells (p-p65-S536/p65, pATM-S1981/ATM, and
p-Kap-S824/Kap1). γH2AX is expressed as ratio of MFI compared to untreated cells. The error bars
represent the SEM calculated from 3 independent experiments (* P≤ 0.05; ** P≤ 0.01; *** P≤ 0.001).