A comprehensive, mechanistically detailed, and executable model of the Cell Division Cycle in *Saccharomyces cerevisiae*

Ulrike Münzner[1,2], Edda Klipp[1] and Marcus Krantz[1]

1. Institute of Biology, Humboldt-Universität zu Berlin, Berlin, Germany

2. Bioinformatics Center, Institute for Chemical Research, Kyoto University, Uji, Japan

marcus.krantz@rxncon.org

ABSTRACT

Understanding how cellular functions emerge from the underlying molecular mechanisms is one of the principal challenges in biology. It is widely recognised that this will require computational methods, and we expect the predictive power of these models to increase as model coverage and precision of formulation increase. The most potent examples to date are the genome-scale metabolic models that revolutionised their field and paved the way for a bacterial whole-cell model. However, to realise the potential of whole-cell models for eukaryotes, we must enable genome-scale modelling of other cellular networks. This has proven particularly challenging for the cellular regulatory networks, i.e. signal transduction, gene regulation and cell cycle control. Here, we present a comprehensive mechanistic model of the molecular network that controls the cell division cycle in *Saccharomyces cerevisiae*. We use rxncon, the reaction-contingency language, to neutralise the scalability issues associated with regulatory networks, and to enable model formulation, visualisation and simulation at the genome-scale. In particular, we use a parameter-free simulation logic to validate the network and to predict genotype-to-phenotype relationships down to residue resolution. The genome-scale offers a new perspective on cell division cycle control: In contrast to conventional understanding, we find that the cell division cycle control network consists of disjunct control modules rather than a cycle, which synchronise three independent replication cycles. Taken together, we demonstrate that it is possible to build, visualise and simulate mechanistic models of cellular regulatory networks at the genome-scale, and that these models provide powerful tools to reconcile molecular biology and physiology - and to predict phenotypic effects of mutations down to the level of residue substitutions. We envisage these results as a key step towards eukaryotic whole-cell models and, in the long run, personalised medicine.

INTRODUCTION

Computational models provide powerful tools to study biological systems (Kitano 2002). In particular, mechanistic models that explain cellular functions and phenotypes from molecular events are powerful tools to assemble knowledge into understanding. When built carefully, they combine three functions: As integrated and internally consistent knowledge bases, as scaffolds for integration, analysis and interpretation of data, and as executable models. The value of these models arguably increases the more mechanistically detailed and comprehensive they are, culminating in the genome-scale mechanistic models of metabolism and the whole-cell model of *Mycoplasma genitalium* (Herrgard, Swainston et al. 2008, Karr, Sanghvi et al. 2012, Thiele, Swainston et al. 2013). These models can be used to explain and predict perturbation responses and genotype-to-phenotype relationships, and whole cell models have the potential to revolutionise biology, biotechnology and medicine. However, to realise this potential, we must be able to scale up modelling of all cellular processes.

This has proven especially challenging for the cellular networks that process information (reviewed in (Hlavacek and Faeder 2009, Münzner, Lubitz et al. 2017)). These networks encode information primarily through reversible state changes in their components, such as bonds or covalent modifications. Typically, a component can interact with multiple partners and be modified at multiple residues, and most of these bonds and/or modifications are not mutually exclusive. Hence, there is a one-to-many relationship between empirical observables (*elemental states*), such as a specific bond or the modification status at a specific residue, and the possible configurations of the protein (*microstates*). This leads to problems in most classical modelling formalism, where the resolution difference leads either to a loss of mechanistic detail (e.g. component level modelling) or to the combinatorial complexity (microstate modelling). The solution is to use a formalism with adaptive resolution, such as rule-based modelling languages (RBMLs (Blinov, Faeder et al. 2004, Danos, Feret et al. 2007)), the Entity Relationship diagrams of the Systems Biology Graphical Notation (SBGN-ER (Le Novere, Hucka et al. 2009)) or rxncon, the reaction-contingency language (Tiger, Krause et al. 2012) (reviewed in (Rother, Munzner et al. 2013, Chylek, Harris et al. 2015)). However, the potential of these formalisms has yet to be realised in a comprehensive mechanistic model of a eukaryotic signalling system.

The eukaryotic cell division cycle (CDC) may be the most interesting - both medically and philosophically - of these systems, as it constitutes the very core of life as we know it. The CDC is best understood in baker's yeast, *Saccharomyces cerevisiae*, where the rise and fall in activity of a single cyclin dependent kinase (CDK) suffices to drive the replication of DNA, nuclear division (including duplication and separation of the spindle pole body (SPB, the yeast centrosome)) and cell morphology and division (Enserink and Kolodner 2010, Howell and Lew 2012). The molecular basis of cell cycle control has been studied since the 1970-ies, when the first *cdc* mutants were identified (Hartwell, Culotti et al. 1970). It has also been a popular target of computational analysis, and has been studied with models ranging in complexity from conceptual oscillators via logical networks to extensive ODE systems that can predict most of the known mutant phenotypes (Goldbeter 1991, Chen, Csikasz-Nagy et al. 2000, Chasapi, Wachowicz et al. 2015, Kraikivski, Chen et al. 2015). However, even the largest of these models are limited in scope and cannot easily be extended to the genome scale. On the other hand, Kaizu and colleagues compiled the state-of-the-art knowledge in an exquisitely detailed map of the molecular network and its connections to a set of cellular signalling pathways (Kaizu, Ghosh et al. 2010), using the process description diagram language (SBGN-PD; (Le Novere, Hucka et al. 2009)). However, these maps are not executable and cannot be used for simulation. To realise the potential of a mechanistic

genome-scale model, we need to combine the features of all three efforts: The mechanistic precision in the molecular biology, the scope of the comprehensive maps, and the executability of the mathematical models.

Here, we present a mechanistic, executable and genome-scale model based on the network that controls and executes the cell division cycle in baker's yeast, *Saccharomyces cerevisiae*. We chose to build this model using rxncon, as it (similarly to RBMLs) has the adaptive resolution required to reconcile the necessary scalability and precision, and as rxncon networks (in contrast to rule-based models (RBMs)) can be compiled into and simulated as bipartite Boolean models (Romers, Thieme et al. 2017). This scalable, parameter-free simulation logic makes it possible to qualitatively simulate the cell cycle network, which is too large and/or has too many unknown parameter values (or truth tables) to be simulated with classical methods.

Based on an in-depth literature review, we compiled a mechanistically detailed model including 357 components and 790 elemental reactions of which 598 have regulatory contingencies. Despite the coverage, we could not verify that the regulatory network forms a closed cycle. Instead, it appears to consist of three disjunct control modules gating the G_1/S, G_2/M and M/G_1 transitions, which control and respond to the progression of three replication and division cycles: DNA replication, nuclear division and cell division (including bud emergence, growth, morphogenesis and cytokinesis). We added three coarse-grained "macroscopic cycles" and an interface to the control network, to close the regulatory gap through a hybrid model. The corresponding parameter-free model reproduces both starvation induced G_0 arrest and ordered progression through the three division cycles in the presence of nutrients, as well as arrest in the presence of pheromone (G_1-phase), hydroxyurea (HU, S-phase), latrunculin-A (LatA, G_2/M-phase) and nocodazole (NOC, G_2/M-phase). To test this model, we analysed a set of 85 mutants, including residue point mutations, and accurately predicted the viability or lethality of 72%. Finally, we visualised the complete regulatory structure in a rxncon regulatory graph, providing a poster-sized overview of the current mechanistic knowledge without any simplifying assumptions. Taken together, we show that it is possible to compile, visualise and analyse comprehensive mechanistic models of eukaryotic signal transduction, and that a qualitative network definition enables systems level predictions on genotype-to-phenotype relationships, down to residue resolution, without parametrisation or model optimisation. We envisage this as a stepping stone towards eukaryotic whole-cell models, and towards truly personalised medicine.

METHODS

The cell division cycle model was created using the second generation rxncon language (Romers and Krantz 2017), as described in the supplementary materials and elsewhere (Romers, Thieme et al. 2018). The rxncon model consists of two types of statements that both correspond directly to empirical data. First, *elemental reactions* define decontextualised reaction events in terms of changes in elemental states. An elemental reaction is essentially a reaction centre in a RBM (Faeder, Blinov et al. 2009). Second, *contingencies* define constraints on the elemental reactions in terms of (Boolean combinations of) elemental states. The complete set of contingencies for a single elemental reaction define the reaction context(s) in a RBM, but in rxncon the context definition is separated over several contingencies that each define the impact of a single elemental state (Romers and Krantz 2017). This gives a one-to-one correspondence between model definition and empirical data, improving composability as model changes and extension have a local impact only. Multiple contingencies can be combined into statements that are arbitrarily complex, defining reactions down to microstate resolution when necessary. Typically, however, single or few contingencies per elemental reactions suffice to capture all the empirically known regulatory mechanisms. In the end, the rxncon model constitutes a molecular biology knowledge base in formal language, which is computer readable, can be automatically visualised and compiled into a uniquely defined executable model.

Parameter-free models were created automatically using the rxncon compiler (Romers, Thieme et al. 2017), and simulated with the R package BoolNet (Mussel, Hopfensitz et al. 2010). The rxncon software can be downloaded from https://github.com/rxncon/rxncon or directly installed from the python package index ("pip install rxncon"). See https://rxncon.org for further instructions.

The cell division cycle model was visualised as rxncon regulatory graphs (Tiger, Krause et al. 2012), using Cytoscape (http://www.cytoscape.org/) and a visual formatting file (rxncon2cytoscape.xml; https://github.com/rxncon/tools).

The cell division cycle model is compiled in an SBtab compatible spreadsheet (Lubitz, Hahn et al. 2016), and is available from https://github.com/rxncon/models (CDC_S_cerevisiae.xls). The model is fully referenced through the "reference" columns in the reaction and contingency sheet, and explained in detail in the supplementary material.

The model is based on the literature listed in Table S3, as specified for individual reactions and contingencies in Table S1.

RESULTS

A comprehensive, mechanistic and executable model of a eukaryotic cell division cycle

We present a comprehensive, mechanistically detailed and executable model of the cell division cycle in baker's yeast (Figure 1A, Table S1, Figure S1 (printable A0)). The model includes regulated expression and degradation of proteins, assembly and regulation of the cyclin dependent protein kinases Cdc28 (Cdk1) and Pho85 (Cdk5), and the regulation of DNA replication, SPB duplication and nuclear division, and cell polarity and morphogenesis. Each of these processes is defined – as far as empirical knowledge allows – down to the role of specific modifications and bonds at particular residues and domains. The model accounts for 357 components. This number includes 229 individual proteins, and the genes and mRNAs of the 44 proteins for which we consider regulated expression and/or degradation. In addition, the model includes 7 multimeric protein complexes (e.g. APC/C), 3 chromosomal features (e.g. the origins of replication), 29 kinase and phosphatase activities as target specific enzymes that remain to be mapped on one or more gene products (e.g. Sfi1PPT as an unidentified phosphatase (PPT) of Sfi1), and one small molecule: phosphatidylinositol (PI). These components take part in 790 elemental reactions that produce and consume 1238 elemental states, and that are regulated by 598 contingencies - several of which correspond to Boolean combinations of elemental states.

In addition to the molecular reaction network (MRN), we encode a coarse-grained model (CGM) of DNA replication, SPB duplication and budding in 12 macroscopic reactions and 12 macroscopic states (Figure 1B). The macroscopic reactions respond to a series of inputs that correspond to outputs (observables) of the MRN. Conversely, the CGM provides outputs that act as inputs to the MRN. The entire regulatory logic is encoded in a single hybrid rxncon model, encompassing 802 reactions and 972 lines of contingencies. The contingencies connect the MRN and CGM to each other, and to the five external inputs we consider: Nutrients, Pheromone, HU, LatA and NOC. The biology and implementation is described and visualised in detail in the supplementary documentation. The model accounts for all components to which we could assign a mechanistic function in the control of cell cycle division, and hence it constitutes a first draft of a genome-scale mechanistic model of eukaryotic cell division.

Building a mechanistic knowledge base at the genome scale

The rxncon model constitutes a biological knowledge base rather than a mathematical model. We used an iterative workflow with information retrieval and interpretation, data formalisation, visual validation, computational validation and finally model analysis (Romers, Thieme et al. 2018). Through in-depth literature review, we extracted and evaluated empirical information and formalised it as reactions and contingencies using statements such as: "Cdc28 phosphorylates Sld2 at Threonine 84" (an *elemental reaction*) and "Cdc28 phosphorylates Sld2 at Threonine 84, only if Cdc28 is bound to Clb5 (or Clb6)" (a *contingency*). This type of mechanistic model puts high demands on data coverage and quality, which cannot always be met. Hence, the model contains gap-filling assumptions that are indicated in the model annotation (Table S1). In addition to these specific assumptions, we also made the general assumptions that reversible states that are produced are reversed (e.g. all phosphorylated residues are dephosphorylated) and that synthesised components (proteins, mRNAs) are degraded. This leads us to introduce a set of 28 undefined phosphatase activities to compensate for the fact that these reaction types are understudied (Tiger, Krause et al. 2012).

We make a few exceptions from these general assumptions when proteins are highly stable (e.g. Mcm2-Mcm7) or when modifications are reversed by degradation (e.g. Sic1 phosphorylation). Furthermore, several proteins in the network are regulated at the level of localisation, which cannot be properly expressed in rxncon. We work around this limitation by mapping the effect directly on the elemental states that are responsible for the localisation changes. For example, the transcription factors Ace2 and Swi5 are regulated at the level of nuclear localisation, which in turn is regulated by phosphorylation in the nuclear localisation signal (NLS). In the model, we make promoter recruitment directly dependent on NLS phosphorylation, bypassing the spatial description without compromising the regulatory logic. However, for most of the network we find biochemical (for reactions) or combinations of biochemical and genetic (for contingencies) data that can be formalised in the rxncon language using 14 elemental reaction types (Table S1), creating a mechanistically detailed knowledge base of the network that controls and executes the cell division cycle.

Insights at the genome scale: Three independent replication programs

The mechanistic genome-scale model presents a new perspective on the cell division cycle. The model summarises the current mechanistic knowledge on information transfer and hence connections in the model represent direct and functional connections *in vivo*. Consequently, the modules that emerge are not artefacts of model assumptions, but reflect known biological function. Conversely, the lack of connection implies that no (known) direct and/or functional connections exist. In particular, there is very little interaction between the three duplication cycles that execute DNA replication, SPB duplication and nuclear division (ND), and budding, morphology and cytokinesis (CD). These processes do not directly interact at the molecular level outside mitosis. Instead, these processes are monitored and synchronised by the regulatory network. While the lack of direct mechanistic connections between the cycles may reflect missing knowledge, the current model predicts that the cell division cycle consists of three distinct programs that can be executed independently: DNA replication, nuclear division (including SPB duplication) and cell division. With the appropriate adaptation of (the state of) the regulatory network, it should be possible to execute these programs separately and hence uncouple DNA replication, nuclear division and cell division. Such uncoupling could generate ploidy shifts, multinucleate cells, or cells without nuclei; which have all been observed in eukaryotic cells. The perhaps most prominent example of the modularity of these processes is meiosis, where a single DNA replication (pre-meiotic S-phase) is followed by two rounds of nuclear division (meiosis I + II) without any cell division (Neiman 2011). Hence, the CDC control network gates the cell division cycle in response to cellular and/or environmental status to synchronise three independent cycles; DNA replication, nuclear division and cell division, during a normal cell division cycle.

Mechanistic modularity: The CDC control network is not a cycle but three major checkpoints

We made a number of striking observations in the model building process. While we only used previously published data, the holistic picture of the assembled knowledge highlighted features that were not obvious from the modules themselves. In particular, the regulatory network controlling the cell division cycle does not in itself form a cycle. Instead, the regulatory network falls apart into three modules, corresponding to three major checkpoints that gate the G_1/S, the G_2/M and the M/G_1 transitions (we found no evidence for a regulatory S/G_2 boundary). Judged on the available data, these three subnetworks only interact indirectly

through the progression of the DNA, ND, and CD cycles (Figure 1C). Arguably, this makes perfect sense, as the control network must be responsive to the state of the cell division cycle. At the same time, this is not widely recognised in the literature, and several current models of the cell division cycle directly link the three regulatory modules into a clock-like network. Our findings here suggest that this is incorrect.

Merging perspectives: A role for the DNA damage response in normal CDC control

The model also connects knowledge from several distinct areas of research. When we combined knowledge from the cell cycle and DNA damage research fields, a clear picture of how DNA replication is monitored and how this prevents mitotic entry emerged. Ongoing DNA replication leads to ssDNA/Rad53 signalling from replication forks (Figure 2). This signal maintains S-phase transcription (through inhibition of the Nrm1 repressor), and inhibits the G2/M transition through transcriptional inhibition of the *CLB2* cluster (through inhibition of the Ndd1 activator and through post-translational inhibition of Cdc28-Clb2 through stabilisation of the inhibitory CDK tyrosine kinase Swe1. While the S-phase checkpoint is well known from studies on DNA damage and repair, their function are again not (widely) recognised in the literature on the cell division cycle control. However, the essential function of Mec1/Rad53 signalling in normal cell cycle control suggests that this may be its primary function.

Executing the logic: Ordered progression of the cell division cycle and nutrient dependent G_0 arrest

We analysed the cell division cycle model through parameter-free simulation. The rxncon network is not directly executable, but compilable into a uniquely defined bipartite Boolean model (bBM; (Romers, Thieme et al. 2017)). We used the bBM to evaluate the completeness of the cell division cycle model in three steps: First, we searched for a point attractor in the absence of nutrients, reflecting G_0 arrest. Second, starting from this G_0 attractor, we released the cell cycle arrest through addition of nutrients and searched for a cyclic attractor with ordered progression through the three macroscopic cycles. Third, we interrupted the cyclic attractor by adding compounds known to halt cell cycle, i.e. Pheromones, HU, LatA or NOC. The first step was to find an appropriate initial state vector for the model. The model has 1403 nodes and hence 2^{1403} ($\sim 10^{422}$) possible state vectors, precluding an exhaustive search. Instead, we used the rxncon default vector where all neutral elemental states and state-less components are True and all other states, all reactions and inputs are False (see (Romers, Thieme et al. 2017)), and released it in the absence of nutrients (setting the placeholder [Histones] input to True). Interestingly, the resulting point attractor is consistent with nutrient dependent G_0 arrest (Fig 3A; Fig S2). Next, we simulated the network from this initial state in the presence of nutrients, and evaluated the simulation trajectory and attractor to identify inconsistencies and gaps in the knowledge base. Most importantly, the crude time concept in the Boolean model caused significant problems, as shorter event chains are faster even if they occur through slower reactions. To resolve this problem, we introduced a time-scale separation that made all transcriptional reactions and DNA replication slower than other reactions (such as post-translational modifications or protein-protein interactions), by requiring their input to be true for twenty consecutive time steps before firing (reflecting the longest signalling path). After introducing the time scale separation, the model accurately reflects the ordered progression through the three macroscopic cycles (Fig 3B; Fig S3), resulting in a cyclic attractor with

period 186. Finally, we examined the arrest point in response to Pheromone, HU, LatA and NOC (Fig 3C; Fig S4), identifying and correcting one final issue in the network: The ability of Swe1 to inhibit mitotic entry in the absence of a proper bud. The changes made in the network validation phase are summarised in Table 1.

Testing the model: Parameter-free simulation accurately predicts genotype-to-phenotype relationship for 63 out of 85 tested mutants

To examine the predictive power of the model, we analysed the predicted arrest point of *cdc* mutants with known phenotypes. We chose 85 (combinations of) deletions, point, and constitutive (over) expression mutants, and examined their cell cycle progression (cyclic attractor) or arrest (point attractor) (Table S2). We observe two types of cyclic attractors; a normal and ordered progression through all three macroscopic cycles (viability) and a partial cyclic attractor passing through DNA replication and nuclear division, but not cell division (resulting in multinucleate cells, scored as lethality). In addition, we observe twelve distinct point attractors (at the level of macroscopic states), which correspond to G_1, S, G_2/M, M, and T arrest (Fig 4). The model correctly predicts all 43 lethal mutants, and 19 out of 42 viable mutants, but also scores 23 viable mutants as inviable. We note that the model is conservative in estimating viability, and look closer at the 23 inconsistent predictions (Table 2). Several of these mutants can be explained by implicit synthetic lethality[1]. I.e., the mutations are synthetically lethal *in vivo* with components for which key mechanisms are not known and hence not included in the model. E.g., almost no mechanistic data exist on the function of Clb3 and Clb4, which can partially compensate for loss of Clb5, explaining why *clb5* mutants are predicted to be lethal in the model (they correspond to the lethal *clb3clb4clb5* triple mutants (Schwob and Nasmyth 1993)). Similarly, no strictly required mechanism for Bck2 is implemented, explaining the lethality of the *cln3* (*bck2cln3* (Epstein and Cross 1994)) mutant. Seven of the 23 inconsistent predictions can be explained through implicit synthetic lethality. Some of the remaining mutants can be explained by strong phenotypes, which could be justified as "lethal" on a binary scale, such as the SBF-mutants (*swi4*, *swi6*, *swi6S4A* and *swi4swi6*) and *pds1*. Others, like *mbp1* and *cdh1*, which are predicted to be lethal but have no or weak reported phenotypes, likely indicate missing redundancy in the model. Taken together, the biological knowledge base uniquely defines a parameter-free model, which accurately predicts the vast majority of the mutant phenotypes that we tested - including point mutants.

---

[1] Synthetic lethality: The combination of two or more mutations that are individually viable results in lethality.

DISCUSSION

We present a genome-scale, mechanistic model of the cell division cycle. It is mechanistic, as the connections within the model correspond to direct biochemical reactions or dependencies of these reactions on the state of the reactants, down to the level of specific residues and domains. It is genome-scale, as it accounts for all components in the cell division cycle for which we could assign a mechanistic function. We visualise the complete knowledge base as a single, connected network, and show that it defines an executable model. We use this model for validation of the network and to predict genotype-to-phenotype relationships down to the resolution of specific residues.

The genome-scale scope allows us to interpret missing features. The most striking observation is the lack of a single cycle: The control network falls apart into three distinct control circuits; G_1/S, G_2/M and M/G_1, which monitor and control three distinct replication cycles: DNA replication, nuclear division and cell division. While we cannot rule out as yet unknown connections between the three control modules and between the three replication cycles, the modularity we observe seems to make perfect sense: The control network needs to be more than a sizer/timer; it must also depend on the replication cycles to be able to control these processes. On the other hand, the three replication cycles need to be distinct in order to explain ploidy shifts, multi-nucleate (or nuclei-free) cells and meiosis. Nevertheless, the regulatory mechanisms that uncouple these cycles remain largely unknown even in baker's yeast.

These findings have implications for how we model the CDC. We and others have previously modelled the regulatory network as a closed cycle, where CDC progression may be nutrient dependent, but not explicitly dependent on the progression of the internal division events. In particular, it has been difficult to find a direct link from the G_1/S transition (Cln1/2, Clb5/6 expression) to mitotic entry (Clb1/2 expression). We and others have modelled this as a gene expression cascade (Clb5/6 -> Clb3/4 -> Clb1/2), but without being able to explain how this cascade responds to e.g. HU-induced S-phase arrest (Linke, Chasapi et al. 2017, Spiesser, Uschner et al. Submitted) . Here, we can explain this by linking DNA damage signalling to cell cycle progression, through inhibition of Clb1/2 at two distinct levels: Transcriptionally through inhibition of Ndd1, and post-translationally through the stabilisation of Swe1. This knowledge was available in the literature, as was the knowledge that *mec1* and *rad53* mutants are impaired in normal CDC-progression, but has previously not been brought together in cell cycle models. Based on the observations we make here, it seems that the interplay between the regulatory network and division events are a necessary component of a mechanistic cell cycle model.

Despite its scope, the model we present here comes with several limitations. First of all, it is primarily a biological knowledge base. We have focussed on the qualitative information, i.e., which state variables change, which reactions are responsible for these changes, and in which molecular context these reactions occur, as this information is more abundant and less uncertain - as it is for metabolic networks (Herrgard, Swainston et al. 2008). Second, the model is limited by literature data coverage and quality. While the CDC of *S. cerevisiae* is exceptionally well known, we nevertheless observe uncertainty and gaps in this knowledge. Here, we make a minimal set of assumptions to make the model functional, and consider these gap-filling additions hypotheses for future experimental validation. However, the mutant analysis also suggests that we are missing redundancies in the network, or that some effects cannot be properly described at the qualitative level only. The latter is particularly true for the macroscopic reactions we introduce to cover DNA replication, spindle pole duplication and

cell growth and cytokinesis. The mono- and bimolecular formulation (in the context of arbitrarily large complexes) is a useful abstraction level to describe signal transduction, but not to describe large cellular structures, polymers or quantitative properties of molecule pools (such as polarity establishment). The macroscopic placeholder reactions of the CGM must be turned into models of their own before we reach a quantitative predictive model of the cell division cycle. Third, the model does not explicitly include spatial aspects, as complex level properties (such as localisation) cannot currently be expressed in the rxncon language. Instead, we directly connect the states that regulate localisation to the reactions that depend on it. Fourth and finally, the Boolean modelling logic is a very crude approximation both quantitatively and temporally, and, indeed, we needed to introduce time-scale separation between transcription and the other signalling events. However, a quantitative simulation approach would require parameters, creating a massively underdetermined parameter-estimation problem. All things considered, we believe the parameter-free simulation - which reflects the qualitative model only - may be more informative, and, indeed, find that the vast majority of the mutant phenotypes (which were not used to validate or improve the model) can be reproduced even by this qualitative simulation method. Interestingly, the discrepancies between model predictions and known phenotypes seem to primarily identify missing redundancies, and hence missing empirical knowledge. Taken together, we present a first draft model, to be refined and extended by the community, which highlights the need for further dedicated work on the experimental characterisation of this fundamental model system.

This work is complementary to quantitative modelling efforts. The qualitative simulation we use here can indeed give a readout for 1403 variables - elemental reactions and states - but the values of these variables are limited to True or False. In addition, we project all this on a binary phenotype: Dead or Alive. In contrast, quantitative models can help us precisely predict quantitative or complex phenotypes, such as cell cycle duration, cell size distribution, and polarity establishment (Chen, Csikasz-Nagy et al. 2000, Giese, Eigel et al. 2015, Kraikivski, Chen et al. 2015). However, the model we present here is much larger in scope than these quantitative models. Until today, only the Kaizu map is similar in scope: This network contains 318 proteins, including 96 signalling components, leaving a cell cycle module of 230 proteins - of which 148 are represented in the model we present here (individually or in one of the protein complexes) (Kaizu, Ghosh et al. 2010). However, the Kaizu map is not executable and can hence be neither computationally validated, nor used for genotype-to-phenotype predictions. In the future, we may be able to reconcile the two: Scope and quantitative modelling. The model here could in principle be converted into a quantitative model, as a rxncon model is semantically defined in terms of a rule-based model in the BioNetGen language (Romers, Thieme et al. 2018). However, to simulate this network as a quantitative model, we need to overcome a number of hurdles: First, there is no clear equivalence to rxncon inputs in a rule based model, so the connections between modules must be hand crafted. Second, we would need to find or estimate values for >790 unique parameters. Third, we would need representations of the DNA replication, nuclear division, and cell division cycles that are compatible with agent based simulation. In the foreseeable future, it seems likely that meaningful simulation of signal transduction networks at this scale will remain qualitative or semi-quantitative, as it does for metabolic networks.

However, we show that we can build and visualise models of information processing networks that are mechanistically detailed, large scale and executable. Until today, this has only been done for metabolic (or mass-transfer) networks, and the mass-transfer logic leads to crippling scalability issues when applied to signal transduction networks (due to microstate enumeration; reviewed in: (Hlavacek and Faeder 2009, Chylek, Harris et al. 2015, Münzner, Lubitz et al. 2017)). This technical issue can be solved by using a formalism with adaptive

resolution, such as a rule-based modelling language (see e.g.:(Blinov, Faeder et al. 2004, Danos, Feret et al. 2007, Chylek, Harris et al. 2015)), SBGN-ER (Le Novere, Hucka et al. 2009) or rxncon (Tiger, Krause et al. 2012). We settled for rxncon, as it is text based rather than graphical, uses a higher-level biological language rather than model code, supports scalable visualisation, and is compilable into a parameter-free model for direct simulation (Romers and Krantz 2017, Romers, Thieme et al. 2017). In particular, we find the iterative visualisation in the regulatory graph invaluable for the model construction process, and we present the final graph in a wall-chart that is inspired by the biochemical pathway maps (Michal 2014). In addition, the ability to go from a pure biological knowledge base to a parameter-free model that can predict system level features from molecular mechanisms is highly non-trivial (Hlavacek and Faeder 2009). Here, we use this feature for network validation - by iteratively analysing and improving the network until the corresponding model reproduces wild-type behaviour - and for genotype-to-phenotype predictions of 85 mutants. Hence, we bring together the mechanistic precision in the molecular biology, the scope of the comprehensive maps, and the executability of the mathematical models.

Taken together, we show that it is possible to build, visualise and simulate mechanistic genome-scale models of eukaryotic signal transduction networks. Here, we do this for the cell cycle network of baker's yeast, and, although the rxncon language must be extended to deal with e.g. alternative splicing and localisation, there are no fundamental differences to higher eukaryotes. A similar high-resolution knowledge base on the information processing network in human cells would be an important step towards a human whole-cell model and, when it accounts for the molecular effect of allele differences and drug perturbations, truly personalised medicine.

FIGURE LEGENDS

Figure 1: The molecular architecture of the CDC control network. (A) The model consists of seven parts: (i) Gene expression, (ii) regulated degradation, (iii) CDK assembly and activation, (iv) DNA replication, (v) SPB duplication and nuclear division, (vi) cell division, and (vii) a set of reactions involving cell cycle components but without (known) regulatory connections to the main network. The outputs of the mechanistic reaction network (MRN) in modules (iv)-(vi) determine the progression of the coarse-grained model (CGM). **(B)** The CGM consists of three parts: (left) SPB duplication and nuclear division, including satellite formation (SPB: neutral (0) -> satellite), duplication plaque formation (SPB: satellite -> duplication plaque), SPB duplication (SPB: duplication plaque -> SPB duplicated), SPB separation (SPB: duplicated -> bipolar), and nuclear division (SPB: bipolar -> neutral). (middle) DNA replication, including licencing (DNA: neutral (0) -> licensed), initiation (DNA: licensed -> replicating), replication (DNA: replicating -> replicated) and separation (DNA: replicated -> neutral). (right) Cell division, including: Bud emergence (Bud: none (0) -> small), bud growth (Bud: small -> large), and cytokinesis (Bud: large -> 0). The MRN and CGM parts communicate via an input/output layer. (C) Schematic representation of the hybrid model architecture. The three independent replication cycles are regulated by three disjunct control modules gating the G_1/S, G_2/M and M/G_1 transitions. The network is responsive to different forms of stress and other external cues, but here we only consider nutrients and four classical CDC arrest drugs: Pheromone, HU, LatA and NOC.

Figure 2: The Mec1/Rad53 pathway senses ongoing DNA replication to inhibit mitotic entry. Ongoing DNA replication leads to exposed ssDNA and ssDNA/dsDNA junctions at the replication forks, which are sensed by the Mec1/Rad53 pathway. Mec1 phosphorylates and stabilises Swe1, a Cdk1-Clb1/2 inhibitor, and Rad53 phosphorylates and inhibits Nrm1, which is needed for *CLB1/2* transcription. The regulatory graph is a bipartite representation where elemental reactions (red) produce or consume elemental states (blue), and elemental states influence - through contingencies - elemental reactions. Complex contingencies are collected in Boolean nodes (AND/OR/NOT; light grey), and Input/Output nodes (dark grey) define the interface between the model and its surroundings - and between the MRN and CGM parts.

Figure 3: The parameter-free model reproduces both cell cycle progression and arrest. Simulation results visualised on the CGM states only. (A) When the model is simulated with nutrients set to False, the cells arrest at G_0: With licensed DNA, SPB-satellite and no bud. (B) When cells are released from the G_0 arrest by setting nutrients to True, they progress through the cell division cycle by replicating their DNA, undergoing nuclear division and rapidly thereafter cell division, and resetting the network for a new division, traversing 186 Boolean steps. (C) The cells arrest again when exposed to pheromone (G_1: licensed DNA, SPB-satellite, no bud), HU (S: replicating DNA, duplicated SPB, large bud), LatA (G_2/M: replicated DNA, duplicated SPB, small bud) and NOC (G_2/M: replicated DNA, duplicated SPB, large bud).

Figure 4: The model predicts the phenotypes of a majority of all tested mutants. Simulations were performed by introducing deletions (setting all protein states (and mRNA and Gene, if appropriate) to False), point mutations (by locking proteins in a certain

modification state) or constitutive overexpression (by setting the transcription reaction to True constantly), and the mutants were analysed through simulation by release from the (modified) G_0 state of the wild-type. (A) The model predicts the correct phenotype (dead or alive) for 62 out of 85 mutants, although closer inspection revealed that several of the 23 inconsistent mutants can be explained by implicit synthetic lethality or strong phenotypes (Table 2). (B) The 62 mutants that are predicted to be dead arrest at thirteen distinct arrest points, that correspond to G_1 (1a-c, 5*, 6*), S (2), G_2/M (3a-e), M (4) and T (5*, 6*) arrest, or progress through a partial cyclic attractor with repeated DNA replication and nuclear division but no cell division, resulting in multinucleate cells. Note that group 5 and 6 contains G1 and T arrested cells that completed nuclear division but arrest before cytokinesis. They can only be distinguished on the DNA/Nuclei count (Table S3).

Table 1: List of gap-filling additions.

Table 2: List of inconsistent mutants. The model predicted 23 viable mutants to be lethal. Of these, seven mutants can be explained by implicit synthetic lethality: While *cln3* and *clb5* are viable, the model lacks implementation of mechanistic functions of Bck2 and Clb3/Clb4, which are necessary in the absence of Cln3 and Clb5, respectively. While other mutants have strong phenotypes in e.g. cell size or growth rate, such as *swi4*, *swi6* and *pds1*, others - like *cdh1* - have no strong phenotypes as single mutations, suggesting that redundancy mechanisms are missing in our current knowledge.

Fig S1: The molecular architecture of the cell division cycle. Available at https://github.com/rxncon/models/ as CDC_S_cerevisiae_poster.pdf.

Table S1: The rxncon model. Available at https://github.com/rxncon/models/ as CDC_S_cerevisiae.xls.

Table S2: List of references.

Table S3: List of mutants examined in this study.

ACKNOWLEDGEMENTS

The authors would like to thank Jesper Romers, Sebastian Thieme and Mathias Wajnberg for close collaboration in the methods development, and Matteo Barberis for critical comments on the network model. MK would like to thank Hiroaki Kitano and Stefan Hohmann for the inspiration and support to tackle the challenge of large-scale signalling networks. This work was supported by the German Federal Ministry of Education and Research via e:Bio Cellemental (FKZ0316193, to MK).

REFERENCES

Blinov, M. L., J. R. Faeder, B. Goldstein and W. S. Hlavacek (2004). "BioNetGen: software for rule-based modeling of signal transduction based on the interactions of molecular domains." Bioinformatics **20**(17): 3289-3291.

Chasapi, A., P. Wachowicz, A. Niknejad, P. Collin, A. Krapp, E. Cano, V. Simanis and I. Xenarios (2015). "An Extended, Boolean Model of the Septation Initiation Network in S.Pombe Provides Insights into Its Regulation." PLoS One **10**(8): e0134214.

Chen, K. C., A. Csikasz-Nagy, B. Gyorffy, J. Val, B. Novak and J. J. Tyson (2000). "Kinetic analysis of a molecular model of the budding yeast cell cycle." Mol Biol Cell **11**(1): 369-391.

Chylek, L. A., L. A. Harris, J. R. Faeder and W. S. Hlavacek (2015). "Modeling for (physical) biologists: an introduction to the rule-based approach." Phys Biol **12**(4): 045007.

Danos, V., J. Feret, W. Fontana, R. Harmer and J. Krivine (2007). Rule-Based Modelling of Cellular Signalling. CONCUR 2007 – Concurrency Theory: 18th International Conference, CONCUR 2007, Lisbon, Portugal, September 3-8, 2007. Proceedings. L. Caires and V. T. Vasconcelos. Berlin, Heidelberg, Springer Berlin Heidelberg**:** 17-41.

Enserink, J. M. and R. D. Kolodner (2010). "An overview of Cdk1-controlled targets and processes." Cell Div **5**: 11.

Epstein, C. B. and F. R. Cross (1994). "Genes that can bypass the CLN requirement for Saccharomyces cerevisiae cell cycle START." Mol Cell Biol **14**(3): 2041-2047.

Faeder, J. R., M. L. Blinov and W. S. Hlavacek (2009). "Rule-based modeling of biochemical systems with BioNetGen." Methods Mol Biol **500**: 113-167.

Giese, W., M. Eigel, S. Westerheide, C. Engwer and E. Klipp (2015). "Influence of cell shape, inhomogeneities and diffusion barriers in cell polarization models." Phys Biol **12**(6): 066014.

Goldbeter, A. (1991). "A minimal cascade model for the mitotic oscillator involving cyclin and cdc2 kinase." Proc Natl Acad Sci U S A **88**(20): 9107-9111.

Hartwell, L. H., J. Culotti and B. Reid (1970). "Genetic control of the cell-division cycle in yeast. I. Detection of mutants." Proc Natl Acad Sci U S A **66**(2): 352-359.

Herrgard, M. J., N. Swainston, P. Dobson, W. B. Dunn, K. Y. Arga, M. Arvas, N. Bluthgen, S. Borger, R. Costenoble, M. Heinemann, M. Hucka, N. Le Novere, P. Li, W. Liebermeister, M. L. Mo, A. P. Oliveira, D. Petranovic, S. Pettifer, E. Simeonidis, K. Smallbone, I. Spasic, D. Weichart, R. Brent, D. S. Broomhead, H. V. Westerhoff, B. Kirdar, M. Penttila, E. Klipp, B. O. Palsson, U. Sauer, S. G. Oliver, P. Mendes, J. Nielsen and D. B. Kell (2008). "A consensus yeast metabolic network reconstruction obtained from a community approach to systems biology." Nat Biotechnol **26**(10): 1155-1160.

Hlavacek, W. S. and J. R. Faeder (2009). "The complexity of cell signaling and the need for a new mechanics." Sci Signal **2**(81): pe46.

Howell, A. S. and D. J. Lew (2012). "Morphogenesis and the cell cycle." Genetics **190**(1): 51-77.

Kaizu, K., S. Ghosh, Y. Matsuoka, H. Moriya, Y. Shimizu-Yoshida and H. Kitano (2010). "A comprehensive molecular interaction map of the budding yeast cell cycle." Mol Syst Biol **6**: 415.

Karr, J. R., J. C. Sanghvi, D. N. Macklin, M. V. Gutschow, J. M. Jacobs, B. Bolival, Jr., N. Assad-Garcia, J. I. Glass and M. W. Covert (2012). "A whole-cell computational model predicts phenotype from genotype." Cell **150**(2): 389-401.

Kitano, H. (2002). "Computational systems biology." Nature **420**(6912): 206-210.

Kraikivski, P., K. C. Chen, T. Laomettachit, T. M. Murali and J. J. Tyson (2015). "From START to FINISH: computational analysis of cell cycle control in budding yeast." NPJ Syst Biol Appl **1**: 15016.

Le Novere, N., M. Hucka, H. Mi, S. Moodie, F. Schreiber, A. Sorokin, E. Demir, K. Wegner, M. I. Aladjem, S. M. Wimalaratne, F. T. Bergman, R. Gauges, P. Ghazal, H. Kawaji, L. Li, Y.

Matsuoka, A. Villeger, S. E. Boyd, L. Calzone, M. Courtot, U. Dogrusoz, T. C. Freeman, A. Funahashi, S. Ghosh, A. Jouraku, S. Kim, F. Kolpakov, A. Luna, S. Sahle, E. Schmidt, S. Watterson, G. Wu, I. Goryanin, D. B. Kell, C. Sander, H. Sauro, J. L. Snoep, K. Kohn and H. Kitano (2009). "The Systems Biology Graphical Notation." Nat Biotechnol **27**(8): 735-741.

Linke, C., A. Chasapi, A. Gonzalez-Novo, I. Al Sawad, S. Tognetti, E. Klipp, M. Loog, S. Krobitsch, F. Posas, I. Xenarios and M. Barberis (2017). "A Clb/Cdk1-mediated regulation of Fkh2 synchronizes CLB expression in the budding yeast cell cycle." NPJ Syst Biol Appl **3**: 7.

Lubitz, T., J. Hahn, F. T. Bergmann, E. Noor, E. Klipp and W. Liebermeister (2016). "SBtab: a flexible table format for data exchange in systems biology." Bioinformatics.

Michal, G. (2014). "Roche Biochemical Pathways (wall chart)."

Münzner, U., T. Lubitz, E. Klipp and M. Krantz (2017). Towards genome-scale models of signal transduction networks. Systems Biology. J. Nielsen and S. Hohmann, Wiley**:** 215-242.

Mussel, C., M. Hopfensitz and H. A. Kestler (2010). "BoolNet--an R package for generation, reconstruction and analysis of Boolean networks." Bioinformatics **26**(10): 1378-1380.

Neiman, A. M. (2011). "Sporulation in the budding yeast Saccharomyces cerevisiae." Genetics **189**(3): 737-765.

Romers, J., S. Thieme, U. Münzner and M. Krantz (2018). Pre-print: Using rxncon to develop rule based models. Modeling Biomolecular Site Dynamics. W. S. Hlavacek, Springer.

Romers, J. C. and M. Krantz (2017). "Pre-print: rxncon 2.0: a language for executable molecular systems biology." bioRxiv.

Romers, J. C., S. Thieme, U. Muenzner and M. Krantz (2017). "Pre-print: A scalable method for parameter-free simulation and validation of mechanistic cellular signal transduction network models." bioRxiv.

Rother, M., U. Munzner, S. Thieme and M. Krantz (2013). "Information content and scalability in signal transduction network reconstruction formats." Mol Biosyst **9**(8): 1993-2004.

Schwob, E. and K. Nasmyth (1993). "CLB5 and CLB6, a new pair of B cyclins involved in DNA replication in Saccharomyces cerevisiae." Genes Dev **7**(7A): 1160-1175.

Spiesser, T. W., F. Uschner, S. O. Adler, U. Münzner, M. Krantz and E. Klipp (Submitted). "A yeast cell cycle model integrating stress, signaling, and physiology."

Thiele, I., N. Swainston, R. M. Fleming, A. Hoppe, S. Sahoo, M. K. Aurich, H. Haraldsdottir, M. L. Mo, O. Rolfsson, M. D. Stobbe, S. G. Thorleifsson, R. Agren, C. Bolling, S. Bordel, A. K. Chavali, P. Dobson, W. B. Dunn, L. Endler, D. Hala, M. Hucka, D. Hull, D. Jameson, N. Jamshidi, J. J. Jonsson, N. Juty, S. Keating, I. Nookaew, N. Le Novere, N. Malys, A. Mazein, J. A. Papin, N. D. Price, E. Selkov, Sr., M. I. Sigurdsson, E. Simeonidis, N. Sonnenschein, K. Smallbone, A. Sorokin, J. H. van Beek, D. Weichart, I. Goryanin, J. Nielsen, H. V. Westerhoff, D. B. Kell, P. Mendes and B. O. Palsson (2013). "A community-driven global reconstruction of human metabolism." Nat Biotechnol **31**(5): 419-425.

Tiger, C. F., F. Krause, G. Cedersund, R. Palmer, E. Klipp, S. Hohmann, H. Kitano and M. Krantz (2012). "A framework for mapping, visualisation and automatic model creation of signal-transduction networks." Mol Syst Biol **8**: 578.

Figure 1

Figure 2

Figure 3

Figure 4

TABLE 1

Observation	Solution	Reference
MBF target genes constitutively expressed	Swi6 phosphorylation strict requirement for MBF activity	Palumbo et al. (2016)
Fkh2, Ndd1, Mcm1 target genes constitutively expressed	Fkh2 phosphorylation strict requirement for Fkh2 activity	Pic-Taylor et al. (2004)
Clb2 and Clb5 not degraded in G1	Ubiqitylation by both APC-Cdc20 and APC-Cdh1	Wäsch and Cross (2002), Lu et al. (2014)
Pcl9 degradation not regulated	Regulated degradation equivalent to Pcl1,2 degradation	Hernandez-Ortega et al. (2013)
Pho85-Pcl9 phosphorylation of Whi5 dominates Cdc28-mediated Whi5 phosphorylation	Pho85-Pcl9 phosphorylation positive influence on Whi5 dissociation	*Hypothesis: Alternatively, Pcl9 expression could be nutrient dependent as for Cln3*
Sic1 deactivation premature	Phosphorylation restricted to Cdc28-Cln1,2	*Hypothesis: No Cdc28-Cln3 phosphosylation of Sic1*
Bud growth module regulation insufficient	Symmetry breaking included as output/input when criteria for polarisation are fulfilled	*Placeholder for quantitative phenomenon. Cannot be described mechanstically with Boolean logic*
Stable complexes in SPB cycle disappeared	[Daughter SPB] present with duplicated and/or separated SPBs	*Technical correction*
Periodic Rad53 transcription and constant degradation lead to Rad53 periodic depletion	Constant Rad53 expression: MBF regulation as positive influence instead of strict requirement	*Hypothesis: Basal Rad53 expression constitutive to account for constant role in DNA damage response*
Discrepancy between Boolean and physiological time	Introduction of delays on transcriptional reactions	*Time scale separation. Solution to technical issue with Boolean networks.*
Dbf2 release premature	Correct spindle positioning blocks phosphorylation of Kin4 by Elm1	Consistent with SPOC activation of phosphorylation; Kurtulmus et al. 2010
No SPB duplication due to premature Spc42 disappearance. MBF transcription very brief.	Introduction of delays on replication	*Time scale separation. Solution to technical issue with Boolean networks.*
SPB positioning premature	Kar9 phosphorylation by Cdc28-Clb1,2,5,6	*Hypothesis*
No (morphological) cell division cycle due to premature activation of APC (due to Mad2 depeletion).	Degradation of Cdc20 block when bound to Mad2	*Hypothesis*
No (morphological) cell division cycle: Cytokinesis doesn't trigger as symmetry breaking and Chs2 release never overlap	Acm1 phosphorylation completely blocks (x) instead of decreases (k-) Acm1 degradation	*Hypothesis*

TABLE 2

Mutant	Arrest point	DNA	Nuclei	Comment	Synthetic Lethality
cln3	lic, sat, 0	1	1	G1-arrest	*cln3bck2*
cln3mbp1	lic, sat, 0	1	1	G1-arrest	*cln3bck2*
cln3swi6	lic, sat, 0	1	1	G1-arrest	*cln3bck2*
cln3swi4	lic, sat, 0	1	1	G1-arrest	*cln3bck2*
swi4	lic, sat, large	1	1	G1-arrest	
swi6	lic, sat, large	1	1	G1-arrest	
swi64A	lic, sat, large	1	1	G1-arrest	
mbp1	lic, sat, large	1	1	G1-arrest	
clb5	lic, sat, large	1	1	G1-arrest	*clb3clb4clb5*
cdc20pds1clb5	lic, sat, large	1	1	G1-arrest	*clb3clb4clb5*
cdc20pds1clb5cdh1SIC1	lic, sat, large	1	1	G1-arrest	*clb3clb4clb5*
spc1102ACDK	lic, sat, large	1	1	G1-arrest	
Sld2A	lic, sat, large	1	1	G1-arrest	
Sld3T600A	lic, sat, large	1	1	G1-arrest	
Sld3S622A	lic, sat, large	1	1	G1-arrest	
Sic17A	lic, sat, large	1	1	G1-arrest	
cdh1	0, sat, 0	1	1	G1-arrest	
cln1cln2cdh1	0, sat, 0	1	1	G1-arrest	
spc1104A	rep'd, dup, large	1	1	G2/M-arrest	
pds1	0, 0, large	2	2	T-arrest	
net1cdc15cdh1	0, sat, large	2	2	T-arrest	
cdc15net1	?, ?, large	*	*	Multinucleate: DNA replication and ND, no cell division	
tem1net1	?, ?, large	*	*	Multinucleate: DNA replication and ND, no cell division	

References

[1] Mohammed Sbia, Emily J Parnell, Yaxin Yu, Aileen E Olsen, Kelsi L Kretschmann, Warren P Voth, and David J Stillman. Regulation of the yeast ace2 transcription factor during the cell cycle. *Journal of Biological Chemistry*, 283(17):11135–11145, 2008.

[2] David Pruyne, Marie Evangelista, Changsong Yang, Erfei Bi, Sally Zigmond, Anthony Bretscher, and Charles Boone. Role of formins in actin assembly: nucleation and barbed-end association. *Science*, 297(5581):612–615, 2002.

[3] Martine Coué, Stephen L Brenner, Ilan Spector, and Edward D Korn. Inhibition of actin polymerization by latrunculin a. *FEBS letters*, 213(2):316–318, 1987.

[4] Sue Biggins. The composition, functions, and regulation of the budding yeast kineto-chore. *Genetics*, 194(4):817–846, 2013.

[5] Jorrit M Enserink and Richard D Kolodner. An overview of Cdk1-controlled targets and processes. *Cell Division*, 5:11, 2010.

[6] Eric L Weiss. Mitotic exit and separation of mother and daughter cells. *Genetics*, 192(4):1165–1202, 2012.

[7] Maria Enquist-Newman, Matt Sullivan, and David O Morgan. Modulation of the mitotic regulatory network by apc-dependent destruction of the cdh1 inhibitor acm1. *Molecular Cell*, 30(4):437–446, 2008.

[8] Denis Ostapenko, Janet L Burton, Ruiwen Wang, and Mark J Solomon. Pseudosubstrate inhibition of the anaphase-promoting complex by acm1: regulation by proteolysis and cdc28 phosphorylation. *Molecular and Cellular Biology*, 28(15):4653–4664, 2008.

[9] Karen Crasta, Phillips Huang, Garry Morgan, Mark Winey, and Uttam Surana. Cdk1 regulates centrosome separation by restraining proteolysis of microtubule-associated proteins. *The EMBO Journal*, 25(11):2551–2563, 2006.

[10] Ralph Wäsch and Frederick R Cross. Apc-dependent proteolysis of the mitotic cyclin clb2 is essential for mitotic exit. *Nature*, 418(6897):556, 2002.

[11] Dan Lu, Jennifer Y Hsiao, Norman E Davey, Vanessa A Van Voorhis, Scott A Foster, Chao Tang, and David O Morgan. Multiple mechanisms determine the order of apc/c substrate degradation in mitosis. *The Journal of Cell Biology*, 207(1):23–39, 2014.

[12] Mackenzie E Malo, Spike DL Postnikoff, Terra G Arnason, and Troy AA Harkness. Mitotic degradation of yeast fkh1 by the anaphase promoting complex is required for normal longevity, genomic stability and stress resistance. *Aging (Albany NY)*, 8(4):810, 2016.

[13] Denis Ostapenko, Janet L Burton, and Mark J Solomon. Identification of anaphase promoting complex substrates in *S. cerevisiae*. *PLoS One*, 7(9):e45895, 2012.

[14] Denis Ostapenko and Mark J Solomon. Anaphase promoting complex–dependent degradation of transcriptional repressors nrm1 and yhp1 in *Saccharomyces cerevisiae*. *Molecular Biology of the Cell*, 22(13):2175–2184, 2011.

[15] Zoe Hilioti, Yin-Shin Chung, Yuko Mochizuki, Christopher FJ Hardy, and Orna Cohen-Fix. The anaphase inhibitor pds1 binds to the apc/c-associated protein cdc20 in a destruction box-dependent manner. *Current Biology*, 11(17):1347–1352, 2001.

[16] VıÌ₳ctor J Cid, Javier Jiménez, MarıÌ₳a Molina, Miguel Sánchez, César Nombela, and Jeremy W Thorner. Orchestrating the cell cycle in yeast: sequential localization of key mitotic regulators at the spindle pole and the bud neck. *Microbiology*, 148(9):2647–2659, 2002.

[17] Liam J Holt, Andrew N Krutchinsky, and David O Morgan. Positive feedback sharpens the anaphase switch. *Nature*, 454(7202):353, 2008.

[18] Jae ook Kim, Alex Zelter, Neil T Umbreit, Athena Bollozos, Michael Riffle, Richard Johnson, Michael J MacCoss, Charles L Asbury, and Trisha N Davis. The ndc80 complex bridges two dam1 complex rings. *eLife*, 6:e21069, 2017.

[19] Erfei Bi and Hay-Oak Park. Cell polarization and cytokinesis in budding yeast. *Genetics*, 191(2):347–387, 2012.

[20] Anne-Christine Butty, Nathalie Perrinjaquet, Audrey Petit, Malika Jaquenoud, Jeffrey E Segall, Kay Hofmann, Catherine Zwahlen, and Matthias Peter. A positive feedback loop stabilizes the guanine-nucleotide exchange factor cdc24 at sites of polarization. *The EMBO Journal*, 21(7):1565–1576, 2002.

[21] Audrey S Howell and Daniel J Lew. Morphogenesis and the cell cycle. *Genetics*, 190(1):51–77, 2012.

[22] Steven M Markus, Katelyn A Kalutkiewicz, and Wei-Lih Lee. Astral microtubule asymmetry provides directional cues for spindle positioning in budding yeast. *Experimental Cell Research*, 318(12):1400–1406, 2012.

[23] Jeffrey K Moore and Rita K Miller. The cyclin-dependent kinase cdc28p regulates multiple aspects of kar9p function in yeast. *Molecular Biology of the Cell*, 18(4):1187–1202, 2007.

[24] Marie Evangelista, David Pruyne, David C Amberg, Charles Boone, and Anthony Bretscher. Formins direct arp2/3-independent actin filament assembly to polarize cell growth in yeast. *Nature Cell Biology*, 4(1):32, 2002.

[25] Marie Evangelista, Kelly Blundell, Mark S Longtine, Clinton J Chow, Neil Adames, John R Pringle, Matthias Peter, and Charles Boone. Bni1p, a yeast formin linking cdc42p and the actin cytoskeleton during polarized morphogenesis. *Science*, 276(5309):118–122, 1997.

[26] H Kohno, K Tanaka, A Mino, M Umikawa, H Imamura, T Fujiwara, Y Fujita, K Hotta, H Qadota, T Watanabe, et al. Bni1p implicated in cytoskeletal control is a putative target of rho1p small gtp binding protein in *Saccharomyces cerevisiae*. *The EMBO Journal*, 15(22):6060, 1996.

[27] Masayuki Iwase, Jianying Luo, Satish Nagaraj, Mark Longtine, Hyong Bai Kim, Brian K Haarer, Carlo Caruso, Zongtian Tong, John R Pringle, and Erfei Bi. Role of a cdc42p effector pathway in recruitment of the yeast septins to the presumptive bud site. *Molecular Biology of the Cell*, 17(3):1110–1125, 2006.

[28] Dane A Mohl, Michael J Huddleston, Therese S Collingwood, Roland S Annan, and Raymond J Deshaies. Dbf2–mob1 drives relocalization of protein phosphatase cdc14 to the cytoplasm during exit from mitosis. *The Journal of Cell Biology*, 184(4):527–539, 2009.

[29] Satoshi Yoshida and Akio Toh-e. Budding yeast cdc5 phosphorylates net1 and assists cdc14 release from the nucleolus. *Biochemical and biophysical research communications*, 294(3):687–691, 2002.

[30] Ramzi Azzam, Susan L Chen, Wenying Shou, Angie S Mah, Gabriela Alexandru, Kim Nasmyth, Roland S Annan, Steven A Carr, and Raymond J Deshaies. Phosphorylation by cyclin b-cdk underlies release of mitotic exit activator cdc14 from the nucleolus. *Science*, 305(5683):516–519, 2004.

[31] Wenying Shou, Jae Hong Seol, Anna Shevchenko, Christopher Baskerville, Danesh Moazed, ZW Susan Chen, Joanne Jang, Andrej Shevchenko, Harry Charbonneau, and Raymond J Deshaies. Exit from mitosis is triggered by tem1-dependent release of the protein phosphatase cdc14 from nucleolar rent complex. *Cell*, 97(2):233–244, 1999.

[32] Benjamin Woods, Chun-Chen Kuo, Chi-Fang Wu, Trevin R Zyla, and Daniel J Lew. Polarity establishment requires localized activation of cdc42. *The Journal of Cell Biology*, 211(1):19–26, 2015.

[33] Karen E Ross, Philipp Kaldis, and Mark J Solomon. Activating phosphorylation of the *Saccharomyces cerevisiae* cyclin-dependent kinase, cdc28p, precedes cyclin binding. *Molecular Biology of the Cell*, 11(5):1597–1609, 2000.

[34] Matthias Peter and Ira Herskowitz. Direct inhibition of the yeast cyclin-dependent kinase cdc28-cln by far1. *Science*, 265(5176):1228–1232, 1994.

[35] Bong-Kwan Han, Lydia M Bogomolnaya, James M Totten, Heidi M Blank, Lawrence J Dangott, and Michael Polymenis. Bem1p, a scaffold signaling protein, mediates cyclin-dependent control of vacuolar homeostasis in saccharomyces cerevisiae. *Genes & Development*, 19(21):2606–2618, 2005.

[36] Derek McCusker, Carilee Denison, Scott Anderson, Thea A Egelhofer, John R Yates, Steven P Gygi, and Douglas R Kellogg. Cdk1 coordinates cell-surface growth with the cell cycle. *Nature Cell Biology*, 9(5):506–515, 2007.

[37] Eric M Mortensen, Wilhelm Haas, Melanie Gygi, Steven P Gygi, and Douglas R Kellogg. Cdc28-dependent regulation of the cdc5/polo kinase. *Current Biology*, 15(22):2033–2037, 2005.

[38] Suzanne Elsasser, Yong Chi, Ping Yang, and Judith L Campbell. Phosphorylation controls timing of cdc6p destruction: a biochemical analysis. *Molecular Biology of the Cell*, 10(10):3263–3277, 1999.

[39] Ee Mei Teh, Chuan Chung Chai, and Foong May Yeong. Retention of chs2p in the er requires n-terminal cdk1-phosphorylation sites. *Cell Cycle*, 8(18):2965–2976, 2009.

[40] Benjamin D Landry, John P Doyle, David P Toczyski, and Jennifer A Benanti. F-box protein specificity for g1 cyclins is dictated by subcellular localization. *PLoS Genetics*, 8(7):e1002851, 2012.

[41] Mart Loog and David O Morgan. Cyclin specificity in the phosphorylation of cyclin-dependent kinase substrates. *Nature*, 434(7029):104–108, 2005.

[42] Aline Pic-Taylor, Zoulfia Darieva, Brian A Morgan, and Andrew D Sharrocks. Regulation of cell cycle-specific gene expression through cyclin-dependent kinase-mediated phosphorylation of the forkhead transcription factor fkh2p. *Molecular and Cellular Biology*, 24(22):10036–10046, 2004.

[43] Vincent Archambault, Emmanuel J Chang, Benjamin J Drapkin, Frederick R Cross, Brian T Chait, and Michael P Rout. Targeted proteomic study of the cyclin-cdk module. *Molecular Cell*, 14(6):699–711, 2004.

[44] Muluye E Liku, Van Q Nguyen, Audrey W Rosales, Kaoru Irie, and Joachim J Li. Cdk phosphorylation of a novel nls-nes module distributed between two subunits of the mcm2-7 complex prevents chromosomal rereplication. *Molecular Biology of the Cell*, 16(10):5026–5039, 2005.

[45] Sue L Jaspersen, Brenda J Huneycutt, Thomas H Giddings, Katheryn A Resing, Natalie G Ahn, and Mark Winey. Cdc28/cdk1 regulates spindle pole body duplication through phosphorylation of spc42 and mps1. *Developmental Cell*, 7(2):263–274, 2004.

[46] David Reynolds, Bu Jun Shi, Cameron McLean, Frosa Katsis, Bruce Kemp, and Stephen Dalton. Recruitment of thr 319-phosphorylated ndd1p to the fha domain of fkh2p requires clbkinase activity: a mechanism for clb cluster gene activation. *Genes & Development*, 17(14):1789–1802, 2003.

[47] Van Q Nguyen, Joachim J Li, et al. Cyclin-dependent kinases prevent dna re-replication through multiple mechanisms. *Nature*, 411(6841):1068–1073, 2001.

[48] Mardo Kõivomägi, Ervin Valk, Rainis Venta, Anna Iofik, Martin Lepiku, David O Morgan, and Mart Loog. Dynamics of cdk1 substrate specificity during the cell cycle. *Molecular Cell*, 42(5):610–623, 2011.

[49] Richelle Sopko, Dongqing Huang, Jeffrey C Smith, Daniel Figeys, and Brenda J Andrews. Activation of the cdc42p gtpase by cyclin-dependent protein kinases in budding yeast. *The EMBO Journal*, 26(21):4487–4500, 2007.

[50] Menattallah Elserafy, Mirela Šarić, Annett Neuner, Tien-chen Lin, Wanlu Zhang, Christian Seybold, Lavanya Sivashanmugam, and Elmar Schiebel. Molecular mechanisms that restrict yeast centrosome duplication to one event per cell cycle. *Current Biology*, 24(13):1456–1466, 2014.

[51] Seiji Tanaka, Toshiko Umemori, Kazuyuki Hirai, Sachiko Muramatsu, Yoichiro Kamimura, and Hiroyuki Araki. Cdk-dependent phosphorylation of sld2 and sld3 initiates dna replication in budding yeast. *Nature*, 445(7125):328, 2007.

[52] Xavier Robellet, Yogitha Thattikota, Fang Wang, Tse-Luen Wee, Mirela Pascariu, Sahana Shankar, Éric Bonneil, Claire M Brown, and Damien D'Amours. A high-sensitivity phospho-switch triggered by cdk1 governs chromosome morphogenesis during cell division. *Genes & Development*, 29(4):426–439, 2015.

[53] Tien-chen Lin, Annett Neuner, Yvonne T Schlosser, Elmar Schiebel, Annette ND Scharf, and Lisa Weber. Cell-cycle dependent phosphorylation of yeast pericentrin regulates γ-tusc-mediated microtubule nucleation. *Elife*, 3:e02208, 2014.

[54] Andrew W Truman, Kolbrun Kristjansdottir, Donald Wolfgeher, Naushaba Hasin, Sigrun Polier, Hong Zhang, Sarah Perrett, Chrisostomos Prodromou, Gary W Jones, and Stephen J Kron. Cdk-dependent hsp70 phosphorylation controls g1 cyclin abundance and cell-cycle progression. *Cell*, 151(6):1308–1318, 2012.

[55] Michael Costanzo, Oliver Schub, and Brenda Andrews. G1 transcription factors are differentially regulated in *Saccharomyces cerevisiae* by the swi6-binding protein stb1. *Molecular and Cellular Biology*, 23(14):5064–5077, 2003.

[56] Steven B Haase and Curt Wittenberg. Topology and control of the cell-cycle-regulated transcriptional circuitry. *Genetics*, 196(1):65–90, 2014.

[57] Michelle V Wagner, Marcus B Smolka, Rob AM De Bruin, Huilin Zhou, Curt Wittenberg, and Steven F Dowdy. Whi5 regulation by site specific cdk-phosphorylation in *Saccharomyces cerevisiae*. *PLoS One*, 4(1):e4300, 2009.

[58] Jamie M Keck, Michele H Jones, Catherine CL Wong, Jonathan Binkley, Daici Chen, Sue L Jaspersen, Eric P Holinger, Tao Xu, Mario Niepel, Michael P Rout, et al. A cell cycle phosphoproteome of the yeast centrosome. *Science*, 332(6037):1557–1561, 2011.

[59] Keiko Kono, Satoru Nogami, Mitsuhiro Abe, Masafumi Nishizawa, Shinichi Morishita, David Pellman, and Yoshikazu Ohya. G1/s cyclin-dependent kinase regulates small gtpase rho1p through phosphorylation of rhogef tus1p in *Saccharomyces cerevisiae*. *Molecular Biology of the Cell*, 19(4):1763–1771, 2008.

[60] Julia A Yaglom, Alfred L Goldberg, Daniel Finley, and Michael Y Sherman. The molecular chaperone ydj1 is required for the p34cdc28-dependent phosphorylation of the cyclin cln3 that signals its degradation. *Molecular and Cellular Biology*, 16(7):3679–3684, 1996.

[61] Matthias Versele and Jeremy Thorner. Septin collar formation in budding yeast requires gtp binding and direct phosphorylation by the pak, cla4. *The Journal of Cell Biology*, 164(5):701–715, 2004.

[62] Hao Wu, Courtney Turner, Jimmy Gardner, Brenda Temple, and Patrick Brennwald. The exo70 subunit of the exocyst is an effector for both cdc42 and rho3 function in polarized exocytosis. *Molecular biology of the cell*, 21(3):430–442, 2010.

[63] Joseph TP Yeeles, Tom D Deegan, Agnieszka Janska, Anne Early, and John FX Diffley. Regulated eukaryotic dna replication origin firing with purified proteins. *Nature*, 519(7544):431–435, 2015.

[64] Gordon Perkins, Lucy S Drury, and John FX Diffley. Separate scf cdc4 recognition elements target cdc6 for proteolysis in s phase and mitosis. *The EMBO Journal*, 20(17):4836–4845, 2001.

[65] Leisa P Jackson, Steven I Reed, and Steven B Haase. Distinct mechanisms control the stability of the related s-phase cyclins clb5 and clb6. *Molecular and Cellular Biology*, 26(6):2456–2466, 2006.

[66] Michael D Mendenhall and Amy E Hodge. Regulation of cdc28 cyclin-dependent protein kinase activity during the cell cycle of the yeast *Saccharomyces cerevisiae. Microbiology and Molecular Biology Reviews*, 62(4):1191–1243, 1998.

[67] Inma Quilis and J Carlos Igual. A comparative study of the degradation of yeast cyclins cln1 and cln2. *FEBS open bio*, 7(1):74–87, 2017.

[68] Anton Gartner, Alexandra Jovanović, Doo-Il Jeoung, Sarah Bourlat, Frederick R Cross, and Gustav Ammerer. Pheromone-dependent g1 cell cycle arrest requires far1 phosphorylation, but may not involve inhibition of cdc28-cln2 kinase, in vivo. *Molecular and Cellular Biology*, 18(7):3681–3691, 1998.

[69] Marc Blondel, Jean-Marc Galan, Yong Chi, Céline Lafourcade, Céline Longaretti, Raymond J Deshaies, and Matthias Peter. Nuclear-specific degradation of far1 is controlled by the localization of the f-box protein cdc4. *The EMBO Journal*, 19(22):6085–6097, 2000.

[70] Piers Nash, Xiaojing Tang, Stephen Orlicky, Qinghua Chen, Frank B Gertler, Michael D Mendenhall, Frank Sicheri, Tony Pawson, and Mike Tyers. Multisite phosphorylation of a cdk inhibitor sets a threshold for the onset of dna replication. *Nature*, 414(6863):514–521, 2001.

[71] Jose-Antonio Rodriguez-Rodriguez, Yolanda Moyano, Soraya Játiva, and Ethel Queralt. Mitotic exit function of polo-like kinase cdc5 is dependent on sequential activation by cdk1. *Cell Reports*, 15(9):2050–2062, 2016.

[72] Ethel Queralt, Chris Lehane, Bela Novak, and Frank Uhlmann. Downregulation of pp2a cdc55 phosphatase by separase initiates mitotic exit in budding yeast. *Cell*, 125(4):719–732, 2006.

[73] Satoshi Asano, Jung-Eun Park, Krisada Sakchaisri, Li-Rong Yu, Sukgil Song, Porntip Supavilai, Timothy D Veenstra, and Kyung S Lee. Concerted mechanism of swe1/wee1 regulation by multiple kinases in budding yeast. *The EMBO Journal*, 24(12):2194–2204, 2005.

[74] Christian Speck, Zhiqiang Chen, Huilin Li, and Bruce Stillman. Atpase-dependent, cooperative binding of orc and cdc6p to origin dna. *Nature Structural & Molecular Biology*, 12(11):965, 2005.

[75] Stephen P Bell and Karim Labib. Chromosome duplication in *Saccharomyces cerevisiae*. *Genetics*, 203(3):1027–1067, 2016.

[76] Rentian Wu, Jiafeng Wang, and Chun Liang. Cdt1p, through its interaction with mcm6p, is required for the formation, nuclear accumulation and chromatin loading of the mcm complex. *J Cell Sci*, 125(1):209–219, 2012.

[77] Jonathan M Scholey, Gul Civelekoglu-Scholey, and Ingrid Brust-Mascher. Anaphase b. *Biology*, 5(4):51, 2016.

[78] Jingyan Fu, Iain M Hagan, and David M Glover. The centrosome and its duplication cycle. *Cold Spring Harbor Perspectives in Biology*, 7(2):a015800, 2015.

[79] David Pruyne, Lina Gao, Erfei Bi, and Anthony Bretscher. Stable and dynamic axes of polarity use distinct formin isoforms in budding yeast. *Molecular Biology of the Cell*, 15(11):4971–4989, 2004.

[80] Shuyan Chen, Milan A de Vries, and Stephen P Bell. Orc6 is required for dynamic recruitment of cdt1 during repeated mcm2–7 loading. *Genes & Development*, 21(22):2897–2907, 2007.

[81] Jérôme Poli, Olga Tsaponina, Laure Crabbé, Andrea Keszthelyi, Véronique Pantesco, Andrei Chabes, Armelle Lengronne, and Philippe Pasero. dntp pools determine fork progression and origin usage under replication stress. *The EMBO Journal*, 31(4):883–894, 2012.

[82] David Norris, Barbara Dunn, and Mary Ann Osley. The effect of histone gene deletions on chromatin structure in *Saccharomyces cerevisiae*. *Science*, 242(4879):759–761, 1988.

[83] Max E Douglas and John FX Diffley. Recruitment of mcm10 to sites of replication initiation requires direct binding to the minichromosome maintenance (mcm) complex. *Journal of Biological Chemistry*, 291(11):5879–5888, 2016.

[84] Marko Lõoke, Michael F Maloney, and Stephen P Bell. Mcm10 regulates dna replication elongation by stimulating the cmg replicative helicase. *Genes & Development*, 31(3):291–305, 2017.

[85] Diana Rüthnick and Elmar Schiebel. Duplication of the yeast spindle pole body once per cell cycle. *Molecular and Cellular Biology*, 36(9):1324–1331, 2016.

[86] Adele L Marston. Chromosome segregation in budding yeast: sister chromatid cohesion and related mechanisms. *Genetics*, 196(1):31–63, 2014.

[87] Chetan K Rane and Audrey Minden. P21 activated kinases: structure, regulation, and functions. *Small GTPases*, 5(1):e28003, 2014.

[88] Angela C Wild, W Yu Jong, Mark A Lemmon, and Kendall J Blumer. The p21-activated protein kinase-related kinase cla4 is a coincidence detector of signaling by cdc42 and phosphatidylinositol 4-phosphate. *Journal of Biological Chemistry*, 279(17):17101–17110, 2004.

[89] Douglas M Cyr and Michael G Douglas. Differential regulation of hsp70 subfamilies by the eukaryotic dnaj homologue ydj1. *Journal of Biological Chemistry*, 269(13):9798–9804, 1994.

[90] Iain M Cheeseman, Scott Anderson, Miri Jwa, Erin M Green, Jung-seog Kang, John R Yates, Clarence SM Chan, David G Drubin, and Georjana Barnes. Phospho-regulation of kinetochore-microtubule attachments by the aurora kinase ipl1p. *Cell*, 111(2):163–172, 2002.

[91] Fabienne Lampert, Peter Hornung, and Stefan Westermann. The dam1 complex confers microtubule plus end–tracking activity to the ndc80 kinetochore complex. *The Journal of Cell Biology*, 189(4):641–649, 2010.

[92] Younghoon Oh, Kuang-Jung Chang, Peter Orlean, Carsten Wloka, Raymond Deshaies, and Erfei Bi. Mitotic exit kinase dbf2 directly phosphorylates chitin synthase chs2 to regulate cytokinesis in budding yeast. *Molecular Biology of the Cell*, 23(13):2445–2456, 2012.

[93] Angie S Mah, Joanne Jang, and Raymond J Deshaies. Protein kinase cdc15 activates the dbf2-mob1 kinase complex. *Proceedings of the National Academy of Sciences*, 98(13):7325–7330, 2001.

[94] Kristen J Holmes, Daniel M Klass, Evan L Guiney, and Martha S Cyert. Whi3, an *S. cerevisiae* rna-binding protein, is a component of stress granules that regulates levels of its target mrnas. *PLoS One*, 8(12):e84060, 2013.

[95] Sara Hernández-Ortega, Samuel Bru, Natalia Ricco, Sara Ramírez, Núria Casals, Javier Jiménez, Marta Isasa, Bernat Crosas, and Josep Clotet. Defective in mitotic arrest 1 (dma1) ubiquitin ligase controls g1 cyclin degradation. *Journal of Biological Chemistry*, 288(7):4704–4714, 2013.

[96] Benjamin Pardo, Laure Crabbé, and Philippe Pasero. Signaling pathways of replication stress in yeast. *FEMS Yeast Research*, 17(2), 2017.

[97] Sheng-hong Chen and Huilin Zhou. Reconstitution of rad53 activation by mec1 through adaptor protein mrc1. *Journal of Biological Chemistry*, 284(28):18593–18604, 2009.

[98] JJ L Miranda, David S King, and Stephen C Harrison. Protein arms in the kinetochore-microtubule interface of the yeast dash complex. *Molecular Biology of the Cell*, 18(7):2503–2510, 2007.

[99] Thibault Legal, Juan Zou, Alicja Sochaj, Juri Rappsilber, and Julie PI Welburn. Molecular architecture of the dam1 complex–microtubule interaction. *Open Biology*, 6(3):150237, 2016.

[100] Daniela Trinca Bertazzi, Bahtiyar Kurtulmus, and Gislene Pereira. The cortical protein lte1 promotes mitotic exit by inhibiting the spindle position checkpoint kinase kin4. *The Journal of Cell Biology*, 193(6):1033–1048, 2011.

[101] Gabriela Alexandru, Frank Uhlmann, Karl Mechtler, Marc-André Poupart, and Kim Nasmyth. Phosphorylation of the cohesin subunit scc1 by polo/cdc5 kinase regulates sister chromatid separation in yeast. *Cell*, 105(4):459–472, 2001.

[102] Rafal Ciosk, Wolfgang Zachariae, Christine Michaelis, Andrej Shevchenko, Matthias Mann, and Kim Nasmyth. An esp1/pds1 complex regulates loss of sister chromatid cohesion at the metaphase to anaphase transition in yeast. *Cell*, 93(6):1067–1076, 1998.

[103] Matthew Sullivan, Christine Lehane, and Frank Uhlmann. Orchestrating anaphase and mitotic exit: separase cleavage and localization of slk19. *Nature Cell Biology*, 3(9):771–777, 2001.

[104] Chong J Park, Jung-Eun Park, Tatiana S Karpova, Nak-Kyun Soung, Li-Rong Yu, Sukgil Song, Kyung H Lee, Xue Xia, Eugene Kang, Ilknur Dabanoglu, et al. Requirement for the budding yeast polo kinase cdc5 in proper microtubule growth and dynamics. *Eukaryotic Cell*, 7(3):444–453, 2008.

[105] Bing He, Fengong Xi, Xiaoyu Zhang, Jian Zhang, and Wei Guo. Exo70 interacts with phospholipids and mediates the targeting of the exocyst to the plasma membrane. *The EMBO Journal*, 26(18):4053–4065, 2007.

[106] Zoulfia Darieva, Richard Bulmer, Aline Pic-Taylor, Kathryn S Doris, Marco Geymonat, Steven G Sedgwick, Brian A Morgan, and Andrew D Sharrocks. Polo kinase controls cell cycle-dependent transcription by direct targeting of a coactivator protein. *Nature*, 444(7118):494, 2006.

[107] Syam Kumar Yelamanchi, Jiri Veis, Dorothea Anrather, Helene Klug, and Gustav Ammerer. Genotoxic stress prevents ndd1-dependent transcriptional activation of g2/m-specific genes in *Saccharomyces cerevisiae*. *Molecular and Cellular Biology*, 34(4):711–724, 2014.

[108] Hiroshi Qadota, Christophe P Python, Shunsuke B Inoue, Mikio Arisawa, Yasuhiro Anraku, Yi Zheng, Takahide Watanabe, David E Levin, and Yoshikazu Ohya. Identification of yeast rho1p gtpase as a regulatory subunit of 1, 3-β-glucan synthase. *Science*, pages 279–281, 1996.

[109] Jessica S Rosenberg, Frederick R Cross, and Hironori Funabiki. Knl1/spc105 recruits pp1 to silence the spindle assembly checkpoint. *Current Biology*, 21(11):942–947, 2011.

[110] Ellen R Edenberg, Kevin G Mark, and David P Toczyski. Ndd1 turnover by scfgrr1 is inhibited by the dna damage checkpoint in *Saccharomyces cerevisiae*. *PLoS Genetics*, 11(4):e1005162, 2015.

[111] Adam M Perez, Gregory C Finnigan, Françoise M Roelants, and Jeremy Thorner. Septin-associated protein kinases in the yeast *Saccharomyces cerevisiae*. *Frontiers in Cell and Developmental Biology*, 4, 2016.

[112] Yuko Nakajima, Randall G Tyers, Catherine CL Wong, John R Yates, David G Drubin, and Georjana Barnes. Nbl1p: a borealin/dasra/csc-1-like protein essential for aurora/ipl1 complex function and integrity in *Saccharomyces cerevisiae*. *Molecular Biology of the Cell*, 20(6):1772–1784, 2009.

[113] Dan Liu, Gerben Vader, Martijn JM Vromans, Michael A Lampson, and Susanne MA Lens. Sensing chromosome bi-orientation by spatial separation of aurora b kinase from kinetochore substrates. *Science*, 323(5919):1350–1353, 2009.

[114] Ayse Koca Caydasi, Bahtiyar Kurtulmus, Maria IL Orrico, Astrid Hofmann, Bashar Ibrahim, and Gislene Pereira. Elm1 kinase activates the spindle position checkpoint kinase kin4. *The Journal of Cell Biology*, 190(6):975–989, 2010.

[115] Daisuke Watanabe, Mitsuhiro Abe, and Yoshikazu Ohya. Yeast lrg1p acts as a specialized rhogap regulating 1, 3-β-glucan synthesis. *Yeast*, 18(10):943–951, 2001.

[116] Judith Zich and Kevin G Hardwick. Getting down to the phosphorylated 'nuts and bolts' of spindle checkpoint signalling. *Trends in Biochemical Sciences*, 35(1):18–27, 2010.

[117] Robertus AM de Bruin, Tatyana I Kalashnikova, and Curt Wittenberg. Stb1 collaborates with other regulators to modulate the g1-specific transcriptional circuit. *Molecular and Cellular Biology*, 28(22):6919–6928, 2008.

[118] Nazareth Bastajian, Helena Friesen, and Brenda J Andrews. Bck2 acts through the mads box protein mcm1 to activate cell-cycle-regulated genes in budding yeast. *PLoS Genetics*, 9(5):e1003507, 2013.

[119] Luis J García-Rodríguez, Giacomo De Piccoli, Vanessa Marchesi, Richard C Jones, Ricky D Edmondson, and Karim Labib. A conserved polϵ binding module in ctf18-rfc is required for s-phase checkpoint activation downstream of mec1. *Nucleic Acids Research*, 43(18):8830–8838, 2015.

[120] Anna Maria Hegnauer, Nicole Hustedt, Kenji Shimada, Brietta L Pike, Markus Vogel, Philipp Amsler, Seth M Rubin, Fred Van Leeuwen, Aude Guénolé, Haico Van Attikum, et al. An n-terminal acidic region of sgs1 interacts with rpa70 and recruits rad53 kinase to stalled forks. *The EMBO Journal*, 31(18):3768–3783, 2012.

[121] Gloria Palou, Roger Palou, Fanli Zeng, Ajay A Vashisht, James A Wohlschlegel, and David G Quintana. Three different pathways prevent chromosome segregation in the presence of dna damage or replication stress in budding yeast. *PLoS Genetics*, 11(9):e1005468, 2015.

[122] Peter Kaiser, Rey AL Sia, Elaine GS Bardes, Daniel J Lew, and Steven I Reed. Cdc34 and the f-box protein met30 are required for degradation of the cdk-inhibitory kinase swe1. *Genes & Development*, 12(16):2587–2597, 1998.

[123] Jeremy M Rock, Daniel Lim, Lasse Stach, Roksana W Ogrodowicz, Jamie M Keck, Michele H Jones, Catherine CL Wong, John R Yates, Mark Winey, Stephen J Smerdon, et al. Activation of the yeast hippo pathway by phosphorylation-dependent assembly of signaling complexes. *Science*, 340(6134):871–875, 2013.

[124] Anjon Audhya and Scott D Emr. Stt4 pi 4-kinase localizes to the plasma membrane and functions in the pkc1-mediated map kinase cascade. *Developmental Cell*, 2(5):593–605, 2002.

[125] Yuya Yamagishi, Takeshi Sakuno, Yuhei Goto, and Yoshinori Watanabe. Kinetochore composition and its function: lessons from yeasts. *FEMS Microbiology Reviews*, 38(2):185–200, 2014.

[126] Anna Travesa, Tatyana I Kalashnikova, Robertus AM de Bruin, Sarah Rose Cass, Charly Chahwan, David E Lee, Noel F Lowndes, and Curt Wittenberg. Repression of g1/s transcription is mediated via interaction of the gtb motifs of nrm1 and whi5 with swi6. *Molecular and Cellular Biology*, 33(8):1476–1486, 2013.

[127] Anna Travesa, Dwight Kuo, Robertus AM De Bruin, Tatyana I Kalashnikova, Marisela Guaderrama, Kevin Thai, Aaron Aslanian, Marcus B Smolka, John R Yates, Trey Ideker, et al. Dna replication stress differentially regulates g1/s genes via rad53-dependent inactivation of nrm1. *The EMBO Journal*, 31(7):1811–1822, 2012.

[128] Masafumi Nishizawa, Masaoki Kawasumi, Marie Fujino, and Akio Toh-e. Phosphorylation of sic1, a cyclin-dependent kinase (cdk) inhibitor, by cdk including pho85 kinase is required for its prompt degradation. *Molecular Biology of the Cell*, 9(9):2393–2405, 1998.

[129] Dongqing Huang, Supipi Kaluarachchi, Dewald van Dyk, Helena Friesen, Richelle Sopko, Wei Ye, Nazareth Bastajian, Jason Moffat, Holly Sassi, Michael Costanzo, et al. Dual regulation by pairs of cyclin-dependent protein kinases and histone deacetylases controls g1 transcription in budding yeast. *PLoS Biology*, 7(9):e1000188, 2009.

[130] Itamar Simon, John Barnett, Nancy Hannett, Christopher T Harbison, Nicola J Rinaldi, Thomas L Volkert, John J Wyrick, Julia Zeitlinger, David K Gifford, Tommi S Jaakkola, et al. Serial regulation of transcriptional regulators in the yeast cell cycle. *Cell*, 106(6):697–708, 2001.

[131] Tata Pramila, Shawna Miles, Debraj GuhaThakurta, Dave Jemiolo, and Linda L Breeden. Conserved homeodomain proteins interact with mads box protein mcm1 to restrict ecb-dependent transcription to the m/g1 phase of the cell cycle. *Genes & Development*, 16(23):3034–3045, 2002.

[132] Tata Pramila, Wei Wu, Shawna Miles, William Stafford Noble, and Linda L Breeden. The forkhead transcription factor hcm1 regulates chromosome segregation genes and fills the s-phase gap in the transcriptional circuitry of the cell cycle. *Genes & Development*, 20(16):2266–2278, 2006.

[133] Christian Linke, Anastasia Chasapi, Alberto González-Novo, Istabrak Al Sawad, Silvia Tognetti, Edda Klipp, Mart Loog, Sylvia Krobitsch, Francesc Posas, Ioannis Xenarios, et al. A clb/cdk1-mediated regulation of fkh2 synchronizes clb expression in the budding yeast cell cycle. *npj Systems Biology and Applications*, 3:1, 2017.

[134] Kim Nasmyth and Léon Dirick. The role of swi4 and swi6 in the activity of g1 cyclins in yeast. *Cell*, 66(5):995–1013, 1991.

[135] Christine E Horak, Nicholas M Luscombe, Jiang Qian, Paul Bertone, Stacy Piccirrillo, Mark Gerstein, and Michael Snyder. Complex transcriptional circuitry at the g1/s transition in *Saccharomyces cerevisiae*. *Genes & Development*, 16(23):3017–3033, 2002.

[136] James M Bean, Eric D Siggia, and Frederick R Cross. High functional overlap between mlui cell-cycle box binding factor and swi4/6 cell-cycle box binding factor in the g1/s transcriptional program in *Saccharomyces cerevisiae*. *Genetics*, 171(1):49–61, 2005.

[137] Raymond J Cho, Michael J Campbell, Elizabeth A Winzeler, Lars Steinmetz, Andrew Conway, Lisa Wodicka, Tyra G Wolfsberg, Andrei E Gabrielian, David Landsman, David J Lockhart, et al. A genome-wide transcriptional analysis of the mitotic cell cycle. *Molecular Cell*, 2(1):65–73, 1998.

[138] Christine N Tennyson, Jinhwa Lee, and Brenda J Andrews. A role for the pcl9-pho85 cyclin–cdk complex at the m/g1 boundary in *Saccharomyces cerevisiae*. *Molecular Microbiology*, 28(1):69–79, 1998.

[139] Kenzie D MacIsaac, Ting Wang, D Benjamin Gordon, David K Gifford, Gary D Stormo, and Ernest Fraenkel. An improved map of conserved regulatory sites for *Saccharomyces cerevisiae*. *BMC Bioinformatics*, 7(1):113, 2006.

[140] P Zheng, DS Fay, J Burton, HONG Xiao, JL Pinkham, and DF Stern. Spk1 is an essential s-phase-specific gene of *Saccharomyces cerevisiae* that encodes a nuclear serine/threonine/tyrosine kinase. *Molecular and Cellular Biology*, 13(9):5829–5842, 1993.

[141] Dunja Knapp, Leena Bhoite, David J Stillman, and Kim Nasmyth. The transcription factor swi5 regulates expression of the cyclin kinase inhibitor p40sic1. *Molecular and Cellular Biology*, 16(10):5701–5707, 1996.

[142] Anne D Donaldson and John V Kilmartin. Spc42p: a phosphorylated component of the *S. cerevisiae* spindle pole body (spb) with an essential function during spb duplication. *The Journal of Cell Biology*, 132(5):887–901, 1996.

[143] RA Sia, Heather A Herald, and Daniel J Lew. Cdc28 tyrosine phosphorylation and the morphogenesis checkpoint in budding yeast. *Molecular Biology of the Cell*, 7(11):1657–1666, 1996.

[144] Leon Y Chan and Angelika Amon. The protein phosphatase 2a functions in the spindle position checkpoint by regulating the checkpoint kinase kin4. *Genes & Development*, 23(14):1639–1649, 2009.

[145] Frank Uhlmann, Friedrich Lottspeich, and Kim Nasmyth. Sister-chromatid separation at anaphase onset is promoted by cleavage of the cohesin subunit scc1. *Nature*, 400(6739):37, 1999.

[146] Esther Bullitt, Michael P Rout, John V Kilmartin, and Christopher W Akey. The yeast spindle pole body is assembled around a central crystal of spc42p. *Cell*, 89(7):1077–1086, 1997.

[147] Sheng-hong Chen, Marcus B Smolka, and Huilin Zhou. Mechanism of dun1 activation by rad53 phosphorylation in *Saccharomyces cerevisiae*. *Journal of Biological Chemistry*, 282(2):986–995, 2007.

[148] Cristina Iftode, Yaron Daniely, and James A Borowiec. Replication protein a (rpa): the eukaryotic ssb. *Critical Reviews in Biochemistry and Molecular Biology*, 34(3):141–180, 1999.

[149] Olga Chilkova, Peter Stenlund, Isabelle Isoz, Carrie M Stith, Pawel Grabowski, Else-Britt Lundström, Peter M Burgers, and Erik Johansson. The eukaryotic leading and lagging strand dna polymerases are loaded onto primer-ends via separate mechanisms but have comparable processivity in the presence of pcna. *Nucleic Acids Research*, 35(19):6588–6597, 2007.

[150] Mitsuhiro Abe, Hiroshi Qadota, Aiko Hirata, and Yoshikazu Ohya. Lack of gtp-bound rho1p in secretory vesicles of *Saccharomyces cerevisiae*. *The Journal of Cell Biology*, 162(1):85–97, 2003.

[151] Olivier Roumanie, Hao Wu, Jeffrey N Molk, Guendalina Rossi, Kerry Bloom, and Patrick Brennwald. Rho gtpase regulation of exocytosis in yeast is independent of gtp hydrolysis and polarization of the exocyst complex. *The Journal of Cell Biology*, 170(4):583–594, 2005.

[152] Nik C Barbet, Ulrich Schneider, Stephen B Helliwell, Ian Stansfield, Mick F Tuite, and Michael N Hall. Tor controls translation initiation and early g1 progression in yeast. *Molecular Biology of the Cell*, 7(1):25–42, 1996.

[153] Bevin Philip and David E Levin. Wsc1 and mid2 are cell surface sensors for cell wall integrity signaling that act through rom2, a guanine nucleotide exchange factor for rho1. *Molecular and Cellular Biology*, 21(1):271–280, 2001.

[154] Karolina Peplowska, Andreas U Wallek, and Zuzana Storchova. Sgo1 regulates both condensin and ipl1/aurora b to promote chromosome biorientation. *PLoS Genetics*, 10(6):e1004411, 2014.

[155] Sarah Elliott, Michael Knop, Gabriel Schlenstedt, and Elmar Schiebel. Spc29p is a component of the spc110p subcomplex and is essential for spindle pole body duplication. *Proceedings of the National Academy of Sciences*, 96(11):6205–6210, 1999.

[156] Andrew S Lyon, Geneviève Morin, Michelle Moritz, King Clyde B Yabut, Tamira Vojnar, Alex Zelter, Eric Muller, Trisha N Davis, and David A Agard. Higher-order oligomerization of spc110p drives γ-tubulin ring complex assembly. *Molecular Biology of the Cell*, 27(14):2245–2258, 2016.

[157] Mark Winey and Kerry Bloom. Mitotic spindle form and function. *Genetics*, 190(4):1197–1224, 2012.

[158] Christian Koch, Alexander Schleiffer, Gustav Ammerer, and Kim Nasmyth. Switching transcription on and off during the yeast cell cycle: Cln/cdc28 kinases activate bound transcription factor sbf (swi4/swi6) at start, whereas clb/cdc28 kinases displace it from the promoter in g2. *Genes & Development*, 10(2):129–141, 1996.

[159] Thomas Moll, Graham Tebb, Uttam Surana, Helmut Robitsch, and Kim Nasmyth. The role of phosphorylation and the cdc28 protein kinase in cell cycle-regulated nuclear import of the *S. cerevisiae* transcription factor swi5. *Cell*, 66(4):743–758, 1991.

[160] Sarah Erlemann, Annett Neuner, Linda Gombos, Romain Gibeaux, Claude Antony, and Elmar Schiebel. An extended γ-tubulin ring functions as a stable platform in microtubule nucleation. *The Journal of Cell Biology*, 197(1):59–74, 2012.

[161] Satoshi Yoshida, Keiko Kono, Drew M Lowery, Sara Bartolini, Michael B Yaffe, Yoshikazu Ohya, and David Pellman. Polo-like kinase cdc5 controls the local activation of rho1 to promote cytokinesis. *Science*, 313(5783):108–111, 2006.

[162] J Michael Dial, Evgeniy V Petrotchenko, and Christoph H Borchers. Inhibition of apccdh1 activity by cdh1/acm1/bmh1 ternary complex formation. *Journal of Biological Chemistry*, 282(8):5237–5248, 2007.

[163] Beidong Liu, Lisa Larsson, Antonio Caballero, Xinxin Hao, David Öling, Julie Grantham, and Thomas Nyström. The polarisome is required for segregation and retrograde transport of protein aggregates. *Cell*, 140(2):257–267, 2010.

[164] Scott C Schuyler, Jenny Y Liu, and David Pellman. The molecular function of ase1p. *The Journal of Cell Biology*, 160(4):517–528, 2003.

[165] Pil Jung Kang, Mid Eum Lee, and Hay-Oak Park. Bud3 activates cdc42 to establish a proper growth site in budding yeast. *The Journal of Cell Biology*, 206(1):19–28, 2014.

[166] Indrani Bose, Javier E Irazoqui, John J Moskow, Elaine SG Bardes, Trevin R Zyla, and Daniel J Lew. Assembly of scaffold-mediated complexes containing cdc42p, the exchange factor cdc24p, and the effector cla4p required for cell cycle-regulated phosphorylation of cdc24p. *Journal of Biological Chemistry*, 276(10):7176–7186, 2001.

[167] Sarah E Lee, Sanne Jensen, Lisa M Frenz, Anthony L Johnson, Didier Fesquet, and Leland H Johnston. The bub2-dependent mitotic pathway in yeast acts every cell cycle and regulates cytokinesis. *Journal of Cell Science*, 114(12):2345–2354, 2001.

[168] Marco Geymonat, Adonis Spanos, Geoffroy de Bettignies, and Steven G Sedgwick. Lte1 contributes to bfa1 localization rather than stimulating nucleotide exchange by tem1. *The Journal of Cell Biology*, 187(4):497–511, 2009.

[169] Hye-Joo Yoon and John Carbon. Participation of bir1p, a member of the inhibitor of apoptosis family, in yeast chromosome segregation events. *Proceedings of the National Academy of Sciences*, 96(23):13208–13213, 1999.

[170] Arthur S Alberts. Identification of a carboxyl-terminal diaphanous-related formin homology protein autoregulatory domain. *Journal of Biological Chemistry*, 276(4):2824–2830, 2001.

[171] Yuqing Dong, David Pruyne, and Anthony Bretscher. Formin-dependent actin assembly is regulated by distinct modes of rho signaling in yeast. *The Journal of cell biology*, 161(6):1081–1092, 2003.

[172] Yingwu Xu, James B Moseley, Isabelle Sagot, Florence Poy, David Pellman, Bruce L Goode, and Michael J Eck. Crystal structures of a formin homology-2 domain reveal a tethered dimer architecture. *Cell*, 116(5):711–723, 2004.

[173] Takeshi Fujiwara, Kazuma Tanaka, Akihisa Mino, Mitsuhiro Kikyo, Kazuo Takahashi, Kazuya Shimizu, and Yoshimi Takai. Rho1p-bni1p-spa2p interactions: Implication in localization of bni1p at the bud site and regulation of the actin cytoskeleton in *Saccharomyces cerevisiae*. *Molecular Biology of the Cell*, 9(5):1221–1233, 1998.

[174] Marco Geymonat, Ad Spanos, Susan JM Smith, Edward Wheatley, Katrin Rittinger, Leland H Johnston, and Steven G Sedgwick. Control of mitotic exit in budding yeast *In vitro* regulation of tem1 gtpase by bub2 and bfa1. *Journal of Biological Chemistry*, 277(32):28439–28445, 2002.

[175] Aurelie Bertin, Michael A McMurray, Patricia Grob, Sang-Shin Park, Galo Garcia, Insiyyah Patanwala, Ho-leung Ng, Tom Alber, Jeremy Thorner, and Eva Nogales. *Saccharomyces cerevisiae* septins: supramolecular organization of heterooligomers and the mechanism of filament assembly. *Proceedings of the National Academy of Sciences*, 105(24):8274–8279, 2008.

[176] Rosella Visintin, Karen Craig, Ellen S Hwang, Susanne Prinz, Mike Tyers, and Angelika Amon. The phosphatase cdc14 triggers mitotic exit by reversal of cdk-dependent phosphorylation. *Molecular Cell*, 2(6):709–718, 1998.

[177] Aljoscha Nern and Robert A Arkowitz. A cdc24p-far1p-g$\beta\gamma$ protein complex required for yeast orientation during mating. *The Journal of Cell Biology*, 144(6):1187–1202, 1999.

[178] Yi Zheng, Richard Cerione, and Alan Bender. Control of the yeast bud-site assembly gtpase cdc42. catalysis of guanine nucleotide exchange by cdc24 and stimulation of gtpase activity by bem3. *Journal of Biological Chemistry*, 269(4):2369–2372, 1994.

[179] JA Hadwiger, C Wittenberg, MD Mendenhall, and SI Reed. The *Saccharomyces cerevisiae* cks1 gene, a homolog of the *Schizosaccharomyces pombe* suc1+ gene, encodes a subunit of the cdc28 protein kinase complex. *Molecular and Cellular Biology*, 9(5):2034–2041, 1989.

[180] Julia Yaglom, MH Linskens, Seth Sadis, David M Rubin, Bruce Futcher, and Daniel Finley. p34cdc28-mediated control of cln3 cyclin degradation. *Molecular and Cellular Biology*, 15(2):731–741, 1995.

[181] Ritu Agarwal and Orna Cohen-Fix. Phosphorylation of the mitotic regulator pds1/securin by cdc28 is required for efficient nuclear localization of esp1/separase. *Genes & Development*, 16(11):1371–1382, 2002.

[182] Kyung S Lee, Satoshi Asano, Jung-Eun Park, Krisada Sakchaisri, and Raymond L Erikson. Monitoring the cell cycle by multi-kinase-dependent regulation of swe1/wee1 in budding yeast. *Cell Cycle*, 4(10):1346–1349, 2005.

[183] Hongyin Wang, Eloi Garí, Emili Verges, Carme Gallego, and Martí Aldea. Recruitment of cdc28 by whi3 restricts nuclear accumulation of the g1 cyclin–cdk complex to late g1. *The EMBO Journal*, 23(1):180–190, 2004.

[184] Jae Hong Seol, RM Renny Feldman, Wolfgang Zachariae, Anna Shevchenko, Craig C Correll, Svetlana Lyapina, Yong Chi, Marta Galova, Jonathan Claypool, Suzanne Sandmeyer, et al. Cdc53/cullin and the essential hrt1 ring–h2 subunit of scf define a ubiquitin ligase module that activates the e2 enzyme cdc34. *Genes & Development*, 13(12):1614–1626, 1999.

[185] Gregory R Hoffman, Nicolas Nassar, and Richard A Cerione. Structure of the rho family gtp-binding protein cdc42 in complex with the multifunctional regulator rhogdi. *Cell*, 100(3):345–356, 2000.

[186] Dorota Skowyra, Karen L Craig, Mike Tyers, Stephen J Elledge, and J Wade Harper. F-box proteins are receptors that recruit phosphorylated substrates to the scf ubiquitin-ligase complex. *Cell*, 91(2):209–219, 1997.

[187] E Elizabeth Patton, Andrew R Willems, Danne Sa, Laurent Kuras, Dominique Thomas, Karen L Craig, and Mike Tyers. Cdc53 is a scaffold protein for multiple cdc34/skp1/f-box protein complexes that regulate cell division and methionine biosynthesis in yeast. *Genes & Development*, 12(5):692–705, 1998.

[188] Roni Koren, Liat Rainis, and Tamar Kleinberger. The scaffolding a/tpd3 subunit and high phosphatase activity are dispensable for cdc55 function in the *Saccharomyces cerevisiae* spindle checkpoint and in cytokinesis. *Journal of Biological Chemistry*, 279(47):48598–48606, 2004.

[189] Sukgil Song and Kyung S Lee. A novel function of *Saccharomyces cerevisiae* cdc5 in cytokinesis. *The Journal of Cell Biology*, 152(3):451–470, 2001.

[190] Junwon Kim, Guangming Luo, Young Yil Bahk, and Kiwon Song. Cdc5-dependent asymmetric localization of bfa1 fine-tunes timely mitotic exit. *PLoS Genetics*, 8(1):e1002450, 2012.

[191] Wenying Shou, Ramzi Azzam, Susan L Chen, Michael J Huddleston, Christopher Baskerville, Harry Charbonneau, Roland S Annan, Steve A Carr, and Raymond J Deshaies. Cdc5 influences phosphorylation of net1 and disassembly of the rent complex. *BMC Molecular Biology*, 3(1):3, 2002.

[192] Luo Feng, Bin Wang, Lingtao Wu, and Ambrose Jong. Loss control of mcm5 interaction with chromatin in cdc6-1 mutated in cdc-ntp motif. *DNA and Cell Biology*, 19(7):447–457, 2000.

[193] Hong Wang, Dou Liu, Yanchang Wang, Jun Qin, and Stephen J Elledge. Pds1 phosphorylation in response to dna damage is essential for its dna damage checkpoint function. *Genes & Development*, 15(11):1361–1372, 2001.

[194] Emily R Hildebrandt, Larisa Gheber, Tami Kingsbury, and M Andrew Hoyt. Homotetrameric form of cin8p, a *Saccharomyces cerevisiae* kinesin-5 motor, is essential for its in vivo function. *Journal of Biological Chemistry*, 281(36):26004–26013, 2006.

[195] Robert F Siegmund and Kim A Nasmyth. The *Saccharomyces cerevisiae* start-specific transcription factor swi4 interacts through the ankyrin repeats with the mitotic clb2/cdc28 kinase and through its conserved carboxy terminus with swi6. *Molecular and Cellular Biology*, 16(6):2647–2655, 1996.

[196] Eloi Garí, Tom Volpe, Hongyin Wang, Carme Gallego, Bruce Futcher, and Martí Aldea. Whi3 binds the mrna of the g1 cyclin cln3 to modulate cell fate in budding yeast. *Genes & Development*, 15(21):2803–2808, 2001.

[197] Laure Crabbé, Aubin Thomas, Véronique Pantesco, John De Vos, Philippe Pasero, and Armelle Lengronne. Analysis of replication profiles reveals key role of rfc-ctf18 in yeast replication stress response. *Nature Structural & Molecular Biology*, 17(11):1391–1397, 2010.

[198] Svetlana I Komarnitsky, Yueh-Chin Chiang, Francis C Luca, Junji Chen, Jeremy H Toyn, Mark Winey, Leland H Johnston, and Clyde L Denis. Dbf2 protein kinase binds to and acts through the cell cycle-regulated mob1 protein. *Molecular and Cellular Biology*, 18(4):2100–2107, 1998.

[199] Franz Meitinger, Saravanan Palani, Birgit Hub, and Gislene Pereira. Dual function of the ndr-kinase dbf2 in the regulation of the f-bar protein hof1 during cytokinesis. *Molecular Biology of the Cell*, 24(9):1290–1304, 2013.

[200] Seiji Tanaka, Yayoi Komeda, Toshiko Umemori, Yumiko Kubota, Haruhiko Takisawa, and Hiroyuki Araki. Efficient initiation of dna replication in eukaryotes requires dpb11/topbp1-gins interaction. *Molecular and Cellular Biology*, 33(13):2614–2622, 2013.

[201] Hiroshi Masumoto, Akio Sugino, and Hiroyuki Araki. Dpb11 controls the association between dna polymerases α and ϵ and the autonomously replicating sequence region of budding yeast. *Molecular and Cellular Biology*, 20(8):2809–2817, 2000.

[202] Leigh Francisco, Wenfang Wang, and CS Chan. Type 1 protein phosphatase acts in opposition to ipl1 protein kinase in regulating yeast chromosome segregation. *Molecular and Cellular Biology*, 14(7):4731–4740, 1994.

[203] Bryce Nelson, Cornelia Kurischko, Joe Horecka, Manali Mody, Pradeep Nair, Lana Pratt, Alexandre Zougman, Linda DB McBroom, Timothy R Hughes, Charlie Boone, et al. Ram: a conserved signaling network that regulates ace2p transcriptional activity and polarized morphogenesis. *Molecular Biology of the Cell*, 14(9):3782–3803, 2003.

[204] Leslie E Stolz, Winnie J Kuo, Jason Longchamps, Mandeep K Sekhon, and John D York. Inp51, a yeast inositol polyphosphate 5-phosphatase required for phosphatidylinositol 4, 5-bisphosphate homeostasis and whose absence confers a cold-resistant phenotype. *Journal of Biological Chemistry*, 273(19):11852–11861, 1998.

[205] Jung-seog Kang, Iain M Cheeseman, George Kallstrom, Soundarapandian Velmurugan, Georjana Barnes, and Clarence SM Chan. Functional cooperation of dam1, ipl1, and the inner centromere protein (incenp)–related protein sli15 during chromosome segregation. *The Journal of Cell Biology*, 155(5):763–774, 2001.

[206] Hongwei Yin, David Pruyne, Tim C Huffaker, and Anthony Bretscher. Myosin v orientates the mitotic spindle in yeast. *Nature*, 406(6799):1013, 2000.

[207] Hiromi Maekawa, Claire Priest, Johannes Lechner, Gislene Pereira, and Elmar Schiebel. The yeast centrosome translates the positional information of the anaphase spindle into a cell cycle signal. *The Journal of Cell Biology*, 179(3):423–436, 2007.

[208] Vladimir Fridman, Adina Gerson-Gurwitz, Ofer Shapira, Natalia Movshovich, Stefan Lakämper, Christoph F Schmidt, and Larisa Gheber. Kinesin-5 kip1 is a bi-directional motor that stabilizes microtubules and tracks their plus-ends in vivo. *J Cell Sci*, 126(18):4147–4159, 2013.

[209] Michael Glotzer. The 3ms of central spindle assembly: microtubules, motors and maps. *Nature Reviews. Molecular Cell Biology*, 10(1):9, 2009.

[210] Mark K Chee and Steven B Haase. B-cyclin/cdks regulate mitotic spindle assembly by phosphorylating kinesins-5 in budding yeast. *PLoS Genetics*, 6(5):e1000935, 2010.

[211] Tatsushi Wakayama, Tae Kondo, Seiko Ando, Kunihiro Matsumoto, and Katsunori Sugimoto. Pie1, a protein interacting with mec1, controls cell growth and checkpoint responses in *Saccharomyces cerevisiae*. *Molecular and Cellular Biology*, 21(3):755–764, 2001.

[212] John Rouse and Stephen P Jackson. Lcd1p recruits mec1p to dna lesions in vitro and in vivo. *Molecular Cell*, 9(4):857–869, 2002.

[213] Annette A Alcasabas, Alexander J Osborn, Jeff Bachant, Fenghua Hu, Petra JH Werler, Kristine Bousset, Kanji Furuya, John FX Diffley, Antony M Carr, and Stephen J Elledge. Mrc1 transduces signals of dna replication stress to activate rad53. *Nature Cell Biology*, 3(11):958–965, 2001.

[214] Huiqiang Lou, Makiko Komata, Yuki Katou, Zhiyun Guan, Clara C Reis, Martin Budd, Katsuhiko Shirahige, and Judith L Campbell. Mrc1 and dna polymerase ϵ function together in linking dna replication and the s phase checkpoint. *Molecular Cell*, 32(1):106–117, 2008.

[215] Xiang-Dong Gao, Stefan Albert, Serguei E Tcheperegine, Christopher G Burd, Dieter Gallwitz, and Erfei Bi. The gap activity of msb3p and msb4p for the rab gtpase sec4p is required for efficient exocytosis and actin organization. *The Journal of Cell Biology*, 162(4):635–646, 2003.

[216] Ulrike Gruneberg, Kirsteen Campbell, Clare Simpson, Joan Grindlay, and Elmar Schiebel. Nud1p links astral microtubule organization and the control of exit from mitosis. *The EMBO Journal*, 19(23):6475–6488, 2000.

[217] Mauricio Valerio-Santiago and Fernando Monje-Casas. Tem1 localization to the spindle pole bodies is essential for mitotic exit and impairs spindle checkpoint function. *The Journal of Cell Biology*, pages jcb–201007044, 2011.

[218] Philipp Müller, Sookhee Park, Erika Shor, Dana J Huebert, Christopher L Warren, Aseem Z Ansari, Michael Weinreich, Matthew L Eaton, David M MacAlpine, and Catherine A Fox. The conserved bromo-adjacent homology domain of yeast orc1 functions in the selection of dna replication origins within chromatin. *Genes & Development*, 24(13):1418–1433, 2010.

[219] Jingchuan Sun, Hironori Kawakami, Juergen Zech, Christian Speck, Bruce Stillman, and Huilin Li. Cdc6-induced conformational changes in orc bound to origin dna revealed by cryo-electron microscopy. *Structure*, 20(3):534–544, 2012.

[220] V Measday, L Moore, R Retnakaran, J Lee, M Donoviel, AM Neiman, and B Andrews. A family of cyclin-like proteins that interact with the pho85 cyclin-dependent kinase. *Molecular and Cellular Biology*, 17(3):1212–1223, 1997.

[221] Yue-Li Juang, James Huang, Jan-Michael Peters, Margaret E McLaughlin, Chin-Yin Tai, and David Pellman. Apc-mediated proteolysis of ase1 and the morphogenesis of the mitotic spindle. *Science*, 275(5304):1311–1314, 1997.

[222] Paul T Spellman, Gavin Sherlock, Michael Q Zhang, Vishwanath R Iyer, Kirk Anders, Michael B Eisen, Patrick O Brown, David Botstein, and Bruce Futcher. Comprehensive identification of cell cycle–regulated genes of the yeast *Saccharomyces cerevisiae* by microarray hybridization. *Molecular Biology of the Cell*, 9(12):3273–3297, 1998.

[223] Olivier Poch, Etienne Schwob, Florence de Fraipont, Alain Camasses, Rémy Bordonné, and Robert P Martin. Rpk1, an essential yeast protein kinase involved in the regulation of the onset of mitosis, shows homology to mammalian dual-specificity kinases. *Molecular and General Genetics MGG*, 243(6):641–653, 1994.

[224] Stefan Kemmler, Manuel Stach, Maria Knapp, Jennifer Ortiz, Jens Pfannstiel, Thomas Ruppert, and Johannes Lechner. Mimicking ndc80 phosphorylation triggers spindle assembly checkpoint signalling. *The EMBO Journal*, 28(8):1099–1110, 2009.

[225] Yu Jiang and James R Broach. Tor proteins and protein phosphatase 2a reciprocally regulate tap42 in controlling cell growth in yeast. *The EMBO Journal*, 18(10):2782–2792, 1999.

[226] Benjamin D Landry, Claudine E Mapa, Heather E Arsenault, Kristin E Poti, and Jennifer A Benanti. Regulation of a transcription factor network by cdk1 coordinates late cell cycle gene expression. *The EMBO Journal*, 33(9):1044–1060, 2014.

[227] Aiyang Cheng, Karen E Ross, Philipp Kaldis, and Mark J Solomon. Dephosphorylation of cyclin-dependent kinases by type 2c protein phosphatases. *Genes & Development*, 13(22):2946–2957, 1999.

[228] Ying-Chou Chen, Jessica Kenworthy, Carrie Gabrielse, Christine Hänni, Philip Zegerman, and Michael Weinreich. Dna replication checkpoint signaling depends on a rad53–dbf4 n-terminal interaction in *Saccharomyces cerevisiae*. *Genetics*, 194(2):389–401, 2013.

[229] Vladimir I Bashkirov, Elena V Bashkirova, Edwin Haghnazari, and Wolf-Dietrich Heyer. Direct kinase-to-kinase signaling mediated by the fha phosphoprotein recognition domain of the dun1 dna damage checkpoint kinase. *Molecular and Cellular Biology*, 23(4):1441–1452, 2003.

[230] Jaime Lopez-Mosqueda, Nancy L Maas, Zophonias O Jonsson, Lisa G DeFazio Eli, James Wohlschlegel, and David P Toczyski. Damage-induced phosphorylation of sld3 is important to block late origin firing. *Nature*, 467(7314):479, 2010.

[231] Steven J Brill and Suzanne Bastin-Shanower. Identification and characterization of the fourth single-stranded-dna binding domain of replication protein a. *Molecular and Cellular Biology*, 18(12):7225–7234, 1998.

[232] Doranelly Philipova, Janet R Mullen, Hina S Maniar, Jian Lu, Chunyan Gu, and Steven J Brill. A hierarchy of ssb protomers in replication protein a. *Genes & Development*, 10(17):2222–2233, 1996.

[233] Wei Guo, Fuyuhiko Tamanoi, and Peter Novick. Spatial regulation of the exocyst complex by rho1 gtpase. *Nature Cell Biology*, 3(4):353–360, 2001.

[234] Xiaolan Zhao, Eric GD Muller, and Rodney Rothstein. A suppressor of two essential checkpoint genes identifies a novel protein that negatively affects dntp pools. *Molecular Cell*, 2(3):329–340, 1998.

[235] K Ozaki, K Tanaka, H Imamura, T Hihara, T Kameyama, H Nonaka, H Hirano, Y Matsuura, and Y Takai. Rom1p and rom2p are gdp/gtp exchange proteins (geps) for the rho1p small gtp binding protein in *Saccharomyces cerevisiae*. *The EMBO Journal*, 15(9):2196, 1996.

[236] Shinya Takahata, Yaxin Yu, and David J Stillman. The e2f functional analogue sbf recruits the rpd3 (l) hdac, via whi5 and stb1, and the fact chromatin reorganizer, to yeast g1 cyclin promoters. *The EMBO Journal*, 28(21):3378–3389, 2009.

[237] Christian G Riedel, Vittorio L Katis, Yuki Katou, Saori Mori, Takehiko Itoh, Wolfgang Helmhart, Marta Gálová, Mark Petronczki, Juraj Gregan, Bulent Cetin, et al. Protein phosphatase 2a protects centromeric sister chromatid cohesion during meiosis i. *Nature*, 441(7089):53–61, 2006.

[238] Michelangelo Foti, Anjon Audhya, and Scott D Emr. Sac1 lipid phosphatase and stt4 phosphatidylinositol 4-kinase regulate a pool of phosphatidylinositol 4-phosphate that functions in the control of the actin cytoskeleton and vacuole morphology. *Molecular Biology of the Cell*, 12(8):2396–2411, 2001.

[239] Josefin Fernius, Olga O Nerusheva, Stefan Galander, Flavia de Lima Alves, Juri Rappsilber, and Adele L Marston. Cohesin-dependent association of scc2/4 with the centromere initiates pericentromeric cohesion establishment. *Current Biology*, 23(7):599–606, 2013.

[240] Stephen M Hinshaw, Vasso Makrantoni, Alastair Kerr, Adèle L Marston, and Stephen C Harrison. Structural evidence for scc4-dependent localization of cohesin loading. *Elife*, 4:e06057, 2015.

[241] Avrom J Caplan, Douglas M Cyr, and Michael G Douglas. Ydj1p facilitates polypeptide translocation across different intracellular membranes by a conserved mechanism. *Cell*, 71(7):1143–1155, 1992.

[242] Satoshi Yoshida, Yoshikazu Ohya, Mark Goebl, Akihiko Nakano, and Yasuhiro Anraku. A novel gene, stt4, encodes a phosphatidylinositol 4-kinase in the pkc1 protein kinase pathway of *Saccharomyces cerevisiae*. *Journal of Biological Chemistry*, 269(2):1166–1172, 1994.

[243] Eva Nogales, Michael Whittaker, Ronald A Milligan, and Kenneth H Downing. High-resolution model of the microtubule. *Cell*, 96(1):79–88, 1999.

[244] Tobias Schmelzle, Stephen B Helliwell, and Michael N Hall. Yeast protein kinases and the rho1 exchange factor tus1 are novel components of the cell integrity pathway in yeast. *Molecular and Cellular Biology*, 22(5):1329–1339, 2002.

[245] G Paul H Heusden, Dominic JF Griffiths, Jon C Ford, Peter AT Schrader, Anthony M Carr, H Yde Steensma, et al. The 14-3-3 proteins encoded by the bmh1 and bmh2 genes are essential in the yeast *Saccharomyces cerevisiae* and can be replaced by a plant homologue. *The FEBS Journal*, 229(1):45–53, 1995.

[246] Uttam Surana, Helmut Robitsch, Clive Price, Tillman Schuster, Ian Fitch, A Bruce Futcher, and Kim Nasmyth. The role of cdc28 and cyclins during mitosis in the budding yeast *S. cerevisiae*. *Cell*, 65(1):145–161, 1991.

[247] Mignon A Keaton, Elaine SG Bardes, Aron R Marquitz, Christopher D Freel, Trevin R Zyla, Johannes Rudolph, and Daniel J Lew. Differential susceptibility of yeast s and m phase cdk complexes to inhibitory tyrosine phosphorylation. *Current Biology*, 17(14):1181–1189, 2007.

[248] Charles B Epstein and Frederick R Cross. Clb5: a novel b cyclin from budding yeast with a role in s phase. *Genes & Development*, 6(9):1695–1706, 1992.

[249] E Schwob and K Nasmyth. Clb5 and clb6, a new pair of b cyclins involved in s phase and mitotic spindle formation in *S. cerevisiae*. *Genes & Development*, 7:1160–1175, 1993.

[250] Gregory J Reynard, William Reynolds, Rati Verma, and Raymond J Deshaies. Cks1 is required for g1cyclin–cyclin-dependent kinase activity in budding yeast. *Molecular and Cellular Biology*, 20(16):5858–5864, 2000.

[251] Helena E Richardson, Curt Wittenberg, Fred Cross, and Steven I Reed. An essential g1 function for cyclin-like proteins in yeast. *Cell*, 59(6):1127–1133, 1989.

[252] Emili Vergés, Neus Colomina, Eloi Garí, Carme Gallego, and Martí Aldea. Cyclin cln3 is retained at the er and released by the j chaperone ydj1 in late g1 to trigger cell cycle entry. *Molecular Cell*, 26(5):649–662, 2007.

[253] Elena Nazarova, Eileen O'Toole, Susi Kaitna, Paul Francois, Mark Winey, and Jackie Vogel. Distinct roles for antiparallel microtubule pairing and overlap during early spindle assembly. *Molecular Biology of the Cell*, 24(20):3238–3250, 2013.

[254] Annelise Sahin, Bertrand Daignan-Fornier, and Isabelle Sagot. Polarized growth in the absence of f-actin in *Saccharomyces cerevisiae* exiting quiescence. *PLoS One*, 3(7):e2556, 2008.

[255] Pasquale Palumbo, Marco Vanoni, Valerio Cusimano, Stefano Busti, Francesca Marano, Costanzo Manes, and Lilia Alberghina. Whi5 phosphorylation embedded in the g1/s network dynamically controls critical cell size and cell fate. *Nature Communications*, 7, 2016.

[256] Robertus AM de Bruin, Tatyana I Kalashnikova, Charly Chahwan, W Hayes McDonald, James Wohlschlegel, John Yates, Paul Russell, and Curt Wittenberg. Constraining g1-specific transcription to late g1 phase: the mbf-associated corepressor nrm1 acts via negative feedback. *Molecular cell*, 23(4):483–496, 2006.

[257] Youmin Shu, Haifeng Yang, Elizabeth Hallberg, and Richard Hallberg. Molecular genetic analysis of rts1p, a b'regulatory subunit of *Saccharomyces cerevisiae* protein phosphatase 2a. *Molecular and Cellular Biology*, 17(6):3242–3253, 1997.

[258] Noriko Kantake, Tomohiko Sugiyama, Richard D Kolodner, and Stephen C Kowalczykowski. The recombination-deficient mutant rpa (rfa1-t11) is displaced slowly from single-stranded dna by rad51 protein. *Journal of Biological Chemistry*, 278(26):23410–23417, 2003.

Table S3

Mutant	Prediction	Arrest point (Fig 4B)	DNA	Nuclei
bub2	viable	N/A		
cdc28Y19F	viable	N/A		
CDC5	viable	N/A		
cdc55	viable	N/A		
clb1	viable	N/A		
clb2	viable	N/A		
clb6	viable	N/A		
cln1	viable	N/A		
cln1cln2	viable	N/A		
cln2	viable	N/A		
cln3bck2whi5	viable	N/A		
cln3whi5	viable	N/A		
lte1	viable	N/A		
spc1102AMps1	viable	N/A		
ssa1	viable	N/A		
whi5	viable	N/A		
whi512A	viable	N/A		
whi512Acln3	viable	N/A		
ydj1	viable	N/A		
cdc28	inviable	1a	1	1
cdc28T169A	inviable	1a	1	1
cln1cln2cln3	inviable	1a	1	1
cln1cln2cln3cdh1	inviable	1a	1	1
cln1cln2cln3sic1	inviable	1a	1	1
cln1cln2cln3whi5	inviable	1a	1	1
cln3	inviable	1a	1	1
cln3bck2	inviable	1a	1	1
cln3bck2sic1	inviable	1a	1	1
cln3mbp1	inviable	1a	1	1
cln3swi4	inviable	1a	1	1
cln3swi6	inviable	1a	1	1
whi54Aswi6S4A	inviable	1a	1	1
bck2swi6sic1	inviable	1b	1	1
cdc20pds1cdh1SIC1	inviable	1b	1	1
cdc20pds1clb5	inviable	1b	1	1
cdc20pds1clb5cdh1	inviable	1b	1	1
cdc20pds1clb5cdh1SIC1	inviable	1b	1	1
cdc34	inviable	1b	1	1
cdc4	inviable	1b	1	1
cdc53	inviable	1b	1	1
cdc7	inviable	1b	1	1
clb5	inviable	1b	1	1
clb5clb6	inviable	1b	1	1
cln1cln2swi4	inviable	1b	1	1
mbp1	inviable	1b	1	1
Sic17A	inviable	1b	1	1
Sld2A	inviable	1b	1	1
Sld32A	inviable	1b	1	1
Sld3S622A	inviable	1b	1	1
Sld3T600A	inviable	1b	1	1
spc1102ACDK	inviable	1b	1	1
swi4	inviable	1b	1	1
swi4mbp1	inviable	1b	1	1
swi4mbp1BCK2	inviable	1b	1	1
swi4swi6	inviable	1b	1	1
swi4swi6sic1	inviable	1b	1	1
swi6	inviable	1b	1	1
swi64A	inviable	1b	1	1
whi512Aswi6S4A	inviable	1b	1	1
cdh1	inviable	1c	1	1
cln1cln2cdh1	inviable	1c	1	1
clb1clb2	inviable	3a	1	1
NOC_CDC5	inviable	3a	1	1
HU_CDC5	inviable	3a	1	1
cdc31	inviable	3c		
spc1104A	inviable	3d	1	1
cdc10	inviable	3e	1	1
cdc11	inviable	3e	1	1
cdc12	inviable	3e	1	1
cdc24	inviable	3e	1	1
cdc3	inviable	3e	1	1
cdc42	inviable	3e	1	1
cdc14	inviable	4	1	1
cdc14SIC1	inviable	4	1	1
cdc20	inviable	4	1	1
cdc5	inviable	4	1	1
esp1	inviable	4		
cdc20pds1	inviable	5	1	1
pds1	inviable	5	2	2
cdc6	inviable	6	1	1
net1cdc15cdh1	inviable	6	2	2
cdc15	inviable	7	*	*
cdc15net1	inviable	7	*	*
tem1	inviable	7	*	*
tem1net1	inviable	7	*	*