A major locus for ivermectin resistance in a parasitic nematode

Stephen R. Doyle[1]*, Christopher J. R. Illingworth[2]*, Roz Laing[3], David J. Bartley[4],

Elizabeth Redman[5], Axel Martinelli[1+], Nancy Holroyd[1], Alison A. Morrison[4], Andrew

Rezansoff[5], Alan Tracey[1], Eileen Devaney[3], Matthew Berriman[1], Neil Sargison[6],

James A. Cotton[1**], John S. Gilleard[5**]

* Joint first authors

** Authors for correspondence

+ Current affiliation: Global Station for Zoonosis Control, Global Institution for Collaborative Research
and Education (GI-CoRE), Hokkaido University, Sapporo, Japan, Biological and Environmental
Sciences and Engineering (BESE) Division, King Abdullah University of Science and Technology
(KAUST), Thuwal, Saudi Arabia

1. Wellcome Sanger Institute, Hinxton, Cambridgeshire, CB10 1SA, United Kingdom
Stephen R. Doyle: sd21@sanger.ac.uk
Alan Tracey: alt@sanger.ac.uk
James A. Cotton: jc17@sanger.ac.uk
Nancy Holroyd: neh@sanger.ac.uk
Matthew Berriman: mb4@sanger.ac.uk
Axel Martinelli: axel.martinelli@gmail.com

2. Department of Genetics, University of Cambridge, Downing Street, Cambridge CB2 1EH,
United Kingdom
Chris J. R. Illingworth: cjri2@cam.ac.uk
(also Department of Applied Maths and Theoretical Physics, Wilberforce Road, Cambridge CB3 0WA)

3. Institute of Biodiversity Animal Health and Comparative Medicine, College of Medical,
Veterinary and Life Sciences, University of Glasgow, Garscube Campus, Glasgow, G61 1QH,
United Kingdom
Roz Laing: Rosalind.Laing@glasgow.ac.uk
Eileen Devaney: Eileen.Devaney@glasgow.ac.uk

4. Moredun Research Institute, Pentlands Science Park, Bush Loan, Penicuik EH26 0PZ, United
Kingdom
David J. Bartley: Dave.Bartley@moredun.ac.uk

Alison A. Morrison: Alison.Morrison@moredun.ac.uk

5. Department of Comparative Biology and Experimental Medicine, Faculty of Veterinary

Medicine, University of Calgary, Calgary, Alberta, Canada

John S. Gilleard: jsgillea@ucalgary.ca

Elizabeth Redman: elmredma@ucalgary.ca

Andrew Rezansoff: amrezans@ucalgary.ca

6. University of Edinburgh, Royal (Dick) School of Veterinary Studies, Edinburgh, EH25 9RG,

United Kingdom

Neil Sargison: Neil.Sargison@ed.ac.uk

Abstract

Helminth infections cause an enormous disease burden in humans, livestock and

companion animals. Helminth control is heavily dependent on the regular

administration of anthelmintic drugs. Resistance to these drugs is widespread in

many livestock systems, and there is concern that mass drug administration

programs to control human helminths may be selecting for resistance. Here, we build

on two previous genetic crosses between ivermectin resistant and sensitive isolates

of the parasitic nematode *Haemonchus contortus*, an economically important

gastrointestinal parasite of small ruminants that is a model for anthelmintic research.

By combining genetic mapping and whole genome sequencing with novel analyses,

we identify the first single genomic locus conclusively associated with ivermectin

resistance in a parasitic nematode. This locus is common between two

geographically and genetically divergent strains and does not include leading

candidate genes. This work represents the first example of a major anthelmintic

resistance locus being identified by genetic mapping in a parasitic nematode and

represents the most comprehensive genetic analysis of this trait in a parasitic

nematode to date.

Introduction

Helminths are an extremely diverse group of parasitic organisms, many of which are responsible for significant morbidity and mortality in their hosts. Control of those considered to be of human and veterinary importance is heavily dependent on the large-scale administration of anthelmintic drugs. Anthelmintic drugs such as the macrocyclic lactone, ivermectin, have been extremely successful in the control of a number of helminths in both humans and animals (Omura and Crump, 2004). The importance of the discovery and development of this drug as an anthelmintic was recently recognised by the award of the 2015 Nobel Prize for Medicine or Physiology. However, these successes are now threatened by the emergence of drug resistance. Anthelmintic drug resistance is already widespread in many species of parasitic nematodes of livestock and rapid acquisition of resistance has been documented against single and multiple drug classes (Kaplan and Vidyashankar, 2012; Rose et al., 2015), and is already a major problem for global agricultural production and animal welfare. There are growing concerns regarding reduced efficacy of compounds used in mass drug administration (MDA) programs for some human helminths, which could herald the emergence of resistance (Crellen et al., 2016; Doyle et al., 2017; Geerts and Gryseels, 2000; Osei-Atweneboana et al., 2011). Consequently, anthelmintic drug resistance is an emerging threat to the sustainable control of human parasites, particularly those targeted by MDA programs. However, in spite of the importance of this issue, remarkably little is known regarding the molecular mechanisms of resistance to most anthelmintic drug groups, with the notable exception of the benzimidazole class. This is, at least in part, due to a lack of

genomic resources, tools and techniques with which to study these experimentally

challenging organisms.

Haemonchus contortus is a parasite of wild and domesticated ruminants of global

distribution, and causes major health and economic impacts in sheep and goats

worldwide. This parasite has a propensity to develop resistance to anthelmintic

drugs, with resistance to almost all of the classes of anthelmintic drugs having been

documented in multiple regions of the world (Yadav et al. 1995; Jabbar et al. 2013;

Echevarria and Trindade 1989; Van den Brom et al. 2015; Howell et al. 2008). *H.*

contortus isolates that are resistant to multiple classes simultaneously are common

and resistance can occur within a just a few years of introduction of a new drug class

(Williamson et al., 2011; van Wyk and Malan, 1988). Partly for these reasons, *H.*

contortus is emerging as a model parasitic nematode for anthelmintic resistance as

well as drug and vaccine discovery research (Geary, 2016; Gilleard, 2013; Nisbet et

al., 2016). Its utility as a model is largely due to a greater amenability to

experimentation than most parasitic nematodes; it is possible to establish and

maintain isolates *in vivo* in the natural host, perform genetic crosses *in vivo*, and

undertake *in vitro* culture for at least part of its life cycle, allowing drug assays and

genetic manipulation such as RNAi to be performed (Britton et al., 2016). The ability

to utilise these molecular approaches is complemented by extensive genetic

information about the genome structure and transcriptional differentiation of the

major life stages (Doyle et al., 2018; Laing et al., 2013; Schwarz et al., 2013).

The genetic characterisation of drug resistance in parasitic nematodes offers the promise of sensitive and specific diagnostic tools for use in research, surveillance and clinical diagnostics to support sustainable parasite management programs. Understanding the mechanisms of resistance to current drugs should also provide important information to help direct the development of new anthelmintic drugs (Hefnawy et al., 2017). However, the identification of genetic loci associated with drug resistance in these organisms remains a significant challenge. The only anthelmintic drug class to date for which the major resistance loci have been unequivocally identified is the benzimidazoles, where three mutations in the the isotype-1 ß-tubulin gene of *H. contortus* and several other trichostrongylid nematodes species are the major causal loci (Elard and Humbert, 1999; Kwa et al., 1994; Rufener et al., 2009; Silvestre and Cabaret, 2002). In the case of other anthelmintic drug classes, the genetic mediators of resistance remain obscure; for example, mutagenesis experiments in the free-living nematode *Caenorhabditis elegans* have identified many genetic loci capable of conferring low levels of ivermectin resistance, however, very few mutations confer high levels of resistance (Hunt, 1995). Concurrent mutation of three genes, *avr-14*, *avr-15* and *glc-1,* each encoding a glutamate-gated chloride channel (GluCl), confers a high level of resistance to ivermectin (Dent et al., 2000); however, mutations in orthologous GluCl genes have not been associated with ivermectin resistance in parasitic nematodes. In the case of parasitic nematodes, largely due to the lack of genomic resources, research into the genetic basis of ivermectin resistance has been dominated by the examination of small numbers of candidates genes (Kotze et al., 2014)). Candidate genes are generally chosen based on their potential roles as drug targets or in drug

efflux and examined for associations with sequence polymorphisms with the resistance phenotype. However, the extremely high levels of genetic diversity of *H. contortus* together with the limited number of well characterised ivermectin resistant isolates has provided a major challenge to such studies. The overall outcome of such candidate gene studies to date has provided at best circumstantial and inconsistent support for the involvement of any of the leading candidate genes in ivermectin resistance.

Genome-wide and genetic mapping approaches are at last becoming feasible for *H. contortus* due to progress in genetic crossing methodologies (Sargison et al., 2018) and an increasing complement of genomic resources. Previous work has undertaken two serial backcrosses between the genome reference susceptible strain MHco3(ISE) and two geographically independent ivermectin resistant isolates, MHco4(WRS) and MHco10(CAVR) (Redman et al., 2012)(**Figure 1 A,B**). These successfully introgressed ivermectin resistance loci from the two resistant strains into the susceptible MHco3(ISE) genomic background. Genetic validation of these backcross experiments, using a panel of microsatellite markers, revealed one such marker, Hcms8a20, to be present in an introgressed region of the genome. This was confirmed by subsequent work that further showed that none of the other leading candidate genes from the literature - *Hco-avr-14, Hco-glc-5, Hco-lgc-37, Hco-pgp-9, Hco-pgp-2* and *Hco-dyf-7* - showed evidence of introgression (Rezansoff et al., 2016). The increasing accessibility to high throughput sequencing of non-model organisms, together with an increasingly high quality reference for *H. contortus*,

offers an opportunity to characterise precisely the genome-wide evidence of

introgression and selection surrounding the Hcms8a20 locus.

Here, we build on these two previous genetic crosses, extending them for further

generations of passage with ivermectin selection (**Figure 1 C**), and generating whole

genome sequencing data from four steps in the experiment. We analyse these data

with novel statistical methods and exploit a chromosome-scale genome assembly to

map ivermectin resistance in the two genetically and geographically distinct strains of

H. contortus. Previous research has highlighted the value of novel methods in

population genetics for analysing experimental cross populations (Illingworth and

Mustonen, 2013). Here, in order to gain the maximum possible insight into the data

collected we developed a bespoke analysis, extending previous statistical work

(Abkallo et al., 2017; Illingworth et al., 2012), to analyse data from the experiment in

a manner that explicitly accounts for the genetic structure of the organisms in the

experiment; this approach was combined with more traditional population genetic

tools. Together, our methods provided a consistent and deep insight into the data,

identifying a single region of genomic introgression under selection for resistance.

This work represents the most comprehensive analysis of the genetics of

anthelmintic resistance in a parasitic nematode to date and demonstrates the power

of genetic crossing and contiguous genome assembly to eliminate false positive

genetic signals typically linked to resistance.

Figure 1. Outline of the initial crosses, backcrosses, *in vivo* passage and selection experiments. A. Ivermectin resistant strains MHco10(CAVR) or MHco4(WRS) ("resistant" haplotypes are depicted as red lines) were crossed with an ivermectin sensitive strain MHco3(ISE) ("susceptible" haplotypes as blue lines) to generate heterozygous F_1 progeny. F_1 eggs were cultured *in vitro* to L_3, before maturation *in vivo* to L_4/immature adults, from which females were used to initiate the backcross. **B.** The first round of backcross was performed by crossing heterozygous females from the initial cross with susceptible MHco3(ISE) males *in vivo*, resulting in F_2 progeny with reduced heterozygosity due to enrichment of MHco3(ISE) haplotypes. Resistance alleles were maintained in the backcross population by selection with ivermectin, before seeding the next round of the backcross with cross-derived heterozygous females and new susceptible MHco3(ISE) males. This process was repeated for four rounds of backcrossing, resulting in the backcrossed population becoming genetically similar to the MHco3(ISE) parental line in all regions of the genome not linked to ivermectin resistance. After four rounds of backcrossing, introgressed L_3 progeny were used to infect a new recipient sheep, resulting in segregation of susceptible and resistant alleles in both haplotypes among the progeny (mixed red/blue haplotypes). Eggs were cultured to L_3 before infecting two sheep, one exposed to ivermectin and one that did not receive drug. Post ivermectin treatment, L_4/immature adults from both sheep were recovered on necropsy for sequencing. **C.** Post-backcross L_3 were further passaged into a worm-naive sheep and treated with ivermectin, after which eggs were recovered and cultured to infective L_3 for reinfection. This process was repeated for four generations of

190   passage with selection (but without backcrossing). L_4/immature adults were
191   recovered after passage three and four for sequence analysis.

192   **Figure Supplement**
193   **Table S1.** Sequencing datasets used in this study

Results

Genetic diversity within and between parental *H. contortus* strains

We have taken a genome-wide approach to characterise genetic diversity within and between two ivermectin resistant strains (MHco10[CAVR] and MHco4[WRS]) and one ivermectin susceptible (MHco3[ISE]) strain of *H. contortus*. This study exploits a greatly improved, chromosomal-scale *H. contortus* reference genome sequence (**Figure 2A**), which was derived from sequencing and assembly of an inbred isolate of the MHco3(ISE) strain used here. Whole genome sequencing of pooled worms from each of the geographically distinct strains (**Figure 2B**) revealed abundant levels of nucleotide diversity throughout the five autosomes of the genome, with almost twice as much diversity on average in the resistant strains (MHco10[CAVR] mean π = 0.035 ± 0.008 standard deviations (SD); MHco4[WRS] mean π = 0.038 ± 0.008 SD) relative to the susceptible strain (MHco3[ISE] mean π = 0.022 ± 0.008 SD) (**Figure 2C; Figure S1A**). In contrast to the autosomes, the X chromosomal scaffolds had significantly less diversity in all three strains analysed (MHco3[ISE] mean π = 0.008 ± 0.008, MHco10[CAVR] mean π = 0.014 ± 0.012, MHco4[WRS] mean π = 0.017 ± 0.014) (**Figure 2C; Figure S1A**). This is consistent with a recent genome-wide analysis that revealed the X chromosome contained as little as 10% of the genetic diversity relative to the autosomes (Doyle et al., 2018). The difference between sensitive and resistant strains is perhaps not surprising given their history; MHco3(ISE) is a partially inbred laboratory strain that has been maintained in a closed, experimental flock, whereas the resistant strains have not been subjected to experimental inbreeding and have been isolated from outbred populations more recently.

We examined short-range haplotype diversity in each strain by measuring linkage

disequilibrium (LD) between pairs of SNPs detected in paired reads (**Figure S1B**).

Although restricted by the poolseq design to pairwise SNP comparisons less than

500 bp apart, i.e., within a read pair, we nevertheless observed considerable decay

in LD over this distance. Moreover, the rate of LD decay was correlated with

nucleotide diversity, such that there was a greatest loss of LD in MHco4(WRS)

followed by MHco10(CAVR) over the 500 bp, and a greater proportion of variants in

LD in the less diverse, susceptible MHco3(ISE) strain.

To understand the extent of genetic diversity that is private to, and thus unlikely to be

shared between, each population, we identified variant positions in the genome

whereby the variant frequency was enriched in one population but not the other two

populations. Of the ~7.6 million biallelic SNPs distributed in the genomes of the

samples analysed, ~514 k were private to MHco3(ISE) (**Figure S2A**), ~960 k to

MHco10(CAVR) (**Figure S2B**) and ~685 k to MHco4(WRS) (**Figure S2C**). Each

strain contained local regions of high diversity that differentiated it from the others.

We also explored larger structural variation within each of the strains, including

putative deletions, duplications and inversions detected by the presence of

discordant and split read pairs among the perfectly paired reads (**Figure S3**). We

note that structural variant calling in pooled individual, short read data will be

insensitive and subject to false positive calls due to mismapping. However, even a

conservative approach examining only high quality homozygous variants revealed

significant variation between strains (**Figure S3 A,C,E**), with a large number of

deletions, and substantially fewer duplications and inversions detected (**Figure S3**

B,D,F).

The genetic diversity between each of the three strains was assessed by pairwise

analysis of genetic diversity (measured by F_{ST} from single nucleotide variants in 10

kb windows), which confirmed significant genome-wide genetic diversity and strain

differentiation (**Figure 2D**). These data are consistent with the hypothesis that this

diversity reflects independent evolutionary histories due to being geographically and

therefore reproductively isolated (**Figure 2B**), and in turn most of this genetic

variation reflects genetic structure between these populations. These data

emphasise the challenge of characterising genetic variation associated with

phenotypes such as drug resistance by simply comparing genetic diversity between

a susceptible and resistant strain without accounting for this background genetic

variation.

Figure 2. Within and between strain genetic diversity of parental isolates used in the genetic cross. A. The 279 Mb genome assembly of *H. contortus* consists of five autosomal and two X chromosome scaffolds, named based on synteny with *Caenorhabditis elegans* chromosomes. The size of each scaffold is indicated, and are presented in order by length (Mb). **B.** Geographic origin of the susceptible MHco3(ISE) and ivermectin resistant MHco10(CAVR) and MHco4(WRS) strains used in the genetic crosses. All strains are archived at the Moredun Research Institute, UK – while MHco3(ISE) has been maintained for decades, it was originally

isolated in East Africa. **C.** Within-strain nucleotide diversity for each of the parental strains, calculated as mean diversity per 100 kb windows throughout the genome using *npstat*. **D.** Between strain diversity, calculated as pairwise F_{ST} in 10 kb windows throughout the genome using *popoolation2*. Colours here and throughout represent chromosomes as described in A.

Figure supplements

Figure S1. Characterisation of within-strain diversity.

Figure S2. Distribution of "private" variant sites per parental population.

Figure S3. Copy number and structural variation in the parental populations.

Genome-wide analysis of genetic diversity reveals the same single large introgressed region in both backcross lines

To utilise the genetic variation that differentiates these strains as described above, a backcross experiment was performed to introgress resistance-conferring alleles from the MHco10(CAVR) and MHco4(WRS) strains into the MHco3(ISE) susceptible genetic background (Redman et al., 2012). Here, we describe whole genome pairwise comparisons of genetic diversity throughout the backcross to assess the degree of introgression of resistance allele(s) into the susceptible genetic background, and to detect evidence of selection associated with resistance (**Figure 3**). Each pairwise comparison made was between particular generations of each backcross and the MHco3(ISE) susceptible parent. We hypothesised that where backcross populations were not subjected to further ivermectin treatment (MHco3/10.BC$_4$.noIVM & MHco3/4.BC$_4$.noIVM), there would be relatively little genetic differentiation across the whole genome between the population and MHco3(ISE). By contrast, where populations were subjected to further ivermectin treatment, we hypothesised that regions of the genome would have variable levels of

genetic differentiation with MHco3(ISE), with high levels of differentiation in regions

of the genome linked to an ivermectin resistance conferring locus and low levels of

differentiation elsewhere.

In both crosses, as we hypothesised, there was little genetic differentiation between

the post-backcross, no selection population and the susceptible parental strain

(**Figure 3**; MHco3/10.BC$_4$.noIVM and MHco3/4.BC$_4$.noIVM). This is despite the fact

that resistant alleles had been selected for in each round of the backcross; post

backcross, susceptible alleles would be segregating at a frequency of ~0.97 due to

the repeated backcrossing with MHco3(ISE), and therefore largely indistinguishable

from the susceptible parent. Upon selection (**Figure 3**; MHco3/10.BC$_4$.IVM and

MHco3/4.BC$_4$.IVM), we observed a region of differentiation from the susceptible

parent, particularly in the MHco3/10 cross, localised to the right arm of chromosome

V. This region was markedly differentiated in both MHco3/10 and MHco3/4 crosses

after *in vivo* selection and passage (**Figure 3**; MHco3/10.BC$_4$.IVM.P$_3$,

MHco3/4.BC$_4$.IVM.P$_3$, MHc3/10.BC$_4$.IVM.P$_4$ and MHc3/10.BC$_4$.IVM.P$_4$). No other

region of the genome was observed to show both a marked and progressive

increase in differentiation throughout the crosses. To test this more explicitly, we

regressed F$_{ST}$ per window throughout the genome over the course of the cross; only

the chromosome V region demonstrated consistent change associated with

introgression over time (**Figure S4 A,B**).

Figure 3. Genome-wide analysis of ivermectin associated loci. A. Pairwise
genetic diversity throughout the MHco3/10 (left plots) and MHco3/4 (right plots)
backcrosses. In both crosses and at each point in the backcross, the experimental
strain was backcrossed against the MHco3(ISE) parental strain – the data is
therefore presented to compare the genetic diversity (F_{ST} measure in 10 kb windows
using *popoolation2*) between the MHco3(ISE) and sampled time point in the cross as
per the cross schema in Figure 1. **B**. Output of the single-locus evolutionary model,
describing sites which were inconsistent with a model of neutral evolution, measured
using a likelihood threshold. Note that the X chromosome is not represented. Colours
represent chromosomes as described in Figure 2A.

Figure supplements

Figure S4. Summary of genome-wide change in F_{ST} throughout the backcross and subsequent
passage.

Figure S5. Fits between the model and the data for each data sample.

Figure S6. Location of significantly non-neutral loci identified using the single-locus population genetic
model.

These data also confirmed that the manner in which the population changed at a

given site was highly correlated between the two crosses (**Figure S4C**). Collectively,

these data suggest that there is a single introgression region on chromosome V that

is under selection upon drug exposure, and that this region is under selection in the

two independent serial backcrosses performed. This conclusion was supported by an

independent evolutionary analysis, which compared the observed frequencies of

variants associated with the resistant parental population with the distribution of

allele frequencies expected under the assumption of selective neutrality (**Figure S6**).

Alleles with frequencies that were strongly inconsistent with the neutral expectation

in all four of the collected samples, suggesting that they had potentially evolved

under the influence of selection, were over-represented in chromosome V (**Figure**

3B), with 71% and 91% of the total number of such alleles being found in this region

for the MHco3/10 (**Figure S6 A)** and MHco3/4 (**Figure S6 B)** crosses, respectively.

This supports the inference of a single selected region in the parasite genome

towards the right hand end of chromosome V.

Detailed analysis of introgression region on chromosome 5

To more precisely characterise the introgression region, we focus on chromosome V

(**Figure 4**). A comparison of genetic diversity between MHco3(ISE) and the end-point

of both crosses (BC$_4$.IVM.P$_4$) revealed a strikingly similar pattern between the

crosses, with a major region of differentiation between 38-42 Mb, with a second but

lesser increase in genetic differentiation between 45-48 Mb that is more prominent in

the MHco3/10 cross but less so in the MHco3/4 cross (**Figure 4A**). Encouragingly,

the major region of introgression lies immediately adjacent to the Hcms8a20

microsatellite marker (located at position 36.16 Mb along chromosome V) that was

shown to be genetically linked to ivermectin resistance in the preliminary analysis of

these backcrosses (Redman et al., 2012).

The F_{ST} data are supported by a comparison of Tajima's D throughout this region

(**Figure 4B**; top plots). Tajima's D test statistic is commonly used to distinguish

neutrally evolving loci (D ~ 0) from non-neutral processes, such as balancing (D > 0)

or positive selection (D < 0). Although variation in Tajima's D was observed

throughout the genome in both crosses (**Figure S7**), the greatest variation between

the susceptible parent and backcrossed lines was observed at approximately 39-40

Mb (**Figure 4B**; bottom plots), which was associated with significantly negative

Tajima's D in the backcrossed lines consistent with strong positive selection. In

contrast to the F_{ST} data, analysis using Tajima's D suggested a single peak,

corresponding to a single site or region under positive selection.

Figure 4. Analysis of chromosome V and the major region of introgression. A.
Genetic differentiation (F_{ST}) measured in 10 kb windows throughout chromosome V
is presented between MHco3(ISE) parent and both MHco3/10.BC$_4$.IVM.P$_4$ (light) and
MHco3/4.BC$_4$.IVM.P$_4$ (dark) representing the final time point of the crosses. Inset
presents the smoothed F_{ST} distribution of the two comparisons. Putative candidate
genes proposed in the literature in or near the introgression region (grey vertical
lines), as well as the original Hcms8a20 microsatellite marker identified by Redman
et al. (2012) as being linked to ivermectin resistance (black vertical line), are
presented. **B.** Comparison of Tajima's D per chromosome between MHco3(ISE)
parent (blue), MHco10(CAVR) (left panels) or MHco4(WRS) (right panels) and
passage 3 (MHco3/10.BC$_4$.IVM.P$_3$ or MHco3/4.BC$_4$.IVM.P$_3$; orange) and passage 4

(MHco3/10.BC$_4$.IVM.P$_4$ or MHco3/4.BC$_4$.IVM.P$_4$; yellow) of the crosses. Tajima's D

was calculated using *npstat* in 100 kb windows spanning the genome. The variance

in the mean value of Tajima's D among MHco3(ISE) and passages 3 and 4 samples

is presented as smoothed line (red). The null expectation is that variance between

these samples will be low in regions of the genome not under selection. **C**. Inferred

likelihoods from the multi-locus population genetic model. The mean likelihood of the

best fitting single-locus model at each locus position is shown by the solid black line;

the light blue interval around this line shows the 5-95% confidence interval for this

statistic calculated by a bootstrapping process. The position of the maximum

likelihood value of the likelihood is shown by the vertical black dotted line; a

confidence interval for this position, calculated from the bootstrapping values, is

shaded in gray. The mean likelihood of the best fitting two-locus model at including

each locus position is shown by the black dashed line.

Figure supplements

Figure S7. Analysis of Tajima's D variation in each chromosome per cross.

Figure S8. Contour maps of log likelihood scores derived from the two locus driver model.

Figure S9. Relative position and location of candidate genes from literature proposed to be

associated with ivermectin resistance in *Haemonchus contortus* and / or *Caenorhabditis elegans.*

Table S1. Coordinates of candidate genes from literature proposed to be associated with ivermectin

resistance in *Haemonchus contortus* and/or *Caenorhabditis elegans*

A multi-locus population genetic model assuming the presence of a single selected

allele in the introgression region inferred the selected resistance conferring allele to

be in a location consistent with the previous analysis. In the first step of this model,

putative segregating sites in chromosome V between the resistant and susceptible

parental populations were identified; a total of 474 such sites were identified in the

MHco3/10 dataset, with 157 sites in the MHco3/4 dataset; calculations were

performed upon data from these loci. A maximum likelihood model inferred the

location of the driver to be in a broadly consistent location in each experiment, with

peaks at 40.10 and 41.31 Mb respectively. However, there was considerable

uncertainty in the location of the selected allele, with confidence intervals of 36.86 to

44.03 Mb and 37.41 to 47.56 Mb in each case (**Figure 4 C**). These broad confidence

intervals were explained by the population structures inferred by the model, which

explicitly considered the location of crossover recombination events between the

parental strains throughout the experiment. While the precise location of crossover

events is stochastic, repeated calculations produced an ensemble of replicate

population structures for each set of model parameters. Example structures

generated using the maximum likelihood parameters for each population show how

selection for resistance drives the accumulation of genetic material from the resistant

parent through the course of the experiment (**Figure 5**); these outputs may be

contrasted with equivalent data generated under a model of selective neutrality

(**Figure S10**). Across 250 replicates, a mean of 177.1 crossover recombination

events were predicted in the final CAVR population (range 120 to 224), giving a

mean length of a block of parental genome of 17.64 Mb; in the final WRS population

a mean of 186.4 recombination events were seen (range 143 to 238), giving a mean

length of a block of parental genome of 17.06 Mb. Under selection, blocks of genome

containing the selected allele are favoured; the large size of each block reduces the

precision with which the location of the allele under selection can be identified.

Figure 5. Haplotype structure of chromosome V in an example output from the multi-locus model. Segments of genome from the parent containing the resistance allele are shown in red, while segments of genome from the susceptible parent are shown in blue. Data shown were generated using the maximum likelihood parameters in each case.

Figure supplements

Figure S10. Haplotype structure of chromosome V in an example output from the two driver model under neutral evolution.

Our analysis suggests that the large region of introgression was expected due to

limited extent of recombination in the population. To test this, we investigated an

extended multi-locus model incorporating selection on alleles at two loci within the

introgression region, as the presence of multiple drivers could explain the second,

smaller distinct peak of differentiation at approximately 45 Mb in the F_{ST} analysis.

Some separation of selected alleles by recombination is required for these alleles to

be distinguished; we therefore applied the constraint that these alleles must be at

least 2 Mb apart in the genome. Under this constraint, the best fitting model identified

variants under selection at the positions 38.59 Mb and 45.68 Mb in the MHco3/10

data (**Figure S8 A)**, and at the positions 41.20 Mb and 44.80 Mb in the MHco3/4

data (**Figure S8 B**). However, under the imposed constraint, the two-driver model

performed worse than the single-driver model shown above, favouring an

explanation in which only one variant is under selection. (We note that removing the

constraint that selected alleles be separated would guarantee a likelihood at least

equal to that of the single-driver model). Due to the limited extent of recombination in

the experiment, we cannot rule out the presence of more than one drug-resistance

variant in the identified region of chromosome V, but we did not find evidence in

favour of a second selected allele.

Many candidate genes have been proposed in the literature as being in association

with ivermectin resistance. We curated a list of genes that have been explored in *H.*

contortus, and/or had been shown to confer ivermectin resistance when mutated in

C. elegans. The current genome assembly is not annotated, so we determined their

location, either via mapping the published gene models from the V1 annotation, or

determining the closest *H. contortus* orthologous gene from *C. elegans* candidates

using Wormbase Parasite (Howe et al., 2017) (**Table S2; Figure S9**). At least one

candidate gene was located in each chromosome; in chromosome V, the location of

six candidate genes were determined. However, none of these genes was found in

the main introgression region defined by the F_{ST} analysis (**Figure 4 A**; vertical

annotated lines). Three genes - *Hco-lgc-55, Hco-avr-15,* and *Hco-pgp-1(9)*, the latter

two which have a strong association with ivermectin resistance - lie to the

telomere-side of the introgression region, however, they are located on the periphery

of the F_{ST} peaks rather than within them, suggesting that they are unlikely candidates

to be under direct selection. Considering the absence of candidate resistance genes

in the F_{ST} peaks, we conclude that the driver(s) of ivermectin resistance in the

MHco4(CAVR) and MHco4(WRS) strains are novel and have not been previously

described in association with ivermectin resistance.

Strength of selection

We further used the single locus model to infer the strength and manner of selection

in favour of the drug resistance allele. Data in each case indicated strong selection

for the resistant allele (**Figure 6**), such that susceptible worms produce multiple

times fewer offspring than the resistant worms under drug treatment. In each case

the value of the dominance coefficient, *h,* was less than one, indicating either an

additive effect for the MHco10(CAVR) case, whereby each copy of the resistance

allele contributes an equal amount of resistance, or of weak dominance for the

MHco4(WRS) data, whereby the first resistance allele contributes slightly more than

the second. This is consistent with the phenotypic characterisation of the initial

crosses, whereby the F_1 individuals that would have been heterozygous for resistant

alleles were partially resistant to ivermectin (Redman et al., 2012). As such, having

two copies of the resistant allele had a greater phenotypic effect than one; these data

therefore have implications for the mechanism of the drug resistance allele, and

evolution of drug resistance in general terms, as discussed further below.

Figure 6: Likelihood surface describing inferred selection parameters for the resistance allele. Likelihoods in each landscape are reported for the case in which the position in the genome of the resistant allele was at the maximum likelihood location for both the MHco10(CAVR) (**A**) and MHco4(WRS) (**B**) parental strains. Within our model the fitness of a genotype is determined by the alleles at a single resistant allele, with homozygous susceptible worms having fitness 1, homozygous resistant worms having fitness 1+*s*, and heterozygous worms having fitness 1+*hs* in the presence of the drug. The black point in each plot gives the location of the maximum likelihood values for selection.

Discussion

Genetic mapping identifies the same major ivermectin resistance locus in two independant *H. contortus* strains derived from different continents.

Given the importance of ivermectin, and other macrocyclic lactones, to human and animal health, the widespread emergence of resistance in multiple parasite species is a major concern. In spite of several decades of research, the molecular genetic basis of ivermectin resistance in parasitic nematodes has remained elusive. This is due to a combination of factors including a lack of genomic resources, the general experimental intractability of parasitic nematodes and the often high levels of genetic variation which complicates genetic analysis. *H. contortus* is the most experimentally tractable and most widely accepted parasite model for anthelmintic resistance research (Gilleard, 2013). Many candidate genes have been proposed to be associated with ivermectin resistance in *H. contortus*, chosen largely on *a priori* assumptions regarding their known or predicted function. Such candidate genes have typically been implicated as involved in ivermectin resistance largely on the basis of studies comparing SNP or haplotype frequencies between small numbers of resistance and susceptible isolates (Blackhall et al. 1998; Eng et al. 2006; Luo et al. 2017; Blackhall et al. 2003; Urdaneta-Marquez et al. 2014) or pre- and post-ivermectin treatment (Raza et al. 2016; Lloberas et al. 2013). However, careful validation of a number of these candidates in different field populations (Laing et al., 2016) or in controlled crosses (Rezansoff et al., 2016) have failed to support the initial associations observed. The discordance among studies reflects the difficulty in identifying real associations between genotype and phenotype simply by comparing resistant and susceptible populations, or individuals, due to the confounding effects

of extremely high levels of genome-wide genetic variation. The lack of consistency

between studies had led to much discussion and debate regarding the complexity of

the genetic basis of ivermectin resistance both in terms of the number of loci involved

and the extent of geographical variation (Kotze et al. 2014; Prichard 2001; Beech et

al. 2011). We have undertaken a genetic mapping approach as a more systematic

and objective way to identify the major ivermectin loci to help resolve the lack of

clarity and debate.

In this paper, we describe the analysis of whole genome sequence data collected

from the most recent phase of a multi-generational crossing experiment conducted in

populations of the parasitic worm *H. contortus*. Our analysis of genome sequence

data unequivocally identifies a single region within chromosome V as containing at

least one major ivermectin resistance allele. This major ivermectin resistance locus is

common to two independent ivermectin strains - MHco10(CAVR) and MHco4(WRS) -

that were originally isolated from different continents; Australia and Africa

respectively. This genomic region is clearly distinguishable in the experimental data,

being highlighted by each of the different population genetic methods applied to our

data. There are no other comparable regions of introgression elsewhere in the

genome in either of the two backcross experiments suggesting this is the most

important ivermectin resistance locus in the MHco10(CAVR) and MHco4(WRS)

strains. There has been much discussion regarding the likely multigenic basis of

ivermectin resistance but, although there could be additional additive loci, these

results strongly suggest a single locus (or potentially multiple closely linked loci) is

likely to be the major effector of ivermectin resistance in these two strains.

While each of our analytical methods gave a consistent identification of the region under selection, our multi-locus population genetic analysis provided an estimate of the uncertainty in the location of the selected allele, inferring its position to lie at a position between 37 Mb and 45 Mb. While the region in which we infer a selected allele to exist is large, our result nevertheless provides some ability to discriminate between potential ivermectin resistance alleles in the worm population. Most of the leading published candidate ivermectin resistance genes including *Hco-glc-5*, *Hco-avr-14*, *Hco-pgp-9*, *Hco-lgc-37*, *Hco-pgp-2* and *Hco-dyf-7* were not located in or near the region of introgression, which is consistent with a recent study using targeted sequencing of these genes in which none of them showed a signal of introgression in these two backcross experiments (Rezansoff 2016). This suggests that none of these candidate genes are a major ivermectin resistance-conferring locus in the MHco10(CAVR) and MHco4(WRS) strains. Although, we identified three other previously described candidate genes - *Hco-lgc-55*, *Hco-avr-15*, and *Hco-pgp-1* - adjacent to the introgression region, none were found within the peak of the region suggesting that these also are unlikely to be major direct targets of ivermectin-mediated selection. We acknowledge that we cannot discount previously described candidate genes as mediators of resistance in other isolates of *H. contortus*, or in other nematode species. Moreover, our data does not exclude a driver of transcriptional regulation within the introgression region that is under selection, which itself influences expression of other genes, including candidate genes, outside of the introgression region.

What might account for the large number of candidate genes previously suggested to be associated with ivermectin resistance? The answer might lie in our direct comparison of genetic diversity between the parental strains (**Figure 2D**). This visualisation of the extent of genetic divergence between these strains makes it easier to understand how particular sequence polymorphisms might be naively attributed to being associated with resistance when in fact they simply represent one of many strain specific genetic variations that occur as a result of the independent evolutionary histories and lack of interbreeding between the strains. This highlights the challenges of interpreting simple direct comparison of strains, even with genomic approaches, when trying to disentangle those genetic polymorphisms underlying resistance from a complex background of genetic variation independent of a strains resistance status. The use of controlled genetic crosses as presented here and elsewhere (Choi et al., 2017; Redman et al., 2012; Rezansoff et al., 2016), whereby population genetic structure can be explicitly accounted for in the analysis, provides a powerful way to mitigate the challenge of discriminating resistance causing alleles from background genetic variation. As such, the locus we have identified on chromosome V represents the only unequivocal major ivermectin resistance locus identified in any parasitic nematode species to date.

Population genetic and evolutionary modelling helps to statistically define the boundaries of the genetically introgressed genomic region linked to the ivermectin resistance conferring locus

Inference of selection from the multi-locus model suggested the presence of stronger selection for the drug resistance allele in the MHco3/10 cross than in the MHco3/4

cross (**Figure 6**). Additionally, although the data suggests that resistance is at least additive in both crosses, i.e., heterozygotes are likely to confer some resistance, but increased resistance is achieved by having two resistant alleles, there is some evidence for dominance in the MHco4(WRS) strain, whereby the second resistant allele confers a lesser fitness advantage. A previous study reported resistance to be a completely dominant trait in the CAVR strain (Le Jambre et al., 2000) on the basis that no difference was observed in the resistance levels of heterozygote and homozygote resistance worms. Our finding of additivity arises from the evolution of the worm populations during passage following the backcross experiment (i.e., in $BC_4.IVM.P_{3/4}$ samples versus the $BC_4.IVM$ data), and particularly the fixation of alleles from the resistant parent at putative segregating sites near the inferred position of resistance. In our model, alleles can reach fixation only where there is selection in favour of the homozygote resistant type over and above that for heterozygote resistant individuals.

We have here utilised a range of population genetic methods to attempt to localise selection. In addition to the standard metrics of F_{ST} and Tajima's D, we have implemented single-locus and multi-locus methods which explicitly account for the history of the cross population. Each approach has limitations. While most models neglect linkage between adjacent sites in the genome, our multi-locus model explicitly accounts for genomic structure during the cross. However, even this model involves the use of heuristic measures to identify segregating sites between the two parental strains, and is based upon a map of recombination in the worm genome (Doyle et al., 2018) that, while the state of the art for parasitic species, was derived

from a limited number of observed crossover events. None of our models accounts

for genomic rearrangements during the experiment, which are inherently difficult to

observe via short read sequencing.

The output of our population genetic model suggests that our ability to resolve the

precise location of the variant under selection is limited by the number of

recombination events occurring in the worm population. Precise identification of the

location of an allele under selection requires that the allele not be linked to other

nearby alleles in at least some fraction of individuals in the population. While the

inherent biological recombination rate here has a role, some of the lack of

recombination events in this case results from bottlenecking of the population during

the course of the experiment due to the limited number of L_4 worms that it was

possible to collect from each sheep and transplant in the successive generation of

the cross. The limited individuals, and hence, limited recombinants, reduced the

resolution of the experiment to resolve selection. Genetic drift induced by successive

population bottlenecks introduces considerable uncertainty in the outcome of the

experiment, such that replicate sets of allele frequencies from our model showed

considerable differences between each other (**Figure S5**). This implies that the data

from the experiment itself should be understood as the output of a stochastic

process; beyond the clear large-scale patterns observable in the data and detailed

above, more minor details of the output might not be seen again were the experiment

to be repeated.

The structure of the experiment thus imposes a limit on our ability to infer the location of a selected allele, and an improved characterisation of selection would likely best be achieved through the use of alternative experimental procedures. Conducting further generations of cross as performed within this experimental framework would induce more recombination events, reducing the mean size of parental genomic blocks. However, such a course of action would be limited in scope were it not accompanied by the use of larger L_4 populations, for example by the simultaneous passage of the population through multiple animals in parallel. Clearly, a larger or longer experiment would have cost and welfare implications: the statistical framework we have developed will help design and justify such an approach.

The importance of chromosomal-scale genome assemblies and the dangers of using fragmented draft genomes in genome-wide genetic mapping and population genomic analyses

The success of identifying a single region of introgression was dependent upon a greatly improved and highly contiguous, chromosomal-scale reference genome assembly for the *H. contortus* isolate, MHco3(ISE).N1. This isolate was used in the original draft assembly of this species (Laing et al., 2013), which was derived via inbreeding of the same MHco3(ISE) strain used in the backcrosses presented. In the absence of a contiguous chromosomal scale genome assembly, interpretation of these analyses can be extremely challenging; true genetic signal(s) can be obscured by technical artifacts associated with a fragmented assembly that can include short contigs lacking spatial orientation, multiple haplotypes present, collapsed paralogs,

and poor resolution of repeat structures. As a consequence, read mapping and

variant calling in fragmented genomes can be suboptimal. These technical

challenges are exacerbated in highly polymorphic species such as *H. contortus* and

Teladorsagia circumcincta (Choi et al., 2017), in which reference genomes were

constructed from pools of individuals that, despite inbreeding, still contained

significant genetic variation. Perhaps the biggest problem with using a fragmented

genome assembly for genome-wide analyses is the inability to determine linkage.

Signals of selection will often be dispersed across multiple scaffolds of a fragmented

draft genome assembly when in reality they are co-located in a single genomic

region. This can potentially can give rise to the erroneous conclusion that multiple

independent loci show a signal of selection when it may be that only a single

selected locus exists. For example, a major effect single locus that has a physically

extensive signature of selection due to a hard selective sweep, and the associated

hitchhiking effect, could easily be interpreted as multiple independent unlinked loci

under selection. These artefacts are evident when examining the sequence data we

have generated on different versions of the *H. contortus* MHco3(ISE).N1 reference

genome assembly. In this case, the introgression region of chromosome V only

became apparent after significant improvement in chromosome contiguity and once

chromosome V was fully assembled (**Figure 7**). In the earlier, more fragmented, draft

genome assemblies, the signature of selection was dispersed across multiple

scaffolds making the number of major loci underlying ivermectin resistance in the

MHco10(CAVR) and MHco4(WRS) strains unclear. This will be an increasingly

important issue as unfinished draft reference genomes are produced for an

increasing number of non-model species and whole genome sequencing becomes

cheaper and more widely available. It will be important for those using genome-wide

data for population genomic and genetic studies to be aware of the limitations of

using unfinished draft reference genomes in various genome-wide analyses.

Figure 7: Demonstrable effect of genome contiguity on the resolution of the
introgression region. The same pairwise comparison - MHco3(ISE) *vs*
MHco3/10.BC$_4$.P$_4$ - is presented using on three versions of the *H. contortus* genome;
the published V1 draft assembly (Laing et al., 2013) (top; N50 = 0.083 Mb, N50(n) =
1,151, n = 23,860), an intermediate improved assembly (middle; N50 = 5.2 Mb,
N50(n) = 16, n = 6668), and the chromosomal-scale assembly presented here and
elsewhere (Doyle et al., 2018)(bottom; N50 = 47.4 Mb, N50(n) = 3, n = 8). The top
and middle plots are coloured light and dark grey to reflect alternate contigs /
scaffolds, whereas the bottom plot is coloured as in Figure 2A.

Our comprehensive genome-wide analysis of parental strain genetic diversity

re-emphasises that *H. contortus* is highly genetically diverse both within and between

strains. It is clear, however, that short read mapping-based analyses that aim to

exploit a single reference genome are almost certain to underestimate the extent of

genetic variation that distinguishes a divergent strain. We have exploited linked

selection of a subset of this variation, i.e., only single nucleotide polymorphisms, to

characterise the introgression event in these analyses; however, understanding the

functional consequences of this strain specific diversity will rely on a more

comprehensive description of all of the variation that defines a strain. As an

alternative to a linear reference genome, the use of population genome graphs - a

non-linear or branching reference that contains alternate paths representing known

genetic variation - may be more suitable (Paten et al., 2017), and allow better

characterisation of known variation that is too complex to be analysed using a linear

reference (Maciuca et al., 2016). This could be made possible once a

comprehensive analysis of population and perhaps global genetic diversity reference

set of genetic variation is made available (Sallé *et al.*, in preparation). Alternatively,

such variation may require geographically diverse isolates to each have a *de novo*

reference sequence, ideally sequenced and assembled using long read sequencing

technology such as Pacbio or Oxford Nanopore sequencing, which will allow for a

more comprehensive description of genetic variation within a population, while

allowing large scale variation between reference strains to be characterised.

Conclusion

Our genome-wide analysis of two previous genetic crosses between ivermectin

resistant and ivermectin sensitive isolates has identified a major ivermectin

resistance locus in two genetically and geographically distinct strains of *Haemonchus contortus*. This result depended on the development of novel population genetic methods to understand the genetics of the crossing procedure, and on the availability of a chromosome-scale genome assembly. More traditional population genetic analysis highlights the high level of variation both between and within the parental populations used to construct the cross, potentially explaining the difficulty in validating previous candidate genes that were identified using approaches that neither experimentally, nor statistically controlled for this genetic diversity. This work represents the most comprehensive analysis of the genetics of anthelmintic resistance in a parasitic nematode to date and demonstrates the power of genetic crossing and contiguous genome assembly to eliminate false positive genetic signals typically linked to resistance. Our results eliminate the possibility that many of the previously proposed candidate genes are involved in ivermectin resistance in these isolates, and should focus future efforts on identifying the causal variant within the region we identify.

Acknowledgement

Work at the Wellcome Sanger Institute was funded by Wellcome (grants 098051 and 206194) and by the Biotechnology and Biological Sciences Research Council (BB/M003949). CJRI was supported by a Sir Henry Dale Fellowship, jointly funded by Wellcome and the Royal Society (grant 101239/Z/13/Z). We thank Pathogen Informatics and DNA Pipelines (WSI) for their support and expertise, and Guillaume Sallé for constructive feedback on the manuscript.

Competing interests

The authors declare no competing interests.

Materials and methods

Background of strains and original backcross

The MHco3(ISE), MHco4(WRS) and MHco10(CAVR) are *H. contortus* strains maintained and stored at the Moredun Institute. The MHco3(ISE) strain was originally derived by multiple rounds of inbreeding of the SE strain (Roos et al. 2004). The precise provenance of the SE strain is not clearly recorded but is thought to be originally obtained from East Africa. MHco10(CAVR) is derived from the Chiswick Avermectin Resistant Strain (CAVR), an ivermectin resistant field strain originally isolated in Australia (Le Jambre et al., 1995). MHco4(WRS) is derived from the ivermectin resistant White River Strain (WRS) originally isolated from South Africa (van Wyk and Malan, 1988).

The experiment described here extends two previously-described backcross experiments. The construction, phenotypic validation, and initial microsatellite analysis of these backcrosses has been described previously (Redman et al., 2012), and is outlined in **Figure 1A and B.** Briefly, two independant crosses were performed in parallel: (i) MHco3(ISE) x MHco10(CAVR), and (ii) MHco3(ISE) x MHco4(WRS). In the first generation of each cross, female worms of the ivermectin resistant parental strains [MHco10(CAVR) or MHco4(WRS)] were crossed with male MHco3(ISE) worms, to generate lines designated MHco3/4 and MHco3/10, respectively (**Figure 1A**). After the first cross, the F_1 generation females were mated to male MHco3(ISE), resulting in backcross generations designated MHco3/10.BC_n and MHco3/4.BC_n, where n is the number of backcross generations (BC). This involved the recovery of immature worms from the abomasum at day 14 post infection by necropsy of donor

sheep, which were washed in physiological saline (0.85% NaCl) after which worm

sex was determined (Denham, 1969), followed by surgical transfer of 45-100 male

MHco3(ISE) and 50-100 female F_1 L_4/immature adult worms into worm-free recipient

lambs all within 2 h of the original collection. This process of backcrossing was

repeated for a total of 4 generations, i.e., MHco3/10.BC_4 and MHco3/4.BC_4 (**Figure**

1B). *In vivo* ivermectin selection was applied after mating and before collection of

eggs to enrich for ivermectin resistant adults to be used in the subsequent round of

the backcross. A controlled efficacy test was performed on the three adult strains

and MHco3/4.BC_4 and MHco3/10.BC_4 strains to determine the initial levels of

ivermectin efficacy, and the degree of resistance acquired in the introgressed lines.

After four rounds of backcrossing, a selection experiment was performed.

MHco3/10.BC_4 and MHco3/4.BC_4 strains were used to infect recipient sheep, after

which eggs were collected and in vitro culture to L_3. These infective larvae were used

to infect two sheep per strain, one that received 0.1 mg/kg ivermectin and one that

remained drug naive. At 7 days post treatment, treated and naive L_4/immature adults

were recovered by necropy and stored for molecular analysis.

The backcrosses described above were extended by performing a further four

rounds of *in vivo* ivermectin selection, during which mating within the BC_4 population

continued (**Figure 1C**). For each cross, progeny of the BC_4 generation were cultured

to L_3 and used to infect a donor sheep, which was subsequently treated with

ivermectin (0.1 mg/kg in round 1, followed by 0.2 mg/kg in following rounds). Eggs

from ivermectin-treated survivor adults were collected post-treatment, cultured to L_3,

and used to infect a new donor sheep. L_4/immature adults were collected by

necropsy after rounds three (i.e., $BC_4.IVM.P_3$) and four (i.e., $BC_4.IVM.P_4$) of selection

and stored for molecular analysis.

Library preparation and sequencing

Whole genome sequencing was performed on the three parental strains

(MHco3(ISE), MHco10(CAVR) & MHco4(WRS)), and from each cross, pre and post

ivermectin treatment following four rounds of backcrossing (MHco3/10.BC_4.noIVM,

MHco3/4.BC_4.noIVM, MHco3/10.BC_4.IVM, MHco3/4.BC_4.IVM), and after 3 and 4

rounds of additional selection after the backcross (MHco3/10.BC_4.IVM.P_3,

MHco3/4.BC_4.IVM.P_3, MHco3/10.BC_4.IVM.P_4, MHco3/4.BC_4.IVM.P_4). Pools of male

and female worms were included for the parental (n = 50-60 worms) and BC_4

samples (n = 25-30 worms), whereas only female worms were used in the passage 3

& 4 samples (n = 60 worms). Genomic DNA from each pooled sample was prepared

using a phenol chloroform extraction protocol as previously described. Sequencing

libraries were prepared using a PCR-free protocol (Kozarewa et al. 2009), and

sequenced as described in **Table S1**. We generated approximately 614×10^9 bp of

sequence data, which equates to approximately 199.65× unmapped genome

coverage per sample. Raw sequence data quality was analysed using FASTQC

(*http://www.bioinformatics.babraham.ac.uk/projects/fastqc/)* and was assessed using

MultiQC (Ewels et al., 2016) prior to further processing.

The raw sequencing data generated during the current study are available in the

European Nucleotide Archive repository (http://www.ebi.ac.uk/ena/) under the study

accession number PRJEB2353 (**Table S1**).

Mapping and variant analysis

Raw sequence data was mapped per lane to the current reference genome

(available here: ftp://ngs.sanger.ac.uk/production/pathogens/Haemonchus_contortus

) using BWA-MEM (Li, 2013)(default parameters with *-k 15*). Samples for which

multiple lanes of mapped data were available were merged, duplicate reads were

marked and removed using *Picard v2.5.0* (https://github.com/broadinstitute/picard),

and mapped perfect read pairs were extracted per sample (*samtools-1.3 view -f 2*).

Genome-wide variants were determined for each sample using samtools-1.3 mpileup

(*-F 0.25 -d 500*). Nucleotide diversity and Tajima's D was determined using *npstat*

(Ferretti et al., 2013), which required pileup files (which we derived from the mpileup

generated above) that were split per sample per chromosome as input. We

determined short-range linkage disequilibrium between pairs of variants in single and

paired reads using *LDx* (Feder et al., 2012). We compared these data with estimates

of LD decay over genetic distance as proposed by Hill and Weir (Feder et al., 2012;

Hill and Weir, 1988). *Popoolation2* (Kofler et al., 2011) was used for the analysis of

pairwise genetic diversity throughout the genome. Briefly, the mpileup file was

converted into a synchronised file (*popoolation2 mpileup2sync.jar --min-qual 20*),

after which indels + 5 bp (popoolation2 *identify-indel-regions.pl −−min-count 2*

−−indel-window 5) and repetitive and difficult to map regions (separately identified by

repeat masking the genome [http://www.repeatmasker.org/]) were excluded

(*popoolation2 filter-sync-by-gtf.pl*). The synchronised file was used as input to

determine pairwise F_{ST} which was calculated in 10 kb windows throughout the

genome (*popoolation2 fst-sliding.pl --pool-size 50 --window-size 10000 --step-size 5000 --min-count 4 --min-coverage 50 --max-coverage 2%*). Similarly, per base comparisons were determined using a Fisher's exact test (*popoolation2 fisher-test.pl --min-count 4 --min-coverage 50 --max-coverage 2% --suppress-noninformative*).

Copy number variation between parental strains was determined using *cnv-seq* (Xie and Tammi, 2009). Best read mapping hits were determined per bam (*samtools-1.3 view -F 4 bam | perl -lane 'print "$F[2]\t$F[3]"' > bam.hits*), before read counts for pairwise comparisons determined in 10 kb sliding windows using *cnv-seq.pl*. The R package *cnv* (v 0.2-8) was used to determine pairwise chromosome-wide normalised $\log_2$ ratios. Characterisation of structural variation in the parental strains was performed using the *speedseq sv* pipeline (Chiang et al., 2015) to identify putative duplications, deletions and inversions in each strain. This approach exploits all reads mapped using BWA-MEM, including split and discordant read pairs, which are subsequently scored using LUMPY and SVTyper. Conservative filtering was applied to retain only homozygous variants (1/1) with a minimum quality score of 100.

Population genetic modelling

Bespoke single-locus and multi-locus models were used for population genetic inference from the data. Common to both methods, variants in the genome potentially associated with resistance were identified. At each locus in the genome, a nucleotide {A,C,G,T} was defined as existing in the susceptible parental population if it was observed at a frequency of at least 1% in that population. Loci in the resistant parental population were then identified for which exactly one nucleotide that did not

exist in the susceptible population was observed at a frequency of at least 1%;

without prejudice as to its phenotypic effect, this was denoted the 'resistant' allele at

that locus, being associated with the resistant parental population.

Single-locus model

A single-locus population genetic model was then used to identify variants with

frequencies that were inconsistent with selective neutrality. The neutral expectation

was calculated using a Wright-Fisher evolutionary model to simulate the progress of

a variant allele, modelling the probability distribution of the frequencies of

homozygote resistant (q_1), heterozygote (q_h), and homozygote susceptible (q_0)

individuals throughout the course of the experiment under the assumption of

selective neutrality. The Wright-Fisher model uses a simple multinomial process for

propagation; if the next generation of the population contains N individuals, the

probability of obtaining n_1 , n_h , and n_0 individuals with each diploid variant is given

by:

$$P(n_1, n_h, n_0) \;=\; \tfrac{N!}{n_1! n_h! n_0!} q_1^{n_1} q_h^{n_h} q_0^{n_0} \tag{1}$$

At the stages in the experiment where the population was comprised of eggs, we

assumed N to be large, such that the processes of crossing and backcrossing could

be described deterministically. Under these circumstances the genotype frequencies

following a cross, denoted q_1' , q_h' , and q_0' , are given by:

857
$$q'_1 = q_1{}^2 + q_1 q_h + \tfrac{1}{4} q_h{}^2$$

858
$$q'_h = q_1 q_h + 2 q_1 q_0 + \tfrac{1}{2} q_h{}^2 + q_h q_0 \qquad [2]$$

859
$$q'_0 = q_0{}^2 + \tfrac{1}{4} q_h{}^2$$

860 while the genotype frequencies following a backcross with the homozygous

861 susceptible strain are given by:

862
$$q'_1 = 0$$

863
$$q'_h = q_1 + \tfrac{1}{2} q_h \qquad [3]$$

864
$$q'_0 = q_0 + \tfrac{1}{2} q_h$$

Evaluating this process gives a probability distribution for the frequency of the

resistant allele at any given point in the experiment, dependent upon the number of

resistant alleles in the initial resistant population. This number of resistant alleles,

which we denote n_r, is unknown; sequence data from the resistant parental strain

were used to generate a prior distribution for this value.

We first suppose that the frequency of the resistant allele in the resistant parental

population is equal to some value, p_r. Given that the initial experimental population

contains 50 resistant worms, with 100 alleles, collected from this larger population,

the distribution of n_r is then given by:

874
$$P(n_r = j) = \binom{100}{j} p_r^j (1 - p_r)^{100-j} \qquad [4]$$

We now suppose that sequencing the resistant parental population gave a_r resistant

alleles at the locus in question out of a total read depth of A_r. The initial frequency

p_r is unknown; however using the data an estimate can be made for this statistic,

expressed as a distribution of the allele frequency. Specifically, under the assumption

of a uniform prior, the underlying probability p_r can be said to be distributed as a

beta distribution with parameters a_r and $A_r - a_r + 1$. We therefore obtain the

following distribution for the statistic n_r:

882
$$P(n_r = j) = \int_0^1 \frac{A_r!}{n!(A_r - a_r!)} p_r^{a_r+1}(1 - p_r)^{A_r - a_r + 1} \binom{100}{j} p_r^j (1 - p_r)^{100-j} dp_r \qquad [5]$$

for any value of j between 0 and 100. Values of this distribution were calculated

using numerical integration. This process generated the neutral expectation of the

resistant allele frequency conditional on the observation of this frequency in the

resistant population. If at the sampling point t, a total of N_t worms were collected,

the distribution of the number of worms n_t with the resistant allele at that point is

given by:

889
$$P(n_t = k) = \sum_j P(n_t|n_r = j)P(n_r = j) \qquad [6]$$

890 for all $0 \le k \le N_t$.

The extent to which observed allele frequencies were consistent with the neutral model was calculated using a likelihood model:

$$L_t \; = \; \sum_{k=0}^{N_t} P(n_r = k)\binom{A_t}{a_t}(k/N_t)^{a_t}(1 - (k/N_t))^{A_t - a_t} \qquad [7]$$

where at time t, a_t resistant alleles from a read depth of A_t were observed. A low likelihood L_t indicates deviation from the neutral expectation; Bonferroni correction was used to identify significance at the 95% level. Data from each sample were analysed; four-fold non-neutral sites, being loci for which a significant likelihood was calculated from all four samples collected throughout each cross experiment, were identified.

Multi-locus model

A multi-locus model was developed to describe the manner in which allele frequencies would be expected to change over the course of the experiment; this model exploits information about the location of variant loci provided in the *H. contortus* reference genome, and the genome-wide map of recombination rate in the worms (Doyle et al., 2018); this map was inferred by characterising recombination breakpoints in F_1 L_3 progeny by whole genome sequencing, from which recombination rates were determined by comparing genetic distances between SNPs against their physical location in the genome. Where fixed genetic differences occur between parental strains, the dynamics of adaptation in the resulting cross are relatively straightforward (Illingworth and Mustonen, 2013; Illingworth et al., 2012);

this has been exploited to identify quantitative trait loci in yeast and malaria cross

populations (Abkallo et al., 2017; Parts et al., 2011). Here a heuristic approach was

used to identify fixed genetic differences between the parental strains before

modelling evolution at these loci to identify the location of alleles conveying drug

resistance.

Three filters were used to identify putatively fixed differences between parental

strains. Firstly, we identified genomic sites for which the frequency of the resistant

allele was 95% or greater in the resistant parent, requiring a read depth of at least

50x for such sites to achieve a good level of statistical certainty. Secondly, we noted

that, following the backcross performed at the start of the experiment, no locus can

be homozygous for the resistant allele and as such, the resistant allele frequency

can be no greater than 50% in the population; by way of reducing noise, we removed

any locus with resistant alleles at a frequency of 60% or greater in the

MHco3/10.BC$_4$.noIVM sequence data. Thirdly, we noted that at homozygous

separating sites in a population following a cross, the allele frequency will change

smoothly over time; where there are N genomes in the population, the allele

frequency, considered as a function of chromosome position, will change by $1/N$

wherever a crossover recombination event occurs within a genome. A diffusion

model of allele frequency change, described in a previous publication (Abkallo et al.,

2017), was used to identify allele frequencies across the genome that were

consistent with this pattern. This analysis fits a posterior distribution to the allele

frequencies across the genome, also inferring the extent of noise in the sequence

data via a beta-binomial model intrinsic to the fitting process. Conservatively, sites no

more than eight standard deviations from the mean of the posterior frequencies were

included in the analysis. These approaches resulted in data from a total of 1,368 loci

in the MHco10(CAVR) dataset, and from 219 loci in the MHco4(WRS) dataset, were

retained; the full set of allele frequencies are shown in Supporting Figure S5.

Considering these data, we used an individual-based Wright-Fisher simulation to

model the outcome of the experiment, taking into account the backcrossing,

selection, and bottlenecking according to the experimental design. This approach

accounted for the stochastic nature of allele frequency changes due to the effect of

repeated population bottlenecking. We suppose that selection acts in favour of the

resistance allele at a given locus when worms are in a sheep being treated with

ivermectin, such that the fitnesses of individuals that are homozygous for the

resistant allele (w_1), heterozygous for the resistant allele (w_h), or homozygous for

susceptible alleles (w_0) are given by:

$$w_1 = 1 + s$$

$$w_h = 1 + hs \qquad\qquad [10]$$

$$w_0 = 1$$

where s is the selection coefficient (fitness advantage of the homozygous resistant

compared to homozygous susceptible) and h is the dominance coefficient

(proportion of fitness change conveyed by a single copy of the allele). We assumed

that all worms have equal fitness in the absence of the drug. The likelihood of any

given inference was calculated using the beta-binomial likelihood function obtained

from the diffusion model above. Accounting for the stochasticity of the system, a set

of at least 250 simulations were generated for each set of parameters, calculating

the mean likelihood fit to the data across the simulations. A bootstrapping process

was applied to quantify the uncertainty in this likelihood.

In an extension to this model, a two-driver was considered, in which the resistant

variant at each of two loci was under selection; to allow the exploration of model

space in this case within reasonable computational time the assumption was made

of additive selection. To distinguish this from the previous model, a requirement was

imposed that the selected alleles be separated by at least 2 Mb in the genome.

Table S1. Sample sequencing data archived at European Nucleotide Archive

repository under the study accession PRJEB2353

-   See accompanying .xlsx document

Figure S1. Characterisation of within-strain diversity. A. Within strain nucleotide
diversity per chromosome, summarising genome-wide data presented in Figure 2C.
Colours represent chromosomes as described in Figure 2A. **B**. Linkage
disequilibrium between variants present in paired reads was estimated using LDx for
each parental strain. Line represents expected LD decay over genetic distance
(Feder et al., 2012; Hill and Weir, 1988).

Figure S2. Distribution of "private" variant sites per parental population. A.
MHco3(ISE). **B**. MHco10(CAVR). **C**. MHco4(WRS). Private sites were defined as
having a frequency greater than 0.05 in the population of interest, but less than 0.05
in the two additional populations.

Figures contain the following tables:

B. MHco3(ISE)

MHco3(ISE)	Deletions	Duplications	Inversions
Count	1455	110	1
Mean length (SD)	1374.65 (1823.59)	7950 (12,441.94)	2594
Range (bp)	3 - 22714	92 - 71290	2594 - 2594

D. MHco10(CAVR)

MHco10(CAVR)	Deletions	Duplications	Inversions
Count	7049	206	11
Mean length (SD)	1413.20 (2466.03)	16389.51 (19434.07)	764.45 (609.80)
Range (bp)	27 - 89216	140 - 94802	305 - 2372

F. MHco4(WRS)

MHco4(WRS)	Deletions	Duplications	Inversions
Count	5800	141	5
Mean length (SD)	1453.03 (2407.22)	12493.09 (15613.98)	584 (374.19)
Range (bp)	3 - 79176	173 - 92931	180 - 1121

Figure S3. Copy number and structural variation in the parental lines. A,B.
MHco3(ISE), **C,D**. MHco10(CAVR), **E,F.** MHco4(WRS). Data per circos plot (**A,C,E**)
is orientated as follows; outer circle: CNV variation between MHco3(ISE) and each
resistant parent. No CNV comparison was made in A; second circle: deletions; third
circle: duplications; inner circle: inversions. The data presented in **A,C,E** is
summarised in the boxplots (feature length distribution) and tables in **B,D,F**.

Figure S4. Summary of genome-wide change in F_{ST} throughout the backcross and subsequent passage. A regression between F_{ST} and the four sampling time points was performed for each 10 kb window sampled across the genome for both MHco3/10 (A) and MHco3/4 (B) lines. The slope of the regression was plotted. C. Correlation in slope (F_{ST} vs backcross progression) between MHco3/10 and MHco3/4. Dashed line represents x=y. Colours represent chromosomes as described in Figure 2A.

Figure S5. Fits between the model and the data for each data sample. Blue dots show filtered allele frequencies for putative segregating sites. The model fit is shown as gray lines; a distinct line is shown for each of the 250 replicate simulations run for the parameters generating the maximum likelihood fit. **A.** MHco3/10.BC$_4$.noIVM. **B.** MHco3/10.BC$_4$.IVM. **C.** MHco3/10.BC$_4$.IVM.P$_3$. **D.** MHco3/10.BC$_4$.IVM.P$_4$. **E.** MHco3/4.BC$_4$.noIVM. **F.** MHco3/4.BC$_4$.IVM. **G.** MHco3/4.BC$_4$.IVM.P$_3$. **H.** MHco3/4.BC$_4$.IVM.P$_4$.

Figure S6. Location of significantly non-neutral loci identified using the single-locus population genetic model. Corresponding peaks in the location of significant sites can be seen in the MHco3/10 (**A**) and MHco3/4 datasets (**B**). A total of 70.6% of significant sites in the MHco3/10 dataset, and 90.6% of significant sites in the MHco3/4 dataset, were found in chromosome V. Data are binned in 1 Mb windows. Colours represent chromosomes as described in Figure 2A.

Figure S7. Analysis of Tajima's D variation in each chromosome per cross.
Comparison of Tajima's D per chromosome between MHco3(ISE) parent (blue),
MHco10(CAVR) (panel column 1; red) or MHco4(WRS) (panel column 3; red) and
passages 3 (orange) and 4 (yellow) of the crosses. Tajima's D was calculated using
npstat in 100 kb windows spanning the genome. The variance in the mean value of
Tajima's D among MHco3(ISE) and passages 3 and 4 – for which an increase in
variance would suggest introgression and evidence of selection – was determined
and is presented as smoothed line (red) in panel columns 2 and 4.

A

B

1014 **Figure S9. Relative position and location of candidate genes from the literature**
1015 **proposed to be associated with ivermectin resistance in *Haemonchus***
1016 ***contortus* and / or *Caenorhabditis elegans.*** Gene coordinates are presented in
1017 Table S2. Colours represent chromosomes as described in Figure 2A.

Table S2. Candidate genes from literature proposed to be associated with

ivermectin resistance in *Haemonchus contortus* and / or *Caenorhabditis*

elegans

Gene	GeneID[3]	Chromosome	Coordinates[4]	Reference
avr-14b / gbr-2 or HcGluClα3 [2]	HCOI00274000, HCOI00274100	I	28352540 - 28362260	(Williamson et al., 2011)
avr-15	HCOI00130400	V	45145104 - 45158307	(Dent et al., 2000)
tbb-1	HCOI01967600	I	7026384 - 7029311	(Eng et al., 2006; de Lourdes Mottier and Prichard, 2008)
che-3 [1]	HCOI01710300	I	11666953 - 11695321	(Dent et al., 2000)
dyf-11 [1]	HCOI01514900	I	20494101 - 20505946	(Dent et al., 2000)
dyf-7	HCOI01065900	Xb	8461163 - 8465904	(Urdaneta-Marquez et al., 2014)
ggr-3	HCOI00098800	II	45626226 - 45639348	(Rao et al., 2009)
glc-1 [1]	NA			(Dent et al., 2000)
glc-2 / GluClb [2]	HCOI00383000	I	1443439 - 1446121	
glc-3	HCOI00543300	V	27643249 - 27655188	(Williamson et al., 2011)
glc-4	HPLM_0001206301	II	36314509 - 36331683	
glc-5 / HcGluCla [2]	HCOI00617300	I	40196936 - 40206359	(Blackhall et al., 1998)
lgc-36	HCOI02054500	V	9008092 - 9028181	
lgc-37 / HG-1 [2]	HCOI00977100	III	15741515 - 15762899	(Blackhall et al., 2003)
lgc-55	HCOI00162900	V	42450788 - 42469177	
osm-1 [1]	HCOI00970500	II	3105809 - 3126440	

osm-3	HCOI00164700, HCOI00673700			
osm-5 [1]	HCOI00498400, HCOI00077900			
pgp-1 / pgp-9 [2]	HCOI00233200	V	47249715 - 47265234	(James and Davey, 2009; Raza et al., 2016; van Wyk and Malan, 1988)
pgp-2 / pgp-a [2]	HCOI00025600	I	5849505 - 5868032	(Raza et al., 2016)
pgp-11	HCOI00622400	II	31321675 - 31344154	(Raza et al., 2016)
unc-38	HCOI01939400	I	25118411 - 25125593	
Unc-7 [1]	HPLM_0000412001			(Dent et al., 2000)
Unc-9 [1]	HCOI02075100	Xb	4410309 - 4424525	(Dent et al., 2000)
che-11	HCOI00062600	II	3237676 - 3245201	
osm-6	HCOI00937900	V	28370676 - 28397433	
pgp-3	HCOI00117000	II	11888372 - 11913805	(Raza et al., 2016)
haf-6	HCOI01007100	IV	44336050 - 44341372	(Raza et al., 2016)
pgp-13 / pgp-12 [1]	HCOI01115500	II	10272077 - 10286519	(David et al., 2018)
mrp-1 [1]	HCOI01904000	Xb	3325452 - 3334565	(James and Davey, 2009)

1. *C. elegans* genes

2. Alternate names in the literature

3. Gene IDs correspond to the annotated gene in the published version of the MHco3(ISE)

genome (Laing et al., 2013).

4. Coordinates are approximate, as they represent the transfer of coordinates from the

published version based on homology, and therefore may have changed depending on the

conservation of the gene structure in the genome between the published and updated

genome version used here.

Figure S10: Haplotype structure of chromosome V in an example output from
the model under neutral evolution. Segments of genome from the resistant parent
are shown in red, while segments of genome from the susceptible parent are shown
in blue. The repeated backcross removes most of the resistant genotypes from the
population.

References

Abkallo, H.M., Martinelli, A., Inoue, M., Ramaprasad, A., Xangsayarath, P., Gitaka, J., Tang, J., Yahata, K., Zoungrana, A., Mitaka, H., et al. (2017). Rapid identification of genes controlling virulence and immunity in malaria parasites. PLoS Pathog. *13*, e1006447.

Blackhall, W.J., Pouliot, J.-F., Prichard, R.K., and Beech, R.N. (1998). Haemonchus contortus: Selection at a Glutamate-Gated Chloride Channel Gene in Ivermectin- and Moxidectin-Selected Strains. Exp. Parasitol. *90*, 42–48.

Blackhall, W.J., Prichard, R.K., and Beech, R.N. (2003). Selection at a γ-aminobutyric acid receptor gene in Haemonchus contortus resistant to avermectins/milbemycins. Mol. Biochem. Parasitol. *131*, 137–145.

Britton, C., Roberts, B., and Marks, N.D. (2016). Functional Genomics Tools for Haemonchus contortus and Lessons From Other Helminths. Adv. Parasitol. *93*, 599–623.

Chiang, C., Layer, R.M., Faust, G.G., Lindberg, M.R., Rose, D.B., Garrison, E.P., Marth, G.T., Quinlan, A.R., and Hall, I.M. (2015). SpeedSeq: ultra-fast personal genome analysis and interpretation. Nat. Methods *12*, 966–968.

Choi, Y.-J., Bisset, S.A., Doyle, S.R., Hallsworth-Pepin, K., Martin, J., Grant, W.N., and Mitreva, M. (2017). Genomic introgression mapping of field-derived multiple-anthelmintic resistance in Teladorsagia circumcincta. PLoS Genet. *13*, e1006857.

Crellen, T., Walker, M., Lamberton, P.H.L., Kabatereine, N.B., Tukahebwa, E.M., Cotton, J.A., and Webster, J.P. (2016). Reduced Efficacy of Praziquantel Against Schistosoma mansoni Is Associated With Multiple Rounds of Mass Drug Administration. Clin. Infect. Dis. *63*, 1151–1159.

David, M., Lebrun, C., Duguet, T., Talmont, F., Beech, R., Orlowski, S., André, F., Prichard, R.K., and Lespine, A. (2018). Structural model, functional modulation by ivermectin and tissue localization of Haemonchus contortus P-glycoprotein-13. Int. J. Parasitol. Drugs Drug Resist.

Denham, D.A. (1969). The Development of Ostertagia circumcincta in Lambs. J. Helminthol. *43*, 299.

Dent, J.A., Smith, M.M., Vassilatis, D.K., and Avery, L. (2000). The genetics of ivermectin resistance in Caenorhabditis elegans. Proc. Natl. Acad. Sci. U. S. A. *97*, 2674–2679.

Doyle, S.R., Bourguinat, C., Nana-Djeunga, H.C., Kengne-Ouafo, J.A., Pion, S.D.S., Bopda, J., Kamgno, J., Wanji, S., Che, H., Kuesel, A.C., et al. (2017). Genome-wide analysis of ivermectin response by Onchocerca volvulus reveals that genetic drift and soft selective sweeps contribute to loss of drug sensitivity. PLoS Negl. Trop. Dis.

11, e0005816.

Doyle, S.R., Laing, R., Bartley, D.J., Britton, C., Chaudhry, U., Gilleard, J.D., Holroyd, N., Mable, B.K., Maitland, K., Morrison, A.A., et al. (2018). A genome resequencing-based genetic map reveals the recombination landscape of an outbred parasitic nematode in the presence of polyploidy and polyandry. Genome Biol. Evol *10*, 396–409.

Echevarria, F., and Trindade, G. (1989). Anthelmintic resistance by Haemonchus contortus to ivermectin in Brazil: a preliminary report. Vet. Rec. *124*, 147–148.

Elard, L., and Humbert, J.F. (1999). Importance of the mutation of amino acid 200 of the isotype 1 β-tubulin gene in the benzimidazole resistance of the small-ruminant parasite Teladorsagia circumcincta. Parasitol. Res. *85*, 452–456.

Eng, J.K.L., Blackhall, W.J., Osei-Atweneboana, M.Y., Bourguinat, C., Galazzo, D., Beech, R.N., Unnasch, T.R., Awadzi, K., Lubega, G.W., and Prichard, R.K. (2006). Ivermectin selection on β-tubulin: Evidence in Onchocerca volvulus and Haemonchus contortus. Mol. Biochem. Parasitol. *150*, 229–235.

Ewels, P., Magnusson, M., Lundin, S., and Käller, M. (2016). MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics *32*, 3047–3048.

Feder, A.F., Petrov, D.A., and Bergland, A.O. (2012). LDx: estimation of linkage disequilibrium from high-throughput pooled resequencing data. PLoS One *7*, e48588.

Ferretti, L., Ramos-Onsins, S.E., and Pérez-Enciso, M. (2013). Population genomics from pool sequencing. Mol. Ecol. *22*, 5561–5576.

Geary, T.G. (2016). Haemonchus contortus: Applications in Drug Discovery. Adv. Parasitol. *93*, 429–463.

Geerts, S., and Gryseels, B. (2000). Drug Resistance in Human Helminths: Current Situation and Lessons from Livestock. Clin. Microbiol. Rev. *13*, 207–222.

Gilleard, J.S. (2013). Haemonchus contortus as a paradigm and model to study anthelmintic drug resistance. Parasitology *140*, 1506–1522.

Hefnawy, A., Berg, M., Dujardin, J.-C., and De Muylder, G. (2017). Exploiting Knowledge on Leishmania Drug Resistance to Support the Quest for New Drugs. Trends Parasitol. *33*, 162–174.

Hill, W.G., and Weir, B.S. (1988). Variances and covariances of squared linkage disequilibria in finite populations. Theor. Popul. Biol. *33*, 54–78.

Howe, K.L., Bolt, B.J., Shafie, M., Kersey, P., and Berriman, M. (2017). WormBase ParaSite − a comprehensive resource for helminth genomics. Mol. Biochem. Parasitol. *215*, 2–10.

Hunt, P. (1995). Identification and characterisation of dominant ivermectin resistant genes in Caenorhabditis elegans. PhD. University of New England.

Illingworth, C.J.R., and Mustonen, V. (2013). Quantifying selection in evolving populations using time-resolved genetic data. J. Stat. Mech: Theory Exp. *2013*, P01004.

Illingworth, C.J.R., Parts, L., Schiffels, S., Liti, G., and Mustonen, V. (2012). Quantifying selection acting on a complex trait using allele frequency time series data. Mol. Biol. Evol. *29*, 1187–1197.

Jabbar, A., Campbell, A.J.D., Charles, J.A., and Gasser, R.B. (2013). First report of anthelmintic resistance in Haemonchus contortus in alpacas in Australia. Parasit. Vectors *6*, 243.

James, C.E., and Davey, M.W. (2009). Increased expression of ABC transport proteins is associated with ivermectin resistance in the model nematode Caenorhabditis elegans. Int. J. Parasitol. *39*, 213–220.

Kaplan, R.M., and Vidyashankar, A.N. (2012). An inconvenient truth: global worming and anthelmintic resistance. Vet. Parasitol. *186*, 70–78.

Kofler, R., Pandey, R.V., and Schlötterer, C. (2011). PoPoolation2: identifying differentiation between populations using sequencing of pooled DNA samples (Pool-Seq). Bioinformatics *27*, 3435–3436.

Kotze, A.C., Hunt, P.W., Skuce, P., von Samson-Himmelstjerna, G., Martin, R.J., Sager, H., Krücken, J., Hodgkinson, J., Lespine, A., Jex, A.R., et al. (2014). Recent advances in candidate-gene and whole-genome approaches to the discovery of anthelmintic resistance markers and the description of drug/receptor interactions. Int. J. Parasitol. Drugs Drug Resist. *4*, 164–184.

Kwa, M.S.G., Veenstra, J.G., and Roos, M.H. (1994). Benzimidazole resistance in Haemonchus contortus is correlated with a conserved mutation at amino acid 200 in β-tubulin isotype 1. Mol. Biochem. Parasitol. *63*, 299–303.

Laing, R., Kikuchi, T., Martinelli, A., Tsai, I.J., Beech, R.N., Redman, E., Holroyd, N., Bartley, D.J., Beasley, H., Britton, C., et al. (2013). The genome and transcriptome of Haemonchus contortus, a key model parasite for drug and vaccine discovery. Genome Biol. *14*, R88.

Laing, R., Maitland, K., Lecová, L., Skuce, P.J., Tait, A., and Devaney, E. (2016). Analysis of putative resistance gene loci in UK field populations of Haemonchus contortus after 6 years of macrocyclic lactone use. Int. J. Parasitol. *46*, 621–630.

Le Jambre, L.F., Gill, J.H., Lenane, I.J., and Lacey, E. (1995). Characterisation of an avermectin resistant strain of Australian Haemonchus contortus. Int. J. Parasitol. *25*, 691–698.

Le Jambre, L.F., Gill, J.H., Lenane, I.J., and Baker, P. (2000). Inheritance of

avermectin resistance in Haemonchus contortus. Int. J. Parasitol. *30,* 105–111.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

de Lourdes Mottier, M., and Prichard, R.K. (2008). Genetic analysis of a relationship between macrocyclic lactone and benzimidazole anthelmintic selection on Haemonchus contortus. Pharmacogenet. Genomics *18,* 129–140.

Luo, X., Shi, X., Yuan, C., Ai, M., Ge, C., Hu, M., Feng, X., and Yang, X. (2017). Genome-wide SNP analysis using 2b-RAD sequencing identifies the candidate genes putatively associated with resistance to ivermectin in Haemonchus contortus. Parasit. Vectors *10,* 31.

Maciuca, S., del Ojo Elias, C., McVean, G., and Iqbal, Z. (2016). A natural encoding of genetic variation in a Burrows-Wheeler Transform to enable mapping and genome inference.

Nisbet, A.J., Meeusen, E.N., González, J.F., and Piedrafita, D.M. (2016). Immunity to Haemonchus contortus and Vaccine Development. In Advances in Parasitology, pp. 353–396.

Omura, S., and Crump, A. (2004). The life and times of ivermectin - a success story. Nat. Rev. Microbiol. *2,* 984–989.

Osei-Atweneboana, M.Y., Awadzi, K., Attah, S.K., Boakye, D.A., Gyapong, J.O., and Prichard, R.K. (2011). Phenotypic evidence of emerging ivermectin resistance in Onchocerca volvulus. PLoS Negl. Trop. Dis. *5,* e998.

Parts, L., Cubillos, F.A., Warringer, J., Jain, K., Salinas, F., Bumpstead, S.J., Molin, M., Zia, A., Simpson, J.T., Quail, M.A., et al. (2011). Revealing the genetic structure of a trait by sequencing a population under selection. Genome Res. *21,* 1131–1138.

Paten, B., Novak, A.M., Eizenga, J.M., and Garrison, E. (2017). Genome graphs and the evolution of genome inference. Genome Res. *27,* 665–676.

Rao, V.T.S., Siddiqui, S.Z., Prichard, R.K., and Forrester, S.G. (2009). A dopamine-gated ion channel (HcGGR3*) from Haemonchus contortus is expressed in the cervical papillae and is associated with macrocyclic lactone resistance. Mol. Biochem. Parasitol. *166,* 54–61.

Raza, A., Kopp, S.R., Bagnall, N.H., Jabbar, A., and Kotze, A.C. (2016). Effects of in vitro exposure to ivermectin and levamisole on the expression patterns of ABC transporters in Haemonchus contortus larvae. Int. J. Parasitol. Drugs Drug Resist. *6,* 103–115.

Redman, E., Sargison, N., Whitelaw, F., Jackson, F., Morrison, A., Bartley, D.J., and Gilleard, J.S. (2012). Introgression of ivermectin resistance genes into a susceptible Haemonchus contortus strain by multiple backcrossing. PLoS Pathog. *8,* e1002534.

Rezansoff, A.M., Laing, R., and Gilleard, J.S. (2016). Evidence from two independent

backcross experiments supports genetic linkage of microsatellite Hcms8a20, but not other candidate loci, to a major ivermectin resistance locus in Haemonchus contortus. Int. J. Parasitol. *46*, 653–661.

Roos, M.H., Otsen, M., Hoekstra, R., Veenstra, J.G., and Lenstra, J.A. (2004). Genetic analysis of inbreeding of two strains of the parasitic nematode Haemonchus contortus. Int. J. Parasitol. *34*, 109–115.

Rose, H., Rinaldi, L., Bosco, A., Mavrot, F., de Waal, T., Skuce, P., Charlier, J., Torgerson, P.R., Hertzberg, H., Hendrickx, G., et al. (2015). Widespread anthelmintic resistance in European farmed ruminants: a systematic review. Vet. Rec. *176*, 546.

Rufener, L., Kaminsky, R., and Mäser, P. (2009). In vitro selection of Haemonchus contortus for benzimidazole resistance reveals a mutation at amino acid 198 of β-tubulin. Mol. Biochem. Parasitol. *168*, 120–122.

Sargison, N.D., Redman, E., Morrison, A.A., Bartley, D.J., Jackson, F., Gijzel, H.N., Holroyd, N., Berriman, M., Cotton, J.A., and Gilleard, J.S. (2018). A method for single pair mating in an obligate parasitic nematode. Int. J. Parasitol. *48*, 159–165.

Schwarz, E.M., Korhonen, P.K., Campbell, B.E., Young, N.D., Jex, A.R., Jabbar, A., Hall, R.S., Mondal, A., Howe, A.C., Pell, J., et al. (2013). The genome and developmental transcriptome of the strongylid nematode Haemonchus contortus. Genome Biol. *14*, R89.

Silvestre, A., and Cabaret, J. (2002). Mutation in position 167 of isotype 1 β-tubulin gene of Trichostrongylid nematodes: role in benzimidazole resistance? Mol. Biochem. Parasitol. *120*, 297–300.

Urdaneta-Marquez, L., Bae, S.H., Janukavicius, P., Beech, R., Dent, J., and Prichard, R. (2014). A dyf-7 haplotype causes sensory neuron defects and is associated with macrocyclic lactone resistance worldwide in the nematode parasite Haemonchus contortus. Int. J. Parasitol. *44*, 1063–1071.

Williamson, S.M., Storey, B., Howell, S., Harper, K.M., Kaplan, R.M., and Wolstenholme, A.J. (2011). Candidate anthelmintic resistance-associated gene expression and sequence polymorphisms in a triple-resistant field isolate of Haemonchus contortus. Mol. Biochem. Parasitol. *180*, 99–105.

van Wyk, J.A., and Malan, F.S. (1988). Resistance of field strains of Haemonchus contortus to ivermectin, closantel, rafoxanide and the benzimidazoles in South Africa. Vet. Rec. *123*, 226–228.

Xie, C., and Tammi, M.T. (2009). CNV-seq, a new method to detect copy number variation using high-throughput sequencing. BMC Bioinformatics *10*, 80.

Yadav, C.L., Kumar, R., Uppal, R.P., and Verma, S.P. (1995). Multiple anthelmintic resistance in Haemonchus contortus on a sheep farm in India. Vet. Parasitol. *60*, 355–360.