1 **A catalog of single nucleotide changes distinguishing modern humans from archaic hominins**
2

3 Martin Kuhlwilm[1], Cedric Boeckx[2,3,4]
4 [1] Universitat Pompeu Fabra, Institute for Evolutionary Biology, Barcelona, Spain
5 [2] ICREA, Barcelona, Spain
6 [3] University of Barcelona, Barcelona, Spain
7 [4] UB Institute of Complex Systems, Barcelona, Spain

8
9
Abstract

Throughout the past decade, studying ancient genomes provided unique insights into human
prehistory, and differences between modern humans and other branches like Neanderthals can
enrich our understanding of the molecular basis of unique modern human traits. Modern human
variation and the interactions between different hominin lineages are now well studied, making it
reasonable to go beyond fixed changes and explore changes that are observed at high frequency in
present-day humans. Here, we identify 571 genes with non-synonymous changes at high frequency.
We suggest that molecular mechanisms in cell division and networks affecting cellular features of
neurons were prominently modified by these changes. Complex phenotypes in brain growth
trajectory and cognitive traits are likely influenced by these networks and other changes presented
here. We propose that at least some of these changes contributed to uniquely human traits, and
should be prioritized for experimental validation.

Corresponding author contact information:
Cedric Boeckx
Email: cedric.boeckx@ub.edu
Phone: +34 93 403 3782
ICREA/Universitat de Barcelona
Gran Via de les Corts Catalanes, 585
08007 Barcelona, Spain

Introduction

Homo sapiens appears to be a "very special primate" (Pääbo 2014). Our position among animal species stands out largely thanks to the composite complexity of our cultures, social structures and communication systems. It seems reasonable that this "human condition" is rooted, at least in part, in the properties of our brain, and that these can be traced to changes in the genome on the modern human lineage. This phenotype in the population called "anatomically modern humans" emerged in Africa likely before the deepest divergence less than 100,000-200,000 years ago (Schlebusch et al. 2012; Kuhlwilm et al. 2016), although complex population structure may reach back up to 300,000 years ago (Hublin et al. 2017; Schlebusch et al. 2017; Skoglund et al. 2017). Except for some early dispersals (Rabett 2018), humans most likely peopled other parts of the world than Africa and the Middle East permanently only after around 65,000 years ago. It has been claimed that the brain of modern humans adopted a specific, apomorphic growth trajectory early in life that gave rise to the skull shape difference between modern humans and extinct branches of the genus *Homo* (Hublin et al. 2015). Importantly, the growth pattern might differ between the populations (Gunz et al. 2010; Neubauer et al. 2018), while the brain size and encephalization of humans and Neanderthals is similar, with slightly larger brains in the latter (Trinkaus and Howells 1979; Schoenemann 2004; Hublin et al. 2015). This ontogenic trajectory, termed the "globularization phase", might have contributed to cognitive changes that underlie behavioral traits in which humans differ from their extinct relatives, despite mounting evidence for their cognitive sophistication (Gunz et al. 2012; Hublin et al. 2015; Wynn et al. 2016; Boeckx 2017; Hoffmann et al. 2018).

We are now in a favorable position to examine the evolution of human biology with the help of the fossil record, in particular thanks to breakthroughs in paleogenomics: The recent reconstruction of the high quality genomes of members of archaic *Homo* populations (Meyer et al. 2012; Prüfer et al. 2014; Prüfer et al. 2017) has opened the door to new comparative genomic approaches and molecular analyses. The split of the lineages leading to modern humans and other archaic forms (Neanderthals and Denisovans) is estimated to around 600,000 years ago (Kuhlwilm et al. 2016), setting the timeframe for truly modern human-specific changes after this split, but before the divergence of modern human populations (Fig. 1). Together with efforts to explore present-day human diversity (Auton et al. 2015), this progress has allowed to narrow down the number of candidate point mutations from ~35 million differences since the split from chimpanzee when comparing only reference genomes (Consortium 2005) to 31,389 fixed human-specific changes in a previous seminal study (Pääbo 2014). Other types of more complex changes like structural variants most likely contributed to human-specific traits. For example, it is well known that since the split from chimpanzees functional differences arose through gene duplications in *ARHGAP11B* and other genes (Florio et al. 2015; Ju et al. 2016), copy number variants in *SRGAP2* and other genes (Dennis et al. 2012; Dumas et al. 2012; Suzuki et al. 2018) or regulatory deletions (McLean et al. 2011). In

these cases, the variants arose before the split of humans and Neanderthals, but the differences in structural variation that exist between the hominin lineages (Chintalapati et al. 2017) need to be explored in more detail, with advancement of technologies in ancient DNA sequencing and computational methods. This will result in complementary lists of changes for understanding the human condition outside the scope of this study. Beyond that, parts of the genome which are complex and not yet examined by conventional sequencing platforms (O'Bleness et al. 2012) possibly harbor important human-specific changes as well.

Some of the single nucleotide changes have been linked to putative functional consequences (Castellano et al. 2014; Pääbo 2014; Prüfer et al. 2014), and evidence is mounting that several molecular changes affecting gene expression in the brain were subject to selective pressures (Green et al. 2010; Somel et al. 2013; Zhou et al. 2015; Racimo 2016; Peyrégne et al. 2017). Furthermore, the genomic impact of interbreeding events is not evenly distributed across the genome. Genes expressed in regions of the brain regarded as critical for certain cognitive functions such as language are depleted in introgressed archaic genetic material (Sankararaman et al. 2014; Vernot and Akey 2014; Sankararaman et al. 2016; Vernot et al. 2016), and introgressed alleles are downregulated in some of these brain regions, suggesting natural selection acting on tissue-specific gene regulation (McCoy et al. 2017). Thus, it seems reasonable to conclude that there were differences between anatomically modern human and Neanderthal brains, and that these underlie at least some of the characteristics of our lineage (Wynn and Coolidge 2004). We want to emphasize that such recent differences are likely to be subtle when compared to those after the split from our closest living relatives on a scale of 6-10 million years (Langergraber et al. 2012), where fundamental changes arose since the divergence from chimpanzees and bonobos (Varki and Altheide 2005; O'Bleness et al. 2012). The observation of recurrent gene flow between modern human and archaic populations also implies a broad overall similarity, yet, such subtle differences may still have contributed to the evolutionary outcome (Wynn et al. 2016). This does not imply a superiority of humans, but specific changes that might have facilitated survival under the given environmental conditions. Obviously, not all human-specific changes are beneficial: While most mutations may be rather neutral and have little effect on the phenotype, some may have had deleterious effects or side-effects, possibly increasing the risks for neurodevelopmental or neurodegenerative disorders in humans (Bufill et al. 2011; Bruner and Jacobs 2013; Bufill et al. 2013).

The goal of this paper is to provide a revised, extended set of recent single nucleotide changes in humans since their split from Neanderthals that could enrich our understanding of the molecular basis of recent human condition. The previous focus on fixed alleles was reasonable given limited data (Pääbo 2014), but having a better grasp of the magnitude of modern human variation and the interaction between different hominin lineages seems a good reason to cast a wider net, and take into account not only fixed differences but also high-frequency changes shared by more than 90%

of present-day individuals. Here, we present a revised list of 36 genes that carry missense substitutions which are fixed across thousands of human individuals and for which all archaic hominin individuals sequenced so far carry the ancestral state. In total, 647 protein-altering changes in 571 genes reached a frequency of at least 90% in the present-day human population. We attempt to interpret this list, as well as some regulatory changes, since it seems very likely that some of these genes would have contributed to the human condition. We discuss some of their known functions, and how these relate to pathways that might have been modified during human evolution from the molecular level to cellular features and more complex phenotypic traits (Fig. 1). We restrict our attention to genes where the literature may allow firm conclusions and predictions about functional effects, since many genes likely have multiple different functions (Gratten and Visscher 2016). Obviously, it cannot be emphasized enough that ultimately, experimental validation will be needed to confirm our hypotheses concerning alterations in specific functions. For example, transcription factors or enzymatically active proteins can be tested using cell cultures or *in vitro* assays, while brain organoids could be used to test differences in neuronal functions (Giandomenico and Lancaster 2017), especially in combination with single-cell RNA sequencing (Camp et al. 2015; Camp and Treutlein 2017). Ultimately, these variants can be introduced into model organisms like mice to test complex features related to cognitive abilities or behavior (Enard et al. 2009). Still, given limitations to the amount of changes that can be tested at once, networks which are modified by multiple changes cannot be tested with current technologies.

Results

Genetic differences between present-day humans and archaic hominins

Using publicly available data on one Denivosan and two Neanderthal individuals and present-day human variation (Methods), we calculated the numbers of single nucleotide changes (SNCs) which most likely arose recently on the respective lineages after their split from each other, and their functional consequences (Table 1). Previously, a number of 31,389 sites has been reported as recently fixed derived in present-day humans, while being ancestral in archaics (Pääbo 2014; Prüfer et al. 2014). We find a smaller number of only 12,027 positions in the genome, in part because of including another archaic individual and different filters, but mainly by a richer picture of present-day human variation. The 1,000 Genomes Project as well as other sources contributing to the dbSNP database now provide data for thousands of individuals, which results in very high allele frequencies for many loci instead of fixation. Indeed, 29,358 positions show allele frequencies larger than 0.995, demonstrating that the level of near-fixation is similar to the level of previously presented fixation. The number of loci with high frequency (HF) changes of more than 90% in

present-day humans is an order of magnitude larger than the number of fixed differences. The three
archaic individuals carry more than twice as many changes than present-day humans; however, we
emphasize that much of this difference is not due to more mutations in archaics, but rather the fact
that data for only three individuals is available, compared to thousands of humans. The variation
across the archaic population is not represented equally well, which makes these numbers not
directly comparable. On the other hand, much less variation is found by the sequencing of each
additional Neanderthal individual compared to humans due to the low diversity of Neanderthals
(Fig. S36 in (Kuhlwilm et al. 2016)). This low diversity across their geographic range suggests that
most alleles observed as ancestral here will be the same state in other individuals. Furthermore, we
take variability into account due to gene flow or errors, decreasing the possibility that positions
ancestral in the archaic individuals studied to date turn out to be derived in most archaic individuals,
hence this extended catalog will likely not undergo drastic changes. However, changes in structural
variants or regions of the genome that are not accessible by current sequencing technologies will
most likely complement our results (O'Bleness et al. 2012).

Present-day humans carry 42 fixed amino acid-changes in 36 genes (Table 2, Fig. 2), while
Neanderthals carry 159 such changes. Additionally, modern humans carry 605 amino acid-changes
at high frequency (human-lineage high-frequency missense changes, referred to as HHMCs),
amounting to a total of 647 such changes in 571 genes (Table S1). Together with 323 SNCs on the
human lineage with low confidence (Methods, Table S2), almost 1,000 putative protein-altering
changes were found across most present-day humans. Generally, synonymous changes are found at
a similar magnitude as missense changes, but only few SNCs altering start and stop codons, and
thousands of changes in putative regulatory and untranslated regions. We admit that some of the
loci presented here are variable across the phylogenetic tree, or less reliable due to low coverage in
the archaics, but we accept this since our intention is retrieve an inclusive picture of possibly
functional recent changes. The 42 protein-altering changes for which the ancestral allele has not
been observed in any present-day human, most of which have been presented before (Pääbo 2014),
constitute without doubt the strongest entry points into a molecular understanding of the human
condition, and should be prime candidates for experimental validation. Only one gene, *SPAG5*,
carries three such SNCs, and four genes (*ADAM18, CASC5, SSH2* and *ZNHIT2*) carry two fixed
protein-coding changes in all modern humans. We identified 15 SNCs (in *AHR, BOD1L1,*
C1orf159, C3, DNHD1, DNMT3L, FRMD8, OTUD5, PROM2, SHROOM4, SIX5, SSH2, TBC1D3,
ZNF106, ZNHIT2) that have not been previously described as fixed differences between humans
and archaics. We note that another 12 previously described (Pääbo 2014) protein-altering
substitutions were not found among the genotypes analyzed here (in *C21orf62, DHX29,*
FAM149B1, FRRS1L, GPT, GSR, HERC5, IFI44L, KLF14, PLAC1L, PTCD2, SCAF11). These
genotype calls are absent from the files provided for the three archaic genomes due to different
genotype calling and filtering procedures compared to the original publication of the Altai

Neanderthal genome (Prüfer et al. 2014; Prüfer et al. 2017). Hence, some potentially relevant
candidate changes were not included here, and future research is necessary to evaluate these as well.
Despite attempting an extended interpretation, our data is thus not fully exhaustive.

It is noteworthy that the number of fixed SNCs decreased substantially, and it is possible that
single individuals will be found to carry some of the ancestral alleles for the remaining fixed sites.
Hence, it is important to focus not only on fixed differences, but also consider variants at high
frequency. When analyzing the 647 HHMCs, 68 genes carry more than one amino acid-altering
change. Among these, *TSGA10IP* (Testis Specific 10 Interacting Protein) and *ABCC12* (ATP
Binding Cassette Subfamily C Member 12) carry four such changes, and seven more genes
(*MUC5B, NPAP1, OR10AG1, OR5M9, PIGZ, SLX4, VCAN*) carry three HHMCs. 1,542 genes carry
at least one HF missense change on the archaic lineage (archaic-lineage high-frequency missense
change, referred to as AHMC, Tables S3, S4). We find an overlap of 122 genes with HHMCs and
AHMCs, which is more than expected considering that among 1,000 sets of random genes of a
similar length distribution, no overlap of this extent was observed. The same genes seem to have
acquired missense changes on both lineages since their divergence more often than expected. We
find a high ratio of HHMCs over synonymous changes for chromosome 21 (1.75-fold), and a very
small ratio (0.18-fold) for chromosome 13. We do not find such extreme ratios for AHMCs and
corresponding synonymous changes, suggesting differences in the distribution of amino acid
changes between both lineages (Fig. S1).

	Fixed human	HF human	Extended human	Fixed archaic	HF archaic	Extended archaic
All	12,027	136,435	83,254	33,498	380,756	983
Non-synonymous	42	647	327	167	1,921	13
Synonymous	41	843	363	193	2,123	14
Start/stop	1	14	10	3	48	2
Splice site	4	23	8	4	54	0
TFBS	28	226	126	87	914	1
Upstream	1,935	19,599	11,235	4,920	55,188	289
5' UTR	180	1,853	1,012	195	2,016	7
3' UTR	77	702	334	506	5,303	19
Downstream	1,922	19,704	11,673	4,956	55,832	281
miRNA	0	1	2	0	4	0
Regulatory element	1,952	20,971	12,320	5,125	59,248	195

198 **Table 1**: Summary of single nucleotide changes. TFBS: Transcription factor binding sites. UTR:
199 Untranslated Region. HF: High frequency. Fixed changes are a subset of HF changes.

200
201
202

Ranking and enrichment

We assessed the impact of mutations for different deleteriousness scores (Table 2), finding 12 genes with deleterious HHMCs according to SIFT, three according to PolyPhen, and 16 when using the Grantham score (>180), measuring the physical properties of amino acid changes. The C-score and GWAVA can be used to rank all mutation classes, and we present the top candidates.

Then, we attempted a ranking of genes by the density of lineage-specific changes in the dataset. As expected, the total number of segregating sites is correlated with gene length (Pearsons' R = 0.93). This correlation is weaker for HF human SNCs (R = 0.73) and fixed human-specific SNCs (R = 0.25), as well as for fixed (R = 0.37) and HF (R = 0.82) SNCs in archaics. We conclude that some genes with a large number of human-specific changes might carry these large numbers by chance, while others are depleted. Indeed, 17,453 (88.9%) of these genes do not carry any fixed human-specific change, and 80.5% do not carry fixed archaic-specific changes. Of note, genes that have attracted attention in the context of traits related to the "human condition" like *CNTNAP2* and *AUTS2* are among the longest genes in the genome, hence changes in these genes should be interpreted with caution as they are not unexpected. We ranked the genes by the number of HF changes in either modern humans or archaics, divided by their genomic lengths, and categorize the top 5% of this distribution as putatively enriched for changes on each lineage (Table S5). We note that 191 genes (30.9%) fall within this category for both human HF changes and archaic HF changes, as a result of differences in mutation density. In order to distinguish a truly lineage-specific enrichment, we calculated the ratios of HF changes for humans and archaics, defining the top 10% of genes in this distribution as putatively enriched (Table S5). Among the genes enriched for changes on the modern human lineage, 18 carry no HF changes on the archaic lineage, and ten of these also fall within the 5% of genes carrying many changes considering their length (*ARSJ, CLUAP1, COL20A1, EPPIN, KLHL31, MKNK1, PALMD, RIC3, TDRD7, UBE2H*). These might be candidates for an accumulation of changes, even though this is not identical to selective sweep signals. Among these, the collagen COL20A1 and the Epididymal Peptidase Inhibitor EPPIN carry HHMCs. *ACAD10, DST* and *TTC40*, which carry two HHMCs, might be other notable genes with a human-specific enrichment.

Gene Ontology (GO) categories are neither enriched for HHMCs on the human lineage in a hypergeometric test, nor for genes carrying AHMCs, HF changes in UTRs or transcription factor binding sites. However, instead of singular changes that might be observed more often in long genes, or genes that are more prone to mutations in hominins, the density of HF changes in a gene might yield a better picture of lineage-specific changes, possibly for cumulative changes. We applied a test for the ratio of the number of gene-wise HF changes on one lineage over the other lineage, finding an enrichment for 12 GO categories on the human lineage (Table S6), with "soft palate development", "negative regulation of adenylate cyclase activity", "collagen catabolic process" and "cell adhesion" in the biological process category. Among the cellular components

category, the "postsynaptic membrane", "spermatoproteasome complex", "collagen trimer", "dendrite" and "cell junction" show enrichment, as well as the molecular functions "calcium ion binding", "histone methyltransferase activity (H3-K27 specific)" and "metallopeptidase activity". We find no GO enrichment for genes with an excess of changes on the archaic lineage. In order to approach a deeper exploration of genes with associated complex traits in humans, we explored the NHGRI-EBI GWAS Catalog (MacArthur et al. 2017), containing 2,385 traits. We performed a systematic enrichment screen, finding 17 unique traits enriched for genes with HHMCs, and 11 for genes with AHMCs (Table S7). Changes in genes associated to "Cognitive decline (age-related)", "Rheumatoid arthritis" or "Major depressive disorder" might point to pathways that could have been influenced by protein-coding changes on the human lineage. In archaics, genes are enriched, among others, for associations to traits related to body mass index or cholesterol levels, which might reflect differences in their physiology.

We find a significant enrichment of protein-protein interactions (P = 0.006) among the gene products of HHMC genes (Fig. S2), meaning that these proteins interact with each other more than expected. Functional enrichment is found for the biological process "cellular component assembly involved in morphogenesis", most strongly for the cellular components cytoskeleton and microtubule, as well as the molecular function "cytoskeletal protein binding". Three proteins have at least 20 interactions in this network and might be considered important nodes: TOP2A, PRDM10 and AVPR2 (Table S8). However, proteins encoded by genes with synonymous changes on the modern human lineage seem to be enriched for interactions as well (P = 0.003), as are proteins encoded by genes with AHMCs (P = 1.68 x 10^{-14}), with an enrichment in GO categories related to the extracellular matrix and the cytoskeleton, and proteins with more than 40 interactions (Table S8). We caution that these networks might be biased due to more mutations and possibly more interactions in longer, multi-domain genes.

Regulatory changes might have been important during our evolution (Wray 2007), hence we tested for an overrepresentation of transcription factors. We find 78 known or putative transcription factors among the HHMC genes (Table S9) on the modern human lineage (Chawla et al. 2013), which is not overrepresented among genes with HHMCs (with 49.2% of random genes sets containing fewer HHMCs). Despite no enrichment as a category, single transcription factors on the modern human lineage might have been important, particularly those with an excess of modern human over archaic HF changes (*AHR, MACC1, PRDM2, TCF3, ZNF420, ZNF516*). Others, like *RB1CC1* (Prüfer et al. 2014) or *PRDM10* and *NCOA6* (Peyrégne et al. 2017) have been found in selective sweep screens, suggesting contributions of individual transcription factors, rather than the class of proteins. We also tested for an enrichment of gene expression in different brain regions and developmental stages (Miller et al. 2014; Grote et al. 2016), using the HF synonymous changes on each lineage as background sets. We find an enrichment of gene expression in the orbital frontal cortex at infant age (0-2 years) for genes with HHMCs, but no enrichment for genes with AHMCs.

277 Furthermore, when testing the genes with HHMCs and using the set of genes with AHMCs as
278 background, "gray matter of forebrain" at adolescent age (12-19 years) is enriched, while no
279 enrichment was found for genes with AHMCs.

Fixed HHMCs	ADAM18, ADSL, AHR, ANKMY1, ANKRD30A, BBIP1, BOD1L1, C1orf159, C3, CASC5, CDH16, DCHS1, DNHD1, DNMT3L, FRMD8, GBP5, GLDC, GREB1L, GRM6, KIF26B, LMNB2, NCOA6, NOTO, OTUD5, PRDM10, PROM2, RFNG, SCAP, SHROOM4, SIX5, SPAG5, SSH2, TBC1D3, ZNF106, ZNF185, ZNHIT2
Selection 2014	C11orf80, CKAP5, GREB1L, HMCN1, NLRX1, PDZD3, PRDM2, RB1CC1
Selection 2015	MSS51, NCOA6, OMD, SPAG17, SPAG5
Selection 2016	ACE, ADSL, ALMS1, ANKRD30A, BZRAP1, DNAH1, GREB1L, KMT2C, NWD1, PROM2, RASA1, STAB1, STARD9, ZNF106
Selection 2017	ADSL, AKAP8, BAP1, BBIP1, BCAR3, CAPN5, CR2, CSMD2, DNAH1, ENTHD1, FAAH, FRMD8, GBP5, GBP7, GPR157, GTF3C5, HERC5, HERC6, HMCN1, HRASLS5, KATNA1, KIF15, KIF18A, LYST, MKL1, MYH3, NAALADL1, NCOA6, PRDM10, PRDM2, PROM2, PTPRC, RNF44, SCAP, SLC12A8, SLC25A45, SLITRK1, TIGD3, TMEM235, TRGV4, TTC6, VOPP1, ZNF501, ZNF502, ZNHIT2
Grantham	ABHD14A-ACY1, ACY1, ABHD14A, CCDC158, CCDC30, DNHD1, EML2, ERI1, GBA3, GREB1, OR1K1, TTC6, UBQLN3, UIMC1, ZBP1, ZNF510, ZNHIT2
SIFT	BEND2, CCT6B, COPA, CUL4B, GBP7, KRTAP10-10, MEPE, NHEJ1, OR1K1, SLC6A15, TPO, ZNF510
PolyPhen-2	FSHR, NLN, TPO
CADD	C11orf80, C5orf66, CCT6B, CDH15, CEP128, CPM, FGF21, FMN2, FUT1, H2AFY, HERC6, KCNK5, KPNA4, KRT33A, KRT8P12, MUM1, NR1H2, OPRM1, PDSS2, ROCK1, RPS15P9, SLC22A31, SUCLG2P4, TMPRSS7, UNC5D
GWAVA	ANK2, COPA, CTRC, CYP2B6, MAPK10, MCTP1, SLC38A6, SYT1, YTHDC1

Table 2: Genes with fixed non-synonymous changes on the human lineage, genes under positive selection
with HHMCs, and deleterious candidate HHMCs. Selection 2014: Prüfer *et al.*, 2014. Selection 2015: Zhou
et al., 2015. Selection 2016: Racimo, 2016. Selection 2017: Peyrégne *et al.*, 2017.

Discussion

The enrichment of broad categories above suggests that traits related to brain functions are
prominently represented by HHMCs. It should be noted that such results would be less clear if we
just focused on completely fixed changes, given the drastically reduced number of genes harboring
such changes. Here, we will further examine the possible impact on the brain that some of these
changes might have, paying special attention to hypotheses formulated in earlier work on modern
human-specific changes. Our extended catalog of changes appears to provide additional support for
some of these hypotheses.

Cell division and the brain growth trajectory

It has been proposed previously that protein-coding changes in cell cycle-related genes are highly
relevant candidates for human-specific traits (Pääbo 2014; Prüfer et al. 2014). Indeed, three genes
(*CASC5*, *SPAG5*, and *KIF18A*) have been singled out as involved in spindle pole assembly during
mitosis (Pääbo 2014). Other genes with protein-coding SNCs (*NEK6* and *STARD9/KIF16A*) turn

out to be implicated in the regulation of spindle pole assembly as well (O'Regan and Fry 2009; Torres et al. 2017). Among the 15 fixed protein-coding changes identified here but absent from previous analyses (Pääbo 2014; Prüfer et al. 2014), some might have also contributed to complex modifications of pathways in cell division, like *AHR* (Puga et al. 2002) or *DNHD1* (Bader et al. 2011) (Supplementary Information 1), as well as other genes with HHMCs, like *CHEK1* (Zachos et al. 2017) or the gene encoding for the protein TOP2A (Yoshida and Azuma 2016), which shows the largest number of interactions with other HHMC-carrying proteins, suggesting a function as interaction hub in the cell division complex (Supplementary Information 1). Taken together, these changes suggest that the cell cycle machinery might have been modified in a specific way in humans compared to other hominins.

It has been claimed (Prüfer et al. 2014) that genes with fixed non-synonymous changes in humans are also more often expressed in the ventricular zone of the developing neocortex, compared to fixed synonymous changes. Since the kinetochore-associated genes *CASC5, KIF18A* and *SPAG5* are among these genes, it has been emphasized that this "may be relevant phenotypically as the orientation of the mitotic cleavage plane in neural precursor cells during cortex development is thought to influence the fate of the daughter cells and the number of neurons generated (Fietz and Huttner 2011)" (Prüfer et al. 2014). Several fixed SNCs on the modern human lineage are observed for *CASC5* (two changes) and *SPAG5* (three changes), which is also among genes with a relatively high proportion of HF changes (Table S5). The changes in *KIF18A, KIF16A* and *NEK6* can no longer be considered as fixed, but occur at very high frequencies (>99.9%) in present-day humans. We attempted to determine whether an enrichment of genes with HHMCs on the human lineage can be observed in the ventricular zone (Miller et al. 2014), but instead find an enrichment in the intermediate zone, where less than 5% of random gene sets of the same size are expressed. However, synonymous HF changes also show an enrichment in this layer, as well as genes with AHMCs (Table S10), suggesting an overrepresentation of genes that carry mutations in the coding regions rather than lineage-specific effects. We were able to broadly recapitulate the observation of an enrichment of expression in the ventricular zone if restricting the test to genes with non-synonymous changes at a frequency greater than 99.9% in present-day humans, which is not observed for corresponding synonymous and archaic non-synonymous changes (Table S10). Among the 28 genes expressed in the ventricular zone that carry almost fixed HHMCs, four might be enriched for HF changes in humans (*HERC5, LMNB2, SPAG5, VCAM1*), and one shows an excess of HF changes on the human compared to the archaic lineage (*AMKMY1*). Other notable genes discussed in this study include *ADSL, FAM178A, KIF26B, SLC38A10*, and *SPAG17*.

The centrosome-cilium interface is known to be critical for early brain development, and centrosome-related proteins are overrepresented in studies on the microcephaly phenotype in humans (Megraw et al. 2011). We find 126 genes (Table S9) with 143 HHMCs that putatively interact with proteins at the centrosome-cilium interface (Gupta et al. 2015). Some of the genes

listed here and discussed in this study, such as *FMR1, KIF15, LMNB2, NCOA6, RB1CC1, SPAG5*
and *TEX2*, harbor not only HHMCs, but an overall high proportion of HF changes on the human
lineage. Although an early analysis suggested several candidate genes associated to microcephaly,
not all of these could be confirmed by high-coverage data. Among eleven candidate genes (Green et
al. 2010), only two (*PCNT, UCP1*) are among the HHMC gene list presented here, while most of
the other changes are not human-specific, and only *PCNT* has been related to microcephaly (Li et
al. 2015). Nevertheless, more changes related to microcephaly are found on both lineages, for
example in *ATRX* (K. Ritchie et al. 2014) or *CASC5* (Genin et al. 2012) (Supplementary
Information 3).

Changes in genes associated with brain growth trajectory differences lead not necessarily to a
decrease but also an increase of brain size (Montgomery et al. 2011), suggesting that the disease
phenotype of macrocephaly might point to genes relevant in the context of brain growth as well.
One of the few genes with several HHMCs, *CASC5*, has been found to be associated with gray
matter volume differences (Shi et al. 2017). It has been claimed that mutations in *PTEN* alter the
brain growth trajectory and allocation of cell types through elevated Beta-Catenin signaling (Chen
et al. 2015). This well-known gene, critical for brain development (Li et al. 2017), has not been
highlighted in the context of human-specific changes, while we find that *PTEN* falls among the
genes with an excess on the modern human over the archaic lineage, suggesting that regulatory
changes in this gene might have contributed to human-specific traits. This is also the case for the
HHMC-carrying transcription factor TCF3, which is known to repress Wnt-Beta-Catenin signaling
and maintain the neural stem cell population during neocortical development (Kuwahara et al.
2014). Changes in these and other genes (Supplementary Information 3) like *CCND2* (Mirzaa et al.
2014), *GLI3* (Jamsheer et al. 2012), or *RB1CC1* (Wang et al. 2013), for which a regulatory SNC
has been suggested to modify transcriptional activity (Weyer and Pääbo 2016) and which carries a
signature of positive selection (Prüfer et al. 2014), could have contributed to the brain growth
trajectory changes hypothesized to give rise to the modern human-specific globular braincase shape
during the past several 100,000 years (Gunz et al. 2012; Hublin et al. 2015; Neubauer et al. 2018).
Finally, we find changes that might have affected the size of the cerebellum, a key contributor to our
brain shape (Kochiyama et al. 2018; Neubauer et al. 2018), such as HF regulatory SNCs in *ZIC1*
and *ZIC4* (Blank et al. 2011), an excess of HF mutations in *AHI1* (Cheng et al. 2012), and a
deleterious HHMC in *ABHD14A*, which is a target of ZIC1 (Hoshino et al. 2003).

Cellular features of neurons

To form critical networks during the early development of the brain, axonal extensions of the
neurons in the cortical region must be sent and guided to eventually reach their synaptic targets.
Studies conducted on avian vocal learners (Pfenning et al. 2014; Wang et al. 2015) have shown a
convergent differential regulation of axon guidance genes of the *SLIT-ROBO* families in the pallial

motor nucleus of the learning species, allowing for the formation of connections virtually absent in
the brains of vocal non-learners. In modern humans, genes with axon-guidance-related functions
such as *FOXP2, SLIT2* and *ROBO2* have been found to lie within deserts of archaic introgression
(Sankararaman et al. 2016; Vernot et al. 2016; Kuhlwilm 2018), suggesting incompatibilities
between modern humans and archaics for these regions. Our dataset contains a fair amount of genes
known to impact brain wiring: Some of the aforementioned microtubule-related genes, specifically
those associated with axonal transport and known to play a role in post-mitotic neural wiring and
plasticity (Lüders 2016), are associated with signals of positive selection, such as *KIF18A*
(McVicker et al. 2016) or *KATNA1* (Ahmad et al. 1999; Karabay et al. 2004). Furthermore, an
interactor of KIF18A, KIF15 (Kevenaar et al. 2017), might have been under positive selection in
modern humans (Peyrégne et al. 2017), and contains two HHMCs. Versican (*VCAN*), which
promotes neurite outgrowth (Wu et al. 2004), carries three HHMCs, and *SSH2* (two HHMCs) might
be involved in neurite outgrowth (Cuberos et al. 2015). *PIEZO1*, which carries a non-synonymous
change that is almost fixed in modern humans, is another factor in axon guidance (Koser et al.
2016), as well as *NOVA1* (Jensen et al. 2000), which is an interactor of *ELAVL4* (Ratti et al. 2008), a
gene that codes for a neuronal-specific RNA-binding protein and might have been under positive
selection in humans (Zhou et al. 2015; Peyrégne et al. 2017). Furthermore, we find one of the most
deleterious regulatory SNCs in the Netrin receptor UNC5D, which is critical for axon guidance
(Takemoto et al. 2011).

We also detect changes in genes associated with myelination and synaptic vesicle endocytosis,
critical to sustain a high rate of synaptic transmission, including *DCX* (Yap et al. 2012), *SCAP*
(Verheijen et al. 2009), *RB1CC1* (Menzies et al. 2015), *ADSL* (Jurecka et al. 2012) and *PACSIN1*
(Widagdo et al. 2016) among others (Supplementary Information 2). It is noteworthy that among
traits associated with cognitive functions such as language or theory of mind, the timing of
myelination appears to be a good predictor of computational abilities (Skeide and Friederici 2016;
Grosse Wiesmann et al. 2017). Computational processing might have been facilitated by some of
the changes presented here, at least in some of the circuits that have expanded in our lineage (Mars
et al. 2018), since subtle maturational differences early in development (Dubois et al. 2016) may
have had a considerable impact on the phenotype. In this context, it is worth mentioning that in our
dataset, several genes carrying HHMCs and associated with basal ganglia functions (critical for
language and cognition) stand out, like *SLITRK1* (Abelson et al. 2005) and *NOVA1* (Jelen et al.
2010; Konopka et al. 2012; Alsiö et al. 2013; Zhou et al. 2015; Popovitchenko et al. 2016)
(Supplementary Information 4). Finally, in the broader context of cognition, we find an enrichment
of HHMCs in genes associated to "Alzheimer's disease (cognitive decline)" and "Cognitive decline
(age-related)", with seven associated genes (*COX7B2, BCAS3, DMXL1, LIPC, PLEKHG1, TTLL2*
and *VIT*). Among genes influencing behavioral traits (Supplementary Information 4) are *GPR153*
(Sreedharan et al. 2011), *NCOA6* (Takata et al. 2018), or the Adenylosuccinate Lyase (*ADSL*) (Fon

et al. 1995), for which the ancestral Neanderthal-like allele has not been observed in 1,000s of
modern human genomes and which has been pointed out before as under positive selection
(Castellano et al. 2014; Racimo et al. 2014; Racimo 2016; Peyrégne et al. 2017) We know that
archaic hominins likely had certain language-like abilities (Dediu and Levinson 2013; Dediu and
Levinson 2018), and hybrids of modern and archaic humans must have survived in their
communities (Fu et al. 2015), underlining the large overall similarity of these populations.
However, genes associated with axon guidance functions, which are important for the refinement of
neural circuits including those relevant for speech and language, are found in introgression deserts
(Jeong et al. 2016; Lei et al. 2017), which seems to be a unidirectional and human-specific pattern
especially in the *FOXP2* region (Kuhlwilm 2018). We suggest that modifications of a complex
network in cognition or learning took place in modern human evolution (Boeckx and Benítez-
Burraco 2014), possibly related to other brain-related (Bastir et al. 2011; Hublin et al. 2015; Boeckx
2017; Bryant and Preuss 2018), vocal tract (Gokhman et al. 2017) or neural changes (Belyk and
Brown 2017).

The craniofacial phenotype

In previous work on ancient genomes changes related to craniofacial morphology have been
highlighted (Castellano et al. 2014; Gokhman et al. 2017), and we find an enrichment of genes with
an excess of HF SNCs on the modern human lineage for soft palate development (Table S6).
Among genes harboring an excess of HF SNCs associated with specific facial features, we find
RUNX2, EDAR, and *GLI3* (Adhikari et al. 2016), *NFATC1* (Kim and Kim 2014), *SPOP* (Cai and
Liu 2016), *DDR2* (Zhang et al. 2011) and *NELL1* (Zhang et al. 2012), possibly carrying changes in
regulatory regions, while mutations in the HHMC-carrying gene encoding for the transcription
factor ATRX cause facial dysmorphism (Moncini et al. 2013). In addition, genes with HHMCs such
as *PLXNA2* (Oh et al. 2012), *EVC2* (Kwon et al. 2018), *MEPE* (Gullard et al. 2016), *OMD*
(Tashima et al. 2015), and *SPAG17* (Teves et al. 2015) are known to affect craniofacial bone and
tooth morphologies. These genes appear to be important in determining bone density, mineralization
and remodeling, hence they may underlie differences between archaic and modern human facial
growth (Lacruz et al. 2015). Some of these facial properties may have been present in the earliest
fossils attributed to *H. sapiens,* like the Jebel Irhoud fossils (Hublin et al. 2017), deviating from
craniofacial features which emerged in earlier forms of *Homo* (Lacruz et al. 2013), and may have
become established before some brain-related changes discussed here (Stringer 2016; Neubauer et
al. 2018). The gene encoding the transcription factor PRDM10 stands out for carrying HHMCs,
being found in selective sweep regions and the second-most interacting protein within the HHMC
dataset. Although little is known about *PRDM10*, it may be related to dendrite growth (Siegel et al.
2002) and neural crest related changes that contributed to the formation of our distinct modern face
(Park and Kim 2010). Other craniofacial morphology-related genes, such as *DCHS2* (Adhikari et al.

2016), *HIVEP2* (Jones et al. 2010), *HIVEP3* (Imamura et al. 2014), *FREM1* (Lee et al. 2017), and

FRAS1 (Talbot et al. 2016) harbor AHMCs, while another bone-related gene, *MEF2C* (Verzi et al.

2007), shows an excess of HF changes on the archaic lineage. These changes may underlie some of

the prominent derived facial traits of Neanderthals (Rak 1986).

Life history and other phenotypic traits

Apart from their consequences for cognitive functions, it has been suggested that changes

involved in synaptic plasticity might be interpreted in a context of neoteny (Somel et al. 2009; Liu

et al. 2012; Peyrégne et al. 2017; Sherwood and Gómez-Robles 2017), with the implication of

delayed maturation in humans (Bednarik 2013) and a longer timeframe for brain development.

However, given their similar brain sizes (Hofman 1983), humans and Neanderthals might both have

needed a long overall maturation time (Ponce de León et al. 2017; Rosas et al. 2017). Accordingly,

notions like neoteny and heterochrony are unlikely to be fine-grained enough to capture differences

between these populations, but early differences in infant brain growth between humans and

Neanderthals (Gunz et al. 2010; Hublin et al. 2015) could have rendered our maturational profile

distinct during limited developmental periods and within specific brain regions, imposing different

metabolic requirements (Bruner et al. 2014). One of the brain regions where such differences are

found is the orbitofrontal cortex (OFC) (Bastir et al. 2011), and we find that the OFC at infant age

(0-2 years) is enriched for the expression of genes that carry HHMCs compared to synonymous

SNCs. We suggest that the development of the OFC in infants might have been subject to subtle

changes since the split from Neanderthals rather than a general developmental delay, which is

particularly interesting given that this brain region has been implied in social cognition (Beer et al.

2006) and learning (Miller et al. 2018).

Genes carrying HHMCs are enriched for expression in the gray matter of the forebrain at the

adolescent age compared to AHMC-carrying genes, hence additional human-specific modifications

during this period might have taken place, possibly linked to changes in myelination described

above. It has been suggested that differences in childhood adolescence time existed between

humans and Neanderthals, after a general developmental delay in the hominin lineage (Smith and

Tompkins 1995; Bock and Sellen 2002). Dental evidence suggests an earlier maturation in

Neanderthals than modern humans (Smith et al. 2010), and it has been claimed that Neanderthals

might have reached adulthood earlier (Ramirez Rozzi and de Castro 2004). Furthermore, an

introgressed indel from Neanderthals causes an earlier onset of menarche in present-day humans

(Chintalapati et al. 2017), supporting at least the existence of alleles for earlier maturation in the

Neanderthal population. Among the genes carrying fixed HHMCs, *NCOA6* has also been linked to

age at menarche and onset of puberty (Day et al. 2017), as well as placental function (Antonson et

al. 2003). This putative transcription factor is enriched in HF changes and has been suggested to

have been under positive selection on the modern human lineage (Racimo et al. 2014; Peyrégne et

al. 2017). The HHMC is located nearby and three 5'-UTR variants within a putatively selected region (Zhou et al. 2015), with an estimated time of selection at around 150 kya (assuming a slow mutation rate). Even though this gene carries an AHMC as well, it remains possible that modern humans acquired subtle differences in their reproductive system through lineage-specific changes in this gene. A delay in reproductive age may influence overall longevity, another trait for which our data set yields an enrichment of genes with HHMCs (*SLC38A10*, *TBC1D22A* and *ZNF516*).

The male reproductive system might have been subject to changes as well, since we find that several proteins in spermatogenesis seem to carry two HHMCs: Sperm Specific Antigen 2 (*SSFA2*), Sperm Associated Antigen 17 (*SPAG17*), *ADAM18* (Zhu et al. 1999) and *WDR52* (Tang et al. 2017), out of which *ADAM18* and *SPAG17* also carry AHMCs. Lineage-specific differences in genes related to sperm function or spermatogenesis might have been relevant for the genetic compatibility between humans and Neanderthals. Another gene harboring a HHMC with similar functions is *EPPIN* (Wang et al. 2005), which shows no HF changes on the archaic, but 27 such SNCs on the modern human lineage. The gene encoding for the Testis Expressed 2 protein (*TEX2*) is enriched for HF changes in both humans and archaics, with one HHMC and five AHMCs, but its function is not yet known. Another possible SNC that might be relevant in this context is a splice site change in *IZUMO4*, since proteins encoded by the IZUMO family form complexes on mammalian sperm (Ellerman et al. 2009). The adjacent exon is not present in all transcripts of this gene, suggesting a functional role of this splice site SNC. Finally, genes in the GO category "spermatoproteasome complex" are enriched for an excess of HF changes on the human lineage.

It has been found that Neanderthal alleles contribute to addiction and, possibly, pain sensitivity in modern humans (Simonti et al. 2016; Dannemann et al. 2017). In this context, an interesting protein-truncating SNC at high frequency in humans is the loss of a stop codon in the opioid receptor OPRM1 (6:154360569), potentially changing the structure of the protein encoded by this gene in some transcripts. Other mutations in this gene are associated to heroin addiction (Shi et al. 2002), and pain perception (Tan et al. 2009), but also sociality traits (Pearce et al. 2017). Interestingly, a recent study found a pain insensitivity disorder caused by a mutation in *ZFHX2* (Habib et al. 2017), which carries an AHMC, and three HHMCs are observed in *NPAP1*, which might be associated with the Prader-Willi syndrome, involving behavioral problems and a high pain threshold (Buiting et al. 2007). Such changes may point to differences in levels of resilience to pain between Neanderthals and modern humans.

Conclusion

The long-term evolutionary processes that led to the human condition (Pääbo 2014) is still subject to debate and investigation, and the high-quality genomes from archaic humans provide opportunities to explore the recent evolution of our species. We want to contribute to an attempt to

unveil the genetic basis of specific molecular events in the time-window after the split from these archaic populations and before the emergence of most of the present-day diversity. We sought to combine different sources of information, from genome-wide enrichment analyses to functional information available for specific genes, to identify threads linking molecular needles in this expanded haystack. In doing so, we have mainly built on existing proposals concerning brain-related changes, but we have divided the observations into different biological levels, from cellular changes through brain organization differences to complex phenotypic traits. Only future experimental work will determine which of the changes highlighted here contributed significantly to making us "fully human". We hope that our characterization and presentation of some new candidate genes will help prioritize inquiry in this area, since the specific type of validation depends on each candidate gene or network.

Methods

We used the publicly available high-coverage genotypes for three archaic individuals: One Denisovan (Meyer et al. 2012), one Neanderthal from the Denisova cave in Altai mountains (Prüfer et al. 2014), and another Neanderthal from Vindija cave, Croatia (Prüfer et al. 2017). The data is publicly available under http://cdna.eva.mpg.de/neandertal/Vindija/VCF/, with the human genome version *hg19* as reference, covering ~1.8 billion base pairs of the genome (Prüfer et al. 2017). We applied further filtering to remove sites with less than 5-fold coverage and more than 105-fold coverage in the Altai Neanderthal or 75-fold coverage in the other archaic individuals, if such cases occurred. We also removed sites with genotype quality smaller than 20, and heterozygous sites with strong allele imbalance (<0.2 minor allele frequency). Although these permissive filters increase power compared to previous studies, we caution that in some cases genotypes might be incorrect. We added the genotype and coverage for the exome and chromosome 21 sequences of the Vindija and El Sidrón Neanderthals from previous studies (Castellano et al. 2014; Kuhlwilm et al. 2016), with 75-fold and 50-fold coverage cutoffs, respectively. These studies provided data for the same Vindija individual (Prüfer et al. 2017).

We applied the Ensembl Variant Effect Predictor VEP (McLaren et al. 2016) in order to obtain inferences for protein-coding and regulatory mutations, scores for SIFT (Kumar et al. 2009), PolyPhen (Adzhubei et al. 2010), CADD (Kircher et al. 2014) and GWAVA (G.R.S. Ritchie et al. 2014), and allele frequencies in the 1000 Genomes and ExAC human variation databases (Auton et al. 2015; Lek et al. 2016). We used the inferred ancestral allele from published data on multiple genome alignments (Paten et al. 2008), and at positions where this information was not available, the macaque reference allele, *rheMac3 (Yan et al. 2011)*. We determined the allele frequencies in present-day humans using the dbSNP database build 147 (Sherry et al. 2001). We retrieved the

counts for each allele type, and summarized the counts of non-reference alleles at each position.

Grantham scores (Grantham 1974) were calculated for missense mutations.

Data processing and database retrieval was performed using bcftools/samtools v1.0 (Li 2011),

bedtools v2.16.2 (Quinlan and Hall 2010), and R/Bioconductor (Huber et al. 2015), with rtracklayer

(Lawrence et al. 2009) and biomaRt (Durinck et al. 2005) packages, and plotting with Rcircos

(Zhang et al. 2013). We analyzed all positions where at least two alleles (human reference and

alternative allele) were observed among the human reference and at least one out of three of the

high-coverage archaic individuals, in at least one archaic chromosome. The 22 autosomal

chromosomes and the X chromosome were analyzed, in the absence of Y chromosome data for the

three female archaic individuals. The data for 4,409,518 segregating sites is available under

[http:tbd.database]. The following subsets were created:

Fixed differences: Positions where all present-day humans carry a derived allele, while at least

two out of three archaics carry the ancestral allele, accounting for potential human gene flow into

Neanderthals.

High-frequency (HF) differences: Positions where more than 90% of present-day humans carry a

derived allele, while at least the Denisovan and one Neanderthal carry the ancestral allele,

accounting for different types of errors and bi-directional gene flow.

Extended high-frequency differences: Positions where more than 90% of present-day humans

carry a derived allele, while one of the following conditions is true: a) Not all archaics have reliable

genotypes, but those that have carry the ancestral allele. b) Some archaics carry an alternative

genotype that is not identical to either the human or the ancestral allele. c) The Denisovan carries

the ancestral allele, while one Neanderthal carries a derived allele, which allows for gene flow from

humans into Neanderthals. d) The ancestral allele is missing in the EPO alignment, but the macaque

reference sequence is identical to the allele in all three archaics.

We also created corresponding lists of archaic-specific changes. Fixed changes were defined as

sites where the three archaics carry the derived allele, while humans carry the ancestral allele at

more than 99.999%. High-frequency changes occur to less than 1% in present-day humans, while at

least two archaic individuals carry the derived allele. An extended list presents high-frequency

changes where the ancestral allele is unknown, but the macaque allele is identical to the present-day

human allele.

A ranking of mutation density was performed for genes with protein-coding sequences and their

genomic regions as retrieved from Ensembl. For each gene, unique associated changes as predicted

by VEP were counted. A ranking on the number of HF changes per gene length was performed for

all genes that span at least 5,000 bp in the genome and carry at least 25 segregating sites in the

dataset (at any frequency in humans or in archaics), in order to remove genes which are very short

or poor in mutations. The top 5% of the empirical distribution was defined as putatively enriched

for changes on each lineage. The ratio of lineage-specific HF changes was calculated for the subset

of genes where at least 20 lineage-specific HF changes were observed on the human and the archaic
lineages combined. The top 10% of the empirical distribution was defined as putatively enriched for
lineage-specific changes.

We performed enrichment tests using the R packages ABAEnrichment (Grote et al. 2016) and
DescTools (Signorell 2017). We used the NHGRI-EBI GWAS Catalog (MacArthur et al. 2017), and
overlapped the associated genes with protein-coding changes on the human and archaic lineages,
respectively. We counted the number of HF missense changes on each lineage and the subset of
those associated to each trait ("Disease trait"), and performed a significance test (G-test) against the
number of genes associated to each trait, and all genes in the genome, with a P value cutoff at 0.1.
This suggests a genome-wide enrichment of changes for each trait. We then performed a G-test
between the numbers of HF missense changes on each lineage, and the subset of each associated to
each trait (P-value cutoff at 0.1), to determine a difference between the two lineages. We then
performed an empirical test by creating 1,000 random sets of genes with similar length as the genes
associated to each trait, and counting the overlap to the lineage-specific missense changes. At least
90% of these 1,000 random sets were required to contain fewer missense changes than the real set
of associated genes. Only traits were considered for which at least 10 associated loci were
annotated.

Gene Ontology (GO) enrichment was performed using the software FUNC (Prüfer et al. 2007),
with a significance cutoff of the adjusted p-value < 0.05 and a family-wise error rate < 0.05. When
testing missense changes, a background set of synonymous changes on the same lineage was used
for the hypergeometric test. When testing genes with relative mutation enrichment, the Wilcoxon
rank test was applied. Enrichment for sequence-specific DNA-binding RNA polymerase II
transcription factors and transcription factor candidate genes from (Chawla et al. 2013), and genes
interacting at the centrosome-cilium interface (Gupta et al. 2015) was tested with an empirical test
in which 1,000 random sets of genes were created that matched the length distributions of the genes
in the test list. The same strategy was applied for genes expressed in the developing brain (Table
S10) (Miller et al. 2014). Protein-protein interactions were analyzed using the STRING online
interface v10.5 (Szklarczyk et al. 2017) with standard settings (medium confidence, all sources,
query proteins only) as of January 2018. The overlap with selective sweep screens considers
HHMCs within 50,000 bp of the selected regions (Prüfer et al. 2014; Zhou et al. 2015; Peyrégne et
al. 2017).
626

Figure Legends

Figure 1: Conceptual summary of this study.

Figure 2: Features discussed in this study. From inside to outside: Genes with HHMCs and
signatures of positive selection (compare Table 2), genes with fixed non-synonymous SNCs on the
human lineage, HHMCs, AHMCs, karyogram of human chromosomes.

Acknowledgments
We thank S. Han and T. Marques-Bonet for helpful discussions, and A. G. Andirkó and P .T. Martins for
help with figures. M.K. is supported by a Deutsche Forschungsgemeinschaft (DFG) fellowship (KU 3467/1-
1). C.B. acknowledges research funds from the Spanish Ministry of Economy and Competitiveness/FEDER
(grant FFI2016-78034-C2-1-P), Marie Curie International Reintegration Grant from the European Union
(PIRG-GA-2009-256413), research funds from the Fundació Bosch i Gimpera, MEXT/JSPS Grant-in-Aid
for Scientific Research on Innovative Areas 4903 (Evolinguistics: JP17H06379), and Generalitat de
Catalunya (Government of Catalonia) – 2017-SGR-341.

Author contributions
M.K. and C.B. analyzed data and wrote the manuscript.

Competing interests statement
The authors declare no competing interests.

References

Abelson JF, Kwan KY, O'Roak BJ, Baek DY, Stillman AA, Morgan TM, Mathews CA, Pauls DL, Rašin M-R, Gunel M, et al. 2005. Sequence Variants in SLITRK1 Are Associated with Tourette's Syndrome. Science (80-.). [Internet] 310:317 LP-320. Available from: http://science.sciencemag.org/content/310/5746/317.abstract

Adhikari K, Fuentes-Guajardo M, Quinto-Sánchez M, Mendoza-Revilla J, Camilo Chacón-Duque J, Acuña-Alonzo V, Jaramillo C, Arias W, Lozano RB, Pérez GM, et al. 2016. A genome-wide association scan implicates DCHS2, RUNX2, GLI3, PAX1 and EDAR in human facial variation. Nat. Commun. [Internet] 7:11616. Available from: http://dx.doi.org/10.1038/ncomms11616

Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS, Sunyaev SR. 2010. A method and server for predicting damaging missense mutations. Nat. Methods 7:248–249.

Ahmad FJ, Yu W, McNally FJ, Baas PW. 1999. An Essential Role for Katanin in Severing Microtubules in the Neuron. J. Cell Biol. [Internet] 145:305 LP-315. Available from: http://jcb.rupress.org/content/145/2/305.abstract

Alsiö JM, Tarchini B, Cayouette M, Livesey FJ. 2013. Ikaros promotes early-born neuronal fates in the cerebral cortex. Proc. Natl. Acad. Sci. [Internet] 110:E716–E725. Available from: http://www.pnas.org/content/110/8/E716.abstract

Antonson P, Schuster GU, Wang L, Rozell B, Holter E, Flodby P, Treuter E, Holmgren L, Gustafsson J-Å. 2003. Inactivation of the Nuclear Receptor Coactivator RAP250 in Mice Results in Placental Vascular Dysfunction. Mol. Cell. Biol. [Internet] 23:1260–1268. Available from: http://mcb.asm.org/content/23/4/1260.abstract

Auton A, Abecasis GR, Altshuler DM, Durbin RM, Bentley DR, Chakravarti A, Clark AG, Donnelly P, Eichler EE, Flicek P, et al. 2015. A global reference for human genetic variation. Nature [Internet] 526:68–74. Available from: http://dx.doi.org/10.1038/nature15393

Bader JR, Kasuboski JM, Winding M, Vaughan PS, Hinchcliffe EH, Vaughan KT. 2011. Polo-like Kinase1 Is Required for Recruitment of Dynein to Kinetochores during Mitosis. J. Biol. Chem. [Internet] 286:20769–20777. Available from: http://www.jbc.org/content/286/23/20769.abstract

Bastir M, Rosas A, Gunz P, Peña-Melian A, Manzi G, Harvati K, Kruszynski R, Stringer C, Hublin J-J. 2011. Evolution of the base of the brain in highly encephalized human species. Nat. Commun. [Internet] 2:588. Available from: http://dx.doi.org/10.1038/ncomms1593

Bednarik GR. 2013. From Human Past to Human Future. Humanit. 2.

Beer JS, John OP, Scabini D, Knight RT. 2006. Orbitofrontal Cortex and Social Behavior: Integrating Self-monitoring and Emotion-Cognition Interactions. J. Cogn. Neurosci. [Internet] 18:871–879. Available from: https://doi.org/10.1162/jocn.2006.18.6.871

Belyk M, Brown S. 2017. The origins of the vocal brain in humans. Neurosci. Biobehav. Rev. [Internet] 77:177–193. Available from: http://www.sciencedirect.com/science/article/pii/S0149763416306583

Blank MC, Grinberg I, Aryee E, Laliberte C, Chizhikov V V, Henkelman RM, Millen KJ. 2011. Multiple developmental programs are altered by loss of Zic1 and Zic4 to cause Dandy-Walker malformation cerebellar pathogenesis. Development [Internet] 138:1207 LP-1216. Available from: http://dev.biologists.org/content/138/6/1207.abstract

Bock J, Sellen DW. 2002. Childhood and the evolution of the human life course. Hum. Nat. [Internet] 13:153–159. Available from: https://doi.org/10.1007/s12110-002-1006-5

Boeckx C. 2017. The language-ready head: Evolutionary considerations. Psychon. Bull. Rev. [Internet] 24:194–199. Available from: https://doi.org/10.3758/s13423-016-1087-5

Boeckx C, Benítez-Burraco A. 2014. Globularity and language-readiness: generating new predictions by expanding the set of genes of interest . Front. Psychol. [Internet] 5:1324. Available from: https://www.frontiersin.org/article/10.3389/fpsyg.2014.01324

Bruner E, Jacobs HIL. 2013. Alzheimer's disease: the downside of a highly evolved parietal lobe? J. Alzheimers. Dis. 35:227–240.

Bruner E, de la Cuétara JM, Masters M, Amano H, Ogihara N. 2014. Functional craniology and brain evolution: from paleontology to biomedicine. Front. Neuroanat. [Internet] 8:19. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3980103/

Bryant KL, Preuss TM. 2018. A Comparative Perspective on the Human Temporal Lobe BT - Digital Endocasts: From Skulls to Brains. In: Bruner E, Ogihara N, Tanabe HC, editors. Tokyo: Springer Japan. p. 239–258. Available from: https://doi.org/10.1007/978-4-431-56582-6_16

Bufill E, Agustí J, Blesa R. 2011. Human neoteny revisited: The case of synaptic plasticity. Am. J. Hum. Biol. [Internet] 23:729–739. Available from: http://dx.doi.org/10.1002/ajhb.21225

Bufill E, Blesa R, Augusti J. 2013. Alzheimer's disease: an evolutionary approach. J. Anthropol. Sci. = Riv. di Antropol. JASS 91:135–157.

Buiting K, Nazlican H, Galetzka D, Wawrzik M, Groβ S, Horsthemke B. 2007. C15orf2 and a novel noncoding transcript from the Prader–Willi/Angelman syndrome region show monoallelic expression in fetal brain. Genomics [Internet] 89:588–595. Available from: http://www.sciencedirect.com/science/article/pii/S0888754306003703

Cai H, Liu A. 2016. Spop promotes skeletal development and homeostasis by positively regulating Ihh signaling. Proc. Natl. Acad. Sci. [Internet] 113:14751–14756. Available from: http://www.pnas.org/content/113/51/14751.abstract

Camp JG, Badsha F, Florio M, Kanton S, Gerber T, Wilsch-Bräuninger M, Lewitus E, Sykes A, Hevers W, Lancaster M, et al. 2015. Human cerebral organoids recapitulate gene expression programs of fetal neocortex development. Proc. Natl. Acad. Sci. U. S. A. [Internet] 112:15672–15677. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4697386&tool=pmcentrez&rendertype=abstract

Camp JG, Treutlein B. 2017. Human organomics: a fresh approach to understanding human development using single-cell transcriptomics. Development [Internet] 144:1584 LP-1587. Available from: http://dev.biologists.org/content/144/9/1584.abstract

Castellano S, Parra G, Sánchez-Quinto FA, Racimo F, Kuhlwilm M, Kircher M, Sawyer S, Fu Q, Heinze A, Nickel B, et al. 2014. Patterns of coding variation in the complete exomes of three Neandertals. Proc. Natl. Acad. Sci. U. S. A. [Internet] 111:6666–6671. Available from: http://www.pnas.org/content/111/18/6666

Chawla K, Tripathi S, Thommesen L, Lægreid A, Kuiper M. 2013. TFcheckpoint: a curated compendium of specific DNA-binding RNA polymerase II transcription factors. Bioinformatics [Internet] 29:2519–2520. Available from: http://dx.doi.org/10.1093/bioinformatics/btt432

Chen Y, Huang W-C, Séjourné J, Clipperton-Allen AE, Page DT. 2015. Pten Mutations Alter Brain Growth Trajectory and Allocation of Cell Types through Elevated β-Catenin Signaling. J. Neurosci. [Internet] 35:10252 LP-10267. Available from: http://www.jneurosci.org/content/35/28/10252.abstract

Cheng Y-Z, Eley L, Hynes A-M, Overman LM, Simms RJ, Barker A, Dawe HR, Lindsay S, Sayer JA. 2012. Investigating Embryonic Expression Patterns and Evolution of AHI1 and CEP290 Genes, Implicated in Joubert Syndrome. PLoS One [Internet] 7:e44975. Available from: https://doi.org/10.1371/journal.pone.0044975

Chintalapati M, Dannemann M, Prüfer K. 2017. Using the Neandertal genome to study the evolution of small insertions and deletions in modern humans. BMC Evol. Biol. [Internet] 17:179. Available from: https://doi.org/10.1186/s12862-017-1018-8

Consortium TCS and A. 2005. Initial sequence of the chimpanzee genome and comparison with the human genome. Nature [Internet] 437:69. Available from: http://dx.doi.org/10.1038/nature04072

Cuberos H, Vallée B, Vourc'h P, Tastet J, Andres CR, Bénédetti H. 2015. Roles of LIM kinases in central nervous system function and dysfunction. FEBS Lett. [Internet] 589:3795–3806. Available from: http://doi.wiley.com/10.1016/j.febslet.2015.10.032

Dannemann M, Prüfer K, Kelso J. 2017. Functional implications of Neandertal introgression in modern humans. Genome Biol. [Internet] 18:61. Available from: https://doi.org/10.1186/s13059-017-1181-7

Day FR, Thompson DJ, Helgason H, Chasman DI, Finucane H, Sulem P, Ruth KS, Whalen S, Sarkar AK, Albrecht E, et al. 2017. Genomic analyses identify hundreds of variants associated with age at menarche and support a role for puberty timing in cancer risk. Nat. Genet. [Internet] 49:834–841. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28436984

Dediu D, Levinson SC. 2013. On the antiquity of language: The reinterpretation of neandertal linguistic capacities and its consequences. Front. Psychol. 4:1–17.

Dediu D, Levinson SC. 2018. Neanderthal language revisited: not only us. Curr. Opin. Behav. Sci. [Internet] 21:49–55. Available from: https://www.sciencedirect.com/science/article/pii/S2352154617301754

Dennis MY, Nuttle X, Sudmant PH, Antonacci F, Graves TA, Nefedov M, Rosenfeld JA, Sajjadian S, Malig M, Kotkiewicz H, et al. 2012. Evolution of Human-Specific Neural SRGAP2 Genes by Incomplete Segmental Duplication. Cell [Internet] 149:912–922. Available from: https://doi.org/10.1016/j.cell.2012.03.033

Dubois J, Poupon C, Thirion B, Simonnet H, Kulikova S, Leroy F, Hertz-Pannier L, Dehaene-Lambertz G. 2016. Exploring the Early Organization and Maturation of Linguistic Pathways in the Human Infant Brain. Cereb. Cortex [Internet] 26:2283–2298. Available from: http://dx.doi.org/10.1093/cercor/bhv082

Dumas LJ, O'Bleness MS, Davis JM, Dickens CM, Anderson N, Keeney JG, Jackson J, Sikela M, Raznahan A, Giedd J, et al. 2012. DUF1220-Domain Copy Number Implicated in Human Brain-Size Pathology and Evolution. Am. J. Hum. Genet. [Internet] 91:444–454. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3511999/

Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, Huber W. 2005. BioMart and Bioconductor: a powerful link between biological databases and microarray data analysis. Bioinformatics [Internet] 21:3439–3440. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16082012

Ellerman DA, Pei J, Gupta S, Snell WJ, Myles D, Primakoff P. 2009. Izumo is part of a multiprotein family whose members form large complexes on mammalian sperm. Mol. Reprod. Dev. [Internet] 76:1188–1199. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3779078/

Enard W, Gehre S, Hammerschmidt K, Hölter SM, Blass T, Somel M, Brückner MK, Schreiweis C, Winter C, Sohr R, et al. 2009. A Humanized Version of Foxp2 Affects Cortico-Basal Ganglia Circuits in Mice. Cell [Internet] 137:961–971. Available from: http://linkinghub.elsevier.com/retrieve/pii/S009286740900378X

Fietz SA, Huttner WB. 2011. Cortical progenitor expansion, self-renewal and neurogenesis—a polarized perspective. Curr. Opin. Neurobiol. [Internet] 21:23–35. Available from: http://www.sciencedirect.com/science/article/pii/S095943881000187X

Florio M, Albert M, Taverna E, Namba T, Brandl H, Lewitus E, Haffner C, Sykes A, Wong FK, Peters J, et al. 2015. Human-specific gene ARHGAP11B promotes basal progenitor amplification and neocortex expansion. Science [Internet] 347:1465–1470. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25721503

Fon EA, Sarrazin J, Meunier C, Alarcia J, Shevell MI, Philippe A, Leboyer M, Rouleau GA. 1995. Adenylosuccinate lyase (ADSL) and infantile autism: Absence of previously reported point mutation. Am. J. Med. Genet. [Internet] 60:554–557. Available from: http://doi.wiley.com/10.1002/ajmg.1320600614

Fu Q, Hajdinjak M, Moldovan OT, Constantin S, Mallick S, Skoglund P, Patterson N, Rohland N, Lazaridis I, Nickel B, et al. 2015. An early modern human from Romania with a recent Neanderthal ancestor. Nature [Internet] 524:216–219. Available from: http://www.nature.com/nature/journal/v524/n7564/abs/nature14558.html

Genin A, Desir J, Lambert N, Biervliet M, Van Der Aa N, Pierquin G, Killian A, Tosi M, Urbina M, Lefort A, et al. 2012. Kinetochore KMN network gene CASC5 mutated in primary microcephaly. Hum. Mol. Genet. [Internet] 21:5306–5317. Available from: http://dx.doi.org/10.1093/hmg/dds386

Giandomenico SL, Lancaster MA. 2017. Probing human brain evolution and development in organoids. Curr. Opin. Cell Biol. [Internet] 44:36–43. Available from: http://www.sciencedirect.com/science/article/pii/S0955067417300030

Gokhman D, Agranat-Tamir L, Housman G, Garcia-Perez R, Nissim-Rafinia M, Mallick S, Nieves-Colón M, Li H, Alpaslan-Roodenberg S, Novak M, et al. 2017. Extensive Regulatory Changes in Genes Affecting Vocal and Facial Anatomy Separate Modern from Archaic Humans. bioRxiv [Internet]. Available from: http://biorxiv.org/content/early/2017/10/03/106955.abstract

Grantham R. 1974. Amino Acid Difference Formula to Help Explain Protein Evolution. Science (80-.). 185.

Gratten J, Visscher PM. 2016. Genetic pleiotropy in complex traits and diseases: implications for genomic medicine. Genome Med. [Internet] 8:78. Available from: https://doi.org/10.1186/s13073-016-0332-x

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W, Fritz MH-Y, et al. 2010. A Draft Sequence of the Neandertal Genome. Science (80-.). [Internet] 328:710–722. Available from: http://www.sciencemag.org/content/328/5979/710.abstract

Grosse Wiesmann C, Schreiber J, Singer T, Steinbeis N, Friederici AD. 2017. White matter maturation is associated with the emergence of Theory of Mind in early childhood. Nat. Commun. [Internet] 8:14692. Available from: http://dx.doi.org/10.1038/ncomms14692

Grote S, Prüfer K, Kelso J, Dannemann M. 2016. ABAEnrichment: an R package to test for gene set expression enrichment in the adult and developing human brain. Bioinformatics [Internet] 32:3201–3203. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5048072/

Gullard A, Gluhak-Heinrich J, Papagerakis S, Sohn P, Unterbrink A, Chen S, MacDougall M. 2016. MEPE Localization in the Craniofacial Complex and Function in Tooth Dentin Formation. J. Histochem. Cytochem. [Internet] 64:224–236. Available from: https://doi.org/10.1369/0022155416635569

Gunz P, Neubauer S, Golovanova L, Doronichev V, Maureille B, Hublin J-J. 2012. A uniquely modern human pattern of endocranial development. Insights from a new cranial reconstruction of the Neandertal newborn from Mezmaiskaya. J. Hum. Evol. [Internet] 62:300–313. Available from: http://www.sciencedirect.com/science/article/pii/S0047248411002314

Gunz P, Neubauer S, Maureille B, Hublin JJ. 2010. Brain development after birth differs between Neandertals and modern humans. Curr. Biol. 20:921–922.

Gupta GD, Coyaud É, Gonçalves J, Mojarad BA, Liu Y, Wu Q, Gheiratmand L, Comartin D, Tkach JM, Cheung SWT, et al. 2015. A Dynamic Protein Interaction Landscape of the Human Centrosome-Cilium Interface. Cell [Internet] 163:1484–1499. Available from: http://dx.doi.org/10.1016/j.cell.2015.10.065

Habib AM, Matsuyama A, Okorokov AL, Santana-Varela S, Bras JT, Aloisi AM, Emery EC, Bogdanov YD, Follenfant M, Gossage SJ, et al. 2017. A novel human pain insensitivity disorder caused by a point mutation in ZFHX2. Brain [Internet]:awx326-awx326. Available from: http://dx.doi.org/10.1093/brain/awx326

Hoffmann DL, Angelucci DE, Villaverde V, Zapata J, Zilhão J. 2018. Symbolic use of marine shells and mineral pigments by Iberian Neandertals 115,000 years ago. Sci. Adv. [Internet] 4. Available from: http://advances.sciencemag.org/content/4/2/eaar5255.abstract

Hofman M a. 1983. Encephalization in hominids: evidence for the model of punctuationalism. Brain. Behav. Evol.

Hoshino J, Aruga J, Ishiguro A, Mikoshiba K. 2003. Dorz1, a novel gene expressed in differentiating cerebellar granule neurons, is down-regulated in Zic1-deficient mouse. Mol. Brain Res. [Internet] 120:57–64. Available from: http://www.sciencedirect.com/science/article/pii/S0169328X03004467

Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, Bravo HC, Davis S, Gatto L, Girke T, et al. 2015. Orchestrating high-throughput genomic analysis with Bioconductor. Nat Meth [Internet] 12:115–121. Available from: http://dx.doi.org/10.1038/nmeth.3252%5Cn10.1038/nmeth.3252

Hublin J-J, Ben-Ncer A, Bailey SE, Freidline SE, Neubauer S, Skinner MM, Bergmann I, Le Cabec A, Benazzi S, Harvati K, et al. 2017. New fossils from Jebel Irhoud, Morocco and the pan-African origin of Homo sapiens. Nature [Internet] 546:289. Available from: http://dx.doi.org/10.1038/nature22336

Hublin J-J, Neubauer S, Gunz P. 2015. Brain ontogeny and life history in Pleistocene hominins. Philos. Trans. R. Soc. B Biol. Sci. [Internet] 370. Available from: http://rstb.royalsocietypublishing.org/content/370/1663/20140062.abstract

Imamura K, Maeda S, Kawamura I, Matsuyama K, Shinohara N, Yahiro Y, Nagano S, Setoguchi T, Yokouchi M, Ishidou Y, et al. 2014. Human Immunodeficiency Virus Type 1 Enhancer-binding Protein 3 Is Essential for the Expression of Asparagine-linked Glycosylation 2 in the Regulation of Osteoblast and Chondrocyte Differentiation. J. Biol. Chem. [Internet] 289:9865–9879. Available from: http://www.jbc.org/content/289/14/9865.abstract

Jamsheer A, Sowińska A, Trzeciak T, Jamsheer-Bratkowska M, Geppert A, Latos-Bieleńska A. 2012. Expanded mutational spectrum of the GLI3 gene substantiates genotype–phenotype correlations. J. Appl. Genet. [Internet] 53:415–422. Available from: https://doi.org/10.1007/s13353-012-0109-x

Jelen N, Ule J, Živin M. 2010. Cholinergic regulation of striatal Nova mRNAs. Neuroscience [Internet] 169:619–627. Available from: http://www.sciencedirect.com/science/article/pii/S0306452210007128

Jensen KB, Dredge BK, Stefani G, Zhong R, Buckanovich RJ, Okano HJ, Yang YYL, Darnell RB. 2000. Nova-1
Regulates Neuron-Specific Alternative Splicing and Is Essential for Neuronal Viability. Neuron [Internet] 25:359–
371. Available from: http://dx.doi.org/10.1016/S0896-6273(00)80900-9
Jeong J-W, Sundaram S, Behen ME, Chugani HT. 2016. Relationship between genotype and arcuate fasciculus
morphology in six young children with global developmental delay: Preliminary DTI stuy. J. Magn. Reson.
Imaging [Internet] 44:1504–1512. Available from: http://doi.wiley.com/10.1002/jmri.25306
Jones DC, Schweitzer MN, Wein M, Sigrist K, Takagi T, Ishii S, Glimcher LH. 2010. Uncoupling of growth plate
maturation and bone formation in mice lacking both Schnurri-2 and Schnurri-3. Proc. Natl. Acad. Sci. [Internet]
107:8254–8258. Available from: http://www.pnas.org/content/107/18/8254.abstract
Ju X-C, Hou Q-Q, Sheng A-L, Wu K-Y, Zhou Y, Jin Y, Wen T, Yang Z, Wang X, Luo Z-G. 2016. The hominoid-specific
gene TBC1D3 promotes generation of basal neural progenitors and induces cortical folding in mice.Gleeson JG,
editor. Elife [Internet] 5:e18197. Available from: https://doi.org/10.7554/eLife.18197
Jurecka A, Jurkiewicz E, Tylki-Szymanska A. 2012. Magnetic resonance imaging of the brain in adenylosuccinate lyase
deficiency: a report of seven cases and a review of the literature. Eur. J. Pediatr. [Internet] 171:131–138. Available
from: https://doi.org/10.1007/s00431-011-1503-9
Karabay A, Yu W, Solowska JM, Baird DH, Baas PW. 2004. Axonal Growth Is Sensitive to the Levels of Katanin, a
Protein That Severs Microtubules. J. Neurosci. [Internet] 24:5778 LP-5788. Available from:
http://www.jneurosci.org/content/24/25/5778.abstract
Kevenaar JT, Bianchi S, van Spronsen M, Olieric N, Lipka J, Frias CP, Mikhaylova M, Harterink M, Keijzer N, Wulf
PS, et al. 2017. Kinesin-Binding Protein Controls Microtubule Dynamics and Cargo Trafficking by Regulating
Kinesin Motor Activity. Curr. Biol. [Internet] 26:849–861. Available from:
http://dx.doi.org/10.1016/j.cub.2016.01.048
Kim JH, Kim N. 2014. Regulation of NFATc1 in Osteoclast Differentiation. J Bone Metab [Internet] 21:233–241.
Available from: http://synapse.koreamed.org/DOIx.php?id=10.11005%2Fjbm.2014.21.4.233
Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. 2014. A general framework for estimating the
relative pathogenicity of human genetic variants. Nat Genet [Internet] 46:310–315. Available from:
https://doi.org/10.1038/ng.2892
Kochiyama T, Ogihara N, Tanabe HC, Kondo O, Amano H, Hasegawa K, Suzuki H, Ponce de León MS, Zollikofer
CPE, Bastir M, et al. 2018. Reconstructing the Neanderthal brain using computational anatomy. Sci. Rep.
[Internet] 8:6296. Available from: https://doi.org/10.1038/s41598-018-24331-0
Konopka G, Friedrich T, Davis-Turak J, Winden K, Oldham MC, Gao F, Chen L, Wang G-Z, Luo R, Preuss TM, et al.
2012. Human-Specific Transcriptional Networks in the Brain. Neuron [Internet] 75:601–617. Available from:
http://dx.doi.org/10.1016/j.neuron.2012.05.034
Koser DE, Thompson AJ, Foster SK, Dwivedy A, Pillai EK, Sheridan GK, Svoboda H, Viana M, Costa L da F, Guck J,
et al. 2016. Mechanosensing is critical for axon growth in the developing brain. Nat. Neurosci. [Internet] 19:1592.
Available from: http://dx.doi.org/10.1038/nn.4394
Kuhlwilm M. 2018. The evolution of FOXP2 in the light of admixture. Curr. Opin. Behav. Sci. 21.
Kuhlwilm M, Gronau I, Hubisz MJMJ, Filippo C de, Prado-Martinez J, Kircher M, Fu Q, Burbano HAHA, Lalueza-
Fox C, Rasilla M de la, et al. 2016. Ancient gene flow from early modern humans into Eastern Neanderthals.
Nature [Internet] 530:429–433. Available from: http://www.nature.com/doifinder/10.1038/nature16544
Kumar P, Henikoff S, Ng PC. 2009. Predicting the effects of coding non-synonymous variants on protein function using
the SIFT algorithm. Nat. Protoc. 4:1073–1081.
Kuwahara A, Sakai H, Xu Y, Itoh Y, Hirabayashi Y, Gotoh Y. 2014. Tcf3 Represses Wnt–β-Catenin Signaling and
Maintains Neural Stem Cell Population during Neocortical Development. PLoS One [Internet] 9:e94408.
Available from: https://doi.org/10.1371/journal.pone.0094408
Kwon EK, Louie K, Kulkarni A, Yatabe M, Ruellas AC de O, Snider TN, Mochida Y, Cevidanes LHS, Mishina Y,
Zhang H. 2018. The Role of Ellis-Van Creveld 2(*EVC2*) in Mice During Cranial Bone Development. Anat. Rec.
[Internet] 301:46–55. Available from: http://doi.wiley.com/10.1002/ar.23692
Lacruz RS, Bromage TG, O'Higgins P, Arsuaga J-L, Stringer C, Godinho RM, Warshaw J, Martínez I, Gracia-Tellez A,
de Castro JMB, et al. 2015. Ontogeny of the maxilla in Neanderthals and their ancestors. Nat. Commun. [Internet]
6:8996. Available from: http://dx.doi.org/10.1038/ncomms9996
Lacruz RS, de Castro JMB, Martinón-Torres M, O'Higgins P, Paine ML, Carbonell E, Arsuaga JL, Bromage TG. 2013.
Facial Morphogenesis of the Earliest Europeans. PLoS One [Internet] 8:e65199. Available from:
https://doi.org/10.1371/journal.pone.0065199
Langergraber KE, Prüfer K, Rowney C, Boesch C, Crockford C, Fawcett K, Inoue E, Inoue-Muruyama M, Mitani JC,
Muller MN, et al. 2012. Generation times in wild chimpanzees and gorillas suggest earlier divergence times in
great ape and human evolution. Proc. … [Internet] 109:15716–15721. Available from:
http://www.pnas.org/content/early/2012/08/08/1211740109.short
Lawrence M, Gentleman R, Carey V. 2009. rtracklayer: an R package for interfacing with genome browsers.
Bioinformatics [Internet] 25:1841–1842. Available from: https://academic.oup.com/bioinformatics/article-lookup/
doi/10.1093/bioinformatics/btp328
Lee MK, Shaffer JR, Leslie EJ, Orlova E, Carlson JC, Feingold E, Marazita ML, Weinberg SM. 2017. Genome-wide
association study of facial morphology reveals novel associations with FREM1 and PARK2. PLoS One [Internet]
12:e0176566. Available from: https://doi.org/10.1371/journal.pone.0176566

Lei H, Yan Z, Sun X, Zhang Y, Wang J, Ma C, Xu Q, Wang R, Jarvis ED, Sun Z. 2017. Axon guidance pathways served as common targets for human speech/language evolution and related disorders. Brain Lang. [Internet] 174:1–8. Available from: http://www.sciencedirect.com/science/article/pii/S0093934X16301869

Lek M, Karczewski KJ, Minikel E V, Samocha KE, Banks E, Fennell T, O'Donnell-Luria AH, Ware JS, Hill AJ, Cummings BB, et al. 2016. Analysis of protein-coding genetic variation in 60,706 humans. Nature [Internet] 536:285–291. Available from: http://dx.doi.org/10.1038/nature19057

Li F-F, Wang X-D, Zhu M-W, Lou Z-H, Zhang Q, Zhu C-Y, Feng H-L, Lin Z-G, Liu S-L. 2015. Identification of two novel critical mutations in PCNT gene resulting in microcephalic osteodysplastic primordial dwarfism type II associated with multiple intracranial aneurysms. Metab. Brain Dis. [Internet] 30:1387–1394. Available from: https://doi.org/10.1007/s11011-015-9712-y

Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics [Internet] 27:2987–2993. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3198575/

Li Y, Muffat J, Omer A, Bosch I, Lancaster MA, Sur M, Gehrke L, Knoblich JA, Jaenisch R. 2017. Induction of Expansion and Folding in Human Cerebral Organoids. Cell Stem Cell [Internet] 20:385–396.e3. Available from: http://dx.doi.org/10.1016/j.stem.2016.11.017

Liu X, Somel M, Tang L, Yan Z, Jiang X, Guo S, Yuan Y, He L, Oleksiak A, Zhang Y, et al. 2012. Extension of cortical synaptic development distinguishes humans from chimpanzees and macaques. Genome Res. [Internet] 22:611–622. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3317144/

Lüders J. 2016. The Microtubule Cytoskeleton. (Lüders J, editor.). Vienna: Springer Vienna Available from: http://link.springer.com/10.1007/978-3-7091-1903-7

MacArthur J, Bowler E, Cerezo M, Gil L, Hall P, Hastings E, Junkins H, McMahon A, Milano A, Morales J, et al. 2017. The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). Nucleic Acids Res. [Internet] 45:D896–D901. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5210590/

Mars RB, Eichert N, Jbabdi S, Verhagen L, Rushworth MFS. 2018. Connectivity and the search for specializations in the language-capable brain. Curr. Opin. Behav. Sci. [Internet] 21:19–26. Available from: https://www.sciencedirect.com/science/article/pii/S235215461730181X

McCoy RC, Wakefield J, Akey JM. 2017. Impacts of Neanderthal-Introgressed Sequences on the Landscape of Human Gene Expression. Cell [Internet] 168:916–927.e12. Available from: http://dx.doi.org/10.1016/j.cell.2017.01.038

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, Flicek P, Cunningham F. 2016. The Ensembl Variant Effect Predictor. Genome Biol. [Internet] 17:122. Available from: https://doi.org/10.1186/s13059-016-0974-4

McLean CY, Reno PL, Pollen AA, Bassan AI, Capellini TD, Guenther C, Indjeian VB, Lim X, Menke DB, Schaar BT, et al. 2011. Human-specific loss of regulatory DNA and the evolution of human-specific traits. Nature [Internet] 471:216–219. Available from: http://www.nature.com/doifinder/10.1038/nature09774

McVicker DP, Awe AM, Richters KE, Wilson RL, Cowdrey DA, Hu X, Chapman ER, Dent EW. 2016. Transport of a kinesin-cargo pair along microtubules into dendritic spines undergoing synaptic plasticity. Nat. Commun. [Internet] 7:12741. Available from: http://dx.doi.org/10.1038/ncomms12741

Megraw TL, Sharkey JT, Nowakowski RS. 2011. Cdk5rap2 exposes the centrosomal root of microcephaly syndromes. Trends Cell Biol. [Internet] 21:470–480. Available from: http://dx.doi.org/10.1016/j.tcb.2011.04.007

Menzies FM, Fleming A, Rubinsztein DC. 2015. Compromised autophagy and neurodegenerative diseases. Nat. Rev. Neurosci. [Internet] 16:345. Available from: http://dx.doi.org/10.1038/nrn3961

Meyer M, Kircher M, Gansauge M-T, Li H, Racimo F, Mallick S, Schraiber JG, Jay F, Prüfer K, de Filippo C, et al. 2012. a High Coverage Genome Sequence From an Archaic Denisovan Individual. Science 338:222–226.

Miller JA, Ding S-L, Sunkin SM, Smith KA, Ng L, Szafer A, Ebbert A, Riley ZL, Aiona K, Arnold JM, et al. 2014. Transcriptional Landscape of the Prenatal Human Brain. Nature [Internet] 508:199–206. Available from: http://dx.doi.org/10.1038/nature13185

Miller KJ, Botvinick MM, Brody CD. 2018. Value Representations in Orbitofrontal Cortex Drive Learning, but not Choice. bioRxiv [Internet]. Available from: http://biorxiv.org/content/early/2018/01/10/245720.1.abstract

Mirzaa GM, Parry DA, Fry AE, Giamanco KA, Schwartzentruber J, Vanstone M, Logan C V, Roberts N, Johnson CA, Singh S, et al. 2014. De novo CCND2 mutations leading to stabilization of cyclin D2 cause megalencephaly-polymicrogyria-polydactyly-hydrocephalus syndrome. Nat. Genet. [Internet] 46:510. Available from: http://dx.doi.org/10.1038/ng.2948

Moncini S, Bedeschi MF, Castronovo P, Crippa M, Calvello M, Garghentino RR, Scuvera G, Finelli P, Venturin M. 2013. ATRX mutation in two adult brothers with non-specific moderate intellectual disability identified by exome sequencing. Meta Gene [Internet] 1:102–108. Available from: http://www.sciencedirect.com/science/article/pii/S2214540013000066

Montgomery SH, Capellini I, Venditti C, Barton RA, Mundy NI. 2011. Adaptive Evolution of Four Microcephaly Genes and the Evolution of Brain Size in Anthropoid Primates. Mol. Biol. Evol. [Internet] 28:625–638. Available from: http://dx.doi.org/10.1093/molbev/msq237

Neubauer S, Hublin J-J, Gunz P. 2018. The evolution of modern human brain shape. Sci. Adv. [Internet] 4. Available from: http://advances.sciencemag.org/content/4/1/eaao5961.abstract

O'Bleness M, Searles VB, Varki A, Gagneux P, Sikela JM. 2012. Evolution of genetic and genomic features unique to the human lineage. Nat. Rev. Genet. [Internet] 13:853–866. Available from: http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=23154808&retmode=ref&cmd=prlinks%5Cnpapers3://publication/doi/10.1038/nrg3336

O'Regan L, Fry AM. 2009. The Nek6 and Nek7 Protein Kinases Are Required for Robust Mitotic Spindle Formation and Cytokinesis. Mol. Cell. Biol. [Internet] 29:3975–3990. Available from: http://mcb.asm.org/content/29/14/3975.abstract

Oh J-E, Kim HJ, Kim W-S, Lee ZH, Ryoo H-M, Hwang SJ, Lee Y, Kim H-H. 2012. PlexinA2 mediates osteoblast differentiation via regulation of Runx2. J. Bone Miner. Res. [Internet] 27:552–562. Available from: http://doi.wiley.com/10.1002/jbmr.1471

Pääbo S. 2014. The Human Condition—A Molecular Approach. Cell [Internet] 157:216–226. Available from: http://www.sciencedirect.com/science/article/pii/S009286741301605X

Park J-A, Kim KC. 2010. Expression patterns of PRDM10 during mouse embryonic development. BMB Rep. 43:29–33.

Paten B, Herrero J, Fitzgerald S, Beal K, Flicek P, Holmes I, Birney E. 2008. Genome-wide nucleotide-level mammalian ancestor reconstruction. Genome Res. 18:1829–1843.

Pearce E, Wlodarski R, Machin A, Dunbar RIM. 2017. Variation in the β-endorphin, oxytocin, and dopamine receptor genes is associated with different dimensions of human sociality. Proc. Natl. Acad. Sci. [Internet] 114:5300 LP-5305. Available from: http://www.pnas.org/content/114/20/5300.abstract

Peyrégne S, Boyle MJ, Dannemann M, Prüfer K. 2017. Detecting ancient positive selection in humans using extended lineage sorting. Genome Res. [Internet] 27:1563–1572. Available from: http://genome.cshlp.org/content/27/9/1563.abstract

Pfenning AR, Hara E, Whitney O, Rivas M V, Wang R, Roulhac PL, Howard JT, Wirthlin M, Lovell P V, Ganapathy G, et al. 2014. Convergent transcriptional specializations in the brains of humans and song-learning birds. Science (80-.). [Internet] 346. Available from: http://science.sciencemag.org/content/346/6215/1256846.abstract

Ponce de León MS, Bienvenu T, Akazawa T, Zollikofer CPE. 2017. Brain development is similar in Neanderthals and modern humans. Curr. Biol. [Internet] 26:R665–R666. Available from: http://dx.doi.org/10.1016/j.cub.2016.06.022

Popovitchenko T, Thompson K, Viljetic B, Jiao X, Kontonyiannis DL, Kiledjian M, Hart RP, Rasin MR. 2016. The RNA binding protein HuR determines the differential translation of autism-associated FoxP subfamily members in the developing neocortex. Sci. Rep. [Internet] 6:28998. Available from: http://dx.doi.org/10.1038/srep28998

Prüfer K, de Filippo C, Grote S, Mafessoni F, Korlević P, Hajdinjak M, Vernot B, Skov L, Hsieh P, Peyrégne S, et al. 2017. A high-coverage Neandertal genome from Vindija Cave in Croatia. Science (80-.). [Internet]. Available from: http://science.sciencemag.org/content/early/2017/10/04/science.aao1887.abstract

Prüfer K, Muetzel B, Do H-HH, Weiss G, Khaitovich P, Rahm E, Pääbo S, Lachmann M, Enard W, Paabo S, et al. 2007. FUNC: a package for detecting significant associations between gene sets and ontological annotations. BMC Bioinformatics [Internet] 8:41. Available from: https://doi.org/10.1186/1471-2105-8-41

Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, Heinze A, Renaud G, Sudmant PHPH, de Filippo C, et al. 2014. The complete genome sequence of a Neanderthal from the Altai Mountains. Nature [Internet] 505:43–49. Available from: http://europepmc.org/abstract/med/24352235

Puga A, Xia Y, Elferink C. 2002. Role of the aryl hydrocarbon receptor in cell cycle regulation. Chem. Biol. Interact. [Internet] 141:117–130. Available from: http://www.sciencedirect.com/science/article/pii/S0009279702000698

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics [Internet] 26:841–842. Available from: https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btq033

Rabett RJ. 2018. The success of failed Homo sapiens dispersals out of Africa and into Asia. Nat. Ecol. Evol. [Internet] 2:212–219. Available from: https://doi.org/10.1038/s41559-017-0436-8

Racimo F. 2016. Testing for Ancient Selection Using Cross-population Allele Frequency Differentiation. Genetics [Internet] 202:733 LP-750. Available from: http://www.genetics.org/content/202/2/733.abstract

Racimo F, Kuhlwilm M, Slatkin M. 2014. A test for ancient selective sweeps and an application to candidate sites in modern humans. Mol. Biol. Evol. [Internet] 31:3344–3358. Available from: http://europepmc.org/abstract/med/25172957

Rak Y. 1986. The Neanderthal: A new look at an old face. J. Hum. Evol. [Internet] 15:151–164. Available from: http://www.sciencedirect.com/science/article/pii/S0047248486800422

Ramirez Rozzi F V, de Castro JM. 2004. surprisingly rapid growth in Neanderthals. Nature [Internet] 428. Available from: https://doi.org/10.1038/nature02428

Ratti A, Fallini C, Colombrita C, Pascale A, Laforenza U, Quattrone A, Silani V. 2008. Post-transcriptional regulation of neuro-oncological ventral antigen 1 by the neuronal RNA-binding proteins ELAV. J. Biol. Chem. [Internet]. Available from: http://www.jbc.org/content/early/2008/01/24/jbc.M706082200.short

Ritchie GRS, Dunham I, Zeggini E, Flicek P. 2014. Functional annotation of noncoding sequence variants. Nat. Methods [Internet] 11:294–296. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24487584

Ritchie K, Watson LA, Davidson B, Jiang Y, Bérubé NG. 2014. ATRX is required for maintenance of the neuroprogenitor cell pool in the embryonic mouse brain. Biol. Open [Internet] 3:1158 LP-1163. Available from: http://bio.biologists.org/content/3/12/1158.abstract

Rosas A, Ríos L, Estalrrich A, Liversidge H, García-Tabernero A, Huguet R, Cardoso H, Bastir M, Lalueza-Fox C, de la Rasilla M, et al. 2017. The growth pattern of Neandertals, reconstructed from a juvenile skeleton from El Sidrón (Spain). Science (80-.). [Internet] 357:1282 LP-1287. Available from: http://science.sciencemag.org/content/357/6357/1282.abstract

Sankararaman S, Mallick S, Dannemann M, Prufer K, Kelso J, Paabo S, Patterson N, Reich D. 2014. The genomic landscape of Neanderthal ancestry in present-day humans. Nature [Internet] 507:354–357. Available from: http://dx.doi.org/10.1038/nature12961

Sankararaman S, Mallick S, Patterson N, Reich D. 2016. The Combined Landscape of Denisovan and Neanderthal Ancestry in Present-Day Humans. Curr. Biol. [Internet] 26:1241–1247. Available from: http://dx.doi.org/10.1016/j.cub.2016.03.037

Schlebusch CM, Malmström H, Günther T, Sjödin P, Coutinho A, Edlund H, Munters AR, Vicente M, Steyn M, Soodyall H, et al. 2017. Southern African ancient genomes estimate modern human divergence to 350,000 to 260,000 years ago. Science (80-.). [Internet]. Available from: http://science.sciencemag.org/content/early/2017/09/27/science.aao6266.abstract

Schlebusch CM, Skoglund P, Sjödin P, Gattepaille LM, Hernandez D, Jay F, Li S, De Jongh M, Singleton A, Blum MGB, et al. 2012. Genomic Variation in Seven Khoe-San Groups Reveals Adaptation and Complex African History. Science (80-.). [Internet] 338:374 LP-379. Available from: http://science.sciencemag.org/content/338/6105/374.abstract

Schoenemann PT. 2004. Brain Size Scaling and Body Composition in Mammals. Brain. Behav. Evol. [Internet] 63:47–60. Available from: https://www.karger.com/DOI/10.1159/000073759

Sherry ST, Ward M-H, Kholodov M, Baker J, Phan L, Smigielski EM, Sirotkin K. 2001. dbSNP: the NCBI database of genetic variation. Nucleic Acids Res. [Internet] 29:308–311. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC29783/

Sherwood CC, Gómez-Robles A. 2017. Brain Plasticity and Human Evolution. Annu. Rev. Anthropol. [Internet] 46:399–419. Available from: https://doi.org/10.1146/annurev-anthro-102215-100009

Shi J, Hui L, Xu Y, Wang F, Huang W, Hu G. 2002. Sequence variations in the mu-opioid receptor gene (OPRM1) associated with human addiction to heroin. Hum. Mutat. [Internet] 19:459–460. Available from: http://doi.wiley.com/10.1002/humu.9026

Shi L, Hu E, Wang Z, Liu J, Li J, Li M, Chen H, Yu C, Jiang T, Su B. 2017. Regional selection of the brain size regulating gene CASC5 provides new insight into human brain evolution. Hum. Genet. [Internet] 136:193–204. Available from: https://doi.org/10.1007/s00439-016-1748-5

Siegel DA, Huang MK, Becker SF. 2002. Ectopic dendrite initiation: CNS pathogenesis as a model of CNS development. Int. J. Dev. Neurosci. [Internet] 20:373–389. Available from: http://www.sciencedirect.com/science/article/pii/S0736574802000552

Signorell A. 2017. DescTools: Tools for Descriptive Statistics. Available from: https://cran.r-project.org/package=DescTools

Simonti CN, Vernot B, Bastarache L, Bottinger E, Carrell DS, Chisholm RL, Crosslin DR, Hebbring SJ, Jarvik GP, Kullo IJ, et al. 2016. The phenotypic legacy of admixture between modern humans and Neandertals. Science (80-.). [Internet] 351:737 LP-741. Available from: http://science.sciencemag.org/content/351/6274/737.abstract

Skeide MA, Friederici AD. 2016. The ontogeny of the cortical language network. Nat. Rev. Neurosci. [Internet] 17:323. Available from: http://dx.doi.org/10.1038/nrn.2016.23

Skoglund P, Thompson JC, Prendergast ME, Mittnik A, Sirak K, Hajdinjak M, Salie T, Rohland N, Mallick S, Peltzer A, et al. 2017. Reconstructing Prehistoric African Population Structure. Cell [Internet] 171:59–71.e21. Available from: http://dx.doi.org/10.1016/j.cell.2017.08.049

Smith BH, Tompkins RL. 1995. Toward A Life History of the Hominidae. Annu. Rev. Anthropol. [Internet] 24:257–279. Available from: https://doi.org/10.1146/annurev.an.24.100195.001353

Smith TM, Tafforeau P, Reid DJ, Pouech J, Lazzari V, Zermeno JP, Guatelli-Steinberg D, Olejniczak AJ, Hoffman A, Radovcic J, et al. 2010. Dental evidence for ontogenetic differences between modern humans and Neanderthals. Proc Natl Acad Sci U S A [Internet] 107:20923–20928. Available from: http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=21078988&retmode=ref&cmd=prlinks%5Cnpapers3://publication/doi/10.1073/pnas.1010906107

Somel M, Franz H, Yan Z, Lorenc A, Guo S, Giger T, Kelso J, Nickel B, Dannemann M, Bahn S, et al. 2009. Transcriptional neoteny in the human brain. Proc. Natl. Acad. Sci. U. S. A. [Internet] 106:5743–5748. Available from: http://www.pnas.org/content/early/2009/03/20/0900544106

Somel M, Liu X, Khaitovich P. 2013. Human brain evolution: transcripts, metabolites and their regulators. Nat. Rev. Neurosci. [Internet] 14:1–16. Available from: http://dx.doi.org/10.1038/nrn3372%5Cnpapers3://publication/doi/10.1038/nrn3372

Sreedharan S, Almén MS, Carlini VP, Haitina T, Stephansson O, Sommer WH, Heilig M, de Barioglio SR, Fredriksson R, Schiöth HB. 2011. The G protein coupled receptor Gpr153 shares common evolutionary origin with Gpr162 and is highly expressed in central regions including the thalamus, cerebellum and the arcuate nucleus. FEBS J. [Internet] 278:4881–4894. Available from: http://dx.doi.org/10.1111/j.1742-4658.2011.08388.x

Stringer C. 2016. The origin and evolution of Homo sapiens Philos. Trans. R. Soc. B Biol. Sci. [Internet] 371. Available from: http://rstb.royalsocietypublishing.org/content/371/1698/20150237.abstract

Suzuki IK, Gacquer D, Van Heurck R, Kumar D, Wojno M, Bilheu A, Herpoel A, Lambert N, Cheron J, Polleux F, et al. 2018. Human-Specific NOTCH2NL Genes Expand Cortical Neurogenesis through Delta/Notch Regulation. Cell [Internet] 173:1370–1384.e16. Available from: https://doi.org/10.1016/j.cell.2018.03.067

Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, Santos A, Doncheva NT, Roth A, Bork P, et al. 2017. The STRING database in 2017: quality-controlled protein–protein association networks, made broadly accessible. Nucleic Acids Res. [Internet] 45:D362–D368. Available from: http://dx.doi.org/10.1093/nar/gkw937

Takata A, Miyake N, Tsurusaki Y, Fukai R, Miyatake S, Koshimizu E, Kushima I, Okada T, Morikawa M, Uno Y, et al. 2018. Integrative Analyses of De Novo Mutations Provide Deeper Biological Insights into Autism Spectrum Disorder. Cell Rep. [Internet] 22:734–747. Available from: http://dx.doi.org/10.1016/j.celrep.2017.12.074

Takemoto M, Hattori Y, Zhao H, Sato H, Tamada A, Sasaki S, Nakajima K, Yamamoto N. 2011. Laminar and Areal Expression of Unc5d and Its Role in Cortical Cell Survival. Cereb. Cortex [Internet] 21:1925–1934. Available from: http://dx.doi.org/10.1093/cercor/bhq265

Talbot JC, Nichols JT, Yan Y-L, Leonard IF, BreMiller RA, Amacher SL, Postlethwait JH, Kimmel CB. 2016. Pharyngeal morphogenesis requires fras1-itga8-dependent epithelial-mesenchymal interaction. Dev. Biol. [Internet] 416:136–148. Available from: http://www.sciencedirect.com/science/article/pii/S0012160616301051

Tan E, Lim ECP, Teo Y, Lim Y, Law H, Sia AT. 2009. Ethnicity and OPRM variant independently predict pain perception and patient-controlled analgesia usage for post-operative pain. Mol. Pain [Internet] 5:32. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2709614/

Tang S, Wang X, Li W, Yang X, Li Z, Liu W, Li C, Zhu Z, Wang L, Wang J, et al. 2017. Biallelic Mutations in CFAP43 and CFAP44 Cause Male Infertility with Multiple Morphological Abnormalities of the Sperm Flagella. Am. J. Hum. Genet. [Internet] 100:854–864. Available from: http://dx.doi.org/10.1016/j.ajhg.2017.04.012

Tashima T, Nagatoishi S, Sagara H, Ohnuma S, Tsumoto K. 2015. Osteomodulin regulates diameter and alters shape of collagen fibrils. Biochem. Biophys. Res. Commun. [Internet] 463:292–296. Available from: http://www.sciencedirect.com/science/article/pii/S0006291X15009699

Teves ME, Sundaresan G, Cohen DJ, Hyzy SL, Kajan I, Maczis M, Zhang Z, Costanzo RM, Zweit J, Schwartz Z, et al. 2015. Spag17 Deficiency Results in Skeletal Malformations and Bone Abnormalities. PLoS One [Internet] 10:e0125936. Available from: https://doi.org/10.1371/journal.pone.0125936

Torres JZ, Summers MK, Peterson D, Brauer MJ, Lee J, Senese S, Gholkar AA, Lo Y-C, Lei X, Jung K, et al. 2017. The STARD9/Kif16a Kinesin Associates with Mitotic Microtubules and Regulates Spindle Pole Assembly. Cell [Internet] 147:1309–1323. Available from: http://dx.doi.org/10.1016/j.cell.2011.11.020

Trinkaus E, Howells WW. 1979. The Neanderthals. Sci. Am. [Internet] 241:118–133. Available from: http://www.jstor.org/stable/24965359

Varki A, Altheide TK. 2005. Comparing the human and chimpanzee genomes: Searching for needles in a haystack. Genome Res. [Internet] 15:1746–1758. Available from: http://genome.cshlp.org/content/15/12/1746.long

Verheijen MHG, Camargo N, Verdier V, Nadra K, de Preux Charles A-S, Médard J-J, Luoma A, Crowther M, Inouye H, Shimano H, et al. 2009. SCAP is required for timely and proper myelin membrane synthesis. Proc. Natl. Acad. Sci. [Internet] 106:21383–21388. Available from: http://www.pnas.org/content/106/50/21383.abstract

Vernot B, Akey JM. 2014. Resurrecting Surviving Neandertal Lineages from Modern Human Genomes. Science (80-.). [Internet] 343:1017–1021. Available from: http://science.sciencemag.org/content/343/6174/1017.abstract

Vernot B, Tucci S, Kelso J, Schraiber JG, Wolf AB, Gittelman RM, Dannemann M, Grote S, McCoy RC, Norton H, et al. 2016. Excavating Neandertal and Denisovan DNA from the genomes of Melanesian individuals. Science (80-.). [Internet] 352:235–239. Available from: http://science.sciencemag.org/content/352/6282/235.abstract

Verzi MP, Agarwal P, Brown C, McCulley DJ, Schwarz JJ, Black BL. 2007. The Transcription Factor MEF2C Is Required for Craniofacial Development. Dev. Cell [Internet] 12:645–652. Available from: http://dx.doi.org/10.1016/j.devcel.2007.03.007

Wang C, Liang C-C, Bian ZC, Zhu Y, Guan J-L. 2013. FIP200 is required for maintenance and differentiation of postnatal neural stem cells. Nat. Neurosci. [Internet] 16:532. Available from: http://dx.doi.org/10.1038/nn.3365

Wang R, Chen C-C, Hara E, Rivas M V, Roulhac PL, Howard JT, Chakraborty M, Audet J-N, Jarvis ED. 2015. Convergent Differential Regulation of SLIT-ROBO Axon Guidance Genes in the Brains of Vocal Learners. J. Comp. Neurol. [Internet] 523:892–906. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4329046/

Wang Z, Widgren EE, Sivashanmugam P, O'Rand MG, Richardson RT. 2005. Association of Eppin with Semenogelin on Human Spermatozoa1. Biol. Reprod. [Internet] 72:1064–1070. Available from: http://dx.doi.org/10.1095/biolreprod.104.036483

Weyer S, Pääbo S. 2016. Functional Analyses of Transcription Factor Binding Sites that Differ between Present-Day and Archaic Humans. Mol. Biol. Evol. [Internet] 33:316–322. Available from: http://dx.doi.org/10.1093/molbev/msv215

Widagdo J, Fang H, Jang SE, Anggono V. 2016. PACSIN1 regulates the dynamics of AMPA receptor trafficking. Sci. Rep. [Internet] 6:31070. Available from: http://dx.doi.org/10.1038/srep31070

Wray GA. 2007. The evolutionary significance of cis-regulatory mutations. Nat. Rev. Genet. [Internet] 8:206. Available from: http://dx.doi.org/10.1038/nrg2063

Wu Y, Sheng W, Chen L, Dong H, Lee V, Lu F, Wong CS, Lu W-Y, Yang BB. 2004. Versican V1 Isoform Induces Neuronal Differentiation and Promotes Neurite Outgrowth. Mol. Biol. Cell [Internet] 15:2093–2104. Available from: http://www.molbiolcell.org/content/15/5/2093.abstract

Wynn T, Coolidge FL. 2004. The expert Neandertal mind. J. Hum. Evol. [Internet] 46:467–487. Available from: http://www.sciencedirect.com/science/article/pii/S0047248404000302

Wynn T, Overmann K, Coolidge F. 2016. The false dichotomy: a refutation of the Neandertal indistinguishability claim. J. Anthropol. Sci. = Riv. di Antropol. JASS [Internet] 94:201–221. Available from: http://europepmc.org/abstract/MED/26708102

Yan G, Zhang G, Fang X, Zhang Y, Li C, Ling F, Cooper DN, Li Q, Li Y, van Gool AJ, et al. 2011. Genome sequencing
and comparison of two nonhuman primate animal models, the cynomolgus and Chinese rhesus macaques. Nat.
Biotechnol. [Internet] 29:1019. Available from: http://dx.doi.org/10.1038/nbt.1992
Yap CC, Vakulenko M, Kruczek K, Motamedi B, Digilio L, Liu JS, Winckler B. 2012. Doublecortin (DCX) Mediates
Endocytosis of Neurofascin Independently of Microtubule Binding. J. Neurosci. [Internet] 32:7439 LP-7453.
Available from: http://www.jneurosci.org/content/32/22/7439.abstract
Yoshida MM, Azuma Y. 2016. Mechanisms behind Topoisomerase II SUMOylation in chromosome segregation. Cell
Cycle [Internet] 15:3151–3152. Available from: https://doi.org/10.1080/15384101.2016.1216928
Zachos G, Black EJ, Walker M, Scott MT, Vagnarelli P, Earnshaw WC, Gillespie DAF. 2017. Chk1 Is Required for
Spindle Checkpoint Function. Dev. Cell [Internet] 12:247–260. Available from:
http://dx.doi.org/10.1016/j.devcel.2007.01.003
Zhang H, Meltzer P, Davis S. 2013. RCircos: an R package for Circos 2D track plots. BMC Bioinformatics [Internet]
14:244. Available from: https://doi.org/10.1186/1471-2105-14-244
Zhang X, Ting K, Pathmanathan D, Ko T, Chen W, Chen F, Lee H, James AW, Siu RK, Shen J, et al. 2012. Calvarial
Cleidocraniodysplasia-Like Defects With ENU-Induced Nell-1 Deficiency. J. Craniofac. Surg. [Internet] 23.
Available from: https://journals.lww.com/jcraniofacialsurgery/Fulltext/2012/01000/
Calvarial_Cleidocraniodysplasia_Like_Defects_With.16.aspx
Zhang Y, Su J, Yu J, Bu X, Ren T, Liu X, Yao L. 2011. An essential role of discoidin domain receptor 2 (DDR2) in
osteoblast differentiation and chondrocyte maturation via modulation of Runx2 activation. J. Bone Miner. Res.
[Internet] 26:604–617. Available from: http://doi.wiley.com/10.1002/jbmr.225
Zhou H, Hu S, Matveev R, Yu Q, Li J, Khaitovich P, Jin L, Lachmann M, Stoneking M, Fu Q, et al. 2015. A
Chronological Atlas of Natural Selection in the Human Genome during the Past Half-million Years. bioRxiv
[Internet]. Available from: http://biorxiv.org/content/early/2015/06/19/018929.abstract
Zhu G-Z, Lin Y, Myles DG, Primakoff P. 1999. Identification of four novel ADAMs with potential roles in
spermatogenesis   and   fertilization.   Gene   [Internet]   234:227–237.   Available   from:
http://www.sciencedirect.com/science/article/pii/S0378111999002085
1193
1194

The *Homo* tree

Changes in humans

~4.4 million segregating sites

136,435 high frequency on human lineage → 12,027 fixed

Nonsense, UTR, Regulatory, etc

647 amino-acid changes

└→ 571 genes
└→ 42 fixed

Prioritizing
Ranking
Enrichment
Networks

Spindles & Cells

Mitotic Spindle   Neurogenesis

Brain & Bones

Brain growth

Craniofacial features

Complex traits

Life history   Language & cognition