Extensive cellular heterogeneity of X inactivation revealed by single-cell allele-specific

expression in human fibroblasts.

Marco Garieri [1, #], Georgios Stamoulis [1, #], Emilie Falconnet [1], Pascale Ribaux [1], Christelle

Borel [1, †, *], Federico Santoni [1,3, †, *] and Stylianos E. Antonarakis [1, 2, 4, †, *]

[1]Department of Genetic Medicine and Development, University of Geneva Medical School,

Geneva, Switzerland.

[2]University Hospitals of Geneva, Switzerland.

[3] Department of Endocrinology, Diabetology and Metabolism, University Hospitals of

Lausanne, Switzerland.

[4]iGE3 Institute of Genetics and Genomics of Geneva, Switzerland.

# These authors contributed equally to this work

†These authors contributed equally to this work

* Corresponding Authors

Address for correspondence:

Stylianos E. Antonarakis, Christelle Borel

Department of Genetic Medicine and Development,

University of Geneva Medical School, Geneva, Switzerland.

1 rue Michel-Servet

1211 Geneva, Switzerland

Tel +41-22-379-5707

Fax +41-22-379-5706

Email christelle.borel@unige.ch, stylianos.antonarakis@unige.ch

Federico Santoni

Department of Endocrinology, Diabetology and Metabolism,

Lausanne University Hospital (CHUV), Lausanne, Switzerland

7 rue de Bugnon,

1111 Lausanne, Switzerland

email: federico.santoni@chuv.ch

Running title: X-inactivation in single cells.

Keywords: single cell, ASE, X-inactivation

ABSTRACT

In eutherian mammals, X chromosome inactivation (XCI) provides a dosage compensation mechanism where in each female cell one of the two X chromosomes is randomly silenced. However, some genes on the inactive X chromosome and outside the pseudoautosomal regions escape from XCI and are expressed from both alleles (escapees). Given the relevance of the escapees in biology and medicine, we investigated XCI at an unprecedented single-cell resolution. We combined deep single-cell RNA sequencing with whole genome sequencing to examine allelic specific expression (ASE) in 935 primary fibroblast and 48 lymphoblastoid single cells from five female individuals. In this framework we integrated an original method to identify and exclude doublets of cells. We have identified 55 genes as escapees including 5 novel escapee genes. Moreover, we observed that all genes exhibit a variable propensity to escape XCI in each cell and cell type, and that each cell displays a distinct expression profile of the escapee genes. We devised a novel metric, the Inactivation Score (IS), defined as the mean of the allelic expression profiles of the escapees per cell, and discovered a heterogeneous and continuous degree of cellular XCI with extremes represented by "inactive" cells, i.e., exclusively expressing the escaping genes from the active X chromosome, and "escaping" cells, expressing the escapees from both alleles. Intriguingly we found that *XIST* is the major genetic determinant of IS, and that XIST expression, higher in G0 phase, is negatively correlated with the expression of escapees, inactivated and pseudoautosomal genes. In this study we use single-cell allele specific expression to identify novel escapees in different tissues and provide evidence of an unexpected cellular heterogeneity of XCI driven by a possible regulatory activity of *XIST*.

INTRODUCTION

In eutherian mammals, X chromosome inactivation (XCI) is a well-described mechanism of dosage compensation for the X chromosome in females (Lyon 1961; Penny et al. 1996; Chow and Heard 2009; Bartolomei and Ferguson-Smith 2011). In female cells, only one X chromosome is transcribed (Xa; X-active), whereas the second X chromosome is silenced (Xi; X-inactive) (Lyon 1961). Marsupials have an imprinted pattern of XCI, and the paternal allele is predominantly inactive (Sharman 1971). In mice, an imprinted form of XCI occurs

through early embryonic developmental stages (4-8 cell stage)(Huynh and Lee 2003; Okamoto et al. 2004; Okamoto et al. 2005; Patrat et al. 2009), followed by inner cell mass reactivation and random XCI in epiblast cells (Mak et al. 2004). In humans, the two X chromosomes are active during post-zygotic stages, achieve gene dosage compensation by dampening their expression up to or even after late blastocyst formation, until one of the X chromosomes is randomly inactivated in each cell (Petropoulos et al. 2016). In female somatic cells, random XCI is stable, resulting in a mosaicism for gene expression on the X chromosome, in which an average of 50% of cells express the active paternal X and 50% the active maternal X alleles. Most of the genes on the Xi chromosome are transcriptionally silenced through epigenetic processes initiated by the X Inactivation Center (XIC) and spread along the Xi chromosome during early embryogenesis (Lyon 1961). The XIC encodes several genes, including *XIST,* a long non-coding RNA (ncRNA) essential for initiating and completing XCI (Brown et al. 1991; Ballabio and Willard 1992; Brown et al. 1992). *XIST* RNA molecules mediate the establishment and maintenance of XCI in subsequent cycles of mitotic division by coating the Xi chromosome and recruiting Polycomb Repressive Complex 2 with repressive chromatin modifiers (Lee and Bartolomei 2013). It has been shown that the coating of the Xi is regulated by the interaction between *XIST* ncRNA and the Lamin B receptor (LBR)(Chen et al. 2016a). This interaction is needed for the recruitment of Xi to the nuclear lamina and the subsequent spread of *XIST* ncRNA to actively transcribed regions (Chen et al. 2016a).

However, not all X-linked genes are inactivated. In females, genes that escape from XCI (escapees) represent 15-25% of the X-linked genes, and a further 10% of escapees differ between individuals and cell types (Carrel and Willard 2005; Prothero et al. 2009; Yang et al. 2010; Cotton et al. 2013; Crowley et al. 2015). Such genes have been associated to sex-specific traits and to clinical abnormalities in patients with X chromosome aneuploidy, such as Turner and Klinefelter Syndromes (Berletch et al. 2011). Pathogenic variants in escapees also contribute to various disease phenotypes in women carriers, including Kabuki syndrome (KABUK1 [MIM 147920])(Lederer et al. 2012; Miyake et al. 2013), intellectual disabilities(van Haaften et al. 2009; Grasso et al. 2012; Jones et al. 2012; Gropman and Samango-Sprouse 2013; Zhang et al. 2013; Dunford et al. 2016). Genes escaping XCI have been previously identified by whole tissue studies using different approaches, such as X-linked gene expression comparisons between males and females (Yasukochi et al. 2010), detecting allelic imbalance in clonal lymphoblast and fibroblast cell lines (Cotton et al. 2013),identifying inactivated and active transcription start sites by methylation profiles

(Cotton et al. 2015)and among female individuals with X chromosome aneuploidies (Sudbrak

et al. 2001)

The ability to capture single cells and to study their allele-specific expression (ASE) (Borel et

al. 2015) provides the opportunity to explore XCI patterns at the single-cell level and to

identify escapee genes. Recent studies on mouse single cells demonstrated the robust nature

of this technology to monitor the dynamics of XCI through differentiation(Chen et al. 2016b),

mouse preimplantation female embryos (Borensztein et al. 2017) and in clonal somatic cells

(Reinius et al. 2016). Recently, (Tukiainen et al. 2017) performed an across-tissue study of X

inactivation and partially validated their observation performing shallow sequencing (1Mio

reads x cell) on 940 single cells from lymphoblasts and dendritic cells. Here, using RNA-Seq

at high sequencing depth (40Mio reads per cell), we studied the X-linked ASE in 983

isolated, unsynchronized single fibroblast and lymphoblast cells and established the degree of

XCI after the removal of potential confounding effects. One of the caveats of allele

expression quantification in single cells is represented by the allele dropout, which randomly

affects the detection of one of the two alleles of poorly expressed genes (Stegle et al. 2015).

However, in this context, the allelic dropout will not induce false positive escapees. Moreover

given the number of cells analyzed in our study, the probability to consistently miss the

capture of the expressed allele from the X-inactive chromosome of a true escapee in all the

cells is extremely low. In this study we identified 55 escapee genes in at least one individual,

out of which 5 were novel. A subset of 22 genes was detected as escapee in at least two

individuals (robust set), including 3 novel escapee genes. Through the analysis of the

expression profile of the genes in the robust escapee set, we further investigated their

propensity to escape XCI in each cell. We discovered that the transcriptional activity of the

escapee genes is highly variable and significantly associated with the cellular *XIST* transcript

abundance.

RESULTS

Identification and elimination of confounding doublets

Genes located on the X chromosome of female cells express one allele from the randomly

active chromosome, while escaping genes express both alleles (Fialkow 1970). Multiple cells

(i.e., doublets) resulting from the simultaneous capture of two or more cells expressing

discordant haplotypes obviously complicate the detection of escapee genes and potentially

increase the number of false positives. A recent study has explored the limits of current

fluidic technology to conduct single-cell RNAseq, demonstrating that a substantial fraction of

the bona fide isolated single cells are doublets (Macosko et al. 2015). After removing 32 cells

because of low mapping quality (Supplemental Figure S1), in order to eliminate all doublets

with discordant haplotypes, we conducted a pairwise correlation analysis of X-linked ASE

among all the cells. We performed hierarchical clustering analysis and obtained three distinct

clusters of cells (Figure 1B and Supplemental Figure S2, Methods). As expected, two clusters

were populated by cells with one mutually inactivated X chromosome. The third cluster

included cells with a biallelic expression pattern for all the X chromosome genes, revealing

the presence of doublets. Considering all 5 individuals (n = 983 cells), we identified a total of

82 doublets (~8% of the total, consistent with Fluidigm manufacturer's expectations), which

were excluded from further analysis (Figure 1B, Supplemental Figure S2 and Supplemental

Table S1). Doublets expressing concordant haplotypes cannot be detected with this our *in*

silico approach; however, they do not inflate the number of false positive escapees and,

consequently, they do not have an impact on escapee gene discovery and XCI analysis.

Identification of escapee genes

To estimate the allelic ratio (AR) for each gene in each cell, we calculated the ratio of the

number of reads supporting the cell-specific expressed haplotypes over the total number of

reads covering all single nucleotide variants (SNV) of a gene (See Methods). Fully

inactivated genes displayed an AR equal to 1. In the relaxed discovery set of escapee genes,

putative escapees were defined as having an AR =< 0.95 in at least one individual. The

rational and the choice of this threshold is explained in the methods. The gene is considered

as inactivated (i.e. exclusively expressed from the active chromosome) otherwise. As a proof

of principle of the reliability of our approach we first confirmed that *XIST* is expressed

exclusively from the inactivated allele by analyzing its AR in all cells from individuals 3 and

4 (i.e monozygotic twin samples), for which we were able to phase the haplotypes from

parental genotyping (Supplemental Figure S3). As an additional control we examined the

allele expression profile of genes in PAR1 and PAR2 regions. As expected, all of these genes

exhibited a balanced ASE across the two X chromosomes except *VAMP7* in PAR2 where

very few cells displayed an expression from the inactive chromosome (Figure 2,

Supplementary Table S2).

As expected, the majority of chrX genes showed an inactivated status in all the cells (Figure 2
and Supplemental Table S2). From a total of 296 genes interrogated in at least one individual,
we identified the relaxed set of 55 escapees (18.5%): 50 of them previously described to
escape XCI in at least one study, and 5 novel escapees (*INE2* (antisense gene) , *STK26,*
UQCRBP1, LINC00630 and *TTC3P1)* (Figure 2, S3). Out of 203 genes with AR information
from at least 2 individuals, we classified as robust escapees 22 genes (10.9%) which exhibit
an escapee status in at least two individuals including 3 novel genes (*INE2*, *STK26* and
TTC3P1). As expected, the power to detect a gene escaping XCI is linearly related with the
respective expression level (Supplemental Figure S4). The number of overlapping escapees
among all individuals is shown in a Venn diagram (Supplemental Figure S5). Results from
both relaxed and robust sets are consistent with the current understanding of XCI, in which is
predicted that 10% to 20% of X chromosome genes escape XCI (Carrel and Willard 2005).
In addition, we analyzed 48 single cells from a lymphoblastoid cell line derived from the
individual 5 to investigate the escapee concordance with the fibroblasts. After quality control
and doublets removal, we were able to classify 9 genes as escapee in lymphoblastoid cells
with 5 of them being known escapees (*DDX3X*, *KDM6A*, *MSL3*, *PUDP*, *ZFX*)*,* and 4 novel
escapee genes (*IDS*, *SLC9A7*, *STAG2*, *STK26*)*.* We observed that, though expressed and
having an informative heterozygous site, the *MSL3, IDS*, *SLC9A7*, and *STAG2* genes were
not classified as escapees in fibroblasts. Conversely several genes detected as escapees in
fibroblasts were not escapees in lymphoblast cells. This could be partially ascribed to the
previously observed tissue specificity of XCI (Deng et al. 2014b; Tukiainen et al. 2017) and
to the recently discovered peculiar maintenance of XCI in lymphoblast cells **(Wang et al.**
2016). Additionally, the relatively small number of lymphoblastoid single cells along with the
absence of informative heterozygous sites for some genes in other individuals, can affect our
ability to detect escapees(Santoni et al. 2017) and to assess and establish their ASE. However
the *STK26* escapee status was clearly confirmed in both fibroblast and lymphoblastoid single
cells.
198

Heterogeneity of escaping XCI

Among five individuals, the 22 robust escapee genes exhibited a heterogeneous AR profile,
being inactivated in some cells and escaping XCI in others (Figure 3). Specifically, we
calculated a cellular escaping ratio per escapee gene as the proportion of cells escaping XCI
with respect to the total number of cells expressing the gene. Some genes displayed a stable
cellular escaping ratio among all the individuals, while others were more variable. For

example, *CA5BP1* showed consistent cellular escaping ratios ranging from 37% to 52% ;
ZFX, a known constitutive escapee (Schneider-Gadicke et al. 1989), presented with cellular
escaping ratios ranging from 85% to 100%, while *DDX3X* had a broader range going from
29% to 61%. Overall, escapees had different cellular escaping ratios, thereby suggesting that
each escapee gene is independently regulated.

The observed cellular pattern of XCI of the escapees (Figure 3) suggests a variable cellular
ability of expressing genes from the inactivated allele. To investigate this hypothesis, we
calculated the Inactivation Score (IS) for each cell defined as the mean AR of the escapee
genes detected per cell (we considered only cells expressing at least two escapees). For each
individual, we ordered the cells according to the respective IS and discovered that the
capacity to express from the inactive chromosome is a continuous variable (Figure 3, Figure
4A). This suggested a cellular stratification that reflect the propensity of each cell to escape
XCI, as confirmed by the proportion of escapee genes per cell (Figure 4B). As expected, the
Inactivation Score is strongly negatively correlated with expression from the inactive X
chromosome (Supplemental Figure S6). Notably, in all five individuals, we observed two
special groups of cells: one in which all the detectable escapees behaved as inactivated and
another where all detectable escapees expressed both X alleles (Figure 4B). These two
extreme cell populations represent on average 15% of the total number of aggregate cells
among the individuals (individual 1: 20%, individual 2: 14%, individual 3: 8%, individual 4:
7%, individual 5: 26%). As a control, we calculated the Inactivation Score of the remaining
inactivated genes per cell (all close to 1, as expected. Figure 4A, right panel). Overall, these
results demonstrate that XCI is a complex intra- and inter-individual heterogeneous process,
and the ability to escape from X inactivation varies from gene to gene, from cell to cell and
also among individuals. The evidence of similar cell stratification in all individuals suggests
the existence of a general regulatory mechanism that controls the propensity of a cell to
express genes from the inactivated allele.

Potential drivers of cellular XCI heterogeneity

We hypothesized that the cellular XCI heterogeneity may be associated with the level of
expression of genes on the X chromosome. To explore this hypothesis, we correlated the X-
linked gene expression (RPKM > 1) with the Inactivation Score. After FDR correction for
multiple testing, genes were ranked based on the adjusted p-value (Figure 5A). Notably the

only gene positively and significantly correlated with IS was *XIST* (nominal p-value = 3.2 x

10^{-5}; adj. p-value = 3.0 x 10^{-3}). *XIST* is a well-known non-coding RNA that regulates the

establishment and the maintenance of XCI (**Brown et al. 1991**). Other significantly albeit

negatively associated genes were *EIF2S3*, *CD99*, *NDUFA1*, *RPL10* and *BCAP31*. We further

investigated the correlation of *XIST* expression with inactivated genes and all escapee genes

(relaxed and robust sets) and genes located in PAR regions. Notably we observed a

significant trend of negative correlations between *XIST* expression and the majority of the

genes for all four categories (H0: mean of distribution of correlation coefficients = 0;

inactivated genes p = 2.06 x 10^{-83} ; PAR genes p= 6.49 x 10^{-4}; Relaxed escapee p= 1.01 x 10^{-6};

Robust escapees p= 4.19 x 10^{-4}, Mann-Whitney two-tailed test, Figure 5B). This suggests a

general repressive *XIST* regulatory effect in all classes of X-linked genes. To further

investigate whether this effect was cell cycle dependent, we used Cyclone (Scialdone et al.

2015) to assign each cell to a specific cell cycle phase according to the expression of the

appropriate gene markers (from CycleBase (Santos et al. 2015)). Cells not expressing *MKI67*

have been previously reported to be in G0 (Sobecki et al. 2017). A statistically significant

difference between the populations of single cells in G0 and G1 was observed regarding *XIST*

expression (Figure 6). In agreement with our previous observations, the Inactivation Score

was also significantly higher in G0 than in G1 cells on average (Figure 6). These results

together support the hypothesis that *XIST* ncRNA tends to be more expressed in the resting

G0 phase than in G1 and, consequently, the expression of the escapees from the inactive

chromosome is reduced in this cell cycle stage. We could not identify any cell-cycle driven

effect for the remaining cell cycle stages, likely due to a limited statistical power (small

number of single cells classified in these cell cycle stages, Figure 6).

DISCUSSION

Our study, using human single-cell RNAseq datasets, points to a pervasive heterogeneity in

escaping XCI. We have shown that escapees have a different allelic expression profile in

single fibroblasts from the same individual. More than 50% of the escapees had the tendency

to be mainly expressed from Xa (Figure 4), while *ZFX* and *PUDP* exhibited an overall

biallelic expression in more than 70% of the cells. We also observed that some escapees

exhibited a relatively stable cellular escaping ratio (proportion of cells in which the gene is

escaping), i.e., *ZFX* and *CA5BP1*, where others, such as *DDX3X*, showed broader variability

(Figure 3). This finding explains, from a single-cell perspective, the previous observations of

heterogeneous gene expression from Xi in cell lines derived from different individuals (Carrel and Willard 1999) and tissues (Cotton et al. 2013; Berletch et al. 2015). A more recent study revealed the contribution of 6 escapee genes (*TRX*, *CNKSR2*, *DDX3X*, *KDM5C*, *KDM6A*, *MAGEC3*) to cancer sex bias (Dunford et al. 2016). Here, we confirmed the escapee status for *DDX3X, KDM5C* and *KDM6A*. We observed that the inter-individual *DDX3X* inactivation profile is highly variable and thereby could potentially be associated with differences in cancer predisposition occurring among female individuals.

Among the genes that we detected as escapees in this study, *HUWE1* encodes for an E3 ubiquitin ligase and microduplications of this gene have been found in individuals affected by Turner type intellectual disability (MRX17 [MIM 300706])(Froyen et al. 2008; Froyen et al. 2012). Moreover a recent study suggested the critical role of *HUWE1* as a colonic tumor suppressor gene (Myant et al. 2017). *RBMX* encodes an RNA binding protein which is involved in pre- and post-transcriptional regulation and alternative splicing of many premature transcripts. Pathogenic variants in *RBMX* have been associated with X-linked intellectual disability (Shashi et al. 2000; Shashi et al. 2015). *STK26* is a member of germinal center kinase subfamily and encodes a serine/threonine kinase expressed highly in human lymphocytes and thymus (Ceccarelli et al. 2011; Jiao et al. 2015). Expression of *STK26* (alternatively named as *MST4*) was recently shown to be significantly lower in patients affected by Graves' disease, an autoimmune disorder characterized by abnormal thyroid function (Guo et al. 2017). It has also been shown that patients affected with Turner Syndrome (XO) exhibit a high prevalence of hypothyroidism (El-Mansoury et al. 2005). We speculate that this phenotype may be related to the reduced expression of the single copy of *STK26*.

Mutations and deletions of the escapee genes identified in the relaxed set such as *EDA*, *FHL1*, *IL1RAPL1*, and *FMR1* have been associated with different pathological conditions. More specifically pathogenic variants of *EDA*, a gene of the tumor necrosis factor family, have been associated with X-linked dominant tooth agenesis and X-linked ectodermal dysplasia (Kere et al. 1996) (Yotsumoto et al. 1998) (Visinoni et al. 2003) (Tao et al. 2006) (Tarpey et al. 2007). *FHL1* encodes a member of the four-and-a-half-LIM-only protein family and contains two highly conserved zinc finger domains. Pathogenic variants in *FHL1* have been associated with X-linked dominant reduced body myopathy with both affected male and female individuals (Liewluck et al. 2007) (Schessl et al. 2008; Shalaby et al. 2009), Emery-Dreifuss muscular dystrophy (Gueneau et al. 2009), and scapuloperoneal myopathy (Quinzii et al. 2008). Pathogenic variants of *IL1RAPL1*, a member of interleukin 1 receptor

family, have been linked to X-linked mental retardation (Tabolacci et al. 2006; Nawara et al. 2008). A trinucleotide repeat expansion (CGG) in the 5' UTR of *FMR1* is the cause of Fragile X syndrome (Kremer et al. 1991). We hypothesize that the phenotypic variability of these diseases may be partially explained by variable penetrance and variable expressivity issues in these gene-disease associations due to the observed variable escapee status of their respective causative genes in the relevant tissues.

It has been recently observed in a mouse study that some genes escapee XCI in a fraction of cells only (Reinius (Reinius et al. 2016)et al .2016). We confirmed and extended this observation with the finding that XCI heterogeneity extends from genes to single cells. Using the Inactivation Score as a metric for cellular propensity to escape XCI, we indeed showed that cells from the same individual have a variable propensity to transcribe the escapee genes from Xi. According to the IS, those cells stratify into three classes: cells where all escapees are expressed from the inactive allele, cells where all escapees are inactivated, and cells with an intermediate profile (Figure 4B). We have shown that the observed cellular heterogeneity is driven by the expression profile of *XIST* in single cells which is in turn strongly and positively correlated with the Inactivation Score. We found five genes negatively associated with IS. *EIF2S3* encodes the gamma subunit of the translation initiation factor 2 which is involved in protein synthesis initiation **(Moortgat et al. 2016)**. This gene has been described to escape X inactivation while hemizygous mutations have been associated with X mental retardation Borck type (Ehrmann et al. 1998) (Moortgat et al. 2016). *CD99* is located in the PAR1 region and is involved in leukocyte migration, and T-cell adhesion (Bernard et al. 1997) (Bernard et al. 2000). Additionally it has been characterized as a potential oncosuppressor in osteosarcoma(Manara et al. 2006). *NDUFA1* **(**NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 1**)** encodes a component of the mitochondrial respiratory chain; hemizygous mutations have been associated with mitochondrial complex I deficiency (Fernandez-Moreira et al. 2007) (Ambros et al. 1997; Potluri et al. 2009). *RPL10* is involved in protein synthesis and is an essential component of large 60s ribosomal subunit (Gong et al. 2009). Hemizygous *RPL10* mutations have been associated with syndromic X-linked mental retardation (Thevenon et al. 2015). *BCAP31* is involved in the export of secreted proteins and hemizygous loss of function mutations have been identified in individuals affected with deafness and severe dystonia**(Cacciagli et al. 2013)**.

Additionally, we demonstrated that *XIST* expression is negatively correlated with the overall expression of all genes in chrX including pseudoautosomal and escapee genes, thereby indicating a general repressive regulatory activity of *XIST* transcripts. Inspired by a previous

report describing the localisation of XIST during the cell cycle (Jonkers et al. 2008), we

explored the hypothesis that *XIST* expression and IS are related to the cell cycle phases of

individual cells. Our data suggest that *XIST* is more expressed during the G0 than the G1

phase and single cells in the G0 phase have an increased IS, indicating less propensity to

express escapees from Xi. Overall, these results revealed a considerable degree of

heterogeneity of XCI in single cells and enhance our understanding of X inactivation. The

framework presented in this study can be applied on an increased number of cells extracted

from various tissues from a large cohort of individuals. Such investigations will enable us to

define the extent of cellular XCI heterogeneity and clarify its possible implication in the

phenotypic variability of X-linked single gene disorders, whole X chromosome aneuploidies

and to the observed female sex bias in cancer.

MATERIAL AND METHODS

Ethical statement

The study was approved by the ethics committee of the University Hospitals of Geneva, and

written informed consent was obtained from both parents of each individual prior to the

study.

Samples

We established 6 different cell lines from 5 female individuals: 5 primary fibroblast cell lines

and one lymphoblastoid cell line (Table S1). We captured 935 single-cell fibroblasts and 48

lymphoblastoid single cells. Lymphoblastoid cells obtained from one of the five female

individuals (Figure 1A, Table S1)(Borel et al. 2015; Santoni et al. 2017).

Cell growth

Cells were cultured in DMEM GlutaMAX™ (Life Technologies) supplemented with 10%

fetal bovine serum (Life Technologies) and 1% penicillin/streptomycin/fungizone mix

(Amimed, BioConcept) at 37°C in a 5% CO_2 atmosphere as described (Borel et al. 2015).

Whole Genome Sequencing

Genomic DNA was extracted for all five individuals using a QIAamp DNA Blood Mini Kit

(Qiagen) and fragmented by Covaris to peak sizes of 300–400 bp. Libraries were prepared

with TruSeq DNA kit (Illumina) using 1 µg of gDNA and sequenced on an Illumina HiSeq

2000 machine with 2 x 100-bp as previously described(Borel et al. 2015). All experiments

were performed using the manufacturer's protocols. For each individual, raw whole genome

DNA sequences were analyzed using an in-house pipeline that utilizes published algorithms

in a sequential manner (BWA) (Li and Durbin 2010) for mapping the reads over the hg19

reference, SAMtools (Li et al. 2009) for detection of heterozygous sites, and (ANNOVAR)

for the annotation (Wang et al. 2010).

Single-cell capture

Single-cell capture was performed using the C1 single-cell auto prep system (Fluidigm)

following the manufacturer's procedure(Borel et al. 2015). The integrated fluidic circuit used

for the study is the C1™ Single-Cell mRNA Seq IFC, 17–25 µm with a capacity of 96

chambers. During the capture, all 96 chambers were inspected under an inverted phase

contrast microscope; only chambers containing a non-damaged single cell were considered

for downstream analysis.

Single-cell RNA-sequencing

SMARTer Ultra Low RNA kit for Illumina sequencing (version 2, Clontech) was used for the

cell lysis and cDNA synthesis. Libraries were prepared with 0.3 ng of pre-amplified cDNA

using Nextera XT DNA kit (Illumina) as described (Borel et al. 2015). Libraries were

sequenced on an Illumina HiSeq2000 sequencer as 100 bp single-ended reads. RNA

sequences were mapped with GEM (Marco-Sola et al. 2012). Uniquely mapping reads were

extracted by filtering for mapping quality (MQ>=150). An in-house algorithm was used to

quantify RPKM expression using GENCODE v26. Cells with less than 1 million uniquely

mapped reads were excluded from further analysis (Supplemental Figure S7).

Allele-specific expression and classification of escapee genes

For each gene on the X chromosome, the aggregate monoallelic ratio (AR) per cell was

calculated by averaging the allelic ratio of the reads covering the respective heterozygous

sites (AR = sum of number of reads from the active X allele / total SNV reads; $0 \leq AR \leq 1$).

Since we do not have the availability of parental genotype for all the individuals, we designed

an algorithm to estimate the active X allele per site based on the assumption that the active X

allele is, on average, more transcribed than the inactive X. We validated this assumption

comparing the prediction of our algorithm with the phasing of twins' X alleles based on

parental information (more details in Supplemental Text S1). According to this metric,

inactivated genes cluster around AR=1 while known escapees appear as been uniformly distributed between 0.5≤AR≤0.95 (linear phase of the cumulative distribution, Supplemental Figure S7). As support of this observation, AR distribution of autosomal genes clearly indicates AR=0.95 as the threshold separating biallelically expressed genes from monoallelic expressed genes (Supplemental Figure S8). Therefore, we consider a gene as escapee in the relaxed set when the aggregate AR is ≤ 0.95 in at least 1 individual and as escapee in the robust set when the aggregate AR is ≤ 0.95 in at least 2 individuals. To reduce the effect of allele dropout, we only consider for the analysis SNV sites covered by at least 5 reads in at least three cells. To reduce sampling bias effects (Deng et al. 2014a) a gene is included in the analysis if detectable in more than 5 different cells and/or SNVs per sample.

Haplotype and multiple cells (doublets) detection

For each cell, the expressed haplotype was estimated by calculating the allelic ratio of each heterozygous site in the X chromosome as identified by whole genome sequencing, excluding sites in the PAR regions (PAR1: chrX:60001-2699520, PAR2: chrX:154931044-155260560) and in known escapee genes (see section Annotation of known escapee genes). The estimated haplotype of each cell was compared to all others through pairwise correlation based hierarchical clustering procedure. A comparison of cells expressing concordant and discordant haplotypes results in a correlation near 1 and -1 respectively. Doublets simultaneously expressing both haplotypes cluster around an absolute correlation of 0.5 are identified and excluded from further analysis.

Annotation of the escapee genes

First, we curated a list of 190 previously observed escapee genes in different cell types and tissues according to the literature (Carrel and Willard 2005) (Johnston et al. 2008; Park et al. 2010; Yasukochi et al. 2010; Sharp et al. 2011) (Cotton et al. 2013; Zhang et al. 2013)(Table S3). Specifically, we investigated the status of 115 known escapee genes with available informative heterozygous sites and being expressed in fibroblast and lymphoblast cell lines (Supplemental Table S3). Second, we have appended the results published in two studies (Balaton et al. 2015; Tukiainen et al. 2017) in Table S2. Genes detected as escapees in our studies, in absence of citation, have been labeled as novel escapee genes. Genes found as escapee in our study and found subject to inactivation in other studies have been labeled variable escapee genes.

Cell cycle phase assignment

G1, S, and G2M cell cycle stage related gene markers were obtained from CycleBase (Santos et al. 2015) Cells not expressing *MKI67* have been considered to be in G0 (Schonk et al. 1989) The remaining cells were assigned to their respective cell cycle phase according to the expression of CycleBase genes with Cyclone (Scialdone et al. 2015)**.**

ACKNOWLEDGMENTS

We thank the laboratory of E. T. Dermitzakis for discussions. This work was supported by grants from the Swiss National Science Foundation (SNF-144082), and the European Research Council (ERC-249968) and the ChildCare foundation to S.E.A.. The computations were performed at the Vital-IT (http://www.vital-it.ch) Center for high-performance computing of the SIB Swiss Institute of Bioinformatics.

ACCESSION NUMBERS

Newly generated RNA and DNA sequencing data are deposited in the European Genome-phenome Archive (EGA, https://www.ebi.ac.uk/ega/) for controlled accesses; the study accession number is (EGASxxxx, to be determined).

LEGENDS TO FIGURES

Figure 1. Identification and elimination of confounding doublets.

(A) Flowchart of the study. Whole genome sequencing for each individual identified the
respective heterozygous sites. Single-cell RNA-seq provided RNA abundance for each single
cell. After QC and removal of the doublets, our framework calculated AR for the X-linked
genes to enable escapees identification and study the cellular XCI. **(B)** Heat map of
unsupervised hierarchical clustering using cell-cell pairwise Pearson correlation coefficients
of AR from single cells from individual 3 before (left) and after (right), excluding the
identified doublets. The heat map separated the cells expressing one haplotype (blue and
green cluster) from cells expressing two haplotypes (black cluster, doublets). Pearson
correlations range from -1 (red) to 1 (white).

Figure 2. Single cell Allelic Ratio profiles with respect to the active haplotype for genes
on female X chromosome. For each individual, the Allelic Ratio for each gene (fibroblasts
or lymphoblasts) is reported for each cell along the x-axis (rectangles with AR ≥ 0.95 (blue)
an AR ≤ 0.95 (red)) according to the genomic location of genes in the human X chromosome
(y-axis). 55 identified escapee genes in at least one individual are annotated on the left of the
X chromosome ideogram. Known escapees are shown in red; novel escapees in black;
escapees from the robust set with an asterisk. *XIST* is shown in green. PAR, pseudoautosomal
regions; LCL, lymphoblastoid cells.

Figure 3. Single cell ASE profile of 22 robust escapee genes in human female fibroblasts.
Composite figure of individual allelic ratios per gene per cell **(Top of the panel)**. Allelic
Ratio profile of robust escapee genes (listed the rows) with a detectable expression in single
cells (ordered along the columns) is shown. Every dot represents the allelic ratio of the
respective gene in a cell. AR ranges from 0 (light orange) to 1 (dark blue). The size of the dot
is proportional to the respective number of reads detected per cell. (%) is the percentage of
cells where the respective gene is escaping XCI. **(Bottom of the panel).** Bar plot representing
the Inactivation Score (see text for details) per cell. IS ranges from 0 (light orange) to 1 (dark
blue).

Figure 4. Cellular propensity to escape XCI.
(A) Cells ranked by the Inactivation Score calculated on all escapees in the robust set (left -
Escapee) and on all inactive genes (right - Inactive) per individual (color coded - see legend).
Each dot represents a fibroblast cell. **(B)** Y-axis - cells ordered according to the percentage of

escapees displaying an inactive status (AR ≥ 0.95, with respect to all detectable escapees) per

individual (x-axis). Each dot represents a fibroblast cell.

Figure 5. XIST association and correlation with IS and its repressive role for X

chromosome genes.

(A) Manhattan-like plot for Pearson correlation of X-linked genes expression with

Inactivation Score. The plot represents -$\log_{10}$ p-value against X chromosome position.

Vertical lines represent PAR region limits. Red horizontal lines represent the significance

threshold for positively and negatively correlated genes, respectively. **(B)** Distributions of

correlations of expression of XIST with inactivated genes (upper left), PAR located genes

(upper right), escapees in robust set (lower left), escapees in relaxed set (lower right). P-

values calculated with Mann-Whitney test (see text for details).

Figure 6. XIST expression and IS dependency on cell cycle

Distribution of *XIST* expression (left) and of Inactivation Scores per single cells (right)

according to cell cycle phases (n=number of cells per stage). P-values calculated with Mann-

Whitney test (see text for details)

Supplemental Data

Supplemental data include eight figures, additional methods and three tables.

References

Ambros IM, Rumpler S, Luegmayr A, Hattinger CM, Strehl S, Kovar H, Gadner H, Ambros
PF. 1997. Neuroblastoma cells can actively eliminate supernumerary MYCN gene
copies by micronucleus formation--sign of tumour cell revertance? *Eur J Cancer* **33**:
2043-2049.
Balaton BP, Cotton AM, Brown CJ. 2015. Derivation of consensus inactivation status for X-
linked genes from genome-wide studies. *Biol Sex Differ* **6**: 35.
Ballabio A, Willard HF. 1992. Mammalian X-chromosome inactivation and the XIST gene.
Curr Opin Genet Dev **2**: 439-447.
Bartolomei MS, Ferguson-Smith AC. 2011. Mammalian genomic imprinting. *Cold Spring
Harbor perspectives in biology* **3**.
Berletch JB, Ma W, Yang F, Shendure J, Noble WS, Disteche CM, Deng X. 2015. Escape
from X inactivation varies in mouse tissues. *PLoS Genet* **11**: e1005079.
Berletch JB, Yang F, Xu J, Carrel L, Disteche CM. 2011. Genes that escape from X
inactivation. *Human genetics* **130**: 237-245.

Bernard G, Breittmayer JP, de Matteis M, Trampont P, Hofman P, Senik A, Bernard A. 1997.
Apoptosis of immature thymocytes mediated by E2/CD99. *J Immunol* **158**: 2543-
2550.
Bernard G, Raimondi V, Alberti I, Pourtein M, Widjenes J, Ticchioni M, Bernard A. 2000.
CD99 (E2) up-regulates alpha4beta1-dependent T cell adhesion to inflamed vascular
endothelium under flow conditions. *Eur J Immunol* **30**: 3061-3065.
Borel C, Ferreira PG, Santoni F, Delaneau O, Fort A, Popadin KY, Garieri M, Falconnet E,
Ribaux P, Guipponi M et al. 2015. Biased allelic expression in human primary
fibroblast single cells. *Am J Hum Genet* **96**: 70-80.
Borensztein M, Syx L, Ancelin K, Diabangouaya P, Picard C, Liu T, Liang JB, Vassilev I,
Galupa R, Servant N et al. 2017. Xist-dependent imprinted X inactivation and the
early developmental consequences of its failure. *Nat Struct Mol Biol* **24**: 226-233.
Brown CJ, Ballabio A, Rupert JL, Lafreniere RG, Grompe M, Tonlorenzi R, Willard HF.
1991. A gene from the region of the human X inactivation centre is expressed
exclusively from the inactive X chromosome. *Nature* **349**: 38-44.
Brown CJ, Hendrich BD, Rupert JL, Lafreniere RG, Xing Y, Lawrence J, Willard HF. 1992.
The human XIST gene: analysis of a 17 kb inactive X-specific RNA that contains
conserved repeats and is highly localized within the nucleus. *Cell* **71**: 527-542.
Cacciagli P, Sutera-Sardo J, Borges-Correia A, Roux JC, Dorboz I, Desvignes JP, Badens C,
Delepine M, Lathrop M, Cau P et al. 2013. Mutations in BCAP31 cause a severe X-
linked phenotype with deafness, dystonia, and central hypomyelination and
disorganize the Golgi apparatus. *Am J Hum Genet* **93**: 579-586.
Carrel L, Willard HF. 1999. Heterogeneous gene expression from the inactive X
chromosome: an X-linked gene that escapes X inactivation in some human cell lines
but is inactivated in others. *Proc Natl Acad Sci U S A* **96**: 7364-7369.
Carrel L, Willard HF. 2005. X-inactivation profile reveals extensive variability in X-linked
gene expression in females. *Nature* **434**: 400-404.
Ceccarelli DF, Laister RC, Mulligan VK, Kean MJ, Goudreault M, Scott IC, Derry WB,
Chakrabartty A, Gingras AC, Sicheri F. 2011. CCM3/PDCD10 heterodimerizes with
germinal center kinase III (GCKIII) proteins using a mechanism analogous to CCM3
homodimerization. *J Biol Chem* **286**: 25056-25064.
Chen CK, Blanco M, Jackson C, Aznauryan E, Ollikainen N, Surka C, Chow A, Cerase A,
McDonel P, Guttman M. 2016a. Xist recruits the X chromosome to the nuclear lamina
to enable chromosome-wide silencing. *Science* **354**: 468-472.
Chen G, Schell JP, Benitez JA, Petropoulos S, Yilmaz M, Reinius B, Alekseenko Z, Shi L,
Hedlund E, Lanner F et al. 2016b. Single-cell analyses of X Chromosome inactivation
dynamics and pluripotency during differentiation. *Genome research* **26**: 1342-1354.
Chow J, Heard E. 2009. X inactivation and the complexities of silencing a sex chromosome.
Curr Opin Cell Biol **21**: 359-366.
Cotton AM, Ge B, Light N, Adoue V, Pastinen T, Brown CJ. 2013. Analysis of expressed
SNPs identifies variable extents of expression from the human inactive X
chromosome. *Genome Biol* **14**: R122.
Cotton AM, Price EM, Jones MJ, Balaton BP, Kobor MS, Brown CJ. 2015. Landscape of
DNA methylation on the X chromosome reflects CpG density, functional chromatin
state and X-chromosome inactivation. *Hum Mol Genet* **24**: 1528-1539.
Crowley JJ, Zhabotynsky V, Sun W, Huang S, Pakatci IK, Kim Y, Wang JR, Morgan AP,
Calaway JD, Aylor DL et al. 2015. Analyses of allele-specific gene expression in
highly divergent mouse crosses identifies pervasive allelic imbalance. *Nature genetics*
47: 353-360.

Deng Q, Ramskold D, Reinius B, Sandberg R. 2014a. Single-cell RNA-seq reveals dynamic,
random monoallelic gene expression in mammalian cells. *Science* **343**: 193-196.
Deng X, Berletch JB, Nguyen DK, Disteche CM. 2014b. X chromosome regulation: diverse
patterns in development, tissues and disease. *Nature reviews Genetics* **15**: 367-378.
Dunford A, Weinstock DM, Savova V, Schumacher SE, Cleary JP, Yoda A, Sullivan TJ,
Hess JM, Gimelbrant AA, Beroukhim R et al. 2016. Tumor-suppressor genes that
escape from X-inactivation contribute to cancer sex bias. *Nature genetics*
doi:10.1038/ng.3726.
Ehrmann IE, Ellis PS, Mazeyrat S, Duthie S, Brockdorff N, Mattei MG, Gavin MA, Affara
NA, Brown GM, Simpson E et al. 1998. Characterization of genes encoding
translation initiation factor eIF-2gamma in mouse and human: sex chromosome
localization, escape from X-inactivation and evolution. *Hum Mol Genet* **7**: 1725-1737.
El-Mansoury M, Bryman I, Berntorp K, Hanson C, Wilhelmsen L, Landin-Wilhelmsen K.
2005. Hypothyroidism is common in turner syndrome: results of a five-year follow-
up. *J Clin Endocrinol Metab* **90**: 2131-2135.
Fernandez-Moreira D, Ugalde C, Smeets R, Rodenburg RJ, Lopez-Laso E, Ruiz-Falco ML,
Briones P, Martin MA, Smeitink JA, Arenas J. 2007. X-linked NDUFA1 gene
mutations associated with mitochondrial encephalomyopathy. *Ann Neurol* **61**: 73-83.
Fialkow PJ. 1970. X-chromosome inactivation and the Xg locus. *Am J Hum Genet* **22**: 460-
463.
Froyen G, Belet S, Martinez F, Santos-Reboucas CB, Declercq M, Verbeeck J, Donckers L,
Berland S, Mayo S, Rosello M et al. 2012. Copy-number gains of HUWE1 due to
replication- and recombination-based rearrangements. *Am J Hum Genet* **91**: 252-264.
Froyen G, Corbett M, Vandewalle J, Jarvela I, Lawrence O, Meldrum C, Bauters M,
Govaerts K, Vandeleur L, Van Esch H et al. 2008. Submicroscopic duplications of the
hydroxysteroid dehydrogenase HSD17B10 and the E3 ubiquitin ligase HUWE1 are
associated with mental retardation. *Am J Hum Genet* **82**: 432-443.
Gong X, Delorme R, Fauchereau F, Durand CM, Chaste P, Betancur C, Goubran-Botros H,
Nygren G, Anckarsater H, Rastam M et al. 2009. An investigation of ribosomal
protein L10 gene in autism spectrum disorders. *BMC Med Genet* **10**: 7.
Grasso CS, Wu YM, Robinson DR, Cao X, Dhanasekaran SM, Khan AP, Quist MJ, Jing X,
Lonigro RJ, Brenner JC et al. 2012. The mutational landscape of lethal castration-
resistant prostate cancer. *Nature* **487**: 239-243.
Gropman A, Samango-Sprouse CA. 2013. Neurocognitive variance and neurological
underpinnings of the X and Y chromosomal variations. *American journal of medical*
genetics Part C, Seminars in medical genetics **163C**: 35-43.
Gueneau L, Bertrand AT, Jais JP, Salih MA, Stojkovic T, Wehnert M, Hoeltzenbein M,
Spuler S, Saitoh S, Verschueren A et al. 2009. Mutations of the FHL1 gene cause
Emery-Dreifuss muscular dystrophy. *Am J Hum Genet* **85**: 338-353.
Guo A, Tan Y, Liu C, Zheng X. 2017. MST-4 and TRAF-6 expression in the peripheral
blood mononuclear cells of patients with Graves' disease and its significance. *BMC*
Endocr Disord **17**: 11.
Huynh KD, Lee JT. 2003. Inheritance of a pre-inactivated paternal X chromosome in early
mouse embryos. *Nature* **426**: 857-862.
Jiao S, Zhang Z, Li C, Huang M, Shi Z, Wang Y, Song X, Liu H, Li C, Chen M et al. 2015.
The kinase MST4 limits inflammatory responses through direct phosphorylation of
the adaptor TRAF6. *Nat Immunol* **16**: 246-257.
Johnston CM, Lovell FL, Leongamornlert DA, Stranger BE, Dermitzakis ET, Ross MT.
2008. Large-scale population study of human cell lines indicates that dosage
compensation is virtually complete. *PLoS Genet* **4**: e9.

Jones DT, Jager N, Kool M, Zichner T, Hutter B, Sultan M, Cho YJ, Pugh TJ, Hovestadt V,
Stutz AM et al. 2012. Dissecting the genomic complexity underlying
medulloblastoma. *Nature* **488**: 100-105.
Jonkers I, Monkhorst K, Rentmeester E, Grootegoed JA, Grosveld F, Gribnau J. 2008. Xist
RNA is confined to the nuclear territory of the silenced X chromosome throughout the
cell cycle. *Mol Cell Biol* **28**: 5583-5594.
Kere J, Srivastava AK, Montonen O, Zonana J, Thomas N, Ferguson B, Munoz F, Morgan D,
Clarke A, Baybayan P et al. 1996. X-linked anhidrotic (hypohidrotic) ectodermal
dysplasia is caused by mutation in a novel transmembrane protein. *Nature genetics*
13: 409-416.
Kremer EJ, Pritchard M, Lynch M, Yu S, Holman K, Baker E, Warren ST, Schlessinger D,
Sutherland GR, Richards RI. 1991. Mapping of DNA instability at the fragile X to a
trinucleotide repeat sequence p(CCG)n. *Science* **252**: 1711-1714.
Lederer D, Grisart B, Digilio MC, Benoit V, Crespin M, Ghariani SC, Maystadt I,
Dallapiccola B, Verellen-Dumoulin C. 2012. Deletion of KDM6A, a histone
demethylase interacting with MLL2, in three patients with Kabuki syndrome. *Am J*
Hum Genet **90**: 119-124.
Lee JT, Bartolomei MS. 2013. X-inactivation, imprinting, and long noncoding RNAs in
health and disease. *Cell* **152**: 1308-1323.
Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler
transform. *Bioinformatics* **26**: 589-595.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
R, Genome Project Data Processing S. 2009. The Sequence Alignment/Map format
and SAMtools. *Bioinformatics* **25**: 2078-2079.
Liewluck T, Hayashi YK, Ohsawa M, Kurokawa R, Fujita M, Noguchi S, Nonaka I, Nishino
I. 2007. Unfolded protein response and aggresome formation in hereditary reducing-
body myopathy. *Muscle Nerve* **35**: 322-326.
Lyon MF. 1961. Gene action in the X-chromosome of the mouse (Mus musculus L.). *Nature*
190: 372-373.
Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR,
Kamitaki N, Martersteck EM et al. 2015. Highly Parallel Genome-wide Expression
Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**: 1202-1214.
Mak W, Nesterova TB, de Napoles M, Appanah R, Yamanaka S, Otte AP, Brockdorff N.
2004. Reactivation of the paternal X chromosome in early mouse embryos. *Science*
303: 666-669.
Manara MC, Bernard G, Lollini PL, Nanni P, Zuntini M, Landuzzi L, Benini S, Lattanzi G,
Sciandra M, Serra M et al. 2006. CD99 acts as an oncosuppressor in osteosarcoma.
Mol Biol Cell **17**: 1910-1921.
Marco-Sola S, Sammeth M, Guigo R, Ribeca P. 2012. The GEM mapper: fast, accurate and
versatile alignment by filtration. *Nat Methods* doi:nmeth.2221 [pii]
10.1038/nmeth.2221.
Miyake N, Mizuno S, Okamoto N, Ohashi H, Shiina M, Ogata K, Tsurusaki Y, Nakashima
M, Saitsu H, Niikawa N et al. 2013. KDM6A point mutations cause Kabuki
syndrome. *Human mutation* **34**: 108-110.
Moortgat S, Desir J, Benoit V, Boulanger S, Pendeville H, Nassogne MC, Lederer D,
Maystadt I. 2016. Two novel EIF2S3 mutations associated with syndromic
intellectual disability with severe microcephaly, growth retardation, and epilepsy. *Am*
J Med Genet A **170**: 2927-2933.
Myant KB, Cammareri P, Hodder MC, Wills J, Von Kriegsheim A, Gyorffy B, Rashid M,
Polo S, Maspero E, Vaughan L et al. 2017. HUWE1 is a critical colonic tumour

suppressor gene that prevents MYC signalling, DNA damage accumulation and
tumour initiation. *EMBO Mol Med* **9**: 181-197.
Nawara M, Klapecki J, Borg K, Jurek M, Moreno S, Tryfon J, Bal J, Chelly J, Mazurczak T.
2008. Novel mutation of IL1RAPL1 gene in a nonspecific X-linked mental
retardation (MRX) family. *Am J Med Genet A* **146A**: 3167-3172.
Okamoto I, Arnaud D, Le Baccon P, Otte AP, Disteche CM, Avner P, Heard E. 2005.
Evidence for de novo imprinted X-chromosome inactivation independent of meiotic
inactivation in mice. *Nature* **438**: 369-373.
Okamoto I, Otte AP, Allis CD, Reinberg D, Heard E. 2004. Epigenetic dynamics of
imprinted X inactivation during early mouse development. *Science* **303**: 644-649.
Park C, Carrel L, Makova KD. 2010. Strong purifying selection at genes escaping X
chromosome inactivation. *Molecular biology and evolution* **27**: 2446-2450.
Patrat C, Okamoto I, Diabangouaya P, Vialon V, Le Baccon P, Chow J, Heard E. 2009.
Dynamic changes in paternal X-chromosome activity during imprinted X-
chromosome inactivation in mice. *Proc Natl Acad Sci U S A* **106**: 5198-5203.
Penny GD, Kay GF, Sheardown SA, Rastan S, Brockdorff N. 1996. Requirement for Xist in
X chromosome inactivation. *Nature* **379**: 131-137.
Petropoulos S, Edsgard D, Reinius B, Deng Q, Panula SP, Codeluppi S, Reyes AP,
Linnarsson S, Sandberg R, Lanner F. 2016. Single-Cell RNA-Seq Reveals Lineage
and X Chromosome Dynamics in Human Preimplantation Embryos. *Cell* **167**: 285.
Potluri P, Davila A, Ruiz-Pesini E, Mishmar D, O'Hearn S, Hancock S, Simon M, Scheffler
IE, Wallace DC, Procaccio V. 2009. A novel NDUFA1 mutation leads to a
progressive mitochondrial complex I-specific neurodegenerative disease. *Mol Genet*
Metab **96**: 189-195.
Prothero KE, Stahl JM, Carrel L. 2009. Dosage compensation and gene expression on the
mammalian X chromosome: one plus one does not always equal two. *Chromosome*
Res **17**: 637-648.
Quinzii CM, Vu TH, Min KC, Tanji K, Barral S, Grewal RP, Kattah A, Camano P, Otaegui
D, Kunimatsu T et al. 2008. X-linked dominant scapuloperoneal myopathy is due to a
mutation in the gene encoding four-and-a-half-LIM protein 1. *Am J Hum Genet* **82**:
208-213.
Reinius B, Mold JE, Ramskold D, Deng Q, Johnsson P, Michaelsson J, Frisen J, Sandberg R.
2016. Analysis of allelic expression patterns in clonal somatic cells by single-cell
RNA-seq. *Nature genetics* **48**: 1430-1435.
Santoni FA, Stamoulis G, Garieri M, Falconnet E, Ribaux P, Borel C, Antonarakis SE. 2017.
Detection of Imprinted Genes by Single-Cell Allele-Specific Gene Expression. *Am J*
Hum Genet **100**: 444-453.
Santos A, Wernersson R, Jensen LJ. 2015. Cyclebase 3.0: a multi-organism database on cell-
cycle regulation and phenotypes. *Nucleic Acids Res* **43**: D1140-1144.
Schessl J, Zou Y, McGrath MJ, Cowling BS, Maiti B, Chin SS, Sewry C, Battini R, Hu Y,
Cottle DL et al. 2008. Proteomic identification of FHL1 as the protein mutated in
human reducing body myopathy. *J Clin Invest* **118**: 904-912.
Schneider-Gadicke A, Beer-Romero P, Brown LG, Nussbaum R, Page DC. 1989. ZFX has a
gene structure similar to ZFY, the putative human sex determinant, and escapes X
inactivation. *Cell* **57**: 1247-1258.
Schonk DM, Kuijpers HJ, van Drunen E, van Dalen CH, Geurts van Kessel AH, Verheijen R,
Ramaekers FC. 1989. Assignment of the gene(s) involved in the expression of the
proliferation-related Ki-67 antigen to human chromosome 10. *Human genetics* **83**:
297-299.

Scialdone A, Natarajan KN, Saraiva LR, Proserpio V, Teichmann SA, Stegle O, Marioni JC, Buettner F. 2015. Computational assignment of cell-cycle stage from single-cell transcriptome data. *Methods* **85**: 54-61.

Shalaby S, Hayashi YK, Nonaka I, Noguchi S, Nishino I. 2009. Novel FHL1 mutations in fatal and benign reducing body myopathy. *Neurology* **72**: 375-376.

Sharman GB. 1971. Late DNA replication in the paternally derived X chromosome of female kangaroos. *Nature* **230**: 231-232.

Sharp AJ, Stathaki E, Migliavacca E, Brahmachary M, Montgomery SB, Dupre Y, Antonarakis SE. 2011. DNA methylation profiles of human active and inactive X chromosomes. *Genome research* **21**: 1592-1600.

Shashi V, Berry MN, Shoaf S, Sciote JJ, Goldstein D, Hart TC. 2000. A unique form of mental retardation with a distinctive phenotype maps to Xq26-q27. *Am J Hum Genet* **66**: 469-479.

Shashi V, Xie P, Schoch K, Goldstein DB, Howard TD, Berry MN, Schwartz CE, Cronin K, Sliwa S, Allen A et al. 2015. The RBMX gene as a candidate for the Shashi X-linked intellectual disability syndrome. *Clin Genet* **88**: 386-390.

Sobecki M, Mrouj K, Colinge J, Gerbe F, Jay P, Krasinska L, Dulic V, Fisher D. 2017. Cell-Cycle Regulation Accounts for Variability in Ki-67 Expression Levels. *Cancer Res* **77**: 2722-2734.

Stegle O, Teichmann SA, Marioni JC. 2015. Computational and analytical challenges in single-cell transcriptomics. *Nature reviews Genetics* **16**: 133-145.

Sudbrak R, Wieczorek G, Nuber UA, Mann W, Kirchner R, Erdogan F, Brown CJ, Wohrle D, Sterk P, Kalscheuer VM et al. 2001. X chromosome-specific cDNA arrays: identification of genes that escape from X-inactivation and other applications. *Hum Mol Genet* **10**: 77-83.

Tabolacci E, Pomponi MG, Pietrobono R, Terracciano A, Chiurazzi P, Neri G. 2006. A truncating mutation in the IL1RAPL1 gene is responsible for X-linked mental retardation in the MRX21 family. *Am J Med Genet A* **140**: 482-487.

Tao R, Jin B, Guo SZ, Qing W, Feng GY, Brooks DG, Liu L, Xu J, Li T, Yan Y et al. 2006. A novel missense mutation of the EDA gene in a Mongolian family with congenital hypodontia. *J Hum Genet* **51**: 498-502.

Tarpey P, Pemberton TJ, Stockton DW, Das P, Ninis V, Edkins S, Andrew Futreal P, Wooster R, Kamath S, Nayak R et al. 2007. A novel Gln358Glu mutation in ectodysplasin A associated with X-linked dominant incisor hypodontia. *Am J Med Genet A* **143**: 390-394.

Thevenon J, Michot C, Bole C, Nitschke P, Nizon M, Faivre L, Munnich A, Lyonnet S, Bonnefont JP, Portes VD et al. 2015. RPL10 mutation segregating in a family with X-linked syndromic Intellectual Disability. *Am J Med Genet A* **167A**: 1908-1912.

Tukiainen T, Villani AC, Yen A, Rivas MA, Marshall JL, Satija R, Aguirre M, Gauthier L, Fleharty M, Kirby A et al. 2017. Landscape of X chromosome inactivation across human tissues. *Nature* **550**: 244-248.

van Haaften G, Dalgliesh GL, Davies H, Chen L, Bignell G, Greenman C, Edkins S, Hardy C, O'Meara S, Teague J et al. 2009. Somatic mutations of the histone H3K27 demethylase gene UTX in human cancer. *Nature genetics* **41**: 521-523.

Visinoni AF, de Souza RL, Freire-Maia N, Gollop TR, Chautard-Freire-Maia EA. 2003. X-linked hypohidrotic ectodermal dysplasia mutations in Brazilian families. *Am J Med Genet A* **122A**: 51-55.

Wang J, Syrett CM, Kramer MC, Basu A, Atchison ML, Anguera MC. 2016. Unusual maintenance of X chromosome inactivation predisposes female lymphocytes for increased expression from the inactive X. *Proc Natl Acad Sci U S A* **113**: E2029-2038.

Wang K, Li M, Hakonarson H. 2010. ANNOVAR: functional annotation of genetic variants
from high-throughput sequencing data. *Nucleic Acids Res* **38**: e164.
Yang F, Babak T, Shendure J, Disteche CM. 2010. Global survey of escape from X
inactivation by RNA-sequencing in mouse. *Genome research* **20**: 614-622.
Yasukochi Y, Maruyama O, Mahajan MC, Padden C, Euskirchen GM, Schulz V, Hirakawa
H, Kuhara S, Pan XH, Newburger PE et al. 2010. X chromosome-wide analyses of
genomic DNA methylation states and gene expression in male and female neutrophils.
Proc Natl Acad Sci U S A **107**: 3704-3709.
Yotsumoto S, Fukumaru S, Matsushita S, Oku T, Kobayashi K, Saheki T, Kanzaki T. 1998.
A novel point mutation of the EDA gene in a Japanese family with anhidrotic
ectodermal dysplasia. *J Invest Dermatol* **111**: 1246-1247.
Zhang Y, Castillo-Morales A, Jiang M, Zhu Y, Hu L, Urrutia AO, Kong X, Hurst LD. 2013.
Genes that escape X-inactivation in humans have high intraspecific variability in
expression, are associated with mental impairment but are not slow evolving.
Molecular biology and evolution **30**: 2588-2601.
810

A

ind 1　ind 2　ind 3　ind 4　ind 5

229 SC　160 SC　192 SC　192 SC　162 SC　48 SC

WGS
Identification of heterozygous sites for each individual

scRNA-seq
983 SC

QC filtering 951 SC

Removal of double cells 869 SC

X chromosome ASE single cell analysis

Study of cellular dynamics of XCI

B

correlation
−1　0　1

CELLS_ individual 3

CELLS_ individual 3

CELLS_ individual 3

CELLS_ individual 3

LCL 5

ind. 1 ind. 2 ind. 3 ind. 4 ind. 5

PAR

AR: [0-0.95], escapee
AR: [0.95-1], inactivated

cells