Metagenomic sequencing provides insights into the location of

microbial detoxification in the gut of a small mammalian herbivore

Running title: *Determining the site of microbial detoxification*

Kevin D. Kohl[1,2], Kelly F. Oakeson[2], Teri J. Orr[2], Aaron W. Miller[2,3], Jennifer Sorensen

Forbey[4], Caleb D. Phillips[5], Colin Dale[2], Robert B. Weiss[6], M. Denise Dearing[2]

[1]Department of Biological Sciences, University of Pittsburgh, 4249 Fifth Ave.,
Pittsburgh, PA, 15260, USA
[2] Department of Biology, University of Utah, 257 South 1400 East, Salt Lake City, UT,
84112, USA
[3] Departments of Urology and Immunology, Cleveland Clinic Foundation, 9500 Euclid
Ave, Cleveland, OH 44195
[4] Department of Biological Sciences, Boise State University, 1910 University Drive,
Boise, Idaho 83725 USA
[5] Department of Biological Sciences, Texas Tech University, 2901 Main Street, Lubbock,
TX, 79409
[6] Department of Human Genetics, University of Utah, 15 North 2030 East, Salt Lake
City, UT, 84112, USA

Corresponding Author:

Kevin Kohl – University of Pittsburgh

Kevin.d.kohl@gmail.com

Abstract – Microbial detoxification of plant defense compounds influences the use of certain plants as food sources by herbivores. The location of microbial detoxification along the gut could have profound influences on the distribution, metabolism, and tolerance to toxic compounds. Stephen's woodrats (*Neotoma stephensi*) specialize on juniper, which is heavily defended by numerous defensive compounds, such as oxalate, phenolics, and monoterpenes. Woodrats maintain two gut chambers harboring dense microbial communities: a foregut chamber proximal to the major site of toxin absorption, and a cecal chamber in their hindgut. First, we conducted a trial with antibiotics to determine whether the gut microbiota is important in allowing *N. stephensi* to consume juniper. Antibiotic-treated animals lost significantly more body mass while feeding on juniper, relative to those feeding on rabbit chow. Next, we measured levels of toxins throughout the gut. Oxalate concentrations were immediately lower in the foregut chamber, while concentrations of terpenes remain high in the foregut, and are lowest in the cecal chamber. Finally, we conducted metagenomic sequencing of the foregut and cecal chambers. We found that the majority of genes associated with detoxification functions were more abundant in the cecal chamber. However, some genes associated with degradation of oxalate and phenolic compounds were more abundant in the foregut. Thus, it seems that microbial detoxification may take place in various chambers depending on the class of chemical compound. We hypothesize that the location of microbial detoxification could impact the tolerance of animals to these compounds, which may have ecological and evolutionary consequences.

Importance – For decades, ecologists have investigated how partnerships with

fermentative microbes have contributed to the success of mammalian herbivores.

The location of these partnerships in the gut (foregut vs. hindgut fermentation) can

have large implications for host ecology, such as limiting body size or dietary niche

breadth. Mammalian herbivores also host microbial communities that aid in the

detoxification of plant defensive compounds, though the location of these functions

in the gut has not been studied thoroughly. Here, we used metagenomic sequencing

to investigate the location of detoxification functions in the gut of a small

mammalian herbivore, Stephen's woodrat, which has both a foregut and a hindgut

chamber. We found that most detoxification genes exhibited higher abundance in

the hindgut, though genes associated with oxalate degradation were enriched in the

foregut. We hypothesize that similar to microbial fermentation, the location of

microbial detoxification may also have implications for host ecology.

INTRODUCTION

Mammalian herbivores are often challenged by the consumption of toxic plant secondary compounds (PSCs) that negatively impact their physiology in various ways (1). In response, mammalian herbivores have evolved myriad adaptations to cope with dietary toxins, such as altered behavior, efflux pumps in the gut, and enhanced liver detoxification (1). Recently, we demonstrated that the gut microbiota of herbivores also play an important role in allowing them to consume PSCs (2). However, it is still unclear where the process of microbial detoxification occurs in the gut.

It has been proposed that microbial detoxification should occur early in the gastrointestinal tract, to facilitate rapid degradation of PSCs and minimize the potential for their absorption in the digestive tract. In fact, it has been hypothesized that the rumen evolved first for detoxification and was later adopted for processes of fermentation (3, 4). Examples of animals in which foregut detoxification has been documented for particular PSCs include reindeer (*Rangifer tarandus*; (5) that have rumen communities that degrade usnic acid, a toxin produced by lichen, and hoatzins (*Opisthocomus hoazin*), that maintain communities of microbes in their foregut capable of metabolizing saponins (6). There is also evidence of microbial detoxification processes occurring in the hindguts of some animals. For example, Sage-grouse (*Centrocercus urophasianus*), specialize on toxic sagebrush (*Artemsia* spp.) and maintain a cecal microbiota enriched in detoxification genes (7). Additionally, the hindguts of koalas (*Phascolarctos cinereus*) harbor microbes capable of degrading phenolic PSCs (8). Understanding the location of microbial

processes in the gut is likely to be important in the context of animal ecology. For

example, the location of fermentation (foregut vs hindgut), seems to facilitate or

constrain evolution of herbivore body size (9) and dietary niche breadth (10). The

location of microbial detoxification could have similar impacts on animal ecology.

In contrast to many above described examples, woodrats (genus *Neotoma*)

are particularly interesting from a digestive standpoint as these herbivorous

rodents possess both a foregut chamber and a large cecal chamber in their hindgut.

Both the foregut and cecal chambers harbor dense communities of microbes and

high concentrations of volatile fatty acids (11), which are the microbial products of

fermentation. Following the hypothesis that the bovine rumen first evolved for

detoxification regarding the rumen (3, 4), we predict that microbial detoxification of

PSCs might occur prior to the small intestine in the woodrat gut such that the animal

experiences limited absorption and systemic exposure to toxic compounds.

To evaluate this hypothesis, we conducted a series of experiments focusing

on a species of woodrat with a highly specialized diet. Stephen's woodrat (*Neotoma*

stephensi) is a dietary specialist, with juniper (*Juniperus monosperma*) composing

~90% of its diet (12). Juniper defends itself from herbivory with various toxic

compounds, including monoterpenes (13), phenolics (14, 15), and oxalate (16). We

have demonstrated that microbes aid in the detoxification of phenolics (2) and

oxalate in other woodrat species (17, 18). For example, some populations of the

white-throated woodrat (*N. albigula*) specialize on cactus (*Opuntia* spp.), which is

rich in oxalate, but lacking in other defensive compounds. The foregut microbial

communities of these *N. albigula* contain several species of oxalate-degrading

microbes (19). There has also been metagenomic evidence obtained from other

systems suggesting that microbes aid in the detoxification of terpenes (mountain

pine beetles: (20); sage grouse: (7), as well as experimental work demonstrating the

detoxification function of the gut microbiota of pine weevils (21). However, the

detoxification contribution of the microbiome has not been compared between the

foregut and hindgut in any wild vertebrate herbivore.

We addressed several questions within the woodrat-juniper system. First, to

determine if the gut microbiota plays a role in facilitating the consumption of

juniper we conducted a feeding trial with *N. stephensi* where we disrupted the

microbial community with antibiotics. We predicted that antibiotic treatments

would significantly impair the ability for *N. stephensi* to feed on juniper. Next, we

measured concentrations of two classes of toxins (oxalate and monoterpenes)

across regions of the gut. We predicted that if detoxification is occurring proximally

in the gut, then toxins concentrations in the foregut would be significantly lower

compared to food material. Last, we conducted a series of metagenomic analyses to

inventory and compare the functions of microbial communities. We first compared

the metagenomes of foregut communities between the juniper specialist *N. stephensi*

and the cactus specialist *N. albigula* and predicted that if the foregut microbiota of *N.*

stephensi were specialized for detoxification of juniper, which contains oxalate,

monoterpenes, and phenolics, we would observe enrichment of detoxification genes

compared to the foregut of *N. albigula* that is exposed primarily to the oxalate in

cactus. We also compared the metagenomes of foregut and cecal chambers within *N.*

stephensi and predicted that the metagenome of the foregut chamber would be

enriched in detoxification genes compared to the cecal communities.

RESULTS

Antibiotic feeding trial

Treatment with antibiotics significantly impaired the ability of *N. stephensi* to

feed on juniper-rich diets. Animals feeding on juniper exhibited a 9.5% decrease in

food intake when placed on antibiotics, while those feeding on a control diet only

decreased food intake by 2.4% (Fig. 1; Kruskal-Wallis: P = 0.03). This difference in

food intake corresponded to a difference in maintenance of body mass, such that

animals being fed a juniper diet lost roughly 6.3% of their body mass when placed

on antibiotics, while those fed the control diet only lost 1.5% of body mass (Fig. 1;

Kruskal-Wallis: P = 0.03).

Concentrations of PSCs across the gut

Water concentration varied over the length of the gut, with a significant

reduction in the large intestine (Fig. 2). Oxalate concentrations in the entire gut

were significantly less than in the food material (Fig 2; P < 0.0001). Oxalate

concentrations also varied significantly across gut regions (P < 0.001), and were

highest in the foregut and lowest in the stomach. Conversely, foregut concentrations

of monoterpenes were not significantly lower than the food material (Fig. 2; P > 0.05

for all). Concentrations of monoterpenes did vary across gut regions, with the

lowest concentrations always being in the cecum (Fig. 2; P < 0.0001 for all). In fact,

camphene and β-pinene were not detectable in any cecal samples.

Metagenomic sequencing

We obtained an average of 3.35 million reads per sample (± 517 thousand).

We first compared the bacterial metagenomes of foregut chambers from *N. stephensi*

(which specializes on juniper) and *N. albigula* (which specializes on cactus). There

were no functions that were significantly enriched in the foregut bacterial

metagenomes of *N. stephensi*. The foregut bacterial communities of *N. albigula* were

enriched in functions related to the metabolism of several amino acids and vitamin

B6 (Fig. 3). Further, when investigating detoxification genes, there were no

detectable differences between the two species.

We next searched for differential functions between the foregut and cecal

chambers of *N. stephensi*. The foregut bacterial metagenome was enriched in genes

associated with the metabolism of nucleotides (purines), amino acids (glutathione,

thiamine), vitamins (riboflavin, nicotinate, folate), and lipoic acid (Fig. 4A). The cecal

bacterial metagenome was enriched for genes associated with the metabolism of

starch and sucrose, as well as the biosynthesis of several essential amino acids (Fig.

4A). The cecum was also enriched in several functional categories related to

detoxification (drug metabolism, chloroalkane and chloroalkene degradation; Fig.

4A).

We also specifically investigated genes associated with "Xenobiotic

degradation and metabolism", "Geraniol degradation", and "Limonene and pinene

degradation". Here, the only gene enriched in the foregut was 4-oxalocrotonate

tautomerase, which was 24× more abundant in the foregut compared to the cecum

(Fig. 4B). There were 5 other detoxification genes that were significantly more

abundant in the cecum compared to the foregut chamber (Fig. 4B). Three of these

genes (carboxylesterase 2, hydroxyatrazine ethylaminohydrolase, and 2-

hydroxycyclohexanecarboxyl-CoA dehydrogenase) were 2-3× more abundant in the

cecal chambers compared to the foregut chambers. Two genes, anthraniloyl-CoA

monooxygenase and dehalogenases, were not detected in the foregut chambers, but

were present in the cecal chambers of the juniper-feeding *N. stephensi* (Fig. 4B).

Last, we focused on a subset of detoxification enzymes highlighted from

previous studies. The gene oxalyl-CoA decarboxylase, which is anticipated to

function in oxalate degradation, did not exhibit differential abundance between the

foregut chambers of *N. stephensi* and *N. albigula*. However, the abundance of this

gene was 4.5× higher in the foregut of *N. stephensi* compared to the cecum (Fig. 5).

Similarly, aryl-alcohol dehydrogenase, which is more abundant in the woodrat

foregut when animals are fed phenolic-rich creosote resin (2), did not exhibit

significant differences between the foregut chambers of *N. stephensi* or *N. albigula*,

but was 41× higher in the *N. stephensi* foregut compared to the cecum (Fig. 5). Last,

we investigated several genes in the "Limonene and pinene degradation" pathway,

which are enriched in other herbivores that feed on terpene-rich plants (7, 20, 21).

Enoyl-CoA hydratase (E.C. 4.2.1.17) was more abundant in the foregut of *N.*

stephensi compared to *N. albigula*, with no difference between the foregut and cecal

chambers of *N. stephensi* (Fig. 5). Additionally, aldehyde dehydrogenase (E.C.

1.2.1.3) exhibited no difference between the foregut chambers of *N. stephensi* and *N.*

albigula, but was 2.5× more abundant in the cecal chambers of *N. stephensi* (Fig. 5)*.*

We found a significant positive relationship between the relative abundances

of functional categories as estimated by PICRUSt and determined by metagenomic

sequencing (Fig. 6; foregut: R^2 = 0.49, slope = 0.922 ± 0.074, P < 0.0001; cecum: R^2 =

0.51, slope = 0.978 ± 0.072, P < 0.0001). Some functional categories, such as those

representing 'Peptidases' and 'Amino acid Related Enzymes', were not well

predicted by PICRUSt, in that they were present on the x-axis (predicted to be

present by PICRUSt but not detected by metagenomic sequencing). However,

categories that exhibited the greatest fold-differences between the two chambers

were also predicted to differ by PICRUSt (Fig. 6; with the exception of Starch and

Sucrose Metabolism). Following assignment into functional community categories it

was determined that benzoate degradation abundance and the composition of its

contributing microbial lineages was significantly conserved (FDR adjusted p-value =

0.048) between the two chambers. Further investigations revealed that conserved

benzoate degradation function across chambers was contributed by a combination

of 28 different bacterial OTUs classified into eight genera (*Supplementary material).*

DISCUSSION

Here, we investigated the nature and location of microbial detoxification in

the guts of Stephen's woodrat, a juniper specialist. We found that microbes are

instrumental in allowing *N. stephensi* to feed on a diet of juniper. The location of

detoxification within the gut seems to vary depending on the chemical class of

defensive compounds. We discuss our findings below, as well as the potential

ecological and evolutionary consequences of foregut vs hindgut detoxification by the

microbiome.

First, we demonstrated that antibiotic treatment significantly impaired the

ability of *N. stephensi* to maintain body mass when feeding on juniper diets. These

results are consistent with studies demonstrating that reduction in the microbiota

impairs the ability of another woodrat species, *N. lepida*, to consume phenolic-rich

creosote resin (2), and the ability of pine weevils to thrive on terpene-rich diets

(21). It should be noted that antibiotic treatment can also have effects on host

physiology and gene expression (22). However, here we compared changes in food

intake and body mass in response to antibiotics of animals being fed either control

diets or juniper diets. Thus, both groups of animals were treated with antibiotics

and experienced the same effects due to the antibiotic treatment, but those feeding

on juniper exhibited greater declines in food intake and body mass. Based on these

results, it is clear that the gut microbiota plays an important role in detoxification

when juniper is consumed by *N. stephensi*.

Next, we compared concentrations of PSCs along the length of the

gastrointestinal tract. We predicted that if toxins were metabolized early in the gut,

their concentrations in the foregut chamber would be lower than those measured in

food material, similar to what has been observed for usnic acid in the reindeer

rumen (5). This was observed in the case of oxalate, where foregut concentrations

were roughly a third of those measured in the diet. Since oxalate is detected only at

very low levels in the urine of woodrats (16) and mammals do not produce the

necessary oxalate-degrading enzymes (23), it seems that oxalate is detoxified

efficiently by microbes in the foregut. Conversely, terpene concentrations in the

foregut were the same as those measured in food material, suggesting that the

foregut is not a significant site for detoxification. Instead, concentrations of terpenes

were lowest in the cecum, where some terpene compounds were not even detected.

These results are similar to those observed in sage-grouse, where terpene

concentrations are also lowest in the cecal chambers (24). *Neotoma stephensi* has

physiological adaptations that reduce terpene absorption in the gut (25), perhaps

allowing terpene metabolism to be delayed until the food reaches the cecum. In the

case of terpenes, concentrations increased in the large intestine compared to the

cecum. While this change seems puzzling, we hypothesize that the selective

absorption of water in this gut region results in higher concentrations of toxins.

Further experiments, perhaps using radiolabeled compounds, are needed to

understand the routes of absorption and detoxification of these compounds in

different regions of the woodrat digestive system.

Interestingly, we did not observe many differences related to detoxification

between the foregut microbial metagenomes of *N. stephensi* and *N. albigula*, even

though these two host species feed on different diets with contrasting chemical

profiles. It could be that the foregut microbiome offers broad detoxification

capacities, while cecal microbial communities are more specialized. That being said,

we did observe a higher abundance of enoyl-CoA hydratase in the foregut of *N.*

stephensi compared to *N. albigula*. This gene plays a role in the ability for microbes

to degrade terpenes (26), and is enriched in the gut metagenomes of sage-grouse (7)

and mountain pine beetles (20). Thus, there does seem to be some functional

enrichment for detoxification in the foregut of *N. stephensi*.

Within *N. stephensi*, the foregut and cecal microbial communities appear to

provide different detoxification functions. In the foregut, we observed higher

abundance of the gene 4-oxalocrotonate tautomerase, which is involved in the

degradation of aromatic hydrocarbons by bacteria (27). The foregut also had higher

abundances of the gene oxalyl-CoA decarboxylase, which degrades oxalate, and aryl-

alcohol dehydrogenase, which degrades aromatic hydrocarbons. The abundance of

aryl-alcohol dehydrogenase increases in the foregut of another woodrat species (*N.*

lepida) when animals are fed diets containing phenolic PSCs (2) and thus may be an

important enzyme that is provided by the foregut microbial community .

We also obtained evidence for enrichment of detoxification enzymes in the

cecal chamber. Higher-level gene categories, such as 'Drug Metabolism' and

'Chloroalkane and Chloroalkene Degradation' were more abundant in the cecal

chambers than the foregut. Specifically, the cecum was enriched for genes such as

hydroxyatrazine ethylaminohydrolase, which deaminates or dechlorinates triazine

compounds (28, 29), and anthraniloyl-CoA monooxygenase, which plays a role in

the catabolism of aromatic compounds (30). Additionally, the aldehyde

dehydrogenase gene was significantly more abundant in the cecum. This gene is

annotated as a component of the 'Limonene and Pinene Degradation' pathway, and

is found to be enriched in the gut metagenomes of sage-grouse (7) and mountain

pine beetles (20).

We found that the frequencies of genes encoding enzymes that are predicted

to function in benzoate degradation was conserved between gut chambers.

Benzoate degradation is hierarchically classified into xenobiotic degradation and

metabolism pathways. Benzoate degradation was classified as a conserved function

between both chambers due to both the similar frequency of this function in both

chambers as well as the observation that the bacterial lineages inferred to

contribute this function were found to have similar compositional profiles in both

chambers. Some of the eight genera observed in this group are well-known

ruminant symbionts (e.g. *Oxalobacter*), acidophiles (e.g. *Alicyclobacillus*), common

gut lineages (e.g. *Pseudomonas*), and chemoheterotrophs (*Sphingomonas)*. This

result suggests that overlapping components of microbiome communities in both

foregut and cecum are selected on the basis of their ability to contribute benzoate

degradation functions to the community.

Finally, we tested whether predictive profiling of metagenomics based on

16S rRNA sequences is appropriate for woodrat gut samples. We found a significant

relationship between the relative abundances of functional categories as estimated

by PICRUSt and determined by metagenomic sequencing. Additionally, results for

PICRUSt and metagenomics were consistent for some of the categories that

exhibited the largest differences between the two chambers. However, PICRUSt did

not predict all of the categories accurately. Many functional categories (e.g.

'Peptidases' and "Amino acid Related Enzymes') were predicted to be present by

PICRUSt analysis, but were not detected by metagenomic sequencing. Interestingly,

the gut microbial communities of herbivores tend to have lower abundances of

genes encoding protease relative to carnivores, likely to conserve these nutrients on

a low protein diet (31). Thus, the use of predictive metagenomic profiling may not

always be feasible for non-model systems, especially given that most microbial

genomes used for predictive profiling are from the human gut. Indeed, PICRUSt

works significantly better for human samples than for samples from nonhuman

mammalian guts (32). The sequencing of additional genomes from nonhuman

mammals would help to improve the accuracy of programs like PICRUSt in non-

model systems. It seems that results from predicting profiling in non-model systems

should perhaps be verified with other methods, such as quantitative PCR.

Overall, we hypothesized that microbial detoxification should occur

proximally in the gastrointestinal tract, so as to facilitate degradation of PSCs before

they would be absorbed later in the gut. However, it seems that detoxification

processes may be spatially structured along the length of the gut. The evidence for

foregut detoxification is strongest for oxalate, where abundances of the

detoxification enzyme (oxalyl-CoA decarboxylase) were high, and concentrations of

oxalate were already much lower than in food material, and abundances of

potentially oxalate-degrading bacteria are highest (19). Terpene degradation may

occur in the cecal chamber, where abundances of terpene-degradation genes were

higher, and concentrations of terpenes were at their lowest. It is somewhat puzzling

why this spatial arrangement would exist, given that terpenes are absorbed very

rapidly due to their small size and high lipophilicity (25). However, *N. stephensi*

exhibits lower terpene absorption than other woodrat species (25), resulting in

relatively higher terpene exposure and resultant detoxification by microbial

communities in the hindgut. Laboratory rats are able to excrete ~25% of absorbed

monoterpenes back into the gut lumen through the biliary duct (33), an excretion

route that may be enhanced in a specialist herbivore such as *N. stephensi*. This

excretion route may facilitate further microbial degradation of terpenes in the

hindgut. While we did not measure concentrations of phenolics through the gut,

various genes associated with degrading aromatic hydrocarbons were enriched in

both the foregut and cecum. Phenolics can bind to dietary protein and digestive

enzymes, thus limiting nutrient availability (34). Therefore, foregut detoxification of

phenolics would be beneficial for avoiding the anti-nutritive effects of PSCs. Again,

labeled chemicals to trace sites of absorption, excretion, and metabolism would

enhance our understanding of the location of these processes.

The location of detoxification within the gut could have important ecological

and evolutionary implications for plants and herbivores. For example, when

comparing several species of rodents, only those with foregut chambers were able

to metabolize significant amounts of oxalate (35). Thus, evolution of this chamber

may be important for the ability to consume oxalate-rich plants. Consumption of

plant toxins might also drive changes in gut anatomy to enhance detoxification by

facilitating a longer duration of residency in various gut chambers. For example,

woodrats feeding on phenolic-rich oak exhibit an increase in cecal size (36), which

could be driven by increased fiber content of oak or to facilitate the detoxification of

PSCs. Interestingly, supplementing diets with isolated phenolic compounds causes

an increase in cecal size in lab rats (37). Alterations of the size or function of gut

chambers in response to toxins could have implications for dietary niche breadth or

energetics. In herbivorous birds, there is evidence suggesting that microbial

detoxification occurs in both foregut (6) and hindgut chambers (7). The guts of birds

are under strong selective pressure to be small so as to facilitate flight (38), and thus

sites of microbial detoxification may be especially important in these species.

Studies on additional and phylogenetically diverse herbivores and classes of PSCs

may help us to better understand the importance of microbial detoxification in the

gut, and how the spatial arrangement of these processes contribute to animal

morphology, performance, ecology, and evolution.

MATERIALS AND METHODS

Antibiotic Feeding Trial

We conducted a feeding trial coupled with antibiotic treatment to investigate

whether microbial communities contribute to the tolerance of a juniper-based diet.

Individuals of *N. stephensi*, as well as juniper foliage for diet experiments, were

collected from near Wupatki National Monument, AZ, USA (35°30' N, 111°27' W).

Animals were brought to the University of Utah animal facility and fed a diet of 50%

ground juniper and 50% rabbit chow for 3 days. Juniper diets (*J. monosperma*) were

prepared as previously described (25). Animals were then divided into either a

'Control Diet' group or a 'Juniper Diet' group, with four individuals in each group.

Animals in the Control diet group were transitioned to a juniper free diet by feeding

them a diet of 25% juniper/75% rabbit chow for 3 days, followed by 100% rabbit

chow for 6 days. Animals in the 'Juniper Diet' group were transitioned to an

increasing amount of juniper by feeding them a diet of 75% juniper/25% rabbit

chow for 3 days and then 90% juniper/10% rabbit chow for 6 days.

For the duration of the experiment, animals were given water containing

sucralose (2.5% w/v) to facilitate drinking. During the last 3 days of the experiment,

an antibiotic (0.5 g neomycin L^{-1}) was also added to the water of all animals.

Neomycin is a broad spectrum antibiotic that is poorly absorbed across the gut

tissue and reduces gut bacterial density by ~90% (39). This treatment has

effectively been used to demonstrate that gut microbes are critical for the ability of

another woodrat species, *N. lepida*, to consume phenolic-rich creosote resin (2).

Body mass was measured daily over the course of the experiment. The mass

of wet food presented was measured daily, and subsamples were used to estimate

the dry weight of the diet. On the following day, leftover food was collected, dried

overnight at 40°C and weighed to measure intake. Food intake was calculated as the

dry mass of food presented – dry mass of leftover food recovered.

We calculated the percent change in food intake and body mass between the

day prior to the beginning of antibiotic treatment to the end of the feeding trial.

These percentages were compared between the two groups using t-tests. This

feeding trial and all experiments below were approved by the University of Utah

Institutional Animal Care and Use Committee (IACUC) under protocols #12-12010

and #16-02011.

Toxin Levels Throughout the Gut

We conducted a feeding trial on *N. stephensi* to document changes in toxin

concentrations across gut regions. Six individuals of *N. stephensi* were fed increasing

concentrations of *J. monosperma* (3 days on 25%, 3 days on 50%, 4 days on 75%).

On the last morning, animals were euthanized using isoflurane. Contents of each gut

region (foregut, stomach, small intestine, cecum, large intestine, feces) were

collected. A small portion of wet gut contents were placed in airtight headspace

autosampler vials and frozen at -20°C until analysis. Concentrations of individual

monoterpenes were determined using headspace gas chromatography with an

Agilent 7694 headspace sampler coupled with an Agilent 6890N gas chromatograph

(GC). One mL of headspace gas was injected into a J&W DB-5 capillary column (30

m × 250 μm × 0.25 μm). Operating conditions for the headspace sampler were: oven

temperature at 100°C, loop temperature at 110°C, transfer line temperature at

120°C, a vial equilibrium time of 20 min, a pressurization time of 0.20 min, a loop fill

time of 0.50 min, a loop equilibrium time of 0.20 min, and an injection time of 0.50

min. Operating conditions for the GC were: splitless injector at 250°C, flame

ionization detector at 300°C, oven temperature at 40°C for 2 min, then increasing

3°C/min to 60°C, then increasing 5°C/min to 120°C, then increasing 20°C/min to

300°C, and held at 300°C for 7 min. The make-up gas was nitrogen and the carrier

gas was helium. The inlet pressure was 80 kPa with a flow rate of 1.0 mL/min.

Retention times and peak areas of individual monoterpenes were calculated using

Agilent OpenLab version A.01.05. We verified the relative retention times of

individual monoterpenes using standards, and estimated the relative concentrations

of individual monoterpenes in the intestinal content material by comparing AUC of

peaks in diet or gut samples to AUC of a known volume of 10 mg/mL monoterpene

standards.

Remaining gut contents were placed in weighing boats and dried at 40°C for

two nights. The dried gut contents were then used to measure oxalate

concentrations using a previously established protocol (40). Roughly 0.4g of dried

gut contents were ground and added to 5 mL of 6 N H_2SO_4 to solubilize the oxalate.

After 15 min, 25 ml of distilled water was added, and the entire solution was filtered

through grade 4 Whatman filter paper. The filtrate was brought up to a pH of 7 with

NaOH, and 0.1 g of $CaCl_2$ was added to precipitate the oxalate. The samples were

centrifuged and decanted. After centrifugation, a volume of distilled water equal to

that recovered after filtration was added, and the samples were titrated.

Sample collection for metagenomics

We compared the metagenomic sequencing results across species and gut

regions to investigate differential abundance of microbial functions. We collected

individuals of the cactus specialist *N. albigula*, from Castle Valley, UT, USA (38°30' N,

109°18' W). Additional individuals of *N. stephensi* were collected from near Wupatki

National Monument, AZ, USA (35°30' N, 111°27' W). Animals were immediately

transported back to the University of Utah animal facility, provided with native

diets: either cactus pads or juniper overnight (for *N. albigula* and *N. stephensi*,

respectively), and euthanized and dissected on the following morning. Contents of

the foregut chamber were collected from three individuals of both species, and the

contents of the cecal chamber were collected from *N. stephensi*. We sequenced the

metagenome of each sample (described below), and conducted several comparisons.

First, we compared the foregut metagenomes of *N. albigula* and *N. stephensi*, with

the prediction that the foregut of *N. stephensi* would be enriched in genes associated

with detoxification of juniper, which contains more chemical classes of defensive

compounds than cactus. Second, we compared the metagenomes of foregut contents

and cecal contents in *N. stephensi* with the prediction that the abundance of

detoxification genes would be higher in the foregut to facilitate detoxification prior

to the major site of absorption in the small intestine.

Comparative metagenomics

Total DNA was isolated from foregut and cecal contents using MoBio

PowerFecal DNA extraction kit (MoBio, Carlsbad, California). Extracted DNA was

sent to Argonne National Laboratory for sequencing. Genomic DNA was sheared

using a Covaris Sonicator, to roughly 150 bp and metagenomic shotgun libraries

were prepared using the Illumina TruSeq DNA sequencing preparation kits.

Libraries were sequenced on the Illumina HiSeq2000 platform using a 2 x 100 bp

run and V3 chemistry, which resulted in overlapping sequences.

Metagenomic sequences were uploaded to the MG-RAST website (41).

Sequences were screened against the genome of *Arabidopsis thaliana* to remove

potential contamination from the plant-based diet. The reads were then filtered

using dynamic trimming with a quality threshold of 15, such that any sequences

with more than 5 low-quality bases were removed. Several foregut metagenomes

contained high abundances of sequences (~30-50%) identified as plant DNA, likely

resulting from the diet. Therefore, we partitioned our analysis by domain by

selecting sequences identified as bacteria using the non-redundant multi-source

protein annotation database, M5nr (42). Abundances of non-plant eukaryotic and

archaeal sequences were too low for sufficient analysis. Sequences identified as

bacterial in origin were then annotated to identify putative functions using the

KEGG Orthology database (43, 44) with the following thresholds: (1) e-values < 1e-

5, (2) a minimum percent identity to database sequences of 60% and (3) a minimum

alignment length of 15 bases. We compared the abundances at Level 3 of the KEGG

hierarchical classification, as well as specific genes associated with "Xenobiotic

degradation and metabolism", "Geraniol degradation", and "Limonene and pinene

degradation". Abundances of functional categories were compared using JMP 12.0

(SAS Institute Inc.) using t-tests and the robust Response Screening function, which

performs the Benjamini-Hochberg false discovery rate (FDR) correction for multiple

tests (45).

We also conducted specific comparisons of the abundances of several

detoxification genes: oxalyl-CoA decarboxylase, aryl alcohol dehydrogenase, and

several genes associated with terpene metabolism. The gene oxalyl-CoA

decarboxylase (*oxc*) degrades the compound oxalate. The gene aryl alcohol

dehydrogenase is upregulated in the woodrat foregut when individuals are fed diets

containing phenolic-rich resin from creosote bush (2). Last, a number of genes

associated with the "Limonene and pinene degradation" are enriched or present in

the gut microbiota of specialist insect and avian herbivores that feed on terpene-rich

plants (7, 20). We compared the abundances of these genes using JMP 12.0 (SAS

Institute Inc.) using pairwise t-tests (foregut *N. stephensi* vs. foregut *N. albigula*, and

N. stephensi foregut vs. cecum) with the Dunn's correction.

Metagenome prediction

We also conducted microbial inventories by sequencing the variable regions

1-3 of 16S rRNA gene following experimental and bioinformatic methods described

in detail elsewhere (46). Taxonomic comparison of the communities between

foregut and cecal chambers has already been conducted with larger sample sizes

(11). In the current study, we used 16S rRNA inventories to predict the

metagenomic content of gut samples using genomic inference via PICRUSt (32). We

compared the relative abundances of functional categories between the shotgun

metagenomic and predicted metagenomics to assess if predictive profiling via

PICRUSt is informative for woodrat gut samples. First, we averaged the relative

abundances of functional categories within a sample type (foregut and cecum), and

transformed the relative abundances using a log(x+1) transformation. We then

conducted a linear regression between transformed PICRUSt abundances and

metagenomic abundances. We also plotted the relative abundances of functional

categories that exhibited the highest fold difference between the foregut and cecal

chambers to qualitatively inspect whether trends were similar for PICRUSt and

metagenomic data. We next investigated how variation in structure of the microbial

community across foregut and cecum was related to variation in functional

abundances across these sites. Using a comparative statistical framework

implemented in the R package FunkyTax (46) we classified functions into functional

community categories (enhanced, divergent, conserved, equivalent; see Phillips et

al. 2017, for further detail) resulting from the assessment of how similarities or

differences in functional frequency were compositionally derived between sites. The

significance of functional assignment rates to community categories were assessed

as binomial probabilities where the observed success rate was the frequency of a

function-category assignment, and the expected success rate was the study-wide

frequency of the function. False discovery was controlled with a Benjamini-

Hochberg p-value adjustment.

Accession Numbers

Metagenomic sequences can be found in the NCBI Short Read Archive (SRA)

database under BioProject PRJNA336354, and on MG-RAST under Metagenome

Project IDs mgp10174 and mgp10296. 16S rRNA sequences from the foregut and

cecal chambers of N. stephensi, which were used to PICRUSt analyses can be found

in the SRA database under BioProject PRJNA449358.

Acknowledgements – We would like to thank Madelina James and KayLene

Yamada for helping with feeding trials and Chelsea Merriman for analysis of

monoterpenes. Funding for this study was provided by the National Science

Foundation (Doctoral Dissertation Improvement Grant, DEB 1210094, to M.D.D. and

K.D.K.; DEB 1342615 to M.D.D.; DEB-1146194 to JSF; and DBI 1400456 to K.D.K).

549
550

References

1. Dearing MD, Foley WJ, McLean S. 2005. The influence of plant secondary metabolites on the nutritional ecology of herbivorous terrestrial vertebrates Ann Rev Ecol Evol Syst 36:169-185.
2. Kohl KD, Weiss RB, Cox J, Dale C, Dearing MD. 2014. Gut microbes of mammalian herbivores facilitate intake of plant toxins. Ecol Lett 17:1238-1246.
3. Hume ID, Warner ACI. 1980. Evolution of microbial digetion in mammals, p 665-684. *In* Ruckebusch Y, Thivend P (ed), Digestive Physiology and Metabolism in Ruminants. MTD Press, Lancaster.
4. Mackie RI. 2014. Mutualistic fermentative digestion in the gastrointestinal tract: diversity and evolution. Integr Comp Biol 42:319-326.
5. Sundset MA, Barboza PS, Green TK, Folkow LP, Blix AS, Mathiesen SD. 2010. Microbial degradation of usnic acid in the reindeer rumen. Naturwissenschaften 97:273-278.
6. Garcia-Amado MA, Michelangeli F, Gueneau P, Perez ME, Dominguez-Bello MG. 2007. Bacterial detoxification of saponins in the crop of the avian foregut fermenter *Opisthocomus hoazin*. J Anim Feed Sci 16:82-85.
7. Kohl KD, Connelly JW, Dearing MD, Forbey JS. 2016. Microbial detoxification in the gut of a specialist avian herbivore, the Greater Sage-Grouse. FEMS Microbiol Lett 363:fnw144.
8. Osawa R. 1992. Tannin-protein complex-degrading enterobacteria isolated from the alimentary tracts of koalas and a selective medium for their enumeration. Appl Environ Microbiol 58:1754-1759.
9. Clauss M, Frey R, Kiefer B, Lechner-Doll M, Loehlein W, Polster C, Rössner GE, Streich WJ. 2003. The maximum attainable body size of herbivorous mammals: morphophysiological constraints on foregut, and adaptations of hindgut fermenters. Oecologia 136:14-27.
10. Alexander RM. 1991. Optimization of gut structure and diet for higher vertebrate herbivores. Phil Trans R Soc 333:249-255.
11. Kohl KD, Miller AW, Marvin JE, Mackie R, Dearing MD. 2014. Herbivorous rodents (*Neotoma* spp.) harbour abundant and active foregut microbiota. Environ Microbiol 16:2869-2878.
12. Vaughan TA. 1982. Stephen's woodrat, a dietary specialist. J Mammal 63:53-62.
13. Schwartz CC, Nagy JG, Regelin WL. 1980. Juniper oil yield, terpenoid concentration, and antimicrobial effects on deer. J Wildl Manage 44:107-113.
14. Dearing MD, McLister JD, Sorensen JS. 2005. Woodrat (*Neotoma*) herbivores maintain nitrogen balance on a low-nitrogen, high-phenolic forage, *Juniperus monosperma*. J Comp Physiol B 175:349-355.
15. Holechek JL, Munshikpu L, Saiwana L, Nunez-Hernandez G, Valdez R, Wallace JD, Cardenas M. 1990. Influences of six shrub diets varying in phenol content on intake and nitrogen retention by goats. Trop Grassl 24:93-98.

595   16.   Justice KE. 1985. Oxalate digestibility in *Neotoma albigula* and *Neotoma*
596   *mexicana*. Oecologia 67:231-234.
597   17.   Miller AW, Oakeson KF, Dale C, Dearing MD. 2016. Microbial community
598   transplant results in increased and long-term oxalate degradation. Microb
599   Ecol 72:470-478.
600   18.   Miller AW, Dale C, Dearing MD. 2017. The induction of oxalate metabolism in
601   vivo is more effective with functional microbial communities than with
602   functional microbial species. mSystems 2:e00088-17.
603   19.   Miller AW, Kohl KD, Dearing MD. 2014. The gastrointestinal tract of the
604   white-throated woodrat (Neotoma albigula) harbors distinct consortia of
605   oxalate-degrading bacteria. Appl Environ Microbiol 80:1595-1601.
606   20.   Adams AS, Aylward FO, Adams SM, Erbilgin N, Aukema BH, Currie CR, Suen G,
607   Raffa KF. 2013. Mountain pine beetles colonizing historical and naive trees
608   are associated with a bacterial community highly enriched in genes
609   contributing to terpene metabolism. Appl Environ Microbiol 79:3468-3475.
610   21.   Berasategui A, Salem H, Paetz C, Santoro M, Gershenzon J, Kaltenpoth M,
611   Schmidt A. 2017. Gut microbiota of the pine weevil degrades conifer
612   diterpenes and increases insect fitness. Mol Ecol 26:4099-4110.
613   22.   Morgun A, Dzutsev A, Dong X, Greer RL, Sexton DJ, Ravel J, Schuster M, Hsiao
614   W, Matzinger P, Shulzhenko N. 2015. Uncovering effects of antibiotics on the
615   host and microbiota using transkingdom gene networks. Gut 64:1732-1743.
616   23.   Miller AW, Dearing MD. 2013. The metabolic and ecological interactions of
617   oxalate-degrading bacteria in the mammalian gut. Pathogens 2:636-652.
618   24.   Kohl KD, Pitman E, Robb BC, Connelly JW, Dearing MD, Forbey JS. 2015.
619   Monoterpenes as inhibitors of digestive enzymes and counter-adaptations in
620   a specialist avian herbivore. J Comp Physiol B 185:425-434.
621   25.   Sorensen JS, Turnbull CA, Dearing MD. 2004. A specialist herbivore (*Neotoma*
622   *stephensi*) absorbs fewer plant toxins than does a generalist herbivore
623   (*Neotoma albigula*). Physiol Biochem Zool 77.
624   26.   Iurescia S, Marconi AM, Tofani D, Gambacorta A, Paternò A, Devirgiliis C, van
625   der Werf MJ, Zennaro E. 1999. Identification and sequencing of β-myrcene
626   catabolism genes from *Pseudomonas* sp. strain M1. Appl Environ Microbiol
627   65:2871-2876.
628   27.   Harayama S, Timmis KN. 1989. Catabolism of aromatic hydrocarbons by
629   *Pseudomonas*, p 152-175. *In* Hopwood DA, Chater KF (ed), Genetics of
630   Bacterial Diversity. Academic Press, London.
631   28.   Boundy-Mills K, de Souza ML, Mandelbaum RM, Wackett LP, Sadowsky MJ.
632   1997. The atzB gene of *Pseudomonas* sp. strain ADP encodes the second
633   enzyme of a novel atrazine degradation pathway. Appl Environ Microbiol
634   63:916-923.
635   29.   Seffernick JL, Shapir N, Schoeb M, Johnson G, Sadowsky MJ, Wackett LP. 2002.
636   Enzymatic degradation of chlorodiamino-s-triazine. Appl Environ Microbiol
637   68:4672-4675.
638   30.   Carmona M, Zamarro MT, Blázquez B, Durante-Rodríguez G, Juárez JF,
639   Valderrama JA, Barragán MJ, Garcia JL, Díaz E. 2009. Anaerobic catabolism of

aromatic compounds: a genetic and genomic view. Microbiol Mol Biol Rev
73:71-133.

31. Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L,
Henrissat B, Knight R, Gordon JI. 2011. Diet drives convergence in gut
microbiome functions across mammalian phylogeny and within humans.
Science 332:970-974.

32. Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA,
Clemente JC, Burkepile DE, Thurber RLV, Knight R, Beiko RG, Huttenhower C.
2013. Predictive functional profiling of microbial communities using 16S
rRNA marker gene sequences. Nat Biotechnol 31:814-821.

33. Igimi H, Nishimura M, Kodama R, Ide H. 1973. Studies on the metabolism of
d-limonene (p-mentha-1,8-diene): I. the absorption, distribution and
excretion of d-limonene in rats. Xenobiotica 4:77-84.

34. Min BR, Barry TN, Attwood GT, McNabb WC. 2003. The effect of condensed
tannins on the nutrition and health of ruminants fed fresh temperate forages:
a review. Anim Feed Sci Technol 106:3-19.

35. Shirley EK, Schmidt-Nielsen K. 1967. Oxalate metabolism in the pack rat,
sand rat, hamster, and white rat. J Nutr 91:496-502.

36. Skopec MM, Haley S, Torregrossa A-M, Dearing MD. 2008. An oak (Quercus
agrifolia) specialist (*Neotoma macrotis*) and a sympatric generalist (Neotoma
lepida) show similar intakes and digestibilities of oak. Physiol Biochem Zool
81:426-433.

37. Zduńczyk Z, Juśkiewicz J, Estrella I. 2006. Cecal parameters of rats fed diets
containing grapefruit polyphenols and inulin as single supplements or in a
combination. Nutrition 22:898-904.

38. Price ER, Brun A, Caviedes-Vidal E, Karasov WH. 2015. Digestive adaptations
of aerial lifestyles. Physiol 30:69-78.

39. Vijay-Kumar M, Aitken JD, Carvalho FA, Cullender TC, Mwangi S, Srinivasan S,
Sitaraman SV, Knight R, Ley RE, Gewirtz AT. 2010. Metabolic syndrome and
altered gut microbiota in mice lacking Toll-like receptor 5. Science 328:228-
231.

40. Miller AW, Oakeson KF, Dale C, Dearing MD. 2016. Effect of dietary oxalate on
the gut microbiota of the mammalian herbivore *Neotoma albigula*. Appl
Environ Microbiol 82:2669-2675.

41. Meyer F, Paarmann D, D'Souza M, Olson R, Glass EM, Kubal M, Paczian T,
Rodriguez A, Stevens R, Wilke A, Wilkening J, Edwards RA. 2008. The
metagenomics RAST server–a public resource for the automatic phylogenetic
and functional analysis of metagenomes. BMC Bioinformatics 9:386.

42. Wilke A, Harrison T, Wilkening J, Field D, Glass EM, Kyrpides N, Mavrommatis
K, Meyer F. 2012. The M5nr: a novel non-redundant database containing
protein sequences and annotations from multiple sources and associated
tools. BMC Bioinformatics 13:141.

43. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a
reference resource for gene and protein annotation. Nucleid Acids Res
44:D457-D462.

44.   Kanehisa M, Goto S. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes.
Nucleid Acids Res 28:27-30.
45.   Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a
practical and powerful approach to multiple testing. J R Soc Stat Soc B
57:289-300.
46.   Phillips CD, Hanson J, Wilkinson JE, Koenig L, Rees E, Webala P, Kingston T.
2017. Microbiome structural and functional interactions across host dietary
niche space. Integr Comp Biol 57:743-755.

     Figure Legends

Fig. 1. Changes in food intake and body mass as a result of antibiotics (AB). Bars

represent means ± 1 s.e.m.

Fig. 2. Concentrations of water and toxins across gut regions. The dotted line on

toxin graphs represents concentrations as measured in the food material presented

to woodrats. Points represent means ± 1 s.e.m.

Fig. 3. Functional categories (KEGG Level 3) that differed significantly between the

foregut metagenomes of *N. albigula* and *N. stephensi.* Bars represent means ± 1

s.e.m.

Fig 4. Differences in the metagenomic content of the foregut and cecal chambers of

N. stephensi. Bars represent means ± 1 s.e.m. A) Functional categories (KEGG Level

3) that differed significantly in abundance between the foregut and cecal

metagenomes. B) Detoxification genes that differed significantly in abundance

between the foregut and cecal metagenomes.

Fig. 5. Relative abundances of specific detoxification enzyme genes associated with

a priori hypotheses from other systems (see Methods). Bars represent means ± 1

s.e.m. Relative abundances were compared using t-tests (foregut *N. stephensi* vs.

foregut *N. albigula*, and *N. stephensi* foregut vs. cecum) with the Dunn's correction.

Fig. 6. A) A comparison of predictive profiling by PICRUSt and metagenomic data.

B) Bar graphs depict the functional categories that exhibited the highest fold-

differences between the foregut and cecal metagenomes. Bars represent means ± 1

s.e.m.