1

**Human Papillomavirus Type 16 L1/L2 VLP Experimental Internalization by Human**

**Peripheral Blood Leukocytes**

**Running Title:** HPV16 L1/L2VLP internalization by human leukocytes.

Aurora Marques Cianciarullo[1,2#], Vivian Szulczewski[1,2], Erica Akemi Kavati[1,2], Tania

Matiko Hosoda[1,2], Elizabeth Leão[4], Primavera Borelli[5], Enrique Boccardo[6], Martin Müller[7],

Balasubramanyam Karanam[8], and Willy Beçak[1,9]

[1]Laboratory of Genetics, Butantan Institute, Secretary of Sao Paulo State for Health, 1500

Dr. Vital Brazil Avenue, Sao Paulo, SP 05503-900, Brazil.

[2]Program of Postgraduate Interunits in Biotechnology USP-IBU-IPT, Institute of

Biomedical Sciences, University of Sao Paulo, 2415 Prof Lineu Prestes Avenue,

Biomedical III Building, University City, Sao Paulo, SP  05508-900, Brazil.

[3]Program of Professional Upgrading at Butantan Institute, Secretary of Sao Paulo State for

Health, 188 Dr. Eneas de Carvalho Aguiar, Sao Paulo, SP 05403-000, Brazil.

[4]ELSME – Dr. Elizabeth Leão Specialized Medical Services S/C Ltd., Sao Joaquim

Hospital, Real and Meritorious Portuguese Beneficence Association, 694 Martiniano de

Carvalho Street, Sao Paulo, SP 01321-000, Brazil.

[5]Laboratory of Experimental and Clinical Haematology, Department of Clinical and

Toxicological Analyses, School of Pharmaceutical Sciences, University of Sao Paulo, 580

B17 Prof Lineu Prestes Avenue, University City, Sao Paulo, SP 05508-900, Brazil.

[6]Laboratory of Oncovirology, Department of Microbiology, Institute of Biomedical

Sciences, University of Sao Paulo, 1374 Prof Lineu Prestes Avenue, Biomedical II

Building, University City, Sao Paulo, SP 05508-900, Brazil.

[7]Tumorvirus-specific Vaccination Strategies, Infection, Inflammation and Cancer Program,

German Cancer Research Centre (DKFZ), 242 Im Neuenheimer Feld, Heidelberg 69120,

Germany.

[8]Department of Pathology, Johns Hopkins University, Room 38, Cancer Research Building

2, 1550 Orleans Street, Baltimore, MD 21231, USA.

[9]Department of Biology, Federal University of the Latin-America Integration – UNILA,

1842 Silvio Americo Sasdelli Avenue, Vila A, Commercial Lorivo Building, Foz do

Iguaçu, PR 85866-000, Brazil.

**#Corresponding author present address:** Aurora M. Cianciarullo, Laboratory of Viral

Biotechnology, Butantan Institute, 1500 Dr. Vital Brazil Avenue, Sao Paulo, SP 05503-

900, Brazil, phone +5511 26279725, email: aurora.cianciarullo@butantan.gov.br

**Footnote:** B.K. present addresses Department of Biology and Cancer Research, 1200 W.

Montgomery Road, Tuskegee University, Tuskegee, AL 36088, USA.

**Authors email:** aurora.cianciarullo@butantan.gov.br, emefjmlisboa@prefeitura.sp.gov.br,

erica.kavati@butantan.gov.br, hosodatania@yahoo.com.br, eleao@ibest.com.br,

borelli@usp.br, eboccardo@usp.br, martin.mueller@dkfz.de,

bkaranam@mytu.tuskegee.edu, wbecak@yahoo.com

**ABBREVIATIONS**

HPV, human papillomavirus

VLP, virus-like particles

L1, major capsid protein

L2, minor capsid protein

HEK293T, human epithelial cell lineage 293T

CD71, transferrin receptor

CD4, T lymphocyte membrane receptor

CD8, T lymphocyte membrane receptor

CD20, B lymphocyte membrane receptor

CD14, monocyte membrane receptor

rCTB, recombinant cholera toxin B subunit is the non-toxic portion of

cholera toxin (CT)

HIV, human immunodeficiency virus

pUF3L1h vector, humanized expression vector for L1 protein

pUF3L2h vector, humanized expression vector for L2 protein

EDTA, anticoagulant

Tf, transferrin

DMEM, Dulbecco's Modified Medium

FBS, fetal bovine serum

RPMI, Roswell Park Memorial Institute cell culture medium

PBMC, peripheral blood mononuclear cells

BSA, bovine serum albumin

TEM, transmission electron microscopy

**ABSTRACT**

Human papillomavirus (HPV) accounts for hundreds of thousands of new cases of cervical

cancer yearly, and half of these women die of this neoplasia. This study investigates the

possibility of HPV16 L1/L2VLP to be internalized by human peripheral blood leukocytes

in *ex vivo* assays. We have developed a leukocyte separation method from heparinized

blood samples aiming cellular integrity and viability. We have expressed humanized L1

and L2 viral capsid proteins in HEK293T epithelial human cells, transiently transfecting

them with vectors encoding humanized HPV16 *L1* and *L2* genes. Recombinant L1/L2

capsid proteins and structured virus-like particles interacted with human peripheral blood

mononuclear cells, lymphocytes and monocytes, and were internalized through a pathway

involving CD71 transferrin receptors. This was observed, at a percentile of about 54% T-

CD4, 47% T-CD8, 48% B-CD20, and 23% for monocytes-CD14. The group of polymorph

nuclear cells: neutrophils-eosinophils-basophils group did not internalize any VLPs.

Blockage assays with biochemical inhibitors of distinct pathways, like chlorpromazine,

rCTB, filipin, nystatin, liquemin, and sodium azide also evidentiated the occurrence of

virus-like particles indiscriminate entrance via membrane receptor on mononuclear cells.

This study shows that HPV16 L1/L2 VLPs can interact with the plasma membrane surface

and successfully enter lymphocytes without requiring a specific receptor.

**Keywords:** HPV16 L1/L2 VLP, VLP internalization by lymphocytes, membrane receptors,

leukocytes, mononuclear cells, cancer risk factor.

**INTRODUCTION**

In 1974, zur Hausen researches pointed to the HPV as the major etiological agent in

cervical cancer and, later, its DNA was detected in tumors found in other anogenital regions

(1-3). Today, there are more than 200 different types of well characterized HPVs, of which

approximately 40 types infect the genital tract. The most frequently found are types 6 and

11 that induce benignant skin warts formation, and types 16 and 18 which are associated to

cancer development. There are at least 12 HPV types that according to the Agency for

Research on Cancer (IARC) are considered oncogenic to humans (4, Papillomavirus

Episteme (PaVE); http://pave.niaid.nih.gov/). Among them, HPV16 and HPV18 are those

responsible for approximately 70% of all cervical cancers (4, 5). In 2025, the projected

global estimate of cervical cancer is expected to rise to 720,415 new cases per year, and it

is expected that half of these women will die of this neoplasia (5). More recently, novel

HPV lineages and sub-lineages were described, found in HIV/HPV-co-infected pregnant

women, with two high-risk types (6). However, the difficulty to obtain enough viable wild

types or recombinant HPV particles has limited researches to distinct aspects of virus

biology (7). All viruses enter host cells to survive, replicate and evade the immune system.

So far, the entry of HPV and its traffic through the host cells are still not completely

elucidate (8). Several studies describe this process like a complex set of interactions among

different pathways, receptors, co-receptors and co-factors. In keratinocytes, HPV seems to

be internalized via clathrin-dependent endocytic mechanisms, but it might use alternative

uptake pathways to enter cells, such as a caveolae-dependent route, among others

depending on viral type (7-10).

Currently, HPV is recognized as one of the main causes of infection-related cancer

worldwide, as well as the causal factor of other diseases. Infection with high-risk HPV

types is the etiological cause of cervical cancer and is strongly associated with a significate

fraction of penile, vulvar, vaginal, anal and oropharyngeal cancers (11, 12). HPVs are

responsible for approximately 88% of anal cancer and 95% of anal intraepithelial neoplasia

grades 2/3 lesions, and 40%-50% of penis and vulvar cancers (11). In oropharyngeal

cancers, HPV DNA was detected in 35%-50% cases (11). In all HPV-positive non-cervical

cancers, HPV16 is the most common HPV type detected, followed by HPV types 18, 31, 33

and 45. Among the non-cancerous HPV-associated conditions, genital warts and recurrent

respiratory papillomatosis are surely linked to HPV6 and 11 (10). In addition, it was

demonstrated recently that HPV16 virus-like particles (VLPs) L1/L2 interact with

hematopoietic precursor stem cells, present in the amniotic fluid from healthy pregnant

women (13). Whereas new pathologies are increasingly being associated to the HPVs; the

responsibility and costs of HPV-associated diseases and cancer remain an important public

health issue in all countries, regardless of their economic developmental level (14). Due to

the continuous worldwide propagation of HPV it is necessary to investigate the possibility

of HPV internalization by different human cell types. In the present study, we addressed the

capacity of HPV16 L1/L2 VLP entrance/uptake of − in peripheral blood leukocytes *ex vivo*.

**MATERIALS AND METHODS**

**Production of HPV16 L1/L2 VLPs**

The production of VLPs was carried out throughout the recombinant protein expression

HPV16 L1 and L2 in epithelial human cells of the HEK293T lineage, cultured as

previously described (15). Transfection of these cells occur in a transient way using

pUF3L1h and pUF3L2h vectors, which are regulated by the human cytomegalovirus

promoter containing complete sequences of genes *L1* and *L2*, a method already described in

detail (15, 16). In this study, besides being used to produce VLPs containing both capsid

proteins, HEK293T cell line was also used as a control factor on internalization assays with

human peripheral blood leukocytes. All assays performed in this study were analyzed in

duplicate, being representative of at least four independent tests.

**Blood collection**

Blood samples from 10 healthy female volunteers, ages ranging from 35 to 55, were

requested based on epidemiological data associated with genital HPV infection (17). The

screening of the volunteers was based on recent blood, Pap smears and colposcopy tests

and on data filled out by the candidate on a written informed consent form and on a

questionnaire, for the purpose of laboratory research. Volunteers who did not present Pap

and colposcopy results within reference values were dismissed. Blood samples were

collected in sterile tubes containing heparin, for internalization assays with leukocytes, and

EDTA for blocking assays, and were processed quickly, within a maximum of 2-hours after

collection. During this interval, they were stored at $4^{\circ}$C.

**Leukocytes separation**

Whole blood was collected in heparin at a concentration of 0.1 mg/ml, centrifuged at 1,000

rpm for 7 minutes at 5$^{o}$C in a Sorvall$^{®}$ RT 6000 Refrigerated Centrifuge (Du Pont,

Wilmington-DE, USA) with a horizontal angle rotor. The plasma supernatant containing

platelets and leukocytes was gently removed and transferred with the aid of a Pasteur

pipette to a new tube. An aliquot was collected for total and differential counts, performed

in a Neubauer chamber, diluted (ratio 1:1) in Trypan blue and smears were stained by May-

Grünwald-Giemsa in order to determine their composition. Leukocytes sedimented were

centrifuged again at 1,200 rpm, for 3 minutes, at 5$^{o}$C, to remove platelets. Leukocytes

precipitates were resuspended in 0.85% saline solution for subsequent tests (18). The

Neubauer chamber counts and smears containing distinct cell types were analyzed by light

microscopy, with a Leica DMIL I microscope (Leica Microsystems GmbH, Vienna, AUT).

**Internalization assays for HEK293T cells and HPV16 L1/L2 VLPs**

HEK293T cells ($2 \times 10^{4}$ cells/ml) were plated and maintained under growth conditions,

washed with PBS and incubated with 120 µg of VLPs in DMEM without FBS, for 4-hours

at 37$^{o}$C and 5% $CO_2$. After this, cells were washed twice with PBS for 3 minutes each in

order to remove non-internalized particles. Cells were then fixed with 2% PFA

(Paraformaldehyde, Sigma-Aldrich) in PBS for 1-hour, at 4$^{o}$C, and washed three times with

PBS, 5 minutes each. Plates were kept at 4$^{o}$C until immunofluorescence assays. Controls

were performed in the absence of VLPs and/or the denaturation thereof, by heating at

100$^{o}$C for 10 minutes (19, adapted).

**Internalization assays for human leukocytes and HPV16 L1/L2 VLPs**

Leukocytes ($2 \times 10^4$ cells/ml) of healthy volunteers were incubated with 120 μg of VLPs

produced in this study and RPMI without FBS for 4-hours at $37^{\circ}$C and 5% $CO_2$, under

gentle agitation. Cells were centrifuged at 1,000 rpm for 7 minutes and cell pellets washed

three times with PBS, 5 minutes each. Then, cells were fixed with 2% PFA in PBS for 1-

hour, at $4^{\circ}$C. After this step, the centrifugation and PBS washing processes were repeated.

Samples were kept at $4^{\circ}$C until immunofluorescence assays. Controls were the same as

described above for HEK293T cells.

**Internalization assays for leukocytes, transferrin and HPV VLPs**

The same protocol described above was employed, with minor changes described below.

Leukocytes were incubated with 120 μg of VLPs in separate samples, as follows: the

HPV16 L1/L2 VLPs produced in this study, and L1 VLPs of HPV6, 11, 16 and 18 from

Gardasil® vaccine, used as control, kindly provided by Merck Sharp & Dohme. RPMI

without FBS, together with transferrin (Tf) conjugated to fluorochrome TexasRed®

(Molecular Probes™), were added (ratio 1:60, Tf:RPMI). Samples were kept for 15, 45, 60

and 120 minutes in an incubator at $37^{\circ}$C and 5% $CO_2$, under gentle agitation. The other

procedures remained unchanged.

**Blocking assays for membrane receptors**

Leukocytes of healthy donors were counted and incubated in RPMI medium overnight over

coverslips containing poly-L-lysine (Sigma-Aldrich), at $37^{\circ}$C and 5% $CO_2$. Then, cells

were washed and a fresh medium was placed together with specific biochemical inhibitors

of ligand uptake (Table 1), used isolated or associated, and incubated for 2 hours (19). After

incubation, cells were washed and 120 μg of HPV16 L1/L2 VLPs (20) were added and

incubated with medium for 4 hours. Cells were washed again and fixed in 2% PFA

solution. Immunofluorescence assays were carried out to detect the VLP-PBMC interaction

by confocal microscopy, using specific antibodies to recognize L1 and L2 proteins. Z-axis

3D images were obtained to confirm the presence of VLPs within cells.

**244    Immunofluorescence assays for internalization analysis**

Leukocytes were washed and fixed as already described. Samples were incubated in PBS

containing 1% BSA for 5 minutes under gentle agitation. Later, cells were incubated with

the primary antibodies (Table 1); diluted in PBS containing 0.01% Tween® 20 and 0.5%

BSA pH 8 for 2 hours, under gentle agitation, at room temperature. They were then washed

three times with PBS for 10 minutes each, followed by incubation with the corresponding

secondary antibodies (Table 2) conjugated with fluorochromes and diluted in PBS

containing 0.01% Tween® 20 and 1.5% BSA for 1 hour, under light stirring at room

temperature. Once more, they were washed three times with PBS for 10 minutes each.

Samples of leukocytes with transferrin and VLPs were labelled with Phalloidin conjugated

AlexaFluor® 594, incubated for 20 minutes. Immediately after that, they were rinsed twice

with PBS for 10 minutes each. Aliquots of cell suspension from both items assayed were

adhered over silanized slides, and mounted with 5 µl Mowiol® and coverslips. Samples

were kept at 4°C until CLSM analysis, using Confocal Laser Scanning Microscope Zeiss

510 Meta of the Butantan Institute (FAPESP Process No. 2000/11624-5; Carl Zeiss GmbH,

Jena, DEU).

**RESULTS**

**Leukocytes identification**

After isolation of leukocytes from healthy donors, control smears stained by May-

Grünwald-Giemsa method showed cell morphology preservation (18), presenting a positive

correlation with the control blood smears. Counts indicated approximately 98%

lymphocytes; 0.7% monocytes; 1.5% polymorph nuclear cells; 0.1% red blood cells and

0.1% platelets. The ultrastructure was well preserved for all cell types analyzed, with well-

defined and intact membranes (18). The efficiency and speed of the method for obtaining

leukocytes, preserving morphology and cell viability was confirmed.

**Analysis of HPV16 L1/L2 VLPs internalization by HEK293T cells**

HPV16 L1/L2 VLPs internalization by HEK293T cells was analyzed through

immunofluorescence. We emphasize that the genome of this cell line does not contain any

HPV DNA sequences, but it contains Adenovirus DNA and SV40 T-Ag. After 4-hours kept

in contact at $37^{o}C$, cells were washed in order to remove any particles or proteins not

internalized by HEK293T cells. Then, cells were fixed and stained with the primary

antibody Camvir-1 (BD Biosciences – Table 1) for L1 proteins and with L1 conformation-

specific anti-VLP antiserum (Biodesign – Table 1). It was therefore possible to observe the

internalization of VLPs in some cells. In these cells VLPs were found in the cytoplasm,

near the core region, stained with anti-VLP. These results were confirmed by the overlap of

images, and by the most thoroughly detailed internalization display on the Z-axis scanning

sections. The morphological evaluations show that probably the VLPs internalization

occurs simultaneously with a large number of particles, similar to the formation of

endocytic vesicle structures. These observations suggest that after 4 hours, structured VLPs,

as well as non-structured VLPs, such as pentameric and monomeric forms of L1 stained

with Camvir-1 (Table 1), were internalized in epithelial cells from the human kidney

(HEK293T), across the cell membrane. The negative controls showed no immunostaining

for L1 and VLPs and no changes in cell morphology were detected.

**Analysis of HPV16 L1/L2 VLPs internalization by human leukocytes**

L1/L2 particles were added to human leukocytes, for 4-hours at 37$^{o}$C, and prepared for

indirect CLSM immunofluorescence analysis, in order to investigate the possibility of

internalization between the leukocytes and the HPV16 L1/L2 VLPs. Results indicate an

internalization of VLPs with peripheral blood mononuclear cells (PBMC), T and B

lymphocytes and monocytes, from healthy women volunteers (Fig. 1). Structured HPV16

L1/L2 VLPs also interacted with and were internalized by leukocytes (Figs. 2 and 3). After

4-hours of interaction, in most leukocytes examined, these particles were found in the cell

cytoplasm, as was the case with HEK293T. These results were confirmed by the images

overlap (Figs. 1 A, B and D; Figs. 2 and 3), and the detailed internalization, as shown on

the Z-axis scanning sections (Fig. 2 C). Fig. 1 C (green arrow) shows VLPs across the cell

membrane. The Z-axis sweep cuts were able to demonstrate the internalization of the VLP

with a larger number of particles similar to the formation of endocytic vesicles structures

(Fig. 2 A, white arrow), like those found in HEK293T cells, a phenomenon endorsed by

morphological evaluations. In Fig. 3, the interaction of leukocytes with the VLPs is

illustrated, showing the colocalization of HPV16 L1/L2 proteins recognized by anti-L1

(green) and anti-L2 (red), respectively. This colocalization was expected in VLPs that are

composed of these two proteins, as it might be seen from the images overlap (Fig. 3). Fig. 3

B shows the internalization of VLPs by leukocytes in a significant amount when compared

311 to other internalizations. At least 15 fields were analyzed per experiment, containing from 4

312 to 10 cells per field. The presence of structures similar to vacuoles can also be observed

313 (Fig. 3 B, blue arrow). Control assays showed no immunostaining, neither with the

314 antibodies used to detect L1 and L2 separately, nor with structured HPV16 L1/L2 VLPs.

**Identification of VLPs internalization by human PBMC**

317 These cells were treated with antibodies in order to recognize specific cell membrane

318 receptors for each type (Table 1 and Table 2), and analyzed by indirect

319 immunofluorescence using CLSM to identify PBMC interactions with HPV16 VLPs, after

320 4-hours incubation at $37^{o}$C. HPV16 VLPs interacted with *ex vivo* PBMC (Figs. 4-6). These

321 results were confirmed by the images overlap (Figs. 4 and 5) and displayed detailed

322 internalization on the Z-axis sections (Fig. 6). T and B lymphocytes showed greater

323 competence to internalize VLPs, at around 47%-52% of cells (Table 3). Through

324 morphological assessments, T-lymphocytes that internalized VLPs and were recognized by

325 anti-CD8 showed endocytic vesicles-like structures (Fig. 4 B, white arrow). Only 23% of

326 monocytes that were identified by anti-CD14 antibody interacted with the VLPs (Table 3).

**Analysis of the colocalization of VLPs and transferrin (Tf) in human PBMC**

329 After 15 minutes, no colocalization between L1 VLPs and exogenous Tf was detected

330 (Fig.7). Moreover, it was possible to show the internalization of VLPs during this period.

331 After 45 minutes, colocalization of VLPs with exogenous Tf in the cytoplasm of human

332 PBMC (Fig. 8 and 9, yellow arrows), and also with the TfR or CD71 was observed (Fig. 9

333 A-B, yellow arrows), suggesting that the iron pathway may be used to HPV internalization.

334 All results were confirmed by the images overlap, respectively (Figs. 7-9).

We compared the kinetics of Tf and TfR (CD71) internalization with the colocalization of

HPV16 L1/L2 VLPs produced in this study and with control HPV6, 11, 16, 18 L1 VLPs

derived from the Gardasil[®] vaccine (Fig. 9). After only 15 minutes kept in contact it was

possible to observe colocalization of TfR and Tf (Fig. 9 A-B, purple arrows) with both

L1/L2 VLPs (Fig. 9 A) as well as with Gardasil L1 VLPs (Fig. 9 B). By means of

morphological analysis, we were able to demonstrate the colocalization of both VLPs L1

and L1/L2, with Tf and TfR in the leukocytes cytoplasm, after a 45-minutes period kept in

contact (Fig. 9 C-D, white arrows). This colocalization among VLPs, Tf and TfR was

found in approximately 50% of analyzed cells. Within 60 minutes in contact, we observed a

feeble colocalization between Tf and VLPs in about 30% of the cells (Fig. 9 E-F, orange

arrows). No colocalization was observed in the tests after 120 minutes. The colocalization

was evidentiated in detail by three-dimensional figures stacked in Z-axis (Fig. 9). The

negative controls showed no signs of L1 and VLPs. However, it was possible to observe

exogenous Tf internalization. Comparative testing of L1/L2 VLPs produced in this study

and of control L1 VLPs from Gardasil[®] vaccine showed no significant differences in the

analyzed results.

**Analysis of the blocking assays of PBMC membrane receptors**

After successful identification of VLPs within leukocytes and their colocalization with Tf

and TfR, a variety of biochemical inhibitors (Table 4), known to inhibit distinct cellular

processes, was used to demonstrate the involvement of these and other different pathways

of VLPs entry in PBMC.

Chlorpromazine inhibits clathrin-mediated endocytosis of various plasma membrane

proteins. Nystatin is a sterol-binding agent that disassembles caveolae in the membrane.

rCTB (Clostridium toxin B) acts by shortening the actin filaments length, by inhibiting its

process *in vitro*. Liquemin is related with the VLPs obstruction of internalization through

the HSPG (Heparan sulphate proteoglycans) pathway. Sodium azide is an ATPase

inhibitor, which blocks particle movement towards the cell body and leads to a diffuse

random movement.

The use of these different biochemical inhibitors was not sufficient to block the HPV16

L1/L2 VLPs entry in the PBMC (Figs. 10-11). Chlorpromazine, although known for

blocking the clathrin pathway through CD71 receptor of Tf, was not able to inhibit VLPs

entrance inside PBMC in a satisfactory manner. The same occurred when rCTB, Filipin and

Nystatin, Liquemin and Sodium azide were used to block actin-dependent cell processes,

the caveolae pathway, the HSPG pathway, and endocytic pathways, respectively, either

assayed separately or associated in an all-inhibitors cocktail. These results suggest that

PBMC makes use of multiple internalization pathways to uptake HPV16 L1/L2 VLPs.

**DISCUSSION**

Differently of many Beta and Gamma papillomavirus, Alpha papillomaviruses developed

immune system evasion strategies from the host, causing persistent and visible squamous

papillomas, which sometimes evolving to cancers (21-23). Some HPV types including 16,

18, 31, 33, 35, 39, 45, 51, 52, 56, 58 and 59 have been classified as human carcinogens by

the IARC, being responsible for at least 4% of all human malignancies (4, 23, 24

Papillomavirus Episteme (PaVE); http://pave.niaid.nih.gov/). Infection by high-risk HPV

types is the major factor for the development of cervical cancer, and HPV DNA is also

found in tumors affecting other anogenital regions. Besides, they are being correlated with

an increasing proportion of oropharyngeal squamous cells carcinomas, and occurrences

affecting mainly tonsil and base of tongue regions are currently being documented in

particular geographic areas (25-27). Cell-virus interaction is the initial step in viral

infection. Often, virus adsorption occurs on the plasma membrane of susceptible cells. In

general, the most common solution is that the pathogen intracellular trafficking occurs

through and existing entry mechanism, mediated via clathrin, caveolin, macropinocytosis,

among others, moving inside cells to reach the location where viral replication occurs (28).

Transcytosis is used to move antigens and protective antibodies across epithelial barriers.

Similar to what seems to occur with HIV (29), the transcytosis of HPV through epithelial

cells, crossing cellular barriers, should also be dependent on trafficking to the endocytic

recycling pathway. The productive life cycle of HPV is directly linked to epithelial

differentiation. It is believed that the papillomavirus is capable of maintaining the

expression of structural genes, under tight control of the transcription and translation

regulatory mechanisms (30). The HPV L1 capsid protein, regardless of the presence or

absence of L2, has the property of forming structures that mimic morphologically authentic

virions, VLPs, which are being used as replacements for HPV infected cell culture *in vitro*

(9, 15, 31, 32). However, the presence of L2 seems to confer greater stability to the virus

and VLP, therefore, the viral capsid indeed seems to contribute to the virus infectivity (33).

Most experimental models explore the process of host cell interaction by molecular or

biochemical methods, using cell lines from different tissues and species in interactions with

HPV L1 VLPs in their studies.

In the present work, we produced HPV L1/L2 VLPs (15, 16) to investigate the possibility

that these particles to be internalized by human leukocytes. We have developed a method

for the isolation of these cells (18), used in suspension at $37^{o}C$, thereby creating an *ex vivo*

system as closely as possible to the natural conditions of human body, thus conceiving a

novel experimental model.

HPV16 recombinant proteins L1, L2 and VLPs were visualized in the perinuclear region of

HEK293T cells, in compliance with data described in the literature (34). The presence of

BPV (bovine papillomavirus) in lymphocytes has been discussed (35, 36). Using PCR

technique (Polymerase Chain Reaction) in samples taken from humans with anogenital

lesions, the presence of HPV16 in peripheral blood samples (37) and in plasma cells (38)

was detected. Besides, DNA of other HPV types has been found in PBMC (39, 40). The

presence of HPV in lymphocytes of patients treated for anogenital cancers suggests that the

virus can remain in these cells even after treatment. However, there is no convincing

evidence supporting that lymphocytes carry/produce infectious HPV viral particles.

Nonetheless, the presence of HPV macromolecules in lymphocytes may play a role in viral

immune response.

The scientific literature discloses that papillomavirus exhibits specific tropism for species

and tissues. However, BPV1 particles isolated from bovine warts, and HPV16 VLPs

interacted successfully with 14 cell lines derived from different species and tissues, as

demonstrated in two different studies (32, 41). Using our model to investigate this, HPV16

L1/L2 VLPs internalized *in vitro* and *ex vivo* with two very distinct human cells, epithelial

kidney cells and PBMC, respectively. After 4-hours of interaction, these particles

internalization occurred, showing a very similar distribution pattern on both cell types. In

both, L1 VLPs were located in the cytoplasm near the nucleus, suggesting that tissue

differences do not affect the ability of HPV16 VLPs to entry in different cell types, and that

pentameric and monomeric forms of L1 may also be internalized. The interaction between

VLPs and monocytes observed in this study showed colocalization with CD14. As

macrophage precursors, apparently these cells are able to internalize particles at a much

larger amount if compared with lymphocytes. The unexpected colocalization of CD14

membrane receptor and VLPs, as well as the morphological changes observed, suggest that

monocytes respond to this interaction as immune cells, in order to eliminate invading

microorganisms. CD14 is a 53-kDa molecule, expressed mainly in monocytes,

macrophages and granulocytes. It is also found as soluble protein in the serum (42).

Tf is a beta globulin responsible for the transport of iron (Fe) through plasma to the interior

of cells. Besides hepatocytes, other cells in the body are able to synthesize it. The TfR is a

membrane protein expressed primarily in T and B lymphocytes, macrophages, and

proliferating cells such as homodimers with N-terminal cytoplasmic tails (42). TfR is

constantly being endocytosed by plasma membrane through clathrin-mediated endocytosis,

with the purpose of carrying ferric iron bound to Tf to the inside of cells. This Tf-TfR

complex is delivered to the early endosome into a lower pH medium, where iron is released

and TfR is recycled to the cell surface (43). When evaluating the interactions between

HPV16 L1/L2 VLPs, HPV6, 11, 16, 18 L1 VLPs (Gardasil® vaccine) and unstructured

HPV16 L1 in PBMCs and in the presence of exogenous Tf, we determined the

colocalization of L1-VLP-Tf and free Tf in these cells, as well as colocalization with TfR,

after 45 minutes of contact. This colocalization between VLP, Tf and TfR was found in

approximately 50% of analyzed cells. These results suggest that HPV16 L1/L2 VLPs,

HPV6, 11, 16, 18 L1 VLPs (Gardasil® vaccine) and unstructured HPV16 L1 can enter cells

by endocytosis through clathrin-dependent, via TfR. It is probable that, along viral infection

processes, interactions between HPV16 (about 55 nm size) and Tf molecules (80-kDa)

might indeed happen, through fusion with Fe-Tf-TfR complexes, mimicking its entry into

the cell via low pH endosomes, thus enabling transcytosis (28, 44-47).

There have been no reports in the literature showing the internalization of HPV VLPs in

human PBMC. So far, this study has showed that PBMCs of healthy female volunteers

were able to internalize HPV16 VLPs. Also, T and B lymphocytes were the main cell types

that further internalized VLPs. It is important to emphasize that the lymphocytes are cells

capable of dividing themselves and it is estimated that the half-life of these inactive cells in

humans is of some years. Furthermore, inactive lymphocytes circulate continuously through

the bloodstream and lymphatic vessels. The entry of HPV VLPs in leukocytes by

independent endocytosis pathways may be due to their ability to recognize, capture and

remove foreign substances from the body. The use of different biochemical inhibitors,

isolated or associated, does not seem to be enough to block the internalization of HPV16

L1/L2 VLPs in the PBMC. However, in order to understand the possible mechanisms

involved in these PBMC interactions in future studies, which in turn cannot be assessed

only by VLP experiments, it will be necessary to drawn new strategies using PsV

(pseudovirion) infectious particles provided with HPV16 viral genome (48-50). Thinking in

another approach, it will be possible to exploit this experimental model in vaccines-cells

interaction studies and the immunological mechanisms involved, for instance, evaluating

the impact of vaccines and the prevalence of determined viral genotypes in distinct

populations (51-54). A recent review discusses interestingly the importance to clarify some

lapses of our understanding about HPV natural history (55).

## CONCLUSIONS

**CONCLUSIONS**

HPV16 L1/L2 VLPs produced in this study interacted with *in vitro* HEK293T cells and *ex*

*vivo* PBMC from healthy women volunteers, and were internalized. This study evinced that

HPV16 L1/L2 VLPs can interact with the plasma membrane surface, been uptake by

lymphocytes. We also demonstrated that HPV16 L1/L2 VLPs might utilize the endocytic

pathway of iron-mediated clathrin as a receptor, which involves multiple membrane

receptors simultaneously and also the classical endocytic pathway of iron, the most

widespread among vertebrates and the animal kingdom as a whole, we have demonstrated

that HPV16 L1/L2 VLPs does not require a specific receptor in order to be internalized by

leukocytes. Complementary studies are required to accurately demonstrate the probable

mechanisms involved.

**AUTHORSHIP**

A.M.C. conceptualized and wrote the manuscript.

**ACKNOWLEDGEMENTS**

The authors would like to express their deepest gratitude to the CAPES Master Degree

Fellowship to V. Szulczewski and E.A. Kavati, from the Biotechnology Interunits

Postgraduate Program USP-IBU-IPT; PAP-SES-FUNDAP Fellowships to T.M. Hosoda;

## ETHICAL COMMITTEE

This work was duly evaluated and approved by the Sao Joaquim Hospital, Real and

Meritorious Portuguese Beneficence Association Ethical Committee on Human Research.

Protocol number: 369-08.

This work is in full compliance with the National Biosafety Law (CTNBio) guidelines and

the present study was approved by the Butantan Institute Institutional Biosafety Committee

(CIBio) and CTNBio – Process Number 01200.004893/1997-93, published in the Brazilian

Official Gazette (DOU) on August 16, 2011.

## GRANT SUPPORT

518  This work was supported by grants from the Foundation for Research Support of the State

519  of Sao Paulo – FAPESP (Process number 2004/15122-5), Butantan Institute and Butantan

520  Foundation, Sao Paulo, SP, Brazil.

## DISCLOSURES

523  The authors disclose no potential conflict of interest with the publication of this article.

**REFERENCES**

1. **zur Hausen H, Meinhof W, Scheiber W, Bornkamm GW.** 1974. Attempts to detect virus-specific DNA sequences in human tumors: I. Nucleic acid hybridizations with complementary RNA of human wart virus. Int J Cancer **13:**650-656. (First negative attempt to find papillomavirus DNA in cancer of the cervix with labelled HPV DNA from plantar warts).

2. **zur Hausen H.** 1976. Condylomata acuminata and human genital cancer. Cancer Res **36:**794. (Hypothesis that papillomaviruses are involved in the etiology of cancer of cervix).

3. **Dürst M, Dzarlieva-Petruseyska RT, Boukamp P, Fusenig NE, Gissmann L.** 1987. Molecular and cytogenetic analysis of immortalized human primary keratinocytes obtained after transfection with human papillomavirus type 16 DNA. Oncogene **1:**251-256.

4. **Viarisio D, Gissmann L, Tommasino M.** 2017. Human papillomaviruses and carcinogenesis: well-established and novel models. Curr Opin Virol **26:**56-62.

5. **Sanjosé S, Serrano B, Castellsagué X, Brotons M, Muñoz J, Bruni L, Bosch FX.** 2012. Human papillomavirus (HPV) and related cancers in the Global Alliance for Vaccines and Immunization (GAVI) countries. AWHO/ICO HPV Information Centre Report. Vaccine **30S:**D1-D83.

6. **Siqueira JD, Alves BM, Prellwitz IM, Furtado C, Meyrelles AR, Machado ES, Seuánez HN, Soares MA, Soares EA.** 2016. Identification of novel human papillomavirus lineages and sublineages in HIV/HPV-coinfected pregnant women by next-generation sequencing. Virology **493:**202-208.

550 7. **Day PM, Lowy DR, Schiller JT.** 2003. Papillomaviruses infect cells via a clathrin-
551   dependent pathway. Virology **307:**1-11.

552 8. **Aksoy P, Gottschalk EY, Meneses PI.** 2017. HPV entry into cells. Mutation Res
553   **772:**13-22.

554 9. **Bousarghin L, Touzé A, Combita-Rojas AL, Coursaget P.** 2003. Positively
555   charged sequences of human papillomavirus type 16 capsid proteins are sufficient to
556   mediate gene transfer into target cells via the heparan sulfate receptor. J Gen Virol
557   **84:**157-164.

558 10. **Hindmarsh PL, Laimins LA.** 2007. Mechanisms regulating expression of the HPV
559   31 L1 and L2 capsid proteins and pseudovirion entry. Virol J **4:**1-12.

560 11. **Bosch FX, Broker TR, Forman D, Moscicki AB, Gillison ML, Doorbar J, Stern**
561   **PL, Stanley M, Arbyn M, Poljak M, Cuzick J, Castle PE, Schiller JT,**
562   **Markowitz LE, Fisher WA, Canfell K, Denny LA, Franco EL, Steben M, Kane**
563   **MA, Schiffman M, Meijer CJ, Sankaranarayanan R, Castellsagué X, Kim JJ,**
564   **Brotons M, Alemany L, Albero G, Diaz M, Sanjosé S**. 2013. Authors of ICO
565   Monograph Comprehensive Control of HPV Infections and Related Diseases.
566   Vaccine **31:**H1-31. Collaborators (98)

567 12. **Alemany L, Saunier M, Alvarado I, Quirós B, Salmeron J, Shin HR, Pirog E,**
568   **Guimerà N, Hernández GA, Felix A, Clavero O, Lloveras B, Kasamatsu E,**
569   **Goodman MT, Hernandez BY, Laco J, Tinoco L, Geraets DT, Lynch CF,**
570   **Mandys V, Poljak M, Jach R, Verge J, Clavel C, Ndiaye C, Klaustermeier J,**
571   **Cubilla A, Castellsagué X, Bravo IG, Pawlita M, Quint W, Muñoz N, Bosch FX,**
572   **Sanjosé S on behalf of the HPV VVAP study group.** 2015. HPV DNA prevalence
573   and type distribution in anal carcinomas worldwide. Int J Cancer **136:**98-107.

13. **Kavati EA, Palumbo ACM, Andrade FB, Marigliani B, Sakauchi D, Leão E,**

**Armbruster-Moraes E, Müller M, Cianciarullo AM.** 2012. Interaction of

HPV16L1L2 VLP with stem cells CD34+/CD117+ of the human amniotic fluid. In:

Current microscopy contributions to advances in science and technology. Editor:

Antonio Méndez-Vilas. Publisher: Formatex Research Center, Badajoz, Spain.

Volume **5:**617-624. ISBN-13: 978-84-939843-5-9.

http://www.formatex.info/microscopy5/book/617-624.pdf

14. **Dunne EF,  Markowitz LE, Saraiya M, Stokley S, Middleman A, Unger ER,**

**Williams A, Iskander J.** 2014. Centers for Disease Control and Prevention. CDC

Grand Rounds: Reducing the Burden of HPV-Associated Cancer and Disease.

MMWR **63:**69-72.

15. **Cianciarullo AM, Szulczewski V, Chaves AAM, Bazan SB, Aires KA, Müller M,**

**Armbruster-Moraes E.** 2010. Production of HPV16 L1L2 VLPs in cultures of

human epithelial cells. In: Microscopy: Science, Technology, Applications and

Education. Editors: Antonio Méndez-Vilas and Jesús Díaz Álvarez. Publisher:

Formatex Research Center, Badajoz, Spain. Volume **4:**1073-1082. ISBN- 13: 978-

84-614-6190. http://www.formatex.info/microscopy4/ 1073-1082.pdf

16. **Marigliani B, Kavati EA, Sakauchi D, Oliveira HB, Canali RA, Sasaki AA,**

**Ferreira Jr JMC, Armbruster-Moraes E, Müller M, Cianciarullo AM.** 2012.

Intracellular distribution of recombinant Human Papillomavirus capsid proteins. In:

Current microscopy contributions to advances in science and technology. Editor:

Antonio Méndez-Vilas. Publisher: Formatex Research Center, Badajoz, Spain.

Volume **5:**678-684. ISBN-13: 978-84-939843-5-9.

http://www.formatex.info/microscopy5/book/678-684.pdf

17. **Nonnenmacher B, Breitenbach V, Villa LL, Prolla JC, Bozzetti MC.** 2002.

Genital human papillomavirus infection identification by molecular biology among

asymptomatic women. Rev Saude Publica **36:**95-100.

18. **Kavati EA, Hosoda TM, Szulczewski V, Leão E, Borelli P, Cianciarullo AM.**

2017.  A simple, fast, inexpensive and efficient method for leukocytes separation

with preservation of morphology and cell viability for use in education and research.

In: Microscopy and imaging science: practical approaches to applied research and

education. Editor: Antonio Méndez-Vilas. Publisher: Formatex Research Center,

Badajoz, Spain. Volume **7:**665-670. ISBN-13: 978-84-942134-9-6.

http://www.microscopy7.org/book/665-670.pdf

19. **Fothergill T, Mcmillan NAJ.** 2006. Papillomavirus virus-like particles activate the

PI3-kinase pathway via alpha-6 beta-4 integrin upon binding. J Virol **352:**319-328.

20. **Van Hamme E, Dewerchin HL, Cornelissen E, Verhasselt B, Nauwynck HJ.**

2008. Clathrin- and caveolae-independent entry of feline infectious peritonitis virus

in monocytes depends on dynamin. J Gen Virol **89:**2147-2156.

21. **Doorbar J,  Egawa N, Griffin H, Kranjec C, Murakami I.** 2015. Human

papillomavirus molecular biology and disease association. Rev Med Virol **25:**2-23.

22. **Viarisio D, Muller-Decker K, Accardi R, Robitaille A, Durst M, Beer M, Jansen**

**L, Flechtenmacher C, Bozza M, Harbottle R, Voegele C, Ardin M, Zavadil J,**

**Caldeira S, Gissmann L, Tommasino M.** 2018. Beta HPV38 oncoproteins act with

a hit and- run mechanism in ultraviolet radiation-induced skin carcinogenesis in

mice. PLoS Pathog **14 (1):**e1006783.

23. **Aro K, Nieminena P, Louvanto K, Jakobsson M, Virtanena S, Lehtinen M,**

**Dillner J, Kalliala I.** 2019. Age-specific HPV type distribution in high-grade

cervical disease in screened and unvaccinated women. Gynecol Oncol **154:** 354-359.

24. **Schellenbacher C, Roden RBS, Kirnbauer R.** 2017. Developments in L2-based

human papillomavirus (HPV) vaccines. Virus Res **231:**166-175.

25. **Näsman A, Nordfors C, Holzhauser S, Vlastos A, Tertipis N, Hammar U,**

**Hammarstedt-Nordenvall L, Marklund L, Munck-Wikland E, Ramqvist T,**

**Bottai M, Dalianis T.** 2015. Incidence of human papillomavirus positive tonsillar

and base of tongue carcinoma: a stabilisation of an epidemic of viral induced

carcinoma? Eur J Cancer **51:**55-61.

26. **Combes JD, Clavel C, Dalstein V, Gheit T, Clifford GM, Tommasino M,**

**Franceschi S, St Guily JL, SPLIT study group.** 2018. Human papillomavirus

detection in gargles, tonsil brushings, and frozen tissues in cancer-free patients. Oral

Oncol **82:**34-36.

27. **Gheit T, Rollo F, Brancaccio RN, Robitaille A, Galati L, Giuliani M, Latini A,**

**Pichi B, Benevolo M, Cuenin C, McKay-Chopin S, Pellini R, Cristaudo A,**

**Morrone A, Tommasino M, Donà MG**. 2020. Oral Infection by Mucosal and

Cutaneous Human Papillomaviruses in the Men Who Have Sex with Men from the

OHMAR Study. Viruses **12(8):**E899.

28. **Gruenberg J, van der Goot FG.** 2006. Mechanisms of pathogen entry through the

endosomal compartments. Nat Rev Mol Cell Biol **7:**495-504.

29. **Kinlock BL, Wang Y, Turner TM, Wang C, Liu B.** 2014. Transcytosis of HIV-1

through vaginal epithelial cells is dependent on trafficking to the endocytic recycling

pathway. PLoS ONE **9:**e96760.

30. **Müller M.** 2005. Human Papilloma Viruses: Methods and Protocols. Humana Press

Inc. In: Davy, C.; Doorbar, J. Methods in Molecular Medicine. Totowa, NJ.

**119:**435-446.

31. **Müller M, Gissmann L, Cristiano RJ, Sun XY, Frazer IH, Jenson AB, Alonso**

**A, Zentgraf H, Zhou J.** 1995. Papillomavirus capsid binding and uptake by cells

from different tissues and species. J Virol **69:**948-954.

32. **Roden RB, Kirnbauer R, Jenson AB, Lowy DR, Schiller JT.** 1994a. Interaction

of papillomaviruses with the cell surface. J Virol **68:**7260-7266.

33. **Kieback E, Müller M.** 2006. Factors influencing subcellular localization of the

human papillomavirus L2 minor structural protein. Virology **345:**199-208.

34. **Leder C, Kleinschmidt JA, Wiethe C, Müller M.** 2001. Enhancement of Capsid

Gene Expression: Preparing the Human Papillomavirus Type 16 Major Structural

Gene L1. J Virol **75:**9201-9209.

35. **Campo MS, Jarrett WF, Barron R, O'Neil BW, Smith KT.** 1992. Association of

bovine papillomavirus type 2 and bracken fern with bladder cancer in cattle. Cancer

Res **52:**6898-6904.

36. **Stocco dos Santos RC, Lindsey CJ, Ferraz OP, Pinto JR, Mirandola RS, Benese**

**FJ, Birgel EH, Bragança-Pereira CA, Beçak W.** 1998. Bovine papillomavirus

transmission and chromosomal aberrations: an experimental model. J Gen Virol

**79:**2127-2135.

37. **Tseng CJ, Lin CY, Wang RL, Chen LJ, Chang YL, Hsieh TT, Pao CC.** 1992.

Possible transplacental transmission of human papillomaviruses. Am. J. Obstet

Gynecol **166:**35-40.

38. **Payne D, Tyring SK, Doherty MG, Daya D, Chan TS.** 1993. Detection of human papillomavirus type 16 in plasma cells. Gynecol Oncol **48:**406-412.

39. **Pao CC, Lin SS, Lin CY, Maa JS, Lai CH, Hsieh TT.** 1991. Identification of human papillomavirus DNA sequences in peripheral blood mononuclear cells. Am J Clin Pathol **95:**540-546.

40. **Karas Z, Porebas E.** 1998. HPV sequences in blood of patients with condyloma acuminata. J Invest Dermatol **110:**843-844.

41. **Roden RB, Weissinger EM, Henderson DW, Booy F, Kirnbauer R, Mushinski JF, Lowy DR, Schiller JT.** 1994b. Neutralization of bovine papillomavirus by antibodies to L1 and L2 capsid proteins. J Virol **68:**7570-7574.

42. **Abbas AK, Lichtman AH.** 2005. Cellular and Molecular Immunology. 5 ed. Philadelphia: Elsevier, 580 p.

43. **Trowbridge IS, Omary MB.** 1981. Human cell surface glycoprotein related to cell proliferation is the receptor for transferrin. Proc Natl Acad Sci USA **78:**3039–3043.

44. **Tuma PL, Hubbard AL.** 2003. Transcytosis: crossing cellular barriers. Physiol Rev **83:**871-932.

45. **Kinlock BL, Wang Y, Turner TM, Wang C, Liu B.** 2014. Transcytosis of HIV-1 through vaginal epithelial cells is dependent on trafficking to the endocytic recycling pathway. PLoS One **9:**e96760.

46. **Anderson DJ.** 2014. Modeling mucosal cell-associated HIV type 1 transmission in vitro. J Infect Dis **210:**S648-S653.

47. **Sakajiri T, Yamamura T, Kikuchi T, Yajima H.** 2009. Computational structure models of apo and diferric transferrin-transferrin receptor complexes. Protein J **28:**407-414.

48. **DiGiuseppe S, Bienkowska-Haba M, Guion LG, Sapp M.** 2017. Cruising the cellular highways: how human papillomavirus travels from the surface to the nucleus. Virus Res **231:**1-9.

49. **Day PM, Thompson CD, Lowy DR, Schiller JT.** 2017. Interferon gamma prevents infectious entry of human papillomavirus 16 via an L2-dependent mechanism. J Virol **91:**e00168-17.

50. **Lipovsky A, Erden A, Kanaya E, Zhang W, Crite M, Clinton B, MacMicking, DiMaio D, Schoggins JW, Iwasaki A.** 2017. The cellular endosomal protein stannin inhibits intracellular trafficking of human papillomavirus during virus entry. J Gen Virol **98:**2821-2836.

51. **Garland SM, Kjaer SK, Muñoz N, Block SL, Brown DR, DiNubile MJ, Lindsay BR, Kuter BJ, Perez G, Dominiak-Felden G, Saah AJ, Drury R, Das R, Velicer C.** 2016. Impact and effectiveness of the quadrivalent human papillomavirus vaccine: a systematic review of 10 years of real-world experience, Clin Infect Dis **63:**519–527.

52. **Wright Jr TC, Parvub V, Stoler MH, Kodsi S, Eckert K, Yanson K, Cooper CK.** 2019. HPV infections and cytologic abnormalities in vaccinated women 21–34 years of age: Results from the baseline phase of the Onclarity trial. Gynecol Oncol **153:**259-265.

53. **Lawton B, Heffernan M, Wurtak G, Steben M, Lhaki P, Cram F, Blas M, Hibma M, Adcock A, Stevenson K, Whop L, Brotherton J, Garland SM.** 2020. IPVS Policy Statement addressing the burden of HPV disease for Indigenous peoples. Papillomavirus Res **9:**100191.

54. **Pontelli MC, Castro IA, Martins RB, Veras FP, La Serra L, Nascimento DC,**

**Cardoso RS, Rosales R, LimaTM, Souza JP, Caetité DB, de Lima MHF,**

**Kawahisa JT, Giannini MC, Bonjorno LP, Lopes MIF, Batah SS, Siyuan L,**

**Assad RL, Almeida SCL, Oliveira FR, Benatti MN, Pontes LLF, Santana RC,**

**Vilar FC, Martins MA, Cunha TM, Calado RT, Alves-Filho JC, Zamboni DS,**

**Fabro A, Louzada-Junior P, Oliveira RDR, Cunha FQ, Arruda E.** 2020.

Infection of human lymphomononuclear cells by SARS-CoV-2. bioRxiv preprint

doi: https://doi.org/10.1101/2020.07.28.225912. This version posted August 7, 2020.

55. **Gravitt PE, Winer RL.** 2017. Natural History of HPV Infection across the

Lifespan: Role of Viral Latency. Viruses **9:**e267.

**Figure Legends**

**Figure 1:** Interactions of human leukocytes with VLPs of HPV16, detected by indirect

immunofluorescence. The leukocyte cells were incubated with the VLPs for 4 h. Following

fixation, they were immunostained with anti-HPV16 L1 antibody and then stained with a

secondary antibody conjugated to FITC (green). In all assays, AlexaFluor[®] 594-conjugated

Phalloidin (red) detected the actin cytoskeleton. The internalization of VLPs (green arrows)

is shown in image C, through the sweep cuts on the Z-axis (C) 9.50 µm thick sections and

each section to 1.60 µm. CLSM Zeiss LSM 510 Meta. Magnification: Objective C-

Apochromatic 63xs /1.4 oil. Bar = 5 µm.

**Figure 2:** Interactions of human leukocytes with VLPs of HPV16, detected by indirect

immunofluorescence. The leukocyte cells were incubated with the VLPs for 4 h. Following

fixation, they were immunostained with the conformational anti-HPV16 VLPs and stained

with a secondary antibody conjugated to FITC (green). AlexaFluor[®] 594-conjugated

Phalloidin (red) detected the actin cytoskeleton. The internalization of VLPs (green arrows)

is shown in image C, through Z-axis sweep cuts (C) 7.80 µm thick sections and each

section to 1.30 µm. In (A) we see a structure similar to an endocytic vesicle (white arrow).

CLSM Zeiss LSM 510 Meta. Magnification: Objective C-Apochromatic 63xs /1.4 oil. Bar

= 5 µm.

**Figure 3:** Interactions of human leukocytes with VLPs of HPV16, detected by indirect

immunofluorescence. After fixation, cells were immunostained with anti-HPV16 L1

antibody and then stained with a secondary antibody conjugated to FITC (green). Then

immunostained with anti-HPV16 L2 antibody and revealed by AlexaFluor[®]555 secondary

antibody (red). The internalization of VLPs (yellow arrows) is shown in confocal images

(merge). In (B) structures similar to vacuoles (blue arrow) might be observed. CLSM Zeiss

LSM 510 Meta. Magnification: Objective C-Apochromatic 63xs /1.4 oil. Bar = 5 μm.

**Figure 4:** Identification of human leukocyte interactions with the HPV16 VLPs analyzed

by indirect immunofluorescence. After fixation, cells were immunostained with the

conformational anti-HPV16 VLPs and revealed by a secondary antibody conjugated to

FITC (green). Then, the cells were treated with anti-CD04 (A) and anti-CD08 (B) and

stained with a secondary antibody AlexaFluor[®]546 (red). VLP internalization (green

arrows) is visible on image overlay (merged images). In (B), a structure similar to an

endocytic vesicle is detectable (white arrow). CLSM Zeiss LSM 510 Meta. Magnification:

Objective C-Apochromatic 63xs /1.4 oil. Bar = 5 μm.

**Figure 5:** Identification of human leukocyte interactions with the HPV16 VLPs, analyzed

by indirect immunofluorescence. After fixation, cells were immunostained with the

conformational anti-HPV16 VLPs and revealed by a secondary antibody conjugated to

FITC (green). Then, the cells were treated with anti-CD20 and anti-CD14 and stained with

a secondary antibody AlexaFluor[®]546 (red). VLP internalization (green arrows) is visible

on image overlay (merged images). In (B) we can observe the colocalization of CD14 and

VLPs (orange arrow) and vacuole-like structures (blue arrow). CLSM Zeiss LSM 510

Meta. Magnification: Objective C-Apochromatic 63xs /1.4 oil. Bar = 5 μm.

**Figure 6:** Identification of human leukocyte interactions with the HPV16 VLPs analyzed

by indirect immunofluorescence. After fixation, cells were immunostained with the

conformational anti-HPV16 VLPs and revealed by a secondary antibody conjugated to

FITC (green). Then, the cells were treated with anti-CD20 (C), anti-CD04 (A), anti-CD08

(B), anti-CD14 (D) and AlexaFluor[®]546 revealed by secondary antibody (red). VLP

internalization (green arrows) is shown in the images through Z-axis sweep cuts.  (A) 9.25

µm of thick sections and each section of 0.84 µm; (B) 8.56 µm thick sections and each

section with 0.95 µm; (C) 9.27 µm thick sections and each section with 1.03 µm; (D) 10,94

µm thick sections and each section having 1.40 µm. CLSM Zeiss LSM 510 Meta.

Magnification: Objective C-Apochromatic 63xs /1.4 oil.

**Figure 7:** Colocalization of VLPs of HPV16 and Tf in human leukocytes, detected by

indirect immunofluorescence. The leukocyte cells were incubated with the VLPs and the

conjugated Tf TexasRed[®] (red). Leukocytes were treated with anti-HPV16 L1 antibody,

revealed by a secondary antibody conjugated to FITC (green). In confocality (merge), the

images reveal colocalization of L1 and Tf (yellow arrows). CLSM Zeiss LSM 510 Meta.

Magnification: Objective C-Apochromatic 63xs /1.4 oil. Bar = 5 µm.

**Figure 8:** Colocalization of HPV16 VLPs and Tf in human white blood cells, detected by

indirect immunofluorescence. The leukocyte cells were incubated with the VLPs and the

conjugated Tf TexasRed[®] (red). Leukocytes were immunostained with anti-HPV16 VLP

antibody and revealed by a secondary antibody conjugated to FITC (green). In confocality

(merge), the images reveal colocalization of VLPs and Tf (yellow arrows). CLSM Zeiss

LSM 510 Meta. Magnification: Objective C-Apochromatic 63xs /1.4 oil. Bar = 5 µm.

**Figure 9:** Three-dimensional imaging of leukocyte interactions with the VLPs produced in

this study or the VLP vaccine Gardasil, Tf and TfR; detected by indirect

immunofluorescence. The leukocyte cells were incubated with the VLPs and the conjugated

Tf TexasRed® (red). After fixation, the leukocytes were immunostained with anti-HPV16

VLP antibody and revealed by a secondary antibody, conjugated to FITC (green), then

treated with anti-CD71 and revealed by AlexaFluor®633 (blue). (A) and (B) colocalization

Tf and TfR (purple arrow), (C) and (D) colocalization VLP / Tf / TfR (white arrows), (E)

and (F) weak colocalization VLP / Tf (orange arrows). CLSM Zeiss LSM 510 Meta.

Magnification: Objective C-Apochromatic 63xs /1.4 oil.

**Figure 10:** Immunofluorescence assay detecting the HPV VLPs entry in PBMC, even in

presence of specific biochemical inhibitors of ligand uptake: (A) Sodium azide; (B)

Chlorpromazine; (C) Liquemin; (D) rCTB; (E) Filipin; (F) Nystatin showing the overlay

(merge) of images detected by stain of L1 protein (green), L2 protein (red), nuclei (blue).

The entry of L1 and L2 proteins in cells was not inhibited. CLSM Zeiss LSM 510 Meta.

Magnification: Objective C-Apochromatic 63xs /1.4 oil. Bar = 5 µm.

**Figure 11.** Assay performed with all inhibitors used in Fig.10 (A-F), in association. (A)

Detection of L1 protein (green); L2 protein within cells (red) and  nuclei (blue) in the

overlay (merge) of images; (B-C) Z-axis showing the L1 (B) and L2 (C) proteins within

cells. CLSM Zeiss LSM 510 Meta. Magnification: Objective C-Apochromatic 63xs /1.4 oil.

Bar = 5 µm.

**Tables**

**Table 1:** List of primary antibodies used in immunocytochemistry reactions.

| Antibody | Dilution | Description | Supplier |
|---|---|---|---|
| Anti-L1 Camvir-1 | 1:500 | anti-mouse $IgG_{2a}$, specific for HPV16 | BD Biosciences |
| Anti-L1 conformational | 1:100 | anti-mouse $IgG_{2a}$, specific for HPV16 | Biodesign |
| Anti-L2 | 1:100 | anti-rabbit IgG | Roden, R. MD/USA |
| Transferrin conjugated | 1:60 | TexasRed® (Ex. 595 nm, Em. 615 nm) | Molecular Probes™ |
| Anti-Tf | 1:750 | anti- rabbit IgG | Dako |
| Anti-Ferritin | 1:750 | anti- rabbit IgG | Dako |
| Anti-CD71 | 1:750 | anti-mouse $IgG_1$; Tf receptor | Dako |
| Anti-CD04 | 1:20 | anti-mouse $IgG_{2a}$; recognizes T lymphocytes "helper" | CALTAG™ Laboratories |
| Anti-CD08 | 1:20 | anti-mouse $IgG_{2a}$; recognizes cytotoxic T lymphocytes | CALTAG™ Laboratories |
| Anti-CD14 | 1:20 | anti-mouse $IgG_{2a}$; recognizes monocytes | CALTAG™ Laboratories |
| Anti-CD20 | 1:20 | anti-mouse $IgG_3$; recognizes B lymphocytes | CALTAG™ Laboratories |

836 **Table 2:** List of secondary antibodies used in immunocytochemistry reactions.

| Antibody | Dilution | Description | Ex/Em. (nm) | Supplier |
|---|---|---|---|---|
| FITC | 1:250 | anti- mouse $IgG_{2a}$ | 495/519 | BD Biosciences |
| AlexaFluor®546 | 1:500 | anti- mouse IgG | 556/573 | Molecular Probes™ |
| AlexaFluor®555 | 1:500 | anti- rabbit IgG | 555/565 | Molecular Probes™ |
| AlexaFluor®594 | 1:500 | anti- mouse $IgG_3$ | 590/617 | Molecular Probes™ |
| AlexaFluor®633 | 1:500 | anti- mouse IgG | 632/647 | Molecular Probes™ |
| AlexaFluor®633 | 1:500 | anti- rabbit IgG | 632/647 | Molecular Probes™ |

**Table 3:** Interactions of HPV16 VLPs produced in this study with PMBC (mononuclear

cells) from peripheral blood of healthy female volunteers.

| PMBC | Receptor | Percentage of cells that interacted with VLPs[1] |
|---|---|---|
| Lymphocyte T | CD04 | 52% ± 3 |
| Lymphocyte T | CD08 | 47% ± 2 |
| Lymphocyte B | CD20 | 48% ± 3 |
| Monocyte | CD14 | 23% ± 5 |

[1.] The percentage of cells that interact with HPV16 VLPs was calculated by the number of

cells recognized by anti-CD antibodies, which have internalized the particles. The result

corresponds to the analysis in duplicate, being representative of at least four trials.

**Table 4:** Biochemical inhibitors used to examine their effects on the entry of HPV VLPs in

PBMC.

| Inhibitors | Inhibition |
|---|---|
| Chlorpromazine | Clathrin pathway |
| rCTB (Clostridium toxin B) | Actin-dependent cells process |
| Filipin and Nystatin | Caveolae pathway |
| Liquemin | HSPG pathway |
| Sodium azide | Endocytic pathways |

850 **Figures**

851 **<u>Figure 1</u>**

857     **Figure 2**

870     **Figure 3**

876    **Figure 4**

884    **Figure 5**

890    **Figure 6**

898    **<u>Figure 7</u>**

905 **Figure 8**

922    **Figure 9**

931    **<u>Figure 10</u>**

942     **<u>Figure 11</u>**

**Legend of the Graphical  Abstract:**

Graphical abstract showing *ex vivo* and *in vitro* internalization between VLPs and host cells.

After leukocytes separation from human whole blood, it was performed the identification of human peripheral blood leukocytes in ex vivo interactions with VLPs. The graph shows that of the cells that interacted with VLPs, 52% corresponded to lymphocytes T-CD4, 47% lymphocytes T-CD8, 48% lymphocytes B-CD20, and only 23% of the monocytes CD14 interacted with these particles. However, monocytes apparently internalized larger amounts of particles when compared to lymphocytes.

It is probable that in some T lymphocytes the amount of internalized particles has been imperceptible to the confocal microscope, since the VLPs produced in this research are around 50 nm in diameter. These results lead to two important implications. First, the interaction of VLPs with lymphocytes may result in the activation of these cells and, consequently, increase the population of these circulating cells, this being crucial in the induction of specific immune response.

In the second implication, these lymphocytes would internalize small amounts of virus, insufficient to activate the immune system. Here it is important to note that lymphocytes are cells capable of dividing and it is estimated that the half-life of these inactive cells in humans is of some years. In addition, as it is known, inactive lymphocytes continually re-circulate through the bloodstream and lymphatic vessels.

The percentage of cells that interacted with the HPV16 L1/L2 VLPs was calculated by the

number of cells recognized by the anti-CD antibodies, which internalized these particles.

The result corresponds to the analysis in duplicates, being representative of at least four

tests.

All images are original and cells were processed by Cianciarullo AM et al., at the Butantan

Institute, Sao Paulo – SP, Brazil.

Electron micrographs of human leukocytes, HEK293T and HPV16 L1/L2 VLPs were

obtained in a Zeiss EM109 transmission electron microscope. The blue color of the VLPs

and colored leukocytes were virtually attributed. Leukocyte and HEK293T present

filamentous actin (red) and HPV16 L1/L2 VLPs (green), by fluorescence in a Confocal

Zeiss LSM 510 Meta Microscope.

1000

1001

1002

1003

Graphical abstract showing *ex vivo* and *in vitro* internalization between VLP and host cells

1004