Acute effects of drugs on *C.elegans* movement reveal complex responses and

plasticity

Diversity of Acute Drug Responses in C. elegans

Mark Spensley[1], Samantha Del Borrello[1,2], Djina Pajkic[1] and Andrew G Fraser[1,2]

[1] Donnelly Centre for Cellular and Biomolecular Research, University of

Toronto, Toronto, Ontario, Canada

[2] Department of Molecular Genetics, University of Toronto, Toronto, Ontario,

Canada

Abstract

Many drugs act very rapidly — they can turn on or off their targets within minutes in a whole animal. What are the acute effects of drug treatment and how does an animal respond to these? We developed a simple assay to measure the acute effects of drugs on *C.elegans* movement and examined the effects of a range of compounds including neuroactive drugs, toxins, environmental stresses and novel compounds on worm movement over a time period of 3 hours. We find that many treatments show complex acute responses — a phase of rapid paralysis is followed by one or more recovery phases. The recoveries are not the result of some generic stress response but are specific to the drug e.g. recovery from paralysis due to a neuroactive drug requires neurotransmitter pathways whereas recovery from a metabolic inhibitor requires metabolic changes. Finally, we also find that acute responses can vary greatly across development and that there is extensive and complex natural variation in acute responses. In summary, acute responses are sensitive probes of the ability of biological networks to respond to drug treatment and these responses can reveal the action of unexplored pathways.

Author Summary

Drugs are powerful tools that let us switch on or off key pathways in whole animals and watch the effects. Here we set up a simple assay to measure how drugs affect the movement of the simple nematode *C.elegans* — crucially, we look how those responses

change over time. We find that worms have complex responses to many different drugs

— they go through clear phases of paralysis and recovery. The recovery from the initial

effect of any drug is not due to a generic stress response but is specific to the way each

drug acts. We find for example that worms can recover from paralysis driven by one

neurotransmitter pathway by activating a different neurotransmitter pathway or from

paralysis caused by loss of one metabolic pathway by turning on a second one. These

complex responses show how the basic genetic networks that are needed for normal

behaviour and function can rewire and respond to the effect of many drugs.

Importantly, the responses can vary in many ways — different aged worms or different

individuals can have different responses and capturing how drug responses change over

time is critical to dissect this complexity.

Introduction

Drugs are extremely powerful research tools. Addition of a drug can turn on or off a specific target protein and the effect on the organism can be followed over time. In model organisms, a well-characterized drug response can form the basis for genetic screens to identify the drug target and to find genes that modulate the effect of the drug. In *C.elegans*, for example, genetic screens for mutants with altered drug responses were key to finding the targets of several major anthelmintics (1–4) as well as to identifying core components of conserved neuronal signaling pathways (5,1,6,7).

Many genetic screens for *C.elegans* mutants with altered drug responses have screened for mutants that escape the effects of chronic exposure to drugs. In these screens, populations of mutant worms are typically exposed to drugs for several days (8–13). However, many drugs act very rapidly on worms yet currently relatively little is known about these acute responses of *C.elegans* to drugs. What do acute responses generally look like? Are acute responses the same at all developmental stages? Is there natural variation in acute responses as there is in mutant phenotypes or chronic drug responses? Do multiple drugs with the same long-term effect have the same acute responses or very different ones? What are the genes that affect the acute responses — are they the same or different genes as those that affect long-term drug effects?

Our goal in this study was to examine acute drug responses in *C.elegans* and address at least some of these questions. We developed an image-based method that accurately measures worm movement at high throughput. This allows us to gather

rich information about the acute effects of drugs on worm movement, measuring

both the effect of a range of doses of drugs and how the effects of any drug changes

across time. We examined a wide range of drugs that were previously shown to be

bioactive as well as a number of environmental stresses. We found that many drugs

have complex acute effects on worms indicating that the animal responds rapidly to

the initial drug effect — these acute responses include both transient and sustained

recovery from initial paralysis. We show that the basis for complex acute responses

is drug-specific and that acute responses can differ greatly between different

developmental stages. Finally, we compare acute responses in two isolates and find

that there is substantial variation in acute responses and that this changes with

dose, time and developmental stage. We thus find that measuring acute responses to

drugs can reveal the action of unexplored pathways and that acute responses vary

extensively and at many levels between different individuals. The ability to measure

acute responses at high throughput will allow both drug screens and genetic screens

to uncover the underlying molecular basis for these complex responses.

Results

Development of a high throughput image-based movement assay

We wanted to measure the acute effects of drugs on worms over the time period of

minutes to hours. We focused on movement as a key phenotype — a very wide

range of compounds affect movement including toxins, neuroactive compounds and

environmental stresses. There are a wide variety of assays to measure the acute

effects of drugs on worm movement, with different levels of throughput and

resolution, and different requirements for specialized equipment such as worm

trackers or microfluidic devices (14–19). Here we focused on creating a simple

assay for worm movement that allows us to follow how worms respond to drug

treatment over time periods ranging from minutes to hours. The assay is simple,

automated, gives quantitative data and has sufficient throughput to efficiently

analyse responses to thousands of drugs.

In outline, our assay measures the movement of worms in 96-well format in liquid.

Worms are continuously bathed in buffer containing the compound and there are

multiple worms in each well (e.g. ~100 L1 larvae). We capture two successive

images of each well, separated by a 500 millisecond interval — by comparing these

two images we identify the extent of worm movement at any particular time point

or drug concentration (see Fig 1A for details) and calculate a fractional mobility

score (FMS) for each well at any individual time point. Worm movement can be

measured at many time points to provide a time-course of the effect of a given drug

concentration on worm movement (Fig 1B, right), at a range of drug concentrations

to yield a dose-response curve (Fig 1B, lower) or (since the throughput is high

enough) to cover both dose and time responses. The assay can thus be used to give a

rich quantitative measurement of the effect of any drug on worm movement.

To validate our assay, and in particular to ensure that detected effects on movement

are specific to the targeted pathway, we examined the response of adult worms to

aldicarb (Ald), a very well characterised neuro-active compound (reviewed in depth

in (20)). Acetylcholine (ACh) is a core neurotransmitter that is critical for normal

movement (21). Following release of ACh at synapses, ACh is rapidly metabolised by

acetylcholinesterases to turn off ACh responses. Aldicarb is a carbamate

acetylcholinesterase inhibitor — Ald thus increases the local concentration of ACh at

the synapse or neuromuscular junction by reducing the rate of degradation of

endogenous released ACh (22). Aldi is known to cause paralysis in *C. elegans*, via the

sustained activation of ionotropic acetylcholine receptors (nAChRs) and ACh

signalling and the effects of both Ald and ACh have been extremely well studied

(reviewed in (20)). We thus wanted to validate our assay by testing whether we can

detect the well-established effects of Ald to induce paralysis and whether any Ald-

induced paralysis is mediated by the well-characterised nAChR subunits, confirming

that the effects we see are specific.

Consistent with previous studies, we found that Ald treatment results in paralysis of

adult worms. Both the extent of paralysis and the rate of onset were dose dependent

(Fig 2A). Aldicarb-induced paralysis is mediated by activation of nAChRs in body

wall muscles; nAChRs are ligand gated ion channels composed of five subunits (23–

26). Several of these subunits are known to be required for ACh-driven paralysis

including UNC-38, UNC-63 and LEV-8. (23–25) To confirm that the effects of drugs

on movement that we detect in our assay are specific to the activation of known

AChRs (and not due to some kind of non-specific toxicity for example), we tested the

response of mutant worms lacking functional copies of each of these subunits. As

expected from previous studies, adult worms homozygous for loss-of-function

mutations in *unc-38*, *unc-63*, *lev-8*, *lev-1* or *unc-29* are markedly less sensitive (Fig

2B, Fig S1). We also find that mecamylamine, a non-competitive nAChR antagonist,

affects the dose-response relationship of aldicarb: in the presence of 600 μM

mecamylamine, the EC50 of aldicarb shifts from 0.76 mM to 1.6 mM, with a slight

decrease in maximal response (Fig 2C). The effects of Ald on worm movement that

we measure in our assay are thus specific to the action of elevated ACh on nAChRs

and not some non-specific effect on movement.

We note that while we focus here on the use of this assay to measure the effects of

drugs on worm movement, it can also be used to measure the effect of genetic

perturbations on movement directly. To illustrate this, we used RNA-mediated

interference (RNAi) to knock down 7 genes that are known to be required for wild-

type movement and measured movement after RNAi. We found robust RNAi

phenotypes for all 7 genes (Fig 2D) showing that our movement assay is applicable

to both genetic and drug based screens. Thus, we expect our method to be

applicable in a range of pharmacological, genetic, and RNAi experiments.

We have thus developed a simple method that allows us to measure the acute effects

of drug treatments on worm movement at high throughput. We showed that our

assay recapitulates the known effects of the well-characterised neuroactive drug Ald

and that this is indeed due to its known mode of action through the activation of

specific receptors. Crucially, our assay allows us to measure both dose responses

and time-resolved responses to drugs. In the rest of this paper we demonstrate that

time-resolved profiles of drug responses can yield new insights into *C. elegans*

biology.

Acute responses to drugs are often complex

We established a simple assay to measure acute responses of worms to drugs over time periods of minutes to hours. We wanted first to use this to explore acute responses in general — what do these look like for a variety of drugs and treatments? We exposed *C.elegans* L1 larvae to a wide range of different compounds and treatments including compounds that inhibit core essential machineries such as the electron transport chain (inhibition by potassium cyanide, KCN, shown in Fig 3A) or the ribosome (effect of cycloheximide in Fig 3B), environmental stresses such as high salt conditions (NaCl shown in Fig 3C), known nematicides (abamectin shown in Fig 3D), as well as compounds that affect movement via altered neurotransmission (e.g. aldicarb Fig 2A). In each case, worms were treated and their movement examined over a time course of 3 hours — at least 3 biological replicates were done for each treatment.

In addition to this panel of characterized compounds, we also tested a range of novel compounds that had been identified as affecting worm growth in chronic exposure, population growth assays (13). We examined the effects of 170 of these on the movement of L1 larvae and find that many (53/170) also have acute effects on worm movement. Crucially, an unexpectedly high proportion (31 of 53 with acute effects; 58%) of drugs tested show complex time-resolved responses illustrated in Fig 3E-G, including either sustained or transient recovery from initial drug effects.

Having looked at acute responses to across a range of different treatments, we found complex acute responses across every type of compound, including neuro-active

compounds, environmental stresses, and toxins that target essential genes. These

responses can be very complex — for example, the novel compound shown in Fig 3G

has two distinct phases of recovery and several structurally related compounds

have similarly complex responses (data not shown). We note that every one of these

complex drug responses can be the basis either for a genetic screen for mutants

with altered responses or for drug screens for compounds that modulate the

response. This complexity of many acute drug responses underlines the need to be

able to measure time-resolved acute effects of drugs on movement.

The complex effects of many compounds over time suggest that the worms are

responding to many drug treatments in some way — for example, several drugs

cause an initial rapid paralysis followed by a recovery phase where the worms begin

to move again at almost normal rates. What underlies this recovery? Is it some

generic stress or xenobiotic response or does the worm respond in specific ways to

each compound? To gain some insights into this we focused on the acute responses

to two well-characterized compounds, Aldicarb and cyanide.

Agonists of acetylcholine signalling have complex effects on worm movement

that change across development

As shown in Fig 2A treatment of adult animals with Ald results in a rapid reduction

in movement and sustained paralysis. Interestingly, we found that L1 animals

behave completely differently— Ald induces rapid paralysis of L1 worms but this is

followed by a more gradual recovery (Fig 4A; S2). To further examine the Ald

response across development, we tested all developmental stages and found the

response is qualitatively very different across development (Fig 4A). L1 and L2

stage larvae show an initial phase where Ald treatment results in reduced

movement, followed by a recovery phase during which paralysis is gradually

relieved. As worms progress through the four larval stages, the initial paralysis

remains similar but this recovery is reduced, becoming minimal by adulthood. What

underlies this recovery in L1 and L2 animals?

To further explor the recovery of L1s from paralysis, we asked whether this is

specific to Ald or whether other drugs that affect ACh signaling show similar acute

responses. Ald affects ACh signalling by preventing the degradation of ACh — it does

not activate ACh receptors directly nor does its action have any specificity for any

specific sub-type of ACh receptor. Nicotine (Nic) activates ACh signaling in a

different way — it is a direct agonist of specific ACh receptors, the nicotinic ligand-

gated ion channels. We compared their acute responses and find that both Ald and

Nic show a rapid initial paralysis followed by a recovery of movement (Fig 4B).

While these two responses have superficial similarities, there is a crucial difference

in the recovery phase. For Nic, while L1 worms recover from the reduction of

movement caused by low doses of Nic, increasing concentrations of Nic result in

greater paralysis and less recovery until at high Nic doses there is no appreciable

recovery (Fig 4D). For Ald, we see the precise opposite — there is no recovery at

low doses of Ald, but recovery increases with increasing Ald (Fig 4C). This suggests

that the recovery from Ald and Nic-induced paralysis is fundamentally different and

that Ald is somehow driving recovery since the more Ald we add, the stronger the

recovery. How could Ald be acting and how could this be different to Nic?

ACh is known to activate two completely different classes of receptor: ligand-gated

ion channels (nAChRs) and muscarinic ACh receptors (mAChRs). nAChRs act rapidly

and are stimulated directly by Nic and indirectly by Ald via its action to increase

ACh. mAChRs have much more complex G-protein coupled signaling — crucially,

they are not activated by Nic but are also activated following Ald treatment. We

hypothesized that the Ald recovery phase might be mediated by mAChRs and test

this below.

Recovery from rapid paralysis induced by Aldicarb occurs via muscarinic receptor

signaling

To test whether recovery from Ald-induced paralysis might be due to the activation

of mAChRs, we examined whether a mAChR antagonist, atropine (Atr), could block

the recovery phase of the Ald response. This is indeed the case (Fig 5A) suggesting

that recovery is the result of an activation of mAChR signaling following increased

ACh levels after Ald treatment. The *C. elegans* genome encodes three muscarinic

acetylcholine receptors, GAR-1, GAR-2 and GAR-3 (27–29) — we obtained deletion

mutants for each of these receptors and tested whether the effect Ald was altered in

these mutant strains. The Ald response is clearly different in the *gar-3(gk305)*

mutant strain: while the paralysis phase appears very similar to wild-type, the

recovery phase is strongly suppressed in the *gar-3(gk305)* mutant, although not

completely abolished (Fig 5B). We note that mutations in *gar-1* and *gar-2* also have

effects on recovery but these are much weaker and we suggest that *gar-3* is the key

mAChR that drives recovery from nAChR stimulation in L1 animals.

We thus propose that Ald causes a complex acute response in L1 worms because the

increased ACh levels have effects on two separate neurosignalling pathways. The

first phase of the acute response is a rapid reduction in movement due to activation

of nAChRs. The second phase is a slower recovery due to activation of mAChRs

which somehow relieve the nAChR-driven paralysis. To further test this, we

examined whether we can drive recovery from Nic-induced paralysis with agonists

of mAChRs. Nic can only stimulate nAChRs and has no activity on mAChRs (30).

Consistent with our model, we find that we can induce recovery from Nic-induced

paralysis either by adding Ald (Fig 5C) or the mAChR agonists, Oxotremorine M

(OxoM) or Arecoline (Are) (Fig 5D; supplementary S3). To confirm that the same

mAChR requirement apply to OxoM-induced recovery as in Ald responses, we tested

the ability of each *gar* mutant to recover from Nic-induced paralysis in the presence

of OxoM (Fig 5E). We found that *gar-3* is indeed the primary mAChR driving OxoM-

induced recovery. Our data suggest that mAChR signalling can drive recovery from

paralysis induced by sustained nAChR activation. Finally, we examined whether

changes in mAChR signalling might underlie the difference in the ability of L1 and

adults to recover from Ald paralysis. We find that stimulation of mAChR signalling

by OxoM in adults cannot override the paralysis response (Fig 5F) and thus, changes

in the mAChR signalling machinery might indeed account for the different responses

across development.

Our data thus suggest that a model in which there is some kind of physiological

cross talk between two types of ACh receptor, nAChRs and mAChRs. This model is

summarized in Fig 6, and suggests that i) sustained activation of nAChRs alone

results in long-term paralysis, ii) sustained activation of mAChRs alone has little

effect on movement, but iii) dual activation of both nAChRs and the mAChR GAR-3,

such as with Ald alone, or Nic + OxoM results in an initial phase of nAChR-driven

paralysis followed by a slower mAChR-driven recovery. We conclude that the

complex acute response of L1 worms to Ald is highly specific to the drug used and is

not a generic stress or xenobiotic response. To examine whether this is true for

other drugs, we next looked at the effects of a very different drug, KCN, on worm

movement.

C.elegans has a complex response to treatment with cyanide.

Cyanide (KCN from here on) has a very well-characterized mode of action: it inhibits

aerobic metabolism by blocking the mitochondrial electron transport chain (ETC)

through binding to complex IV (reviewed in (31)). KCN is toxic to *C.elegans*

following prolonged exposure — here we examined the acute responses of worms

to KCN exposure.

We immediately noticed that the dose response at early time points were very

unusual — while intermediate doses of KCN caused almost complete paralysis

movement, high doses appeared to have little effect (Fig 7A). This was most

pronounced in DRCs between 15-30minutes — by ~60 minutes the DRCs appeared

much more normal, with increased effects being seen with increased dose.

How can the effect of low concentrations of KCN be greater than the effect of high

concentrations? We reasoned that this may be due to rewiring of metabolic

pathways — worms might respond to high KCN concentrations by rapidly switching

to use alternative metabolic pathways for generating ATP. An obvious candidate for

such a pathway was anaerobic glycolysis which is the major pathway for ATP

generation in anaerobic conditions in many animals — indeed we found that

addition of 2-deoxy-D-glucose, an inhibitor of phosphoglucose isomerase, a critical

enzyme in anaerobic glycolysis, caused a profound change in KCN DRCs. Worms can

no longer continue moving in high doses of KCN when 2DG is also present — instead

we find that increased concentrations of KCN result in decreased movement at all

time points (Fig 7A). This suggests that the unusual DRCs seen for KCN alone were

due to a shift to utilization of anaerobic glycolysis at early timepoints in high KCN

(Fig 7B).

This underscores the importance of measuring drug responses across time — if we

had only measured KCN DRCs at 80 minutes, they would have looked completely

unremarkable. It also confirms that the way that worms respond to acute drug

treatment is highly specific to the drug itself — while the aldicarb response involves

crosstalk between nAChRs and mAChRs, the KCN response involves metabolic

switches. The rapid responses of worms to compounds thus do not appear to be

generic stress or xenobiotic responses but are specific responses to the specific drug

effects. They reveal how the organism can rewire and adapt to sudden inhibition of

a key pathway.

Natural variation in acute drug response is very complex

Drug responses are known to vary within natural populations. In *C.elegans*, natural

variation in drug responses has been seen for several drugs including abamectin

and etoposide (32,33). In a previous study we compared the effects of KCN on two

natural isolates, the N2 and CB4856 isolate (34). These differ by around

1SNP/800bp (35), a similar degree of variation as that between two human

genomes. In our previous study we simply examined one time point (90 minutes

after drug treatment) and only looked at a single larval stage, the L1 stage — we

wanted to expand this analysis to compare the effects of KCN on N2 and CB4856

across development. We decided to examine how a range of concentrations of KCN

affect movement of all 4 larval stages of either N2 or CB4856 and in each case to

examine how this response evolves over 3hr (Fig 8).

The results are complex and striking. In keeping with our previous study (34), we

find that N2 are more affected by KCN at the L1 stage. However, we find the exact

reverse at the L4 stage — CB4856 are substantially more sensitive here. At the L3

stage, the picture is even more complex — while KCN appears to affect CB4856

more rapidly (CB4856 L3 animals are more severely affected at early timepoints),

the same dose of KCN has a more severe effect on N2 later timepoints. Taken

together, these data show clearly that comparisons of drug sensitivity in natural

populations are complex. It is impossible to make a blanket statement that isolate A

is 'more sensitive' than isolate B — assessing the effect at a different dose, a

different timepoint or a different developmental stage could give precisely the

reverse conclusion. These data underscore the need to measure drug responses as

comprehensively as possible before making general conclusions — without

measuring a range of doses and capturing these responses across time, we would

have missed much of this complexity.

341 Discussion

Drugs have extremely rapid effects on their targets — they can switch on or off their

target within minutes even in a whole animal. What is the effect on the animal and

how does it respond? We looked at the effects of drugs on *C.elegans* movement to

measure acute responses to drugs in a whole animal and find several unexpected

results that have important implications for drug screens in animals.

First, we found that around half of all the bioactive compounds that we found to

affect *C.elegans* movement show a complex response: the worms are rapidly

affected, but after an early phase of near-paralysis, the animals recover movement

— this recovery may be transient or longer lived. In some cases, like the responses

to compound shown in Fig 3E-F, we see multiple phases of paralysis and recovery

over time — studying these complex acute responses of worms over time will drive

exciting new insights. That worms can recover from the rapid effects of many drugs

indicates that the worms are responding in some way. We did not expect to see this

kind of recovery so frequently and to such a wide range of drugs and treatments, or

responses of such complexity — that surprise drove this study.

Second, at least in the examples we examined, the mechanisms underlying complex

acute responses appear to be drug specific and not due to a generic stress or

xenobiotic response. For example, recovery from paralysis due to signaling by one

set of acetylcholine receptors (AChRs) is due to signaling from a different set of

AChRs; recovery from paralysis due to inhibition of aerobic respiration is driven by

the use of anaerobic pathways. We see complex acute responses to a wide range of

drugs and thus worms can respond to many different acute perturbations of their

core networks by rewiring and modulating the activity of specific pathways. This

suggests a tremendous complexity of homeostatic mechanisms and sensors and this

molecular biology can be revealed through studying acute drug responses. For

example, worms recover from paralysis induced by activation of nAChRs through

signaling from mAChRs — this cross talk isn't evident in chronic drug assays but can

be easily measured using acute responses. Acute responses are thus excellent

reporters for the rewiring of genetic networks in response to the effects of a drug.

Finally, we find that many factors affect acute drug responses. The acute response to

the same drug can be very different at different developmental stages, at different

doses of drug, or in different individuals — measuring only a single dose, a single

time point, or a single developmental stage can greatly affect conclusions. For

example, we found that L1 larvae recover readily from Aldicarb-induced paralysis

but adults cannot recover. The recovery of L1s from Ald-induced paralysis is highly

dose-dependent — L1s do not recover at 2mM Ald but recover readily at the higher

concentration of 6mM (Fig 4A). This kind of complexity means one cannot simply

say '*C.elegans* can recover from Ald paralysis' or '*C.elegans* cannot recover from Ald

paralysis'. We see a similar complexity when we look at natural variation in acute

responses. We compared the sensitivity to KCN of two natural isolates of *C.elegans,*

N2 and CB4856 and find a complex picture — while N2 is more sensitive at the L1

stage, CB4856 is more sensitive at the L4 stage. Again, we cannot make a blanket

statement of 'N2 is more sensitive than CB4856' — dose, time point, and

developmental stage drive major differences in the drug responses and these factors

are often neglected. For example a GWAS to identify QTLs that affect 'drug

sensitivity' would likely find different variants depending on drug dose and

developmental stage assayed. Crucially, this means that to really understand acute

responses, we need to measure them over a range of doses and for each

developmental stage. The simple assay we report here allows us to gather this kind

of rich data.

Taken together, our results show that worms have complex acute responses to a

wide variety of drugs and that these complex responses are due to the specific

rewiring of the pathways underlying worm biology. Acute responses are sensitive

readouts of the pathways that are required after rewiring — some of these are

largely unexplored. For example, activation of mAChR signaling alone only has very

subtle effects on worm biology (36–40) but acute responses reveal a clear effect of

mAChR signaling in driving recovery from nAChR-induced paralysis. Genetic screens

to find mutants with altered acute drug responses can thus dissect otherwise

intractable pathways and acute responses can also be the basis for sensitive drug

screens. Understanding the genetics that underpin complex acute responses like

those shown in Fig 3 will reveal novel insights into how genetic networks rewire

and change following insult.

Importantly, we also show that acute drug responses vary in many ways — different

drug doses, different developmental stages, and different genetic backgrounds can

give very different acute responses. Drawing broad conclusions about drug

responses such as '*C.elegans* recovers from Aldicarb-induced paralysis' or 'the N2

isolate is more sensitive to KCN than the CB4856' is thus very difficult — different

doses or different developmental stages often give completely different results.

Studies on drugs in whole animals need to be aware of this complexity and interpret

results with care. We believe that high throughput quantitative assays for drug

responses of the kind we describe here are key — they allow researchers to build up

a rich view of a drug response rather than focus on an arbitrary dose, time point or

developmental stage. Studying the full complexity of acute responses opens up a

whole new set of whole animal phenotypes for drug and genetic screens and will

reveal how animals rapidly rewire their genetic networks in response to drugs,

toxins, and environmental stresses.

Materials and Methods

Worm Maintenance and Strains

All worm stocks were maintained at 20ºC on NGM agar plates seeded with *E.coli*

strain OP50 as described elsewhere (41). In addition to the classical laboratory

strain N2, we describe work using strains *unc-38* (*e264)*, *lev-8* (*x15*), *lev-1* (*e211*),

unc-63 (*x26*), *unc-29* (*e1072*), *acr-8* (*ok1240*), *acr-16* (*ok789*), *gar-1* (*ok755*), *gar-2*

(*ok520*) and *gar-3* (*gk305*). All strains were acquired from the Caenorhabditis

Genetics Center.

To prepare L1 worms, animals were washed from the agar plates with M9 buffer

(41) and rinsed in one buffer change of M9. L1s were then isolated by filtration

though 11 μm nylon mesh filter plates (Milipore Multiscreen). L1 larvae for

immediate use in drug response assays were diluted to approximately 1.2 worms

per microlitre in either M9 buffer or modified NGM buffer (50 mM NaCl, 1 mM $CaCl_2$,

1mM $MgSO_4$, 200 mM KH2PO4, 50 mM K_2HPO4).

To assay drug response over different developmental stages, worm populations

were first developmentally synchronized by filtration of L1s, as described above.

Filtered L1s were returned to fresh NGM agar OP50 plates and incubated at 20ºC

until they had reached the required developmental stage (L1: no further incubation,

L2: 18 hr, L3: 32 hr, L4: 42 hr, young adult: 54 hrs). Worms were then washed from

the plates and rinsed as above. To further synchronize developmental stages, worms

were dispensed into assay wells using a COPAS Biosort machine (Union Biometrica),

which can selectively dispense individual worms on the developmentally-correlated

criteria of size and optical density (L1 worms: 100 / well, L2: 80 / well, L3:, 50 /

well, L4: 30 / well, young adult : 30 / well). Sorting parameters for each stage were

determined empirically, as per the manufacturer's instructions. The volume of liquid

dispensed into the wells was determined and adjusted to 100 μl with buffer.

Drug preparation

Solutions of potassium cyanide (Sigma Aldrich 60178-25G), sodium chloride

(BioShop SOD002.205), cycloheximide (Sigma Aldrich C7698-1G) abamectin (Sigma

Aldrich 31732) and 2-dexoxy-D-glucose (Sigma Aldrich D8375) were prepared fresh

to twice the working concentration in M9 buffer, 1.6% DMSO. Stock solutions of

aldicarb (Sigma Aldrich, 33386) were prepared to 1 M in DMSO. Solutions of

levamisole hydrochloride (Acros Organics, 16595-80-5), oxotremorine M (Sigma

Aldrich, O100) and mecamylamine hydrochloride (Sigma Aldrich, M9020) were

prepared to 100 mM in water. These solutions were divided into single use aliquots

and frozen. Stocks of arecoline hydrobromide (Acros Organics, 250130050) and

atropine sulphate monohydrate (Sigma Aldrich, A0257) were prepared fresh on the

day of use in M9 buffer. (-)-Nicotine (Sigma Aldrich, N3876) was stored frozen,

undiluted in single use aliquots.

Preparation of drug response assays:

Assays were assembled in a final volume of 200 µl in flat-bottomed, polystyrene 96-

well plates (Corning 3997) by combining 100 µl of worms in buffer with an equal

volume of drug solution. To control for any confounding effects of drug solvent, all

experiments were prepared to contain DMSO at 0.8 % v/v, regardless of whether

the drug solution was prepared in DMSO or water. All assays were assembled in M9

buffer apart from the screen for compounds that modify the response to aldicarb,

which were assembled in modified NGM, which we found to cause fewer problems

of drug precipitation. After assembling the assay, plates were sealed with

transparent, self-adhesive films before imaging. The point at which worms and

drugs were combined marked the start of the assay.

Preparation of RNAi assays

RNAi-by-feeding was performed as previously described (42). After 4 days, RNAi-

treated and mock-treated worms were filtered to purify L1 worms and washed in

two buffer changes of M9. Purified L1 samples were diluted to approximately 1.2

worms per microlitre and 100 μl samples were transferred to microtitre plates for

use in mobility assays.

Image acquisition, analysis and calculation of fractional mobility score:

Images were acquired using a Nikon Ti Eclipse inverted microscope with 2x

objective lens and DS-Qi1Mc camera. Image capture was automated in Nikon NIS-

Elements AR software to capture two images of each well, separated by a 500 ms

interval. This procedure was run in a 5 minute loop for the 3 hour duration of the

experiments. A sample NIS-elements script is available as supplementary

information.

Since our microscope control software stores files in a proprietary image stack

format (Nikon ND2), we began our image analysis with a file format conversion

process to convert nd2 files to a collection of 8-bit TIFF images, with structured

filenames identifying the timepoint and well coordinates of each image. The first

step of this process, defined in the Python file *export_nd2.py*, converted the ND2 file

to a series of TIFF files, using the Open Microscopy Environment Bio-Formats tool

(43). This conversion yields a series of 12-bit TIFF files with names encoding the

timepoint and order of acquisition of each image. The second step of the conversion

process invoked the ImageMagick package to convert these files from 12-bit to 8-bit

TIFF, modify filenames to include the well location, rather than order of acquisition

and group files for each timepoint into separate subfolders. This file structure forms

the input for the image analysis script.

Our image analysis pipeline was prepared with the python programming language

using the scikit-image library (44). All images were first processed with a Gaussian

blur to reduce noise levels. Further image analysis consisted of two sets of processes

to find moving and non-background, non-moving parts of the images.

To identify non-background, non-moving pixels, images were processed using a

Sobel filter, which effectively produces the first derivative of the image, thus edges

are highlighted and sharper edges appear brighter than shallow edges. An adaptive

threshold was applied to the Sobel images to remove the background and yield a

binary image. These binary images contained the outlines of the well and any

objects, such as worms, within the well. Binary images were further processed with

two iterations of binary closing to fill the pixels between sufficiently close edges.

This process was hand-tuned to fill in the area occupied by worms but minimize

overfilling into the areas between worms. A circular binary mask was applied to

remove the edges of the wells from each binary image and one of these images was

designated as a reference image for each well (Figure 1A, Image B).

Pixels associated with movement were identified by calculating the absolute

difference of the two consecutive images of a well (Figure 1A, Image A). A threshold

was applied to the difference image to create a binary image (Figure 1A, Image B).

This binary image contained the moving pixels from both parent images, showing

double the number of movement-associated pixels. To correct for this, we multiplied

the binary difference image by the reference image described above revealing only

the non-background, movement associated pixels from a single frame (Figure 1A,

Image C). Finally to identify pixels that did not move between wells, the reference

image was multiplied by the complement of Image C (Figure 1A, Image D).

A fractional mobility score was calculated as the ratio of movement-associated

pixels to total non-background pixels. This process is contained in the Python script

quantify_movement.py. A second Python script, *quantify_movement_adult.py*, was

used to analyse images of adult worms. This script contains an additional step to

mask out objects below a certain number of pixels in size, to remove eggs from the

analysis, since some chemicals stimulate egg-laying.

All our python scripts and an example dataset are available in our repository,

https://github.com/fraser-lab-UofT/acute_assay

Data analysis and curve-fitting:

All drug response data shown were normalised by dividing by the fractional

mobility score of drug-free control wells for each time point. Fitted lines were

determined by least squares minimisation of either a three-parameter logistic

function:

531
$$f(x) = A + \frac{1 - A}{1 + e^{-B(M-x)}}$$

532   or an exponential function:

533
$$f(x) = Me^{-x} + A$$

Where *x* denotes drug concentration and *M*, *A* and *B* denote parameters to be

optimized. The choice of function was determined by the range of the dataset.

Proportional recovery scores were calculated using the formula:

$$\frac{(FMS\ after\ recovery - FMS\ at\ maximum\ paralysis)}{1 - FMS\ at\ maximum\ paralysis}$$

The value of FMS after recovery was calculated as the mean FMS over the period

between 150 and 180 minutes. Maximum paralysis was determined as the minimum

FMS value over the three hour course of the experiment.

References

1.   Lewis JA, Wu CH, Berg H, Levine JH. The genetics of levamisole resistance in

the nematode Caenorhabditis elegans. Genetics. 1980 Aug;95(4):905–28.

2.   Driscoll M, Dean E, Reilly E, Bergholz E, Chalfie M. Genetic and molecular

analysis of a Caenorhabditis elegans beta-tubulin that conveys benzimidazole

sensitivity. J Cell Biol. 1989 Dec;109(6 Pt 1):2993–3003.

3.   Dent JA, Smith MM, Vassilatis DK, Avery L. The genetics of ivermectin

resistance in Caenorhabditis elegans. Proc Natl Acad Sci U S A. 2000 Mar

14;97(6):2674–9.

4.   Jones AK, Buckingham SD, Sattelle DB. Chemistry-to-gene screens in

Caenorhabditis elegans. Nat Rev Drug Discov. 2005 Apr;4(4):321–30.

5.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974

May;77(1):71–94.

6.   Nguyen M, Alfonso A, Johnson CD, Rand JB. Caenorhabditis elegans mutants

resistant to inhibitors of acetylcholinesterase. Genetics. 1995;140(2):527–35.

7.   Miller KG, Alfonso A, Nguyen M, Crowell JA, Johnson CD, Rand JB. A genetic

selection for Caenorhabditis elegans synaptic transmission mutants. Proc Natl

Acad Sci U S A. 1996 Oct 29;93(22):12593–8.

8.   Rogalski TM, Golomb M, Riddle DL. Mutant Caenorhabditis elegans RNA

polymerase II with a 20,000-fold reduced sensitivity to alpha-amanitin.

Genetics. 1990 Dec;126(4):889–98.

9.   Kwok TCY, Ricker N, Fraser R, Chan AW, Burns A, Stanley EF, et al. A small-

molecule screen in C. elegans yields a new calcium channel antagonist. Nature.

2006 May 4;441(7089):91–5.

10. Burns AR, Kwok TCY, Howard A, Houston E, Johanson K, Chan A, et al. High-

throughput screening of small molecules for bioactivity and target

identification in Caenorhabditis elegans. Nat Protoc. 2006 Nov;1(4):1906–14.

11. Kwok TCY, Hui K, Kostelecki W, Ricker N, Selman G, Feng Z-P, et al. A Genetic

Screen for Dihydropyridine (DHP)-Resistant Worms Reveals New Residues

Required for DHP-Blockage of Mammalian Calcium Channels. Frankel WN,

editor. PLoS Genet. 2008 May 9;4(5):e1000067.

12. Luciani GM, Magomedova L, Puckrin R, Urbanus ML, Wallace IM, Giaever G, et

al. Dafadine inhibits DAF-9 to promote dauer formation and longevity of

Caenorhabditis elegans. Nat Chem Biol. 2011 Dec 6;7(12):891–3.

13. Burns AR, Luciani GM, Musso G, Bagg R, Yeo M, Zhang Y, et al. Caenorhabditis

elegans is a useful model for anthelmintic discovery. Nat Commun. 2015 Jun

25;6:7485.

14.   Buckingham SD, Sattelle DB. Fast, automated measurement of nematode

swimming (thrashing) without morphometry. BMC Neurosci. 2009 Jul

20;10(1):84.

15.   Brown AEX, Yemini EI, Grundy LJ, Jucikas T, Schafer WR. A dictionary of

behavioral motifs reveals clusters of genes affecting Caenorhabditis elegans

locomotion. Proc Natl Acad Sci. 2013 Jan 8;110(2):791–6.

16.   Yemini E, Kerr RA, Schafer WR. Tracking Movement Behavior of Multiple

Worms on Food. Cold Spring Harb Protoc. 2011 Dec

1;2011(12):pdb.prot067025.

17.   Yemini E, Jucikas T, Grundy LJ, Brown AEX, Schafer WR. A database of

Caenorhabditis elegans behavioral phenotypes. Nat Methods. 2013 Sep

14;10(9):877–9.

18.   Larsch J, Ventimiglia D, Bargmann CI, Albrecht DR. High-throughput imaging

of neuronal activity in Caenorhabditis elegans. Proc Natl Acad Sci. 2013 Nov

5;110(45):E4266–73.

19.   Swierczek NA, Giles AC, Rankin CH, Kerr RA. High-throughput behavioral

analysis in C. elegans. Nat Methods. 2011;8(7):592-U112.

20.   Rand J. Acetylcholine. The C. elegans Research Community, editor.

WormBook. 2007. 1-21 p.

21.   Richmond JE, Jorgensen EM. One GABA and two acetylcholine receptors

function at the. Nat Neurosci. 1999;2(9):791–8.

22.   Opperman CH, Chang S. Effects of Aldicarb and Fenamiphos on

Acetycholinesterase and Motility of Caenorhabditis elegans. J Nematol. 1991

Jan;23(1):20–7.

23.   Fleming JT, Squire MD, Barnes TM, Tornoe C, Matsuda K, Ahnn J, et al.

604   Caenorhabditis elegans levamisole resistance genes lev-1, unc-29, and unc-38

605   encode functional nicotinic acetylcholine receptor subunits. J Neurosci. 1997

606   Aug 1;17(15):5843–57.

607   24.   Culetto E, Baylis HA, Richmond JE, Jones AK, Fleming JT, Squire MD, et al. The

608   Caenorhabditis elegans unc-63 gene encodes a levamisole-sensitive nicotinic

609   acetylcholine receptor alpha subunit. J Biol Chem. 2004 Oct 8;279(41):42476–

610   83.

611   25.   Towers PR, Edwards B, Richmond JE, Sattelle DB. The Caenorhabditis elegans

612   lev-8 gene encodes a novel type of nicotinic acetylcholine receptor alpha

613   subunit. J Neurochem. 2005 Apr;93(1):1–9.

614   26.   Boulin T, Gielen M, Richmond JE, Williams DC, Paoletti P, Bessereau J-L. Eight

615   genes are required for functional reconstitution of the Caenorhabditis elegans

616   levamisole-sensitive acetylcholine receptor. Proc Natl Acad Sci.

617   2008;105(47):18590–5.

618   27.   Lee YS, Park YS, Chang DJ, Hwang JM, Min CK, Kaang BK, et al. Cloning and

619   expression of a G protein-linked acetylcholine receptor from Caenorhabditis

620   elegans. J Neurochem. 1999 Jan;72(1):58–65.

621   28.   Hwang JM, Chang DJ, Kim US, Lee YS, Park YS, Kaang BK, et al. Cloning and

622   functional characterization of a Caenorhabditis elegans muscarinic

623   acetylcholine receptor. Receptors Channels. 1999;6(6):415–24.

624   29.   Lee YS, Park YS, Nam S, Suh SJ, Lee J, Kaang BK, et al. Characterization of GAR-

625   2, a novel G protein-linked acetylcholine receptor from Caenorhabditis

626   elegans. J Neurochem. 2000 Nov;75(5):1800–9.

627 30. Ishii M, Kurachi Y. Muscarinic acetylcholine receptors. Curr Pharm Des.

628   2006;12(28):3573–81.

629 31. Hanstein WG. Uncoupling of oxidative phosphorylation. Biochim Biophys

630   Acta. 1976 Sep 27;456(2):129–48.

631 32. Ghosh R, Andersen EC, Shapiro J a, Gerke JP, Kruglyak L. Natural Variation in a

632   Chloride. Science (80-). 2012;335(February):574–8.

633 33. Zdraljevic S, Strand C, Seidel HS, Cook DE, Doench JG, Andersen EC. Natural

634   variation in a single amino acid substitution underlies physiological responses

635   to topoisomerase II poisons. Macdonald SJ, editor. PLoS Genet. 2017 Jul

636   12;13(7):e1006891.

637 34. Vu V, Verster AJAJ, Schertzberg M, Chuluunbaatar T, Spensley M, Pajkic D, et

638   al. Natural Variation in Gene Expression Modulates the Severity of Mutant

639   Phenotypes. Cell. 2015;162(2):391–402.

640 35. Wicks SR, Yeh RT, Gish WR, Waterston RH, Plasterk RH. Rapid gene mapping

641   in Caenorhabditis elegans using a high density polymorphism map. Nat Genet.

642   2001 Jun 1;28(2):160–4.

643 36. Steger KA, Avery L. The GAR-3 muscarinic receptor cooperates with calcium

644   signals to regulate muscle contraction in the Caenorhabditis elegans pharynx.

645   Genetics. 2004 Jun 1;167(2):633–43.

646 37. Liu Y, LeBoeuf B, Garcia LR. G alpha(q)-coupled muscarinic acetylcholine

647   receptors enhance nicotinic acetylcholine receptor signaling in

Caenorhabditis elegans mating behavior. J Neurosci. 2007 Feb 7;27(6):1411–

21.

38.   Dittman JS, Kaplan JM. Behavioral impact of neurotransmitter-activated G-

protein-coupled receptors: muscarinic and GABAB receptors regulate

Caenorhabditis elegans locomotion. J Neurosci. 2008;28(28):7104–12.

39.   Chan JP, Staab TA, Wang H, Mazzasette C, Butte Z, Sieburth D. Extrasynaptic

muscarinic acetylcholine receptors on neuronal cell bodies regulate

presynaptic function in Caenorhabditis elegans. J Neurosci. 2013 Aug

28;33(35):14146–59.

40.   Hendricks M, Ha H, Maffey N, Zhang Y. Compartmentalized calcium dynamics

in a C. elegans interneuron encode head movement. Nature.

2012;487(7405):99–103.

41.   Stiernagle T. Maintenance of C. elegans. In: The C. elegans Research

Community, editor. Wormbook. 2006.

42.   Lehner B, Crombie C, Tischler J, Fortunato A, Fraser AG. Systematic mapping

of genetic interactions in Caenorhabditis elegans identifies common modifiers

of diverse signaling pathways. Nat Genet. 2006;38(8):896–903.

43.   Moore J, Linkert, Melissa Blackburn, Colin Carroll, Mark Ferguson RK, Flynn H,

Gillen K, Leigh R, Li S, et al. OMERO and Bio-Formats 5: flexible access to large

bioimaging datasets at scale. Proc SPIE 9413, Med Imaging 2015 Image

Process. 2015;941307.

44.   van der Walt S, Schönberger J, Nunez-Iglesias, J Boulogne F, Warner J, Yager N,

Gouillart E, et al. scikit-image: Image Processing in Python. PeerJ.

2014;(2):e423.

Figure Captions

Figure 1: Outline of an image-based assay for worm mobility

A: Worms are placed in a 96 well plate in M9 buffer and treated with drugs as required. Two images of each well (Im1 and Im2) are captured at an interval of 500 ms at each timepoint. Images are processed as described in Materials and Methods to identify portions of the worms that move between the two images (green) and parts that remain stationary (red). This allows the calculation of a raw fractional mobility score (F.M.S.) as the fraction of total worm-associated pixels (red and green) that moved (green). This processing is illustrated in a magnified region of Im1 and Im2 (defined by blue box). **B: The image-based assay produces a rich view of drug effects on worm movement.** L4 worms were treated with a range of doses of aldicarb and movement measured at 5 minute time points over a 3 hour time course. The heat map shows the F.M.S. for each drug dose at each timepoint — white indicates full movement, red indicates lack of movement. Either dose response curves at a given time point (pink arrow) or kinetic responses to a single drug dose (red and blue arrows) can be extracted from these rich data. Data are means of four independent experiments. Error bars show standard error of the mean.

Figure 2: Assay characterisation.

A: Effect of aldicarb on adult worms: Adult worms were treated with aldicarb at a range of concentrations and FMS was calculated at 5 minute intervals. FMS was

plotted as a time-course of response for each concentration. **B: Adult worms**

homozygous for a loss-of-function *unc-38*(*e264*) mutation are insensitive to

aldicarb. A-C show means of 4 independent experiments; error bars show standard

error of the mean. All data are normalised to drug-free controls.**C: The dose-**

response relationship for aldicarb is affected by addition of mecamylamine,

an antagonist of nAChRs. EC50 values were calculated by fitting a 3-parameter

logistic function to the data (see Materials and Methods). **D: Assay detects effects**

of targeting genes required for normal movement using RNAi. L1 worms were

grown for 4 days in the presence of *E. coli* strains expressing dsRNA that target

individual genes known to be required for normal movement. L1 progeny from

these cultures were isolated by filtration and their mobility quantified using our

assay. Mobility scores are normalized to a non-targeting RNAi control (dsRNA

targeted against GFP). Data represent the mean of 8 wells, containing 100 worms

per well.

Figure 3: A wide range of compounds elicit movement responses.

L1 worms were exposed to a range of doses of various compounds and movement

quantified. **A: Effect of potassium cyanide on L1 worms.** Data are means of 4

independent replicates. **B: Effect of cycloheximide on L1 worms.** Data are means

of 3 independent replicates. **C: Effect of sodium chloride on L1 worms**. Data are

means of 4 independent replicates. **D: Effect of abamectin on L1 worms.** Data are

means of 4 independent replicates. **E-G: Novel compounds elicit a range of**

dynamic responses. Time-resolved responses for a number of novel compounds

elicited a range of different response dynamics. L1 worms were continuously

exposed to 40 μM of each compound. Data are means of 3 independent replicates.

(**A-G)** Error bars show standard error of the mean.

Figure 4: Larval worms show dose-dependent recovery from cholinergic

paralysis.

A: Timecourse of response to 4mM aldicarb at different developmental stages.

Worms from each developmental stage were purified and dispensed using a COPAS

biosort worm sorter before exposure to aldicarb as in Figure 2. Early larval stages

show a robust recovery from paralysis, whereas L4 and adult worms show little

recovery. Data are means of 3 independent experiments. **B: Response of L1 worms**

to nicotine. L1 worms were treated with a range of nicotine concentrations and

FMS measured over a 3hr time course. Data are means of 4 independent

experiments. **C: Recovery of L1 worms from Ald-induced paralysis increases**

with higher doses. For each concentration shown, a proportional recovery value

was calculated as described in Material and Methods. Data are means of four

independent experiments. **D: Recovery of L1 worms from Nic-induced paralysis**

decrease with higher doses. Proportional recovery scores were calculated as

above. Data are means of three independent experiments. **A-D:** Error bars show

standard error of the mean.

Figure 5: L1 cholinergic recovery can be modified by compounds with

muscarinic activity.

A: The recovery from Ald-paralysis is suppressed by the mAChR antagonist

Atr. L1 worms were treated with 4mM Ald along with a range of doses of Atropine

(Atr); FMS was measured every 5mins over a 3hr time course.

B: *gar-3* mutants show greatly reduced recovery from Ald-induced paralysis.

Wild-type animals, or worms homozygous for mutations in either *gar-1, gar-2,* or

gar-3 were exposed to 4mM Ald and FMS measured across a 3hr time course.

C: Aldicarb induces recovery from nicotine-induced paralysis in L1 larvae.

Various concentrations of aldicarb were combined with a concentration of nicotine

sufficient to cause sustained paralysis (10 mM, shown in blue). Increasing

concentrations of aldicarb stimulated increasing degrees of recovery from paralysis.

D: The mAChR-specific agonist OxoM similarly stimulates recovery from Nic-

induced paralysis. L1 animals were treated with 10mM Nic combined with a range

of concentrations of OxoM. **E: OxoM-induced recovery requires *gar-3*:** Wild-type

animals, or worms homozygous for mutations in either *gar-1, gar-2,* or *gar*-3 were

exposed to 10mM Nic and 10mM OxoM and FMS measured across a 3hr time course.

The shaded portion represents the range of responses of N2, *gar-1, gar-2* and *gar-3*

to nicotine, in the absence of OxoM. **(A-E)** All data are means of 4 independent

experiments, normalized to no drug controls. Error bars show standard error of the

mean. **F: OxoM-does not induce recovery from Ald paralysis in adult worms:**

requires *gar-3*: Adult worms were exposed to 5mM Ald, with or without 10mM

OxoM. The presence of OxoM does not induce recovery from Ald-mediated paralysis

in adult worms. Data are means of three independent experiments; error bars show

standard error of the mean.

Figure 6: Conceptual model of cholinergic responses in L1 *C. elegans*

A: Nic alone causes paralysis through binding to nAChRs. **B:** The paralysis caused by

the action of Nic at nAChRs in relieved by the action of OxoM at mAChRs. **C:** By

elevating levels of ACh, Ald treatment leads to transient paralysis by activation of

nAChRs, followed by recovery from paralysis, mediated by the action of ACh on

mAChRs.

Figure 7 Dose sensitivity to KCN is complex.

A: L1 worms were treated with a range of KCN doses and their movement

measured over time. The Fractional Mobility Scores (FMS) are shown at various

time points. At early time points the dose response curves are complex —

intermediate doses cause more severe movement defects than high doses. **B:**

Decreased sensitivity to KCN at higher doses is dependent on a metabolic

transition to glycolysis. Inhibition of glycolysis with 2-dexoy-D-glucose causes

higher-doses of KCN to be more toxic than lower doses, in contrast to the dose-

response relationship of KCN alone. This suggests that the relatively mild effects of

higher KCN concentrations seen in panel A is at least partially attributable to KCN

stimulating a shift in metabolic pathways from oxidative phosphorylation to

glycolysis. Data shown are at the 20 minute time point. **A-B:** Curves show mean of 3

repeats; error bars are standard error of the mean.

Figure 8 Variation in dose sensitivity to KCN changes across development.

A: Worms of specific stages of either N2 or CB4856 isolates were purified using a worm sorter. Worms are sorted based on length and darkness — the exact windows chosen are indicated by coloured boxes and examples of the sorted worms shown.

B: N2 or CB4856 worms of specific developmental stages were exposed to a range of doses of KCN and their movement measured over a 3hr time course. The fractional mobility score (F.M.S.) was calculated at each time point. The figure shows the difference in F.M.S. scores for N2 and CB4856 — blue colours indicate that N2 has a lower score, red colours that CB4856 is moving less.

Figure S1: Response of nAChR subunit mutants to aldicarb in L1 worms

Several nAChR subunits are required for wild-type sensitivity of L1 worms to 8mM aldicarb, including UNC-29, UNC-63, LEV1, LEV-8 and UNC-38. ACR-16 and ACR-8 are not required for Ald-sensitivity in adults. All data are means of 3 independent replicates. Error bars are standard error of the mean.

Figure S2: Further characterisation of mAChR recovery

Arecoline (Are), a mAChR agonist induces recovery from paralysis under 10 mM Nic in the same manner as OxoM. Data are means of 4 wells. Error bars are standard error of the mean.

Figure S3: Time course of response of L1 worms to a range of Ald concentrations. Data are means of 4 independent experiments. Error bars are standard error of the mean.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

B

Figure 8

A

B

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3