Genomic basis of multidrug-resistance, mating, and virulence in *Candida auris* and related emerging species

José F. Muñoz[1], Lalitha Gade[2], Nancy A. Chow[2], Vladimir N. Loparev[3], Phalasy Juieng[3], Elizabeth L. Berkow[2], Rhys A. Farrer[1], Anastasia P. Litvintseva[2]*, Christina A. Cuomo[1]*

[1] Broad Institute of MIT and Harvard, Cambridge, MA USA. [2] Mycotic Diseases Branch, Centers for Disease Control and Prevention, Atlanta, GA USA. [3] Biotechnology Core Facility Branch, Centers for Disease Control and Prevention, Atlanta, GA USA.

*Corresponding authors: Christina A. Cuomo cuomo@broadinstitute.org; Anastasia P. Litvintseva, frq8@cdc.gov

Abstract

Candida auris is an emergent fungal pathogen of rising public health concern due to increasing reports of outbreaks in healthcare settings and resistance to multiple classes of antifungal drugs. While distantly related to the more common pathogens *C. albicans* and *C. glabrata*, *C. auris* is closely related to three rarely observed and often multidrug-resistant species, *C. haemulonii, C. duobushaemulonii and C. pseudohaemulonii*. Here, we generated and analyzed near complete genome assemblies and RNA-Seq-guided gene predictions for isolates from each of the four major *C. auris* clades and for *C. haemulonii, C. duobushaemulonii and C. pseudohaemulonii*. Our analyses mapped seven chromosomes and revealed chromosomal rearrangements between *C. auris* clades and related species. We found conservation of genes involved in mating and meiosis and identified both *MTL***a** and *MTL*α *C. auris* isolates, suggesting the potential for mating between clades. Gene conservation analysis highlighted that many genes linked to drug resistance and virulence in other pathogenic *Candida* species are conserved in *C. auris* and related species including expanded families of transporters and lipases, as well as mutations and copy number variants in *ERG11* that confer drug resistance. In addition, we found genetic features of the emerging species that likely underlie differences in virulence and drug response between these and other *Candida* species, including genes involved in cell wall structure. To begin to characterize the species-specific genes important for antifungal response, we profiled the gene expression of *C. auris* in response to voriconazole and amphotericin B and found induction of several transporters and metabolic regulators that may play a role in drug resistance. This study provides a comprehensive view of the genomic basis of drug resistance, potential for mating, and virulence in this emerging fungal clade.

Introduction

Candida auris is an emerging fungal pathogen of increasing concern due to high drug resistance and high mortality rates[1,2]. In addition, outbreaks have been reported in hospital settings, suggesting healthcare transmission[2,3]. C. auris clinical isolates are typically multidrug-resistant (MDR), with common resistance to fluconazole and variable susceptibility to other azoles, amphotericin B, and echinocandins[2]. C. auris causes bloodstream and other invasive and superficial infections, similar to a group of rarely observed, phylogenetically related species including C. haemulonii, C. duobushaemulonii and C. pseudohaemulonii[4,5]. These species also display MDR, most commonly to amphotericin B and also reduced susceptibility to azoles and echinocandins[4,5]. Together, C. auris and these closely related species represent an emerging clade of invasive fungal pathogens, which are not only difficult to treat, but also difficult to identify using standard laboratory methods[6], generally requiring molecular methods for proper identification.

Initial whole genome analysis of C. auris isolates from Pakistan, India, South Africa, Japan and Venezuela identified four clades that are specific to each geographic region, suggesting that each of these clades emerged nearly simultaneously in different regions of the world[2]. Including data from other recent studies, the current representation of each clade is as follows: clade I comprises isolates from India, Pakistan and the United Kingdom[2,3,7], clade II from Japan and South Korea[2,8], clade III from South Africa, and clade IV from Venezuela[2]. Analyses of SNPs identified from whole genome sequence[2,9] and of multilocus sequence typing (MLST)[10,11] found low genetic diversity between isolates within each C. auris clade. Little is known about whether phenotypic differences exist among isolates from different C. auris clades; one noted difference is that isolates from India and South Africa are capable of assimilating N-acetylglucosamine while isolates from Japan and South Korea were not able to process this compound[2].

Phylogenetic studies revealed that Candida auris belongs to the C. haemulonii clade and is distantly related to the more common human pathogenic species including C. albicans and C. glabrata [4,5,12,13]. However, these prior studies have not clearly resolved the relationships of species within the C. haemulonii clade, including C. haemulonii, C. duobushaemulonii and C. pseudohaemulonii[4,5,12,13]. C. lusitaniae, a rarely observed cause of infection[14], is a sister clade to this group of emerging species[13,15]. Despite the fact that C. auris is highly divergent from other Saccharomycetales yeasts from the CTG clade, which includes the common human pathogens C. albicans, C. tropicalis and C. parapsilosis, most of our limited current knowledge of C. auris

resistance and virulence had been inferred based on conservation of genes associated with
drug resistance and virulence in *C. albicans* or in *C. glabrata*, which is part of the distantly
related *Nakaseomyces* clade. Initial comparative analysis of the gene content of one isolate of
C. auris with *C. albicans* found that some orthologs associated with antifungal resistance are
present in *C. auris*, including drug transporters, secreted proteases and manosyl transferases[13].
In addition, preliminary genomic studies showed that the targets of several classes of antifungal
drugs are conserved in *C. auris*, including the azole target lanosterol 14 α-demethylase
(*ERG11*), the echinocandin target 1,3-beta-glucan synthase (*FKS1*), and the flucytosine target
uracil phosphoribosyl-transferase (*FUR1*)[2,16]. Furthermore, point mutations associated with drug
resistance in other species are observed in many clinical isolates and are associated with *C.*
auris clades[2,7,16]. Recently, two *ERG11* mutations (Y132F and K143R) in *C. auris* were found to
increase resistance to fluconazole[17], however the role of other genes in drug resistance and
virulence has not been reported for this group of emerging MDR species.
81
While efforts to sequence *C. auris* genome provided an initial view of genome content[9,13],
available assemblies in GenBank are highly fragmented, inconsistently annotated, and do not
provide a complete representation of all *C. auris* clades. The related species *C. haemulonii* and
C. duobushaemulonii were recently sequenced [18,19]. Here, we generated and annotated highly
complete genome assemblies for isolates from each of the clades of *C. auris* as well as for the
related species *C. haemulonii*, *C. duobushaemulonii* and *C. pseudohaemulonii*. Comparison of
these genomes to other sequenced *Candida* species revealed that *C. auris* has notable
expansions of genes linked to drug resistance and virulence in *C. albicans*, including families of
oligopeptide transporters, siderophore-based iron transporters, and secreted lipases. We used
RNA-Seq to examine the response of two isolates of *C. auris* to antifungal drugs and detected
the upregulation of transporters and metabolic regulators that have been previously associated
with drug resistance in *C. albicans*. In addition, we also observed several genes that were either
unique or expanded in *C. auris* and related species were upregulated in response to azole and
amphotericin B exposure. We also found evidence that *C. auris* may be capable of mating and
meiosis, based on the identification of both mating types in the populations, conservation of
genes involved in mating and meiosis, and detection of chromosomal rearrangements between
two clades. These results revealed fundamental insights into the evolution of drug resistance
and pathogenesis in *C. auris* and closely related species and the potential for mating and
recombination between the *C. auris* clades.
Results

Genome characteristics of *Candida auris* and closely related species

102
We generated highly complete genome assemblies for four *Candida auris* isolates and
for three phylogenetically related species, *C. haemulonii*, *C. duobushaemulonii,* and *C.*
pseudohaemulonii. Most of these isolates displayed increased resistance to fluconazole and
also increased resistance to voriconazole or amphotericin B (**Table S1**). The *Candida auris*
assemblies represent each of the four major clades[2], including an updated assembly for strain
B8441 (clade I) and the first representatives of clade II (strain B11220), clade III (strain B11221)
and clade IV (strain B11243). As *C. auris* is distantly related to other previously sequenced
Candida species, we also sequenced genomes of the closely related species *C. haemulonii*
(strain B11899), *C. duobushaemulonii* (strain B09383)*,* and *C. pseudohaemulonii* (strain
B12108) to enable comparative genomic analysis. The genome assemblies of *C. auris* (B8441
and B11221), *C. haemulonii*, and *C. duobushaemulonii* were sequenced using PacBio and
Illumina, whereas the other two *C. auris* strains and *C. pseudohaemulonii* were sequenced only
with Illumina (**Table 1**; **Methods**). The genome assemblies of *C. auris* range from 12.1 Mb in
B11220 to 12.7 Mb in B11221. The *C. auris* B8441 and B11221 genome assemblies were
organized in 15 and 20 scaffolds, respectively, of which 7 scaffolds included most of the
sequenced bases in both strains (**Table 1**). This represents an improvement upon the
previously generated genome assembly of *C. auris* strain 6684 (clade I)[13], which consists of 99
scaffolds (759 contigs) that include contig gaps and has an inflated number of predicted genes,
based on our analysis (see below; **Methods**). The genome assemblies of Illumina data for *C.*
auris B11220 (clade II) and B11243 (clade IV) are less contiguous than the assemblies that
included long reads, consisting of 324 or 285 contigs, respectively (**Table 1**). The genome
assemblies of *C. haemulonii*, *C. duobushaemulonii,* and *C. pseudohaemulonii* were also highly
contiguous with 11, 7 and 36 scaffolds, respectively (**Table 1**). Comparison of these four *C.*
auris genomes and those of the three related species revealed that the genome sizes are very
similar, ranging in size from 12.1 Mb in *C. auris* B11220 to 13.3 Mb in *C. haemulonii*, similar to
that of *C. lusitaniae* (12.1 Mb) and other *Candida* species (**Table S2**).

129
The assemblies of the *C. auris* clades are highly identical, sharing an average pairwise
nucleotide identity of 98.7% across all four clades (I, II, III and IV); clades II and III appear the
most similar and share 99.3% identity. By contrast, comparison of nucleotide genome identity
between species highlights greater interspecies genetic divergence: 88% between *C. auris* and
each of the three related species (*C. haemulonii*, *C. duobushaemulonii,* and *C.*
pseudohaemulonii), 89% between *C. haemulonii* and *C. duobushaemulonii*, and 92% between

C. duobushaemulonii and *C. pseudohaemulonii*. To provide a quantitative measure of the
degree of genetic variation within the *C. auris* population, we calculated the genome-wide
nucleotide diversity (π) using SNPs identified from the isolate sequences of Lockhart *et al*[2]. The
estimated π is lowest for *C. auris* clade I 0.00050, and slightly higher for clades III and IV
(0.00079 and 0.00075, respectively); π for all *C. auris* isolates is 0.0039, ~17-fold higher than
the intra-clade levels. Comparing to other human fungal pathogens, this level of diversity is
lower than that reported in *C. neoformans* var. *grubii*, (π = 0.0074, [20])**,** but higher than the
extreme clonality reported for *Trichophyton rubrum* (π = 0.00054, [21]). These comparisons
support that the different clades of *C. auris* comprise a single species well separated from the
related MDR species.
146
As an independent assessment of the genome assembly size and structure, we generated
optical maps of the *C. auris* B8441 and B11221 isolates (**Figure S1**). Consistent with the
assemblies of these isolates, the maps had seven linkage groups; nearly all of the genome
assemblies were anchored to the optical maps (98.8 % of B8441 and 97.2 % of B11221; **Figure**
S1; Table 1). This supports the presence of seven chromosomes in *C. auris*, consistent with the
chromosome number found in previous studies using electrophoretic karyotyping by pulsed-field
gel electrophoresis (PFGE)[22]. While the genomes of *C. auris* are highly syntenic, we found
evidence of a few large chromosomal rearrangements between *C. auris* B8441 and B11221
based on comparison of the assemblies and the optical maps (**Figures 1**, **S1**). We confirmed
that the junctions of these rearrangements are well supported in each assembly; these regions
show no variation in the depth of PacBio and Illumina aligned reads and there is no evidence of
assembly errors across these rearrangement breakpoints. We additionally independently
identified structural variants based on the read alignments to the B8441 and B11221 assemblies
and recovered each of the rearrangements present in these assemblies. These large
chromosomal rearrangements included one inversion of 136 kb between B8441 sc01 and
B11221 sc01, a 274 kb translocation between B8441 sc08 and B11221 sc03, and a 300 kb
translocation between B8441 sc10 and B11221 sc01 (**Figure 1**). The genomes of *C. auris*, *C.*
haemulonii, *C. duobushaemulonii,* and *C. pseudohaemulonii* showed limited chromosomal
rearrangements between each other, mostly intra-chromosomal inversions between *C. auris*
and *C. haemulonii, C. duobushaemulonii,* and *C. pseudohaemulonii,* and large chromosomal
translocations between *C. haemulonii*, *C. duobushaemulonii,* and *C. pseudohaemulonii* (**Figure**
1a). These rearrangements between *C. auris* clades, and between species, could potentially
prevent genetic exchange between these groups, since some crossover events will generate

missing chromosomal regions or other aneuploidies and may result in nonviable progeny.

Evolution of mating-type locus in the emerging multidrug-resistant species including *C.*

auris

We characterized the mating type locus in *C. auris* and identified representatives of both

mating types. The mating locus structure is highly conserved compared to closely related

species including *C. lusitaniae* and other Saccharomycetales yeasts from the CTG clade

Candida (**Figure 2**). Many *Candida* species, including diploid asexual species, have

heterothallic *MTL* idiomorphs, and mating occurs between cells of opposite mating type, *MTL***a**

and *MTL*α[23]. We found that the genes flanking the *MTL* locus in some species from the CTG

clade *Candida*, namely the phosphatidylinositol kinase gene (*PIK1*), the oxysterol binding

protein gene (*OBP1*) and the poly(A) polymerase gene (*PAP1*) were adjacent in all sequenced

genomes of the emerging MDR clade (**Figure 2a**). Previous work reported that the genome of

C. auris 6684 included the *MTL* flanking genes but did not identify either *MTL***a** or *MTL*α genes

at this locus[13]. In the chromosome level genome assemblies of *C. auris* (B8441 and B11221),

the *PIK1/OBP1/PAP1* genes are present at a single locus in chromosome 3 in B8441 (sc05)

and B11221 (sc03). Notably, we found all *C. auris* isolates contained either the *MTL***a** and *MTL*α

idiomorphs at this locus and that gene order was conserved compared to the *C. lusitaniae*

MTL[24] (**Figure 2a**; **Table S3**). We found that the *MTL***a** is present in *C. auris* B8441 and 6684

(clade I), B11243 (clade IV), and *C. pseudohaemulonii*, spanning 14.9 kb (**Figure 2a**). By

contrast, *C. auris* B11220 (clade II), B11221 (clade III), and *C. haemulonii* and *C.*

duobushaemulonii contain *MTL*α, spanning 14.3 kb (**Figure 2a**). Phylogenetic analysis of the

non-mating flanking genes (*PIK1/OBP1/PAP1*) supports the inheritance of idiomorphs of these

genes with the *MTL***a** and *MTL*α genes (**Figure 2b**). Upon further examination and manual

annotation, we determined that **a**1*/***a**2 are present in *MTL***a** isolates, and α1 is present in *MTL*α

isolates; as in *C. lusitaniae*, the MTL α locus is missing the α2 gene[24] (**Figure 2a**). RNA-Seq

data was used to guide gene prediction of **a**1 and **a**2, and further established that both genes

are expressed in B8441 (**Figure S2**), supporting the hypothesis that these genes could be

functional, and that the MTL locus could be used to classify *C. auris* isolates. To further

characterize the evolution of mating type in the population, we examined the *MTL* locus using

50 isolates from Lockhart *et al.*[2] and found that all isolates from clade I and IV have *MTL***a**,

whereas all isolates from clade II and III had *MTL*α (**Figures 2c**, **S3**). The fact both **a** and α *MTL*

alleles are present and expressed in *C. auris* suggests this species may be capable of mating,

either in the ancestral population of the outbreak clades or if the expanding outbreak results in
the presence of isolates from both mating types in a geographic area.

We examined the conservation of genes involved in meiosis to provide additional support for
potential mating in *C. auris*.　We found that many of the key meiotic genes are similarly
conserved between *C. lusitaniae*, *C. guilliermondii*, *C. auris*, *C. haemulonii*, *C.*
duobushaemulonii and *C. pseudohaemulonii* (**Table S3**). Some genes involved in meiosis in *S.*
cerevisiae appeared absent in these species including the recombinase *DMC1* and cofactors
(*MEI5* and *SAE3*), synaptonemal-complex proteins (*ZIP1* and *HOP1*), and genes involved in
crossover interference (*MSH4* and *MSH5*; **Table S3**). A small number of genes involved in
meiosis were present in *C. lusitaniae* but absent in *C. auris*, *C. haemulonii*, *C.*
duobushaemulonii, and *C. pseudohaemulonii*. In *C. auris*, the DNA recombination and repair
genes *RAD55* and *RAD57* appear to be absent, however *RAD55* is widely absent in the CTG
Candida clade while the *RAD51* paralog of the *RAD55*/*RAD57* complex is present. *C. lusitaniae*,
despite the loss of many of the same meiotic genes, undergoes meiosis during sexual
reproduction involving diploid intermediates[24]. The fact that most components of the mating and
meiosis pathways are similarly conserved in *C. auris* and closely related species including *C.*
lusitaniae suggests these species may have the ability to mate and undergo meiosis as
observed for *C. lusitaniae*.

Phylogenetic position of *C. auris, C. haemulonii, C. duobushaemulonii,* and *C.*
pseudohaemulonii
Using the complete genomes, we estimated a strongly supported phylogeny of *C. auris*,
C. haemulonii, *C. duobushaemulonii,* and *C. pseudohaemulonii*, relative to other species from
the order Saccharomycetales including *C. lusitaniae*, *C. tropicalis*, *C. albicans* and *C. glabrata*
(**Table S2**). Based on a concatenated alignment of 1,570 single copy core genes, a well
supported maximum likelihood tree placed *C. auris*, *C. haemulonii*, *C. duobushaemulonii,* and
C. pseudohaemulonii as a single clade, confirming the close relationship of these species
(100% of bootstrap replicates; **Figure 3a**). The *C. auris* clades appear more recently diverged
based on short branch lengths within this species. Previous phylogenetic analyses had shown
conflicting relationships between *C. auris*, *C. haemulonii*, *C. duobushaemulonii,* and *C.*
pseudohaemulonii[4,5,12,13]. Our phylogenetic analysis strongly supports that *C. duobushaemulonii*
and *C. pseudohaemulonii* are most closely related to each other, and form a sister group to *C.*
haemulonii, which appeared as the more basally branching species (**Figure 3a**). The most

closely related species to this MDR clade is *C. lusitaniae*, which is the more basally branching
member of this group (**Figure 3a**).

Gene family expansions supported mechanisms of drug resistance and virulence

Gene annotation of the *C. auris* genomes was performed using RNA-Seq paired-end
reads to improve gene structure predictions (**Methods**). The predicted gene number was highly
similar across all *C. auris* genomes as well as in *C. haemulonii*, *C. duobushaemulonii,* and *C.*
pseudohaemulonii. In *C. auris*, the number of protein-coding genes varied between 5,421 in
B8441 and 5,601 in B11243. For *C. haemulonii*, *C. duobushaemulonii,* and *C.*
pseudohaemulonii the numbers were very similar, ranging from 5,288 to 5,410 predicted genes
(**Table 1**; **Figure S4a**). High representation of core eukaryotic genes provides evidence that
these genomes are nearly complete; 96-98% of these conserved genes are found in all
annotated genome assemblies (**Figure S4b**). By examining orthologous genes in *C. auris*, *C.*
haemulonii, *C. duobushaemulonii,* *C. pseudohaemulonii*, and twelve additional
Saccharomycetales genomes, including *C. lusitaniae*, *C. tropicalis*, *C. albicans* and *C. glabrata*,
we found a total of 2,379 core ortholog clusters had representative genes from all twenty
analyzed genomes (**Figure S4**). We found a small number of unique genes in the *C. auris*
clades ranging from 15 (B8441; clade I) to 54 (B11221; clade III) genes, and higher numbers in
the three related species (203 in *C. haemulonii*, 83 in *C. duobushaemulonii,* and 88 *C.*
pseudohaemulonii) (**Figure S5**). The unique genes in *C. auris* clades include oligopeptide and
ABC transporters (**Table S4**). While unique glycophosphatidylinositol (GPI)-anchored proteins
were identified in *C. auris*, we found high conservation across *C. auris*, *C. haemulonii*, *C.*
duobushaemulonii, C. pseudohaemulonii of seven (GPI)-anchored proteins (*PLB3*, *IFF4*,
PGA52, *PGA26*, *CSA1*, *HYR3* and *PGA7*) that were upregulated during *C. auris* biofilm
formation and are associated with antifungal resistance and biofilm mechanisms in *C.*
albicans[25](**Table S4**).

To characterize changes in gene content that may play a role in the evolution of multidrug-
resistance and virulence in *C. auris*, *C. haemulonii*, *C. duobushaemulonii,* and *C.*
pseudohaemulonii, we searched for expansions or contractions in functionally classified genes
compared to other related species (**Table S2**). We identified PFAM domains that were
significantly enriched or depleted (**Methods**; **Figures 3b, S4**). Domains associated with
transmembrane transporters (OPT, MFS) and secreted lipases (LIP) were enriched in *C. auris*,
C. haemulonii, *C. duobushaemulonii,* and *C. pseudohaemulonii* compared to other genomes (*q-*

value < 0.05, Fisher's exact test; **Figure 3b**). We therefore further classified transmembrane transporters using the Transporter Classification Database (TCDB) and found that the higher copy number of transporters in the emergent MDR clade could be attributed to oligopeptide transporters (*OPT*) and siderophore iron transporters (*SIT*). In *C. albicans*, *OPT* transporters enable uptake of small peptides; the expression of some *OPT* transporters are up-regulated by azole drugs[26,27]. While most of the 14 *OPT* genes found in *C. auris* had orthologs in *C. albicans* (*OPT1-8*), a subset of these transporters appear recently duplicated in the MDR emergent clade, including an expansion of three *OPT1*-like transporters, and five transporters similar to *OPT2*, *OPT3*, and *OPT4* (**Figures 4a**, **S6**). Most of the *OPT* genes (up to 8) are located in a conserved locus among emerging MDR species encompassing 296 kb of chromosome 6; this gene family appears to have expanded by tandem duplication (**Figures 4c**). In *C. albicans*, iron transporters include the siderophore transporter *SIT1* and the iron permeases *FTR1* and *FTR2*; a subset of these transporters is uniquely expanded in *C. auris* and closely related species, including the expansion of 14 ortholog groups in *C. auris* related to *C. albicans SIT1* (**Figures 4b**, **S6**). Secreted lipases are also expanded in the genomes of *C. auris* and closely related species (*q*-value < 0.05, Fisher's exact test; **Figure 3b**). The *C. auris* clade has similar counts of lipases relative to *C. albicans* and *C. dubliniensis*, however, these proteins are expanded relative to more closely related human pathogenic species, including *C. lusitaniae*, *C. guilliermondii*, *C. krusei*, and *C. glabrata* (**Figure 3b**). Phylogenetic analysis suggested independent evolutionary trajectories of secreted lipases in emerging *C. auris* and related species, where the most recent ortholog family of lipases includes *C. albicans LIP4*, *5*, *8* and *9* (**Figure S7**). The secretion of lipases may be important during infection for nutrient acquisition, adaptation, virulence and immune evasion[28]; we identified a predicted secretion signal in all lipases encoded by *C. auris* and related species, supporting an extracellular role in these emergent MDR species.

Conservation of known drug resistance and pathogenesis-associated genes

The isolates selected for genome sequencing display increased resistance to antifungal drugs. Increased resistance to fluconazole is most commonly observed in *C. auris* with some isolates also displaying increased resistance to voriconazole or amphotericin B (**Table S1;**[2]). The three related species all displayed increased amphotericin B resistance, and two were also resistant to fluconazole. All isolates appeared sensitive to the echinocandins tested, anidulafungin and caspofungin (**Table S1**). Most of the genes associated with drug resistance and pathogenesis in *C. albicans* are conserved in *C. auris*, *C. haemulonii*, *C. duobushaemulonii,*

and *C. pseudohaemulonii*. We identified orthologs of genes noted to confer drug resistance in
C. albicans, either by acquiring point mutations, increasing transcription, or copy number
variation. The annotated genome assemblies of *C. auris*, *C. haemulonii*, *C. duobushaemulonii,*
and *C. pseudohaemulonii* contain a single copy of the *ERG11* azole target and the *UPC2*
transcription factor that regulates expression of genes in the ergosterol pathway, as well as all
the gene components of the ergosterol biosynthesis pathway (**Table S5**). Several of the sites in
ERG11 subject to drug resistant mutations in *C. albicans* are similarly mutated in drug resistant
C. auris isolates (at positions Y132, K143, and F126, as reported previously[2]). Analysis of the
annotated genome assemblies of *C. auris* isolates agrees with prior SNP analysis[2], with the
exception of F126L mutation in B11221 (clade III), initially reported as F126T for this isolate
though this was recently corrected to F126L[29]. The F126L mutation is also observed in a
separate genomic study of *C. auris*[7] and in drug resistant *C. albicans*[30] (**Figure S8**). The Y132F
mutation observed in two *C. auris* clades is also found in *C. pseudohaemulonii*, suggesting this
site may contribute to the increased azole resistance of this *C. pseudohaemulonii* isolate (**Table
S1**). Other sites in *ERG11* that confer drug resistance in *C. albicans* do not display drug
resistant mutations in *C. auris*, *C. haemulonii*, *C. duobushaemulonii,* and *C. pseudohaemulonii*
(**Figure S8**). In addition, we found that not all *C. auris* isolates have a single copy of *ERG11*.
Using copy number variation (CNV) analysis of the Illumina read depth of 47 previously
sequenced isolates, we found a total of 6 large duplicated regions (CNVnator *p*-value < 0.01)
ranging in size from 12 to 153 kb (**Table S6, Methods**). The largest duplicated region of 153 kb
is present in two isolates (B11227 and B11229) from clade III and encompassed 62 genes
including *ERG11* (**Figure S9**; **Table S6**). These two isolates also display increased resistance
to fluconazole (**Table S1**). Long read assemblies of these isolates could be used to examine the
chromosomal context of this duplication.
329
Additionally, we identified orthologs of transporters from the ATP binding cassette (*ABC*) and
major facilitator superfamily (*MFS*) classes of efflux proteins that are involved in clinical
antifungal resistance in *C. albicans* by the overexpression of ABC transporter family *CDRs*
(*CDR1* and *CDR2*) and the MFS transporter *MDR1* [31,32]. We identified a single copy of the
multidrug efflux pump *MDR1* in all sequenced isolates (**Table 2**). We further characterized
presence of *CDR* genes and *MDR1* in the *C. auris* population by examining CNV and gene
conservation across the 47 isolates from Lockhart *et al.*[2], and did not detect any copy number
variation of these transporters across this set (**Table S7**). Candidate multidrug transporters
similar to *CDR1*, *CDR2*, and related genes include 5 genes in most genomes (*C. auris* B8441,

B11220, and B11243, *C. haemulonii*, and *C. pseudohaemulonii*), 6 genes in *C. auris* B11221,

and 4 genes in *C. duobushaemulonii* (**Table 2**; **Figure S10**). Phylogenetic analysis of these

ABC transporters showed that one of the genes in *C. auris* is related to *CDR1*/*CDR2*/*CDR11*,

two genes are related to *CDR4*, and two genes are related to *SNQ2*; *C. auris* B11221 has an

additional copy of *SNQ2* (**Figure S10**). The *TAC1* transcription factor that regulates expression

of *CDR1* and *CDR2* in *C. albicans* is present in two tandem copies in *C. auris*, *C. haemulonii*, *C.*

duobushaemulonii, and *C. pseudohaemulonii* (**Table 2**).

While many gene families involved in pathogenesis in *C. albicans* are present in similar

numbers in *C. auris*, *C. haemulonii*, *C. duobushaemulonii,* and *C. pseudohaemulonii*, there are

some notable differences such as of cell wall and transmembrane proteins. We identified similar

numbers of the secreted aspartyl proteases, lipases and oligopeptide transporters (*OPT*), and

only one copy of the *ALS* cell surface family of *C. albicans* (**Table 2**). We also examined

whether other genes involved in the *C. albicans* core filamentation response were conserved in

the emerging multidrug-resistant species (**Table S4**). While most of these genes are conserved

in *C. auris* and closely related species, two genes are absent, candidalysin (*ECE1*) and the

hyphal cell wall protein (*HWP1*), both of which are highly expressed in *C. albicans* hyphae[33,34].

Thus, we additionally assessed if any other cell surface families of proteins are enriched in *C.*

auris and closely related species. We found a total of 75 genes with a predicted GPI anchor,

including genes that were found only in the emerging multidrug resistant clade, including one

unique family expanded in *C. auris* (**Table S4**). The most represented protein family domains in

these genes included the N-terminal cell wall domain, the aspartyl protease domain, and the

fungal specific cysteine rich (CFEM) domain (**Table S4**). The shared profile of these genes

across *C. auris* and other MDR species suggests that the more rarely observed species may be

similarly primed to become more common human pathogens.

Transcriptional analysis to voriconazole and amphotericin B in *C. auris* susceptible and

resistant strains

To investigate which *C. auris* genes are involved in antifungal resistance, we carried out

RNA-Seq of *C. auris* strains B8441 and B11210 (clade I) to profile gene expression changes

after exposure to two antifungal drugs, voriconazole (VCZ) and amphotericin B (AMB). B11210

is resistant to AMB and exhibits an elevated MIC to VCZ, while B8441 is susceptible to AMB

and displays a low MIC to VCZ (**Table S1**; [2]). We identified differentially expressed genes

(DEGs) in B8441 and B11210 after 2 and 4 hours of drug exposure (**Methods; Table S8**;

Figure S11a). The response of the sensitive isolate B8441 to AMB or VCZ involved increased
expression of small sets of genes: 39 genes were induced in response to AMB, 21 in response
to VCZ, of which 14 were induced by both drugs (fold change (FC) > 2; false discovery rate
(FDR) < 0.05; **Figure S11b**). Genes induced upon AMB exposure were enriched in small
molecule biosynthetic process and iron homeostasis (enriched GO terms corrected-*P* < 0.05,
hypergeometric distribution with Bonferroni correction; **Table S9**). Notably, this set includes
genes involved in the *C. albicans* transcriptional response to AMB associated with transport and
with lipid, fatty acid, and sterol metabolism[35] including genes involved in arginine synthesis
(*ARG1/ARG3*), ergosterol biosynthesis (*ERG24*), fatty-acid metabolism (*FAS1/FAS2*), GPI-
linked surface proteins (*PGA7* and *RBT5*), and several iron transporters (class *FTH1* and *SIT1*)
(**Table S9; Figure S11c**). Three of these genes (*SIT1*, *PGA7* and *RBT5*) have also been found
to be upregulated during *C. auris* biofilm formation[25], suggesting that cell wall reorganization may
be part of the drug response. Genes induced in B8441 in response to both AMB and VCZ were
enriched in transmembrane transport and iron transport categories (enriched GO terms
corrected-*P* < 0.05, hypergeometric distribution with Bonferroni correction; **Table S8**), including
a ferric reductase (*FRP1*), a high affinity iron transporter (*FTH1*), a glucose transporter (*HGT7*),
a N-acetylglucosamine transporter (*NGT1*) and an oligopeptide transporter (*OPT1-like*; **Table
S9**). Genes only induced with VCZ included the oligopeptide transporter *PTR22.* We also
examined expression differences in the resistant isolate B11210. As a large set of genes was
identified as differentially expressed compared to the control sample, we focused on the most
highly induced or repressed genes (FC > 4; FDR < 0.001; **Figure S11a**). A total of 106 genes
were induced in response to AMB and 41 genes in response to VCZ, of which 40 were
commonly induced (**Figure S11b**). Genes induced in response to AMB were enriched in
transcription and translation processes and in sterol biosynthetic process (enriched GO terms
corrected-*P* < 0.05, hypergeometric distribution with Bonferroni correction). Notably, 5 of the 21
genes involved in the ergosterol biosynthesis pathway were highly induced in B11210, including
MVD, *ERG2*, *ERG1*, *ERG6*, and *ERG13* (**Table S9**; **Figure S11b**; **Figure S12**). This correlates
with the maintenance of cell membrane stability, as previously noted for the transcriptional
response of *C. albicans* to AMB, since AMB binds to ergosterol in the cell membrane[35]. Genes
induced by both drugs in B11210 are enriched in amide biosynthetic process and translation,
and genes only induced by VCZ are enriched in RNA processing and transcription (enriched GO
terms corrected-*P* < 0.05, hypergeometric distribution with Bonferroni correction; **Table S9**).
405
Since *C. auris* isolates B8441 and B11210 (clade I) had disparate resistance phenotypes

predominantly in response to AMB, we further examined expression changes between these
two strains. Despite the fact that B8441 and B11210 are from the same clade (clade I) only 8
genes were induced in both strains upon AMB treatment, including *ARG1*, *CSA1* and *MET15*,
and one *OPT1*-like transporter (B9J08_001998) (**Table S10**). In addition, comparison of both
isolates before treatment revealed that the resistant strain B11210 has higher expression of
genes previously noted to be involved in the *C. albicans* transcriptional response to drug
exposure relative to B8441 even in the absence of drug (**Table S11**). This set of genes
encompasses the D-xylulose reductase (*XYL2*)[35], Phosphoenolpyruvate carboxykinase
(*PCK1*)[26,35], and a large set of transporters (*FRP1*, *FTH1*, *HGT13*, *HGT7*, *HSP70*, *NGT1*, *OPT1*,
PTR22, *SEC26*)[35], which suggests intrinsic expression of genes associated with polyene
resistance in B11210 (**Table S10**). In addition, we observed that the expression levels of
homologs of multidrug transporters (*CDR1*, *CDR4a*, *CDR4b*, *MDR1*, *SNQ2*) did not significantly
vary across exposure to either drug or between isolates. Of these, *CDR4a* appears more highly
transcribed relative to other multidrug transporters (*CDR1*, *MDR1* or *SNQ2*). This supports
previous findings of intrinsic higher expression of transporters as a possible mechanism of drug
resistance in *C. auris*[12].

We compared these results to SNP variants identified between B8441 and B11210 to evaluate if
any may explain the phenotypic difference in drug resistance in these strains. We found a total
of 1,148 SNPs and 430 insertion or deletion events (**Table S11**). A total of 15 genes display a
loss of function mutation (frame-shift, stop gained or stop lost), including the stationary phase
protein (SNZ1) and the putative GPI-anchored adhesin-like protein (*HYR3*). We did not find loss
of function mutations in genes previously associated with amphotericin B resistance in *C.*
albicans, however, we did identify non-synonymous mutations in genes associated with drug
resistance and response to amphotericin B, including *ERG11*, *ERG4*, *CDR1, C5_05230C_A,*
NHP6A and other transporters such as *HGT7* and *YCF1,* (**Table S11**). These variants suggest
candidate mutations that could explain phenotypic variation in drug resistance, however a wider
association study of mutations and drug resistance levels would help identify the major genes
involved across the different clades.

While many genes associated with antifungal resistance are conserved across *C. auris* and
CTG species, unique genes and gene duplication in *C. auris* may also contribute to the
underlying emergence of multidrug resistant phenotypes. Some genes induced by AMB or VCZ
in *C. auris* B8441 and B11210 (**Table S10**) were specific to *C. auris* or to the emerging clade of

C. auris, *C. haemulonii*, *C. duobushaemulonii,* and *C. pseudohaemulonii*. This includes five
ortholog families unique to *C. auris*, comprising three putative GPI-anchored cell wall proteins,
two homologs of *IFF6* (a putative GPI-anchored adhesin-like protein), one homolog of *PGA54*,
and an aspartyl protease similar to *SAP8* (**Table S9**). In addition, this set included gene families
that are expanded in *C. auris* or in closely related species (**Figures 3b** and **4**), including
transporters (four *SIT*-like, one *FTR*-like, and two *OPT*-like class), cell wall adhesins, and
several predicted secreted proteins (**Table S9**). Notably, while other Saccharomycetales
species including *C. albicans* only have one copy of *SIT1* that is induced during AMB treatment,
C. auris and closely related species had up to 11 *SIT1*-like genes, of which 4 are induced during
AMB treatment (**Figures 3b**, **4**, **S6**). Together, these transcriptional changes highlighted shared
and *C. auris*-specific genes that might contribute to the MDR phenotype observed in *C. auris*
isolates and provided candidate genes to further investigate *C. auris* multidrug-resistance.
453
454

Discussion
As an emerging pathogen, *Candida auris* has not been well studied to date, highlighting
the need for rapidly closing this knowledge gap to respond to the increasing number of fatal
infections. There is also a limitation on how much of the biology of *C. auris* we can infer from
related *Candida* species; *C. auris* is distantly related to the two most commonly observed
pathogenic *Candida* species, *C. albicans* and *C. glabrata*, as well as other sequenced species.
Our comparative genomic analyses, incorporating new genomic data for more closely related,
multidrug-resistant species, revealed the recent evolution of this group of emerging pathogens
including shared properties that underlie antifungal resistance and virulence. In addition to *C.*
auris clades, we generated annotated genomes for three other closely related species rarely
observed as infecting humans. Building on prior studies of individual loci[4,5,12,13], the phylogenetic
relationship of these species was more clearly resolved by whole genome comparisons; *C.*
haemulonii, *C. duobushaemulonii*, and *C. pseudohaemulonii* are more closely related to each
other than to *C. auris*, with the closest relationship being between *C. duobushaemulonii*, and *C.*
pseudohaemulonii. Our phylogenetic analysis integrating the genomes of these species with
other *Candida* highlight the distant relationship of this group to other pathogenic *Candida*
species and the placement of these species within the CTG clade.
472

To characterize mechanisms that may contribute to virulence and drug resistance, we
compared the gene content between the emerging multidrug resistant species and other related

Candida. Recent work found that virulence in *C. auris* appears similar to *C. albicans* and *C.*
glabrata[36], suggesting that shared gene content could play a role. *C. auris* shares some notable
gene family expansions described in *C. albicans* and related pathogens[23], including of
transporters and secreted lipases. While an expansion of transporters is shared, species-
specific expansions have contributed to the diversification of transporters in *C. auris* and closely
related species. Similarly, the expansion of lipases suggests this could be part of a shared
mechanism of virulence; however, the roles of specific genes will need to be investigated. In
contrast, expansions of cell wall families detected in *C. albicans* and related pathogens are not
found in *C. auris* and related species. For example, the *ALS* family is represented by two to four
copies in *C. auris* and related MDR species, compared to the eight copies present in *C.*
albicans. Examining the predicted cell wall proteins in *C. auris* did not reveal any highly
expanded families; perhaps the longer history of association with humans has impacted
diversification of such cell wall protein families in the more commonly observed pathogenic
Candida species.

Drug resistance in *C. auris* likely involves mechanisms previously described in *C. albicans*,
however the specific transporters involved in drug response are less well conserved. All four
species are resistant to antifungal drugs, and the Y132F mutation in the azole drug target
ERG11, previously described in *C. auris*, and was also detected in *C. pseudohaemulonii*. The
direct link between these mutations and azole resistance in *C. auris* is supported by recent work
that found that expression of either the Y132F and K143R *C. auris ERG11* allele led to an
increase in azole resistance in a heterologous system[17]. In addition, we find evidence of
increased copy number of *ERG11* in two *C. auris* isolates, but little evidence of multiple isolates
with regions of copy number variation, and no evidence of aneuploidy. This suggests that
increased copy number of *ERG11* may be a mechanism of drug resistance in *C. auris*, as has
been described in *C. albicans*[37]. However other mechanisms of drug resistance may vary
between species and strains. For example, we found that efflux proteins such as the *CDR* family
were not induced during voriconazole treatment, instead other transporters were up-regulated,
including those that are either expanded or unique in *C. auris* and closely multidrug emergent
species, highlighting that different molecular mechanisms are likely involved in the drug
response.

Our analysis identified representative *C. auris* isolates for each of the two mating types,
suggesting that this species has the potential to undergo mating and meiosis. Each of the four

described *C. auris* clades consists of isolates that are all *MTLa* or *MTLα;* as the clades are
geographically restricted, this suggests that there is a geographic barrier to opposite sex mating.
This expectation would change if wider sampling demonstrated the establishment of both
mating types within a geographic area. The potential for mating within this species is supported
by the conservation of genes involved in mating and meiosis; these patterns are similar to that
of the related species *C. lusitaniae*, for which mating and recombination have been
demonstrated[24]. Isolates from clades of opposite mating types could be directly tested for
mating and production of progeny. One potential barrier to mating between clades is the
presence of chromosomal rearrangements, as some recombination events may result in
inviable progeny.
519
Analyses of these genomes have revealed fundamental aspects of these emerging multidrug
resistant fungi. The role of specific genes in mating, drug resistance or pathogenesis needs to
be directly tested, utilizing gene deletion technologies recently adapted for *C. auris* (e.g. [38]).
Further analysis of this data will not only advance our understanding of the basis of drug
resistance and virulence of this pathogen but can also inform development of fungal diagnostics
for accurate tracking of these emerging pathogens.
526
527

Methods
Selected isolates and genome sequencing
Strains used in this study were described previously[2,18,19]; brief details are presented in
Table 1. Isolates were grown on Sabouard Dextrose media supplemented with chloramphenicol
and gentamycin and incubated for 24-48 hours at 37°C. For Illumina sequencing, genomic DNA
was extracted using the *Quick*-DNA™ (ZR) Fungal/Bacterial Miniprep Kit (Zymo Research,
Irvine, CA, USA). Genomic libraries were constructed and barcoded using the NEBNext Ultra
DNA Library Prep kit (New England Biolabs, Ipswich, MA, USA) by following manufacturer's
instructions. Genomic libraries were sequenced using either Illumina HiSeq 2500 with HiSeq
Rapid SBS Kit v2 or Illumina MiSeq platform using MiSeq Reagent Kit v2 (Illumina, San Diego,
CA, USA). For PacBio sequencing, DNA was extracted using MasterPure™ Yeast DNA
Purification Kit (Epicenter, Madison, WI, USA). Single-molecule real-time (SMRT) sequencing
was done using the PacBio RS II SMRT DNA sequencing system (Pacific Biosciences, Menlo
Park, CA, USA). Specifically, 20-kb libraries were generated with the SMRTbell Template Prep
Kit 1.0 (Pacific Biosciences). Libraries were bound to polymerase using the DNA/Polymerase

Binding Kit P6v2 (Pacific Biosciences), loaded on two SMRTcells (Pacific Biosciences), and
sequenced with C4v2 chemistry (Pacific Biosciences) for 360 min movies.

Genome assemblies

The genomes of B8441 and B11221 *C. auris* isolates were assembled as described[2]. *C.*
haemulonii and *C. duobushaemulonii* genomes were assembled using Canu v1.6[39]. The
resultant contigs were checked for further joins and circularity using Circlator v1.5[40]. The final
contigs were polished using Quiver, part of SmrtAnalysis suite v2.3 (Pacific Biosciences)[41]. The
sequence order for the chromosomes was verified using restriction enzyme AflII Whole Genome
Mapping (OpGen, Gaithersburg, MA).

The sequenced Illumina reads of *C. auris* (B11220 and B11243), and *C. pseudohaemulonii*
(strain B12108) were assembled using the SPAdes assembler v3.1.1[42]. Next, Pilon v1.16[43] was
used to polish the best assembly of each isolate, resolving single nucleotide errors (SNPs),
artifactual indels and local mis-assemblies. All genome assemblies were evaluated using the
GAEMR package (http://software.broadinstitute.org/software/gaemr/), which revealed no
aberrant regions of coverage, GC content or contigs with sequence similarity suggestive of
contamination. Scaffolds representing the mitochondrial genome were separated out from the
nuclear assembly. All genome assemblies have been deposited at deposited at
DDBJ/EMBL/GenBank (see ***Data availability statement***). To address if any strain representing
each *C. auris* clade could be uniformly diploid, we examined candidate heterozygous positions
predicted by Pilon v1.12[43] using mapped Illumina data. The low frequency and absence of such
positions supported that all sequenced genomes are homozygous haploids.

Optical mapping

Two strains of *C. auris* (B11221 and B8441) were compared by the OpGen optical
mapping platform (OpGen, Inc., Gaithersburg, Maryland). High molecular weight genomic DNA
from overnight grown cells were purified with Argus HMW DNA Isolation Kit (OpGen, Inc.) and
examined for quality and concentration using the ARGUS QCards(OpGen, Inc.). The software
program Enzyme Chooser (OpGen, Inc.) identified *BamHI* restriction endonuclease to be
optimal for optical map production, because its cleavage of reference genomes would result in
fragments that average 6–12 kbp in size, with no fragments larger than 80 kbp. Single genomic
DNA fragments were loaded onto a glass surface of a MapCard (OpGen, Inc.) using the
microfluidic device, washed and then digested with *BamHI* restriction enzyme, and stained with

JOJO-1 dye through the ARGUS MapCard Processor (OpGen, Inc.). Map cards were scanned
and analyzed by automated fluorescent microscopy using the ARGUS Whole Genome Mapper
(OpGen, Inc.). The single molecule restriction map collections were then tiled according to
overlapping fragment patterns to produce a consensus whole genome map. This map was
imported into MapSolver (OpGen, Inc.) along with predicted *in silico* maps of contigs derived
from WGS, using the same restriction enzyme for ordering and orientation of contigs during
genome circularization. *In-silico* predicted optical maps of complete genomes were scaled
according to the size of sequenced genomes to show identity with Optical maps (**Figures S1**).
585

RNA-Seq of B8441 and B11210 during drug treatment
For RNA extraction *C. auris* cells were grown in YPD broth medium (Difco Laboratories,
Sparks, MD) at 30°C in a shaking incubator at 300 rpm. After 18 hours, the stationary phase
cells were diluted with the equal volume of fresh YPD broth and incubated for two hours at 37°
to induce growth. After that, the cells were treated with amphotericin B at the final concentration
of 0.25 µg/mL or voriconazole at the final concentration of 1 µg/mL (Sigma-Aldrich, St. Lois,
MO) and incubated at 30°C for additional 2 and 4 hours in the presence of drug. Cells were
centrifuged for 2 minutes at 12000xg, pellets were flash frozen in dry ice/ethanol bath and
stored at -80°C. RNA was isolated using RiboPure™-Yeast rapid RNA isolation kit (Life
technologies, Carlsbad, CA) using the manufacturer's protocol. RNA was adapted for
sequencing using the RNAtag-Seq approach[44], with the modification that the yeast RiboZero
reagent was used for rRNA depletion. For each condition, two biological replicates were
performed, and the read counts per transcript were highly correlated between replicates (R>
0.90).
600

Gene annotation
Gene annotation in *C. auris* was performed using RNA-Seq paired-end reads to improve
gene calling and structure predictions. Briefly, we mapped RNA-Seq reads to the genome
assembly using Tophat2, and use the alignments to predict genes using BRAKER1[45], that
combines GeneMark-ET[46] and AUGUSTUS[47], incorporating RNA-Seq data into unsupervised
training and subsequently generates *ab initio* gene predictions. Additionally, we re-annotated
the genome of the 6684 strain[13] improving its gene set and predicted gene structures. tRNAs
were predicted using tRNAscan[48] and rRNAs predicted using RNAmmer[49]. Genes containing
PFAM domains found in repetitive elements or overlapping tRNA/rRNA features were removed.
Genes were named and numbered sequentially. For the protein coding-gene name assignment

we combined HMMER PFAM/TIGRFAM, Swissprot and Kegg products. For comparative

analysis genes were functionally annotated by assigning PFAM domains, GO terms, and KEGG

classification. HMMER3[50] was used to identify PFAM domains using release 27. GO terms were

assigned using Blast2GO[51], with a minimum e-value of 1×10^{-10}. Protein kinases were identified

using Kinannote[52] and transporter families using TCDB version 01-05-2017[53]. To evaluate the

completeness of predicted gene sets, the representation of core eukaryotic genes was analyzed

using CEGMA genes[54] and BUSCO[55].

Copy number variation and nucleotide diversity

To identify regions in *C. auris* that exhibit copy number variation (CNV) we analyzed

Illumina read depth of 47 previously sequenced isolates. We identified genomic windows (1 kb)

showing significant variation (*p*-value < 0.01) in normalized read depth using CNVnator v0.3[56].

To measure nucleotide diversity we used SNPs identified from the isolate sequences of

Lockhart *et al*[2]. We computed genome-wide π using VCFtools v0.1.12[57] for non-overlapping

sliding windows of 5 kb for *C. auris* clade I, II, II and all isolates.

Comparative genomics and phylogenomic analysis

To examine the phylogenetic relationship of the emerging multidrug-resistance clade,

including *C. auris*, *C. haemulonii*, *C. duobushaemulonii,* and *C. pseudohaemulonii* we identified

single copy orthologs in these sequenced genomes and twelve related species using OrthoMCL

v1.4[58] (Markov index 1.5; maximum e-value 1e-5). A total of 1,570 protein-coding genes that

have one single copy and are conserved in all 20 genomes were aligned using MUSCLE, and a

phylogeny was estimated from the concatenated alignments using RAxML v7.7.8[59] with model

PROTCATWAG with a total of 1,000 bootstrap replicates. To compare gene family expansion

and contractions, we used orthologous gene clusters we classified as core, auxiliary and

unique. We then searched for expansions or contractions in functionally classified genes by

assigning PFAM domains, GO terms, and KEGG classification. Using a matrix of gene class

counts for each classification type, we identified enrichment comparing the emerging multidrug-

resistance clade with all the other related species using Fisher's exact test. Fisher's exact test

was used to detect enrichment of PFAM, KEGG, GO terms, and transporter families between

groups of interest, and p-values were corrected for multiple comparisons[60]. Significant

(corrected *P*-value < 0.05) gene class expansions or depletions were examined for different
comparisons.

Transcriptional analysis of *C. auris* RNA-Seq

RNA-Seq reads were aligned to the transcript sequences of *C. auris* B8441 or B11221
using Bowtie2[61]. Transcript abundance was estimated using RSEM (RNA-Seq by expectation
maximization; v.1.2.21) as transcripts per million (TPM). TPM-normalized 'transcripts per million
transcripts' (TPM) for each transcript were calculated, and differentially expressed transcripts
were identified using edgeR[62], all as implemented in the Trinity package version 2.1.1[63]. Genes
were considered differentially expressed only if they had a 2-fold change difference (> 2 FC) in
TPM values and a false discovery rate below or equal to 0.05 (FDR < 0.05), unless specified
otherwise. To determine major patterns of antifungal-response specific we clustered gene
expression patterns by *k-means*. To identify functional enrichment of differentially expressed
genes, we used functional gene assignations from PFAM, GO terms and KEGG (see **Gene**
annotation), and then performed comparisons with Fisher's exact test. To identify possible
functions of the gene products of significantly differentially expressed drug resistance genes,
protein homologs were assigned based on orthology, functional assignments (GO, PFAM,
TIGRFAM) and experimental evidence from *Candida* genome database
(http://www.candidagenome.org).

Antifungal susceptibility testing

Antifungal susceptibility testing was performed on isolates according to Clinical and
Laboratory Standards Institute (CLSI) guidelines[64]. Custom prepared microdilution plates (Trek
Diagnostics, Oakwood Village, OH, USA) were used for generating both azole and
echinocandin susceptibilities while Etest® strips (BioMerieux, Marcy l'Etoile, France) were used
for amphotericin B susceptibilities. Interpretive breakpoints for *C. auris* and related species do
not exist but have been conservatively based on those breakpoints established for other
Candida species.

Data availability statement

All genome assemblies and gene annotations have been deposited at
DDBJ/EMBL/GenBank under the following accession numbers: *C. auris* B8441 PEKT00000000;
C. auris B11221 PGLS00000000, *C. auris* B11220 PYFR00000000, *C. auris* B11243
PYGM00000000, *C. haemulonii* B11899 PKFO00000000, *C. duobushaemulonii* B09383

PKFP00000000, *C. pseudohaemulonii* B12108 PYFQ00000000. All whole genome sequence
data have been deposited in NCBI under the following BioProjects: *C. auris* PRJNA328792, *C.*
haemulonii PRJNA421961, *C. duobushaemulonii* PRJNA421966, *C. pseudohaemulonii*
PRJNA438484.The RNA-Seq data from *C. auris* has been deposited at NCBI under BioProject
PRJNA445471.

Acknowledgements

We thank Sarah Young and Paul Cao for their assistance with genome annotation, the
Broad Technology labs for RNAtag-Seq library construction, and the Broad Genomics Platform
for Illumina sequencing. This project has been funded in whole or in part with Federal funds
from the National Institute of Allergy and Infectious Diseases, National Institutes of Health,
Department of Health and Human Services, under award N°: U19AI110818. The use of product
names in this manuscript does not imply their endorsement by the US Department of Health and
Human Services.

Disclaimer

The finding and conclusions in this article are those of the authors and do not
necessarily represent the views of the CDC.

Author contributions

Conceived and designed the study: APL CAC. Performed experiments: LG NAC VNL
ELB PJ. Performed the assembly and annotation: JFM. Analyzed the data: JFM VNL PJ RAF
CAC. Wrote the paper: JFM CAC.

Additional information

Competing interests: The authors declare no competing interests.

 Tables and Figures

707   **Table 1.** Genome assembly statistics of *Candida auris* and closely related species*

Species	*Candida auris*				*C. hae*	*C. duo*	*C. pseu*
Strain	B8441	B11221	B11220	B11243	B11899	B09383	B12108
Clade	I	III	II	IV	.	.	
Country of origin	Pakistan	South Africa	Japan	Venezuela	Israel	USA	Venezuela
Total assembly size (Mb)	12.4	12.7	12.1	12.3	13.3	12,3	12.6
Chromosomes	7	7	.	.	.	.	
Assembly anchored (%)	98.8	97.2	.	.	.	.	
Scaffolds	15	20	324	240	11	7	36
Contigs	18	23	324	265	11	7	41
Scaffold N50 (Mb)	1.1	2.4	0.06	0.09	1.7	3.3	0.64
Scaffold N90 (kb)	777	949	19.5	27.1	952	788	227
GC content (%)	45.2	45.3	45.0	45.0	45.3	46.9	47.2
Protein coding genes	5,421	5,527	5,546	5,601	5,410	5,331	5,288

708   **C. hae = C. haemulonii; C. duo = C. duobushaemulonii; C. pseu = C. pseudohaemulonii*

Table 2. Conservation of genes involved in pathogenesis and drug resistance*

Category	C. albicans	Candida auris (B8441 - I)	Candida auris (B11221 - III)	Candida auris (B11220 - II)	Candida auris (B11243 - IV)	Candida auris (6684 - I)	Candida haemulonii	Candida pseudohaemulonii	Candida duobushaeumulonii	Candida lusitaniae	Debaryomyces hansenii	Candida guilliermondii	Candida tropicalis	Candida albicans	Candida dubliniensis	Candida orthopsilosis	Candida parapsilosis	Lodderomyces elongisporus	Candida krusei	Candida glabrata	Saccharomyces cerevisiae
Drug resistance	ERG11	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	TAC1	2	2	2	2	2	2	2	2	2	2	1	1	1	1	1	1	1	1	1	0	0
	UPC2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2
	MDR1	1	1	1	1	1	1	1	1	1	3	6	2	1	1	3	2	4	0	1	1
	SNQ2 CDR4 CDR2 CDR11 CDR1	5	6	5	5	5	5	5	4	3	6	8	2	5	5	3	6	5	9	5	8
Secreted aspartyl proteinases	SAP1 SAP2 SAP8 SAP3	3	3	2	3	3	2	2	2	2	0	4	3	4	4	3	2	1	0	0	0
	SAP9	1	1	1	1	1	1	1	1	1	2	1	1	1	1	2	5	1	2	1	3
Secreted lipases	LIP4, LIP9 LIP5, LIP8 LIP2, LIP1 LIP10, LIP6 LIP3	8	8	8	8	8	9	9	9	0	1	1	2	9	9	2	2	1	0	0	0
Cell wall adhesins	ALS2, ALS5 ALS1, ALS9 ALS4, ALS3 ALS7, ALS6	1	1	1	1	1	2	3	3	1	0	2	4	8	6	1	2	0	0	0	0
	ALS-like	2	2	1	2	2	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0

712 * Roman numerals represent *C. auris* clades.

Figure 1. Whole genome conservation, structure and synteny. **(a)** Genome wide gene synteny among *Candida auris*, *C. haemulonii*, *C. duobushaemulonii*, and *C. pseudohaemulonii*. To determine synteny regions of conserved gene order we identified orthologs between these isolates and then plotted chains of orthologs (grey lines). Isolate names are shown to the left of their genomes, which are represented by horizontal black lines, with vertical lines indicating scaffold borders, and their identifiers listed above (+/−□=□orientation). **(b)** Shared synteny regions based on whole genome alignments between *C. auris* B8441 and B11221.

Figure 2. Identification of Mating-type loci (*MTL*) in *Candida auris* and closely related species. **(a)** Synteny schema depicting orientation and conservation of the color-coded *MTL* idiomorphs and genes adjacent to the *MTL*. The putative MTL in *C. auris*, *C. haemulonii*, *C. duobushaemulonii*, and *C. pseudohaemulonii* are shown in comparison with the *MTL**a*** and *MTL*α idiomorphs from *C. lusitaniae*. **(b)** Phylogenetic analysis of the non-mating flanking genes (*PIK1/OBP1/PAP1*) showing the inheritance of idiomorphs of these genes within the *MTL**a*** and *MTL*α loci. Branch lengths indicate the mean number of changes per site. **(c)** Phylogenetic tree of *C. auris* isolates from Lockhart et al. 2016. Isolates are color-coded according the clades (I, II, III and IV) and mating type (*MTL**a*** and *MTL*α). Figure S3 shows isolates, origin and the normalized depth read coverage of mapped positions for all isolates aligned to B8441 (*MTL**a***) and B11221 (*MTL*α), supporting the classification into *MTL**a*** or *MTL*α.

Figure 3. Phylogenomic and gene family changes across *Candida auris* and closely related species. **(a)** Maximum likelihood phylogeny using 1,570 core genes based on 1,000 replicates, among 20 annotated genome assemblies, including *Candida auris, C. haemulonii* (B11899), *C. duobushaemulonii* (B09383), and *C. pseudohaemulonii* (B12108), and closely related species. Branch lengths indicate the mean number of changes per site. **(b)** Heatmap depicting results of protein family enrichment analysis (PFAM domains; corrected *p-value* < 0.05) comparing the gene content of *C. auris* strains representing each clade, *C. haemulonii*, *C. duobushaemulonii*, and *C. pseudohaemulonii*, and other closely related species, including *C. lusitaniae*, *C. albicans*, *C. krusei* and *C. glabrata*. Values are colored along a blue (low counts) to red (high counts) color scale, with color scaling relative to the low and high values of each row. Each protein family domain has a color code (right) indicating whether expanded or depleted.

Figure 4. Phylogenetic relationships of oligopeptide transporters (*OPT*) and siderophore iron
transporters (*SIT*) families. Phylogenetic trees estimated by maximum likelihood with RAxML
showing expansion of *OPT* **(a)** and *SIT* **(b)** transporter families in *C. auris* and *C. haemulonii*, *C.*
duobushaemulonii and *C. pseudohaemulonii*. Each species has a color code and lineage-
specific expansions (blue branches) can be seen in *C. auris* and closely related species relative
to the close ancestor *C. lusitaniae* and *C. albicans*. Orthologs of *OPT* and *SIT* transporters in *C.*
albicans are depicted alongside each tree. **(c)** Chromosome view depicting genes and
orientation located in chromosome 6 (B8441 scaffold05). This region highlights expansion and
tandem duplication of eight *OPT* class transporters (in red).
759

References

1. Clancy, C. J. & Nguyen, M. H. Emergence of *Candida auris*: An International Call to Arms.

Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am. **64,** 141–143 (2017).

2. Lockhart, S. R. *et al.* Simultaneous Emergence of Multidrug-Resistant *Candida auris* on 3

Continents Confirmed by Whole-Genome Sequencing and Epidemiological Analyses. *Clin*

Infect Dis (2016). doi:10.1093/cid/ciw691

3. Schelenz, S. *et al.* First hospital outbreak of the globally emerging *Candida auris* in a

European hospital. *Antimicrob. Resist. Infect. Control* **5,** (2016).

4. Cendejas-Bueno, E. *et al.* Reclassification of the *Candida haemulonii* Complex as *Candida*

haemulonii (*C. haemulonii* Group I), *C. duobushaemulonii* sp. nov. (*C. haemulonii* Group II),

and *C. haemulonii var. vulnera var. nov.*: Three Multiresistant Human Pathogenic Yeasts. *J.*

Clin. Microbiol. **50,** 3641–3651 (2012).

5. Kumar, A. *et al. Candida haemulonii* species complex: an emerging species in India and its

genetic diversity assessed with multilocus sequence and amplified fragment-length

polymorphism analyses. *Emerg. Microbes Infect.* **5,** e49 (2016).

6. Kathuria, S. *et al.* Multidrug-Resistant *Candida auris* Misidentified as *Candida haemulonii*:

Characterization by Matrix-Assisted Laser Desorption Ionization–Time of Flight Mass

Spectrometry and DNA Sequencing and Its Antifungal Susceptibility Profile Variability by

Vitek 2, CLSI Broth Microdilution, and Etest Method. *J. Clin. Microbiol.* **53,** 1823–1830

(2015).

7. Rhodes, J. *et al.* Genomic epidemiology of the UK outbreak of the emerging human fungal

pathogen *Candida auris*. *Emerg. Microbes Infect.* **7,** 43 (2018).

8. Borman, A. M., Szekely, A. & Johnson, E. M. Isolates of the emerging pathogen *Candida*

auris present in the UK have several geographic origins. *Med. Mycol.* **55,** 563–567 (2017).

9. Sharma, C., Kumar, N., Pandey, R., Meis, J. F. & Chowdhary, A. Whole genome

sequencing of emerging multidrug resistant *Candida auris* isolates in India demonstrates

low genetic variation. *New Microbes New Infect.* **13,** 77–82 (2016).

10. Chowdhary, A. *et al.* Multidrug-resistant endemic clonal strain of *Candida auris* in India. *Eur.*

J. Clin. Microbiol. Infect. Dis. **33,** 919–926 (2014).

11. Prakash, A. *et al.* Evidence of genotypic diversity among *Candida auris* isolates by

multilocus sequence typing, matrix-assisted laser desorption ionization time-of-flight mass

spectrometry and amplified fragment length polymorphism. *Clin. Microbiol. Infect. Off. Publ.*

Eur. Soc. Clin. Microbiol. Infect. Dis. **22,** 277.e1–9 (2016).

12. Ben-Ami, R. *et al.* Multidrug-Resistant *Candida haemulonii* and *C. auris*, Tel Aviv, Israel.

Emerg. Infect. Dis. **23,** 195–203 (2017).

13. Chatterjee, S. *et al.* Draft genome of a commonly misdiagnosed multidrug resistant

pathogen *Candida auris. BMC Genomics* **16,** 686 (2015).

14. Gargeya, I. B., Pruitt, W. R., Simmons, R. B., Meyer, S. A. & Ahearn, D. G. Occurrence of

Clavispora lusitaniae, the teleomorph of *Candida lusitaniae*, among clinical isolates. *J. Clin.*

Microbiol. **28,** 2224–2227 (1990).

15. Gabaldón, T., Naranjo-Ortíz, M. A. & Marcet-Houben, M. Evolutionary genomics of yeast

pathogens in the Saccharomycotina. *FEMS Yeast Res.* **16,** (2016).

16. Chowdhary, A. *et al.* A multicentre study of antifungal susceptibility patterns among 350

Candida auris isolates (2009–17) in India: role of the ERG11 and FKS1 genes in azole and

echinocandin resistance. *J. Antimicrob. Chemother.* (2018). doi:10.1093/jac/dkx480

17. Healey, K. R. *et al.* Limited ERG11 mutations identified in isolates of *Candida auris* directly

contribute to reduced azole susceptibility. *Antimicrob. Agents Chemother.* (2018).

doi:10.1128/AAC.01427-18

18. Chow, N. A. *et al.* Genome Sequence of a Multidrug-Resistant *Candida haemulonii* Isolate

from a Patient with Chronic Leg Ulcers in Israel. *Genome Announc.* **6,** e00176-18 (2018).

19. Chow, N. A. *et al.* Genome Sequence of the Amphotericin B-Resistant *Candida*

duobushaemulonii Strain, B09383. *Genome Announc.* **6,** e00204-18 (2018).

20. Desjardins, C. A. *et al.* Population genomics and the evolution of virulence in the fungal

pathogen Cryptococcus neoformans. *Genome Res.* **27,** 1207–1219 (2017).

21. Persinoti, G. F. *et al.* Whole Genome Analysis Illustrates Global Clonal Population Structure

of the Ubiquitous Dermatophyte Pathogen *Trichophyton rubrum. Genetics*

genetics.300573.2017 (2018). doi:10.1534/genetics.117.300573

22. Oh, B. J. *et al.* Biofilm formation and genotyping of *Candida haemulonii*, *Candida*

pseudohaemulonii, and a proposed new species (*Candida auris*) isolates from Korea. *Med.*

Mycol. **49,** 98–102 (2011).

23. Butler, G. *et al.* Evolution of pathogenicity and sexual reproduction in eight *Candida*

genomes. *Nature* **459,** 657–62 (2009).

24. Reedy, J. L., Floyd, A. M. & Heitman, J. Mechanistic Plasticity of Sexual Reproduction and

Meiosis in the *Candida* Pathogenic Species Complex. *Curr. Biol.* **19,** 891–899 (2009).

25. Kean, R. *et al.* Transcriptome Assembly and Profiling of *Candida auris* Reveals Novel

Insights into Biofilm-Mediated Resistance. *mSphere* **3,** e00334-18 (2018).

26. Copping, V. M. S. *et al.* Exposure of *Candida albicans* to antifungal agents affects

expression of SAP2 and SAP9 secreted proteinase genes. *J. Antimicrob. Chemother.* **55,**

645–654 (2005).

27. Reuß, O. & Morschhäuser, J. A family of oligopeptide transporters is required for growth of

Candida albicans on proteins. *Mol. Microbiol.* **60,** 795–812 (2006).

28. Hube, B. *et al.* Secreted lipases of *Candida albicans*: cloning, characterisation and

expression analysis of a new gene family with at least ten members. *Arch. Microbiol.* **174,**

362–374 (2000).

29. Erratum. *Clin. Infect. Dis.* **67,** 987–987 (2018).

30. Perea, S. *et al.* Prevalence of Molecular Mechanisms of Resistance to Azole Antifungal

Agents in *Candida albicans* Strains Displaying High-Level Fluconazole Resistance Isolated

from Human Immunodeficiency Virus-Infected Patients. *Antimicrob. Agents Chemother.* **45,**

2676–2684 (2001).

31. Gaur, M. *et al.* MFS transportome of the human pathogenic yeast *Candida albicans. BMC*

Genomics **9,** 579 (2008).

32. Prasad, R. & Goffeau, A. Yeast ATP-binding cassette transporters conferring multidrug

resistance. *Annu. Rev. Microbiol.* **66,** 39–63 (2012).

33. Sharkey, L. L., McNemar, M. D., Saporito-Irwin, S. M., Sypherd, P. S. & Fonzi, W. A. HWP1

Functions in the Morphological Development of *Candida albicans* Downstream ofEFG1,

TUP1, and RBF1. *J. Bacteriol.* **181,** 5273–5279 (1999).

34. Birse, C. E., Irwin, M. Y., Fonzi, W. A. & Sypherd, P. S. Cloning and characterization of

ECE1, a gene expressed in association with cell elongation of the dimorphic pathogen

Candida albicans. Infect. Immun. **61,** 3648–3655 (1993).

35. Liu, T. T. *et al.* Genome-wide expression profiling of the response to azole, polyene,

echinocandin, and pyrimidine antifungal agents in *Candida albicans. Antimicrob. Agents*

Chemother. **49,** 2226–2236 (2005).

36. Fakhim, H. *et al.* Comparative virulence of *Candida auris* with *Candida haemulonii*, *Candida*

glabrata and *Candida albicans* in a murine model. *Mycoses* (2018). doi:10.1111/myc.12754

37. Selmecki A., Gerami☐Nejad M., Paulson C., Forche A. & Berman J. An isochromosome

confers drug resistance in vivo by amplification of two genes, ERG11 and TAC1. *Mol.*

Microbiol. **68,** 624–641 (2008).

38. Grahl, N., Demers, E. G., Crocker, A. W. & Hogan, D. A. Use of RNA-Protein Complexes for

Genome Editing in Non-*albicans Candida* Species. *mSphere* **2,** e00218-17 (2017).

39. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptivek-mer

weighting and repeat separation. *Genome Res.* **27,** 722–736 (2017).

40. Hunt, M. *et al.* Circlator: automated circularization of genome assemblies using long

sequencing reads. *Genome Biol.* **16,** 294 (2015).

41. Chin, C.-S. *et al.* Nonhybrid, finished microbial genome assemblies from long-read SMRT

sequencing data. *Nat. Methods* **10,** 563–569 (2013).

42. Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to

single-cell sequencing. *J Comput Biol* **19,** 455–77 (2012).

43. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

and genome assembly improvement. *PLoS One* **9,** e112963 (2014).

44. Shishkin, A. A. *et al.* Simultaneous generation of many RNA-seq libraries in a single

reaction. *Nat Methods* **12,** 323–5 (2015).

45. Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M. & Stanke, M. BRAKER1:

Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS.

Bioinformatics **32,** 767–769 (2016).

46. Lomsadze, A., Burns, P. D. & Borodovsky, M. Integration of mapped RNA-Seq reads into

automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res.* **42,** e119–e119

(2014).

47. Stanke, M., Diekhans, M., Baertsch, R. & Haussler, D. Using native and syntenically

mapped cDNA alignments to improve de novo gene finding. *Bioinformatics* **24,** 637–644

(2008).

48. Lowe, T. M. & Eddy, S. R. tRNAscan-SE: a program for improved detection of transfer RNA

genes in genomic sequence. *Nucleic Acids Res.* **25,** 955–964 (1997).

49. Lagesen, K. *et al.* RNAmmer: consistent and rapid annotation of ribosomal RNA genes.

Nucleic Acids Res. **35,** 3100–3108 (2007).

50. Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7,** e1002195 (2011).

51. Conesa, A. *et al.* Blast2GO: a universal tool for annotation, visualization and analysis in

functional genomics research. *Bioinformatics* **21,** 3674–6 (2005).

52. Goldberg, J. M. *et al.* Kinannote, a computer program to identify and classify members of

the eukaryotic protein kinase superfamily. *Bioinformatics* **29,** 2387–94 (2013).

53. Saier, M. H., Jr., Tran, C. V. & Barabote, R. D. TCDB: the Transporter Classification

Database for membrane transport protein analyses and information. *Nucleic Acids Res* **34,**

D181-6 (2006).

54. Parra, G., Bradnam, K. & Korf, I. CEGMA: a pipeline to accurately annotate core genes in

eukaryotic genomes. *Bioinformatics* **23,** 1061–7 (2007).

55. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31,** 3210–3212 (2015).

56. Abyzov, A., Urban, A. E., Snyder, M. & Gerstein, M. CNVnator: an approach to discover,

genotype, and characterize typical and atypical CNVs from family and population genome

sequencing. *Genome Res.* **21,** 974–984 (2011).

57. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27,** 2156–8 (2011).

58. Li, L., Stoeckert, C. J., Jr. & Roos, D. S. OrthoMCL: identification of ortholog groups for

eukaryotic genomes. *Genome Res* **13,** 2178–89 (2003).

59. Stamatakis, A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with

thousands of taxa and mixed models. *Bioinformatics* **22,** 2688–90 (2006).

60. Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies. *Proc Natl Acad*

Sci U A **100,** 9440–5 (2003).

61. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9,**

357–359 (2012).

62. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinforma. Oxf. Engl.* **26,**

139–140 (2010).

63. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the

Trinity platform for reference generation and analysis. *Nat Protoc* **8,** 1494–512 (2013).

64. *Reference method for broth dilution antifungal susceptibility testing of yeasts: approved*

standard. (National Committee for Clinical Laboratory Standards, 2008).