Within-subject combination of diffusion tensor MRI and CLARITY reveals alterations in the social brain network resulting from α-neurexin II deletion

Eleftheria Pervolaraki Ph.D[1]*, Adam L. Tyson MRes[2,3,4]*, Francesca Pibiri MSc[5], Steven L. Poulter Ph.D[5], Amy C. Reichelt Ph.D[6], R. John Rodgers Ph.D[7], Steven J. Clapcote Ph.D[1], Colin Lever Ph.D[5], Laura C. Andreae MBBS Ph.D[2,3]*, James Dachtler Ph.D[1,5]*

[1]School of Biomedical Sciences, University of Leeds, LS2 9JT, UK

[2]Centre for Developmental Neurobiology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London SE1 1UL, UK

[3]MRC Centre for Neurodevelopmental Disorders, King's College London, London SE1 1UL, UK

[4]Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London SE5 8AF, UK

[5]Department of Psychology, Durham University, South Road, Durham, DH1 3LE, UK

[6]School of Health and Biomedical Sciences, RMIT University, VIC 3083, Australia

[7]School of Psychology, University of Leeds, LS2 9JT, UK

*These authors contributed equally.

Address for Correspondence: Dr J Dachtler, Department of Psychology, Durham University, South Road, Durham, DH1 3LE, UK.

Short title: Social brain changes revealed by DTI and CLARITY

Conflicts of Interest: The Authors declare no competing conflicts of interest.

Abstract: 241 words; **Introduction:** 619 words; **Discussion:** 1019 words; **Figures:** 5; **Pages:** 24; **Supplemental Material:** 5 figures, 3 tables.

Pervolaraki et al.

Abstract

Of the many genetic mutations known to increase the risk of autism spectrum disorder, a large proportion cluster upon synaptic proteins. One such family of presynaptic proteins are the neurexins (NRXN), and recent genetic and mouse evidence has suggested a causative role for *NRXN2* in generating altered social behaviours. Autism has been conceptualised as a disorder of atypical inter-regional connectivity, yet how single-gene mutations affect such connectivity remains under-explored. For the first time, we have developed a within-subject, quantitative analysis of white matter microstructure and connectivity leveraging diffusion tensor MRI (DTI) with high-resolution 3D imaging in optically cleared (CLARITY) brain tissue in the same mouse, applied here to the *Nrxn2α* knockout (KO) model. DTI revealed decreases in fractional anisotropy and increases in apparent diffusion coefficient in the amygdala (including the basolateral nuclei), the anterior cingulate cortex, the orbitofrontal cortex and the hippocampus. Radial diffusivity of the anterior cingulate cortex and orbitofrontal cortex was significantly increased in Nrxn2α KO mice, as were tracts between the amygdala and the orbitofrontal cortex. Using CLARITY, we find significantly altered axonal orientation in the amygdala, orbitofrontal cortex and the anterior cingulate cortex, which was unrelated to cell density. Our findings demonstrate that deleting a single neurexin gene (*Nrxn2α*) induces atypical connectivity within the social brain, likely underlying autism-related impairments. More generally, our combined within-subject DTI-plus-CLARITY approach presents a new, more sensitive method of revealing hitherto undetectable differences in the autistic brain.

Introduction

Autism is a highly heritable, common neurodevelopmental disorder. Although genetic studies have identified hundreds of implicated risk genes[1, 2], it is now clear that many of these cluster upon proteins relating to synaptic signaling[3]. A family of presynaptic proteins garnering recent interest have been the neurexins (*NRXNs*).

Mutations within all three *NRXN* genes have been linked to autism[3]. Heterozygous deletions within *NRXN2* were identified in a number of individuals with autistic phenotypes. These include an autistic boy and his father (who had severe language delay but not autism) who both had a frameshift mutation within exon 12 of *NRXN2*[4]; a 570-kb de novo deletion of 24 genes at chromosome 11q13.1, including *NRXN2*, in a 21-year old man displaying a clinical phenotype including autistic traits[5]; and a 1.6Mb deletion at chromosome region 11q12.3-11q13.1, including *NRXN2*, in a 23-year old man with intellectual disability and behavioral problems [6]. A recent de novo frameshift mutation has also been identified in a Chinese man with autism spectrum disorder (ASD)[7]. Consistent with these association studies, we and others have previously found that homozygous or heterozygous deletion of *Nrxn2α* induces impairment in social approach and social recognition[8-10].

Currently it is unknown whether deletion of *Nrxn2α* changes the brain's microstructure and connectivity. One previous study found coarse alterations to cell layer thickness within the hippocampus of *Nrxn2α* homozygous KOs[11]. However, cell density measurements are unlikely to reveal the true extent of changes within the autistic brain. Within the current study, we have addressed this by developing a dual imaging approach (DTI-plus-CLARITY) that quantifies the alignment and density of white matter, applied here to the social brain in a mouse model of autism.

Diffusion tensor MRI (or DTI) is based upon the movement of water molecules, a measure that is termed fractional anisotropy (FA). Apparent diffusion coefficient (ADC) is similar to FA, but quantifies diffusion restriction

as opposed to the spatial symmetry of diffusion. This approach has been used to explore neuropathological markers in autistic patients; alterations in myelination, axonal abundance, size and orientation all modify FA and ADC values[12-14]. Using the preferred direction of the diffusion of tensors between brain regions can be used to explore their potential connection. Quantification of those computed streamlines by FA and axial and/or radial diffusion can indicate impairments in regional connectivity. Since aberrant brain connectivity is likely a core feature of autism[15], we reasoned that the candidate method for probing the autistic brain should combine tractographic techniques. Accordingly, here, we combined high resolution imaging of labelled neuronal tracts in brains rendered transparent by CLARITY with DTI.

CLARITY is a recent development that renders tissue optically transparent and macromolecule permeable[16]. This permits antibody staining and imaging of much larger tissue volumes than possible under traditional immunofluorescence techniques. By examining fiber orientation without sectioning-related artefacts and biases, axonal staining in cleared tissue affords a deeper understanding of the microstructure and connectivity of a brain region.

Given the strong social impairments found within Nrxn2α mice, we sought to examine those brain regions most closely linked with social behavior (See Extended Data). Briefly, we identified four regions of interest (ROIs): the amygdala, and three brain regions strongly and directly connected to the amygdala; the hippocampus, orbitofrontal cortex (OFC), and anterior cingulate cortex (ACC). As predicted, connectivity was abnormal in Nrxn2α mice.

Pervolaraki et al.

Methods

Ethics

All procedures were approved by the University of Leeds and Durham University Animal Ethical and Welfare Review Boards and were performed under UK Home Office Project and Personal Licenses in accordance with the Animals (Scientific Procedures) Act 1986.

Animals

Full details of the animals, their background, genotyping and housing can be found elsewhere[8]. In brief, male B6;129-*Nrxn3tm1Sud*/*Nrxn1tm1Sud*/*Nrxn2tm1Sud*/J mice (JAX #006377) were purchased from the Jackson Laboratory and outbred once to the C57BL/6NCrl strain (Charles River, Margate, United Kingdom) to obtain mice that were individually Nrxn2α KO heterozygotes. Subsequently, HET knockout males were bred with HET females (cousin mating).

Experimental animals

6 adult wild-type males (Charles River, Margate, UK) and 6 age matched littermate Nrxn2α KO homozygotes (71 days ± 6 days old (SEM)) were perfused-fixed with 4% paraformaldehyde (PFA) in 0.1 M phosphate buffer saline (PBS) and the brains extracted. The brains were immersed in 4% PFA/0.1 M PBS for a minimum of 48 hours prior to imaging. During imaging, the samples were placed in custom-built MR-compatible tubes containing Fomblin Y (Sigma, Poole, Dorset, UK).

Data Acquisition

Image acquisition has been described elsewhere[17]. Each brain was 3D imaged using the protocol TE: 35 ms, TR: 700 ms and 10 signal averages. The field of view was set at 128 x 128 x 128, with a cubic resolution of 100 μm/pixel and a b value of 1200 s/mm^2. Further details can be found in Supplemental Materials.

Regions of Interest

Pervolaraki et al.

Our DTI approach was to undertake an *a posteriori* analysis of neural organization in regions of interest (ROIs) identified by previous literature as socially-relevant (see Supplemental Materials). We identified a canonical coronal slice (100 µm) for a given ROI from the standard mouse brain atlas (Figure 1A-D)[18]. We analysed three coronal slices centred on the canonical slice, totalling 300 µm in anterior/posterior extent.

CLARITY

Following MR imaging, the brains were washed in PBS to remove all Fomblin Y and then incubated for 7 days in hydrogel solution at 4°C prior to polymerisation at 37°C for 3.5 hours. The tissue was cut into 1.5mm coronal sections using a custom 3D-printed brain matrix[19] and incubated in clearing buffer for 24 days at 37°C with shaking. The cleared tissue was then washed in PBSTN3 (0.1% TritonX-100 and 1.5mM sodium azide in PBS) for 24 hours at room temperature and incubated in primary antibody solution (neurofilament (Aves NF-H) 1:100 in PBSTN$_3$) at 37°C with shaking for 13 days. Samples were washed, and then incubated in secondary antibody (AlexaFluor 488 goat anti-chicken IgY) as per the primary. Sections were washed again, and incubated in 3.6 µM DAPI (4',6-diamidino-2-phenylindole) followed by 85% glycerol in PBS for refractive index matching.

Cleared samples were imaged using a Zeiss 7MP multiphoton microscope at 770nm using a 20x objective lens (W Plan-Apochromat, NA 1.0, WD 1.7mm). Images (512 x 512 x 512 μm with an isotropic resolution of 520 nm) were acquired in ACC, basolateral (BLA) and basomedial amygdala and OFC) in both hemispheres. DAPI and neurofilament signal was segmented into cell nuclei and axons, and the resulting binary images were used to generate values for cell density, axonal density and axonal alignment.

Full CLARITY methodological details are available within Supplemental Materials.

Data Analysis

All data are expressed as mean ± standard error of the mean (SEM). To assess differences between genotypes within a single brain structure, data was analyzed by repeated measures two-way ANOVAs, with Sidak multiple corrections employed, or unpaired T-tests. Non-significant statistical results, particularly hemisphere comparisons, can be found in Extended Data. Statistical testing and graphs were made using GraphPad Prism version 6.

Pervolaraki et al.

Results

Nrxn2α deletion disrupts DTI measures of microstructure in social brain regions

To assess whether *Nrxn2α* deletion alters gross morphology, we quantified whole brain volume using DTI. We found total brain volume for wild-types and *Nrxn2*α KOs was similar (456.0 ± 14.76 vs. 466.2 ± 11.0 mm^3 (respectively); $t_{(10)}$ = 0.55, p = 0.59). Thus, Nrxn2α deletion does not change total brain size.

To quantitatively measure DTI, we examined FA and ADC. FA analyses changes in the linear orientation (i.e. along an axonal tract), whereas ADC (mean diffusivity) averages diffusion in all directions (i.e. the X, Y and Z axis), which is sensitive to changes such as altered alignment. The amygdala is critically important for social behaviours. To assess whether amygdalar alterations might account for social impairments in Nrxn2α KO mice, we segmented the whole amygdala structure and the basolateral nuclei along the anterior-posterior axis.

Both the anterior and posterior amygdala showed a significant reduction in FA in *Nrxn2α* KO mice (Figure 2A-B) (anterior: genotype ($F_{(1, 10)}$ = 5.81, p = 0.037); posterior: genotype ($F_{(1, 10)}$ = 11.2, p = 0.007)). This FA reduction was also observed specifically in the BLA, a region strongly associated with social behaviours (Figure 2C; genotype ($F_{(1, 10)}$ = 6.31, p = 0.03)). ADC was not significantly altered in the anterior amygdala, posterior amygdala or BLA (Figure 2D-F; all genotype: $F_{(1, 10)}$ <1).

We conducted the same analysis for the two prefrontal regions implicated in social behaviour and autism: the OFC and ACC. The pattern of results was similar for both regions: FA was not altered, while ADC was increased in the OFC (Figure 3A-B) and the ACC (Figure 3C-D). FA for the OFC was not significantly altered (genotype: ($F_{(1, 10)}$ = 3.04, p = 0.11)) but ADC was significantly increased (genotype: ($F_{(1, 10)}$ = 8.20, p = 0.017)). The ACC was unaltered in FA ($t_{(10)}$ = 1.70, p = 0.12) but had significantly increased ADC ($t_{(10)}$ = 7.52, p < 0.0001).

We sought to examine whether changes in the amygdala (Supp. Figure 1), OFC or ACC FA and ADC were driven by diffusion in the primary axis (λ_1) or the radial orientations (λ_2 and λ_3) by characterisation of AD (primary) and RD (radial). Within the OFC (Figure 3E-F), AD was significantly increased (genotype: ($F_{(1, 10)}$ = 10.74, p = 0.0083)), as was RD (genotype: ($F_{(1, 10)}$ = 18.26, p = 0.002)), suggesting that both along-tract diffusion and tract branching were affected. However, in the ACC (Figure 3G-H), only RD was significantly increased ($t_{(10)}$ = 5.65, p = 0.0002), with no alteration in AD ($t_{(10)}$ = 1.69, p = 0.123). Increased RD is thought to reflect demyelination or changes in axonal density or orientation[20].

DTI reveals altered hippocampal microstructure in *Nrxn2α* KO mice

The hippocampus has recently been associated with social motivation and social recognition. Since the specific contributions of the dorsal and ventral hippocampal poles remain unclear, we segmented the whole hippocampus into anterior (Bregma -1.94 mm), middle (Bregma -2.46 mm) (both dorsal) and posterior (Bregma -3.28 mm) (incorporating ventral regions) levels.

FA values in the anterior, mid and posterior hippocampus were not significantly altered (see Supp. Table 1 for statistics and Supp. Figure 2A-C). Similarly, ADC was unaltered for the anterior and mid hippocampus (Supp. Table 1 for statistics and Supp. Figure 2D-E), but was significantly increased in *Nrxn2α* KO mice in the posterior hippocampus (genotype: ($F_{(1, 10)}$ = 8.80, p = 0.014); Supp. Figure 2F). There were no significant genotype differences in AD in any of the hippocampal regions (Supp. Figure 3A-C). However, RD was significantly increased in the posterior hippocampus in *Nrxn2α* KO mice (genotype: ($F_{(1, 10)}$ = 10.83, p = 0.008; Supp. Figure 3F) but not in anterior and mid regions (see Supp. Table 1; Supp. Figure 3D-E). In summary, the microstructural measures most altered by *Nrxn2α* deletion were increases in ADC and RD, and both these alterations occurred in the posterior hippocampus, in line with recent work suggesting a role for ventral hippocampus in social memory[21].

DTI tractography reveals *Nrxn2α* deletion affects connectivity between the amygdala and orbitofrontal cortex

The amygdala is strongly and bidirectionally connected to both the hippocampus[22] and the OFC[23]. As all three regions are themselves important for social behaviour, and autism is thought to be related to abnormal connectivity[15], we performed tractography analysis between the amygdala (and specifically the BLA) and the hippocampus, and between the amygdala and the OFC.

From the anterior amygdala, we examined the diffusivity (AD and RD) of connections to the anterior and posterior hippocampus (Supp. Figure 4). We did not observe differences in RD for amygdala to hippocampal connections (see Supp. Table 2 for non-significant statistics). Although AD between the anterior amygdala and anterior hippocampus did not differ by genotype, there was a significant interaction between genotype and hemisphere (genotype x hemisphere ($F_{(1, 10)}$ = 12.12, p = 0.0059; Figure 4A), which post hoc analysis shows was driven by larger right-vs-left hemisphere values within the *Nrxn2α* KOs only (p = 0.012). This difference could be driven by the BLA; there was increased AD in both the BLA and anterior hippocampus connection (genotype x hemisphere ($F_{(1, 10)}$ = 10.53, p = 0.009) and the BLA and posterior hippocampus connection (genotype x hemisphere ($F_{(1, 10)}$ = 12.97, p = 0.005), which again was related to larger right-vs-left hemisphere values in the *Nrxn2α* KOs (p = 0.004 and p = 0.001, respectively (Figure 4C-D)) but not the wild-type (p = 0.87, p = 1.00, respectively). These results indicate that there are substantial differences for the connectivity of the amygdala to the hippocampus within the left and right hemisphere in *Nrxn2α* KO mice, with increased diffusivity in the right hemisphere. This finding is particularly interesting, as hemispheric differences in functional connectivity, particularly affecting connections from the right amygdala, have been found children with ASD[24, 25].

Pervolaraki et al.

Finally, we tested connections between the amygdala and the orbitofrontal cortex. For AD, wild-type and Nrxn2α KO mice did not differ by genotype, although there was a main effect of hemisphere (Figure 4E: genotype: ($F_{(1, 10)}$ = 2.85, p = 0.122), hemisphere: ($F_{(1, 10)}$ = 6.38, p = 0.03). RD was strikingly higher in *Nrxn2α* KO mice (Figure 4F: genotype: ($F_{(1, 10)}$ = 26.06, p = 0.005)), again indicative of a change in demyelination, axonal density or orientation[20].

CLARITY reveals fibre disruption in *Nrxn2α* KO mice in the amygdala, orbitofrontal cortex, and anterior cingulate cortex

To further explore the differences as revealed by DTI, we performed CLARITY on the same brain tissue used in DTI, and stained with neurofilament and DAPI to label axons and cell bodies, respectively. We were then able to derive both the axonal alignment and density of the stained fibers, in addition to the cell density.

The pattern of results was broadly similar for both the prefrontal cortical ROIs. That is, first, axonal alignment was increased in *Nrxn2α* KO mice in both the ACC (Figure 5D: genotype: ($F_{(1, 10)}$ = 16.06, p = 0.003) and the OFC (Figure 5G: genotype: ($F_{(1, 10)}$ = 5.56, p = 0.04). Second, this could not be explained by a difference in cell density, since that was similar between the KO and wild-type mice in both the ACC (Figure 5F: genotype: ($F_{(1, 10)}$ <1), hemisphere: ($F_{(1, 10)}$ = 1.73, p = 0.22) and the OFC (Figure 5H: genotype: ($F_{(1, 10)}$ = 3.09, p = 0.11). An increase in axonal density in *Nrxn2α* KO mice was reliable in the ACC (Figure 5E: genotype: ($F_{(1, 10)}$ = 14.64, p = 0.003), but not in the OFC (Figure 5H: genotype: ($F_{(1, 10)}$ = 3.09, p = 0.11).

We further examined two regions of the anterior amygdala, the BLA and basomedial (BMA) nuclei, where altered social cellular responses have been reported in human autism[26]. We did not observe any significant differences for axonal alignment or fibre density in the BLA (see Supp. Figure 5A-C), but the above prefrontal cortical pattern of alteration was observed, whereby axonal alignment (Figure 5J, genotype: $F_{(1, 10)}$ = 7.70, p = 0.02) and axonal density (Figure 5K: genotype: ($F_{(1, 10)}$ = 6.10, p = 0.033), were increased in *Nrxn2α* KO

Pervolaraki et al.

mice in the basomedial nuclei, while cell density was unaffected (Figure 5L: genotype: ($F_{(1, 10)}$ <1). Alterations in axonal alignment and density as directly revealed by CLARITY could explain the increases in diffusivity and RD in the prefrontal regions, as measured by DTI.

To confirm the specificity of these alterations, we examined three further brain regions; the primary motor cortex (M1; Supp. Figure 5D-F), the primary somatosensory cortex (S1; Supp. Figure 5H-J) and the barrel field (BF; Supp. Figure 5K-M). Although there were differences between the hemispheres, there were no statistical differences between the genotypes or genotype x hemisphere interactions for any measure (Supp. Table 3).

In summary, in both the prefrontal ROIs, namely the OFC, and the ACC, DTI showed that ADC and RD were increased in *Nrxn2α* KO mice, likely related to complementary analysis from CLARITY showing that axonal alignment was altered in *Nrxn2α* KO mice in both prefrontal ROIs.

Pervolaraki et al.

Discussion

The *Nrxn2α* deletion captures many of the key aspects of human ASD. In terms of behaviour, three studies have now found social deficits associated with *Nrxn2α* KO, and in terms of brain structure, as summarised below, the *Nrxn2α* KO mouse model shows similar altered connectivity patterns in 'social brain' regions.

A DTI approach has been used for some time to explore neuropathological markers in autistic patients; alterations in myelination, axonal abundance, size and orientation all modify FA and ADC values[12, 27], specifically by reducing amygdala FA[14, 27], and have been used as a quantitative measure of changes to brain white matter integrity[14, 15]. Increased cortical ADC typically indexes reduced functionality: e.g. increased hippocampal ADC is associated with mild cognitive impairment[28, 29] and predicts verbal and visuospatial memory in old healthy controls[30]. Furthermore, both increased RD of various white matter tracts[31, 32] and increased whole-brain AD[32] have been observed in ASD. The *Nrxn2α* KO mouse encapsulates many of these specific changes, including reduced FA and increases in ADC, AD and RD. Whole brain ADC, AD and RD but not FA were increased in ASD children, as was ADC and RD in frontal cortex tracts[32]. This is in agreement with other studies noting frontal ADC increases but not for FA[27]. Likewise, FA was reduced in the amygdala in ASD children and adolescents[33], and remarkably, right-sided lateralisation of amygdala/hippocampal connections has been noted in high-functioning adolescents/adults with autism[34].

Our findings corroborate these quantifications of clinical autism, but highlights the question of what do the different measures of ADC, FA, AD and RD represent? Importantly, we observed these microstructural changes in various social brain regions against a background of unchanged cell density in all our study's ROIs. Unexpectedly, this highlights the power of our new approach. Dudanova et al. (2007) attempted to define the contribution of NRXN2 to the development of normal brain structure by cell counting and measuring cortical cell layer thickness and concluded that NRXN2 played little role in normal brain development [11]. Indeed, in earlier studies, it was suggested that deletion

of all Nrxns was unlikely to affect synaptic development but instead disrupts synaptic function[35]. We propose that coarse measures such as two-dimensional cell counting may be underestimating the impact of genetic mutations upon normal development. By staining cleared brain tissue with a nuclear marker and performing automated three-dimensional cell counting, we found no effect of *Nrxn2α* deletion on cell density in any region of interest examined. But this belies the clear effects upon microstructure integrity across multiple regions as measured by both DTI and CLARITY, and its specificity; only social brain regions and not primary sensory or motor regions were disrupted. Future studies will benefit from employing more sensitive measures of brain structure and connectivity to determine the relevance of genetic mutations in development.

FA and ADC can be influenced by changes in axonal density and alignment (e.g. by myelination, demyelination, axonal damage, loss of white matter coherence[36]). It is likely that the axonal alignment metric used to quantify CLARITY more closely reflects the ADC measure of DTI, given that ADC (or mean diffusivity) equally weights diffusion across all eigenvectors and does not bias the primary eigenvector as FA does. Thus, it is likely that alterations in the properties of axons in Nrxn2α KO mice are driving these changes in FA and ADC. Given we mostly see differences in RD, thought to reflect tract branching and myelination (as it measures λ_2 and λ_3), it is possible that the orientation in the perpendicular not parallel orientation of fibres is mostly affected. Given the differences in the amygdala, OFC and ACC, it is possible that even though neuronal densities are similar in the Nrxn2α KO brain, it is the connections between neurones and brain regions that are perturbed. This would be consistent with the idea that connectivity disruption may represent a core feature of autism[37].

Here we have developed a new application of CLARITY to quantitatively investigate disease models by combining DTI with high resolution 3D imaging and automated analysis of axonal fibres in a within subject study. Inevitably, there are some technical limitations that will require future refinement as this technology matures.

First, while we used CLARITY and immunolabelling to identify axons, we cannot know whether axon-related changes alone reflects all the changes we observed for our DTI measures. Second, whilst we can segment entire brain regions for DTI analysis, it was not practical to image larger brain areas at the necessary resolution. While it is theoretically possible that we may bias sampling of each brain region by picking ROIs for multiphoton imaging, this was done using atlas-defined coordinates and by an experimenter blind to the DTI results, so minimising any bias. However, within the current study, we were only able to apply the CLARITY approach to the amygdala, OFC and ACC. It was not practical to apply this methodology to the hippocampus, due to its extremely heterogeneous structure. The small cubic ROIs could not be reproducibly positioned, and larger ROIs to average across larger areas of the hippocampus were not possible. Although imaging of fibre tracts in large volumes of cleared tissue is possible[38], fluorescent labelling limitations make it impractical for a study of this nature. Despite this, as the adoption of the CLARITY technique increases, we hope that the use of DTI and CLARITY to study connectivity across spatial scales will become commonplace.

As yet, no one DTI protocol has emerged as the standard for in vivo or ex vivo imaging. Indeed, there has been debate regarding the best number of diffusion gradients to use, among other parameters[39]. Whilst some claim a minimum of 30 directions are required to estimate anisotropy[39], others claim that the benefits of using more than 6 are limited[40-43]. A limitation of these studies is that they have not followed up using anatomical staining. Our study provides further evidence against the notion that more directions are necessarily better. Where we found DTI differences, these were corroborated by the quantification of CLARITY.

In summary, our combined use of DTI-plus-CLARITY has revealed changes in microstructure and connectivity of the social brain that was previously inconceivable using classical experimental approaches. We envisage this approach will deliver a new level of detail in the connectivity hypothesis of autism.

Pervolaraki et al.

Acknowledgements

This work was supported by the Guy's and St. Thomas' Charity Prize PhD scholarship to AT, a Medical Research Council (UK) grant (G0900625) to SJC and RJR, a University of Leeds Wellcome Trust ISSF (UK) Fellowship, a Royal Society (UK) grant (RG130316), an Alzheimer's Society Fellowship (AS-JF-15-008) to JD, a British Pharmacological Society (UK) grant to JD and CL, a BBSRC grant to LCA (BB/P000479/1) and a BBSRC grant to CL (BB/M008975/1). We acknowledge financial support from the Innovative Medicines Initiative Joint Undertaking under grant agreement no. 115300, resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007–2013) and EFPIA companies' in kind contribution, the Mortimer D Sackler Foundation and the Sackler Institute for Translational Neurodevelopment (ALT and LCA). Some analysis scripts were provided to AT at the Computational Image Analysis in Cellular and Developmental Biology course at the Marine Biological Laboratory (Woods Hole, MA, USA), funded by National Institutes of Health (R25 GM103792-01).

References

1.   Cardno AG, Gottesman, II. Twin studies of schizophrenia: from bow-and-arrow concordances to star wars Mx and functional genomics. *American journal of medical genetics* 2000; **97**(1)**:** 12-17.

2.   Ronald A, Happe F, Bolton P, Butcher LM, Price TS, Wheelwright S*, et al*. Genetic heterogeneity between the three components of the autism spectrum: a twin study. *Journal of the American Academy of Child and Adolescent Psychiatry* 2006; **45**(6)**:** 691-699.

3.   Sahin M, Sur M. Genes, circuits, and precision therapies for autism and related neurodevelopmental disorders. *Science* 2015; **350**(6263).

4.   Gauthier J, Siddiqui TJ, Huashan P, Yokomaku D, Hamdan FF, Champagne N*, et al*. Truncating mutations in NRXN2 and NRXN1 in autism spectrum disorders and schizophrenia. *Human genetics* 2011; **130**(4)**:** 563-573.

5.   Mohrmann I, Gillessen-Kaesbach G, Siebert R, Caliebe A, Hellenbroich Y. A de novo 0.57 Mb microdeletion in chromosome 11q13.1 in a patient with speech problems, autistic traits, dysmorphic features and multiple endocrine neoplasia type 1. *European journal of medical genetics* 2011; **54**(4)**:** e461-464.

6.   Boyle MI, Jespersgaard C, Nazaryan L, Ravn K, Brondum-Nielsen K, Bisgaard AM*, et al*. Deletion of 11q12.3-11q13.1 in a patient with intellectual disability and childhood facial features resembling Cornelia de Lange syndrome. *Gene* 2015; **572**(1)**:** 130-134.

7.   Li J, Wang L, Gou H, Shi L, Zhang K, Tang M*, et al*. Targeted sequencing and functional analysis reveal brain-size-related genes and their networks in autism spectrum disorders. *Molecular Psychiatry* 2017; **22:** 1282–1290.

8.   Dachtler J, Glasper J, Cohen RN, Ivorra JL, Swiffen DJ, Jackson AJ*, et al*. Deletion of alpha-neurexin II results in autism-related behaviors in mice. *Translational psychiatry* 2014; **4:** e484.

9.   Born G, Grayton HM, Langhorst H, Dudanova I, Rohlmann A, Woodward BW*, et al*. Genetic targeting of NRXN2 in mice unveils role in excitatory cortical synapse function and social behaviors. *Frontiers in synaptic neuroscience* 2015; **7:** 3.

10.   Dachtler J, Ivorra JL, Rowland TE, Lever C, Rodgers RJ, Clapcote SJ. Heterozygous deletion of alpha-neurexin I or alpha-neurexin II results in behaviors relevant to autism and schizophrenia. *Behavioral neuroscience* 2015; **129**(6)**:** 765-776.

11.   Dudanova I, Tabuchi K, Rohlmann A, Sudhof TC, Missler M. Deletion of alpha-neurexins does not cause a major impairment of axonal pathfinding or synapse formation. *The Journal of comparative neurology* 2007; **502**(2)**:** 261-274.

12. Beaulieu C. The basis of anisotropic water diffusion in the nervous system - a technical review. *NMR in biomedicine* 2002; **15**(7-8)**:** 435-455.

13. Barnea-Goraly N, Kwon H, Menon V, Eliez S, Lotspeich L, Reiss AL. White matter structure in autism: preliminary evidence from diffusion tensor imaging. *Biological psychiatry* 2004; **55**(3)**:** 323-326.

14. Noriuchi M, Kikuchi Y, Yoshiura T, Kira R, Shigeto H, Hara T*, et al*. Altered white matter fractional anisotropy and social impairment in children with autism spectrum disorder. *Brain research* 2010; **1362:** 141-149.

15. Belmonte MK, Allen G, Beckel-Mitchener A, Boulanger LM, Carper RA, Webb SJ. Autism and abnormal development of brain connectivity. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2004; **24**(42)**:** 9228-9231.

16. Chung K, Wallace J, Kim SY, Kalyanasundaram S, Andalman AS, Davidson TJ*, et al*. Structural and molecular interrogation of intact biological systems. *Nature* 2013; **497**(7449)**:** 332-337.

17. Pervolaraki E, Anderson RA, Benson AP, Hayes-Gill B, Holden AV, Moore BJ*, et al*. Antenatal architecture and activity of the human heart. *Interface focus* 2013; **3**(2)**:** 20120065.

18. Paxinos G, Franklin K. *The Mouse Brain in Stereotaxic Coordinates*, Third edn. Academic Press, 2008.

19. Tyson AL, Hilton ST, Andreae LC. Rapid, simple and inexpensive production of custom 3D printed equipment for large-volume fluorescence microscopy. *International journal of pharmaceutics* 2015; **494**(2)**:** 651-656.

20. Alexander AL, Lee JE, Lazar M, Field AS. Diffusion tensor imaging of the brain. *Neurotherapeutics : the journal of the American Society for Experimental NeuroTherapeutics* 2007; **4**(3)**:** 316-329.

21. Okuyama T, Kitamura T, Roy DS, Itohara S, Tonegawa S. Ventral CA1 neurons store social memory. *Science* 2016; **353**(6307)**:** 1536-1541.

22. Pitkanen A, Pikkarainen M, Nurminen N, Ylinen A. Reciprocal connections between the amygdala and the hippocampal formation, perirhinal cortex, and postrhinal cortex in rat. A review. *Annals of the New York Academy of Sciences* 2000; **911:** 369-391.

23. Cavada C, Company T, Tejedor J, Cruz-Rizzolo RJ, Reinoso-Suarez F. The anatomical connections of the macaque monkey orbitofrontal cortex. A review. *Cerebral cortex* 2000; **10**(3)**:** 220-242.

24. Abrams DA, Lynch CJ, Cheng KM, Phillips J, Supekar K, Ryali S*, et al*. Underconnectivity between voice-selective cortex and reward circuitry in

children with autism. *Proceedings of the National Academy of Sciences of the United States of America* 2013; **110**(29)**:** 12060-12065.

25.   Murphy ER, Foss-Feig J, Kenworthy L, Gaillard WD, Vaidya CJ. Atypical Functional Connectivity of the Amygdala in Childhood Autism Spectrum Disorders during Spontaneous Attention to Eye-Gaze. *Autism research and treatment* 2012; **2012:** 652408.

26.   Rutishauser U, Tudusciuc O, Wang S, Mamelak AN, Ross IB, Adolphs R. Single-neuron correlates of atypical face processing in autism. *Neuron* 2013; **80**(4)**:** 887-899.

27.   Sundaram SK, Kumar A, Makki MI, Behen ME, Chugani HT, Chugani DC. Diffusion tensor imaging of frontal lobe in autism spectrum disorder. *Cerebral cortex* 2008; **18**(11)**:** 2659-2665.

28.   Fellgiebel A, Wille P, Muller MJ, Winterer G, Scheurich A, Vucurevic G*, et al*. Ultrastructural hippocampal and white matter alterations in mild cognitive impairment: a diffusion tensor imaging study. *Dementia and geriatric cognitive disorders* 2004; **18**(1)**:** 101-108.

29.   Stebbins GT, Murphy CM. Diffusion tensor imaging in Alzheimer's disease and mild cognitive impairment. *Behavioural neurology* 2009; **21**(1)**:** 39-49.

30.   Carlesimo GA, Cherubini A, Caltagirone C, Spalletta G. Hippocampal mean diffusivity and memory in healthy elderly individuals: a cross-sectional study. *Neurology* 2010; **74**(3)**:** 194-200.

31.   Ameis SH, Fan J, Rockel C, Voineskos AN, Lobaugh NJ, Soorya L*, et al*. Impaired structural connectivity of socio-emotional circuits in autism spectrum disorders: a diffusion tensor imaging study. *PloS one* 2011; **6**(11)**:** e28044.

32.   Shukla DK, Keehn B, Muller RA. Tract-specific analyses of diffusion tensor imaging show widespread white matter compromise in autism spectrum disorder. *Journal of child psychology and psychiatry, and allied disciplines* 2011; **52**(3)**:** 286-295.

33.   Jou RJ, Jackowski AP, Papademetris X, Rajeevan N, Staib LH, Volkmar FR. Diffusion tensor imaging in autism spectrum disorders: preliminary evidence of abnormal neural connectivity. *The Australian and New Zealand journal of psychiatry* 2011; **45**(2)**:** 153-162.

34.   Conturo TE, Williams DL, Smith CD, Gultepe E, Akbudak E, Minshew NJ. Neuronal fiber pathway abnormalities in autism: an initial MRI diffusion tensor tracking study of hippocampo-fusiform and amygdalo-fusiform pathways. *Journal of the International Neuropsychological Society : JINS* 2008; **14**(6)**:** 933-946.

35.   Missler M, Zhang W, Rohlmann A, Kattenstroth G, Hammer RE, Gottmann K, *et al*. Alpha-neurexins couple Ca2+ channels to synaptic vesicle exocytosis. *Nature* 2003; **423**(6943)**:** 939-948.

36.   Alba-Ferrara LM, de Erausquin GA. What does anisotropy measure? Insights from increased and decreased anisotropy in selective fiber tracts in schizophrenia. *Frontiers in integrative neuroscience* 2013; **7:** 9.

37.   Girgis RR, Minshew NJ, Melhem NM, Nutche JJ, Keshavan MS, Hardan AY. Volumetric alterations of the orbitofrontal cortex in autism. *Progress in neuro-psychopharmacology & biological psychiatry* 2007; **31**(1)**:** 41-45.

38.   Ye L, Allen WE, Thompson KR, Tian Q, Hsueh B, Ramakrishnan C, *et al*. Wiring and Molecular Features of Prefrontal Ensembles Representing Distinct Experiences. *Cell* 2016; **165**(7)**:** 1776-1788.

39.   Jones DK. The effect of gradient sampling schemes on measures derived from diffusion tensor MRI: a Monte Carlo study. *Magnetic resonance in medicine* 2004; **51**(4)**:** 807-815.

40.   Lebel C, Benner T, Beaulieu C. Six is enough? Comparison of diffusion parameters measured using six or more diffusion-encoding gradient directions with deterministic tractography. *Magnetic resonance in medicine* 2012; **68**(2)**:** 474-483.

41.   Hasan KM, Parker DL, Alexander AL. Comparison of gradient encoding schemes for diffusion-tensor MRI. *Journal of magnetic resonance imaging : JMRI* 2001; **13**(5)**:** 769-780.

42.   Yao X, Yu T, Liang B, Xia T, Huang Q, Zhuang S. Effect of increasing diffusion gradient direction number on diffusion tensor imaging fiber tracking in the human brain. *Korean journal of radiology* 2015; **16**(2)**:** 410-418.

43.   Ni H, Kavcic V, Zhu T, Ekholm S, Zhong J. Effects of number of diffusion gradient directions on derived diffusion tensor imaging indices in human brain. *AJNR American journal of neuroradiology* 2006; **27**(8)**:** 1776-1781.

Figure Legends

Figure 1

Quantification of CLARITY imaging. **A** Sections of DTI-scanned brain were segmented at different Bregma levels for (i) the orbitofrontal cortex, (ii) the anterior hippocampus and amygdala, (iii) the mid hippocampus and posterior amygdala and (iv) the posterior hippocampus. **B-D** DTI-scanned brains were computed for tracts. Tissue from wild-type and *Nrxn2α* KO mice were cleared and stained for neurofilament and DAPI (**E**). **F** Automated MATLAB scripts were used to segment the DAPI (blue) and neurofilament (purple) channels such that cell density and axonal density and orientation could be calculated. **G** is representative of a CLARITY-derived 3D stacked image of a DAPI and neurofilament of a region of interest, with **H** being the corresponding segmented image. Scale bar: 100 μm.

Figure 2

Deletion of *Nrxn2α* reduces amygdala fractional anisotropy (FA) but not apparent diffusion coefficient (ADC). DTI images of the amygdala was segmented at two regions; anterior (Bregma -1.94 mm) and posterior (Bregma -2.46 mm). FA of the whole amygdala structure was significantly reduced at both anterior (**A**) and posterior (**B**) regions. FA was also significantly reduced in the anterior basolateral amygdala (BLA) (**C**). However, ADC was similar between the genotypes in the anterior (**D**), posterior (**E**) and BLA (**F**). **=$P<0.01$, *=$P<0.05$. Error bars represent s.e.m. Wild-type n=6, *Nrxn2α* KO n=6.

Figure 3

Nrxn2α KO mice have increased apparent diffusion coefficient (ADC) and axial (AD) and radial diffusivity (RD) in the orbitofrontal cortex (OFC) and the anterior cingulate cortex (ACC). Although reduced, there was no significant difference between wild-types and *Nrxn2α* KO mice for FA in the OFC (**A**) and ACC (**C**), but ADC was significantly increased in *Nrxn2α* KO mice in both prefrontal regions (**B** and **D**). The OFC has significantly increased AD and RD (**E-F**), whereas only RD was increased in the ACC (**G-H**). ***=$P=0.0005$,

Pervolaraki et al.

***=P<0.001, **=P<0.01, *=P<0.05. Error bars represent s.e.m. Wild-type n=6, *Nrxn2α* KO n=6.

Figure 4

Tractographic analysis of amygdala-hippocampus and amygdala-orbitofrontal cortex (OFC) connectivity. Amygdala-hippocampal connections are characterised by greater right hemisphere axial diffusivity (AD) *Nrxn2α* KO mice (**A**) but not radial diffusivity (RD) (**B**). Specific to the BLA, connections to the anterior hippocampus (**C**) and posterior hippocampus (**D**) have greater right hemisphere AD. Although the amygdala-OFC connection was similar between the genotypes for AD (**E**), *Nrxn2α* KO mice had significantly increased RD (**F**). *=P<0.05, ***=P<0.001. Error bars represent s.e.m. Wild-type n=6, Nrxn2α KO n=6.

Figure 5

CLARITY reveals differences in axonal alignment and fibre density in *Nrxn2α* KO mice. (**A-C**) Representative images of the CLARITY-treated brain, with ROI defined for the anterior cingulate cortex (ACC), orbitofrontal cortex (OFC), basomedial amygdala (BMA) and basolateral amygdala (BLA). For the ACC, the axonal alignment (**D**) and axon density (**E**) were significantly altered in KO mice, but cell density was unaltered (**F**). Within the medial OFC, only axonal alignment was significantly altered in KOs (**G**), with axon density (**H**) and cell density (**I**) being similar. For the BMA, both the axonal alignment (**J**) and axon density (**K**) were significantly increased, whilst cell density was unaltered (**L**). *=P<0.05, **=P<0.01. Error bars represent s.e.m. Wild-type n=6, *Nrxn2α* KO n=6.

Figure 1

i   ii   iii   iv

Bregma +2.58 mm   Bregma -1.94 mm   Bregma -2.46 mm   Bregma -3.28 mm

Figure 2

Figure 3

Figure 4

Figure 5