Representation of natural image contents by sparsely active neurons in visual cortex

Takashi Yoshida[1,2,3,*] and Kenichi Ohki[1,2,3,4,*]

1. Department of Physiology, The University of Tokyo School of Medicine, Tokyo, Japan.

2. Department of Molecular Physiology, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

3. CREST, Japan Science and Technology Agency, Tokyo, Japan

4. International Research Center for Neurointelligence (WPI-IRCN), The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

*Correspondence: Kenichi Ohki, kohki@m.u-tokyo.ac.jp

Takashi Yoshida, takashiy@m.u-tokyo.ac.jp

Abstract

Natural scenes sparsely activate neurons in the primary visual cortex (V1). However, whether sparsely active

neurons sufficiently represent natural image contents and what proportion of neurons are involved in the

natural image processing have not been revealed. We recorded the visual responses of mouse V1 neurons to

natural images. Small numbers of neurons highly responded to single natural images, whereas most neurons

were activated by at least one image. We reconstructed the natural images from the neuronal activities. Visual

contents of single natural images were linearly decodable from surprisingly small numbers (~20) of highly

responsive neurons. This was achieved by diverse visual features encoded by the small number of responsive

neurons. This representation allows the downstream areas to easily access information contents encoded in

sparsely active V1 neurons.

Introduction

Sensory information is thought to be represented by relatively small numbers of active neurons in the sensory cortex. This sparse representation has been observed in several cortical areas[1-9] and is postulated to represent an efficient coding of the statistical features in sensory inputs[4, 10]. However, it has not been determined whether and how small numbers of active neurons represent sufficient information about sensory inputs.

In the primary visual cortex (V1), a type of neuron termed a simple cell has a receptive field (RF) structure that is spatially localized, oriented, and has a bandpass filter property of specific spatial frequency. This RF structure is modelled by a two-dimensional (2D) Gabor function[11]. According to theoretical studies, single natural images are represented by relatively small numbers of neurons using Gabor-like RFs, whereas information about multiple natural scenes is distributed across the neuronal population[10,12,13]. Indeed, V1 neurons respond sparsely to natural scenes at the single cell level[2, 3, 5-9] and population level[3,5,14]. Population activity with higher sparseness exhibits greater discriminability between natural scenes[5].

What types of information from natural scenes are represented in sparsely active neuronal populations? The visual contents of natural scenes or movies are reconstructed from populations of single unit activities in the lateral geniculate nucleus (LGN) collected from several experiments[15] and functional magnetic resonance imaging (fMRI) data from the visual cortices[16-19]. However, it has not been addressed whether the visual contents of natural images are represented by small numbers of sparsely active neurons in populations of single neurons.

Furthermore, how the sparsely active neurons are distributed among neurons in a local population has not been determined. It has been reported that subset of neurons are 'unresponsive' to visual stimuli (e.g., a responsive rate for visual stimuli in mouse V1 of 26–68%)[14, 20-25]. This may partly because stimulus properties do not completely cover receptive field properties of all neurons. Thus, there are two extreme possibilities; sparsely active neurons distributed among all neurons in a local population, or only a specific subset of cells process the natural images. What proportion of neurons are actually involved in information processing has been debated[26, 27].

Here, we examined whether a small number of highly responsive V1 neurons was sufficient for the

representation of natural image contents and what proportion of V1 neurons were involved in the natural

image processing. Using two-photon Ca^{2+} imaging, we recorded visual responses to natural images from local

populations of single neurons in V1 of anaesthetized mice. A small number of neurons (<3%) highly

responded to each natural image, which was sparser than that predicted by linear encoding model. On the other

hand, approximately 90% of neurons were activated by at least one of the natural images, revealing that most

neurons in a local population are involved in natural image processing, consistent with the sparse, distributed

representation[10]. We reconstructed the natural images from the neuronal activities to estimate the information

about the visual contents represented by local populations. Visual contents of single natural images were

linearly decodable from small numbers (~20) of highly responsive neurons. The individual responsive neurons

encoded unique visual information, which helped small numbers of neurons represent complex natural images.

Furthermore, the responsive neurons were only slightly overlapped between images, and many natural images

were represented by the combinations of responsive neurons in a population. Finally, the features represented

by a local population of all responsive neurons were sufficient to represent features in all the natural images

we used. Based on these results, most neurons in a local population are involved in natural image processing,

and visual contents of single natural images are linearly decodable from surprisingly small number of neurons.

This representation allows the downstream areas to easily access information contents in the small number of

V1 neurons.

Preliminary results of this study have been published in an abstract form[28]

Results

The main purpose of this study is to examine whether and how the visual contents of natural images are

represented in the sparse representation scheme. We first confirmed the sparse response to natural images in

our dataset. Next, we examined the visual features that were encoded in the responses of individual neurons.

Finally, using these features, the visual contents of a natural image were reconstructed from a relatively small

number of responsive neurons.

Sparse visual responses to natural images in mouse V1

We presented flashes of natural images as visual stimuli (Fig. 1a, see Methods) and simultaneously recorded

the activities of several hundreds of single neurons from layers 2/3 and 4 of mouse V1 using two-photon

calcium (Ca^{2+}) imaging (560 [284–712] cells/plane, median [25–75th percentiles], n = 24 planes from 14 mice,

260–450 microns in depth, Fig. 1b for representative response traces). Fig. 1c presents plots of significant

visual response events for all images (x-axis) across all neurons (y-axis) in a plane (n = 726 cells, depth: 360

microns from the brain surface). Significant response for each image was defined as an evoked response which

was significantly different from 0 (p < 0.01 using the signed-rank test) and whose normalized response

amplitude (z-score, see Methods) was greater than 1. Hereafter, we call these significant visual responses

highly responsive. A few to 10% of neurons were highly responsive to a single image (5.1% [3.9–6.7%]

cells/image, Fig. 1c bottom panel), indicating sparse visual responses to natural images. In contrast, nearly all

neurons (98%, 711/726 cells) responded to at least one image (each cell responded 4.5% [2.5–7.5%] images,

Fig. 1c right panel). Across planes, 2.7% cells were activated by a single image ([1.8–3.2%], Fig. 1f), whereas

almost all cells responded to at least one image (90% [86–93%], Fig. 1g). This low responsive rate to each

image was not due to poor recording conditions. The same neurons responded well to moving gratings (27%

[22–34%] for one direction, and 75% [66–79%] for at least one of 8 directions, Fig. 1h and i).

The highly responsive neurons only slightly overlapped between images. Fig. 1d presents representative

activity patterns for three natural images (Fig. 1d, left column). Each image activated different subsets of

neurons that exhibited small overlaps between images (Fig. 1d, right column). Of the responsive cells, 4.8%

exhibited overlap between two images (25–75th percentiles for 24 planes: 4.0–5.5%, Fig. 1j). We further

computed the distributions of the response amplitudes to single images (Fig. 1e). Small numbers of neurons

exhibited visual responses with greater amplitudes, which is a characteristic property of a sparse representation

(Fig. 1e). Population sparseness[2, 3], a measure of sparse representation, was comparable to a previous report for

mouse V1[5] (0.36 [0.30–0.42], Fig. 1k, see Methods). Thus, each natural image activated a relatively small

number of neurons, whereas most neurons in a local population were activated by at least one of the images,

suggesting the sparsely distributed representation of natural image in V1 that was originally proposed in a

previous study[10]. The latter result also provides the first report that most neurons in mouse V1 are visually

responsive to natural image stimuli[26, 27].

Small overlapping representations of visual features among local V1 populations

We created encoding models for the visual response of individual neurons to examine the visual features

represented by each neuron. We used a set of Gabor wavelet filters (1248 filters, Supplementary Fig. 1a and b,

see Methods) to extract the visual features from the natural images. Natural images were applied to each

Gabor filter and transformed into sets of feature values (Gabor feature values). For each neuron, we first

selected the Gabor features that exhibited strong correlations with the visual response. The correlation

threshold for the selected feature was adjusted to maximize the visual response prediction (Supplementary Fig.

1c–e, see Methods). Then, the visual response was represented by a linear regression of the selected feature

values followed by a non-linear scaling (Fig. 2a, see Methods). The visual response prediction of the model

was estimated with a different dataset from the dataset used in the regression (see Methods).

Visual responses of individual neurons were represented by small numbers of Gabor features. In the

example cells (Fig. 2b and c), the correlation coefficients between the observed responses and the responses

predicted by the model were 0.76 and 0.89. These neurons were represented by 19 and 13 Gabor features,

respectively (Fig. 2b and c, right panels), and their encoding filters (weighted sums of the Gabor filters) were

spatially localized (Fig. 2b and c, insets in the right panels). In the representative plane presented in Fig. 1, the

median of the prediction performance of the encoding model (i.e., the correlation coefficient between the

observed and predicted responses) was 0.34 (25–75[th] percentiles: 0.16–0.52, n = 726 cells, Supplementary Fig.

1f), and the median performance of all cells across planes was 0.24 (25–75[th] percentiles: 0.07–0.45, n = 12755

cells across 24 planes, Supplementary Fig. 1i). An examination of the non-linear scaling function revealed that

this step suppressed weak predicted responses and enhanced strong predicted responses (see Fig. 2d and e for a

representative cell and average across planes, respectively), suggesting that this non-linear step enhanced the

sparseness of the predicted response obtained from the linear step (i.e., linear regression by feature values). On

average, 2.0% of the features (25/1248 features, 25–75th percentiles: 2.0–2.1%) were represented in each cell

of the example plane (upper panels in Fig. 2f and Supplementary Fig. 1g), and 2.1% were represented in each

cell of all recorded cells across all planes (~26/1248 features, 25–75th percentiles: 0.9–4.9%, n = 12755 cells,

Fig. 2h and Supplementary Fig. 1k). These features were related to the RF structure of each cell

(Supplementary Fig. 2). The RF structure of each cell was estimated using the regularized inverse method[29-31]

(see Methods). The regression weights of the Gabor features in the encoding model were positively correlated

with the similarity between the corresponding Gabor filter and the RF structure (Supplementary Fig.2a–d).

The Gabor features encoded in one cell slightly overlapped with those of other cells in a local population

(Fig. 2i). Among 19 and 13 Gabor features represented by the two example cells (Fig. 2b and c), only two

features overlapped. For all cell pairs across all planes, the median overlap was 3.4% (25–75th percentile:

0.0–9.6% relative to features represented by each cell, Fig. 2i and Supplementary Fig. 1h and 1l). The feature

overlap between neurons was positively correlated with the similarity of RF structure (Supplementary Fig.

2e–j). Based on these findings, the Gabor features encoded by neurons in a local population were highly

diverse.

The analysis of the encoding model also revealed how the individual Gabor features were encoded across

neurons (upper left and bottom panels in Fig. 2f and g). As the spatial frequency (SF) of the Gabor filter

increased (i.e. the scale decreased), the corresponding feature contributed to the visual responses of fewer

neurons (Fig. 2g). This pattern likely reflected the fact that Gabor filters with a low SF (i.e., a large scale)

covered more of the neuron's RF, whereas Gabor filters with a high SF (i.e. a small scale) affected the

responses of fewer neurons. Furthermore, almost all features contributed to the responses of at least one cell

(100% in the plane presented in Fig. 2f and 100% [99.4–100%] across all planes, median [25–75th percentiles],

Fig. 2j).

Image reconstruction from the activities of the neuronal population

The encoding model revealed the Gabor features represented by each neuron. We next examined whether the

features encoded in a local population of neurons were sufficient to represent the visual contents of the natural

images. We reconstructed stimulus images from the neuronal activities to evaluate information about visual

contents in the population activity[15-19]. Using the same Gabor features as in the encoding model, each Gabor

feature value was subjected to a linear regression of the neuronal activities of multiple neurons (Fig. 3a and

Supplementary Fig. 3a). Each Gabor feature value was independently reconstructed. Then, the sets of

reconstructed feature values were transformed into images (Fig. 3a, see Methods). The reconstruction

performance was estimated with a different dataset from the dataset used in the regression (see Methods).

We first used all simultaneously recorded neurons to reconstruct the image. In the examples of the plane

(n = 726 neurons, presented in Figs. 1 and 2), the rough structures of the stimulus images were reconstructed

from the population activities ("All-cells" in Fig. 3b). The reconstruction performances (pixel-to-pixel

correlations between stimulus images and reconstructed images) were 0.45 [0.36–0.56] (median [25–75th

percentiles] of 200 images) in the representative plane (n = 726 cells, Fig. 3c upper panel) and 0.36

[0.31–0.38] across all planes (n = 24 planes, "All cells" in Fig. 3d). Thus, the visual contents of natural images

were extracted linearly from the neuronal activities of local population in V1

The encoding model used in the previous section revealed how each neuron encodes the Gabor features

(Fig. 2f). We next examined whether these encoded features were sufficient for the representation of visual

contents. In this analysis, each Gabor feature value was reconstructed with a subset of neurons selected using

the encoding model (cell-selection model, Supplementary Fig. 3a, and see Methods). In this model, different

subsets of neurons were used to reconstruct different features (Fig. 2f). Across all features, almost all neurons

were used to reconstruct at least one feature (Fig. 2j). The examples of the reconstructed images from the

cell-selection model are presented in Fig. 3b (Cell-selection). The reconstruction performance of the

cell-selection model was comparable to or even slightly higher than the model using all cells (R = 0.49

[0.37–0.59] for the representative plane, Fig 3c lower panel, and 0.36 [0.32–0.39] for all planes, median

[25–75th percentiles], p = 4.0×10^{-4} using the signed-rank test, Fig. 3d). Thus, the Gabor features encoded in

individual cells in a population captured sufficient information about the visual contents of the natural image.

When the neurons were selected to maximize the reconstruction of each feature, the image reconstruction

performance was only slightly improved (Supplementary Fig. 3b–h). Thus, main information about the visual

contents was captured by the cell-selection model.

Visual contents of natural images are linearly decodable from small numbers of responsive neurons

Single natural images activated small numbers of neurons in a local population (Fig. 1). We next examined

whether these small numbers of highly responsive neurons contained sufficient information about the visual

contents of an image. In the cell-selection model (Fig. 3), cells were selected for each feature, and almost all

cells were used for at least one feature. Thus, almost all cells, including both the responsive and

non-responsive cells were used to reconstruct each image, and the same set of cells was used across all images.

In contrast, only a small number of neurons were highly responsive to each image (Fig. 1f). In the analysis

used in this section (Fig. 4), only highly responsive cells for each image were used in the reconstructions to

estimate the information contained in the highly responsive cells for each image. The parameters (weights and

biases) of the cell-selection model were used in the reconstruction model of this analysis.

Representative results are presented in Fig. 4a–c. In each image, cells were sorted by visual response

amplitude (descending order) first among the highly responsive neurons (red dots in Fig. 4a–c) and then

among the remaining cells (black dots in Fig. 4a–c). The image was reconstructed by top N cells (N = 1–726

cells), and the reconstruction performances were plotted against the numbers of cells used (Fig. 4a–d). Based

on the results of this analysis, all highly responsive cells or even fewer cells were sufficient to reconstruct the

image to a level that was fairly comparable to the reconstruction created with all cells (Fig. 4a–d). In summary,

the performance of the highly responsive cells was slightly better than the performance of all cells in the

representative plane (R = 0.52 [0.40–0.64] for the responsive cells and 0.49 [0.37–0.59] for all cells, Fig. 4f)

and across planes (R = 0.38 [0.34–0.44] for the responsive cells and 0.35 [0.31–0.40] for all cells, median

[25-75th percentiles], p = 3.2×10^{-4} using the signed-rank test, n = 24 planes, Fig. 4g). Furthermore, only

approximately 20 cells were sufficient to achieve 95% of the peak performance (vertical line in Fig. 4d). Thus,

the visual contents of the single natural images were linearly decodable from small numbers of highly

responsive neurons.

The features represented by individual neurons should not be highly redundant to represent diverse

features in a natural image using a small number of neurons. Fig. 4e illustrates how individual responsive

neurons contributed to the image reconstruction in the case presented in Fig 4a. Each neuron had a specific

pattern of contributions (reverse filter: sum of Gabor filters × weights, see Methods), and the contribution

patterns varied across neurons (Fig. 4e top panels). In neuron pairs that were highly responsive to the same

image, the number of overlapping Gabor features were slightly increased compared to all pairs, but the

percentage was still less than 10% (7.1% [1.0–16%] of features for the all pairs and 8.1% [6.3–10%] of

features for 24 planes, Fig. 4h–j, cf. Fig. 2g). These small overlaps and diversity in the represented features

between neurons should be useful for the representation of natural images by relatively small numbers of

highly responsive neurons.

Finally, we examined how multiple natural images were represented in a population of responsive

neurons (Fig. 5a–c). Figs. 5a and b provide an example of the representative plane shown in the previous

figures (n = 726 cells). Natural images were sorted by reconstruction performance (y-axis in Fig. 5a), and the

cells responding to each image are plotted in each row. First, as the number of images increased, new

responsive cells are added and the total number of responsive cells used for the reconstructions quickly

increased (right end of the plot on each row, Fig. 5a). At approximately 50 images, the number of newly added

responsive cells quickly decreased, and the increase in the total number of responsive cells slowed, indicating

that the newly added image was represented by a combination of the already plotted responsive cells (i.e., the

neurons that responded to other images), which was due to the small overlap in responsive cells between

images (Fig. 1j). These findings are summarized in Figures 5b and c in which the number of newly added cells

225 quickly decreased to zero as the number of images increased (red lines in Fig. 5b and c for the representative

226 case and for all planes, respectively). Therefore, although only 4.8% responsive neurons overlapped between

227 images (Fig. 1j), this small overlap is useful for the representation of many natural images by a limited number

228 of responsive neurons.

229  We also analyzed whether the features represented by the local populations of the responsive neurons

230 were sufficient to represent all features of the natural images. If the features in a local population are sufficient

231 to represent all natural images, all features of the natural images should be accurately represented by the

232 combination of features in the individual cells in a population. We represented a set of features in each image

233 by linear regression of weights (i.e., features) of all responsive cells from the reconstruction model

234 (cell-selection model) and computed the fitting errors (see Methods, Fig. 5d). The median error was less than

235 10% for all images and all planes (8.2% [4.5–15.2%] for all image cases and 5.7% [4.9–16%], n = 24 for the

236 planes, Fig. 5e and f). Based on this result, features that sufficiently represent the visual contents of natural

237 images are encoded in neurons in a local population.

239 **Discussion**

240 In the mouse V1, single natural images activated small numbers of neurons (2.7%) which was sparser than that

241 predicted by the linear model. The Gabor features represented in the individual cells only slightly overlapped

242 between neurons, indicating diverse response properties across individual neurons. Visual contents of natural

243 images were linearly decodable from relatively small number of neurons (about 20 neurons) at a level

244 comparable to that from all recorded neurons. The neurons that responded to the same image contributed to

245 different visual features with low redundancy. The small overlaps of highly responsive neurons between the

246 images helped a limited number of the responsive neurons to represent multiple natural images. Finally, the

247 visual features represented by all the responsive neurons provided a good representation of the original visual

248 features in the natural images.

249  Visual responses to natural images or movies in V1 are sparse at the single cell level (high lifetime

sparseness)[2, 3, 5-9] and at the level of populations (population sparseness)[3, 5, 6]. Recently, recordings of local

population activity using two-photon Ca^{2+} imaging have enabled us to precisely evaluate the population

sparseness of local populations[5, 32]. We confirmed that single natural images activated only small numbers of

neurons. Encoding model analysis indicated that visual responses in individual neurons were sparser than that

predicted from a linear model (Fig. 2d, e). Here, this sparse activity was shown to contain sufficient

information to represent the visual contents of natural images. Image reconstruction is useful for evaluating the

information contents represented by the neuronal activity and is widely used to analyze populations of single

unit activities in LGN[15] and fMRI data in several visual cortical areas[16-19]. The former[15] study used

"pseudo-population" data collected from several experiments, and the latter studies[16-19] used blood oxygen

level-dependent (BOLD) signals that indirectly reflect the average local neuronal activity. Thus, it has not been

examined whether and how the visual contents of natural images are represented in simultaneously recorded

populations of single neurons in the cortex. We applied image reconstruction to two-photon Ca^{2+} imaging data

that completely covered activities of a local population at a single cell resolution, and revealed that visual

contents of single natural images were linearly decodable from relatively small numbers of responsive neurons

in a local population. It has been proposed that information is easy to be read out from the sparse

representation[4]. Indeed, the sparse population activity increases the discriminability of two natural scenes by

rendering the representations of the two scenes separable[5]. Our results extend this in that information about

visual contents encoded in sparsely active neurons is linearly accessible, suggesting that downstream areas are

easy to read out visual contents of natural image from the sparse representation in V1.

The visual features encoded by individual neurons should be diverse so that a small number of active

neurons represent the complex visual features of the image. Although RF structures in the local population of

the mouse V1 have already been reported[20, 21, 30, 31], their diversity has not been analyzed with respect to

natural image representation. The overlapping features were small in the present study, even among neurons

that were responsive to the same image. Furthermore, the visual features represented by sparsely active

neurons were sufficiently diverse to represent visual contents of natural images.

Our analysis also revealed how multiple natural images are represented in a local population of

responsive neurons. A single natural image activated specific subsets of neurons, whereas most neurons in a

local population responded to at least one of the images, supporting sparse, distributed code proposed in a

previous study[10]. The overlap of responsive neurons between images involved only 4.8% of the responsive

cells (Fig. 1i). However, due to this small overlap, many natural images were represented by a limited number

of responsive neurons (Fig. 5a–c). Furthermore, the features of all responsive neurons in a local population

were sufficient to represent all the natural images used in the present study (Fig. 5d–f). Based on these

findings, any natural image could be represented by a combination of responsive neurons in a local population.

In summary, this work highlighted how the visual contents of natural images are represented in sparsely

active neurons in local populations of V1 neurons. The visual contents of single natural images are linearly

decodable from relatively small numbers of responsive neurons in which the encoded features are sufficiently

diverse to represent the complex images. The responsive neurons are distributed among most of the neurons in

a local population, and natural images are represented by the combination of the sparsely responsive neurons

in which the responses exhibited slight overlap between images. This representation allows downstream areas

to easily read out information about visual contents encoded in the small number of V1 neurons.

Acknowledgements

We thank Ms. T. Inoue, Y. Sono, A. Ohmori, A. Honda, M. Nakamichi for animal care, and all members of

Ohki laboratory for support and discussions. This work was supported by grants from Core Research for

Evolutionary Science and Technology (CREST)—Japan Agency for Medical Research and Development

(AMED) (to KO), Japan Society for the Promotion of Science (JSPS) KAKENHI (Grant number 25221001

and 25117004 to KO and 15K16573, 17K13276 to T.Y.), International Research Center for Neurointelligence

(WPI-IRCN), JSPS (to K.O.), Brain Mapping by Integrated Neurotechnologies for Disease Studies

(Brain/MINDS)—AMED (to K.O.), Strategic International Research Cooperative Program (SICP)—AMED

(to K.O.), grants from the Ichiro Kanehara Foundation for the Promotion of Medical Sciences and Medical

Care, and the Uehara Memorial Foundation (to T.Y.).

Author contributions

T.Y. and K.O. designed the research. T.Y. performed experiments. T.Y. and K.O. analyzed data and wrote the

manuscript. K.O. supervised the project.

Competing financial interests.

We declare no competing financial interests.

References

1.   Rolls, E.T. & Tovee, M.J. Sparseness of the neuronal representation of stimuli in the primate
311   temporal visual cortex. *J Neurophysiol* **73**, 713-726 (1995).

312   2.   Vinje, W.E. & Gallant, J.L. Sparse Coding and Decorrelation in Primary Visual Cortex
313   During Natural Vision. *Science* **287**, 1273-1276 (2000).

314   3.   Weliky, M., Fiser, J., Hunt, R.H. & Wagner, D.N. Coding of natural scenes in primary visual
315   cortex. *Neuron* **37**, 703-718 (2003).

316   4.   Olshausen, B.A. & Field, D.J. Sparse coding of sensory inputs. *Curr Opin Neurobiol* **14**,
317   481-487 (2004).

318   5.   Froudarakis, E. *et al.* Population code in mouse V1 facilitates readout of natural scenes
319   through increased sparseness. *Nat Neurosci* **17**, 851-857 (2014).

320   6.   Yen, S.C., Baker, J. & Gray, C.M. Heterogeneity in the responses of adjacent neurons to
321   natural stimuli in cat striate cortex. *J Neurophysiol* **97**, 1326-1341 (2007).

322   7.   Yao, H., Shi, L., Han, F., Gao, H. & Dan, Y. Rapid learning in cortical coding of visual scenes.
323   *Nat Neurosci* **10**, 772-778 (2007).

324   8.   Tolhurst, D.J., Smyth, D. & Thompson, I.D. The sparseness of neuronal responses in ferret
325   primary visual cortex. *J Neurosci* **29**, 2355-2370 (2009).

326   9.   Willmore, B.D., Mazer, J.A. & Gallant, J.L. Sparse coding in striate and extrastriate visual
327   cortex. *J Neurophysiol* **105**, 2907-2919 (2011).

328   10.   Field, D.J. What Is the Goal of Sensory Coding. *Neural Comput* **6**, 559-601 (1994).

329   11.   Jones, J.P. & Palmer, L.A. An evaluation of the two-dimensional Gabor filter model of simple
330   receptive fields in cat striate cortex. *J Neurophysiol* **58**, 1233-1258 (1987).

331   12.   Olshausen, B.A. & Field, D.J. Emergence of simple-cell receptive field properties by learning
332   a sparse code for natural images. *Nature* **381**, 607-609 (1996).

333   13.   Bell, A.J. & Sejnowski, T.J. The "independent components" of natural scenes are edge filters.
334   *Vision Research* **37**, 3327-3338 (1997).

335   14.   Kampa, B.M., Roth, M.M., Gobel, W. & Helmchen, F. Representation of visual scenes by local
336   neuronal populations in layer 2/3 of mouse visual cortex. *Front Neural Circuits* **5**, 18 (2011).

337   15.   Stanley, G.B., Li, F.F. & Dan, Y. Reconstruction of natural scenes from ensemble responses in
338   the lateral geniculate nucleus. *J Neurosci* **19**, 8036-8042 (1999).

339   16.   Miyawaki, Y. *et al.* Visual Image Reconstruction from Human Brain Activity using a
340   Combination of Multiscale Local Image Decoders. *Neuron* **60**, 915-929 (2008).

341   17.   Naselaris, T., Prenger, R.J., Kay, K.N., Oliver, M. & Gallant, J.L. Bayesian reconstruction of
342   natural images from human brain activity. *Neuron* **63**, 902-915 (2009).

343   18.   Nishimoto, S. *et al.* Reconstructing visual experiences from brain activity evoked by natural
344   movies. *Curr Biol* **21**, 1641-1646 (2011).

345   19.   Horikawa, T., Tamaki, M., Miyawaki, Y. & Kamitani, Y. Neural decoding of visual imagery
346   during sleep. *Science* **340**, 639-642 (2013).

347   20.   Smith, S.L. & Hausser, M. Parallel processing of visual space by neighboring neurons in
348   mouse visual cortex. *Nat Neurosci* **13**, 1144-1149 (2010).

349   21.   Bonin, V., Histed, M.H., Yurgenson, S. & Reid, R.C. Local diversity and fine-scale
350   organization of receptive fields in mouse visual cortex. *J Neurosci* **31**, 18506-18521 (2011).

351   22.   Marshel, J.H., Garrett, M.E., Nauhaus, I. & Callaway, E.M. Functional specialization of
352   seven mouse visual cortical areas. *Neuron* **72**, 1040-1054 (2011).

353   23.   Ko, H. *et al.* Functional specificity of local synaptic connections in neocortical networks.
354   *Nature* **473**, 87-91 (2011).

355   24.   Rikhye, R.V. & Sur, M. Spatial Correlations in Natural Scenes Modulate Response
356   Reliability in Mouse Visual Cortex. *J Neurosci* **35**, 14661-14680 (2015).

357   25.   Miller, J.e.K., Ayzenshtat, I., Carrillo-Reid, L. & Yuste, R. Visual stimuli recruit intrinsically
358   generated cortical ensembles. *Proceedings of the National Academy of Sciences* **111**,
359   E4053-E4061 (2014).

360   26.   Olshausen, B.A. & Field, D.J. How Close Are We to Understanding V1? *Neural Comput* **17**,
361   1665-1699 (2005).

362   27.   Shoham, S., O'Connor, D.H. & Segev, R. How silent is the brain: is there a "dark matter"
363   problem in neuroscience? *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **192**,
364   777-784 (2006).

28.   Yoshida, T. & Ohki, K. Visual image reconstruction from neuronal activities in the mouse

primary visual cortex. *Program No. 415.17. 2015 Neuroscience Meeting Planner. Chicago,*
IL: Society for Neuroscience, 2015. Online. (2015).

29. Smyth, D., Willmore, B., Baker, G.E., Thompson, I.D. & Tolhurst, D.J. The Receptive-Field
Organization of Simple Cells in Primary Visual Cortex of Ferrets under Natural Scene
Stimulation. *The Journal of Neuroscience* **23**, 4746-4759 (2003).

30. Ko, H. *et al.* The emergence of functional microcircuits in visual cortex. *Nature* **496**, 96-100
(2013).

31. Cossell, L. *et al.* Functional organization of excitatory synaptic strength in primary visual
cortex. *Nature* **518**, 399-403 (2015).

32. Greenberg, D.S., Houweling, A.R. & Kerr, J.N. Population imaging of ongoing neuronal
activity in the visual cortex of awake rats. *Nat Neurosci* **11**, 749-751 (2008).

33. Ohki, K., Chung, S., Ch'ng, Y.H., Kara, P. & Reid, R.C. Functional imaging with cellular
resolution reveals precise micro-architecture in visual cortex. *Nature* **433**, 597-603 (2005).

34. Hagihara, K.M., Murakami, T., Yoshida, T., Tagawa, Y. & Ohki, K. Neuronal activity is not
required for the initial formation and maturation of visual selectivity. *Nat Neurosci* **18**,
1780-1788 (2015).

35. Mank, M. *et al.* A genetically encoded calcium indicator for chronic in vivo two-photon
imaging. *Nat Methods* **5**, 805-811 (2008).

36. van Hateren, J.H. & van der Schaaf, A. Independent component filters of natural images
compared with simple cells in primary visual cortex. *Proc Biol Sci* **265**, 359-366 (1998).

37. Olmos, A. & Kingdom, F.A. A biologically inspired algorithm for the recovery of shading and
reflectance images. *Perception* **33**, 1463-1473 (2004).

38. Peirce, J.W. Generating Stimuli for Neuroscience Using PsychoPy. *Front Neuroinform* **2**, 10
(2008).

39. Nimmerjahn, A., Kirchhoff, F., Kerr, J.N. & Helmchen, F. Sulforhodamine 101 as a specific
marker of astroglia in the neocortex in vivo. *Nat Methods* **1**, 31-37 (2004).

40. Kerlin, A.M., Andermann, M.L., Berezovskii, V.K. & Reid, R.C. Broadly tuned response
properties of diverse inhibitory neuron subtypes in mouse visual cortex. *Neuron* **67**, 858-871
(2010).

41. Treves, A. & Rolls, E.T. What determines the capacity of autoassociative memories in the
brain? *Network: Computation in Neural Systems* **2**, 371-397 (1991).

42. Lee, T.S. Image Representation Using 2D Gabor Wavelets. *IEEE Trans. Pattern Anal. Mach.*
Intell. **18**, 959-971 (1996).

43. Kay, K.N., Naselaris, T., Prenger, R.J. & Gallant, J.L. Identifying natural images from
human brain activity. *Nature* **452**, 352-355 (2008).

44.   Bishop, C.M. *Pattern Recognition and Machine Learning (Information Science and*
Statistics). (Springer-Verlag New York, Inc., 2006).

Methods

All experimental procedures were approved by the local Animal Use and Care Committee of Kyushu

University.

Animal preparation for two-photon imaging

C57BL/6 mice (male and female) were used (Japan SLC Inc., Shizuoka, Japan). Mice were anaesthetized with

isoflurane (5 % for induction, 1.5 % for maintenance during surgery, ~0.5% during imaging with a sedation of

< 0.5mg/kg chlorprothixene, Sigma-Aldrich, St. Louis, MO, USA). The head skin was removed from the head,

and the skull over the cortex was exposed. A custom made metal plate for head fixation was attached with

dental cement (Super Bond, Sun Medical, Shiga, Japan), and a craniotomy (~3mm in diameters) was

performed over the primary visual cortex (center position: 0–1 mm anterior to lambda, +2.5–3mm lateral to

the midline). A mixture of 0.8 mM Oregon Green BAPTA1-AM (OGB1, Life Technologies, Grand Island, NY,

USA) dissolved with 10% Pluronic (Life Technologies) and 0.025 mM sulforhodamine 101 (SR101,

Sigma-Aldrich) was pressure-injected with a Picospritzer III (Parker Hannifin, Cleveland, OH, USA) at the

depth of 300–500 µm from the brain surface. The cranial window was sealed with a coverslip and dental

cement. The imaging experiment began at least one hour after the OGB1 injection.

Two-photon Ca^{2+} imaging.

Imaging was performed with a two-photon microscope (A1R MP, Nikon, Tokyo, Japan) equipped with a 25×

objective (NA 1.10, PlanApo, Nikon) and Ti:sapphire mode-locked laser (MaiTai Deep See, Spectra Physics,

Santa clara, CA, USA)[33, 34]. OGB1 and SR101 were excited at a wave length of 920 nm. Emission filters of

525/50nm and 629/56nm were used for the OGB1 and SR101 signals, respectively. The fields of view (FOVs)

were 338 × 338 µm (10 planes from 7 mice) and 507 × 507 µm (14 planes from 7 mice) at 512 × 512 pixels.

The sampling frame rate was at 30Hz using resonant scanner.

Visual stimulation.

Before beginning the recording session, the retinotopic position of the recorded FOV was determined using

moving grating patches (lateral or upper directions, 99.9% contrast, 0.04 cycle/degrees, 2 Hz temporal

frequency, 20 and 50 degrees in diameter) while monitoring the changes in signals over the entire FOV. The

lateral or upper directions of the grating were used to activate many cells because the preferred directions of

mouse V1 neurons are slightly biased towards the cardinal directions[34, 35]. First, the grating patch of 50 degrees

in diameter was presented in one of 15 (5 × 3) positions that covered the entire monitor to roughly determine

the retinotopic position. Then, the patch of 20 degrees in diameter was presented on the 16 (4 × 4) positions

covering an 80 × 80-degree space to finely identify the retinotopic position. The stimulus position that induced

the maximum visual response of the entire FOV was set as the centre of the retinotopic position of the FOV.

A set of circular patches of grey-scaled, contrast-enhanced natural images (200 image types) was used as

the visual stimuli for response prediction and natural image reconstruction (60 degrees in diameter, 512 × 512

pixels, with a circular edge (5 degrees) that was gradually mixed to grey background). Each natural image was

adjusted to almost full contrast (99.9%). The mean intensity across pixels in each image was adjusted to an

approximately 50% intensity. Original natural images were obtained from the van Hateren's Natural Image

Dataset (http://pirsquared.org/research/#van-hateren-database)[36] and the McGill Calibrated Colour Image

Database (http://tabby.vision.mcgill.ca/html/welcome.html)[37]. During image presentation, one image type was

consecutively flashed three times (three 200-ms presentations interleaved with 200 ms of grey screen), and the

presentation of the next image was initiated after the presentation of grey screen for 200 ms. Images were

presented in a pseudo-random sequence in which each image was presented once every 200 image types. Each

image was presented at least 12 times (i.e., 12 trials) in a total recording session. We did not set a long interval

between image flashes to reduce the total recording time and increase the number of repetitions. In this design,

the tail of the Ca^{2+} response to one image invaded the time window of the next image presentation (Fig. 1b).

Although this overlap may have affected the visual responses between two adjacent images, many trial

repetitions (> 11 times for each image) in the pseudo-random order and the sparse responses to natural images

(Fig. 1) minimized the effects of response contamination between two consecutive images.

Moving, square gratings (8 directions, 0.04 cycle/degrees, 2 Hz temporal frequency, 60-degree patch

diameter) were presented at the same position as the natural image on the screen. Each direction was presented

for 4 sec interleaved by 4 sec of grey screen. The sequence of directions was pseudo-randomized, and each

direction was presented 10 times in a recording session.

All stimuli were presented with PsychoPy[38] on a 32-inch LCD monitor (Samsung, Hwaseong, South

Korea) with a 60-Hz refresh rate, and the timing of the stimulus presentation was synchronized with the timing

of image acquisition using a TTL pulse counter (USB-6501, National Instruments, Austin, TX, USA).

The entire recording session for one plane was divided into several recording sessions (4–6

trials/sub-session and 15–25 min for each session). Each recording session was interleaved by approximately

5–10 minutes of rest time during which the slight drift of the FOV was manually corrected. Every two or three

sessions, the retinotopic position of the FOV was checked with the grating patch stimuli during the resting

period. The recording was terminated, and then data were discarded if the retinotopic position was shifted

(probably due to eye movement). The recordings were performed in one to three planes of different depths

and/or positions in each animal (1.7 ± 0.8 planes, mean ± standard deviation).

Data analysis.

All data analysis procedures were performed using Matlab (Mathworks, Natick, MA, USA). Recorded images

were phase-corrected and aligned between frames. The averaged image across frames was used to determine

the region of interests (ROIs) of individual cells. After removing slow SF component (obtained with a

Gaussian filter with a sigma of approximately five times the soma diameter), the frame-averaged image was

subjected to a template matching method in which two dimensional difference of Gaussian (sigma1: 0.26 ×

soma diameter that was adjusted for zero-crossing at the soma radius, sigma2: soma diameter) was used as a

template for the cell body. Highly correlated areas between the frame-averaged image and the template were

detected as ROIs for individual cells. ROIs were manually corrected via visual inspection. SR101-positive

cells (putative astrocytes[39]) were removed from the ROI. Time course of calcium signal in each cell was

computed as an average of all pixels intensities within an ROI. Signal contamination from out of focus plane

was removed by a previously reported method[34, 40]. Briefly, a signal from ring-shaped area surrounding each

ROI was multiplied by a factor (contamination ratio) and subtracted from the signal of each cell. The

contamination ratio was determined to minimize the difference between the signal from blood vessel and the

surrounding ring shape region multiplied by the contamination ratio. The contamination ratios were computed

for several blood vessels in the FOV, and the mean value for several blood vessels was used for all cells in the

FOV.

The average response of 200-ms grey screen period just before each image was subtracted from the

average response of the time course during the last 200 ms of the stimulus period (during 3rd flash of each

image at approximately time of the peak Ca^{2+} transient) to compute visually evoked responses. The evoked

response was normalized for each cell by dividing by the standard deviation across all visual responses (200

images $\times$ trials, z-scored response). If the z-scored response to one image was significantly different from 0 (p

< 0.01 using signed-rank test across trials) and across-trial average of the z-scored response was greater than 1,

the response was considered significant for the image. The population sparseness (s) was computed using the

equation described in previous studies[2, 3, 41] as follows: s = $[1-(\sum R_i)^2/(N\sum R_i^2)]/(1-1/N)$, where R_i is the

evoked response of ith cell, and N is the number of cells (i = 1–N).

A square position (50 $\times$ 50 degrees) of the center of natural image patch was extracted and down-sampled

to 32 $\times$ 32-pixel image. The down-sampled image was used to analyze the Gabor features, response prediction

and image reconstruction.

Gabor features.

A set of spatially overlapping Gabor filter wavelets was prepared to extract the visual features of the natural

images[10, 42, 43]. The down-sampled images were first subjected to the set of Gabor filters to obtain Gabor

feature values. Single feature value corresponds to a single wavelet filters.

Gabor filters have four orientations (0, 45, 90, and 135 degrees), two phases, and 4 sizes (8×8, 16×16,

32×32, and 64×64pixels) located on 11×11, 5×5, 3×3, and 1×1 grids (Supplementary Fig. 1a and b).

Therefore, the three smaller scale filters were spatially overlapped with each other. The spatial frequencies of

the four scale sizes of the Gabor wavelets were 0.13, 0.067, 0.033, and 0.016 cycle/degrees (cpd). This filter

set was almost self-inverting, i.e., the feature values obtained by applying an image to the wavelet set, were

transformed back to the image by summing the filters after multiplying by the feature values[42]. The Gabor

filters and the transformations were based on an open source program (originally written by Drs. Daisuke Kato

and   Izumi   Ohzawa,   Osaka   University,   Japan,

https://visiome.neuroinf.jp/modules/xoonips/detail.php?item_id=6894).

Encoding model.

In the encoding model for response prediction, single-cell responses were predicted using a linear regression

analysis of selected Gabor feature values (Fig. 2a and Supplementary Fig. 1a–e). The encoding model was

created independently for each cell. First, Pearson's correlation coefficients between the response and each

feature value were computed. Then, using one of preset values for the correlation coefficient as a threshold (12

points ranging from 0.05 to 0.35, Supplementary Fig.1c–e), only the more strongly correlated features were

selected (feature selection) and used in the regression analysis. The weight and bias parameters of the

regression were estimated by Bayesian linear regression with an expectation-maximization algorithm which is

almost equivalent to linear regression with L2 regularization[44]. After the regression analysis, the non-linearity

of predicted response was adjusted via a rectification step using the following equation[31], predicted response =

$A/[1 + \exp(\alpha x + \beta)]$, where x is the output of the regression and A, α, and β are parameters to be estimated.

This step only scaled the regression output without changing the regression parameters (i.e., weights and

biases). The response prediction of the model was estimated by 10-fold cross validations (CVs) in which the

response data for 180 images were used to estimate the parameters, and the remaining data for 20 images were

used to evaluate the prediction. In the 10 fold CVs, all images were used once as test data. The prediction

performances were estimated using Pearson's correlation coefficients between the observed (trial-average) and

predicted responses. Encoding models were created for all preset threshold values for feature selection, and the

model that exhibited the best prediction performance was selected as the final model. In the analysis of

weights (i.e., feature) overlap between the two cells, the percentage of overlapping weights relative to the

number of non-zero weights was computed for each cell and averaged between the two cells in the pair.

Using the same dataset as used in the encoding model, The RF structure was estimated for each cell using

a regularized inverse method[29-31]. The regularized inverse method uses one hyper-parameter (regularized

parameter). In the 10-fold CVs, the RF structure was estimated with the training dataset using one of preset

regularized parameters (13 logarithmically spaced points between 10^{-3} and 10^{3}). The visual response was

predicted using the estimated RF and test dataset. The Prediction performance of visual response was

estimated by determining Pearson's correlation coefficients between the observed and the predicted responses.

RFs were estimated for all values of the pre-set regularized parameters, and the value that resulted in the best

response prediction was selected for the final RF model.

Image reconstruction.

In the image reconstruction, each Gabor feature value was linearly regressed by the single-trial activities of

multiple cells. In the 10 CVs, the weights and a bias parameter were estimated using the same algorithm as in

the encoding model with the training dataset (see above), and each Gabor feature value was reconstructed from

the visual response in the test dataset. After each Gabor feature was independently reconstructed, sets of

reconstructed feature values were transformed into images as described above (**Gabor features** section, also

see Fig. 3a). Reconstruction performance was evaluated by determining pixel-to-pixel Pearson's correlation

coefficient between the stimulus and reconstructed images. In the cell-selection model (Fig. 3), each feature

value was reconstructed with the subset of cells that was selected using the encoding model (Fig. 2f and

Supplementary fig. 3a), and almost all cells were used across features (Fig. 2j). In the encoding model, each

cell was represented by subset of features that affected the cell's response. Thus, in the cell-selection model,

each feature was only reconstructed by the cells that encoded information about the reconstructed feature

(Supplementary Fig. 3a).

In the analysis of the weights (i.e., feature) overlap between cells, the percentage of overlapping weights

relative to the number of non-zero weights was computed for each cell and averaged between the two cells in

the pair. The feature values of each image was linearly regressed with the weights of image reconstruction

model (cell-selection model) in all responsive cells in a local population to examine whether all the features of

natural images were represented by the features of the responsive cells (Fig. 5d–f). The fitting error rate (%

error)　was　computed　in　each　image　using　a　following　equation,　%　error　=

$\sum(F_{fitted}-F_{image})^2/\sum(F_{image}-F_{mean})^2\times100$, where F_{fitted} is the set of fitted features, F_{image} is the set of features of

the natural image, and F_{mean} is the mean of F_{image}.

Statistical analyses.

All data are presented as the median and 25–75Th percentiles, unless indicated otherwise. The significant

level was set to 0.05, with the exception of the criteria of significant visual response (0.01). When more than

two groups were compared, the significant level was adjusted with the Bonferroni correction. Two-sided test

was used in all analyses. The experiments were not performed in a blind manner. The sample sizes were not

predetermined by any statistical methods, but are comparable to the sample size of other reports in the field.

Data availability.

The data sets of the current study and the code used to analyze them are available from the corresponding

authors on reasonable request.

 Legends

Figure 1. Sparse visual response to a natural image in mouse V1

a. Experimental schema. Natural image flashes were presented as visual stimuli, and the neuronal activities of

single cells in the mouse V1 were recorded using two-photon Ca^{2+} imaging.

b. Trial-averaged time courses of visual responses to 10 natural images (image# 21–30 in a row) in 10

representative cells (cells# 1–10 in a column). Three lines for each response indicate the mean and the mean ±

the standard error across trials. Black: significant responses, grey: non-significant response, red line: stimulus

periods during which each image type was flashed for three times.

c. Plot of significant responses of all cells in a representative plane (n = 726 cells, upper left panel). Responses

shaded by the red line in the upper left panel correspond to responses presented in b. Percentage of responsive

cells for each image (bottom) and percentage of images to which each cell responded (right) are shown as line

graphs. Red lines (bottom and right) indicate median values. Cell numbers (cell #) were sorted by the

percentage of images to which they responded, and image numbers (image #) were sorted by the percentages

of cells that responded to each image in descending order. Single images activated relatively fewer neurons

(bottom).

d. Examples of population response patterns to three images. (Left column) Natural image stimuli and the

spatial distributions of responsive cells in an imaging area (side length: 507 microns). The red filled and grey

open circles indicate the highly responsive and remaining cells, respectively. (Right column) Histograms of the

visual responses of neurons in a local population. In the top panel, cells are divided into responsive (red bars)

and the remaining groups (black bars) and are sorted by the response amplitude of each group to the natural

image presented in the upper left panel (descending order). Visual responses to other images are plotted in the

middle and bottom panels. The cell # order was fixed among the three histograms. Only small numbers of

responsive neurons are duplicated among the three images.

e. Distribution of the amplitude of responses to single images. The cell # is sorted by the amplitude of the

response to each image and averaged across images in a plane. After normalizing cell# (x-axis), data were

collected across planes (n = 24 planes). The median (thick line) and 25–75th percentiles (thin lines) are shown.

Small percentages of neurons exhibited higher response amplitudes.

f and **g**. Response rate to natural images. The percentages of cells responding to a single natural image (f) and

to at least one image (g). Small percentage of cells responded to single natural image, whereas almost all cells

responded to at least one of images.

h and **i**. Response rate to moving gratings. Percentages of cells that responded to one direction (h) and to at

least one direction (i) of moving grating.

j. Percentages of overlapping responsive cells between the natural images. Only a small percentage of cells

exhibited overlapping significant responses between images, indicating that the cells responding to each image

were distributed in populations.

k. Population sparseness.

f–k. Each dot indicates data from one plane, and the medians of 24 planes are shown as bars

Figure 2. Small overlap in the encoded visual features among cells in a local population

a. Scheme of encoding model for a single cell's visual response. The visual response is represented by

weighted sum of the selected Gabor feature values obtained from a set of Gabor filters. The predicted visual

response to ith image (R_i) is represented by following equation, $R_i = f(\sum W_j \times F_{ji})$, where f is non-linear scaling

function, W_j is weight for jth Gabor feature, and F_{ji} is the feature value for the jth Gabor filter (G_j) obtained

from ith image (Si). Gabor feature was selected based on the correlation between its feature values and visual

response (see Methods).

b. and **c**. Examples of response predictions for two neurons. (Left panels) Blue and red lines indicate the

observed and predicted responses, respectively. (Right panels) Weight parameters of the example neurons

presented in the left panels. The weights of one of 10 models (each model corresponds to one of 10 CVs) are

shown. The number of non-zero weights (i.e., number of used feature) is shown above the panels. Encoding

filters (weighted sums of Gabor filters) are shown in the insets (red and blue indicate positive and negative

values, respectively).

d. Comparison of response predicted by only the linear step (regression of Gabor feature values) and the

observed response in the example neuron shown in Fig. 2c. Each dot indicates a response to one image. The

red curve indicates non-linear (NL) scaling function curve (see Methods). The NL step resulted in an

enhancement of the sparse visual responses. The black line indicates y = x line.

e. NL scaling function curves across planes. Each grey curve was obtained by averaging the NL scaling curves

across cells in each plane. Red curve indicates the averaged curve across planes (n = 24 planes). The black line

indicates y = x line.

f. Upper left panel: raster plot of the weights in the plane illustrated in Fig. 1c (red: positive weight, blue:

negative weight). The median values for the models of the 10-fold CVs are shown. Right panel: Percentage of

features used for each cell. Bottom panel: Percentage of cells in which each feature was involved in the

response prediction. The coloured bar under the x-axis indicates spatial frequency of the Gabor filter

corresponding to each feature. Red lines in the bottom and right panels indicate median values. Only half of

the Gabor features (624/1248 which have one of two phase sets) are shown for visibility, but the remaining

features were included in the data shown in the right panel.

g. Participation rate of each feature in the response predictions for a population. Features were divided in terms

of spatial frequency (SF) and averaged in each SF. Mean ± standard errors across all planes (n = 24 planes) are

shown.

h. Distribution of percentages of features used in each cell (n = 12755 cells across 24 planes).

i. Distribution of percentages of features that overlapped between cells (n = 3993653 cell pairs across 24

planes).

j. Percentage of features used in at least one cell's response prediction.

Figure 3. Image reconstruction based on population activity

a. Scheme of image reconstruction model. Each Gabor feature value (F_{ji}, i: image #, j: Gabor feature #) was

independently linearly regressed (weights: H_{jn}, n: cell #) by multiple cell responses (R_{ni}) to ith image (Si) ($F_{ji} =$

$\sum(H_{jn} \times R_{ni})$). Then, a set of reconstructed features (F_{1i}, F_{2i}, …., F_{ji}) are transformed into an image (S'$_i$). In the

all-cells model, each feature value was reconstructed with all cells. In the cell-selection model, only cells that

were selected by the encoding model were used to reconstruct each feature value (Cell selection, see Methods).

Thus, each feature value was reconstructed from different set of cells. The flow of the reconstruction model is

represented by black arrows from the bottom to top.

b. Representative reconstructed images. Stimulus images (Stim. Image), images that were reconstructed using

the all-cells (All-cells) and using the cell-selection model (Cell-selection) are shown. Each reconstructed

image is trial-averaged. The reconstruction performance (pixel-to-pixel Pearson's correlation coefficients

between the stimulus and reconstructed images, R) were computed in each trial and then averaged. The

trial-averaged R is presented above each reconstructed image.

c. Distributions of Rs of the all-cell model (upper panel) and cell-selection model (bottom panel) in the

representative plane shown in Figs. 1 and 2 (n = 200 images reconstructed using 726 cells from a plane).

d. R across planes. *: p = 4.0×10^{-4} using the signed-rank test (n = 24 planes). Reconstruction performance of

the cell-selection model was comparable with that of the all-cells model.

Figure 4. Sparse responsive cells encode sufficient information to represent the visual contents of

natural images.

a–c. Examples of images reconstructed from only the highly responsive cells (top panels), and reconstruction

performances (R) plotted against number of cells used for the reconstructions (bottom panels). In this analysis,

using the parameters of cell-selection model, the number of cells used to reconstruct each image increased one

by one. The cells were first collected from the responsive cells and then from the remaining cells. (Top panels),

Stimulus image (1st panel) and reconstructed images (trial-average) from a subset of cells (2nd–4th panels). R

and the number of cells used for the reconstruction are shown under the panel. (Bottom panels), Responsive

cells (red dots) and the remaining cells (black dots) were separately sorted by response amplitude (descending

order), and added cells one by one from responsive cells with higher response amplitude to the remaining cells

with lower response amplitude. The horizontal lines indicate 95% peak performances, and the numbers of cells

for which the performance curves crossed the 95% level are indicated by the vertical lines. In each case, the

responsive cells alone were sufficient to reconstruct the image nearly as well as all cells.

d. Averaged performance curve (n = 24 planes) plotted against cell number. The thick black line and grey lines

indicate the means and the means ± standard errors, respectively. The horizontal line indicates the 95 % peak

performance, and the vertical line indicates the first crossing point of the performance curve on the 95% line.

e. Contribution of the top 18 responsive cells to the image reconstruction shown in (a). (Top panels) Reverse

filters (weighted sum of Gabor filters) multiplied by the visual responses reveal the spatial patterns of

individual cell's contributions to the reconstructed image. The patterns vary among cells. (Bottom panels)

Reconstructed images were gradually changed by consecutively adding single cells.

f. Distributions of R for all cells (top, same as bottom of Fig. 3c) and the highly responsive cells (bottom) in

the representative plane (726 cells). N = 200 images. Red vertical lines indicate medians.

g, R for all cells and only the responsive cells. *: $P = 4.0\times10^{-4}$ by signed- rank test (n = 24 planes). Each dot

indicates data for each plane, and bars indicate median. The Rs were comparable between the two models.

h–j. Overlapping weights (i.e., features) between the cells highly responding to the same image.

h, Schema of the analysis.

i, Distribution of percentage of overlapping features in all cell pairs responding to the same image collected

across planes.

j. The median of the percentages of overlapping features in the cell pairs responding to the same image. Each

dot indicates the median in each plane (n = 24 planes) and the bar indicates median across planes. The

percentages of overlapping features were still low even in the cell pairs responding to the same image.

Figure 5. Visual features of natural images are distributed among most neurons in a population

a. Raster plots of highly responsive cells for each image in the representative plane shown in the previous

figures (n = 711/726 responsive cells). The image # is sorted by the image reconstruction performance

(descending order, right panel). In each line, cells that did not respond to the previously plotted images are

added on the right side. As image # increased, the number of newly added cells decreased, and then, cell #

quickly reached a plateau level, indicating that many images are represented by the combination of cells that

responded to other images. Thus, most images could be represented with some degree of accuracy by the

combination of subsets of responsive cells of the population.

b. The numbers of responsive cells (black line) and numbers of newly added responsive cells (red line) are

plotted against image # for the case shown in (a). Again, the number of newly added cells quickly decreased as

the image # increased.

c. The numbers of responsive cells (black line) and numbers of newly added responsive cells (red line) are

plotted against image #. N = 24 planes. Three lines in each colour indicate the mean and the mean ± the

standard errors.

d, Schema of the analysis. The feature set of each natural image was linear regressed by the weights from the

cell-selection model of all the responsive cells in each plane, and the fitting error (% error, see Methods) was

computed in each image. If the features encoded in all the responsive cells are sufficient to represent natural

images, the weights of the responsive cells work as basis functions to represent visual features of the natural

images.

e. Distributions of the errors of all images collected across planes.

f. The median of % error across planes (bar, n = 24 planes). Each dot indicates the median in each plane.

Supplementary figure legends

Supplementary Figure 1. Gabor filter sets and visual response predicted by the encoding model

a. Scheme of the process used to transform natural image features with Gabor filters. Each natural image was

subjected to Gabor filters to obtain the corresponding Gabor feature values. Conversely, a set of Gabor feature

values were transformed into an image by summing the Gabor filters after multiplying by the corresponding

Gabor feature values.

b. Scheme of the Gabor filter set. Four orientations, 2 phases, 4 scales (or spatial frequencies) of Gabor filters

were used. The 4 scales of Gabor filters (spatial frequency: 0.016, 0.033, 0.067, and 0.13 cpd) were positioned

on a 1×1, 3×3, 5×5, and 11×11 grids. A total of 1248 filters were used.

c. Effect of feature selection on the response prediction performance. The response prediction performances

(correlation coefficients between the observed and predicted responses) are plotted against the threshold values

of the feature selection for 30 representative cells. Pearson's correlation coefficient (corr. R) between each

feature values and response was computed, and the features with corr. R greater than the pre-set threshold

values were used for the regression analysis of the encoding model. The threshold value for the final model

was selected from the pre-set values to maximize the response prediction.

d and **e**. Averaged prediction performance plotted against the feature selection threshold among all cells in the

representative plane shown in the main figures (n = 726 cells from one plane, d) and among all planes (n = 24

planes, e). As the threshold increased, the prediction performance increased slightly and reached a plateau

level. The threshold was independently adjusted for each cell such that the response prediction was maximized.

Means ± standard errors are shown.

f–h. Results of the response predictions for the representative plane shown in the main figures (n = 726 cells

from one plane). Distributions of the response prediction performances (n = 200 images, f), percentages of

features used for each cell (n = 726 cells. g), and percentages of overlapping features between cells (n =

263175 pairs, h). In each cell pair, the number of overlapping features was divided by the number of features

used in the encoding model for each cell and averaged between the two cells in a pair.

i. Distribution of the response prediction performance of all cells across all planes (n = 12755 cells from 24

planes).

j–l. Results of the response prediction performance across planes (n = 24 planes). Distributions of the response

prediction performances (j), percentages of features used for each cell (k), and percentages of overlapping

features between cells (l) are shown.

Supplementary Figure 2. Relationship between the weights of the encoding model and the RF structure

a–d. Relationship between the weights of the Gabor features and the RF structure.

a and **b**. (a) Schema of the analysis. In each cell, the RF structure was determined using a pseudo-inverse method (see Methods), and pixel-to-pixel Pearson's correlation coefficients between the RFs and Gabor filters were computed (R1 and R2 in (a); R (RF vs. Gabor)). (b) Then, Pearson's correlation coefficient (R) between the R (RF vs. Gabor) and weight values (W1 and W2 in (a)) was computed for each cell. In the example cell shown in (b), the R (RF vs. Gabor) and weight values were positively correlated (R = 0.83), indicating that Gabor filters similar to the RF were assigned high weight value.

c and **d**. Distributions of R between R(RF vs. Gabor) and weights for all cells in the representative plane (c) and for all planes (d).

e–i. Relationship between the spatial overlap of the encoding filter and percentage of overlapping features between cells.

e. Examples of the response prediction performances of the two representative cells. The black and coloured lines indicate the observed and predicted responses, respectively.

f. Weights and encoding filters of the two cells shown in (e). In these cells, the percentage of overlapping non-zero feature (% overlapping feature) was 12%, and the spatial similarity of the encoding filters (pixel-to-pixel correlation, R (filter)) was 0.26. The encoding filter was obtained by computing the sum of Gabor filters multiplied by the weights of the encoding model. The red and blue colours of the encoding filter indicate positive and negative values, respectively.

g. Relationship between the % overlapping feature and absolute value of the R (filter). These two values were positively correlated, indicating that the % overlapping feature reflects the spatial similarity of the encoding filters. All pairs in the representative plane are illustrated by colour code.

h. Diverse structure of the encoding filters. The absolute R (filter) was relatively small for all pairs (median: 0.05), which indicated the diverse structure of the encoding filters between cells.

i. Distribution of the correlation coefficient between the % overlapping feature and absolute value of R (filter)

(|R(filter)|, N = 24 planes).

Supplementary Figure 3. Image reconstructions based on three models: all-cells, cell-selection

(encoding-optimized) and image-optimized models

a. Relationship between the encoding model (cyan arrows, Fig. 2 in the main text) and the image

reconstruction model (red arrows, Fig.3 in the main text). In the image reconstruction model, each Gabor

feature value was independently linearly regressed by the visual responses of multiple neurons. In the

"all-cells" model of image reconstruction, each feature value was reconstructed using all cells (not shown in

the figure). In the "cell-selection" model of image reconstruction, each feature was reconstructed using a

subset of cells (thin red arrows) selected by the encoding model (thin cyan arrows).

b–c. The image reconstruction performances (R) were compared across the three models, i.e., all-cells,

cell-selection (encoding-optimized) and image-optimized model. The performances were almost comparable

between the all-cells and cell-selection models (see the main text). The performance was only slightly

improved when the cell selection for each Gabor feature was performed to maximize the reconstruction of the

feature value ("image optimized" model in Supplementary Fig. 3b and c). In the image-optimized model, cells

were first selected based on the correlation between the cell's response and the feature value, similar to the

feature selection in the encoding model (see Methods). Then, each feature was linearly regressed only by the

selected cells. Threshold of the correlation coefficient for cell selection was determined such that the

reconstruction performance for each feature value was maximized.

b. Representative reconstructed images. Top panels: stimulus images; second row panels: reconstructed images

from the all-cells model; third row panels: reconstructed images from the cell-selection model (also called

"encoding-optimized" model hereafter in the legends); fourth row panels: reconstructed images from the

image-optimized models; bottom panels: reconstructed images using overlap of the cells for each feature

(overlap) between the encoding-optimized and image-optimized models. In the overlap model, each feature

value was reconstructed using the cells employed in both the encoding-optimized and image-optimized

models.

c. Comparison of R. Mean ± standard errors (n = 24 planes) are shown. Each dot indicates the average of all

images in each plane. The image-optimized model exhibited an only slightly better R than the other models. P

= 2.4×10^{-3} (all-cells vs. encoding-optimized), 1.1×10^{-4} (all-cells vs. image-optimized), 2.1×10^{-4} (all-cells vs.

overlap), 1.1×10^{-4} (encoding-optimized vs. image-optimized), 1.1×10^{-4} (encoding-optimized vs. overlap), and

1.1×10^{-4} (image-optimized vs. overlap), using signed rank test with the Bonferroni correction.

d. Percentages of cells used for each feature are plotted against the spatial frequencies of Gabor filters. The

image-optimized model (blue) used more cells to reconstruct each feature value than the encoding-optimized

model (i.e., cell-selection model, red). The plot of the encoding-optimized model is the same as the plot shown

in Fig.2g in the main text.

e. Overlap of the cells between the image-optimized and encoding-optimized models. In the reconstruction of

each feature value, the cells used in the encoding-optimized model were almost included in the cells used in

the image-optimized model. Image-only: percentage of cells used only in the image-optimized model.

Encoding-only: percentage of cells used only in the encoding-optimized model. Overlap: percentage of cells

used in both models. The results in (c–e) indicate that the overlapping cells mainly contributed to the image

reconstruction and the image-only cells contributed to the slight increase in R of the image-optimized model.

f–h. Comparison of reverse filters among the overlap, image-only and encoding-only cells.

f. Examples of reverse filters (reconstruction filters, i.e., the sums of Gabor filters multiplied by the weights in

each cell) in the overlap cells (upper panels), image-only cells (middle), and encoding-only cells (lower) used

to reconstruct the feature whose Gabor filter is shown in the left panel (reconstruction target). The center

panels presents reverse filters of single cells, and the right-most column presents averaged reverse filters

among the cells. The reverse filters of the overlap cells appear to be similar to the Gabor filter of the

reconstructed feature. In the image-only cells, the reverse filters of the individual cells do not appear to be

similar to the Gabor filter of reconstructed feature, whereas the average of reverse filters resembles the Gabor

filters. This suggests that, the overlapping cells represent the reconstructed feature at the single-cell level,

whereas the image-only cells represent the feature as a population.

g. Correlation between the reverse filters of single cells and the Gabor filters of the reconstructed feature. At

the single-cell level, the reverse filters of the overlapping cells (black line) are more similar to the Gabor filters

of reconstructed features than the reverse filters of the image-only and encoding-only cells. The similarity of

the reverse filter to the Gabor filter of reconstructed feature was indistinguishable between the image-only and

encoding-only cells (red and blue lines). Mean $\pm$ s.e. are shown. P = 1, 1, 0.087 and 0.080 (0.016–0.13 cpd) for

encoding-only vs. image-only; 2.4×10^{-3}, 2.2×10^{-4}, 2.2×10^{-4}, and 2.2×10^{-4} for encoding-only vs. overlap;

2.2×10^{-4}, 2.2×10^{-4}, 2.2×10^{-4}, and 2.2×10^{-4} for image-only vs. overlap; using signed rank test with the

Bonferroni correction.

h. Correlation between the averaged reverse filters and Gabor filters of the reconstructed feature. As a

population, the similarities of the average reverse filters of the image-only cells (blue) to the Gabor filters are

slightly less than the similarity of the overlapping cells (black) but greater than the encoding-only cells (red).

Based on these results, the image-only cells were responsible for the slight increase in R by representing the

reconstructed feature as a population. Mean $\pm$ s.e. are shown. P = 2.4×10^{-3}, 2.2×10^{-4}, 2.5×10^{-4}, and 2.5×10^{-4}

(0.016–0.13 cpd) for encoding-only vs. image-only; 2.4×10^{-3}, 2.2×10^{-4}, 2.2×10^{-4}, and 2.2×10^{-4} for

encoding-only vs. overlap; 4.1×10^{-4}, 2.2×10^{-4}, 2.2×10^{-4}, and 2.5×10^{-4} for image-only vs. overlap; using

signed rank test with the Bonferroni correction.

In summary, the cells used in both the encoding- and image-optimized models (i.e., overlapping cells)

likely represented the reconstructed feature at the single-cell level, whereas the additional cells that were used

only in the image-optimized model (i.e., image-only cells) probably represented the reconstructed feature as a

population. Thus, the main information about visual contents was obtained from the encoding-optimized

model (i.e., cell-selection model), and additional information was extracted when more cells that were not

captured by the cell-selection using encoding model were integrated into the reconstruction model.

Yoshida and Ohki, Figure 3

a

Stim.   Reconstructed

S_i S'_i

Gabor filter

Feature values F_{1i} F_{2i} F_{3i} $\cdots$

Weights H_{jn} H_{11} Σ H_{13} Σ $\cdots$ H_{1n}

Cell selection R_{1i} R_{2i} R_{3i} R_{4i} $\cdots$ R_{ni}

Responses
to ith image R_{1i} R_{2i} R_{3i} R_{4i} $\cdots$ R_{ni}

c n=200 images (1plane)

All-cells

# of images

Cell
selection

# of images

R

d

R

*

All-cells   Cell
selection

b

Stim. Image R: 0.60 0.64 0.71 0.56 0.49 0.59 0.73 0.80 0.70 0.73 0.66 0.52

All-cells R: 0.57 0.71 0.79 0.62 0.54 0.60 0.75 0.79 0.67 0.74 0.57 0.50

Cell
selection

a R: 0.62 R: 0.60 R: 0.57
   18 cells 57 cells 726 cells

b R: 0.74 R: 0.76 R: 0.71
   13 cells 27 cells 726 cells

c R: 0.61 R: 0.64 R: 0.60
   28 cells 83 cells 726 cells

d N = 24 planes

a Response prediction (Encoding)
Image reconstruction (Image reconstruction)

Natural image ⟶ Feature values ⟶ Cell's visual responses

F_1 — R_1
F_2 — R_2
⋮ — ⋮
F_i — R_j

⟶ Feature selection in encoding model
⟵ Cell selection by encoding model

b
Image
All cells
Cell-selection (encoding optimized)
Image optimized
Overlap

c
R
All cells
Encoding optimized
Image. optimized
Overlap

d
% cells/features
Image. optimized
Encoding optimized
Spatial frequency(cpd)

e
% cells/features
Image only
Overlap
Encoding. only
Image optimized
Encoding. optimized
Spatial frequency(cpd)

f
Reverse filters
Single cell filter   Averaged filter
Reconstruction target
Overlap — Ave.
Image. only — Ave.
Encoding only — Ave.

g Reverse filter (single cell) vs. Gabor filter (target)
R
Spatial frequency (cpd)

h Reverse filter (average) vs. Gabor filter (target)
Overlap
Image only
Encoding. only
Spatial frequency (cpd)