Epigenetic changes induced by *Bacteroides fragilis* toxin (BFT)

Jawara Allen[1], Stephanie Hao[2], Cynthia L. Sears[1,3]*, Winston Timp[2]*

1: Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA

2: Department of Biomedical Engineering, Johns Hopkins University School of Medicine,

Baltimore, MD USA

3: Bloomberg-Kimmel Institute for Immunotherapy and Department of Oncology, Sidney

Kimmel Comprehensive Cancer Center, Johns Hopkins Medicine Institutions, Baltimore, MD,

USA

*Co-corresponding authors

Running head: BFT-induced Epigenetic changes

Address correspondence to Cynthia L. Sears (csears@jhmi.edu) or Winston Timp
(wtimp@jhu.edu)
15
16   Cynthia L. Sears, M.D.
17   Johns Hopkins University School of Medicine
18   1550 Orleans Street
19   CRB2 Building, Suite 1M.05
20   Baltimore, MD 21231
21

Winston Timp
Johns Hopkins University
Department of Biomedical Engineering
3400 N. Charles St.
Clark 118A
Baltimore, MD 21218
28

Abstract (limit:250):

Enterotoxigenic *Bacteroides fragilis* (ETBF) is a gram negative, obligate anaerobe member of the

gut microbial community in up to 40% of healthy individuals. This bacterium is found more

frequently in people with colorectal cancer (CRC) and causes tumor formation in the distal

colon of mice heterozygous for the <u>a</u>denomatous <u>p</u>olyposis <u>c</u>oli gene ($Apc^{+/-}$); tumor formation

is dependent on ETBF-secreted *Bacteroides fragilis* toxin (BFT). Though some of the immediate

downstream effects of BFT on colon epithelial cells (CECs) are known, we still do not

understand how this potent exotoxin causes changes in CECs that lead to tumor formation and

growth. Because of the extensive data connecting alterations in the epigenome with tumor

formation, initial experiments attempting to connect BFT-induced tumor formation with

methylation in CECs have been performed, but the effect of BFT on other epigenetic processes,

such as chromatin structure, remains unexplored. Here, the changes in chromatin accessibility

(ATAC-seq) and gene expression (RNA-seq) induced by treatment of HT29/C1 cells with BFT for

24 and 48 hours is examined. Our data show that several genes are differentially expressed

after BFT treatment and these changes correlate with changes in chromatin accessibility. Also,

sites of increased chromatin accessibility are associated with a lower frequency of common

single nucleotide variants (SNVs) in CRC and with a higher frequency of common differentially

methylated regions (DMRs) in CRC. These data provide insight into the mechanisms by which

BFT induces tumor formation. Further understanding of how BFT impacts nuclear structure and

function in vivo is needed.

Importance (limit:150):

Colorectal cancer (CRC) is a major public health concern; there were approximately 135,430

new cases in 2017, and CRC is the second leading cause of cancer-related deaths for both men

and women in the US (1). Many factors have been linked to CRC development, the most recent

of which is the gut microbiome. Pre-clinical models support that enterotoxigenic *Bacteroides*

fragilis (ETBF), among other bacteria, induce colon carcinogenesis. However, it remains unclear

if the virulence determinants of any pro-carcinogenic colon bacterium induce DNA mutations or

changes that initiate clonal CEC expansion. Using a reductionist model, we demonstrate that

BFT rapidly alters chromatin structure and function consistent with capacity to contribute to

CRC pathogenesis.

Introduction (limit for body: 5000)

Bacteroides fragilis is a gram-negative, obligate anaerobe that is found consistently, but in low

numbers, in the gut microbial community (2). ETBF is a particular subtype of *Bacteroides*

fragilis, characterized by its production of *Bacteroides fragilis* toxin (BFT) and its association

with diarrhea, inflammatory bowel disease and colon cancer(3, 4). Since the identification of

ETBF, many studies have examined the carriage rate of this potentially pathogenic bacterium.

Some studies have probed the mucosa for the bacterium and found an asymptomatic carriage

rate as high as 67% (4), but most studies examine the stool and report rates of 10%-12% (3).

So, while we do not yet know exactly how many people are colonized with ETBF, it is clear that

a significant fraction of the population likely experiences prolonged exposure to the bacterium,

and thus BFT.

The *bft* gene codes for the BFT pre-pro-protein that is processed by ETBF, yielding a secreted

active 20 kD zinc-dependent metalloprotease (3). There are three isotypes of the BFT protein,

each produced by a different *bft* gene (3), with the most potent of the BFT isotypes being BFT2.

Expression of BFT2 is required for ETBF-induced tumorigenesis in multiple intestinal neoplasia

($Apc^{min/+}$) mice (5). When interacting with colon epithelial cells (CECs), BFT2 binds to an

unidentified cell surface receptor which leads to rapid, but indirect, cleavage of E-cadherin that,

in turn enhances barrier permeability *in vivo* (6). Because a pool of β–catenin is associated with

the intracellular domain of E-cadherin, cleavage of E-cadherin triggers release of β–catenin and

its subsequent nuclear localization with upregulation of c-Myc expression and enhanced cellular

proliferation (7).

Though we know some of the immediate downstream effects of BFT on CECs, we still do not

understand how this potent exotoxin causes lasting changes in CECs that lead to tumor

formation and growth. A possible explanation is an epigenetic progenitor model to cancer, in

which epigenetic dysregulation enables tumor cell survival (8). Epigenetics is the study of

heritable phenotypic variation that is not caused by base pair changes in the genome.

Epigenetic changes have been linked to various cancers; in colorectal cancer specifically,

hypermethylation of CpG islands and global hypomethylation have both been implicated in

tumor development and progression (9) and global dysregulation of the methylome has been

observed in clinical colon cancer samples (10).

Initial experiments exploring the impact of BFT on the epigenome have been promising. By

tracking the localization of specific epigenetic regulators, one study showed that inoculation of

C57BL/6J mice with ETBF causes an upregulation of gene silencing complexes at CpG islands

(11). Subsequent studies have shown that inoculation of Apc$^{min/+}$ mice with ETBF also causes

recruitment of DNA methyltransferase 1 (DNMT1), a recruitment potentially mediated by DNA

mismatch repair proteins (12). While these studies have established a role for gene regulation

and DNA methylation in the effect of ETBF on CECs, studies do not report on how ETBF affects

other epigenetic processes, such as chromatin structure. This is of particular importance

because chromatin state as measured by changes in various histone marks has been linked to

colon cancer development (9), and differences in chromatin structure have been shown to

impact mutation rates along the genome (13)(14).

Herein, we further test the hypothesis that the epigenome is at least one mechanism by which

BFT acts to increase colon tumor formation. To test our hypothesis, we use a reductionist

model system to investigate the effect of BFT2 on CEC chromatin structure. Specifically, we

examined whether BFT2 alters chromatin accessibility and subsequently, gene expression, using

a model human colon epithelial cell line, HT29/C1, that is known to be uniformly sensitive to

BFT2 and for which extensive prior data exist (3). We extended our analyses to seek to

determine if BFT2-induced chromatin changes were associated with functional consequences

by correlation with transcription factor binding motifs and known cancer-causing mutations in

human CRC.

Results

BFT2 induces dynamic changes in chromatin accessibility throughout the genome

We initially examined time points of 24 and 48 hours after BFT treatment (fig 1A), performed in

biological triplicate. To assay chromatin state, we performed ATAC-seq (Assay for Transposase-

Accessible Chromatin using sequencing) on BFT2-treated and untreated cells at each time point

(15). Briefly, ATAC-seq leverages the accessibility of chromatin to preferentially "attack" open

chromatin using a transposase (Nextera; Illumina), resulting in a tagmentation reaction.

Subsequent preferential amplification and downstream sequencing results in a distribution of

sequencing reads which reveals areas of open versus closed chromatin by examination of the

depth of coverage (i.e., the number of reads which align to each area of the genome). We

utilized software tools like MACS2 (see Methods) to detect peaks in the coverage that indicate

open chromatin regions.

We first quantified the total number of peaks of chromatin accessibility in our baseline samples

(untreated cells) and BFT2-treated samples. We then classified these peaks into 2 types at 24

and 48 hours: BFT2-opened peaks (peaks of chromatin accessibility detected only after BFT2

treatment), and BFT2-closed peaks (peaks of chromatin accessibility that disappeared after

BFT2 treatment). This initial analysis revealed that only a fraction (40% at 24 hours and 35% at

48 hours) of the peaks identified by ATAC-seq are affected by BFT2 treatment, and that BFT2

treatment appears more likely to open chromatin than to close it (~11:3 ratio at 24 hours; ~5:3

ratio at 48 hours) (fig 1B). We next analyzed the overlap of our BFT2-opened peaks and BFT2-

closed peaks between 24 and 48 hours in order to determine if BFT2-induced changes in

chromatin accessibility are stable. We found that only 13.4% (4989 peaks) of 24 hours and 48

hours BFT2-opened peaks overlap, and 4.55% (743 peaks) of 24 hours and 48 hours BFT2-closed

peaks overlap (fig 1B). These results emphasize the dynamic nature of BFT2-induced changes of

chromatin accessibility.

To explore how BFT2 affects peak distribution throughout the genome, we performed a

detailed analysis on the BFT2-opened and BFT2-closed chromatin regions (i.e. the regions of the

genome affected by BFT2 treatment). When we analyze the distribution of these peaks, we see

that more of BFT2-opened and BFT2-closed peaks are located within intergenic regions and

coding regions than promoter regions (Table 1). To determine if this distribution was a

byproduct of the ATAC-seq assay or a specific effect of BFT2, we compared those peak

distributions to the distribution present in our baseline and BFT2-treated samples. This analysis

reinforced that more of the BFT2-opened and BFT2-closed peaks are located within intergenic

and coding regions, while less are located within promoter regions (Table 1). Although we

expect that the effect in all three regions is important, these results suggest that chromatin

accessibility is affected differentially across the genome by BFT2.

BFT2 treatment induces changes in gene expression that are more pronounced at 24 hours

than 48 hours

To examine the relationship between gene expression and chromatin state, we performed

RNA-seq on the same samples used for ATAC-seq analysis. While the effect of BFT2 treatment

on select genes at relatively early time points has been studied (16, 17), how this toxin affects

genome wide expression and whether these effects persist is still unknown. After treatment

with BFT2 for 24 hours, 70 genes were differentially expressed (P-value < 0.01). Of these genes,

41 showed a decrease in gene expression, while 29 showed an increase in gene expression (fig

2A, supplementary table 1). After BFT2 treatment for 48 hours, we found only 16 differentially

expressed genes (P-value < 0.01); 3 showed a decrease in gene expression and 13 showed an

increase in gene expression (fig 2B, supplementary table 2). This result follows a trend seen

throughout our analyses and the literature (7, 16, 18) in which BFT2 has a more potent effect

on CECs at 24 hours after toxin treatment that seems to diminish by 48 hours. A lineplot of the

top 16 differentially expressed genes at 24 and 48 hours shows that gene expression at the two

time points tends to differ; genes differentially expressed at 24 hours tend to move closer to

baseline levels by 48 hours, while genes differentially expressed at 48 hours tend to have

expression levels closer to baseline at 24 hours (supplementary fig 1). Notably, none of the

genes differentially expressed at 24 hours show statistically significant differential expression at

48 hours as well. These data reinforce the observation that BFT2-induced gene expression

changes are highly dynamic, and time specific, similar to BFT2-induced changes in chromatin

accessibility.

To perform a Gene Set Enrichment (GSE) analysis on the list of genes that were differentially

expressed at 24 hours and 48 hours after BFT2 treatment, the cutoff for significance was

relaxed to a P-value of less than 0.05. The GSE analysis revealed several processes that were

overrepresented in our list of upregulated and downregulated genes (fig 2c). Specifically, at 24

hours after BFT2 treatment, expression of transcription factors involved in nucleic acid binding

were downregulated. At 48 hours after BFT2 treatment, genes related to the cell cycle,

chromosome organization and response to DNA damage, among others, were upregulated (fig

2C).

Chromatin accessibility correlates with gene expression at baseline and at 24 hours after BFT2

treatment

We next tested the hypothesis that BFT2-induced changes in chromatin accessibility correlate

with BFT2-induced changes in gene expression. First, we classified genes by promoter

chromatin accessibility status in our 24hr untreated (baseline) sample. Only promoter peaks

were used because of the intuitive connection between open promoter chromatin and changes

in gene expression. Those genes that contained an ATAC-seq peak overlapping their promoter

region were categorized as "open" and those without a peak were categorized as "closed". We

then extracted normalized gene expression values (transcripts per million- TPM) for each of

these genes to determine if the average gene expression values were different for the two

groups. We found that on average, gene expression was statistically higher among genes that

were "open" in our 24hr baseline sample compared to those that were "closed" (P-value < 2e⁻

[16]) (fig 3A).

Next, we examined the correlation between changes in chromatin accessibility and changes in

gene expression at the same time points. To do this, we again focused on BFT-opened and BFT-

closed peaks which overlapped with gene promoter regions. We then calculated the

logarithmic fold-change (Log2FC) expression values for each of these genes at 24 or 48 hours

after BFT2 treatment. We found that genes with a BFT2-opened peak overlapping their

promoter had a three-fold higher Log2FC in gene expression after 24 hours of BFT2 treatment

than genes with a BFT2-closed peak at the same time point. At 48 hours after BFT2 treatment,

we found no difference in Log2FC gene expression (fig 3 B-C). A similar analysis can be

performed for the 86 genes that show statistically significant differences in gene expression

after BFT2 treatment (fig 2 A-B, Supplementary tables 1, 2). Among those, 7 contained BFT2-

opened peaks overlapping their promoter (6 at 24 hours and 1 at 48 hours). Of these genes, 5

showed an increase in gene expression, while 2 showed a decrease. No BFT2-closed peaks

overlapped with the promoters of these 86 genes at either time point (supplementary tables 1,

2). Taken together, these results suggest that at 24 hours after BFT2 treatment, changes in

chromatin structure largely correlate with changes in gene expression, but this relationship

disappears by 48 hours.

Treatment with BFT2 causes increased chromatin accessibility at transcription factor binding

sites

Gene expression is regulated via multiple mechanisms, including the binding of transcription

factors to enhancers and gene promoters. Therefore, we also wanted to test the hypothesis

that BFT2-induced changes in chromatin accessibility impact transcription factor binding sites

and thus gene expression. This can be done by looking for specific transcription factor motifs at

sites of altered chromatin accessibility, or by calculating the overlap of sites of altered

chromatin accessibility with sites of transcription factor binding. This latter approach is only

available for a handful of transcription factors, such as CTCF, which have been extensively

profiled via Chip-seq experiments in various cell types (19).

We first used the haystack_bio bioconda package to query our ATAC-seq data for specific

transcription factor motifs. This analysis revealed several transcription factor motifs that are

enriched in our BFT2-treated or baseline samples (Tables 2,3; Supplementary tables 5,6). In

contrast to the BFT2-dependent dynamic chromatin changes identified in our earlier analyses

(fig 1B, Table 1), many of the most enriched transcription factor motifs identified are present

after both 24 and 48 hours of BFT2 treatment. Notably, these include JUND, JDP2, FOSL1, JUNB,

FOS, FOS:JUN and ZBTB33, many of which are downstream of mitogen-activated protein kinase

(MAPK) pathways previously shown to be modulated by BFT2 (18, 20, 21). In subsequent

analyses, we specifically examined FOSL1, as it was the only transcription factor whose

expression was upregulated 24 hours after BFT2 treatment. We noted that while several

FOSL1-regulated genes were also upregulated after BFT2 treatment, a near equal proportion

were also downregulated (supplementary tables 3,4). For FOSL1 to augment gene expression, it

needs to bind to an accessible DNA-binding site, so we next examined the FOSL1-regulated

genes and classified them according to accessibility 24 hours after BFT2 treatment, as

determined by ATAC-seq. We found that 6 of 7 genes with accessible DNA binding sites

showed an increase in expression after BFT2 treatment, and only 1 of 6 genes without an

accessible DNA binding site showed an increase in expression after BFT2 treatment

(supplementary tables 3,4).

While changes in promoter chromatin structure have been most extensively investigated, and

provide many avenues for exploration, our data indicate that BFT2-induced changes in

chromatin structure occur more frequently in intergenic regions. We hypothesized that these

intergenic chromatin accessibility changes may affect chromatin architecture more globally and

sites of CTCF binding specifically. CTCF is a DNA binding protein that regulates global chromatin

structure by mediating chromatin looping via binding to and bringing together distant regions

of the genome (19). The CTCF motif was enriched in our ATAC-seq data after 24 hours, but not

48 hours, of BFT2 treatment (Tables 2, 3). To further investigate this enrichment, we calculated

a jaccard index for the overlap of CTCF binding sites in CACO-2 cells and HT29/C1 cells (the

latter using our ATAC-seq data); both cell lines are human transformed colon epithelial cell

lines. We found that after 24 hours of BFT2 treatment, but not 48 hours, the HT29/C1/CACO-2

jaccard index was increased by 1.13 fold (data not shown). These results again suggest that

sites in the genome that show increases in chromatin accessibility after BFT2 treatment are

more likely to also be sites of CTCF binding.

Chromatin accessibility is associated with differential DNA methylation and DNA mutation

The previous analyses help us better understand how BFT2 alters chromatin accessibility in CECs and to connect chromatin changes with anticipated gene expression changes. However, they do not help us understand how changes in chromatin accessibility may contribute to common DNA-modifications found in CRC. To explore this, we looked for a correlation between BFT2-induced changes in chromatin accessibility and single nucleotide variants (SNVs) and common sites of differential methylation (DMRs) in CRC. SNVs and DMRs were extracted from the COSMIC database. We calculated the proportion of peaks in each sample that overlapped with a common SNV or DMR. We then used a chi-square test to compare our treated and untreated samples in order to determine if the proportion of peaks overlapping a SNV or DMR differed after treatment of HT29/C1 cells with BFT2.

Correlating our chromatin accessibility changes with SNVs in CRC allows us to explore the effects of BFT2-induced chromatin accessibility changes in coding regions of the genome, where specific chromatin structure states have been linked to decreased mutation rates in cancer (13). In contrast, examining DMRs allows us to explore whether BFT2-induced chromatin accessibility is associated with hypermethylation of CpG islands in promoter regions, or hypomethylation of short interspersed nucleotide elements (SINEs) or long interspersed nucleotide elements (LINEs) in intergenic regions.

Our SNV analysis revealed a smaller proportion of peaks overlapping with common SNVs in CRC

24 hours [OR(95%CI): 0.819(0.794-0.844)] and 48 hours [OR(95%CI): 0.964(0.936-0.993)] after

BFT2 treatment (Table 4). Similar to our prior data, the effect of BFT2 was more pronounced 24

hours after cell treatment. Thus, contrary to our hypothesis, these data suggest that regions of

the genome with BFT2-induced increases in chromatin accessibility are *less* likely to contain

SNVs commonly found in CRC.

In contrast, our DMR analysis found that a larger proportion of BFT2-induced chromatin

accessibility peaks overlap with common DMRs in CRC 48 hours [OR(95%CI): 1.094(1.018-

1.176)], but not 24 hours [OR(95% CI): 1.037(0.965-1.114)] after BFT2 treatment (Table 4). This

result supports a commonly held paradigm that chromatin changes come first, followed by

methylation changes (22). When we further parsed the data to analyze regions of

hypermethylation and hypomethylation individually, the results differed slightly, but

consistently showed a larger proportion of peaks overlapping with common DMRs in CRC after

BFT2 treatment (Table 4). Thus, analysis of DMRS supported our hypothesis that BFT2-induced

regions of increased chromatin accessibility are *more* likely to contain sites of differential

methylation found in CRC.

Discussion

Our data support that acute BFT2 treatment of colon epithelial cells (CECs) leads to rapid onset

and dynamic, but limited changes in chromatin accessibility. These changes in chromatin

accessibility could be associated with BFT2-induced changes in gene expression. We observed

changes in both chromatin accessibility and gene expression over a 48-hour period, with the

strongest changes occurring at 24 hours, then relaxing to baseline by 48 hours. Our results

corroborate previous observations about the acute onset of BFT2 action on epithelial cells in

vitro and in vivo (7, 16, 18).

Our RNA-seq data identifies several genes and pathways that are differentially regulated after

BFT2 treatment. The pathways identified, particularly at 48 hours, include those involved in the

DNA damage response and cell cycle regulation, both germane to previously reported BFT-

induced CEC changes in vitro and in vivo. Namely, BFT2 has been shown to induce DNA double

strand breaks and modulate apoptosis *in vitro* (17, 20) and ETBF similarly induces DNA double

strand breaks (16) as well as colon carcinogenesis *in vivo* (23). In addition, our study also

identified several new genes that may explain how BFT2 contributes to tumorigenesis by

allowing other bacteria to invade the mucus layer and bind to CECs. Specifically, BFT2 causes

upregulation of CEACAM6 and MUC16, and downregulation of MUC2. CEACAM6 acts as a

receptor for adherent-invasive *E.coli*, a particular strain of *E. coli* that has recently been

implicated in the development/progression of CRC in individuals with FAP (24) and has

previously been associated with sporadic CRC (25). MUC2, the major MUC protein present in

human colonic mucus, is notably downregulated after BFT2 treatment, potentially contributing

to mucus invasion by tumorigenic *E.coli*. The upregulation of MUC16, a particular MUC variant

that is not normally expressed in the colon, may represent a compensatory mechanism (i.e., an

attempt to protect CECs after MUC2 downregulation). Collectively, these results may begin to

explain how ETBF works with other bacteria to promote tumor formation and growth.

Several previous studies have shown that chromatin structure in gene promoter regions

correlates with gene expression (26, 27). Our study took this a step further by showing that

acute changes in chromatin accessibility correlate with acute changes in gene expression. 24

hours after BFT2 treatment, increases in chromatin accessibility generally correlated with

increases in gene expression while decreases in chromatin accessibility correlated with

decreases in gene expression. On the other hand, at 48 hours, no significant association is seen

between changes in chromatin accessibility and gene expression. This lack of association is

likely due to the limited gene expression changes that remain 48 hours after BFT2 treatment

and underscores a trend that is seen throughout most of our data whereby changes due to

BFT2 treatment are greater 24 hours after treatment and approach baseline levels by 48 hours.

This pattern is also corroborated by previous *in vitro* data that has shown that the effect of

BFT2 on gene expression in CECs is rapid and appears to decay quickly (7, 16, 18), although

biologic effects such as CEC proliferation persist for at least 72 hours *in vitro* (7).

Areas of increased chromatin accessibility after BFT2 treatment are enriched in binding motifs

for many transcription factors belonging to the AP-1/ATF family of transcription factors,

including FOSL1, JUN and JDP. These transcription factors function downstream of the JNK

pathway known to be activated by BFT2 (18, 20, 21) . After BFT2 treatment, we expected to see

increased expression of genes which are regulated by these particular transcription factors, but

chromatin opening alone appeared insufficient to drive changes in gene expression of these JNK

pathway genes. Hence, we looked for transcription factors that had both an enriched binding

motif and increased gene expression after BFT2 treatment. FOSL1 was the primary candidate

revealed by our analysis. We found that though presence of a FOSL1 binding site alone was

unable to predict differential gene expression after BFT2 treatment, when coupled to promoter

chromatin accessibility, an excellent concordance with increased gene expression was detected.

This finding exemplifies how integration of chromatin accessibility and gene expression data

can be applied to understand how BFT2 acts on CECs. We needed to use both types of data to

correctly surmise that transcription factor upregulation, enrichment of transcription factor

binding motifs, and the presence of open chromatin in the promoter region all contribute to

BFT2-induced gene regulation.

Because CRC usually results from a series of genetic mutations, and can be transformed by

changes in methylation patterns, any potential interaction between BFT2, chromatin

accessibility, and common SNVs/DMRs in CRC is important. Herein, we found a BFT2-induced

increase in chromatin accessibility at sites of common DMRs in CRC and a *decrease* at sites of

common SNVs in CRC. Increased methylation after inoculation of mice with ETBF has been

suggested previously. Specifically, studies by O'Hagan and colleagues have shown that binding

of DNMT1 (DNA methyltransferase 1) at promoters of low-expressing genes is upregulated

after ETBF murine colonization (11). While this study implicates ETBF, and thus BFT, in DNA

hypermethylation, no studies to date have explored the connection between BFT and

hypomethylation. Our data suggest that BFT2-induced changes in chromatin accessibility may

have dual effects on chromatin methylation patterns (fig 4). We hypothesize that increased

chromatin accessibility is, at least, one step critical to chromatin methylation pattern changes in

response to BFT2. This is important because hypermethylation can lead to silencing of tumor

suppressor genes and hypomethylation can lead to upregulation of transposable elements that

may reintegrate into tumor suppressor or proto-oncogenes. In the future, bisulfite sequencing

experiments can be conducted to definitively correlate sites of methylation change after BFT2

treatment with sites of altered chromatin accessibility.

Several experiments, conducted across a variety of species, have also shown that chromatin

accessibility may be inversely correlated with rates of specific DNA mutations (28–30). One

study in particular showed that DNase accessible euchromatin was protected from UV-induced

DNA damage, while lamina-associated heterochromatin was more susceptible to damage (14).

This study is of particular importance because it investigated carcinogenesis-related DNA

damage. Our data suggest that BFT2-induced decreases in chromatin accessibility are

correlated with frequently mutated areas, suggesting that BFT2-induced chromatin changes

may increase the chances of mutations. Such a hypothesis would require significant further

testing to determine its accuracy – this could be done by sequencing BFT2-treated and

untreated cells in order to determine if BFT2 treatment increases the mutation rate. If so, sites

of increased mutation frequency could be identified and correlated with sites of BFT2-induced

chromatin accessibility change.

Our study has several limitations that provide avenues for future experiments. Most notably,

our experiments were performed on HT29/C1 cells, a colon carcinoma cell line. These cells

contain several mutations that may lead them to respond differently to BFT2 than normal CECs.

As a result, these experiments should be confirmed in a primary cell culture system, or ideally,

in vivo in a mouse model or from human samples with and without ETBF colonization. Our

study, by design, was also performed over only an early time window. *In vitro* studies with BFT

best mimic the earliest *in vivo* events whereby ETBF induces CEC changes and colitis by ~24

hours after murine colonization (31). In contrast, the persistent CEC exposure to BFT afforded

by chronic ETBF colonization *in vivo* cannot be reliably modeled *in vitro*, in part, because of CEC

uptake and degradation of BFT (32). Thus, additional *in vivo* studies are warranted since ETBF

murine colonization is persistent and associated with ongoing IL-17-dominant inflammation and

CEC hyperplasia after one year in C57Bl/6 mice (31, 33). Ultimately, we want to know which

BFT-induced chromatin and gene expression changes persist and contribute to tumor

formation. Verifying these experiments in an in vivo system will help answer some of the

questions proffered by these data.

The experiments reported herein provide additional insight into the effect of BFT2, a bacterial

exotoxin linked to CRC pathogenesis, on CECs. We have identified new associations between

BFT2-induced chromatin accessibility and gene expression changes, and also correlated these

changes with previously published data on BFT2-induced signal transduction and DNA

modifications that may contribute to tumor formation or growth. We need to better

understand how BFT2 affects the genome and epigenome of CECs to determine if

asymptomatic carriers of ETBF are at increased risk for colon tumorigenesis.

Acknowledgements: This work was supported by the Bloomberg Philanthropies (CLS), National

Institutes of Health R01CA179440 (CLS), the Johns Hopkins Department of Medicine (CLS), and

Howard Hughes Medical Institute (JA)

All authors declare no completing interests.

Works Cited

1.   National Cancer Institute NI of H. Cancer of the Colon and Rectum - Cancer Stat Facts.

2.   Wexler HM. 2007. Bacteroides: The good, the bad, and the nitty-gritty. Clin Microbiol

Rev.

3.   Sears CL. 2009. Enterotoxigenic Bacteroides fragilis: A rogue among symbiotes. Clin

Microbiol Rev.

4.   Boleij A, Hechenbleikner EM, Goodwin AC, Badani R, Stein EM, Lazarev MG, Ellis B,

Carroll KC, Albesiano E, Wick EC, Platz EA, Pardoll DM, Sears CL. 2015. The bacteroides

fragilis toxin gene is prevalent in the colon mucosa of colorectal cancer patients. Clin

Infect Dis 60:208–215.

5.   Chung L, Thiele-Orberg E, Geis A. 2017. Bacteroides Fragilis Toxin Coordinates a Pro-

Carcinogenic Inflammatory Cascade via Targeting of Colonic Epithelial Cells. Submitted.

6.   Wu S, Lim KC, Huang J, Saidi RF, Sears CL. 1998. Bacteroides fragilis enterotoxin cleaves

the zonula adherens protein, E-cadherin. Proc Natl Acad Sci U S A 95:14979–84.

7.   Wu S, Morin PJ, Maouyo D, Sears CL. 2003. Bacteroides fragilis enterotoxin induces c-

Myc expression and cellular proliferation. Gastroenterology 124:392–400.

8.   Timp W, Feinberg AP. 2013. Cancer as a dysregulated epigenome allowing cellular

growth advantage at the expense of the host. Nat Rev Cancer 13:497–510.

9.   Esteller M. 2008. Epigenetics in Cancer. N Engl J Med 358:1148–1159.

10. Hansen KD, Timp W, Bravo HC, Sabunciyan S, Langmead B, McDonald OG, Wen B, Wu H,

Liu Y, Diep D, Briem E, Zhang K, Irizarry RA, Feinberg AP. 2011. Increased methylation

variation in epigenetic domains across cancer types. Nat Genet 43:768–775.

11. O'Hagan HM, Wang W, Sen S, DeStefano Shields C, Lee SS, Zhang YW, Clements EG, Cai Y,

Van Neste L, Easwaran H, Casero RA, Sears CL, Baylin SB. 2011. Oxidative Damage Targets

Complexes Containing DNA Methyltransferases, SIRT1, and Polycomb Members to

Promoter CpG Islands. Cancer Cell 20:606–619.

12. Maiuri AR, Peng M, Sriramkumar S, Kamplain CM, DeStefano Shields CE, Sears CL,

O'Hagan HM. 2017. Mismatch Repair Proteins Initiate Epigenetic Alterations during

Inflammation-Driven Tumorigenesis. Cancer Res 77:3467–3478.

13. Schuster-Böckler B, Lehner B. 2012. Chromatin organization is a major influence on

regional mutation rates in human cancer cells. Nature 488:504–507.

14. García-Nieto PE, Schwartz EK, King DA, Paulsen J, Collas P, Herrera RE, Morrison AJ. 2017.

Carcinogen susceptibility is regulated by genome architecture and predicts cancer

mutagenesis. EMBO J e201796717.

15.   Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. 2013. Transposition of native

chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding

proteins and nucleosome position. Nat Methods 10:1213–1218.

16.   Goodwin AC, Destefano Shields CE, Wu S, Huso DL, Wu X, Murray-Stewart TR, Hacker-

Prietz A, Rabizadeh S, Woster PM, Sears CL, Casero RA. 2011. Polyamine catabolism

contributes to enterotoxigenic Bacteroides fragilis-induced colon tumorigenesis. Proc

Natl Acad Sci U S A 108:15354–9.

17.   Kim JM, Lee JY, Kim Y-J. 2008. Inhibition of apoptosis in Bacteroides fragilis enterotoxin-

stimulated intestinal epithelial cells through the induction of c-IAP-2. Eur J Immunol

38:2190–9.

18.   Wu S, Powell J, Mathioudakis N, Kane S, Fernandez E, Sears CL. 2004. Bacteroides fragilis

enterotoxin induces intestinal epithelial cell secretion of interleukin-8 through mitogen-

activated protein kinases and a tyrosine kinase-regulated nuclear factor-??B pathway.

Infect Immun 72:5832–5839.

19.   Holwerda SJB, de Laat W. 2013. CTCF: the protein, the binding partners, the binding sites

and their chromatin loops. Philos Trans R Soc B Biol Sci 368:20120369–20120369.

20.   Ko SH, Rho DJ, Jeon JI, Kim YJ, Woo HA, Lee YK, Kim JM. 2016. Bacteroides fragilis

enterotoxin upregulates heme oxygenase-1 in intestinal epithelial cells via a mitogen-

activated protein kinase- and NF-κB-dependent pathway, leading to modulation of

apoptosis. Infect Immun 84:2541–2554.

21.   Hess J. 2004. AP-1 subunits: quarrel and harmony among siblings. J Cell Sci 117:5965–

5973.

22.   Flavahan WA, Gaskell E, Bernstein BE. 2017. Epigenetic plasticity and the hallmarks of

cancer. Science 357:eaal2380.

23.   Chung L, Thiele Orberg E, Geis AL, Chan JL, Fu K, DeStefano Shields CE, Dejea CM, Fathi P,

Chen J, Finard BB, Tam AJ, McAllister F, Fan H, Wu X, Ganguly S, Lebid A, Metz P, Van

Meerbeke SW, Huso DL, Wick EC, Pardoll DM, Wan F, Wu S, Sears CL, Housseau F. 2018.

Bacteroides fragilis Toxin Coordinates a Pro-carcinogenic Inflammatory Cascade via

Targeting of Colonic Epithelial Cells. Cell Host Microbe 23:203–214.e5.

24.   Dejea CM, Fathi P, Craig JM, Boleij A, Taddese R, Geis AL, Wu X, DeStefano Shields CE,

Hechenbleikner EM, Huso DL, Anders RA, Giardiello FM, Wick EC, Wang H, Wu S, Pardoll

DM, Housseau F, Sears CL. 2018. Patients with familial adenomatous polyposis harbor

colonic biofilms containing tumorigenic bacteria. Science 359:592–597.

25.   Arthur JC, Perez-Chanona E, Mühlbauer M, Tomkovich S, Uronis JM, Fan T-J, Campbell BJ,

Abujamel T, Dogan B, Rogers AB, Rhodes JM, Stintzi A, Simpson KW, Hansen JJ, Keku TO,

Fodor A a, Jobin C. 2012. Intestinal inflammation targets cancer-inducing activity of the

microbiota. Science (80-) 338:120–3.

26.   Ackermann AM, Wang Z, Schug J, Naji A, Kaestner KH. 2016. Integration of ATAC-seq and

RNA-seq identifies human alpha cell and beta cell signature genes. Mol Metab 5:233–

244.

27.   Davie K, Jacobs J, Atkins M, Potier D, Christiaens V, Halder G, Aerts S. 2015. Discovery of

transcription factors and regulatory regions driving in vivo tumor development by ATAC-

seq and FAIRE-seq open chromatin profiling. PLoS Genet 11:e1004994.

28.   Prendergast JGD, Semple CAM. 2011. Widespread signatures of recent selection linked to

nucleosome positioning in the human lineage. Genome Res 21:1777–1787.

29.   Prendergast JG, Campbell H, Gilbert N, Dunlop MG, Bickmore WA, Semple CA. 2007.

Chromatin structure and evolution in the human genome. BMC Evol Biol 7:72.

30.   Chen X, Chen Z, Chen H, Su Z, Yang J, Lin F, Shi S, He X. 2012. Nucleosomes Suppress

Spontaneous Mutations Base-Specifically in Eukaryotes. Science (80-) 335:1235–1238.

31.   Rhee K-J, Wu S, Wu X, Huso DL, Karim B, Franco AA, Rabizadeh S, Golub JE, Mathews LE,

Shin J, Sartor RB, Golenbock D, Hamad AR, Gan CM, Housseau F, Sears CL. 2009.

Induction of Persistent Colitis by a Human Commensal, Enterotoxigenic Bacteroides

fragilis, in Wild-Type C57BL/6 Mice. Infect Immun 77:1708–1718.

32.   Wu S, Shin J, Zhang G, Cohen M, Franco A, Sears CL. 2006. The Bacteroides fragilis toxin

binds to a specific intestinal epithelial cell receptor. Infect Immun 74:5382–90.

33.   Wick EC, Rabizadeh S, Albesiano E, Wu X, Wu S, Chan J, Rhee K-J, Ortega G, Huso DL,

Pardoll D, Housseau F, Sears CL. 2014. Stat3 Activation in Murine Colitis Induced by

Enterotoxigenic Bacteroides fragilis. Inflamm Bowel Dis 20:821–834.

34.   Buenrostro JD, Wu B, Chang HY, Greenleaf WJ. 2015. ATAC-seq: A method for assaying

chromatin accessibility genome-wide. Curr Protoc Mol Biol 2015:21.29.1-21.29.9.

35.   Lee J, Christoforo G, Christoforo G, Foo C, Probert C, Kundaje A, Boley N, kohpangwei,

Dacre M, Kim D. 2016. kundajelab/atac_dnase_pipelines: 0.3.3.

36.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods

9:357–359.

37.   Feng J, Liu T, Zhang Y. 2011. Using MACS to identify peaks from ChiP-seq data. Curr

Protoc Bioinforma 1–14.

38.   Zhu LJ, Gazin C, Lawson ND, Pagès H, Lin SM, Lapointe DS, Green MR. 2010.

ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip data. BMC

Bioinformatics 11:237.

39.   Pinello L, Xu J, Orkin SH, Yuan G-C. 2014. Analysis of chromatin-state plasticity identifies

cell-type-specific regulators of H3K27me3 patterns. Proc Natl Acad Sci 111:E344–E353.

40.   Haider S, Waggott D, Lalonde E, Fung C, Liu F-F, Boutros PC. 2016. A bedr way of genomic

interval processing. Source Code Biol Med 11:14.

41.   Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq

quantification. Nat Biotechnol 34:525–527.

42.   Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. 2017. Differential analysis of RNA-

seq incorporating quantification uncertainty. Nat Methods 14:687–690.

Figure Legends

Figure 1: **BFT2 induces dynamic changes in chromatin accessibility.** A.) HT29/C1 cells were

treated with BFT2 or left untreated for 24 or 48 hours. After each time point, DNA and RNA

were isolated. DNA was used to perform ATAC-seq, a measure of genome-wide chromatin

accessibility. RNA was used to perform RNA-seq, a measure of genome-wide gene expression.

B.) Venn diagrams showing the overlap of areas of accessible chromatin in untreated cells

(baseline) and BFT2-treated cells at 24 hours (top) and 48 hours (bottom). Areas of accessible

chromatin detected after BFT2 treatment at 24 or 48 hours, but not in untreated cells, were

categorized as "BFT2-opened" while areas of accessible chromatin detected in untreated cells,

but not after BFT2 treatment, were categorized as "BFT2- closed." Left and right Venn diagrams

show the overlap of BFT2-opened (left) or BFT2-closed (right) regions of the genome at 24 and

48 hours. For all Venn diagrams, the number of open peaks (chromatin accessible) detected and

the percentage of those peaks in each sector of the Venn diagrams are shown. Also, labels

(bold) at the top of each Venn diagram describe the entire circle, both overlapping and non-

overlapping regions, for each experimental condition.

Figure 2: **The effects of BFT2 on gene expression are more pronounced 24 hours after**

treatment than 48 hours. RNA-seq was performed on untreated cells and those treated with

BFT2 in order to determine if BFT2 alters gene expression in HT29/C1 cells. A) Heatmap

showing the change in gene expression at 24 hours after BFT2 treatment for those genes with

P-values < 0.01. B) Heatmap showing the changes in gene expression at 48 hours after BFT2

treatment for those genes with P-values < 0.01. For both heatmaps, colors represent Log2FC

(fold change) values of gene expression in BFT2-treated samples compared to gene expression

in baseline samples. Full list of differentially expressed genes after BFT2 treatment are in

supplementary tables 1 and 2. C.) Dotplot showing the output from a gene set enrichment

analysis looking for gene ontology (GO) and reactome gene sets that are upregulated or

downregulated at 24 or 48 hours after BFT2 treatment. Only gene sets with an FDR qval < 0.25

are shown.

Figure 3: **Chromatin accessibility and gene expression correlate at baseline and 24 hours after**

BFT2 treatment. A.) Stacked histogram showing the correlation between gene expression and

chromatin accessibility at baseline. Genes were first sorted by the presence of a peak in their

promoter in the 24hr baseline sample. Genes with a promoter peak were considered "open"

while those without a peak were considered "closed." Log transcripts per million (TPM) gene

expression values were then extracted and compared. The P-value was calculated by comparing

the gene expression of genes with a promoter peak to those without a promoter peak in the

24hr baseline sample. B-C) MA plot analyzing gene expression change after BFT2 treatment at

24 and 48 hours. Genes are colored according to the presence of a BFT2-opened peak or a

BFT2-closed peak in their promoter region. The dotted lines represent a log2 fold change in

expression of +/- O.5. The P-values were calculated by comparing the fold change in gene

expression for genes with a promoter BFT2-opened peak to those with a promoter BFT2-closed

peak. For all graphs, P-values were calculated using the Wilcoxon rank sum test.

Materials and Methods

Culture of HT-29/C1 cells: HT29/C1 cells were cultured at 37°C and 10% CO2. Cells were plated

at 20% confluency 4 days before addition of BFT2 toxin. At day 0, HT29/C1 cells were washed

five times with PBS, placed in minimal media (DMEM) without FBS or Pen/Strep and toxin was

added one time at a concentration of 100ng/ml. After toxin addition, cells were allowed to

grow at 37°C and 10% CO2 for 24 or 48 hours.

ATAC-seq transposition reaction: For all experiments, cell counts were obtained by trypsinizing

cells in 0.5% trypsin for 10 minutes at 37°C and then counting using a hemocytometer. Non-

trypsinized cells were then scraped and 50,000 cells were added to an Eppendorf tube. ATAC-

seq was performed using the protocol outlined by Buenrostro et al. (2015). Briefly, cells were

washed with 50uL cold 1x PBS buffer, then centrifuged at 500g for 5 minutes at 4°C. Cells were

then resuspended in 50uL cold lysis buffer (10 mM Tris-HCl [pH 7.4], 10 mM NaCl, 3 mM $MgCl_2$,

0.1% IGEPAL CA-630), gently lysed to preserve cell nuclei, and centrifuged again at 500g for 10

minutes at 4°C. Cells were then washed 3x with wash buffer (lysis buffer without igepal). After

each wash, cells were centrifuged at 500g for 5 minutes at 4°C. Cell nuclei were then

resuspended in the transposition reaction mix and incubated for 30 minutes at 37°C. DNA was

purified using a Qiagen MinElute PCR Purification Kit and eluted in 10uL elution buffer. DNA

was stored at -20°C until fragments were amplified via PCR.

PCR amplification following transposition: All of the DNA purified following transposition was

PCR amplified. To do so, the following were combined in a 0.2mL PCR tube: 10uL transposed

DNA, 10uL nuclease-free H2O, 2.5uL 25uM custom Nextera PCR primer 1, 2.5uL 25uM custom

Nextera barcoded PCR primer 2, 25uL NEBNext high-fidelity 2x PCR master mix. The

components were amplified as follows: 1 cycle of 72°C for 5 min, 98°C for 30 sec; 5 cycles of

98°C for 10 sec, 63°C for 30 sec, 72°C for 1 min. After initial amplification, the number of

additional cycles to run was determined using qPCR. For this, the following were combined in a

0.2mL PCR tube: 5uL of previously PCR-amplified DNA, 4.5uL of nuclease-free H2O, 0.25uL of

25uM custom Nextera PCR primer 1, 0.25uL of 25uM custom Nextera PCR primer 2, 5uL KAPA

SYBR FAST qPCR master mix (2x) (total reaction 15 ul). The components were amplified as

follows: 1 cycle of 98°C for 30 sec; 20 cycles of 98°C for 10 sec, 63°C for 30 sec, 72°C for 1 min.

To calculate the number of additional cycles of PCR needed, linear Rn versus cycle was plotted

and the cycle number that corresponds with 1/3 of the maximum fluorescent intensity was

determined. After the number of additional PCR cycles was determined, the remaining 45uL of

PCR product was run as follows: 1 cycle of 98°C for 30 sec; N cycles (as determined via qPCR) of

98°C for 10 sec 63°C for 30 sec, 72°C for 1 min. Finally, the amplified library was purified using

0.9x Ampure beads at room temperature, and eluted in 20uL RNAse-free H2O.

ATAC-seq data analysis: The 4nM pooled ATAC-seq library was sequenced on an Illumina HiSeq

using 50bp paired-end sequencing. For each sample, three biological replicates were

sequenced. After sequencing, the data was analyzed using the pipeline developed by the

Kundaje lab, as outlined on ENCODE (35). Briefly, reads were trimmed, aligned and filtered

using Bowtie2(36) and peaks were called using MACS2(37). The three replicate peak files were

combined in order to create one consensus file for each treatment condition using an

irreproducible discovery rate (IDR) threshold of 0.1. For all analyses, the "optimal set"

consensus file was used. Peak files for samples before and after BFT2 treatment were then

compared in order to create BFT2-opened peaks (peaks present after BFT2 treatment, but not

before) and BFT2-closed peak (peaks present before BFT2 treatment, but not after) files at 24

and 48 hours after treatment. The ChIPpeakAnno package was used to separate peaks into

specific regions of the genome (promoter, coding, intergenic) (38). The ChIPpeakAnno package

was also used to associate peaks located in promoter regions with their nearest downstream

genes. The haystack_bio package was used to identify transcription factor binding motifs (39).

For this analysis, C-G correction was turned off. For both the 24 hours and 48 hours analyses,

the BFT2-treated peak file was compared to the baseline peak file. A ratio greater than 1

represents transcription factor binding motifs that are present more frequently in the BFT2-

treated peak file, while a ratio of less than 1 represents transcription factor binding motifs that

are present more frequently in the baseline peak file. The bedR package was used to calculate

the jaccard index for the overlap between identified peaks in our dataset and CTCF binding

peak data in Caco-2 cells, taken from the UCSC genome browser (40).

RNA-seq assay: Cells were washed once with PBS, collected using a cell scraper, and RNA was

extracted from cells using the Qiagen RNeasy mini kit. After RNA extraction, the RNA pellet was

flash frozen using 100% ethanol and dry ice. Samples were then stored at -80ºC. For library

preparation, samples were removed from storage, and mRNA was enriched using the NEBNext

Poly(A) mRNA Magnetic Isolation Module. Afterwards, a non-directional RNA-seq library was

constructed using the NEBnext Ultra RNA Library Prep kit from Illumina. The 2nM pooled RNA-

seq library was sequenced using the Illumina HiSeq. For one sample, 50bp paired-end

sequencing was performed. For the other two samples, 50bp single-end sequencing was

performed.

RNA-seq data analysis: After sequencing, Kallisto was used to perform pseudoalignment of the

raw RNA-seq data (41). Then, Sleuth was used to quantify gene expression and perform

differential expression analyses (42).

Gene Set Enrichment Analysis: Gene set enrichment analysis was performed using all genes

with a P-value < 0.05 at 24 or 48 hours after BFT2 treatment. For this analysis, a ranklist was

first created by sorting the differentially expressed genes using the following formula: sign of

fold change $\times$ inverse P-value. This created a rank ordered list with the most significantly

upregulated genes at the top and the most significantly downregulated genes at the bottom.

The GSEA Pre-ranked tool was then run, using 1000 permutations, a classic (instead of

weighted) analysis, a minimum gene set size of 15 and a maximum gene set size of 1500. The

output of this analysis was then converted into graphical and tabular formats.

Statistics: To calculate the correlation between chromatin accessibility (ATAC-seq) and gene

expression (RNA-seq) at baseline and after BFT2 treatment, the Wilcoxon Rank Sum test was

used with an alpha of 0.05. To calculate the enrichment of transcription factor binding motifs in

BFT2-treated and untreated samples, a Fisher's exact test with an alpha of 0.01 was used. To

determine if there was a correlation between the location of BFT2-treated peaks and common

SNVs and DMRs in CRC, a chi-square test with an alpha of 0.05 was used in which BFT2-treated

peak files were compared to baseline peak files.

1   Table 1: Distribution of ATAC-seq peaks throughout the genome

	24hr Baseline (48,982)[a]	24hr BFT2 (69,415)	48hr Baseline (58,147)	48hr BFT2 (66,945)	24hr BFT2-Opened (24,967)	24hr BFT2-Closed (6,754)	48hr BFT2-Opened (17,459)	48hr BFT2-Closed (10,417)
Promoter (0.5%)[b]	41.6%	33.7%	37.9%	34.9%	17.4%	18.2%	16.0%	16.7%
Coding (44.3%)	26.8%	31.5%	28.7%	30.7%	40.2%	37.4%	41.1%	39.4%
Intergenic (55.0%)	31.6%	34.8%	33.4%	34.4%	42.4%	44.4%	42.9%	43.9%

The coding region includes 5' UTR, introns and exons
[a] Number of peaks per sample in parentheses
[b] Percent of the human genome represented by each region in parentheses

1   Table 2: Transcription factor binding motifs enriched after cell treatment with BFT2 for 24 hours

Motif Name	Presence in target	Presence in background	Ratio	P-value
JUNB	12.68%	10.17%	1.22	1.90E-40
FOS	12.40%	9.97%	1.22	8.14E-39
JUN(var.2)	12.89%	10.43%	1.22	1.71E-38
JUND	11.97%	9.59%	1.22	2.45E-38
FOSL1	12.50%	10.09%	1.22	1.27E-37
FOS::JUN	12.11%	9.75%	1.22	2.74E-37
JDP2	10.17%	8.01%	1.24	8.26E-37
BATF::JUN	10.91%	8.82%	1.21	2.48E-32
NFE2	10.47%	8.43%	1.22	7.61E-32
ZBTB33	12.90%	15.27%	0.85	4.76E-31
[a]CTCF	37.76%	36.92%	1.02	3.38E-03

Only top 10 motifs are shown. Full list in supplementary table 5
P-values calculated using Fisher's exact test
Ratio < 1.0 represents enrichment in baseline sample
[a] included due to physiological relevance

1 Table 3: Transcription factor binding motifs enriched after cell treatment with BFT2 for 48 hours

Motif Name	Presence in target	Presence in background	Ratio	P-value
ZBTB33	13.59%	14.79%	0.92	1.84E-09
FOS	11.26%	10.26%	1.09	1.49E-08
JUNB	11.43%	10.43%	1.09	1.51E-08
JUND	10.78%	9.82%	1.09	2.32E-08
FOS::JUN	10.95%	10.01%	1.09	7.19E-08
JDP2	9.20%	8.34%	1.09	7.80E-08
FOSL1	11.27%	10.35%	1.08	1.60E-07
E2F4	21.39%	22.60%	0.95	2.60E-07
JUN(var.2)	11.70%	10.82%	1.07	1.06E-06
NFE2	9.60%	8.80%	1.08	1.21E-06

Only top 10 motifs are shown. Full list in supplementary table 5
P-values calculated using Fisher's exact test
Ratio < 1.0 represents enrichment in baseline sample

Table 4: Association between BFT2-induced chromatin accessibility changes and common Single
Nucleotide Variants (SNVs)/Differentially Methylated Regions (DMRs) in CRC

	Number of overlapping peaks	Number of non-overlapping peaks	Total number of peaks	Odds[a]	Odds ratio[b] (95% CI[c])
SNVs					
24hr BFT2	11063	58352	69415	.0281	0.818 (0.794-0.844)
24hr Baseline	9210	39772	48982	.0271	
48hr BFT2	11147	55798	66945	.0259	0.964 (0.936-0.993)
48hr Baseline	9981	48166	58147	.0237	
DMRs					
24hr BFT2	1900	67515	69415	.0281	1.037 (0.965-1.114)
24hr Baseline	1294	47688	48982	.0271	
48hr BFT2	1692	65253	66945	.0259	1.094 (1.018-1.177)
48hr Baseline	1346	56801	58147	.0237	
DMRs hypermethylated					
24hr BFT2	944	68471	69415	.0138	0.979 (0.887-1.082)
24hr Baseline	680	48302	48982	.0141	
48hr BFT2	953	65992	66945	.0144	1.101 (1.000-1.212)
48hr Baseline	753	57394	58147	.0131	
DMRs hypomethylated					
24hr BFT2	993	68422	69415	.0145	1.112 (1.006-1.230)
24hr Baseline	631	48351	48982	.0131	
48hr BFT2	757	66188	66945	.0114	1.081 (0.971-1.203)
48hr Baseline	609	57538	58147	.0106	

[a] Odds represent the number of overlapping peaks divided by the number of non-overlapping peaks
[b] Odds ratios represent the odds in BFT2-treated cells divided by the odds in untreated cells
[c] 95% confidence intervals were calculated using a chi-square test
Common DMRs and SNVs in CRC were extracted from the COSMIC database