The relationship between transmission time and clustering methods in

***Mycobacterium tuberculosis* epidemiology**

Conor J Meehan[1,*], Pieter Moris[1,2,3], Thomas A. Kohl[4,5], Jūlija Pečerska[6], Suriya Akter[1],

Matthias Merker [4,5], Christian Utpatel[4,5], Patrick Beckert[4,5], Florian Gehre[1,7,8], Pauline

Lempens[1], Tanja Stadler[6], Michel K. Kaswa[1,10], Denise Kühnert[9], Stefan Niemann[4,5,&], Bouke

C de Jong[1,&]

Affiliations

[1]Unit of Mycobacteriology, Biomedical Sciences, Institute of Tropical Medicine, Antwerp,

2000, Belgium

[2]Advanced Database Research and Modelling (ADReM), Department of Mathematics and

Computer Science, University of Antwerp, Antwerp, 2020, Belgium

[3]Biomedical Informatics Research Network Antwerp (biomina), University of Antwerp,

Antwerp, 2020, Belgium

[4]German Center for Infection Research, Partner Site Hamburg-Lübeck-Borstel-Riems, D-

23845 Borstel, Germany

[5]Molecular and Experimental Mycobacteriology, Priority Area Infections, Research Center

Borstel, D-23845 Borstel, Germany

[6]Department of Biosystems Science and Engineering, ETH Zürich, 4058 Basel, Switzerland

[7]Vaccines and Immunity Theme, Medical Research Council Unit The Gambia, Serekunda, The

Gambia

[8]Bernhard Nocht Institute for Tropical Medicine, 20359 Hamburg, Germany

[9]University Hospital Zurich, 8091 Zurich, Switzerland

[10]National Tuberculosis Program, B.P 1197, Kinshasa, Democratic Republic of Congo

&equal contribution; *corresponding author: cmeehan@itg.be; +32-32476318

Keywords

Abstract

Background: Tracking recent transmission is a vital part of controlling widespread

pathogens such as *Mycobacterium tuberculosis*. Multiple methods with specific

performance characteristics exist for detecting recent transmission chains, usually by

clustering strains based on genotype similarities. With such a large variety of methods

available, informed selection of an appropriate approach for determining transmissions

within a given setting/time period is difficult.

Methods: This study combines whole genome sequence (WGS) data derived from 324

isolates collected 2005-2010 in Kinshasa, Democratic Republic of Congo (DRC), a high

endemic setting, with phylodynamics to unveil the timing of transmission events posited by

a variety of standard genotyping methods. Clustering data based on Spoligotyping, 24-loci

MIRU-VNTR typing, WGS based SNP (Single Nucleotide Polymorphism) and core genome

multi locus sequence typing (cgMLST) typing were evaluated.

Findings: Our results suggest that clusters based on Spoligotyping could encompass

transmission events that occurred over 70 years prior to sampling while 24-loci-MIRU-VNTR

often represented two or more decades of transmission. Instead, WGS based genotyping

applying low SNP or cgMLST allele thresholds allows for determination of recent

transmission events in timespans of up to 10 years e.g. for a 5 SNP/allele cut-off.

Interpretation: With the rapid uptake of WGS methods in surveillance and outbreak

tracking, the findings obtained in this study can guide the selection of appropriate clustering

methods for uncovering relevant transmission chains within a given time-period. For high

resolution cluster analyses, WGS-SNP and cgMLST based analyses have similar

clustering/timing characteristics even for data obtained from a high incidence setting.

Research in context

<u>Evidence before this study</u>

For nearly 30 years, molecular genotyping tools have been used to define transmission

chains/clusters of *Mycobacterium tuberculosis* strains. A variety of tools are used for such

analysis e.g. the presence/absence of spacers sequences (Spoligotyping), the length of

tandem repeat patterns (24-loci-MIRU-VNTR) or, more recently, nearly the complete

genome by whole genome sequencing (WGS). Each method has been proposed as the gold

standard genotyping technique for detecting transmission events in a certain timeframe and

selection of the optimal method for a given question is difficult as important parameters

(e.g. the time span a particular outbreak can encompass) are not well defined. Based on

inferred mutation rates, there have been some time scales proposed for clusters based on

WGS SNP-based methods, supported by contact tracing data to confirm epidemiological

links. However, there is uncertainty around these timing estimates for SNP-based

techniques, limited timing estimates available for classical genotyping techniques and no

such estimates for cgMLST approaches. This makes it very difficult for researchers, public

health workers and clinicians to correctly interpret reported clustering data. This is

especially the case as WGS based methods are becoming rapidly ingrained in surveillance

and clinical workflows.

<u>Added value of this study</u>

This study is the first to perform a comparative evaluation of cluster data defined by both

classical and WGS-based *M. tuberculosis* genotyping approaches, especially with regard to

transmission timing. While many studies have put forward various methods as the gold

standard for *M. tuberculosis* transmission detection, we have tested clustering data

generated by the different methods in a Bayesian statistical framework to elucidate the true

fraction of recent transmission each approach is detecting. When specifically looking at

recent transmission (e.g. less than 10 years previous), our results indicate that classical

genotyping methods vastly over estimate recent transmission events. This solidifies the

need for WGS-based methods when searching for recent outbreaks of *M. tuberculosis*.

<u>Implications of all the available evidence</u>

Our study allows researchers and public health officials to select the appropriate genotyping

method for assessing transmission with respect to the epidemiological setting and a given

time-period. We also suggest the incorporation of particular genotyping methods in a

cascade system with increasing resolution for various levels of surveillance e.g. from multi-

country surveillance down to recent transmission and outbreak analyses. This is particularly

important as each method comes with specific costs, infrastructure and computational

requirements, human resources, and, last but not least interpretation complexities – all of

which might not be feasible at any site or at any scale. Accordingly, our study can aid a

cost/benefit analysis for selection of genotyping techniques in a cascade system, that might

especially be used in high incidence, low resource settings.

Introduction

Despite the large global efforts at curbing the spread of *Mycobacterium tuberculosis*

complex (Mtbc) strains, 10.4 million new patients develop tuberculosis (TB) every year[1]. In

addition, the prevalence of multidrug resistant (MDR) Mtbc strains is increasing[1],

predominantly through ongoing transmission within large populations[2,3]. The tracking and

timing of recent transmission chains allows TB control programs to effectively pinpoint

transmission hotspots and employ targeted intervention measures. This is especially

important for the transmission of drug resistant strains as it appears that drug resistance

may be transmitted more frequently than acquired[2]. Thus, interrupting transmission is key

for the control of MDR-TB[3,4]. For the development of the most effective control strategies,

there is a strong need for (i) appropriate identification of relevant transmission chains, risk

factors and hotspots and (ii) robust timing of when outbreaks first arose.

Epidemiological TB studies often apply genotyping methods to Mtbc strains to determine

whether two or more patients are linked within a transmission chain (molecular

epidemiology)[5]. Contact tracing is the primary non-molecular epidemiological method for

investigating transmission networks of TB, mainly based on patient interviews[6]. Although

this method is often seen as a gold standard of transmission linking, it does not always

match the true transmission patterns, even in low incidence settings[7] and misses many

connections[8]. The implementation of molecular genotyping and epidemiological approaches

has overcome these limitations and is often used as the main approach for transmission

analyses. Classical genotyping has involved IS*6110* DNA fingerprinting[9], Spoligotyping[10], and

variable-number tandem repeats of mycobacterial interspersed repetitive units (MIRU-

VNTR)[11] which is the most common method at the moment[5]. The latter method is based on

copy numbers of a sequence in tandem repeat patterns derived from 24 distinct loci within

the genome[12]. If two patients have the same classical genotyping pattern such as a 24-loci

MIRU-VNTR pattern (or up to one locus difference[12]) they are considered to be within a

local transmission chain. The combination of spoligotyping and MIRU-VNTR-typing, where

patterns must match in both methods to be considered a transmission link, is often

considered the molecular gold standard for transmission linking and genotyping[12]. However,

examples of unlinked patients with identical patterns have been observed, suggesting that

this threshold covers too broad a genetic diversity and timespan between infections[7].

The application of (whole genome) sequence (WGS)-based approaches for similarity analysis

of Mtbc isolates and cluster determination is known to have high discriminatory power

when assessing transmission dynamics[7,13–16], either using core genome multi-locus

sequence typing (cgMLST)[17,18] or SNP distances[7,14,15,19]. WGS-based approaches compare the

genetic relatedness of the genomes of the clinical strains under consideration, albeit usually

excluding large repetitive portions of the genome, with the assumption that highly similar

strains are linked by a recent transmission event[7,14]. Although many SNP cut-offs for linking

isolates have been proposed[20], the most commonly employed is based on the finding that a

5 SNP cut-off will cluster the genomes of strains from the majority of epidemiologically

linked TB patients, with an upper bound of 12 SNPs between any two linked isolates[14]. The

emerging widespread use of WGS has quickly pushed these cut-offs to be considered the

new molecular gold standard of recent transmission linking, although SNP distances may

vary for technical reasons (e.g. assembly pipelines or filter criteria[21]) and between study

populations e.g. high and low incidence settings[19].

In addition to cluster detection, uncovering the timing of transmission events within a given

cluster is highly useful information for TB control e.g. for assessing the impact of

interventions on the spread of an outbreak or uncovering when MDR-TB transmission first

emerged in a particular setting. Accordingly, knowledge of the rate change associated with

different genotyping methods is essential for correct timing. The whole genome mutation

rate of Mtbc strains has been estimated by several studies as between 10^{-7} and 10^{-8}

substitutions per site per year or ~0·3-0·5 SNPs per genome per year[7,14,22–24], while the rate

of change in the MIRU-VNTR loci specifically is known to be quicker (~10^{-3})[25]. Since these

mutation rates have been shown to also vary by lineage[23,26] and over short periods of

time[22], such variation needs to be accounted for when estimating transmission times, e.g.

by using Bayesian phylogenetic dating techniques[3,22,25].

Considering the multiple genotyping methods currently available, many of them proposed

as a "gold standard", there is an urgent need to precisely define the individual capacity of

each method to accurately detect recent transmission events and perform timing of

outbreaks. To provide this essential information, this study harnesses the power of WGS-

based phylogenetic dating methods to assign timespans onto Mtbc transmission chains

encompassed by the different genotypic clustering methods commonly used in TB

transmission studies.

Materials and Methods

<u>Dataset, ethical approval and sequencing</u>

A set of 324 isolates from Kinshasa, Democratic Republic of Congo were collected from

consecutive retreatment TB patients between 2005 and 2010 at TB clinics, servicing an

estimated 30% of the population of Kinshasa. All isolates were phenotypically resistant to

rifampicin (RR-TB) and the majority are also isoniazid resistant (i.e. MDR-TB). Use of the

stored isolates without any linked personal information was approved by the health

authorities of the DRC and the Institutional Review Board of the ITM in Antwerp (ref no

945/14). Libraries for whole genome sequencing were prepared from extracted genomic

DNA with the Illumina Nextera XT kit, and run on the Illumina NextSeq platform in a

2x151bp run according to manufacturer's instructions. Illumina read sets are available on

the ENA (https://www.ebi.ac.uk/ena) under the accession number PRJEB27847.

Genome reconstruction

The MTBseq pipeline[27] was used to detect the SNPs for each isolate using the H37Rv

reference genome (NCBI accession number NC000962.3)[28]. Sites known to be involved in

drug resistance (as outlined in the PhyResSE list of drug mutations v27[29]) were excluded

from the alignment and additional filtering of sites with ambiguous calls in >5% of isolates

and those SNPs within a 12bp window of each other was also applied.

Transmission cluster estimation methods

Five standard transmission clustering approaches were chosen for comparison and analysis

(four additional methods are outlined in the supplementary methods). Clustering due to

convergent genotypic patterns were removed as outlined in the supplementary methods.

For each method, the total SNP distances were calculated to investigate the range of

variability encompassed within each cluster. Maximum SNP distances were derived from

pairwise comparisons of isolates within the SNP alignment using custom python scripts. A

clustering rate was calculated for each method using the formula $(n_c - c)/n$, where n_c is the

total number of isolates clustered by a given method, c is the number of clusters, and n is

the total number of isolates in the dataset (n=324).

Spoligotyping and MIRU-VNTR

Spoligotype patterns were obtained from membranes following the previously published

protocol[10]. Isolates were said to be clustered if all 43 spacers matched. Genotyping by

MIRU-VNTR was undertaken as previously described[12]. 2µl of DNA was extracted from

cultures and amplified using the 24 loci MIRU-VNTR typing kit (Genoscreen, Lille, France).

Analysis of patterns was undertaken using the ABI 3500 automatic sequencer (Applied

Biosystems, California, USA) and Genemapper software (Applied Biosystems). Isolates were

said to be clustered if all 24 loci matched. Mixed MIRU-VNTR patterns were observed in 18

isolates although this mixing was not observed in the WGS data, likely due to subculturing

for sequencing.

<u>SNP cut-off clustering</u>

In this study, we employed the widely used 5 SNP (proposed by Walker *et al.* [14] as the likely

boundary for linked transmission) and 12 SNP cut-offs (proposed maximum boundary) for

cluster definition. Additionally, we employed a lower cut-off of 1 SNP to look for clusters of

very highly related isolates. Pairwise SNP distances were calculated between all isolates. A

loose cluster definition was used, where every isolate in a cluster at most the SNP cut-off

from at least 1 other isolate in the cluster.

<u>Estimation of transmission times</u>

To estimate the age and timespan of potential transmission clusters, SNP alignments were

created from the convergence-free version of the three primary clustering types:

Spoligotyping, MIRU-VNTR, and 12 SNP clusters.

A Bayesian approach to transmission time estimation was then undertaken. The SNP

alignments were created as above for the three high-level clustering types. Each cluster

method alignment was separately input to BEAST-2 v2.4.7[30] to create a time tree for those

isolates. These phylogenies were built using the following priors: GTR+GAMMA substitution

model, a log-normal relaxed molecular clock model to account for variation in mutation

rates[31] and coalescent constant size demographic model[32], both of which have been found

to be suitable for lineage 4 isolates in a previous study[19]. The MCMC chain was run six times

independently per alignment with a length of at least 400 million, sampled every 40,000[th]

step (MIRU & Spoligotyping: 600 million; 12 SNP: 500 million;). A log normal prior (mean

1.5×10^{-7}; variance 1·0) was used for the clock model to reflect the previously estimated

mutation rate of *M. tuberculosis* lineage 4[7,14,22–24], while allowing for variation as previously

suggested[22]. A 1/X non-informative prior was selected for the population size parameter of

the demographic model. Isolation dates were used as informative heterochronous tip dates

and the SNP alignment was augmented with a count of invariant sites for each of the four

nucleotide bases to avoid ascertainment bias[33]. Tracer v1.6 was used to determine

adequate mixing and convergence of chains (ESS >150) after a 25% burn-in. The chains were

combined via LogCombiner v2.4.8[30] to obtain a single chain for each clustering type with

high (>1000) effective sample sizes. The tree samples were combined in the same manner

and resampled at a lower frequency to create thinned samples of (minimum) 20,000 trees.

The algorithm for estimating the timespan of transmission events encompassed by each

method is outlined in Supplemental Figure 1. Briefly, for each cluster created by the given

method, we defined the MRCA node as the internal node that connects all taxa in that

cluster. The youngest node was then defined as the tip that is furthest from this MRCA

within the clade (i.e. the tip descendant from that node that was sampled closest to the

present time). To better account for changes in the mutation rate over short periods[22], all

trees estimated and sampled during the Bayesian MCMC process were used instead of only

a single summary phylogeny. For each retained tree in the MCMC process, the difference in

age between the MRCA node and youngest node was calculated. This gave a distribution of

likely maximum transmission event times within that cluster. For each method, these per-

cluster aggregated ages were then combined across all clusters to give a per-method

distribution of transmission event times represented by the clusters. The 95% Highest

Posterior Density (HPD) interval of these distributions was calculated with the

LaplacesDemon p.interval function in R v3.4.0 and the distribution within this interval for

each method along with the mean based upon this interval were then visualized in violin

plots per clustering method using ggplot2 in R.

Results

In this study, we assessed five different approaches for generating putative *M. tuberculosis*

transmission clusters: Spoligotyping, MIRU-VNTR, and SNP-based clustering using a 12, 5

and 1 SNP cut-off, using a dataset of 324 isolates collected 2005-2010 in Kinshasa,

Democratic Republic of Congo (DRC). The dataset contained 309 L4 and 15 L5 isolates, with

a maximum of 1671 SNPs between any two isolates. Bayesian phylodynamic dating

approaches implemented in BEAST-2[30] were then utilised to assign timespans to the

transmission events estimated by each genotyping method.

As expected, classical genotyping methods clustered many strains, with the lowest

resolution (i.e. highest clustering rate) (Figure 1, Table 1). Convergent evolution (defined as

the same pattern observed in unrelated strains; see supplemental methods) was found to

affect 39% (12) of Spoligotyping-based clusters. Although MIRU-VNTR performed far better

than spoligotyping, 16% (6) of clustering patterns were influenced by convergence in this

study (Table 1, Figure 1). Spoligotyping and MIRU-VNTR patterns that were affected by

convergence were removed from further timespan analyses. WGS-based methods had by

far the highest discriminatory power and low SNP cut-offs grouped isolates into smaller

clusters (e.g. 2-10 isolates per cluster for a 5 SNP cut-off) (Table 1, Figure 1). The high

percentage of strains in a 12 SNP cluster (75%) suggests high levels of transmission in this

population, making is suitable for further transmission analyses.

Bayesian phylogenetic dating of the timeframe associated with particular transmission

chains showed large differences in estimated cluster ages between the different genotyping

approaches used (Table 1, Figure 2), correlating well with the difference in discriminatory

power. Cluster ages are defined here as the most ancient transmission event that links any

two isolates within a specific cluster (see methods and supplementary figure 1). Thus, in

phylogenetic terms, the cluster age is the difference in time between when the most recent

common ancestor (MCRA) of the entire cluster existed and the date of isolation of the

furthest isolate from this ancestor.

The aggregate median ages of clusters derived from Spoligotyping were found to often be

several decades old (median 77 years (95% HPD: 1-823)) (Table 1, Figure 2). MIRU-VNTR

clustering encompassed more recent transmission events than Spoligotyping, but were still

found to be often over 2 decades old (median 26 years (95% HPD: 0-162)). The combination

of MIRU-VNTR and Spoligotyping resulted in cluster ages similar to MIRU-VNTR alone

(supplemental table 1). Clusters based on SNP cut-offs correlated to 23 years using a 12

SNP cut-off (95% HPD: 0-103), 11 years using a 5 SNP cut-off (95% HPD: 0-47), and 4 years

using a 1 SNP cut-off (95% HPD: 0-24) (Table 1, Figure 2). For comparison, the cgMLST

method was also performed and discriminatory power, cluster sizes and ages based on

cgMLST alleles were similar to the SNP-based clusters (see supplemental methods,

supplemental table 1).

Discussion

The term 'recent transmission' is often applied to gain a better understanding of the current

transmission dynamics of pathogens in a given population. However, little data is available

on how recent a likely transmission event occurred when measured with different

genotyping methods. To get a better understanding of the discriminatory power of different

classical genotyping techniques and WGS-based approaches in relation to outbreak timing,

this study has performed an in-depth comparison of clustering rates and dated phylogenies

obtained in a collection of 324 Mtbc strains from a high incidence setting (Kinshasa, DRC).

With a whole genome phylodynamic approach employed as a gold standard, our study

demonstrates that each genotyping method was associated with a specific discriminatory

power resulting in clusters representing vastly different time periods of transmission events

(Table 1 and Figure 2). This has significant implications for data interpretations e.g. when

selecting and utilising different genotyping/clustering approaches for epidemiological

studies and assessing the effectiveness of public health intervention strategies.

As the most extreme example, Spoligotyping-derived clusters were associated with

transmission events that can be several decades old. This low discriminatory power coupled

with the high rate of convergent evolution (the same spoligotype pattern found in

phylogenetically distant isolates) adds weight to the previous suggestion that this technique

is not suitable for recent transmission studies[34], although may be of use as a low-cost

method of sorting Mtbc strains into the seven primary lineages[35,36].

In line with previous findings[34,37], convergent evolution of 24-loci MIRU-VNTR patterns was

rarer than observed for Spoligotyping, but did occur in 16% of MIRU-VNTR-based clusters.

Additionally, the transmission times encompassed by MIRU-VNTR clusters often spanned

over two decades (Table 1, Figure 2), confirming previous studies showing over-estimation

of recent transmission with this method[7,13,19,38].

For defining transmission events that occurred in more recent time frames before sampling,

WGS-based methods were found to be better suited than classical genotyping methods

(Table 1, Figure 2). The 12 SNP cut-off, currently the recommended upper bound for

clustering isolates, often defines transmission events that occurred on average two decades

prior to sampling, similar in age to clusters estimated by MIRU-VNTR. This suggests that the

12 SNP cluster method may be a good replacement for MIRU-VNTR as it detects larger

transmission networks spanning similar transmission time periods but is less affected by

convergent evolution. Isolates clustered at a low (5 SNP) or nearly identical (1 SNP) cut-off

were found to represent transmission events occurring over a time span of up to ten years.

These findings correlate well with previous studies where confirmed contact tracing-based

epidemiological links were found between patients that were two[15,39] and three[7] SNPs

apart. The original paper that proposed the 5 and 12 SNP cut-offs found that serial isolates

that were 10 years apart differed by, on average, 6 SNPs, also agreeing with the findings

presented here [14]. Comparisons between the SNP-based (using almost all genomic

differences) and the cgMLST-based (using a defined core set of genes) methods

demonstrated that the latter approach gives similar estimates to full SNP approaches. This

supports the use of low SNP or cgMLST differences for detection or exclusion of very recent

transmission, although this low variability between isolates makes robust identification of

transmission direction difficult, especially during short timespans.

The mutation rate of *M. tuberculosis* has been estimated to be between 10^{-7} and 10^{-8}

substitutions per site per year[3,7,23]. Within the Bayesian analysis employed here, the

mutation rate was free to vary between these values but was found to strongly favour

~3×10^{-8} (ESS > 1000 for all runs; 95% HDP: 4×10^{-9} - 8×10^{-8}), translating to approximately 0·3

SNPs per genome per year. While the mutation rate used here is in line with previous

estimates for lineage 4 [23] (which most of this dataset is comprised of), it may be similar in

other lineages, although this has only been shown for lineage 2[3,23]. Thus, per-lineage

estimates are required for all seven lineages to ensure similar transmission times are linked

to genotyping methods across the whole diversity of the Mtbc.

While this study has many advantages due to its five year population based design in an

endemic setting coupled with the application of three different genotyping methods

(Spoligotyping, 24-locus MIRU-VNTR and WGS), future confirmatory studies could address

the following drawbacks that are inherent to genomic epidemiology[16,21]: 1) studies

employing contact tracing and/or digital epidemiology[40] in conjunction with these

genotyping methods can help confirm transmission times associated with different clusters

(although these methods also have many limitations); 2) as outlined above, strains of other

lineages of the Mtbc should be analysed in a similar fashion to ensure transferability of

findings across the entire complex; 3) a broad range of drug resistance profiles should be

included to fully assess the impact of such mutations on transmission estimates; 4)

improved WGS methods, such as directly from clinical samples to help reduce culture

biases[41] and longer reads (e.g. PacBio SMRT or Nanopore MinION) to capture the entire

genome, including repetitive regions such as PE/PPE genes known to impact genome

remodelling[42,43], will ensure that the maximum diversity between isolates is captured; 5)

extensive panels of Spoligotyping and MIRU-VNTR results paired with WGS data will help

assess the extent of convergence in these methods and better correlate their clusters with

those of low SNP thresholds and 6) standardised SNP calling pipelines appropriate across all

lineages, with high true positive/low false negative rates, will ensure that Mtbc molecular

epidemiology can be uniformly implemented and comparable across studies.

Since each method was found to represent different timespans and clustering definitions,

they can be used in a stratified manner in an integrated epidemiological and public health

investigation addressing the transmission of Mtbc strains. For instance, although

Spoligotyping clusters represented potentially very old transmission events, the low

associated cost and its ability to be applied directly on sputum helps reduce culture bias and

thus robustly assign lineages. Thus, Spoligotyping and/or MIRU-VNTR would serve well as

first-line surveillance of potential transmission events in the population, guiding further

investigations and resource allocations.

These potential transmission hotspots could be further investigated with contact tracing

and/or WGS. Employment of different cut-offs and clustering approaches to WGS data can

then address several questions. The 12 SNP/cgMLST allele cluster approaches serve well for

high level surveillance targeting larger (older) transmission networks, akin to what is

currently often done using MIRU-VNTR (e.g.[15,44]). Recent transmission events can then be

detected through employment of low SNP cut-offs (e.g. 5 SNPs for transmission in the past

10 years or 1 SNPs for transmission in the past 5 years). In high incidence/low diversity

settings where amalgamation of clusters may inadvertently obscure distinct hotspots of

transmission at different time points, subdivision into distinct time-dependant clusters can

be undertaken using the algorithm presented in such a study in East Greenland[19].

Overall, phylodynamic approaches applied to whole genome sequences, as undertaken

here, are recommended to fully investigate the specific transmission dynamics within a

study population to account for setting-specific conditions, such as low/high TB incidence,

low/high pathogen population diversity, sampling fractions and social factors influencing

transmission. As WGS methods become more commonplace and easier to implement in a

variety of settings, each genotyping method can be employed as part of an overall evidence

gathering program for transmission, placing molecular epidemiological approaches as an

integral part in tracking and stopping the spread of TB.

Acknowledgements

The authors would like to thank Armand Van Deun and Koen Vandelannoote for valuable

discussion and input and Cecile Uwizeye for aid with spoligotyping.

Funding sources

This work was supported by an ERC grant [INTERRUPTB; no. 311725] to BdJ, FG and CJM; an

ERC grant to TS [PhyPD; no. 335529]; an FWO PhD fellowship to PM [grant number

1141217N]; the German Centre for Infection Research (DZIF) for TAK, MM, CU, PB and SN; a

SNF SystemsX grant (TBX) to JP and TS and a Marie Heim-Vögtlin fellowship granted to DK

by the Swiss National Science Foundation. The computational resources and services used in

this work were provided by the VSC (Flemish Supercomputer Center), funded by the

Research Foundation - Flanders (FWO) and the Flemish Government – department EWI.

Declaration of interests

The authors declare there are no conflicts of interest attached to this work.

Author contributions

CJM, FG and BCdJ conceived the study. MKK and BCdJ oversaw collection of isolates and

ethical approval. TAK, SA, MM, PB and SN undertook classic genotyping and sequencing of

isolates. CJM, PM, TA, CU and PL undertook WGS assembly and data preparation. CJM

undertook all convergence and clustering analyses. CJM, PM, JP, MM, TS and DK undertook

all phylodynamics. CJM, PM, SN and BCdJ wrote the manuscript. All authors read and

revised the manuscript and approved its final form.

References

1.   WHO. Global tuberculosis report 2017 [Internet]. 2018. Available from:

      http://www.who.int/tb/publications/global_report/archive/en/

2.   Kendall EA, Fofana MO, Dowdy DDW, WHO, Dye C, Garnett G, et al. Burden of

      transmitted multidrug resistance in epidemics of tuberculosis: a transmission

      modelling analysis. Lancet Respir Med [Internet]. 2015 Nov 12 [cited 2015 Nov

      23];3(12):963–72. Available from:

      http://linkinghub.elsevier.com/retrieve/pii/S2213260015004580

3.   Merker M, Blin C, Mona S, Duforet-Frebourg N, Lecher S, Willery E, et al. Evolutionary

      history and global spread of the Mycobacterium tuberculosis Beijing lineage. Nat

      Genet [Internet]. 2015 Jan 19 [cited 2017 Sep 22];47(3):242–9. Available from:

      http://www.nature.com/doifinder/10.1038/ng.3195

4.   Shah NS, Auld SC, Brust JCM, Mathema B, Ismail N, Moodley P, et al. Transmission of

      Extensively Drug-Resistant Tuberculosis in South Africa. N Engl J Med [Internet]. 2017

      Jan 19 [cited 2017 Jul 24];376(3):243–53. Available from:

      http://www.nejm.org/doi/10.1056/NEJMoa1604544

5.   Merker M, Kohl TA, Niemann S, Supply P. The Evolution of Strain Typing in the

      Mycobacterium tuberculosis Complex. In: Advances in experimental medicine and

      biology [Internet]. 2017 [cited 2018 Apr 10]. p. 43–78. Available from:

      http://www.ncbi.nlm.nih.gov/pubmed/29116629

6.   Fox GJ, Barry SE, Britton WJ, Marks GB. Contact investigation for tuberculosis: a

      systematic review and meta-analysis. Eur Respir J [Internet]. 2012 [cited 2017 Sep

      6];41(1). Available from: http://erj.ersjournals.com/content/41/1/140.short

7.   Roetzer A, Diel R, Kohl TA, Rückert C, Nübel U, Blom J, et al. Whole genome

sequencing versus traditional genotyping for investigation of a Mycobacterium

tuberculosis outbreak: a longitudinal molecular epidemiological study. Neyrolles O,

editor. PLoS Med [Internet]. 2013 Jan 12 [cited 2014 Mar 31];10(2):e1001387.

Available from: http://dx.plos.org/10.1371/journal.pmed.1001387

8.   Bjorn-Mortensen K, Lillebaek T, Koch A, Soborg B, Ladefoged K, Sørensen HCF, et al.

Extent of transmission captured by contact tracing in a tuberculosis high endemic

setting. Eur Respir J [Internet]. 2017 [cited 2017 Sep 13];49(3). Available from:

http://erj.ersjournals.com/content/49/3/1601851

9.   Thierry D, Cave MD, Eisenach KD, Crawford JT, Bates JH, Gicquel B, et al. IS6110, an

IS-like element of Mycobacterium tuberculosis complex. Nucleic Acids Res [Internet].

1990 Jan 11 [cited 2017 Sep 19];18(1):188. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/2155396

10. Kamerbeek J, Schouls L, Kolk A, van Agterveld M, van Soolingen D, Kuijper S, et al.

Simultaneous detection and strain differentiation of Mycobacterium tuberculosis for

diagnosis and epidemiology. J Clin Microbiol [Internet]. 1997 Apr [cited 2017 May

30];35(4):907–14. Available from: http://www.ncbi.nlm.nih.gov/pubmed/9157152

11. Supply P, Magdalena J, Himpens S, Locht C. Identification of novel intergenic

repetitive units in a mycobacterial two-component system operon. Mol Microbiol

[Internet]. 1997 Dec [cited 2017 Jul 7];26(5):991–1003. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/9426136

12. Supply P, Allix C, Lesjean S, Cardoso-Oelemann M, Rusch-Gerdes S, Willery E, et al.

Proposal for Standardization of Optimized Mycobacterial Interspersed Repetitive

Unit-Variable-Number Tandem Repeat Typing of Mycobacterium tuberculosis. J Clin

Microbiol [Internet]. 2006 Dec 1 [cited 2017 May 3];44(12):4498–510. Available from:

460 http://www.ncbi.nlm.nih.gov/pubmed/17005759

461 13. Wyllie D, Davidson J, Walker T, Rathod P, Peto T, Robinson E, et al. A quantitative

462   evaluation of MIRU-VNTR typing against whole-genome sequencing for identifying

463   Mycobacterium tuberculosis transmission: A prospective observational cohort study.

464   bioRxiv [Internet]. 2018 Jan 24 [cited 2018 Mar 19];252734. Available from:

465   https://www.biorxiv.org/content/early/2018/01/24/252734

466 14. Walker TM, Ip CLC, Harrell RH, Evans JT, Kapatai G, Dedicoat MJ, et al. Whole-genome

467   sequencing to delineate Mycobacterium tuberculosis outbreaks: a retrospective

468   observational study. Lancet Infect Dis [Internet]. 2013 Feb [cited 2014 Feb

469   23];13(2):137–46. Available from:

470   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3556524&tool=pmcentr

471   ez&rendertype=abstract

472 15. Walker TM, Merker M, Knoblauch AM, Helbling P, Schoch OD, van der Werf MJ, et al.

473   A cluster of multidrug-resistant Mycobacterium tuberculosis among patients arriving

474   in Europe from the Horn of Africa: a molecular epidemiological study. Lancet Infect

475   Dis [Internet]. 2018 Jan 8 [cited 2018 Mar 20]; Available from:

476   https://www.sciencedirect.com/science/article/pii/S1473309918300045

477 16. Comas I. Genomic Epidemiology of Tuberculosis. In Springer, Cham; 2017 [cited 2018

478   Mar 20]. p. 79–93. Available from: http://link.springer.com/10.1007/978-3-319-

479   64371-7_4

480 17. Kohl TA, Diel R, Harmsen D, Rothgänger J, Walter KM, Merker M, et al. Whole-

481   genome-based Mycobacterium tuberculosis surveillance: a standardized, portable,

482   and expandable approach. J Clin Microbiol [Internet]. 2014 Jul 1 [cited 2017 Sep

483   18];52(7):2479–86. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24789177

484   18.   Kohl TA, Harmsen D, Rothgänger J, Walker T, Diel R, Niemann S. Harmonized Genome

485   Wide Typing of Tubercle Bacilli Using a Web-Based Gene-By-Gene Nomenclature

486   System. EBioMedicine. 2018;

487   19.   Bjorn-Mortensen K, Soborg B, Koch A, Ladefoged K, Merker M, Lillebaek T, et al.

488   Tracing Mycobacterium tuberculosis transmission by whole genome sequencing in a

489   high incidence setting: a retrospective population-based study in East Greenland. Sci

490   Rep [Internet]. 2016 Sep 12 [cited 2017 May 3];6:33180. Available from:

491   http://www.ncbi.nlm.nih.gov/pubmed/27615360

492   20.   Hatherell H-A, Colijn C, Stagg HR, Jackson C, Winter JR, Abubakar I. Interpreting whole

493   genome sequencing for investigating tuberculosis transmission: a systematic review.

494   BMC Med [Internet]. 2016 Dec 23 [cited 2017 Sep 13];14(1):21. Available from:

495   http://www.biomedcentral.com/1741-7015/14/21

496   21.   Guthrie JL, Gardy JL. A brief primer on genomic epidemiology: lessons learned from

497   Mycobacterium tuberculosis. Ann N Y Acad Sci [Internet]. 2017 Jan 1 [cited 2018 Apr

498   16];1388(1):59–77. Available from: http://doi.wiley.com/10.1111/nyas.13273

499   22.   Bryant JM, Schürch AC, van Deutekom H, Harris SR, de Beer JL, de Jager V, et al.

500   Inferring patient to patient transmission of Mycobacterium tuberculosis from whole

501   genome sequencing data. BMC Infect Dis [Internet]. 2013 Feb 27 [cited 2014 Feb

502   13];13(1):110. Available from:

503   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3599118&tool=pmcentr

504   ez&rendertype=abstract

505   23.   Duchêne S, Holt KE, Weill F-X, Le Hello S, Hawkey J, Edwards DJ, et al. Genome-scale

506   rates of evolutionary change in bacteria. Microb genomics [Internet]. 2016 Nov [cited

507   2017 Jul 7];2(11):e000094. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/28348834

24.   Eldholm V, Monteserin J, Rieux A, Lopez B, Sobkowiak B, Ritacco V, et al. Four

decades of transmission of a multidrug-resistant Mycobacterium tuberculosis

outbreak strain. Nat Commun [Internet]. 2015 May 11 [cited 2016 Aug 17];6:7119.

Available from: http://www.nature.com/doifinder/10.1038/ncomms8119

25.   Wirth T, Hildebrand F, Allix-Béguec C, Wölbeling F, Kubica T, Kremer K, et al. Origin,

Spread and Demography of the Mycobacterium tuberculosis Complex. Achtman M,

editor. PLoS Pathog [Internet]. 2008 Sep 26 [cited 2018 Apr 10];4(9):e1000160.

Available from: http://dx.plos.org/10.1371/journal.ppat.1000160

26.   Ford CB, Shah RR, Maeda MK, Gagneux S, Murray MB, Cohen T, et al. Mycobacterium

tuberculosis mutation rate estimates from different lineages predict substantial

differences in the emergence of drug-resistant tuberculosis. Nat Genet [Internet].

2013 Jul [cited 2017 Aug 30];45(7):784–90. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/23749189

27.   Kohl TA, Utpatel C, Schleusener V, De Filippo MR, Beckert P, Cirillo DM, et al. MTBseq:

A comprehensive pipeline for whole genome sequence analysis of Mycobacterium

tuberculosis complex isolates. Prep. 2018;

28.   Lew JM, Kapopoulou A, Jones LM, Cole ST. TubercuList – 10 years after. Tuberculosis

[Internet]. 2011 [cited 2017 May 3];91(1):1–7. Available from:

http://www.sciencedirect.com/science/article/pii/S1472979210001113

29.   Feuerriegel S, Schleusener V, Beckert P, Kohl TA, Miotto P, Cirillo DM, et al. PhyResSE:

a Web Tool Delineating Mycobacterium tuberculosis Antibiotic Resistance and

Lineage from Whole-Genome Sequencing Data. Carroll KC, editor. J Clin Microbiol

[Internet]. 2015 Jun [cited 2017 May 3];53(6):1908–14. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/25854485

30.   Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, et al. BEAST 2: A Software

Platform for Bayesian Evolutionary Analysis. Prlic A, editor. PLoS Comput Biol

[Internet]. 2014 Apr 10 [cited 2017 May 3];10(4):e1003537. Available from:

http://dx.plos.org/10.1371/journal.pcbi.1003537

31.   Drummond AJ, Ho SYW, Phillips MJ, Rambaut A, Rambaut A. Relaxed Phylogenetics

and Dating with Confidence. Penny D, editor. PLoS Biol [Internet]. 2006 Mar 14 [cited

2017 May 3];4(5):e88. Available from:

http://dx.plos.org/10.1371/journal.pbio.0040088

32.   Drummond AJ, Rambaut A, Shapiro B, Pybus OG. Bayesian Coalescent Inference of

Past Population Dynamics from Molecular Sequences. Mol Biol Evol [Internet]. 2005

Feb 9 [cited 2017 May 3];22(5):1185–92. Available from:

https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msi103

33.   Leaché AD, Banbury BL, Felsenstein J, de Oca A nieto-M, Stamatakis A, K. S, et al.

Short Tree, Long Tree, Right Tree, Wrong Tree: New Acquisition Bias Corrections for

Inferring SNP Phylogenies. Syst Biol [Internet]. 2015 Nov [cited 2017 May

5];64(6):1032–47. Available from: https://academic.oup.com/sysbio/article-

lookup/doi/10.1093/sysbio/syv053

34.   Comas I, Homolka S, Niemann S, Gagneux S. Genotyping of Genetically Monomorphic

Bacteria: DNA Sequencing in Mycobacterium tuberculosis Highlights the Limitations

of Current Methodologies. PLoS One [Internet]. 2009 Nov 12 [cited 2018 Apr

10];4(11):e7815. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19915672

35.   Kato-Maeda M, Gagneux S, Flores LL, Kim EY, Small PM, Desmond EP, et al. Strain

classification of Mycobacterium tuberculosis: congruence between large sequence

polymorphisms and spoligotypes. Int J Tuberc Lung Dis [Internet]. 2011 Jan [cited

2017 Sep 6];15(1):131–3. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/21276309

36.   Filliol I, Motiwala AS, Cavatore M, Qi W, Hazbón MH, Bobadilla del Valle M, et al.

Global phylogeny of Mycobacterium tuberculosis based on single nucleotide

polymorphism (SNP) analysis: insights into tuberculosis evolution, phylogenetic

accuracy of other DNA fingerprinting systems, and recommendations for a minimal

standard SNP set. J Bacteriol [Internet]. 2006 Jan 15 [cited 2017 Sep 6];188(2):759–

72. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16385065

37.   Scott AN, Menzies D, Tannenbaum T-N, Thibert L, Kozak R, Joseph L, et al. Sensitivities

and specificities of spoligotyping and mycobacterial interspersed repetitive unit-

variable-number tandem repeat typing methods for studying molecular epidemiology

of tuberculosis. J Clin Microbiol [Internet]. 2005 Jan [cited 2017 Jul 7];43(1):89–94.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/15634955

38.   Stucki D, Ballif M, Egger M, Furrer H, Altpeter E, Battegay M, et al. Standard

Genotyping Overestimates Transmission of Mycobacterium tuberculosis among

Immigrants in a Low-Incidence Country. J Clin Microbiol [Internet]. 2016 Jul 1 [cited

2017 Jul 21];54(7):1862–70. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/27194683

39.   Walker TM, Lalor MK, Broda A, Ortega LS, Morgan M, Parker L, et al. Assessment of

Mycobacterium tuberculosis transmission in Oxfordshire, UK, 2007-12, with whole

pathogen genome sequences: an observational study. Lancet Respir Med [Internet].

2014 Apr [cited 2018 Apr 10];2(4):285–92. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/24717625

40.   Salathé M, Bengtsson L, Bodnar TJ, Brewer DD, Brownstein JS, Buckee C, et al. Digital

581   Epidemiology. PLoS Comput Biol [Internet]. 2012 Jul 26 [cited 2018 Apr

582   10];8(7):e1002616. Available from: http://dx.plos.org/10.1371/journal.pcbi.1002616

583   41.   Sanoussi CN, Affolabi D, Rigouts L, Anagonou S, de Jong B. Genotypic characterization

584   directly applied to sputum improves the detection of Mycobacterium africanum West

585   African 1, under-represented in positive cultures. Picardeau M, editor. PLoS Negl Trop

586   Dis [Internet]. 2017 Sep 1 [cited 2018 Mar 22];11(9):e0005900. Available from:

587   http://dx.plos.org/10.1371/journal.pntd.0005900

588   42.   Phelan JE, Coll F, Bergval I, Anthony RM, Warren R, Sampson SL, et al. Recombination

589   in pe/ppe genes contributes to genetic variation in Mycobacterium tuberculosis

590   lineages. BMC Genomics [Internet]. 2016 Feb 29 [cited 2018 Mar 16];17:151.

591   Available from: http://www.ncbi.nlm.nih.gov/pubmed/26923687

592   43.   Ates LS, Dippenaar A, Ummels R, Piersma SR, van der Woude AD, van der Kuij K, et al.

593   Mutations in ppe38 block PE_PGRS secretion and increase virulence of

594   Mycobacterium tuberculosis. Nat Microbiol [Internet]. 2018 Feb 15 [cited 2018 Mar

595   16];3(2):181–8. Available from: http://www.nature.com/articles/s41564-017-0090-6

596   44.   Guthrie JL, Kong C, Roth D, Jorgensen D, Rodrigues M, Hoang L, et al. Molecular

597   Epidemiology of Tuberculosis in British Columbia, Canada: A 10-Year Retrospective

598   Study. Clin Infect Dis [Internet]. 2018 Mar 5 [cited 2018 Mar 20];66(6):849–56.

599   Available from: http://www.ncbi.nlm.nih.gov/pubmed/29069284

600   45.   Asnicar F, Weingart G, Tickle TL, Huttenhower C, Segata N. Compact graphical

601   representation of phylogenetic data and metadata with GraPhlAn. PeerJ [Internet].

602   2015 Jun 18 [cited 2017 Aug 23];3:e1029. Available from:

603   http://www.ncbi.nlm.nih.gov/pubmed/26157614

Table legends

Table 1: Clustering method overview.

For each clustering method, the general features are outlined in the table. Median ages and 95% HPD ranges are based upon the BEAST2

estimates of clade heights (see methods).

Method	Strains in clusters	Number of clusters	Percent of strains in clusters	Cluster sizes	Maximum SNP distances	Clustering rate	Median Timespan	Timespan 95% HPD
Spoligotype	118	21	36.42	2-28	0-189	0.2994	76.51	0.81 - 823.21
MIRU-VNTR	121	32	37.35	2-11	0-48	0.2747	26.08	0 - 162.27
12 SNP cluster	242	47	74.69	2-34	0-23	0.6019	23.63	0 - 102.58
5 SNP cluster	147	40	45.37	2-27	0-10	0.3302	10.86	0 - 47.07
1 SNP cluster	74	29	22.84	2-6	0-2	0.1389	3.91	0 - 23.54

<u>Figure 1: Clustering of *M. tuberculosis* isolates.</u>

For each approach the inclusion of an isolate into a cluster is outlined in the surrounding

circles using GraPhlAn[45]. If an isolate is in a cluster not affected by convergence, it is

highlighted in black for the given method. If an isolate is in a cluster affected by

convergence, it is shown in grey. The ML phylogenetic tree is rooted between L4 and L5

isolates.

<u>Figure 2: Timespans associated with transmission clusters</u>

For each clustering method, the timespan associated with a cluster was estimated using

BEAST-2. The ages of each cluster (Y-axis) was aggregated per clustering method (X-axis).

Violin plots show the mean (black dot) for timespans along with the proportion of clusters

with a given age (coloured kernel plots). Note the y-axis is cut at 200 years. See Table 1 for

full 95% HPD for each method.

15