# The relationship between transmission time and clustering methods in *Mycobacterium tuberculosis* epidemiology

Conor J Meehan[1,*], Pieter Moris[1,2,3], Thomas A. Kohl[4,5], Jūlija Pečerska[6], Suriya Akter[1], Matthias Merker [4,5], Christian Utpatel[4,5], Patrick Beckert[4,5], Florian Gehre[1,7,8], Pauline Lempens[1], Tanja Stadler[6], Michel K. Kaswa[1,10], Denise Kühnert[9], Stefan Niemann[4,5,&], Bouke C de Jong[1,&]

**Affiliations**

[1]Unit of Mycobacteriology, Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

[2]Advanced Database Research and Modelling (ADReM), Department of Mathematics and Computer Science, University of Antwerp, Antwerp, Belgium

[3]Biomedical Informatics Research Network Antwerp (biomina), University of Antwerp, Antwerp, Belgium

[4]German Center for Infection Research, Partner Site Hamburg-Lübeck-Borstel-Riems, Borstel, Germany

[5]Molecular and Experimental Mycobacteriology, Priority Area Infections, Research Center Borstel, Borstel, Germany

[6]Department of Biosystems Science and Engineering, ETH Zürich, Switzerland

[7]Vaccines and Immunity Theme, Medical Research Council Unit The Gambia

[8]Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany

[9]University Hospital Zurich, Zurich, Switzerland

[10]National Tuberculosis Program, Kinshasa, Democratic Republic of Congo

[&]equal contribution; *corresponding author: cmeehan@itg.be

**Abstract**

Tracking recent transmission is a vital part of controlling widespread pathogens such as *Mycobacterium tuberculosis*. Multiple approaches exist for detecting recent transmission chains, usually by clustering strains based on the similarity of their genotyping results. However, each method gives varying estimates of transmission cluster sizes and inferring when transmission events within these clusters occurred is almost impossible. This study combines whole genome sequence (WGS) data derived from a high endemic setting with phylodynamics to unveil the timing of transmission events posited by a variety of standard genotyping methods. Our results suggest that clusters based on spoligotyping could encompass transmission events that occurred hundreds of years prior to sampling while 24-loci-MIRU-VNTR often represented decades of transmission. Instead, WGS based genotyping applying a low SNP thresholds allows for estimation of recent transmission events. These findings can guide the selection of appropriate clustering methods for uncovering relevant transmission chains within a given time-period.

**Introduction**

Despite the large global efforts at curbing the spread of *Mycobacterium tuberculosis* complex (Mtbc) strains, 10.4 million new patients develop tuberculosis (TB) every year (WHO, 2018). In addition, the prevalence of multidrug resistant Mtbc strains (MDR-TB) is increasing (WHO, 2018), predominantly through ongoing transmission within large populations (Kendall et al., 2015; Merker et al., 2015). The tracking and timing of recent transmission chains allows TB control programs to effectively pinpoint transmission hotspots and employ targeted intervention measures. This is especially important for the transmission of drug resistant strains as it appears that drug resistance may be transmitted more frequently than acquired

(Kendall et al., 2015; Trauer, Denholm, & McBryde, 2014). Thus, interrupting transmission is

key for the control of MDR-TB (Klopper et al., 2013; Merker et al., 2015; Shah et al., 2017).

For the development of the most effective control strategies, there is a strong need for (i)

appropriate identification of relevant transmission chains, risk factors and hotspots and (ii)

robust timing of when outbreaks first arose.

Epidemiological TB studies often apply genotyping methods to Mtbc strains to determine

whether two or more patients are linked within a transmission chain (molecular

epidemiology) (Merker, Kohl, Niemann, & Supply, 2017). Contact tracing is a primary

epidemiological method for investigating transmission networks of TB, mainly based on

patient interviews (Fox, Barry, Britton, & Marks, 2012). Although this method is often seen as

a gold standard of transmission linking, it does not always match the true transmission

patterns, even in low incidence settings (Behr et al., 1998; Diel et al., 2002; Roetzer et al.,

2011, 2013; Small et al., 1994) and misses many connections (Bjorn-Mortensen et al., 2017;

Vluggen et al., 2017). The implementation of molecular epidemiological approaches has

overcome these limitations and is often used as the main approach for cluster analysis.

Classical genotyping has involved IS*6110* DNA fingerprinting (Thierry et al., 1990; van Embden

et al., 1993), spoligotyping (Goguet de la Salmonière et al., 1997; Guernier, Sola, Brudey,

Guégan, & Rastogi, 2008; Kamerbeek et al., 1997), and variable-number tandem repeats of

mycobacterial interspersed repetitive units (MIRU-VNTR) (Supply, Magdalena, Himpens, &

Locht, 1997) which is the most common method at the moment (Merker et al., 2017). The

latter method is based on copy numbers of a sequence in tandem repeat patterns derived

from 24 distinct loci within the genome (Supply et al., 2006). If two patients have the same

classical genotyping pattern such as a 24-loci MIRU-VNTR pattern (or up to one locus

difference (Jonsson et al., 2014; Supply et al., 2006)) they are considered to be within a local

transmission chain. The combination of spoligotyping and MIRU-VNTR-typing, where patterns

must match in both methods to be considered a transmission link, is often considered the

molecular gold standard for transmission linking and genotyping (Supply et al., 2006).

However, examples of unlinked patients with identical patterns have been observed,

suggesting that this threshold covers too broad a genetic diversity and timespan between

infections (Gardy et al., 2011; Roetzer et al., 2013).

The application of (whole genome) sequence-based approaches for similarity analysis of Mtbc

isolates and cluster determination is known to have high discriminatory power when

assessing transmission dynamics (Comas, 2017; Roetzer et al., 2013; Walker et al., 2013, 2018;

Wyllie et al., 2018). Single nucleotide polymorphisms (SNPs) in the *pncA* gene are associated

with resistance to pyrazinamide (PZA) and can be used to improve the discriminatory power

of spoligotyping in a method referred to as SpoNC (Said et al., 2016). However, this is limited

by the low occurrence of PZA resistance, even in MDR-TB isolates (Kurbatova, Cavanaugh,

Dalton, S. Click, & Cegielski, 2013; Ngabonziza et al., 2017; Xu et al., 2016; Zignol et al., 2016).

The advent of widespread whole genome sequencing (WGS) capabilities has allowed for

highly discriminatory analyses of Mtbc strains either using core genome multi-locus sequence

typing (cgMLST) (Thomas A. Kohl et al., 2014) or SNP distances (Bjorn-Mortensen et al., 2016;

Gardy et al., 2011; Roetzer et al., 2013; Walker et al., 2013, 2018). WGS-based approaches

compare the genetic relatedness of the genomes of the clinical strains under consideration,

albeit usually excluding large repetitive portions of the genome, with the assumption that

highly similar strains are linked by a recent transmission event (Roetzer et al., 2013; Walker

et al., 2013). Although many SNP cut-offs for linking isolates have been proposed (Hatherell

et al., 2016), the most commonly employed is based on the finding that a 5 SNP cut-off will

cluster the genomes of strains from the majority of epidemiologically linked TB patients, with

an upper bound of 12 SNPs between any two linked isolates (Walker et al., 2013). The

widespread use of WGS has quickly pushed these cut-offs to be considered the new molecular

gold standard of recent transmission linking, although SNP distances may vary for technical

reasons (e.g. assembly pipelines or filter criteria (Guthrie & Gardy, 2017)) and between study

populations e.g. high and low incidence settings (Bjorn-Mortensen et al., 2016).

In addition to cluster detection, uncovering the timing of transmission events within a given

cluster is highly useful information for TB control e.g. for assessing the impact of interventions

on the spread of an outbreak. Accordingly, knowledge of the rate change associated with

different genotyping methods is essential for correct timing. The whole genome mutation

rate of Mtbc strains has been estimated by several studies as between $10^{-7}$ and $10^{-8}$

substitutions per site per year or ~0.3-0.5 SNPs per genome per year (Bryant et al., 2013;

Duchêne et al., 2016; Eldholm et al., 2015; Eldholm & Balloux, 2016; Roetzer et al., 2013;

Walker et al., 2013) while the rate of change in the MIRU-VNTR loci specifically is known to

be quicker (~$10^{-3}$) (Ragheb et al., 2013; Wirth et al., 2008). Since these mutation rates have

been shown to also vary by lineage (Duchêne et al., 2016; Ford et al., 2013) and over short

periods of time (Bryant et al., 2013), such variation needs to be accounted for, e.g. in Bayesian

phylogenetic dating techniques (Bryant et al., 2013; Merker et al., 2015; Wirth et al., 2008).

Considering the multiple genotyping methods currently available, many of them proposed as

a "gold standard", there is an urgent need to precisely define the individual capacity of each

method to accurately detect recent transmission events and perform timing of outbreaks. To

provide this essential information, this study harnesses the power of WGS-based

phylogenetic dating methods to assign timespans onto Mtbc transmission chains

encompassed by the different genotypic clustering methods commonly used in TB

transmission studies.

**Results**

In this study, we assessed 20 different approaches for generating putative *M. tuberculosis*

transmission clusters (see methods for approaches and naming schemes) using a dataset of

324 phenotypically rifampicin resistant isolates collected 2005-2010 from retreatment cases

in Kinshasa, Democratic Republic of Congo (DRC). These 20 sets of clustering patterns were

then characterised using whole genome sequence data and the propensity for convergence

of clustering patterns was estimated (see methods). Bayesian phylodynamic approaches

implemented in BEAST-2 (Bouckaert et al., 2014) were then utilised to assign timespans to

the transmission events estimated by each genotyping method.

As expected, both the genome- and membrane-based spoligotyping approaches (named Gen-

Spo and Mem-Spo respectively), clustered the most strains, with the lowest resolution (i.e.

highest clustering rate) (Figure 1, Table 1). Convergent evolution (defined as the same pattern

observed in unrelated strains; see methods) was found to affect 39% (12) of Mem-Spo

clusters and 25% (7) of Gen-Spo clusters. Additionally, some discrepancies between the Mem-

Spo and Gen-Spo patterns of each isolate were observed, with 291 isolates (90%) having the

same pattern in both Mem-Spo and Gen-Spo approaches with 1 mismatch allowed

(Supplementary table 1). The remaining 33 isolates mismatched with 2 to 17 spacers (average

of 5 spacers). Although MIRU-VNTR performed far better than spoligotyping, 16% (6) of

clustering patterns were influenced by convergence in this study (see methods) (Table 1,

Figure 1). Mixed MIRU-VNTR patterns were observed in 18 isolates although this mixing was

not observed in the WGS data.

WGS-based methods had by far the highest discriminatory power and low SNP cut-offs

grouped isolates into smaller clusters (e.g. 2-10 isolates per cluster for a 5 SNP cut-off) (Table

1, Figure 1). When the clusters were expanded to better represent transmission chains using

the novel phylogenetic inclusion method implemented here (see methods), the resulting SNP

clusters often did not increase dramatically in size (Table 1). Discriminatory power and cluster

sizes based on cgMLST alleles were similar to the SNP-based clusters (Table 1, Figure 1).

Table 1: Clustering method overview.

For each clustering method, the general features are outlined in the tables. a) All clusters for each method affected by convergence. b) Clusters

derived only from non-convergent patterns. c) SNP- and cgMLST-based methods Mean ages and 95% HPD ranges are based upon the BEAST2

estimates of clade mean heights.

**a)**

| Method | Strains in clusters | Number of clusters | Percent of strains in clusters | Cluster sizes | Maximum SNP distances | Clustering rate |
|---|---|---|---|---|---|---|
| Gen-Spo | 293 | 29 | 90.43 | 2-42 | 1-653 | 0.8148 |
| GenSpo-MIRU | 190 | 39 | 58.64 | 2-27 | 0-48 | 0.466 |
| Gen-SpoNC | 76 | 23 | 23.46 | 2-10 | 0-195 | 0.1636 |
| Mem-Spo | 276 | 33 | 85.19 | 2-39 | 1-685 | 0.75 |
| MemSpo-MIRU | 174 | 36 | 53.7 | 2-25 | 0-611 | 0.4259 |
| Mem-SpoNC | 64 | 18 | 19.75 | 2-10 | 0-21 | 0.142 |
| MIRU-VNTR | 207 | 38 | 63.89 | 2-30 | 0-611 | 0.5216 |
| MIRU-NC | 59 | 17 | 18.21 | 2-9 | 0-21 | 0.1296 |

**b)**

| Method | Strains in clusters | Number of clusters | Percent of strains in clusters | Cluster sizes | Maximum SNP distances | Clustering rate | Mean Timespan | Timespan 95% HPD |
|---|---|---|---|---|---|---|---|---|
| Gen-Spo | 191 | 22 | 58.95 | 2-37 | 1-322 | 0.5216 | 382.8101 | 0.96 - 1893.15 |
| GenSpo-MIRU | 77 | 22 | 23.77 | 2-10 | 0-48 | 0.1698 | 63.91188 | 0 - 278.77 |
| Gen-SpoNC | 34 | 11 | 10.49 | 2-6 | 0-14 | 0.071 | 21.52556 | 0.16 - 94.95 |
| Mem-Spo | 118 | 21 | 36.42 | 2-28 | 0-189 | 0.2994 | 141.1556 | 0.81 - 823.21 |
| MemSpo-MIRU | 50 | 12 | 15.43 | 2-10 | 2-48 | 0.1173 | 48.80688 | 0.8 - 216.31 |
| Mem-SpoNC | 15 | 5 | 4.63 | 2-4 | 0-14 | 0.0309 | 21.38239 | 1.03 - 97.91 |
| MIRU-VNTR | 121 | 32 | 37.35 | 2-11 | 0-48 | 0.2747 | 37.97812 | 0 - 162.27 |
| MIRU-NC | 25 | 9 | 7.72 | 2-3 | 1-11 | 0.0494 | 15.45935 | 0.77 - 58.38 |

**c)**

| Method | Strains in clusters | Number of clusters | Percent of strains in clusters | Cluster sizes | Maximum SNP distances | Clustering rate | Mean Timespan | Timespan 95% HPD |
|---|---|---|---|---|---|---|---|---|
| 0 SNP cluster | 54 | 25 | 16.67 | 2-4 | 0 | 0.0895 | 4.309937 | 0 - 15.9 |
| 1 SNP cluster | 74 | 29 | 22.84 | 2-6 | 0-2 | 0.1389 | 5.698197 | 0 - 23.54 |
| 5 SNP cluster | 147 | 40 | 45.37 | 2-27 | 0-10 | 0.3302 | 13.4115 | 0 - 47.07 |
| 12 SNP cluster | 242 | 47 | 74.69 | 2-34 | 0-23 | 0.6019 | 28.95219 | 0 - 102.58 |
| 0 SNP clade | 66 | 21 | 20.37 | 2-9 | 0-9 | 0.1389 | 5.746077 | 0 - 23.96 |
| 1 SNP clade | 80 | 27 | 24.69 | 2-9 | 0-9 | 0.1636 | 6.104103 | 0 - 25.74 |
| 5 SNP clade | 149 | 40 | 45.99 | 2-28 | 0-11 | 0.3364 | 13.48716 | 0 - 47.41 |
| 12 SNP clade | 253 | 45 | 78.09 | 2-39 | 0-27 | 0.642 | 29.73941 | 0 - 104.64 |
| 0 allele cgMLST | 51 | 24 | 15.74 | 2-4 | 0-1 | 0.0833 | 4.231405 | 0.03 - 15.48 |
| 1 allele cgMLST | 80 | 31 | 24.69 | 2-6 | 0-4 | 0.1512 | 6.371668 | 0 - 24.65 |
| 5 allele cgMLST | 173 | 42 | 53.4 | 2-28 | 0-22 | 0.4043 | 17.54352 | 0 - 68.53 |
| 12 allele cgMLST | 254 | 45 | 78.4 | 2-39 | 0-51 | 0.6451 | 30.08732 | 0 - 112.25 |

<u>Figure 1: Clustering of *M. tuberculosis* isolates.</u>

For a representative approach of each of the main methods (Mem-Spo, Gen-Spo, MIRU-VNTR,

5 SNP cut-off, 5 SNP clade and 5 cgMLST) the inclusion of an isolate into a cluster is outlined

in the surrounding circles using GraPhlAn(Asnicar, Weingart, Tickle, Huttenhower, & Segata,

2015). If an isolate is in a cluster not affected by convergence, it is highlighted in black for the

given method. If an isolate is in a cluster affected by convergence, it is shown in grey. The

clustering based on all 20 approaches is shown in Supplementary Figure 1.

Statistical estimation of the timeframe associated with particular transmission chains showed

large differences in estimated cluster ages between the genotyping approaches used (Table

1, Figure 2), correlating well with the difference in discriminatory power. Cluster ages are

defined here as the most ancient transmission event that links any two isolates within a

specific cluster. Thus, in phylogenetic terms, the cluster age is the difference in time between

when the most recent common ancestor (MCRA) of the entire cluster existed and the date of

isolation of the furthest isolate from this ancestor. The aggregate mean ages of clusters

derived from spoligotyping approaches were found to often be several hundreds of years old

(Gen-Spo: 383 years ago (95% HPD: 1-1893); Mem-Spo: 141 years ago (95% HPD: 1-823))

(Table 1b, Figure 2a). The addition of MIRU-VNTR or *pncA* mutation data to spoligotyping

resulted in clusters that, on average, originated less than 100 years ago (Table 1b, Figure 2a).

MIRU-VNTR alone gave similar cluster ages as to when combined with spoligotyping (MIRU-

VNTR: 38 (0-162); GenSpo-MIRU: 64 (0-279); MemSpo-MIRU: 49 (1-216)) (Table 1b, Figure

2a).

Clusters based on SNP cut-offs correlated to 4 years of transmission using a 0 SNP cut-off (95%

HPD: 0-16), 6 years using a 1 SNP cut-off (95% HPD: 0-24), 13 years using a 5 SNP cut-off (95%

HPD: 0-47), and 29 years using a 12 SNP cut-off (95% HPD: 0-103) (Table 1c, Figure 2b).

Extension on the tree using the phylogenetic inclusion approach to form SNP clades did not

greatly increase the lengths of transmissions encompassed by clusters (one year increase, on

average) (Table 1c). Similar findings were obtained when clusters were based on allele

differences in the cgMLST method: 4 years of transmission using a 0 cgMLST cut-off (95% HPD:

0-15), 6 years using a 1 cgMLST cut-off (95% HPD: 0-25), 18 years using a 5 cgMLST cut-off

(95% HPD: 0-68), and 30 years using a 12 cgMLST cut-off (95% HPD: 0-112) (Table 1c, Figure

2b)

<u>Figure 2: Timespans associated with transmission clusters</u>

For each clustering method, the timespan associated with a cluster was estimated using

BEAST-2. The ages of each cluster (Y-axis) was aggregated per clustering method (X-axis).

Violin plots show the mean (black dot) for timespans along with the proportion of clusters

with a given age (coloured kernel plots). Methods are split as follows: A) Spoligotype-based

(Gen-Spo-based (red), Mem-Spo-based (orange)) and MIRU-VNTR-based (yellow), B) SNP-

based (blue) and cgMLST-based (green). Note the y-axis is different for each and panel A) is

cut at 400 years.

**Discussion**

The term 'recent transmission' is often applied to gain a better understanding of the current transmission dynamics of pathogens in a given population. However, little data is available on how recent a likely transmission event occurred when measured with different genotyping methods. To get a better understanding of the discriminatory power of different classical genotyping techniques and WGS-based approaches in relation to outbreak timing, this study has performed an in-depth comparison of clustering rates and dated phylogenies obtained in a collection of 324 Mtbc strains from a high incidence setting (Kinshasa, DRC). With a whole genome phylodynamic approach employed as a gold standard, our study demonstrates that each genotyping method was associated with a specific discriminatory power resulting in clusters representing vastly different time periods of transmission events (Table 1 and Figure 2). This has significant implications for data interpretations e.g. when selecting and utilising different genotyping methods/clustering approaches for epidemiological studies and assessing the effectiveness of public health intervention strategies.

As the most extreme example, spoligotyping-derived clusters were associated with transmission events that can be hundreds of years old. This low discriminatory power coupled with the high rate of convergent evolution (the same spoligotype pattern found in phylogenetically distant isolates) in both Mem-Spo and Gen-Spo add weight to the previous suggestion that these techniques are not suitable for recent transmission studies (Comas, Homolka, Niemann, & Gagneux, 2009), although Mem-Spo may be of use as a low-cost method of sorting Mtbc strains into the seven primary lineages (Filliol et al., 2006; Kato-Maeda et al., 2011). Differences between Mem-Spo and Gen-Spo patterns from the same isolate were observed for 10% of isolates in this study, even after rechecking of patterns,

requiring more investigation into which method is closer to the 'true' spoligotyping pattern

within a genome (Coll et al., 2012; Mokrousov et al., 2016; Warren et al., 2002; Xia, Teo, &

Ong, 2016).

In line with previous findings (Comas et al., 2009; Scott et al., 2005), convergent evolution of

24-loci MIRU-VNTR patterns was rarer than observed for spoligotyping, but did occur in 16%

of MIRU-VNTR-based clusters. Additionally, the transmission times encompassed by MIRU-

VNTR clusters spanned several decades (Table 1b, Figure 2a), confirming previous studies

showing over-estimation of recent transmission with this method (Bjorn-Mortensen et al.,

2016; Roetzer et al., 2013; Stucki et al., 2016; Wyllie et al., 2018).

The combination of MIRU-VNTR or spoligotyping with *pncA* mutations (MIRU-NC and Gen-

SpoNC/Mem-SpoNC) appeared to reflect true clusters of PZA resistance transmission based

on the relatively young ages of such transmission clusters (Table 1b). Thus, as discussed

before (den Hertog, Sengstake, & Anthony, 2015; Sengstake et al., 2017), although

transmission of *pncA* mutations seems to occur, further investigation is needed to find out

whether *pncA* mutants are less transmissible than those with a wildtype gene.

For defining transmission events that occurred in more recent time frames before sampling,

WGS-based methods (SNP or cgMLST) were found to be better suited than classical

genotyping methods (Table 1, Figure 2). The 12 SNP cut-off, currently the recommended

upper bound for clustering isolates, likely defines transmission events that occurred on

average three decades prior to sampling, similar in age to clusters estimated by MIRU-VNTR.

This suggests that the 12 SNP cluster method may be a good replacement for MIRU-VNTR as

it detects larger transmission networks spanning similar transmission time periods but is less

affected by convergent evolution. Isolates clustered at identical (0 SNP) or nearly identical (1

SNP) cut-offs were found to represent transmission events occurring four to six years

previous. These findings correlate well with previous studies where confirmed contact

tracing-based epidemiological links were found between patients that were two (Walker et

al., 2014), three (Roetzer et al., 2013) or five (Walker et al., 2013) SNPs apart. Indeed, a recent

study of a cross-country MDR-TB outbreak found only a maximum of two SNP differences

between all 29 isolates involved in the origin of the outbreak (Walker et al., 2018). Although

this supports their use for detection or exclusion of very recent transmission, this low

variability between isolates makes robust identification of transmission direction impossible,

especially during short timespans.

Comparisons between the SNP-based (using almost all genomic differences) and the cgMLST-

based cluster detection (using a defined core set of genes) demonstrated that the latter

approach gives similar estimations to full SNP approaches. However, as current SNP assembly

pipelines for Illumina data exclude repetitive region such as PE/PPE genes, larger differences

between cgMLST and full SNP estimation may be seen once all aspects of the genome can be

utilised.

Different clustering approaches can be applied when grouping isolates by SNP distance. Two

partitional clustering methods are primarily utilised: either the creation of tight clusters

(where the maximum pairwise distance between isolates in a cluster is less than the SNP cut-

off; e.g. (Thomas A. Kohl et al., 2014)) or loose clusters (where each isolate is less than the

SNP cut-off distance from at least one other isolate in the cluster; e.g. (Walker et al., 2014)).

Tight clusters ensure high connectivity within clusters, but may result in isolates belonging to

multiple groups, making interpretation and delineation of transmission events difficult. Loose

clusters (the definition used in this study), separate isolates into non-overlapping clusters, but

may result in low connectivity within clusters. Here we present an extension of the loose

cluster, termed the phylogenetic inclusion method, which adds all other isolates with the

same phylogenetically defined common ancestor to the cluster, potentially identifying larger

circulating genotypes. Tight, loose and phylogenetic inclusion clusters each aim to define

different levels of connectivity through time, an aspect that should be considered when

selecting the appropriate clustering approach.

The mutation rate of *M. tuberculosis* has been estimated to be between $10^{-7}$ and $10^{-8}$

substitutions per site per year (Duchêne et al., 2016; Merker et al., 2015; Roetzer et al., 2013).

Within the Bayesian analysis employed here, the mutation rate was free to vary between

these values but was found to strongly favour ~$3 \times 10^{-8}$ (ESS > 1000 for all runs), translating to

approximately 0.3 SNPs per genome per year. While the mutation rate used here is primarily

applicable for lineage 4 (which most of this dataset is comprised of) and in line with previous

estimates for this lineage (Duchêne et al., 2016), it may be similar in other lineages, although

this has only been shown for lineage 2 (Duchêne et al., 2016; Merker et al., 2015). Thus, per-

lineage estimates are required for all seven lineages to ensure similar transmission times are

linked to genotyping methods across the whole population diversity of the Mtbc.

While this study has many advantages due to its five year population based design in an

endemic setting coupled with the application of three different genotyping methods

(membrane based spoligotyping analysis, 24-locus MIRU-VNTR and WGS), future

confirmatory studies could address the following drawbacks that are inherent to genomic

epidemiology (Comas, 2017; Guthrie & Gardy, 2017): 1) studies employing contact tracing

and/or digital epidemiology (Salathé et al., 2012) in conjunction with these genotyping

methods can help confirm transmission times associated with different clusters; 2) as outlined

above, strains of other lineages of the Mtbc should be analysed in a similar fashion to ensure

transferability of findings across the entire complex; 3) a broad range of drug resistance

profiles should be included to fully assess the impact of such mutations on transmission

estimates; 4) improved WGS methods, such as directly from clinical samples to help reduce

culture biases (Sanoussi, Affolabi, Rigouts, Anagonou, & de Jong, 2017) and longer reads (e.g.

PacBio SMRT or Nanopore MinION) to capture the entire genome, including repetitive regions

such as PE/PPE genes known to impact genome remodelling (Ates et al., 2018; Phelan et al.,

2016), will ensure that the maximum diversity between isolates is captured and 5)

standardised SNP calling pipelines appropriate across all lineages, with high true positive/low

false negative rates, will ensure that Mtbc molecular epidemiology can be uniformly

implemented and comparable across studies.

In conclusion, since each method was found to represent different timespans and clustering

definitions, they can be used in a stratified manner in an integrated epidemiological and

public health investigation addressing the transmission of Mtbc strains. For instance,

although spoligotyping clusters represented potentially very old transmission events, the low

associated cost and its ability to be applied directly on sputum helps reduce culture bias and

thus robustly assign lineages. Thus, spoligotyping and/or MIRU-VNTR would serve well as

first-line surveillance of potential transmission events in the population, guiding further

investigations and resource allocations.

These potential transmission hotspots could be further investigated with contact tracing

and/or WGS. Employment of different cut-offs and clustering approaches to WGS data can

then address several questions. The 12 SNP cluster/clade or 12 allele cgMLST approaches

serve well for high level surveillance targeting larger (older) transmission networks, akin to

what is currently often done using MIRU-VNTR (e.g. (Guthrie et al., 2018; Walker et al., 2018)).

Recent transmission events can then be detected through employment of low SNP or cgMLST-

based cut-offs (e.g. 5 SNPs for transmission in the past 15 years or 0-1 SNPs for transmission

in the past 5 years). These clusters can then be linked to historical isolates or other clusters

through employment of the phylogenetic inclusion method to resolve the local circulating

genotypes. This is especially useful if bursts of sampling are undertaken such as in drug

resistance surveys (WHO, 2015), which are increasingly employing WGS approaches (Cabibbe

& Cirillo, 2016; Zignol et al., 2016, 2018). Alternatively, in high incidence/low diversity settings

where amalgamation of clusters may inadvertently obscure distinct hotspots of transmission

at different time points, subdivision into distinct time-dependant clusters can be undertaken

using the algorithm presented in such a study in East Greenland (Bjorn-Mortensen et al.,

2016).

Overall, phylodynamic approaches applied to whole genome sequences, as undertaken here,

are recommended to fully investigate the specific transmission dynamics within a study

population to account for setting-specific conditions, such as low/high TB incidence, low/high

pathogen population diversity, sampling fractions and social factors influencing transmission.

Thus, each genotyping method can be employed as part of an overall evidence gathering

program for transmission, placing molecular epidemiological approaches as an integral part

in tracking and stopping the spread of TB.

**Materials and Methods**

<u>Dataset and sequencing</u>

A set of 324 isolates from Kinshasa, Democratic Republic of Congo were collected from

consecutive retreatment TB patients between 2005 and 2010 at TB clinics, servicing an

estimated 30% of the population of Kinshasa. All isolates were phenotypically resistant to

rifampicin (RR-TB) and the majority are also isoniazid resistant (i.e. MDR-TB). Use of the

stored isolates without any linked personal information was approved by the health

authorities of the DRC and the Institutional Review Board of the ITM in Antwerp (ref no

945/14). Libraries for whole genome sequencing were prepared from extracted genomic DNA

with the Illumina Nextera XT kit, and run on the Illumina NextSeq platform in a 2x151bp run

according to manufacturer's instructions. Illumina read sets will be available at ReSeqTB

(platform.reseqtb.org) upon publication.

<u>Genome reconstruction and maximum likelihood phylogeny estimation</u>

The MTBseq pipeline(Thomas Andreas Kohl et al., 2018) was used to detect the SNPs for each

isolate using the H37Rv reference genome (NCBI accession number NC000962.3) (Lew,

Kapopoulou, Jones, & Cole, 2011; Médigue, Cole, Camus, & Pryor, 2002). Sites known to be

involved in drug resistance (as outlined in the PhyResSE list of drug mutations v27 (Feuerriegel

et al., 2015)) were excluded from the alignment and additional filtering of sites with

ambiguous calls in >5% of isolates and those SNPs within a 12bp window of each other was

also applied.

The SNP alignment of all isolates was used as the basis for creating a maximum likelihood

(ML) phylogeny. RAxML-NG version 0.5.1b (Kozlov, 2017) was used to reconstruct the

phylogeny from this alignment using a GTR+GAMMA model of evolution, accounting for

ascertainment bias(Lewis, 2001) with the Stamatakis reconstituted DNA approach (Leaché et

al., 2015) and site repeat optimisation (Kobert, Stamatakis, & Flouri, 2016) with 20 different

starting trees and 100 bootstraps. All subsequent topology visualisation was undertaken using

FigTree version 1.4.3 (Rambaut, 2016) and GraPhlAn (Asnicar et al., 2015).

Transmission cluster estimation methods

Several standard transmission clustering approaches were chosen for comparison and

analysis. For each method, the total SNP distances were calculated to investigate the range

of variability encompassed within each cluster. Maximum SNP distances were derived from

pairwise comparisons of isolates within the SNP alignment using custom python scripts. A

clustering rate was calculated for each method using the formula $(n_c - c)/n$, where $n_c$ is the

total number of isolates clustered by a given method, c is the number of clusters, and n is the

total number of isolates in the dataset (n=324).

Spoligotyping

Spoligotype patterns were estimated by 2 methods: membrane-based and genome-based.

Membrane-based patterns were obtained following the previously published protocol

(Kamerbeek et al., 1997). This method is referred to as Mem-Spo. Genome-based

spoligotyping was derived from the Illumina reads of each isolate using SpoTyping v2.1 (Xia

et al., 2016). Reads (both forward and reverse) were input to SpoTyping with default

parameters and the 43 spacer values were extracted from the output. This method is referred

to as Gen-Spo. For both methods, isolates were said to be clustered if all 43 spacers matched.

<u>MIRU-VNTR</u>

Genotyping by MIRU-VNTR was undertaken as previously described (Supply et al., 2006). 2 µl

of DNA was extracted from cultures and amplified using the 24 loci MIRU-VNTR typing kit

(Genoscreen, Lille, France). Analysis of patterns was undertaken using the ABI 3500 automatic

sequencer (Applied Biosystems, California, USA) and Genemapper software (Applied

Biosystems). Isolates were said to be clustered if all 24 loci matched. MIRU-VNTR patterns

were also combined with spoligotyping patterns for additional refinement of clusters. Isolates

were clustered if both the spoligotyping pattern and the 24 loci MIRU-VNTR pattern matched.

These clustering methods are referred to as MemSpo-MIRU and GenSpo-MIRU.

<u>SpoNC</u>

Transmission estimation using spoligotyping has been shown to be improved if combined

with *pncA* mutations (Said et al., 2016). This method, referred to as SpoNC, was applied to

both Mem-Spo (Mem-SpoNC) and Gen-Spo (Gen-SpoNC). Mutations in *pncA* were extracted

from the MTBseq tabular output for each isolate. All mutations were selected, regardless of

drug resistance association, as is done in the SpoNC approach. The upstream promoter region

of *pncA* did not reveal any mutations in this dataset. Isolates were said to be clustered if all

43 spacers matched and the *pncA* mutation was the same in both isolates. MIRU-VNTR

patterns were combined with *pncA* mutations in a similar manner. This is referred to as MIRU-

NC.

<u>SNP cut-off clustering</u>

The advent of whole genome reconstruction has allowed for genome-based comparisons for

transmission clustering. Previous work has suggested that linked and recent transmission can

be estimated by comparison of SNP differences between isolates. The cut-offs proposed by

Walker *et al.* (Walker et al., 2013) are the most widely used and have been employed in

multiple studies (Casali et al., 2016; Tessema et al., 2017; Witney et al., 2016). In this study,

we employed both the 5 SNP (proposed by Walker *et al*. as the likely boundary for linked

transmission) and 12 SNP cut-offs (proposed maximum boundary) for cluster definition.

Additionally, we employed lower cut-offs of 0 and 1 SNPs to look for clusters of very highly

related isolates. Pairwise SNP distances were calculated between all isolates. A loose cluster

definition was used, where every isolate in a cluster at most the SNP cut-off from at least 1

other isolate in the cluster.

Phylogenetic information was used to extend these SNP-based clusters to include any other

isolates that share the same most recent common ancestor (MRCA). These isolates may

exceed the SNP cut-off but should be included as, through sharing an MRCA, they are

intrinsically within the same putative transmission chain. The MRCA is defined here as the

internal node in a phylogenetic tree that is shared by all the isolates within the putative SNP-

based cluster. This extension was achieved by mapping each SNP cluster onto the ML

phylogenetic tree and the MRCA (shared internal node) of all isolates was found using

DendroPy v4.0.3 (Sukumaran & Holder, 2010). Any additional taxa with the same MRCA were

then added to the transmission cluster (Supplemental Figure 2). In other words, all leaf nodes

of the MRCA internal node were labelled as being part of the putative transmission cluster.

We call this approach the phylogenetic inclusion method and extended clusters are hereafter

referred to as extended SNP clades to distinguish them from SNP clusters as created by the

standard non-phylogenetic method above. The python script that implements this method

can be found at https://github.com/conmeehan/pathophy.

<u>cgMLST</u>

An alternative approach to clustering using WGS data is the concept of core genome MLST

(cgMLST) patterns (Thomas A. Kohl et al., 2014). Since SNP detection can be variable between

assembly pipelines, SNP clusters between studies may be difficult to compare. The cgMLST

approach standardises comparisons by ensuring the same core genes are always compared.

BAM files for all isolates are input into Ridom SeqSphere$^{+}$ software (Ridom GmbH, Münster,

Germany) to compile an allelic distance matrix based on the cgMLST v2 scheme consisting of

2,891 core Mtbc genes. Loose clusters were then defined as above using allelic differences of

0, 1, 5 and 12 as cut-offs. These methods are referred to as 0/1/5/12 cgMLST respectively.

<u>Detection of convergent evolution</u>

Convergent evolution towards identical patterns may occur for Spoligotyping, MIRU-VNTR

and *pncA* mutations (Driscoll, 2009; Miotto et al., 2014; Scott et al., 2005; Warren et al., 2002).

Convergence was detected and cross-checked with two methods. Firstly, Mtbc lineage and

sub-lineage numbering (Coll et al., 2014) was applied to all isolates based on the PhyResSE

lineage-defining SNP list v27 (Feuerriegel et al., 2015). If the same clustering pattern was

observed in two different sub-lineages, with other patterns seen in-between, this was flagged

as potential convergence. Additional convergence confirmation was also undertaken using

phylogenetic distances, as estimated by DendroPy. If the phylogenetic distance (combined

branch lengths that separate 2 isolates) between two isolates with identical clustering

patterns was greater than 0.0005, this was flagged as potential convergence. Any isolates

flagged by both methods (lineage-based and distance-based) were marked as clustered by

convergence. For example, if isolates with the same spoligotyping pattern appeared in lineage

4,1 and 4,6 with different patterns in-between and these isolates were distant on the tree

(distance greater than 0.0005), this was confirmed as a convergent pattern. Convergence was

checked for all approaches except the SNP cut-off clusters/clades, which, by definition, could

not be convergent. Clustering methods that combined two other methods (e.g. Gen-SpoNC)

were first checked separately for convergence and then combined to create the final clusters.

<u>Estimation of transmission times</u>

To estimate the age and timespan of potential transmission clusters, SNP alignments were

created from the convergence-free version of the five primary clustering types: Gen-Spo,

Mem-Spo, MIRU-VNTR, extended 12 SNP clades and 12 allele cgMLST. All other methods are

sub-clustering methods of at least one of these five methods (e.g. Mem-SpoNC clusters are

inherently included in any Mem-Spo clusters, and all SNP-based clusters are sub-clusters of

the 12 SNP clades).

A Bayesian approach to transmission time estimation was then undertaken. The SNP

alignments were created as above for the five high-level clustering types. Each cluster method

alignment was separately input to BEAST-2 v2.4.7 (Bouckaert et al., 2014) to create a time

tree for those isolates. These phylogenies were built using the following priors: GTR+GAMMA

substitution model, a log-normal relaxed molecular clock model to account for variation in

mutation rates (Alexei J Drummond, Ho, Phillips, Rambaut, & Rambaut, 2006) and coalescent

constant size demographic model (A. J. Drummond, Rambaut, Shapiro, & Pybus, 2005), both

of which have been found to be suitable for lineage 4 isolates in a previous study (Bjorn-

Mortensen et al., 2016). The MCMC chain was run six times independently per alignment with

a length of at least 400 million, sampled every 40,000$^{th}$ step (Gen-Spo: 400 million; extended

12 SNP & cgMLST: 500 million; MIRU & MemSpo: 600 million). A log normal prior (mean

$1.5x10^{-7}$; variance 1.0) was used for the clock model to reflect the previously estimated

mutation rate of *M. tuberculosis* lineage 4 (Bryant et al., 2013; Duchêne et al., 2016; Eldholm

et al., 2015; Eldholm & Balloux, 2016; Roetzer et al., 2013; Walker et al., 2013), while allowing

for variation as previously suggested (Bryant et al., 2013). A 1/X non-informative prior was

selected for the population size parameter of the demographic model. Isolation dates were

used as informative heterochronous tip dates and the SNP alignment was augmented with a

count of invariant sites for each of the four nucleotide bases to avoid ascertainment

bias(Leaché et al., 2015). Tracer v1.6 (Rambaut, Suchard, Xie, & Drummond, 2013) was used

to determine adequate mixing and convergence of chains (ESS >150) after a 25% burn-in. The

chains were combined via LogCombiner v2.4.8 (Bouckaert et al., 2014) to obtain a single chain

for each clustering type with high (>1000) effective sample sizes. The tree samples were

combined in the same manner and resampled at a lower frequency to create thinned samples

of (minimum) 20,000 trees.

The timespan of transmission events estimated by each method was then calculated as

follows: for each cluster created by the given method, we defined the MRCA node as the

internal node that connects all taxa in that cluster. The youngest node was then defined as

the tip that is furthest from this MRCA within the clade (i.e. the tip descendant from that node

that was sampled closest to the present time). For each retained tree in the MCMC process,

the difference in age between the MRCA node and youngest node was calculated. This gave

a distribution of likely maximum transmission event times within that cluster. For each

method, these per-cluster aggregated ages were then combined across all clusters to give a

per-method distribution of transmission event times represented by the clusters. The 95%

HPD interval of these distributions was calculated with the LaplacesDemon p.interval function

(Statisticat, 2016) in R v3.4.0 (R Core Team, 2017) and the distribution within this interval for

each method along with the mean based upon this interval were then visualized in violin plots

per clustering method using ggplot2 (Wickham, 2009) in R.

**Acknowledgements/Funding**

The authors would like to thank Armand Van Deun and Koen Vandelannoote for valuable

discussion and input and Cecile Uwizeye for aid with spoligotyping. This work was supported

by an ERC grant (INTERRUPTB; no. 311725) to BdJ, FG and CJM; an ERC grant to TS (PhyPD;

no. 335529); an FWO PhD fellowship to PM (grant number 1141217N); the German Centre

for Infection Research (DZIF) for TAK, MM and SN; a SNF SystemsX grant (TBX) to JP and TS

and a Marie Heim-Vögtlin fellowship granted to DK by the Swiss National Science Foundation.

The computational resources and services used in this work were provided by the VSC

(Flemish Supercomputer Center), funded by the Research Foundation - Flanders (FWO) and

the Flemish Government – department EWI;

**Competing interests**

The authors declare there are no competing interests attached to this work.

**References**

Asnicar, F., Weingart, G., Tickle, T. L., Huttenhower, C., & Segata, N. (2015). Compact

graphical representation of phylogenetic data and metadata with GraPhlAn. *PeerJ*, *3*,

e1029. https://doi.org/10.7717/peerj.1029

Ates, L. S., Dippenaar, A., Ummels, R., Piersma, S. R., van der Woude, A. D., van der Kuij, K.,

… Bitter, W. (2018). Mutations in ppe38 block PE_PGRS secretion and increase

virulence of Mycobacterium tuberculosis. *Nature Microbiology*, *3*(2), 181–188.

https://doi.org/10.1038/s41564-017-0090-6

Behr, M. A., Hopewell, P. C., Antonio Paz, E., Masae Kawamura, L., Schecter, G. F., & Small,

P. M. (1998). Predictive Value of Contact Investigation for Identifying Recent

Transmission of Mycobacterium tuberculosis. *American Journal of Respiratory and*

*Critical Care Medicine*, *158*(2), 465–469.

https://doi.org/10.1164/ajrccm.158.2.9801062

Bjorn-Mortensen, K., Lillebaek, T., Koch, A., Soborg, B., Ladefoged, K., Sørensen, H. C. F., …

Andersen, A. B. (2017). Extent of transmission captured by contact tracing in a

tuberculosis high endemic setting. *European Respiratory Journal*, *49*(3). Retrieved from

http://erj.ersjournals.com/content/49/3/1601851

Bjorn-Mortensen, K., Soborg, B., Koch, A., Ladefoged, K., Merker, M., Lillebaek, T., … Kohl, T.

A. (2016). Tracing Mycobacterium tuberculosis transmission by whole genome

sequencing in a high incidence setting: a retrospective population-based study in East

Greenland. *Scientific Reports*, *6*, 33180. https://doi.org/10.1038/srep33180

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., … Drummond, A. J.

(2014). BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. *PLoS*

*Computational Biology*, *10*(4), e1003537.

https://doi.org/10.1371/journal.pcbi.1003537

Bryant, J. M., Schürch, A. C., van Deutekom, H., Harris, S. R., de Beer, J. L., de Jager, V., … van

Soolingen, D. (2013). Inferring patient to patient transmission of Mycobacterium

tuberculosis from whole genome sequencing data. *BMC Infectious Diseases*, *13*(1), 110.

https://doi.org/10.1186/1471-2334-13-110

Cabibbe, A. M., & Cirillo, D. M. (2016). Best approaches to drug-resistance surveillance at

the country level. *International Journal of Mycobacteriology*, *5*, S40–S41.

https://doi.org/10.1016/j.ijmyco.2016.09.010

Casali, N., Broda, A., Harris, S. R., Parkhill, J., Brown, T., & Drobniewski, F. (2016). Whole

Genome Sequence Analysis of a Large Isoniazid-Resistant Tuberculosis Outbreak in

London: A Retrospective Observational Study. *PLOS Medicine*, *13*(10), e1002137.

https://doi.org/10.1371/journal.pmed.1002137

Coll, F., Mallard, K., Preston, M. D., Bentley, S., Parkhill, J., McNerney, R., … Clark, T. G.

(2012). SpolPred: rapid and accurate prediction of Mycobacterium tuberculosis

spoligotypes from short genomic sequences. *Bioinformatics (Oxford, England)*, *28*(22),

2991–3. https://doi.org/10.1093/bioinformatics/bts544

Coll, F., McNerney, R., Guerra-Assunção, J. A., Glynn, J. R., Perdigão, J., Viveiros, M., … Clark,

T. G. (2014). A robust SNP barcode for typing Mycobacterium tuberculosis complex

strains. *Nature Communications*, *5*, 4812. https://doi.org/10.1038/ncomms5812

Comas, I. (2017). Genomic Epidemiology of Tuberculosis (pp. 79–93). Springer, Cham.

https://doi.org/10.1007/978-3-319-64371-7_4

Comas, I., Homolka, S., Niemann, S., & Gagneux, S. (2009). Genotyping of Genetically

Monomorphic Bacteria: DNA Sequencing in Mycobacterium tuberculosis Highlights the

Limitations of Current Methodologies. *PLoS ONE*, *4*(11), e7815.

https://doi.org/10.1371/journal.pone.0007815

den Hertog, A. L., Sengstake, S., & Anthony, R. M. (2015). Pyrazinamide resistance in

Mycobacterium tuberculosis fails to bite? *Pathogens and Disease*, *73*(6), ftv037.

https://doi.org/10.1093/femspd/ftv037

Diel, R., Schneider, S., Meywald-Walter, K., Ruf, C.-M., Rusch-Gerdes, S., & Niemann, S.

(2002). Epidemiology of Tuberculosis in Hamburg, Germany: Long-Term Population-

Based Analysis Applying Classical and Molecular Epidemiological Techniques. *Journal of*

*Clinical Microbiology*, *40*(2), 532–539. https://doi.org/10.1128/JCM.40.2.532-539.2002

Driscoll, J. R. (2009). Spoligotyping for Molecular Epidemiology of the Mycobacterium

tuberculosis Complex. In *Methods in molecular biology (Clifton, N.J.)* (Vol. 551, pp. 117–

128). https://doi.org/10.1007/978-1-60327-999-4_10

Drummond, A. J., Ho, S. Y. W., Phillips, M. J., Rambaut, A., & Rambaut, A. (2006). Relaxed

Phylogenetics and Dating with Confidence. *PLoS Biology*, *4*(5), e88.

https://doi.org/10.1371/journal.pbio.0040088

Drummond, A. J., Rambaut, A., Shapiro, B., & Pybus, O. G. (2005). Bayesian Coalescent

Inference of Past Population Dynamics from Molecular Sequences. *Molecular Biology*

*and Evolution*, *22*(5), 1185–1192. https://doi.org/10.1093/molbev/msi103

Duchêne, S., Holt, K. E., Weill, F.-X., Le Hello, S., Hawkey, J., Edwards, D. J., … Holmes, E. C.

(2016). Genome-scale rates of evolutionary change in bacteria. *Microbial Genomics*,

*2*(11), e000094. https://doi.org/10.1099/mgen.0.000094

Eldholm, V., & Balloux, F. (2016). Antimicrobial Resistance in Mycobacterium tuberculosis:

The Odd One Out. *Trends in Microbiology*. https://doi.org/10.1016/j.tim.2016.03.007

Eldholm, V., Monteserin, J., Rieux, A., Lopez, B., Sobkowiak, B., Ritacco, V., … Lemey, P.

(2015). Four decades of transmission of a multidrug-resistant Mycobacterium

tuberculosis outbreak strain. *Nature Communications*, *6*, 7119.

https://doi.org/10.1038/ncomms8119

Feuerriegel, S., Schleusener, V., Beckert, P., Kohl, T. A., Miotto, P., Cirillo, D. M., …

Fellenberg, K. (2015). PhyResSE: a Web Tool Delineating Mycobacterium tuberculosis

Antibiotic Resistance and Lineage from Whole-Genome Sequencing Data. *Journal of*

*Clinical Microbiology*, *53*(6), 1908–1914. https://doi.org/10.1128/JCM.00025-15

Filliol, I., Motiwala, A. S., Cavatore, M., Qi, W., Hazbón, M. H., Bobadilla del Valle, M., …

Alland, D. (2006). Global phylogeny of Mycobacterium tuberculosis based on single

nucleotide polymorphism (SNP) analysis: insights into tuberculosis evolution,

phylogenetic accuracy of other DNA fingerprinting systems, and recommendations for

a minimal standard SNP set. *Journal of Bacteriology*, *188*(2), 759–72.

https://doi.org/10.1128/JB.188.2.759-772.2006

Ford, C. B., Shah, R. R., Maeda, M. K., Gagneux, S., Murray, M. B., Cohen, T., … Fortune, S. M.

(2013). Mycobacterium tuberculosis mutation rate estimates from different lineages

predict substantial differences in the emergence of drug-resistant tuberculosis. *Nature*

*Genetics*, *45*(7), 784–90. https://doi.org/10.1038/ng.2656

Fox, G. J., Barry, S. E., Britton, W. J., & Marks, G. B. (2012). Contact investigation for

tuberculosis: a systematic review and meta-analysis. *European Respiratory Journal*,

*41*(1). Retrieved from http://erj.ersjournals.com/content/41/1/140.short

Gardy, J. L., Johnston, J. C., Sui, S. J. H., Cook, V. J., Shah, L., Brodkin, E., … Tang, P. (2011).

Whole-Genome Sequencing and Social-Network Analysis of a Tuberculosis Outbreak.

*New England Journal of Medicine*, *364*(8), 730–739.

https://doi.org/10.1056/NEJMoa1003176

Goguet de la Salmonière, Y. O., Li, H. M., Torrea, G., Bunschoten, A., van Embden, J., &

Gicquel, B. (1997). Evaluation of spoligotyping in a study of the transmission of

Mycobacterium tuberculosis. *Journal of Clinical Microbiology*, *35*(9), 2210–4. Retrieved

from http://www.ncbi.nlm.nih.gov/pubmed/9276389

Guernier, V., Sola, C., Brudey, K., Guégan, J.-F., & Rastogi, N. (2008). Use of cluster-graphs

from spoligotyping data to study genotype similarities and a comparison of three

indices to quantify recent tuberculosis transmission among culture positive cases in

French Guiana during a eight year period. *BMC Infectious Diseases*, *8*(1), 46.

https://doi.org/10.1186/1471-2334-8-46

Guthrie, J. L., & Gardy, J. L. (2017). A brief primer on genomic epidemiology: lessons learned

from *Mycobacterium tuberculosis*. *Annals of the New York Academy of Sciences*,

*1388*(1), 59–77. https://doi.org/10.1111/nyas.13273

Guthrie, J. L., Kong, C., Roth, D., Jorgensen, D., Rodrigues, M., Hoang, L., … Gardy, J. L.

(2018). Molecular Epidemiology of Tuberculosis in British Columbia, Canada: A 10-Year

Retrospective Study. *Clinical Infectious Diseases*, *66*(6), 849–856.

https://doi.org/10.1093/cid/cix906

Hatherell, H.-A., Colijn, C., Stagg, H. R., Jackson, C., Winter, J. R., & Abubakar, I. (2016).

Interpreting whole genome sequencing for investigating tuberculosis transmission: a

systematic review. *BMC Medicine*, *14*(1), 21. https://doi.org/10.1186/s12916-016-

0566-x

Jonsson, J., Hoffner, S., Berggren, I., Bruchfeld, J., Ghebremichael, S., Pennhag, A., &

Groenheit, R. (2014). Comparison between RFLP and MIRU-VNTR Genotyping of

Mycobacterium tuberculosis Strains Isolated in Stockholm 2009 to 2011. *PLoS ONE*,

*9*(4), e95159. https://doi.org/10.1371/journal.pone.0095159

Kamerbeek, J., Schouls, L., Kolk, A., van Agterveld, M., van Soolingen, D., Kuijper, S., … van

Embden, J. (1997). Simultaneous detection and strain differentiation of Mycobacterium

tuberculosis for diagnosis and epidemiology. *Journal of Clinical Microbiology*, *35*(4),

907–14. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9157152

Kato-Maeda, M., Gagneux, S., Flores, L. L., Kim, E. Y., Small, P. M., Desmond, E. P., &

Hopewell, P. C. (2011). Strain classification of Mycobacterium tuberculosis: congruence

between large sequence polymorphisms and spoligotypes. *The International Journal of*

*Tuberculosis and Lung Disease : The Official Journal of the International Union against*

*Tuberculosis and Lung Disease*, *15*(1), 131–3. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/21276309

Kendall, E. A., Fofana, M. O., Dowdy, D. D. W., WHO, Dye, C., Garnett, G., … al., et. (2015).

Burden of transmitted multidrug resistance in epidemics of tuberculosis: a transmission

modelling analysis. *The Lancet Respiratory Medicine*, *3*(12), 963–972.

https://doi.org/10.1016/S2213-2600(15)00458-0

Klopper, M., Warren, R. M., Hayes, C., Gey van Pittius, N. C., Streicher, E. M., Müller, B., …

Trollip, A. P. (2013). Emergence and Spread of Extensively and Totally Drug-Resistant

Tuberculosis, South Africa. *Emerging Infectious Diseases*, *19*(3), 449–455.

https://doi.org/10.3201/eid1903.120246

Kobert, K., Stamatakis, A., & Flouri, T. (2016). Efficient Detection of Repeating Sites to

Accelerate Phylogenetic Likelihood Calculations. *Systematic Biology*, *66*(2), syw075.

https://doi.org/10.1093/sysbio/syw075

Kohl, T. A., Diel, R., Harmsen, D., Rothgänger, J., Walter, K. M., Merker, M., … Niemann, S.

(2014). Whole-genome-based Mycobacterium tuberculosis surveillance: a

standardized, portable, and expandable approach. *Journal of Clinical Microbiology*,

*52*(7), 2479–86. https://doi.org/10.1128/JCM.00567-14

Kohl, T. A., Utpatel, C., Schleusener, V., De Filippo, M. R., Beckert, P., Cirillo, D. M., &

Niemann, S. (2018). MTBseq: A comprehensive pipeline for whole genome sequence

analysis of Mycobacterium tuberculosis complex isolates. *In Preparation*.

Kozlov, A. (2017). RAxML-NG.

680 https://doi.org/https://zenodo.org/record/888146#.Wm8r2Uso9TY

681 Kurbatova, E. V., Cavanaugh, J. S., Dalton, T., S. Click, E., & Cegielski, J. P. (2013).

682 Epidemiology of Pyrazinamide-Resistant Tuberculosis in the United States, 1999–2009.

683 *Clinical Infectious Diseases*, *57*(8), 1081–1093. https://doi.org/10.1093/cid/cit452

684 Leaché, A. D., Banbury, B. L., Felsenstein, J., de Oca, A. nieto-M., Stamatakis, A., K., S., … J.,

685 W. (2015). Short Tree, Long Tree, Right Tree, Wrong Tree: New Acquisition Bias

686 Corrections for Inferring SNP Phylogenies. *Systematic Biology*, *64*(6), 1032–1047.

687 https://doi.org/10.1093/sysbio/syv053

688 Lew, J. M., Kapopoulou, A., Jones, L. M., & Cole, S. T. (2011). TubercuList – 10 years after.

689 *Tuberculosis*, *91*(1), 1–7. https://doi.org/10.1016/j.tube.2010.09.008

690 Lewis, P. O. (2001). A likelihood approach to estimating phylogeny from discrete

691 morphological character data. *Systematic Biology*, *50*(6), 913–25. Retrieved from

692 http://www.ncbi.nlm.nih.gov/pubmed/12116640

693 Médigue, C., Cole, S. T., Camus, J.-C., & Pryor, M. J. (2002). Re-annotation of the genome

694 sequence of Mycobacterium tuberculosis H37Rv. *Microbiology*, *148*(10), 2967–2973.

695 https://doi.org/10.1099/00221287-148-10-2967

696 Merker, M., Blin, C., Mona, S., Duforet-Frebourg, N., Lecher, S., Willery, E., … Wirth, T.

697 (2015). Evolutionary history and global spread of the Mycobacterium tuberculosis

698 Beijing lineage. *Nature Genetics*, *47*(3), 242–249. https://doi.org/10.1038/ng.3195

699 Merker, M., Kohl, T. A., Niemann, S., & Supply, P. (2017). The Evolution of Strain Typing in

700 the Mycobacterium tuberculosis Complex. In *Advances in experimental medicine and*

701 *biology* (Vol. 1019, pp. 43–78). https://doi.org/10.1007/978-3-319-64371-7_3

702 Miotto, P., Cabibbe, A. M., Feuerriegel, S., Casali, N., Drobniewski, F., Rodionova, Y., …

703 Cirillo, D. M. (2014). Mycobacterium tuberculosis pyrazinamide resistance

determinants: a multicenter study. *MBio*, *5*(5), e01819-14.

https://doi.org/10.1128/mBio.01819-14

Mokrousov, I., Chernyaeva, E., Vyazovaya, A., Sinkov, V., Zhuravlev, V., & Narvskaya, O.

(2016). Next-Generation Sequencing of Mycobacterium tuberculosis. *Emerging*

*Infectious Diseases*, *22*(6), 1127–9. https://doi.org/10.3201/eid2206.152051

Ngabonziza, J. C. S., Diallo, A. B., Tagliani, E., Diarra, B., Kadanga, A. E., Togo, A. C. G., …

group",  the "Union short M.-T. regimen study. (2017). Half of rifampicin-resistant

Mycobacterium tuberculosis complex isolated from tuberculosis patients in Sub-

Saharan Africa have concomitant resistance to pyrazinamide. *PLOS ONE*, *12*(10),

e0187211. https://doi.org/10.1371/journal.pone.0187211

Phelan, J. E., Coll, F., Bergval, I., Anthony, R. M., Warren, R., Sampson, S. L., … Clark, T. G.

(2016). Recombination in pe/ppe genes contributes to genetic variation in

Mycobacterium tuberculosis lineages. *BMC Genomics*, *17*, 151.

https://doi.org/10.1186/s12864-016-2467-y

R Core Team. (2017). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. Retrieved from http://www.r-project.org/

Ragheb, M. N., Ford, C. B., Chase, M. R., Lin, P. L., Flynn, J. L., & Fortune, S. M. (2013). The

mutation rate of mycobacterial repetitive unit loci in strains of M. tuberculosis from

cynomolgus macaque infection. *BMC Genomics*, *14*(1), 145.

https://doi.org/10.1186/1471-2164-14-145

Rambaut, A. (2016). Figtree. Retrieved May 3, 2017, from

http://tree.bio.ed.ac.uk/software/figtree/

Rambaut, A., Suchard, M. A., Xie, D., & Drummond, A. (2013). Tracer. Retrieved December

11, 2013, from http://beast.bio.ed.ac.uk/Tracer

Roetzer, A., Diel, R., Kohl, T. A., Rückert, C., Nübel, U., Blom, J., … Niemann, S. (2013). Whole

genome sequencing versus traditional genotyping for investigation of a Mycobacterium

tuberculosis outbreak: a longitudinal molecular epidemiological study. *PLoS Medicine*,

*10*(2), e1001387. https://doi.org/10.1371/journal.pmed.1001387

Roetzer, A., Schuback, S., Diel, R., Gasau, F., Ubben, T., di Nauta, A., … Niemann, S. (2011).

Evaluation of Mycobacterium tuberculosis typing methods in a 4-year study in

Schleswig-Holstein, Northern Germany. *Journal of Clinical Microbiology*, *49*(12), 4173–

8. https://doi.org/10.1128/JCM.05293-11

Said, H. M., Kushner, N., Omar, S. V, Dreyer, A. W., Koornhof, H., Erasmus, L., … Ismail, N. A.

(2016). A Novel Molecular Strategy for Surveillance of Multidrug Resistant Tuberculosis

in High Burden Settings. *PloS One*, *11*(1), e0146106.

https://doi.org/10.1371/journal.pone.0146106

Salathé, M., Bengtsson, L., Bodnar, T. J., Brewer, D. D., Brownstein, J. S., Buckee, C., …

Vespignani, A. (2012). Digital Epidemiology. *PLoS Computational Biology*, *8*(7),

e1002616. https://doi.org/10.1371/journal.pcbi.1002616

Sanoussi, C. N., Affolabi, D., Rigouts, L., Anagonou, S., & de Jong, B. (2017). Genotypic

characterization directly applied to sputum improves the detection of Mycobacterium

africanum West African 1, under-represented in positive cultures. *PLOS Neglected*

*Tropical Diseases*, *11*(9), e0005900. https://doi.org/10.1371/journal.pntd.0005900

Scott, A. N., Menzies, D., Tannenbaum, T.-N., Thibert, L., Kozak, R., Joseph, L., … Behr, M. A.

(2005). Sensitivities and specificities of spoligotyping and mycobacterial interspersed

repetitive unit-variable-number tandem repeat typing methods for studying molecular

epidemiology of tuberculosis. *Journal of Clinical Microbiology*, *43*(1), 89–94.

https://doi.org/10.1128/JCM.43.1.89-94.2005

Sengstake, S., Bergval, I. L., Schuitema, A. R., Beer, J. L., Phelan, J., Zwaan, R., … Anthony, R.

M. (2017). Pyrazinamide resistance-conferring mutations in pncA and the transmission

of multidrug resistant TB in Georgia. *BMC Infectious Diseases 2017 17:1*, *17*(1), 491.

https://doi.org/10.1186/s12879-017-2594-3

Shah, N. S., Auld, S. C., Brust, J. C. M., Mathema, B., Ismail, N., Moodley, P., … Gandhi, N. R.

(2017). Transmission of Extensively Drug-Resistant Tuberculosis in South Africa. *New*

*England Journal of Medicine*, *376*(3), 243–253.

https://doi.org/10.1056/NEJMoa1604544

Small, P. M., Hopewell, P. C., Singh, S. P., Paz, A., Parsonnet, J., Ruston, D. C., … Schoolnik, G.

K. (1994). The Epidemiology of Tuberculosis in San Francisco -- A Population-Based

Study Using Conventional and Molecular Methods. *New England Journal of Medicine*,

*330*(24), 1703–1709. https://doi.org/10.1056/NEJM199406163302402

Statisticat. (2016). LaplacesDemon: Complete Environment for Bayesian Inference.

Bayesian-Inference.com. R package version 16.0.1. Retrieved from

https://web.archive.org/web/20150206004624/http://www.bayesian-

inference.com/software

Stucki, D., Ballif, M., Egger, M., Furrer, H., Altpeter, E., Battegay, M., … Fenner, L. (2016).

Standard Genotyping Overestimates Transmission of Mycobacterium tuberculosis

among Immigrants in a Low-Incidence Country. *Journal of Clinical Microbiology*, *54*(7),

1862–70. https://doi.org/10.1128/JCM.00126-16

Sukumaran, J., & Holder, M. T. (2010). DendroPy: a Python library for phylogenetic

computing. *Bioinformatics*, *26*(12), 1569–1571.

https://doi.org/10.1093/bioinformatics/btq228

Supply, P., Allix, C., Lesjean, S., Cardoso-Oelemann, M., Rusch-Gerdes, S., Willery, E., … van

Soolingen, D. (2006). Proposal for Standardization of Optimized Mycobacterial

Interspersed Repetitive Unit-Variable-Number Tandem Repeat Typing of

Mycobacterium tuberculosis. *Journal of Clinical Microbiology*, *44*(12), 4498–4510.

https://doi.org/10.1128/JCM.01392-06

Supply, P., Magdalena, J., Himpens, S., & Locht, C. (1997). Identification of novel intergenic

repetitive units in a mycobacterial two-component system operon. *Molecular*

*Microbiology*, *26*(5), 991–1003. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/9426136

Tessema, B., Nabeta, P., Valli, E., Albertini, A., Collantes, J., Lan, N. H., … Dolinger, D. L.

(2017). FIND Tuberculosis Strain Bank: a Resource for Researchers and Developers

Working on Tests To Detect Mycobacterium tuberculosis and Related Drug Resistance.

*Journal of Clinical Microbiology*, *55*(4), 1066–1073.

https://doi.org/10.1128/JCM.01662-16

Thierry, D., Cave, M. D., Eisenach, K. D., Crawford, J. T., Bates, J. H., Gicquel, B., & Guesdon,

J. L. (1990). IS6110, an IS-like element of Mycobacterium tuberculosis complex. *Nucleic*

*Acids Research*, *18*(1), 188. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/2155396

Trauer, J. M., Denholm, J. T., & McBryde, E. S. (2014). Construction of a mathematical model

for tuberculosis transmission in highly endemic regions of the Asia-pacific. *Journal of*

*Theoretical Biology*, *358*, 74–84. https://doi.org/10.1016/j.jtbi.2014.05.023

van Embden, J. D., Cave, M. D., Crawford, J. T., Dale, J. W., Eisenach, K. D., Gicquel, B., … al.,

et. (1993). Strain identification of Mycobacterium tuberculosis by DNA fingerprinting:

recommendations for a standardized methodology. *Journal of Clinical Microbiology*,

*31*(2), 406–9. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8381814

Vluggen, C., Soetaert, K., Groenen, G., Wanlin, M., Spitaels, M., Arrazola de Oñate, W., …

Mathys, V. (2017). Molecular epidemiology of Mycobacterium tuberculosis complex in

Brussels, 2010–2013. *PLOS ONE*, *12*(2), e0172554.

https://doi.org/10.1371/journal.pone.0172554

Walker, T. M., Ip, C. L. C., Harrell, R. H., Evans, J. T., Kapatai, G., Dedicoat, M. J., … Peto, T. E.

A. (2013). Whole-genome sequencing to delineate Mycobacterium tuberculosis

outbreaks: a retrospective observational study. *The Lancet Infectious Diseases*, *13*(2),

137–46. https://doi.org/10.1016/S1473-3099(12)70277-3

Walker, T. M., Lalor, M. K., Broda, A., Ortega, L. S., Morgan, M., Parker, L., … Conlon, C. P.

(2014). Assessment of Mycobacterium tuberculosis transmission in Oxfordshire, UK,

2007-12, with whole pathogen genome sequences: an observational study. *The Lancet.*

*Respiratory Medicine*, *2*(4), 285–292. https://doi.org/10.1016/S2213-2600(14)70027-X

Walker, T. M., Merker, M., Knoblauch, A. M., Helbling, P., Schoch, O. D., van der Werf, M. J.,

… Witschi, M. (2018). A cluster of multidrug-resistant Mycobacterium tuberculosis

among patients arriving in Europe from the Horn of Africa: a molecular epidemiological

study. *The Lancet Infectious Diseases*. https://doi.org/10.1016/S1473-3099(18)30004-5

Warren, R. M., Streicher, E. M., Sampson, S. L., van der Spuy, G. D., Richardson, M., Nguyen,

D., … van Helden, P. D. (2002). Microevolution of the direct repeat region of

Mycobacterium tuberculosis: implications for interpretation of spoligotyping data.

*Journal of Clinical Microbiology*, *40*(12), 4457–65. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/12454136

WHO. (2015). *Guidelines for surveillance of drug resistance in tuberculosis* (5th ed.). Geneva.

WHO. (2018). *Global tuberculosis report 2017*. Retrieved from

http://www.who.int/tb/publications/global_report/archive/en/

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Retrieved from http://ggplot2.org

Wirth, T., Hildebrand, F., Allix-Béguec, C., Wölbeling, F., Kubica, T., Kremer, K., … Niemann,

S. (2008). Origin, Spread and Demography of the Mycobacterium tuberculosis Complex.

*PLoS Pathogens*, *4*(9), e1000160. https://doi.org/10.1371/journal.ppat.1000160

Witney, A. A., Cosgrove, C. A., Arnold, A., Hinds, J., Stoker, N. G., & Butcher, P. D. (2016).

Clinical use of whole genome sequencing for Mycobacterium tuberculosis. *BMC*

*Medicine*, *14*(1), 46. https://doi.org/10.1186/s12916-016-0598-2

Wyllie, D., Davidson, J., Walker, T., Rathod, P., Peto, T., Robinson, E., … Campbell, C. (2018).

A quantitative evaluation of MIRU-VNTR typing against whole-genome sequencing for

identifying Mycobacterium tuberculosis transmission: A prospective observational

cohort study. *BioRxiv*, 252734. https://doi.org/10.1101/252734

Xia, E., Teo, Y.-Y., & Ong, R. T.-H. (2016). SpoTyping: fast and accurate in silico

Mycobacterium spoligotyping from sequence reads. *Genome Medicine*, *8*(1), 19.

https://doi.org/10.1186/s13073-016-0270-7

Xu, P., Wu, J., Yang, C., Luo, T., Shen, X., Zhang, Y., … Gao, Q. (2016). Prevalence and

transmission of pyrazinamide resistant Mycobacterium tuberculosis in China.

*Tuberculosis*, *98*, 56–61. https://doi.org/10.1016/j.tube.2016.02.008

Zignol, M., Cabibbe, A. M., Dean, A. S., Glaziou, P., Alikhanova, N., Ama, C., … Raviglione, M.

C. (2018). Genetic sequencing for surveillance of drug resistance in tuberculosis in

highly endemic countries: a multi-country population-based surveillance study. *The*

*Lancet Infectious Diseases*. https://doi.org/10.1016/S1473-3099(18)30073-2

Zignol, M., Dean, A. S., Alikhanova, N., Andres, S., Cabibbe, A. M. A., Cirillo, D. M., … Valafar,

F. (2016). Population-based resistance of Mycobacterium tuberculosis isolates to

pyrazinamide and fluoroquinolones: results from a multicountry surveillance project.

*The Lancet Infectious Diseases*. https://doi.org/10.1016/S1473-3099(16)30190-6

Supplementary legends

<u>Supplementary Table 1: Spoligotyping patterns derived from membrane- and genome-based</u>

<u>methods</u>

Spoligotyping patterns were estimated from both membrane (Mem-Spo) and genome (Gen-

Spo) approaches. Patterns are shown from both methods per isolates with the number of

mismatches between patterns recorded.

| Isolate | Mem-Spoligotype | Gen-Spoligotype | Mismatches |
|---|---|---|---|
| DRC-052577 | 111111100000111111111101000000000010001111 | 111111100000111111111101000000000110001111 | 1 |
| DRC-052578 | 111111100000000000000000111111100001111111 | 111111100000000000000000111111100001111111 | 0 |
| DRC-052654 | 111111111111111111000001111111100001111111 | 110111111111111111101111111011100001101111 | 7 |
| DRC-052667 | 111111110001111111000000000000000000001100 | 111111111111000011100000000000000000111100 | 9 |
| DRC-052750 | 111111100000111111110110000000000010001111 | 111111100000111111110100000000000110001111 | 3 |
| DRC-052956 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-052958 | 111111111111111111111111110111100001110111 | 111111111111111111111111110111100001110111 | 0 |
| DRC-052959 | 111111111111111111111111111111100001111111 | 111111111111111111111111111111100001111111 | 1 |
| DRC-052964 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-060116 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-060118 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-060120 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-060131 | 110111111111111111000011111111100001111111 | 110111111111111111000011111111100001111111 | 0 |
| DRC-060237 | 111111111111111111111111111111100001111111 | 111111111111111111111111111111100001111111 | 0 |
| DRC-060244 | 000000000000000000000000111111100001111111 | 000000000000000000000000111111100001110111 | 3 |
| DRC-060248 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-060253 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-060260 | 111111000000000000000000111111100001011111 | 111111000000000000000000111111100101111111 | 2 |
| DRC-060360 | 111110100000111111111110000000000010000011 | 111111100000111111111110000000000110000011 | 2 |
| DRC-060361 | 110111111111111111000011111111100001111111 | 110111111111111111000011111111100001111111 | 0 |
| DRC-060362 | 111111100000111111111101100000000010001111 | 111111100000111111111101000000000110001111 | 2 |
| DRC-060366 | 111111000000000000000000111111100001011111 | 111111000000000000000000111111100001111111 | 1 |
| DRC-060375 | 110000001111111011100001111111100001111111 | 110000001111111011100001111111100001111111 | 2 |
| DRC-060376 | 110111100011111111110000111111100000011111 | 110111111111111111110000111111100001111111 | 5 |

| | | | |
|---|---|---|---|
| DRC-060381 | 11111111000111111110000000000000000000011111 | 11111111000111111110000000000000000000011111 | 0 |
| DRC-060382 | 11111100000000000000000001111111100001111111 | 11111100000000000000000001111111100001111111 | 0 |
| DRC-060514 | 11000000111111101110000111111110000111111 | 11000000111111101110000111111110000111111 | 0 |
| DRC-060526 | 11111111000111111110000000000000000000011111 | 11111111000111111110000000000000000000011111 | 0 |
| DRC-060527 | 11111111111110000111100001111111100001011111 | 11111111111110000111100001111111100001011111 | 0 |
| DRC-060528 | 11111100000000000000000001111111100001011111 | 11111100000000000000000001111111100001111111 | 1 |
| DRC-060549 | 10000011111111111011111111111100001110111 | 10000011111111111111111111111111100001111111 | 2 |
| DRC-060562 | 11111111111111111111111111111111100001000011 | 11111111111111111111111111111111100001000011 | 0 |
| DRC-060565 | 11111111111111111111111111111111100001110111 | 11111111111111111111111111111111100001110111 | 0 |
| DRC-060756 | 11111100000000000000000001111111100001111111 | 11111100000000000000000001111111100001111111 | 0 |
| DRC-060757 | 11111111111111111111111111101111100001110111 | 11111111111111111111111111101111100001110111 | 0 |
| DRC-060772 | 11111110000011111111101000000000010001111 | 11111110000011111111101000000000110001111 | 1 |
| DRC-060778 | 11111111000111111110000000000000000000011111 | 11111111000111111110000000000000000000011111 | 0 |
| DRC-060856 | 11111111111100001100000000000000000111100 | 11111111111100001100000000000000000111100 | 0 |
| DRC-060858 | 11111111111111111110000111111110000101111 | 11111111111111111111111111101111100001110111 | 7 |
| DRC-060859 | 11111111111111111110000111111110000111111 | 11111111111111111110000111111110000111111 | 0 |
| DRC-061069 | 11111110000011111111111000000000010001111 | 11111110000011111111111000000000110001111 | 1 |
| DRC-061123 | 11111111111111111110000111111110000111111 | 11111111111111111110000111111110000111111 | 0 |
| DRC-061124 | 11011111111111111110000111111110000111111 | 11111111111111111110000111111110000111111 | 1 |
| DRC-061127 | 11111110000000000000000001111111100001111111 | 11111110000000000000000001111111100001111111 | 0 |
| DRC-061136 | 11111100000000000000000001111111100001011111 | 11111100000000000000000001111111100001111111 | 1 |
| DRC-061137 | 11111100000000000000000001111111100001011111 | 11111100000000000000000001111111100001111111 | 1 |
| DRC-061138 | 11111111000111111110000000000000000000011111 | 11111111000111111110000000000000000000011111 | 0 |
| DRC-061142 | 11111111111111111110000111111110000111111 | 11111111111111111110000111111110000111111 | 0 |
| DRC-061431 | 11111111000111111110000000000000000000011111 | 11111111000111111110000000000000000000011111 | 0 |
| DRC-061493 | 11111111111111111111111111111111100001110111 | 11111111110111111111111111111111100001110111 | 1 |
| DRC-062295 | 11111111111111000011111111111111100001110111 | 11111111111111000011111111111111100001110111 | 0 |
| DRC-062298 | 11111111111100001100000000000000000111100 | 11111111111100001100000000000000000111100 | 0 |
| DRC-062303 | 11000000111111101110000111111110000111111 | 11000000111111101110000111111110000111111 | 2 |
| DRC-062444 | 11111111000111111110000000000000000000011111 | 11111111000111111110000000000000000000011111 | 0 |
| DRC-062446 | 11111111111111111110000111101110000111111 | 11111111111111111110000111101110000111111 | 0 |
| DRC-062447 | 11000000111111101110000111111110000111111 | 11000000111111101110000111111110000111111 | 0 |
| DRC-062495 | 11111111111111111111111111111111100001111111 | 11111111111111111111111111111111100001111111 | 0 |
| DRC-062502 | 11111111111111111110000111111110000101111 | 11111111111111111110000111111110000101111 | 0 |
| DRC-062503 | 11111111111111111111111111111111100001111111 | 11111111111111111111111111111111100001110111 | 1 |

| | | | |
|---|---|---|---|
| DRC-062822 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-062827 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-062840 | 111111111111111111111111111111100001111111 | 111111111111111111111111111111100001111111 | 0 |
| DRC-062841 | 110111111111111111100000110111100001111111 | 111111111111111111100000110111100001111111 | 2 |
| DRC-062844 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-062849 | 111111111101111111100001111111100001111111 | 111111111101111111100001111111100001111111 | 0 |
| DRC-063522 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-063540 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-063541 | 111111100000000000000001111111100001111111 | 111111100000000000000001111111100001111111 | 0 |
| DRC-063545 | 111111100000111111111010000000000010001111 | 111111100000111111111010000000000110001111 | 1 |
| DRC-063556 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-063557 | 111111000000000000000001111111100001111111 | 111111000000000000000001111111100001111111 | 0 |
| DRC-063645 | 111111111111111111100001111111100001111111 | 111111111110111111100001110111100001111111 | 3 |
| DRC-063667 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-063803 | 111111111111111111100000110111100001111111 | 111111111111111111100000110111100001111111 | 1 |
| DRC-063887 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-063888 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-063889 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-063896 | 111111111111111111111111111111100001111111 | 111111111111111111111111111111100001111111 | 0 |
| DRC-063898 | 111111111111111111100001111011100001111111 | 111111111111111111100001111011100001111111 | 0 |
| DRC-064081 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 1 |
| DRC-064085 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-064090 | 111111000000000000000001111111100001111111 | 111111000000000000000001111111100001111111 | 0 |
| DRC-064101 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-064208 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-070012 | 101111111010111111111111111111100001110111 | 101111111010111111111111111111100001110111 | 1 |
| DRC-070038 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-070043 | 111000011111111111110000000000111111111 | 111000011111111111110000000000111111111 | 0 |
| DRC-070049 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-070777 | 111111111111111111100001111011100001111111 | 111111111111111111100001111011100001111111 | 0 |
| DRC-070779 | 111111000000000000000001111111100001111111 | 111111000000000000000001111111100001111111 | 0 |
| DRC-070781 | 111111111111011111111111111111100001110111 | 111111111111011111111111111111100001110111 | 0 |
| DRC-070783 | 111111111111111111100000000000000001111111 | 111111111110111111110000000000000001111111 | 1 |
| DRC-070786 | 111111100000111111111010000000000010001111 | 111111100000111111111010000000000110001111 | 1 |
| DRC-070788 | 111100001111000011111111111111100001111111 | 111100001111000011111111111111100001111111 | 0 |

| DRC-070790 | 11111111111111111111000011111111100001111111 | 11111111111111111111000011111111100001111111 | 0 |
| DRC-070792 | 11111111111111111111100000001100001111111 | 11111111111111111111100000001100001111111 | 0 |
| DRC-070794 | 10000011111111111111111111111111100001111111 | 10000011111111111111111111111111100001111111 | 0 |
| DRC-070804 | 11111111111111111110000111111100001011111 | 11111111111111111110000111111100001011111 | 0 |
| DRC-070807 | 1111111111111111111111111011110000111011 | 1111111111111111111111111011110000111011 | 0 |
| DRC-070808 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-070809 | 11011111111111111110000111111110000111111 | 11011111111111111110000111111110000111111 | 0 |
| DRC-071052 | 11111100000000000000000011111110000111111 | 11111100000000000000000011111110000111111 | 0 |
| DRC-071057 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-071074 | 11111111111111111110000111111100001111111 | 11111111111111111110000111111100001111111 | 0 |
| DRC-071075 | 11111100000000000000000011111110000111111 | 11111100000000000000000011111110000111111 | 0 |
| DRC-071076 | 11111110000011111111010000000000010001111 | 11111110000011111111010000000000110001111 | 1 |
| DRC-071458 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-071460 | 111111111111111111111000111111100001110111 | 111111111111111111111000111111100001110111 | 0 |
| DRC-071472 | 11111111111111111110000111111100001111111 | 11111111111111111110000111111100001111111 | 0 |
| DRC-071479 | 1111111111111111111111111111101000011111111 | 1111111111111111111111111111111100001111111 | 1 |
| DRC-071686 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-071691 | 1101111111110111110000111111110000111111 | 1111111111011111100001111111100001111111 | 3 |
| DRC-071696 | 11111111111101111111111111111111100001110111 | 11111111111101111111111111111111100001110111 | 0 |
| DRC-071697 | 111111111111111111100011111010000111111 | 11111111111111111111000111111100001111111 | 1 |
| DRC-071712 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-071960 | 11011111111111111110000111111100001111111 | 11011111111111111110000111111100001111111 | 0 |
| DRC-071968 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-071972 | 11111111111100001100000000000000000111100 | 11111111111100001100000000000000000111100 | 0 |
| DRC-071973 | 11111111000111111100000000000000000011111 | 11111111000111111100000000000000000011111 | 0 |
| DRC-071975 | 11111111111101111111111111111111100001110111 | 11111111111101111111111111111111100001110111 | 0 |
| DRC-071988 | 1111111111111111110000001101110000001111 | 1111111111111111110000011101110000111111 | 3 |
| DRC-071990 | 11111100000000000000000011111110000111111 | 11111100000000000000000011111110000111111 | 0 |
| DRC-071998 | 11111111000111111100000000000000000011111 | 11111111000111111100000000000000000011111 | 0 |
| DRC-072061 | 11111100000000000000000011111110000111111 | 11111100000000000000000011111110000111111 | 0 |
| DRC-072379 | 11111111000111111100000000000000000011111 | 11111111000111111100000000000000000011111 | 0 |
| DRC-072384 | 1111101000001111111111010000000010000011 | 1111111000001111111111100000000000110000011 | 3 |
| DRC-072390 | 11011111111111111110000111111100001111111 | 11011111111111111110000111111100001111111 | 0 |
| DRC-072603 | 11111111111111101111111111111100001111111 | 11111111111111101111111111111100001111111 | 0 |
| DRC-072604 | 111110000000000000000000111111100001110111 | 111110000000000000000000111111100001110111 | 0 |

| | | | |
|---|---|---|---|
| DRC-072605 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-072606 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-072608 | 111111111110111111111111111111100001110111 | 111111111110111111111111111111100001110111 | 0 |
| DRC-072609 | 111111111111111111111111111111100001110111 | 110111111110111111100001111111100001111111 | 7 |
| DRC-072613 | 110111111101111111100001111111100001111111 | 111111111101111111100001111111100001111111 | 1 |
| DRC-072615 | 111111100000000000000000111111100001111111 | 111111100000000000000000111111100001111111 | 0 |
| DRC-072618 | 111111111111000011000000000000000000111100 | 111111111111000011000000000000000000111100 | 0 |
| DRC-072625 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-072818 | 110111111111111111100001111111100001111111 | 111111111111000011000000000000000000111100 | 17 |
| DRC-072819 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-072820 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-072824 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-072825 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-072827 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-072829 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-072830 | 111111111110000110000000000000000000111100 | 111111111110000110000000000000000000111100 | 0 |
| DRC-072834 | 111111111110111111111111111111100001110111 | 111111111110111111111111111111100001110111 | 0 |
| DRC-072836 | 111111111111111111101100111111100001010111 | 111111111111111111101100111111100001110111 | 1 |
| DRC-072837 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-072943 | 111111111111000011000000000000000000111100 | 111111111111000011000000000000000000111100 | 0 |
| DRC-072945 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-072957 | 110111111111011111100001111111100001111111 | 111111111111111111111111111111100001110111 | 7 |
| DRC-072959 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-072963 | 111111111111111101111111110111100001111111 | 111111111111111111111111110111100001111111 | 1 |
| DRC-072974 | 111111111111111111100001111111100001011111 | 111111111111111111100001111111100001011111 | 0 |
| DRC-072986 | 111111111111111111100001100001100001011111 | 111111111111111111100001100001100001111111 | 1 |
| DRC-072988 | 111111000000000000000000111111100001110111 | 111111000000000000000000111111100001110111 | 0 |
| DRC-073093 | 111111100000111111111011000000000010001111 | 111111100000111111111010000000000110001111 | 2 |
| DRC-073099 | 111110100001111111111111000000000010000011 | 111111100001111111111110000000000110000011 | 3 |
| DRC-073168 | 111111111110000110000000000000000000111100 | 111111111110000110000000000000000000111100 | 0 |
| DRC-073169 | 111111111111111111100001111011100001110111 | 111111111111111111100001111011100001110111 | 0 |
| DRC-073170 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-073173 | 110111111101111111100001111111100001111111 | 110111111101111111100001111111100001111111 | 0 |
| DRC-073175 | 110111111111111111111111111111100001110111 | 110111111111111111111111111111100001110111 | 0 |
| DRC-073183 | 111111111111111101111111111111100001111111 | 111111111111111111111111111111100001111111 | 1 |

| | | | |
|---|---|---|---|
| DRC-073186 | 1111111111111111111111111110111100001110111 | 1111111111111111111111111110111100001110111 | 0 |
| DRC-073413 | 1111111111111111111100001110110001111111 | 1111111111111101110000111011100001111111 | 1 |
| DRC-073414 | 0000000000000000000000001111111100001110111 | 0000000000000000000000001111111100001110111 | 0 |
| DRC-073415 | 0000000000000000000000001111111100001110111 | 0000000000000000000000001111111100001110111 | 0 |
| DRC-073420 | 11111111111111111110000111111110000111111 | 11111111111111111110000111111110000111111 | 0 |
| DRC-073422 | 11111111111111111110000111111110000111111 | 11111111111111111110000111111110000111111 | 0 |
| DRC-073423 | 11111111111111110111111111111100001100001 | 11111111111111110111111111111100001100001 | 0 |
| DRC-073424 | 11111111111111111110000111111110000111111 | 11111111110111111110000111111110000111111 | 1 |
| DRC-073429 | 11111100000000000000000111111100001111111 | 11111100000000000000000111111100001111111 | 0 |
| DRC-073430 | 1111111111111111111111111111110000111 0111 | 1111111111111111111111111111110000111 0111 | 0 |
| DRC-073432 | 1111111111111111111111111111110000111 0111 | 111111111101111111111111111111110000111 0111 | 1 |
| DRC-073434 | 11111111111011110111111111111000011 10111 | 11111111111011110111111111111000011 10111 | 0 |
| DRC-073443 | 1111111111110000111000000000000000000111100 | 1111111111110000111000000000000000000111100 | 0 |
| DRC-073515 | 11111100000000000000000111111100001111111 | 11111100000000000000000111111100001111111 | 0 |
| DRC-073712 | 11000000111111101110000111111100001111111 | 11000000111111101110000111111100001111111 | 0 |
| DRC-073716 | 1101111111111111111000011111110000111 1111 | 1101111111111111111000011111110000111 1111 | 0 |
| DRC-080008 | 11111110001111111000000000000000000011111 | 11111110001111111000000000000000000011111 | 0 |
| DRC-080016 | 1111111111111101111111111111111111000011 10111 | 1111111111111111111111111111111111000011 10111 | 1 |
| DRC-080021 | 11111100000000000000000111111100001111111 | 11111100000000000000000111111100001111111 | 0 |
| DRC-080028 | 11111111000111111100000000000000000011111 | 11111111000111111100000000000000000011111 | 0 |
| DRC-080040 | 11111100000000000000000111111100001111111 | 11111100000000000000000111111100001111111 | 0 |
| DRC-080046 | 00000000000000000000000111111 1100001110111 | 00000000000000000000000111111 1100001110111 | 0 |
| DRC-080545 | 111111111111111111110001111111000011 1111 | 111111111111111111110001111111000011 1111 | 0 |
| DRC-080548 | 11111110000011111111011000000000010001111 | 11111110000011111111010000000000110001111 | 3 |
| DRC-080552 | 11111111101111111110000111111110000101 1111 | 11111111101111111110000111111110000101 1111 | 0 |
| DRC-080553 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-080561 | 11111111111111111110000111111100001011111 | 11111111111111111110000111111100001011111 | 0 |
| DRC-080567 | 11111111111111111111111110111100001110111 | 11111111111111111111111110111100001110111 | 0 |
| DRC-081290 | 11111111101111111111111111111110000111 1111 | 11111111101111111110000111111100001111111 | 4 |
| DRC-081527 | 1111111111111111110000111111110000111 1111 | 1111111111111111110000111111110000111 1111 | 0 |
| DRC-081585 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-081588 | 11000000111111101110000111111100001111111 | 11000000111111101110000111111100001111111 | 0 |
| DRC-081589 | 1110111111110000111000000000000000000111100 | 1110111111110000111000000000000000000111100 | 0 |
| DRC-081592 | 11111111111111011110000111111110000111 1111 | 11111111111111111110000111111110000111 1111 | 1 |
| DRC-081593 | 11111111111111111111111111111100001110111 | 11111111111111111111111111111100001110111 | 0 |

| | | | |
|---|---|---|---|
| DRC-081595 | 111111111111111111111111111111111100001110111 | 111111111111111111111111111111111100001110111 | 0 |
| DRC-081600 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-081602 | 1111111111110111111111111111111100001110111 | 1111111111110111111111111111111100001110111 | 0 |
| DRC-081610 | 1111111100111111111111111111111100001111111 | 1111111100111111111111111111111100001111111 | 0 |
| DRC-081622 | 1111111111111111110000110000110000111111 | 1111111111111111110000110000110000111111 | 0 |
| DRC-081722 | 1101111111111111111100001111111100001111111 | 1101111111111111111100001111111100001111111 | 0 |
| DRC-081873 | 1111111111111111111100001111111100001011111 | 1111111111111111111100001111111100001011111 | 0 |
| DRC-081879 | 1111111111111111011111111111111100001100001 | 1111111111111111011111111111111100001100001 | 0 |
| DRC-081881 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-081888 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-081985 | 1111111111111111011111111111111100001111111 | 1111111111111111011111111111111100001111111 | 0 |
| DRC-081986 | 1111111111110101111111111111111100001110111 | 11111111111011111111111111111111100001110111 | 1 |
| DRC-081989 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-081990 | 000000000000000000000000000000000111111111 | 000000000000000000000000000000000111111111 | 0 |
| DRC-082120 | 11111111111111111111111111111111100001110111 | 11111111111111111111111111111111100001110111 | 0 |
| DRC-082121 | 11111111111111111111111111111111100001110111 | 11111111111111111111111111111111100001110111 | 0 |
| DRC-082130 | 11111111111111111110000110000110000110000110001001111 | 01111111111111111110000110000110000110000110001001111 | 1 |
| DRC-082131 | 1111111111111111011111111111111100001111111 | 1111111111111111011111111111111100001111111 | 0 |
| DRC-082317 | 1111111111111111111011001111111100001010111 | 11111111111111111111111111111111111100001110111 | 4 |
| DRC-082318 | 111111000000000000000000111111100001111111 | 111111000000000000000000111111100001111111 | 0 |
| DRC-082327 | 1111111111111011111111111110111100001110111 | 1111111111111111111111111110111100001110111 | 1 |
| DRC-082328 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-082330 | 1101111111111111111100001111111100001111111 | 1101111111111111111100001111111100001111111 | 0 |
| DRC-082584 | 11111111111111111111111111111111100001110111 | 11111111111111111111111111111111100001110111 | 0 |
| DRC-082592 | 1111111111111111111011001111111100001010111 | 1111111111111111111011001111111100001110111 | 1 |
| DRC-082639 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-082656 | 1111111111111111111100001111011100001111111 | 1111111111111111111100001111011100001111111 | 0 |
| DRC-082700 | 1111111111111111111100001111111100001111111 | 1111111111111111111100001111111100001111111 | 0 |
| DRC-082919 | 1100000011111110111000011111111100001111111 | 1100000011111110111000011111111100001111111 | 0 |
| DRC-082923 | 11111110000011111111101100000000010001111 | 11111110000011111111101000000000110001111 | 2 |
| DRC-082939 | 1111111111111111111100001111111100001111111 | 1111111111111111111100001111111100001111111 | 0 |
| DRC-082940 | 111111110001111111000000000000000000011111 | 111111110001111111000000000000000000011111 | 0 |
| DRC-082945 | 1111111111111111111100001111111100001111111 | 1111111111111111111100001111111100001111111 | 0 |
| DRC-082946 | 11111111111111111111111111111111100001110111 | 11111111111111111111111111111111100001110111 | 0 |
| DRC-082948 | 1111111111111111111100001111011100001110111 | 1111111111111111111100001111011100001110111 | 0 |

| | | | |
|---|---|---|---|
| DRC-082949 | 1111111111111101111111111111111100001111111 | 1111111111111111111111111111111100001111111 | 1 |
| DRC-083047 | 1111111111111111111000011110111000011111111 | 1111111111111111111000011110111000011111111 | 0 |
| DRC-083048 | 1111111111111111111000001101110000111111111 | 1111111111111111111000001101110000111111111 | 1 |
| DRC-083050 | 1111111111111111111000011110111000010011111 | 1111111111111111111000011110111000010011111 | 0 |
| DRC-083051 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-083052 | 1111111111111111111111000111111100001111111 | 1111111111111111111111000111111100001111111 | 0 |
| DRC-083066 | 1111111111110000110000000000000000000111100 | 1111111111110000110000000000000000000111100 | 0 |
| DRC-083067 | 1111111111110111111111111111111100001110111 | 1111111111110111111111111111111100001110111 | 0 |
| DRC-083190 | 1111111111111111111000011110111000011111111 | 1111111111111111111000011111111100001111111 | 1 |
| DRC-083191 | 1111111111111111111000011110111000011111111 | 1111111111111111111000011110111000011111111 | 0 |
| DRC-083200 | 1111110000000000000000011111111000011111111 | 1111110000000000000000011111111000011111111 | 0 |
| DRC-083201 | 1000001111111111110000011111111000011111111 | 1000001111111111111111111111111100001111111 | 5 |
| DRC-083202 | 1111111111111111111000011110111000010011111 | 1111111111111111111000011110111000010011111 | 0 |
| DRC-083204 | 1111111100011111110000000000000000000011111 | 1111111100011111110000000000000000000011111 | 0 |
| DRC-083226 | 1111110101111111111111110000000000000000000 | 1111110101111111111111110000000000000000000 | 1 |
| DRC-083250 | 1111111111111111111111111111110100001111111 | 1111111111111111111111111111111100001111111 | 1 |
| DRC-083736 | 1111111111111111111000011111111000011111111 | 1111111111111111111000011111111000011111111 | 0 |
| DRC-083801 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-084109 | 1111111111111111111000011111111000011111111 | 1111111111111111111000011111111000011111111 | 0 |
| DRC-084204 | 1101111101111111111000011111111000011110111 | 1101111101111111111000011111111000011110111 | 0 |
| DRC-084210 | 1111111100011111110000000000000000000011111 | 1111111100011111110000000000000000000011111 | 0 |
| DRC-090010 | 1111111111111111111000011110111000011111111 | 1111111111111111111000011110111000011111111 | 0 |
| DRC-090012 | 1101111111111111111000011111111000011111111 | 1101111111111111111000011111111000011111111 | 0 |
| DRC-090016 | 1111111111111111111000011110111000011111111 | 1111111111111111111000011110111000011111111 | 0 |
| DRC-090022 | 1111111111101111111111111111111100001110111 | 1111111111101111111111111111111100001110111 | 0 |
| DRC-090037 | 1111111111111111111000011000011000011111111 | 1111111111111111111000011000011000011111111 | 0 |
| DRC-090066 | 1111111000001111111110110000000000010001111 | 1111111000001111111110100000000000110001111 | 2 |
| DRC-090069 | 1111111111111111111000011111111000011111111 | 1111111111111111111000011111111000011111111 | 0 |
| DRC-090077 | 1111111111111111111000011111111000011111111 | 1111111111111111111000011111111000011111111 | 0 |
| DRC-090078 | 1111111111111111111000011111111000010111111 | 1111111111111111111000011111111000010111111 | 0 |
| DRC-090165 | 1111111111111111111111111111111100001110111 | 1111111111111111111111111111111100001110111 | 0 |
| DRC-090172 | 1111111111110000110000000000000000000111100 | 1111111111110000110000000000000000000111100 | 0 |
| DRC-090299 | 1101111111111111111000011111111000011111111 | 1101111111111111111000011111111000011111111 | 0 |
| DRC-090308 | 1111110000000000000000011111111000011111111 | 1111110000000000000000011111111000011111111 | 0 |
| DRC-090309 | 1111111111110111111000011000011000011111111 | 1111111111110111111000011000011000011111111 | 0 |

| | | | |
|---|---|---|---|
| DRC-090310 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-090432 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-090451 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-090452 | 111111110001111111100000000000000000011111 | 111111110001111111100000000000000000011111 | 0 |
| DRC-090456 | 111111111111111111111111110111100001110111 | 111111111111111111111111110111100001110111 | 0 |
| DRC-090702 | 111111000000000000000011111110000011111111 | 111111000000000000000011111110000011111111 | 0 |
| DRC-090708 | 111111111111111101111111111111100001100001 | 111111111111111101111111111111100001100001 | 0 |
| DRC-090710 | 111111111110000110000000000000000000111100 | 111111111110000110000000000000000000111100 | 0 |
| DRC-090909 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-090913 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-090947 | 111111110001111111100000000000000000011111 | 111111110001111111100000000000000000011111 | 0 |
| DRC-090976 | 111111111111111111100001100001100001111111 | 111111111111111111100001100001100001111111 | 0 |
| DRC-090977 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-090999 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-091003 | 111111111111111111111111110111100001110111 | 111111110111111111111111110111100001110111 | 1 |
| DRC-091004 | 111111000000000000000011111110000011111111 | 111111000000000000000011111110000011111111 | 0 |
| DRC-091005 | 111111111110101111111111111111100001111111 | 111111000000000000000011111110000011111111 | 16 |
| DRC-091168 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-091171 | 111111111111111111100001111111100001111111 | 111111111111111111100001111011100001111111 | 1 |
| DRC-091172 | 111111111111111111111111110111100001110111 | 111111111111111111111111110111100001110111 | 0 |
| DRC-091696 | 111111111101111111100001111111100001111111 | 111111111101111111100001111111100001111111 | 0 |
| DRC-091821 | 111111111111111111111111110111100001110111 | 111111111111111111111111110111100001110111 | 0 |
| DRC-092401 | 111111111111111111110001111111100001111111 | 111111111111111111110001111111100001111111 | 0 |
| DRC-092631 | 111111111111111111100001100001100001110000 | 111111111111111111100001100001100001110000 | 0 |
| DRC-092748 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-092750 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-093215 | 110111101111111111100001111111100001110111 | 110111101111111111100001111111100001110111 | 0 |
| DRC-093480 | 111111111111111111111111111111100001110111 | 111111111111111111111111111111100001110111 | 0 |
| DRC-093482 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-093488 | 110111111111111111100001111111100001111111 | 110111111111111111100001111111100001111111 | 0 |
| DRC-093493 | 111111111111111111100001111111100001111111 | 111111111111111111100001111111100001111111 | 0 |
| DRC-093495 | 111111111111111111111111110111100001110111 | 111111111111111111100001111111100001111111 | 6 |
| DRC-093664 | 111111000000000000000011111110000011111111 | 111111000000000000000011111110000011111111 | 0 |
| DRC-093666 | 111111111110111111100010111100000001111111 | 111111111110111111100010111100000001111111 | 1 |
| DRC-093674 | 111111111111111111111111111111100001111111 | 111111111111111111111111111111100001111111 | 0 |

| DRC-100033 | 1111111111111111101111111111111100001111111 | 1111111111111111101111111111111100001111111 | 0 |
|---|---|---|---|
| DRC-100035 | 1111111111110111111111111111111100001110111 | 1111111111110111111111111111111100001110111 | 0 |
| DRC-100037 | 1101111111110111111000011111111100001111111 | 1101111111110111111000011111111100001111111 | 0 |
| DRC-100099 | 1111110000000000000000000111111100001111111 | 1111110000000000000000000111111100001111111 | 0 |
| DRC-100122 | 1111111111111111111111111111001100001110111 | 1111111111111111111111111111111100001110111 | 2 |
| DRC-100299 | 1111111111111111111000011111111100001011111 | 1111111111111111111000011111111100001011111 | 0 |
| DRC-100300 | 1111111100011111110000000000000000000011111 | 1111111100011111110000000000000000000011111 | 0 |
| DRC-100301 | 1111111111111111111000011111111100001011111 | 1111111111111111111000011111111100001011111 | 0 |
| DRC-100307 | 1111111111111111111000011000011000011111111 | 1111111111111111111000011000011000011111111 | 0 |
| DRC-100310 | 1111111111111000001111111111111100001110111 | 1111111111111000001111111111111100001110111 | 0 |
| DRC-100503 | 1111111111111111111111111111111111100001110111 | 1111111111111111111111111111111111100001110111 | 0 |
| DRC-100740 | 1111111111111111111000011111111100001111111 | 1111111111111111111000011111111100001111111 | 0 |
| DRC-100753 | 1111111111111111111111111110111100001111111 | 1111111111111111111111111111111100001110111 | 2 |
| DRC-100759 | 1101111111111111111000011111111100001111111 | 1101111111111111111000011111111100001111111 | 0 |
| DRC-100923 | 1111111111111111111000011110111100001111111 | 1111111111111111111000011111111100001011111 | 2 |
| DRC-100929 | 1111111100011111110000000000000000000011111 | 1111111100011111110000000000000000000011111 | 0 |
| DRC-100931 | 1101111111111111111000011111111100001111111 | 1101111111111111111000011111111100001111111 | 0 |
| DRC-101072 | 1111111111111111111000011111111100001111111 | 1111111111111111111000011111111100001111111 | 0 |
| DRC-101306 | 1111111111111111111000011110111100001111111 | 1111111111111111111000011110111100001111111 | 0 |
| DRC-101308 | 1101111111111111111000011111111100001111111 | 1101111111111111111000011111111100001111111 | 0 |

861 <u>Supplementary Figure 1: Clustering of *M. tuberculosis* isolates based on 20 methods.</u>

862 For each method, the inclusion of an isolate into a cluster is outlined in the surrounding

863 circles using GraPhlAn (Asnicar et al., 2015). If an isolate is in a cluster not affected by

864 convergence, it is highlighted in black for the given method. If an isolate is in a cluster

865 affected by convergence, it is shown in grey.

Supplementary Figure 2: Algorithm of the phylogenetic inclusion method

An extension of the SNP-based clustering methods is used to detect complete chains of transmission that may be missed by pairwise SNP comparisons. In this example: A) shows the inferred phylogenetic sub-tree for five taxa and associated SNPs for each branch; B) a SNP cut-off is applied, which places 2 taxa (Taxon 1 and Taxon 4) together into a cluster; the most recent common ancestor (MRCA) of all taxa in this cluster (i.e. Taxon 1 & Taxon 4) is found; D) all descendants of this MRCA are added to the putative transmission chain.