The mechanical impact of *col11a2* loss on joints; *col11a2* mutant zebrafish

show changes to joint development and function, which leads to early onset

osteoarthritis.

Elizabeth A Lawrence[1]*, Erika Kague[1]*, Jessye A Aggleton[2], Robert L Harniman[3],

Karen A Roddy[1] and Chrissy L Hammond[1]

1. School of Physiology, Pharmacology and Neuroscience, University of Bristol

BS8 1TD

2. School of Anthropology and Archaeology, University of Bristol, BS8 1UU

3. School of Chemistry, University of Bristol, BS8 1TS

Corresponding author email

Chrissy.hammond@bristol.ac.uk

*These authors contributed equally to the work

Keywords (max 6)

Zebrafish, biomechanics, collagen, cartilage, Stickler syndrome, development

Abstract (max 200 words)

Collagen is the major structural component of cartilage and mutations in the genes encoding Type XI collagen are associated with severe skeletal dysplasias (Fibrochondrogenesis and Stickler syndrome) and early onset osteoarthritis. The impact of the lack of Type XI collagen on cell behaviour and mechanical performance during skeleton development is unknown. We studied a zebrafish mutant for *col11a2* and evaluated cartilage, bone development and mechanical properties to address this. We show that in *col11a2* mutants Type II collagen is made but is prematurely degraded in maturing cartilage and ectopically expressed in the joint. These changes are correlated with increased stiffness of both bone and cartilage; quantified using Atomic Force Microscopy. In the mutants, the skeletal rudiment terminal region in the jaw joint are broader and the interzone smaller. These differences in shape and material properties impact on joint function and mechanical performance, which we modelled using Finite Element Analyses. Finally, we show that *col11a2* heterozygous carriers reach adulthood but show signs of severe early onset osteoarthritis. Taken together our data demonstrate a key role for Type XI collagen in maintaining the properties of cartilage matrix; which when lost leads to alterations to cell behaviour that give rise to joint pathologies.

Introduction

Articular cartilage is a highly specialised connective tissue, which provides a smooth, lubricated surface for articulation and load transmission with low joint friction. Collagen is a major constituent of cartilage, accounting for around 75% of its dry weight (1). Type II collagen makes up 90-95% percent of the collagen network, while the remaining 5-10% is comprised of other collagens such as Type IX and XI, with studies of chick articular cartilage showing association of these three collagen types in a tight D-periodic array (2,3). These minor collagens help to organise and stabilize the Type II collagen fibril network that, along with proteoglycans, water and other proteins, form a dense extracellular matrix (ECM) in which chondrocytes are dispersed (4). The tight fibrillar structure from collagens and the water content, from interaction with glycosaminoglycans (GAGs), govern the mechanical properties of the cartilage. Type XI collagen belongs to the fibril-forming class of collagens; it is formed as a heterotrimer of three chains each encoded by a different gene: *COL11A1* (5), *COL11A2*, and *COL2A1*(6,7). While the α1 chain of Type XI collagen is expressed in both cartilaginous and ocular tissue, the α2 chain is predominantly expressed in cartilage.

Given the close interaction between Type II and Type XI collagens, mutations that affect either can cause similar destabilization of cartilage organisation, as observed in Marshall Stickler Syndrome. Stickler syndrome, which affects around 1 in 7500 new-borns, encompasses a hereditary group of conditions caused by defective Type II, IX or XI Collagen (8) and is divided in to three phenotypes: depending on the collagen mutation present. Type III is associated with mutations to the Type XI gene *COL11A2* (9) (10). Type III Stickler syndrome is characterised by skeletal, orofacial, and auditory abnormalities including: scoliosis; hearing loss; cleft

palate; joint hypermobility; (11), multiple hereditary exostoses (10), and premature osteoarthritis (OA) in 75% of patients before the age of 30 (8). The majority of mutations linked to Stickler syndrome lead to truncated proteins lacking the c-terminal domain of the peptide, disturbing the association of the alpha helices to form procollagens and consequentially the formation of collagen fibrils and fibres (12). Mutations in genes encoding Type XI collagens are also associated with other skeletal dysplasias, including the severe developmental condition Fibrochondrogenesis (13), and Weissenbacher-Zymuller syndrome (9).

Mutant mice for *Col11a1* (*Cho-/-*), are neonatally lethal and show decreased limb bone length, cleft palate and short snouts (14), and thicker, less uniform collagen fibrils in the cartilage ECM (15). Type II collagen degradation (16) and early onset OA were reported in *Cho/+* heterozygous mice (17). Additionally, mice haploinsufficient for *Col11a1* display altered susceptibility to load induced damage (18). While *Col11a2* mutant mice have been reported to show hearing loss, their skeletal phenotype has not been described (19). The interaction of Type XI collagen with Type II is important for the maintenance of the spacing and diameter of Type II collagen fibrils (20). As Type II collagen is the major collagen in cartilage, changes to its organisation can impact the mechanical performance of the cartilage. Computational modelling has shown that spacing and interconnectivity between collagen fibrils has a significant effect on the mechanical performance of cartilage (21). Cartilage is an intrinsically mechanically sensitive tissue, and changes to cartilage biomechanical performance have been extensively described during development (22), ageing and disease (23). It is also increasingly well understood that subtle changes to skeletal morphology and joint shape can increase susceptibility to joint conditions such as osteoarthritis later in life (24). A recent large

GWAS on hip shape identified *COL11A1* as a contributor to hip shape (25). Joint

shapes seen in development and disease have been shown to have significant

impact on the biomechanical performance of joints (24,26). What is less clear is the

sequence of events within the joint, do changes to shape precede changes to

cartilage structure and mechanical performance or vice versa, and what is the

relative impact of each change?

Zebrafish are an attractive model for studying the effect of genetic lesions on

skeletal development. The larvae are translucent which, twinned with fluorescent

reporter transgenic lines, enables dynamic imaging of skeletal cells (27,28) and the

development of the zebrafish craniofacial skeleton is well documented (29–31). The

zebrafish jaw joint is synovial (32) and requires mechanical input to form normally

(33,34). Acute knockdown of *col11a1* in zebrafish using morpholinos has been

shown to affect chondrocyte maturation (35), but no stable mutants for *col11a1* or

col11a2 have previously been reported.

Here, we show that larval zebrafish carrying a *col11a2* mutation display a

variety of phenotypes including alterations to: joint shape, cartilage composition, cell

organisation and the material properties of the cartilage during development. These

changes impact on the biomechanical and functional performance of the joint. The

mutant fish go on to display phenotypes consistent with Stickler syndrome such as

altered face shape and early onset osteoarthritis. Taken together these data suggest

that mechanical and cellular changes to the developing skeleton explain the

predisposition of people with mutations in Type XI Collagens to early onset

osteoarthritis.

Methods

Zebrafish husbandry and transgenic lines

Zebrafish were maintained as described previously (36), all experiments were approved by the local ethics committee and performed under a UK Home Office Project Licence. Transgenic lines *Tg(col2a1aBAC:mcherry),* *Tg(col10a1aBAC:citrine)^{hu7050}* (37) and *Tg(smyhc1:GFP)* (38), have been described previously. *col11a2^{sa18324}* mutant zebrafish was generated by the Zebrafish Mutation Project (Sanger Institute) and acquired from the European Zebrafish Resource Centre (EZRC). It carries a nonsense mutation (C>A base pair change at position 228aa, zv9 chr19: 7834334), leading to a premature stop codon which shortens the polypeptide to about 1/3 of the triple helical domain of the α2 chain of collagen XI.

DNA extraction and genotyping

Fins were clipped from anaesthetised or fixed zebrafish and incubated in base solution (25 mM NaOH,0.2 mM EDTA) before the addition of neutralisation solution (40 mM Tris HCl, Ph5.0). For genotyping we used KASP (LGC) genotyping or PCR followed by Sanger sequencing (*col11a2* F-GGTGGCCTGATTCTGACCA; *col11a2* R-TATCTCACACCAGGATGCCG). Mutants were identified by C>A base pair change at position 228aa.

Wholemount immunohistochemistry

Performed as previously described (37). Primary antibodies and dilutions used were: rabbit pAb to collagen II, (*abcam* ab34712), 1:500; mouse pAb to collagen II, (*DHSB* II-II6B3), 1:500, rabbit pAb to collagen I, (*abcam* ab23730), 1:100. Secondary antibodies were Dylight 488 or 550 (*Thermo Scientific*) used at a dilution of 1:500.

For imaging, larvae were mounted ventrally in 1% agarose and imaged on a Leica

SP5 confocal microscope with a 10x objective.

3D render generation, joint measurements, quantification of exostoses and cell

circularity

3D volume renders, surface models and measurements were acquired using Amira

6.0 (FEI). Surface models were generated manually by segmenting jaw joints using

the segmentation tool. Measurements, as depicted in Figure 1C-D and 3A were

taken using the 3D perspective measurement tool. To better visualise exostoses, 3D

volume renders were created, and the grayscale range of colour applied. Exostoses

were quantified in each lower jaw element from single confocal image stacks in

ImageJ (39) using the multi-point tool. Cell circularity was measured from confocal

image stacks of Type II collagen immunostained zebrafish larvae at 5dpf. The

freehand selection tool in ImageJ was used to outline chondrocytes in three distinct

jaw regions (shown in Figure 2H) and the measure function was used to analyse the

circularity of each cell. This was done for 10 cells in each region, in 3 wt and 3

col11a2 mutant zebrafish.

Live imaging of transgenic fish

Live larvae at 5dpf were anaesthetised in 0.1mg/ml MS222 and mounted ventrally in

0.3% agarose with tricaine prior to being imaged on a Leica SP5II confocal

microscope with a 10x objective. The number of slow muscle fibres and *col10a1a*

expressing cells were quantified manually in ImageJ from confocal images of double

transgenic *Tg(smyhc1:GFP);(Col2a1aBAC:mcherry)* and *Tg(col10a1aBAC:citrine);*

(col2a1aBAC:mCherry) zebrafish at 5dpf, respectively.

Alcian blue and alizarin red staining

5dpf and 7dpf wt and *col11a2* mutant larvae were stained following a previously

described protocol (40) and imaged on a Leica MZ10F stereo microscope prior to

genotyping.

***In situ* hybridisation**

In situ hybridisation was performed as described (41) using a previously described

col11a2 probe (42). Larvae were imaged on GXM-L3200 B upright microscope.

Nanoscale surface morphology and Young's Moduli

Atomic Force Microscopy (AFM) was performed on adult (1 year) bone and larval

(7dpf) cartilage from wt and *col11a2* mutant fish. A multi-mode VIII AFM with

Nanoscope V controller and PeakForce control mechanism were used and the force-

curves measured for means of set-point control in the PeakForce system and

analysed in real-time to provide quantitative nanomechanical mapping (QNM) of the

samples. QNM analysis was conducted with both Nusense SCOUT cantilevers

[NuNano, Bristol, UK], (nominal tip radius 5 nm, spring constants 21 – 42 N m-1) and

RTESPA-300 cantilevers [Bruker, CA, USA], (nominal tip radius 8 nm and spring

constants 30 – 60 N m-1), whilst high resolution imaging of topographic features was

conducted using SCANASYST-AIR-HR cantilevers [Bruker, CA, USA] (nominal tip

radius of 2 nm). The system was calibrated for measurement of Young's modulus

(YM) fitting with DMT models, using the relative method and samples of known YM

(highly oriented pyrolytic graphite (HOPG) (18 GPa) and PDMS-SOFT-1-12M (2.5

MPa) [Bruker, CA, USA]), for bone and cartilage measurements respectively. Bone

was investigated in ambient environment whilst cartilage was maintained in a

hydrated state post-dissection to minimise structural changes from drying. A root-

mean-square (RMS) mean was calculated for 65536 measurements taken over a

500 nm x 500 nm region, 3 repeats were performed per sample; repeated on 3 fish

per genotype.

Finite Element (FE) models

Single specimens that were representative of the confocal z-stacks of 7dpf wt or

mutant larvae dataset and their relative morphology were selected for the meshes.

Cartilage elements were segmented in Scan IP using Otsu segmentation

(Supplemental Figure 2A), then a solid geometry created using the interpolation and

3D wrap tool. Smoothing filters (recursive Gaussian at 1px3) were used on the

meshes to blend any rough small element clusters.

Cartilage geometry close to the joint was separated from the main cartilage in a

duplicate mesh using the 3D editing tool allowing us to assign different material

properties to hypertrophic chondrocytes and immature chondrocytes (Supplemental

Figure 2B). The mesh of the cartilage near the joints was subtracted from the original

cartilage mesh using a Boolean operation. The meshes were added to a model and

each part assigned their respective elastic isotropic material properties based on

AFM measurements; values in Supplemental Figure 2, Table 1.

The models were imported into Abaqus and 2 steps created: 1 to simulate jaw

closure and 2 for jaw opening. Boundary conditions were applied to these steps, with

the jaw constrained in all axes of motion at the ceratohyal to anchor it in space, and

in y and z at the base of the palatoquadrate. Muscle forces, direction of

opening/closure and muscle attachment points were as previously described (43).

The datum tool in Abaqus was used to create a custom rectangular datum

coordinate system for each muscle; then used as the coordinate system for force

direction between each muscle's insertion and origin to ensure force travelled along

the same vector from one end to the other. A job was created and executed for the

model, and the output analysed for stress, strain and displacement.

Measurement of jaw displacement and movement frequency

High-speed movies were made of jaw movements in wt and *col11a2* mutants;

frames corresponding to maximum jaw displacements were selected, imported into

ImageJ and the difference, in µm, between resting and open states at points shown

in Figure 5A recorded. The number of mouth movements in 1000 frames was

recorded from 7 wild type (wt) and 7 *col11a2* mutant fish as previously described in

(30).

Micro-Computed Tomography (µCT)

Three *col11a2 +/-* and three wt adult fish of the same age (1 year old) were fixed

in 4% PFA for 1 week followed by sequential dehydration to 70% ethanol. Heads

were scanned using an XT H 225ST micro-CT scanner (Nikon) with a voxel size

of 5 µm, x-ray source of 130 kV, 53 µA and without additional filters. Images were

reconstructed using CT Pro 3D software (Nikon).

Histology

Three 1 year old *col11a2 +/-* and three wild types, were decalcified in 1M EDTA

solution for 20 days. Samples were dehydrated in ethanol, embedded in paraffin

and sagittally sectioned at 8 µm, relevant joint sections were de-waxed and

stained with 1% Alcian blue 8GX (pH 2.5, then counterstained with Haematoxylin

and Eosin. We adapted the OARSI cartilage OA histopathology grading system

(44) to grade severity of OA. Five sections per jaw joint (per fish n=3 fish) were

scored. PicroSirius red staining was performed using 0.1% Sirius red F3B in

saturated aqueous Picric acid, washed in acidified water, dehydrated and

mounted under coverslips with DPX, then imaged using polarising filters.

Second Harmonic Generation (SHG)

SHG images were acquired from histological sections of wt and *col11a2 +/-* (n=3

fish for each genotype) using 25x 0.3 NA water dipping lens, 880nm laser

excitation and simultaneous forward and backward detection (440/20) in Leica

SP8 AOBS confocal laser scanning microscope attached to a Leica DM6000

upright epifluorescence microscope with multiphoton lasers and confocal lasers

allowing fluorescent and SHG acquisition of the same sample and z-stack.

Microscope parameters for SHG acquisition were set as described previously

(45). Maximum projection pictures were assembled using LAS AF Lite software

(Leica).

Results

col11a2 and col2a1 are co-expressed in the zebrafish lower jaw

To establish the extent of *col11a2* expression in cartilage we performed *in situ* hybridisation in larval zebrafish. Strong *col11a2* expression could be seen throughout the craniofacial cartilages including the Meckel's cartilage, palatoquadrate, ceratohyal and ethmoid plate (Supplementary Figure 1A). At 3dpf, the expression pattern of *col11a2* largely overlapped the expression of the Type II collagen gene *col2a1a* visualised with the *Tg(col2a1aBAC:mCherry)* reporter zebrafish (Supplementary Figure 1B). The domain of *col11a2* expression labelled more of the joint than was labelled by the *col2a1a* transgene, and expression of both *col11a2* and *col2a1a* preceded that of the mature Type II protein, visualised by immunostaining, such that immature cells at the jaw joint and Meckel's symphysis express *col11a2* and *col2a1a* RNA at 3dpf but not the mature Type II protein (Supplementary Figure 1C).

col11a2 mutants show atypical Type II collagen localisation as they develop

As Type XI collagen has previously been reported in the core of Type II collagen fibrils (3) and is thought to have a role in the stability of Type II collagen (46), we wanted to test whether loss of *col11a2* in zebrafish would impact Type II collagen stability. For this we studied the *col11a2* [sa18324] mutant which carries a nonsense mutation that introduces a premature stop codon at amino acid 228 (of 1877). We observed non-sense mediated decay in mutants *in situ* hybridised for the *col11a2* probe (data not shown), therefore it represents a null mutant. This mutant was crossed with the Tg(*col2a1a:mCherry)* to visualise expression of *col2a1a* and we studied its expression in craniofacial cartilages from 3-7dpf. We saw no differences

in the position, timing or extent of *col2a1* expression between mutants and their

siblings at 3dpf suggesting that loss of *col11a2* has no impact on the expression of

col2a1a, although alterations to craniofacial skeletal shape in the mutant were

detectable from 5dpf (Supplementary Figure 2). We next used immunostaining to

detect Type II collagen protein in mutant and wild type (wt) larvae. At 3dpf we could

not detect any differences between wt and mutant larvae (Figure 1A, B). However,

by 5dpf clear differences in the distribution of Type II collagen were seen in the lower

jaw (denoted by asterisks in Figure 1A, B). In wt fish, Type II collagen can be seen in

the extracellular matrix surrounding each chondrocyte in the lower jaw cartilages,

whereas in mutants, protein expression is concentrated in the perichondrium and

reduced between chondrocytes (dashed insets in Figure 1B). Alongside this

reduction of Type II collagen in the more mature matrix towards the middle of the

cartilage elements, small pieces of immunostained material were seen separate from

the main elements (red asterisks in Figure 1B) and ectopic expression of Type II

collagen in the ligament was observed (red arrow in Figure 1B). Changes to the

shape and size of the cartilage elements also became apparent at 5dpf, with mutant

larvae displaying thicker, shorter cartilage structures, with less definition. At 7dpf

these shape discrepancies were maintained. A pronounced reduction and

disorganisation of Type II collagen became clear by 7dpf (Figure 1B) and was more

obvious in the cartilages that make up the lower jaw, with the lateral cartilage of the

ear more preserved (Figure 1A, B). Taken together these data suggest that

expression and synthesis of *col2a1a* are unaffected by loss of *col11a2,* but that

maintenance of Type II collagen protein is impaired in the mutants.

Zebrafish with a *col11a2* mutation have altered jaw and joint morphology

during development

Humans carrying mutations in Type XI collagen show alterations to craniofacial

shape, including midface hypoplasia and micrognathia (47,48). From Type II

collagen immunostaining and alizarin red/alcian blue staining, we observed that

mutant zebrafish also show altered craniofacial morphology (Figures 1B and 3D). At

3dpf there was no significant difference in jaw morphology, but at 5 and 7dpf mutants

had significantly shortened, wider jaws (Figure 2A, B), in line with the broader, flatter

face shape observed in patients with Stickler syndrome.

As people with mutations in Type XI collagens also display abnormal joint shape (49)

and increased susceptibility to osteoarthritis (8), we generated 3D surface models of

the joint between the Meckel's cartilage and the palatoquadrate (jaw joint) at 3, 5 and

7dpf in wt and mutants (Figure 2C, D). At 3dpf, there was no significant change to

joint morphology, however at 5 and 7dpf, mutants show enlarged terminal regions of

the skeletal rudiments. Specifically, these changes were seen at the Meckel's

cartilage neck and head adjacent to the joint (Figure 2E, F, position of

measurements shown by red and green lines in Figure 2C). A reduction in the joint

space (Figure 2G, position of measurement shown by white line in Figure 2C), such

that the interzone is no longer clearly defined in the renders was also observed,

likely due to the increased local deposition of Type II collagen (Figure 1B and 2D).

These results show that the abnormal pattern of Type II collagen deposition seen in

col11a2 zebrafish mutants at 5 and 7dpf leads to altered joint shape.

Mutation of *col11a2* leads to altered chondrocyte cell behaviour

Using confocal imaging of a transgenic reporter for *col2a1a* we were also able to

observe alterations to chondrocyte behaviour in mutant fish at 7dpf. In wt larvae all

cells expressing *col2a1a* are located within the cartilage element, however, in

mutants we frequently observed chondrocytes located outside the main body of the cartilage element (Figure 2H). We quantified the number of these cells in the lower jaw cartilages of wt and mutant larvae during early development. Prior to 5dpf these spurs, which resemble hereditary multiple exostoses (50), are not observed (Figure 2I). However, at 5dpf and 7dpf they are present in the Meckel's cartilage and ceratohyal of *col11a2* mutants, suggesting a failure of chondrocyte progenitor cells to fully intercalate into the cartilage element prior to expression of *col2a1a*.

In addition to these protrusions, we observed differences to chondrocyte morphology within the jaw elements of mutants at 5dpf. Chondrocyte circularity is a measure of maturation, as in zebrafish chondrocytes become less rounded over time and form tightly packed stacks as they mature towards hypertrophy and show organisation reminiscent of the mammalian cartilage growth plate. Chondrocytes in the centre of cartilage elements (representing the most mature chondrocytes) of *col11a2* mutants show increased circularity over those in wt cartilage (Figure 2J). These results suggest that the maturation of chondrocytes in *col11a2* mutants is disrupted, potentially due to the loss of Type II collagen from the ECM, which could provide an explanation for the thicker appearance of the cartilage elements in mutants.

Larval and adult *col11a2* mutant zebrafish have altered material properties in the craniofacial skeleton

To test whether loss of Type II collagen and alterations to chondrocyte behaviour led to changes to the material properties of the cartilage, we performed atomic force microscopy (AFM) on dissected lower jaw cartilages from 7dpf wt and mutants. We tested the properties in regions containing immature cells close to the jaw joint and Meckel's symphysis (Figure 3A) and observed a significant increase in Young's

modulus (YM) from an average of 4.15 to 7.4 MPa (Figure 3B). In more mature, intercalated cells towards the centre of the Meckel's cartilage (in which we saw loss of Type II collagen) the difference in YM was around 4 times greater than that of comparable regions in wt (Figure 3B). This suggests that loss of Type II collagen as a result of the *col11a2* mutation leads to stiffening of the cartilage ECM. As we observed altered material properties in larvae at pre-skeletonised stages we wanted to test whether this would persist to adulthood and impact bone properties. We dissected jaw bones and operculae from wt and mutant fish and used AFM to establish YM. As in the larval cartilage, we observed that the bone from *col11a2* mutant fish had a significantly higher YM than siblings (Figure 3C).

Type II collagen loss is not accompanied by changes to GAGs, Type X or Type I collagen in *col11a2* mutant zebrafish.

As Type II Collagen was prematurely lost from maturing chondrocytes, we sought to examine whether glycosaminoglycans (GAG)s, were similarly reduced. We stained wt and mutant larvae with Alcian blue (to mark cartilage GAGs) and Alizarin red (to mark bone) at 5 and 7dpf. We saw no reduction of GAG reactivity in the mutants at 5 or 7dpf (Figure 3D). Additionally, we saw no dramatic changes to Alizarin red, with dermal and chondral bones in mutants mineralising at a similar rate to wt fish.

During cartilage maturation, a change in collagens from Type II to Type X collagen is associated with chondrocyte hypertrophy (51). In teleosts such as zebrafish *col10a1a* marks hypertrophic chondrocytes, but also osteoblasts and ligament cells (52,53). To test whether there was any change to the extent of chondrocyte hypertrophy or to the number of osteoblasts or skeletal connective tissue cells we crossed the *col11a2* mutant into the *col10a1a* transgenic reporter (Figure 3E).

Quantification of *col10a1a* positive cells in wild type and mutant fish revealed no

differences to the number of hypertrophic chondrocytes at 7dpf (Fig 3F green arrows

and graphs Fig 3.G,H), we also saw no difference in the number of osteoblasts in the

dentary which is located directly adjacent to the MC (red arrows Fig 3F,

quantification in 3G) or to the number of cells in the IOM ligament (purple arrows in

3F, quantification in 3G). This suggests that the onset of hypertrophy is not disrupted

in *col11a2* fish, despite the 'less mature' appearance of their chondrocytes.

During cartilage degeneration, such as in osteoarthritis, a switch of collagens is

commonly reported with a reduction of Type II and increase in Type I collagen,

associated with stiffer matrix (54). To test whether loss of Type II collagen led to

replacement with Type I, we performed immunostaining in wt and mutant larvae at 5

and 7dpf. In wt fish Type I collagen was present in the jaw joint space, Meckel's

symphysis and at a low level in the cartilage ECM, this pattern was unchanged in

mutants at 5dpf, however by 7 dpf there was a reduction of Col I in the joint interzone

of mutants but no change within the cartilage elements themselves (Supplementary

Figure 4).

Taken together, these results suggest that loss of Type II collagen in *col11a2*

mutants does not lead to loss of GAG, nor to compensatory increases of Col I or

col10a1, and that hypertrophy is unaffected by the *col11a2* mutation. As a result, the

increase in cartilage stiffness observed from AFM cannot be attributed to alterations

in these components.

Both shape and material properties impact the biomechanical performance of

the zebrafish lower jaw

We have previously modelled the biomechanics of zebrafish jaw opening and

closure during early ontogeny using Finite Element Analysis (FEA) (43) and shown

that paralysis and the accompanying changes to joint shape impact the strain pattern

in the developing joint (33). Therefore, we used FEA to model how the changes to

shape and material properties observed in *col11a2* mutants would affect the

biomechanical performance of the lower jaw. Meshes were generated of wt and

mutant larvae at 7dpf (meshes shown in Supplementary Figure 5B). We applied

muscle forces as per Roddy et al (55) and used the material properties established

from AFM (Figure 3B).

We first modelled the wt and the mutant jaw shapes using the material properties

established from each genotype and modelled a 2 step process for jaw movement

with Step 1 denoting jaw closure and Step 2 jaw opening (as per (43,55)) and

visualised the maximum principal (EMax) and minimum principal (EMin) strains for

jaw opening (Figure 4) and closure (Supplementary figure 6). In the wt model,

tensional strains (EMax) are located laterally around the joint and either side of the

Meckel's symphysis, focused on the muscle insertion points with the strain spreading

widely through the element. By contrast, in mutants, maximum principal strain is

concentrated on the joint interzone, with little spread through the cartilaginous

element (note blue colour throughout the cartilage of mutants, c.f. greens and

yellows in wt) (Figure 4A, Supplementary videos 1,2). Comparison of the Meckel's

symphysis joint interzones for strains (EMax) showed that the difference between the

types of models (wt models and the mutant models) was far larger than the

difference between each type of model with different material properties

(Supplementary Figure 5C, D). In the mutant model with mutant values the joints

averaged 1.63E-02 maximum strains (EMax), and in the mutant model with wt

values the joint strain averaged 1.51E-02, whereas in the wt models, the joint strains

averaged 4.41E-02 (EMax) in the wt with wt values model, and 5.14E-06 in the wt

with mutant values model. Between each type of model (wt against mutant), the

average maximum principal strains have not changed substantially despite change in

material properties. In wt, compressional strains (EMin) are at the Meckel's

symphysis, the medial surface of the anterior MC and on the dorsolateral side of the

jaw joint (Figure 4A, Supplementary Figure 6A). In mutants, again, the minimum

principal strains are more focal than in wt larvae (Figure 4A, Supplementary Figure

6A).

Changing the material properties of the models affects patterns of strain and

displacement. This influences the displacement on the overall jaw morphology as

seen in the jaw opening stage when both models are given the stiffer mutant material

properties. In these models, the jaw shows less displacement in the opening

movement compared to the situation in which models are given the less stiff wildtype

material property values.

To test whether the change to the strain pattern was predominantly caused by the

alteration to the shape of the jaw elements or the changes to YM, we next modelled

the effect of mutant properties in the wt shape, and wt properties in the mutant shape

(Figure 4B, Supplementary Figure 6B). We observed that changing YM in the wt

shape to the mutant values decreased the spread of max and min principal strains,

such that the pattern was intermediate between the mutant and wt. Likewise,

inserting the wt values for cartilage into the mutant shape lead to an increase in the

extent of both tension (EMax) and compression (EMin). However, it did not fully

'rescue' the pattern, leading us to conclude that while both shape and material

properties play a role in the mechanical performance of the tissue, the effect of

shape is greater than that of material properties.

col11a2 zebrafish mutants show impaired jaw function

As patients with Stickler syndrome suffer from joint hypermobility (56), and as

col11a2 zebrafish mutants show aberrant joint morphology, we looked at jaw

function at 5dpf. Zebrafish have two joints within the lower jaw and make distinct

movements for feeding and breathing (57). By taking movies and quantifying jaw

movement, we observed that mutants make significantly fewer total movements than

wt (Figure 5B, Supplementary videos 3,4). This was due to a reduction in the number

of movements involving the jaw joint, as we observed no significant difference in the

frequency of movements involving the buccal joint (Figure 5B). However, mutant

zebrafish show an increased range of motion at the buccal joint, which appears to

dislocate (Figure 5A, C). To rule out the possibility that this change to movement was

caused by altered muscle patterning, we quantified the number of slow twitch fibres

in the jaw at 5dpf and saw no difference in fibre number between wt and mutants

(Supplementary Figure 7A,B). We also measured the diameter of the

intermandibularis posterior and interhyoideus muscles in the lower jaw from

birefringence and found no change in diameter between wt and mutants

(Supplementary Figure 7C). Taken together this suggests that the alterations to joint

shape observed in *col11a2* mutants are the cause of abnormal joint function.

Premature OA is observed in adult *col11a2* heterozygous fish

Due to the abnormalities in joint shape, mechanical performance and function in

mutants, and since aberrant joint loading is highly associated with OA risk (58), we

wanted to test whether adult mutants would develop premature osteoarthritis. To

address this question, we analysed 1-year old heterozygous fish (*col11a2+/-*) and wt

siblings using micro-computed tomography (μCT). Craniofacial abnormalities were

observed in *col11a2*+/-, including jaw protrusion and hypoplasia of the fronto-nasal

bone (Figure 6A). Changes in joint shape were observed in *col11a2+/-* accompanied

by narrowing of the inter-joint space (Figure 6B). To identify the histopathological

changes related to OA, we stained wt and *col11a2+/-* joint sections for Alcian Blue

and H&E (Figure 6C). While in wt sections a defined cartilaginous layer lines the

joint, in *col11a2+/-* the cartilage shows signs of degradation. Grading of 5 sections

per joint per fish (n=3 fish) using the criteria in the OARSI scoring system (44)

showed an average score of 6 in the *col11a2+/-* sections which is characterised by

deformation and change in the contour of the articular surface, compared to an intact

surface and normal cartilage with average score of 0 in the siblings (Figure 6C).

Osteophytes were not observed. We analysed collagen organisation using

PicroSirius red staining and Second Harmonic Generation (SHG) (Figure 6D and E).

In wt jaws, the cartilaginous layer at the joint shows organised collagen fibres with a

distinct orientation from those of the underlying bone (Figure 6D, note change in

colour from red to green in PicroSirius red staining). However, in the *col11a2+/-* the

transition from cartilage to bone is lost and the overall organization is perturbed

(Figure 6D). Thicker collagen bundles and fibres displaying abnormal orientations

were seen through SHG in *col11a2+/-* samples (Figure 6E). Taken together these

data demonstrate that loss of *col11a2* leads to early onset of osteoarthritis-like

changes in adults.

Discussion

Mutations in the Type XI collagen genes *col11a1* and *col11a2* have previously

been linked to numerous skeletal dysplasias, such as Stickler syndrome and

Fibrochondrogenesis, which are associated with cartilage destabilisation, and

abnormal skeletal shape and properties. Here, we describe the impact of loss of

col11a2 in zebrafish and show changes to ECM composition, material properties,

craniofacial shape, mechanical performance, chondrocyte behaviour, and joint

function in larval and adult fish.

Type XI collagen is important for the protection of Type II collagen from

degradation (59); our data suggest that, while transcription and secretion of Type II

collagen is unaffected at early stages of larval development, the assembly of Type II

collagen fibrils may be altered in mutants, making them more susceptible to

degradation. This idea is given weight by the identification of fragments of Type II

positive material seen surrounding the cartilage elements. What happens to those

degraded collagen fragments is still unclear. Potentially, they may be cleared by the

phagocytic cells of the innate immune system either with a rapid resolution, or

alternatively, continued accumulation of these fragments could lead to the low-level

inflammation associated with osteoarthritis (60,61). Loss or breakdown of Type II

collagen also occurs as the chondrocytes mature, such that the matrix between the

chondrocytes almost completely lacks Type II collagen, while the matrix of the

perichondrium is relatively preserved. We have tested effects of *col11a2* loss on the

material properties of cartilage and bone, and our data shows an increase in YM in

both tissues, with the greatest difference seen in mature chondrocytes. It may be

noted that Young's modulus for zebrafish cartilage is higher than that from other

species (4.15MPa in fish vs 0.45Mpa in human articular cartilage (62)). One likely explanation is the variation in relative ratio of cells to matrix during development and across species in evolution. In mature human articular cartilage the ratio is approximately 10:90 cells to matrix compared to 80:20 in zebrafish (63). The higher YM in mutants was not explained by any obvious increased calcification, accumulation of Type X or Type I collagen or loss of GAG. In rodent models increased matrix stiffness has been described as chondrocytes mature in the growth plate (64,65). The stiffness of collagen matrix is controlled by several factors including fibre diameter and the density of intra-fibrillar cross-links, and abnormalities in collagen fibrillar assembly have been related to changes to mechanical properties of the cartilage during progression of OA (66) and ageing (67).

It has previously been reported that patients with Stickler syndrome develop premature OA (1) but the mechanism by which this occurs is unclear. We and others have previously shown that, despite living in an aquatic environment, zebrafish can also develop alterations to the joint that strongly resemble OA (68,69). Interestingly, we see premature development of osteoarthritis in *col11a2* heterozygous adult zebrafish. This is manifested by abnormal collagen organisation, degeneration of joint cartilage and loss of joint space. During OA, proteoglycans are lost from the cartilage prior to the degradation of the collagen network in the extracellular matrix (2). This change to the organisation and content of collagen in the cartilage leads to changes in its material properties (3), including its stiffness and tensile strength (4). It has previously been demonstrated that in OA, cartilage stiffness is often reduced (17,70)while we saw a dramatic increase in cartilage matrix stiffness in the *col11a2* mutants these measurements were taken from larvae. We saw increased YM in adult bone, both of dermal (operculum) and chondral (jaw)

bone, albeit less dramatically than in the cartilage. Potentially, stiffer bone could exacerbate OA pathogenesis; as subchondral bone thickening accelerates the degradation of articular cartilage (71). Alternatively, and perhaps more likely, changes to joint loading from the abnormal shape and function throughout life may be the driver for the development of pathogenic osteoarthritis-like changes in the joint. To test this more fully it would be desirable to follow the development of the pathology throughout the life course of the fish.

As joint mechanical performance is impacted by its shape and the material properties of the tissues, we explored the relative impact of each by testing the impact of altering material properties in the wild type and mutant shapes. From this we deduced that while both contribute to the strain pattern, the larger impact comes from joint architecture. However, questions remain to the exact sequence of events; are the increases in YM in immature chondrocytes sufficient to drive local changes to cell behaviour within the joint? If so subtle changes to joint morphology, could impact joint mechanics upon onset of function, leading to further, more significant changes to skeletal cell behaviour. Movement of joints has been shown to be required for their correct specification in the majority of joint types in all species studied (72–74,26,75). Interestingly, at the earliest stages we studied (3dpf), prior to the onset of joint movement, the mutants are barely distinguishable from wild types, despite the *col11a2* gene being expressed throughout the cartilage from 2dpf. Following the onset of movement changes between wild type and mutants become more pronounced, these include the degradation of Type II collagen from the mature matrix, and the loss of the joint space. This loss of correct joint spacing and the enlargement of the rudiments could be explained by premature differentiation of the immature cells of the interzone. A requirement for normal movement has been

demonstrated in chick, mouse and fish to maintain joint space and to prevent ectopic

expression of Type II collagen (33,34,76,77). Alternatively, it could represent a failure

to maintain local gdf5 signalling; it has recently been shown that there is a

requirement for the continued influx of Gdf5 positive cells for correct joint

specification (78).

It is likely that by changing the mechanical performance of the joint, that

mechanosensitive genes will be differentially activated, and these likely control the

cellular changes we describe. Candidates that could be differentially activated in the

mutants could include the Piezo ion channels which have been shown to play a role

in OA (79). Another candidate could be the YAP pathway; YAP is implicated in

negative control of chondrogenesis (80,81). Or the genes in the Wnt signalling

pathway. The Wnt pathway has been implicated in developmental skeletal

mechanosensation in mice, chicks (82) and zebrafish (83), and could potentially be

acting in combination with BMP regulatory genes such as Smurf1 (84). We have

shown in zebrafish that wnt16 controls chondrocyte proliferation and migration in the

joint region. Wnt16 is also linked to hip geometry (85) altered cortical bone thickness

(86,87) the response of chondrocytes to injury and to osteoarthritis (88,89).

Following the onset of movement, we also see the appearance of cells

located outside the cartilage anlage, which bear some resemblance to multiple

hereditary extoses (MHE). Stickler syndrome is associated with MHEs (90). It has

been reported in a zebrafish model that the development of MHE is driven by

changes to the matrix from loss of the Extosin genes, that, while dispensable for

early chondrocyte differentiation are required for chondrocyte maturation,

hypertrophy and intercalation and which encode genes lead to matrix sulfation (50).

Potentially, the loss of Type II collagen in the *col11a2* mutants could perturb

sulfation. Alternatively, these cells could fail to intercalate then be extruded due to altered joint function, as paralysis has been shown to control chondrocyte intercalation in zebrafish (91). The failure of these cells to fully intercalate leads to shorter, thicker elements in *col11a2* mutants.

Taken together our findings show that loss of *col11a2* in zebrafish leads to changes to matrix phenotype, and cell behaviour that impact the biomechanical and functional performance of the developing joint leading to premature OA. By making use of the detailed dynamic imaging unique to small translucent models like the zebrafish we were able to follow the alterations to the developing skeleton at cellular resolution, identifying changes to cell behaviour that go some way to explaining how loss of a relatively minor collagen subtype can have such a profound effect on the human skeleton in diseases such as Stickler syndrome and Fibrochondrogenesis.

The authors declare no competing interests.

EL, KR and EK carried out the molecular lab work, participated in data analysis,

participated in the design of the study and drafted the manuscript; JA generated the

FE models; RH performed AFM; CLH conceived the study, designed the study,

coordinated the study and helped draft the manuscript. All authors gave final

approval for publication.

Acknowledgements

The authors would like to thank Stephen Cross and the Wolfson Biomaging facility

staff for help with image acquisition and analysis.

Funding

CLH and EK were funded by Arthritis Research UK grants 21211 and 19947. KR
was funded by the MRC (MR/L002566/1). EL is funded by a Wellcome Trust
Dynamic Molecular Cell Biology PhD programme. PeakForce AFM was carried with
equipment funded by the EPSRC (EP/K035746/1).

 Figure legends

Figure 1: *col11a2* zebrafish mutant larvae show progressively altered Type II

collagen protein localisation in jaw cartilage. A, B) Maximum projection of ventral

and lateral confocal image stacks from wild type (wt) (A) and homozygous mutant

(*col11a2-/-*) (B) larvae immunostained for Type II collagen at three time points (3, 5

and 7dpf). White arrows indicate areas of change in Type II collagen distribution in

the extracellular matrix. Dashed insets show single stack images of regions with

reduced deposition (white asterisks represent areas where Type II collagen is

maintained in mutant fish and red asterisks show fragments of Type II collagen

positive material outside the main cartilage elements). Red arrows show

interoperculomandibular (IOM) ligament. Scale bar = 100µm.

Figure 2: *col11a2* mutant zebrafish develop altered morphology and joint

spacing in the lower jaw. A, B) Lower jaw shape quantification (n = 3 for all),

location of measurements shown to the left of graphs. C) Representation of

measurements taken of joint neck (red line), joint head (green line) and joint space

(white line) (Meckel's cartilage = light blue, palatoquadrate = dark blue). Orientation

compass: A = anterior, L = lateral, M = medial, P = posterior. D) 3D surface renders

of jaw joint from confocal images of wt and *col11a2 -/-* at 3, 5 and 7dpf. Red

arrowheads = areas of change. E - G) Quantification of joint morphology at the

Meckel's cartilage neck at joint (E), Meckel's cartilage head at joint (F) and joint

space (G) (n = 3 for all). H) 3D volume renders of wt and *col11a2-/-* zebrafish at

7dpf. Dashed insets show Meckel's symphysis at higher magnification (red

arrowheads = protruding cells). I) Quantification of protruding cells in wt and

col11a2-/- zebrafish at 3 - 7dpf (n = 3, 3, 4, 4, 13, 6, 8, 6). J) Quantification of cell

circularity in the Meckel's cartilage in 5dpf wt and *col11a2-/-* fish (n=3 for all).

Location of measurements shown in H (red box = Meckel's symphysis, blue box =

mid element, green box = jaw joint). Student's unpaired t-tests performed in A, B, E -

G, I and J: data is mean with SEM (J shows mean with no SEM, t-tests performed

between mean values). *P≤0.05 P**≤0.01 P***≤0.001

Figure 3: *col11a2* mutants have altered material properties in more mature

cartilage which is not explained by increased mineralisation or hypertrophy.

Location of AFM measurements taken from larvae shown in (A). Measurements for

immature chondrocytes taken from either of the two areas marked by black boxes,

measurements for hypertrophic chondrocytes taken from area marked by red box. B,

C) Young's modulus values for (B) immature and hypertrophic chondrocytes in wt

and *col11a2-/-* (n=3 for both) at 7dpf and (C) adult bone from the operculum and jaw

in wt (n=8 and 3 respectively) and *col11a2-/-* (n=6 and 3 respectively). D) Ventral

and lateral views of alizarin red alcian blue staining show glycosaminoglycans in

cartilage (stained in blue) and mineralisation (stained in red) in wt and *col11a2-/-* fish

at 5 and 7dpf. Red asterisks indicate areas of bone formation. MC = Meckel's

cartilage, PQ = palatoquadrate, C = ceratohyal, BA = branchial arches, HS =

hyosymplectic, OC = otic capsule, OP = operculum, CL = cleithrum, PS =

parasphenoid, NT = notochord tip, O = otoliths. Scale bar = 200μm. E)

col10a1aBAC:citrine;col2:mCherry transgenic line shows Type X (yellow) and Type II

(red) collagen in wt and *col11a2-/-* zebrafish at 7dpf. Scale bar = 100μm. F)

Quantification of col10a1 expressing cells in hypertrophic chondrocytes, IOM

ligament cells and osteoblasts in the lower jaw at 7dpf (position of each cell type

shown by green, purple and orange arrows in (E) respectively) (n=3 for all). G)

Quantification of *col10a1* expressing hypertrophic chondrocytes in 7dpf wt and

col11a2-/- fish (n = 3 for all) (M = Meckel's cartilage, PQ - palatoquadrate, C =

ceratohyal). Student's unpaired t-tests were performed in B, C, F and G, data is

mean with SEM (B shows mean with no SEM). *P≤0.05 P**≤0.01 P***≤0.001

Figure 4: Shape changes in *col11a2* zebrafish mutants have a greater effect on

jaw biomechanics than material property changes. A, B) Finite Element (FE)

models of maximum (E. Max) and minimum (E. Min) principal strain during mouth

opening in 7dpf wt and *col11a2-/-* zebrafish. Red arrowheads = areas of high strain,

black arrowheads = areas of low strain, black asterisks = jaw joint. A) wt jaw shape

with wt material properties and *col11a2-/-* shape with *col11a2-/-* material properties.

B) wt shape with *col11a2-/-* material properties and *col11a2-/-* shape with wt material

properties. Ventral and lateral views shown for each condition.

Figure 5: ***col11a2* mutant zebrafish have abnormal jaw movement at 5dpf**. A)

Stills from high speed movies show range of jaw movement in wt and *col11a2-/-* B,

C) Analysis of (B) total jaw movements and (C) range of movement at two locations

shown in (A): red line = mouth, blue line= buccal joint (n= 7 for all). Student's

unpaired t-tests performed for B and C, data is mean with SEM. *P≤0.05 ***P≤0.001.

Figure 6: Mutations in zebrafish *col11a2* result in changes that trigger

premature OA. A, B) 3D renders from µCTs of 1-year old wt and *col11a2*

heterozygous mutant (*col11a2+/-*). A) Yellow arrow = jaw joint, dashed green arrow =

region of jaw protrusion in *col11a2+/-*, green arrow = region of hypoplasia in fronto-

nasal skeleton. B) Higher magnification image of joint region where dashed yellow

line = inter-joint space. C, D) Paraffin sections of the jaw joint stained with (C) Alcian

blue and Haematoxylin/Eosin and (D) PicroSirius red. Dashed black line = cartilage

layer, black arrows = underlying bone, dashed white line = cartilage, white arrow =

bone (green). E) Second Harmonic Generation (SHG), asterisks pointing to areas of

thinner fibres not detected by SHG, Red arrows = thicker collagen bundles on abnormal orientation. Scale bars = 50μm.

1. Weiss C, Rosenberg L, Helfet AJ. An ultrastructural study of normal young adult human articular cartilage. J Bone Joint Surg Am. 1968 Jun;50(4):663–74.

2. Vaughan L, Mendler M, Huber S, Bruckner P, Winterhalter K, Irwin M, et al. D-periodic distribution of collagen IX along cartilage fibrils. J Cell Biol. 1988;106:991–7.

3. Mendler M, Eich-Bender SG, Vaughan L, Winterhalter KH, Bruckner P. Cartilage Contains Mixed Fibrils of Collagen Types II, IX, and XI.

4. Fox AJS, Bedi A, Rodeo SA. The Basic Science of Articular Cartilage: Structure, Composition, and Function. Sports Health. 2009;1(6):461.

5. Burgeson RE, Hollister DW. Collagen heterogeneity in human cartilage: Identification of several new collagen chains. Biochem Biophys Res Commun. 1979;(87):1124–31.

6. Eyre DR. The Collagens of Articular Cartilage. Semin Arthritis Rheum. 1991;21(3):2–11.

7. Hafez A, Squires R, Pedracini A, Joshi A, Seegmiller RE, Oxford JT. Col11a1 Regulates Bone Microarchitecture during Embryonic Development. J Dev Biol. 2015;3(4):158.

8. Couchouron T, Masson C. Early-onset progressive osteoarthritis with

hereditary progressive ophtalmopathy or Stickler syndrome. Jt Bone Spine.

2011 Jan;78(1):45–9.

9.   Vikkula M, Madman EC., Lui VC., Zhidkova NI, Tiller GE, Goldring MB, et al.

Autosomal dominant and recessive osteochondrodysplasias associated with

the COL11A2 locus. Cell. 1995 Feb 10;80(3):431–7.

10.   Sirko-Osadsa DA, Murray MA, Scott JA, Lavery MA, Warman ML, Robin NH.

Stickler syndrome without eye involvement is caused by mutations in

COL11A2, the gene encoding the α2(XI) chain of type XI collagen. J Pediatr.

1998 Feb 1;132(2):368–71.

11.   Acke FR, Malfait F, Vanakker OM, Steyaert W, De Leeneer K, Mortier G, et al.

Novel pathogenic COL11A1/COL11A2 variants in Stickler syndrome detected

by targeted NGS and exome sequencing. Mol Genet Metab. 2014;

12.   Kuivaniemi H, Tromp G, Prockop DJ. Genetic causes of aortic aneurysms.

Unlearning at least part of what the textbooks say. J Clin Invest.

1991;88(5):1441.

13.   Tompson SW, Faqeih EA, Ala-Kokko L, Hecht JT, Miki R, Funari T, et al.

Dominant and recessive forms of fibrochondrogenesis resulting from mutations

at a second locus, COL11A2. Am J Med Genet A. 2012 Feb;158A(2):309–14.

14.   Seegmiller RE, C. Fraser F, Sheldon H. A new chondrodystrophic mutant in

mice. Vol. 48, The Journal of Cell Biology. 1971. 580-593 p.

15.   Fernandes RJ, Weis M, Scott MA, Seegmiller RE, Eyre DR. Collagen XI chain

misassembly in cartilage of the chondrodysplasia (cho) mouse. Matrix Biol.

2007;

16.   Rodriguez RR, Seegmiller RE, Stark MR, Bridgewater LC. A type XI collagen

mutation leads to increased degradation of type II collagen in articular

cartilage11Supported by NIH Grants #AR46362 and #AR48839. Osteoarthr

Cartil. 2004 Apr;12(4):314–20.

17.   Xu L, Flahiff CM, Waldman BA, Wu D, Olsen BR, Setton LA, et al.

Osteoarthritis-like changes and decreased mechanical function of articular

cartilage in the joints of mice with the chondrodysplasia gene (cho). Arthritis

Rheum. 2003 Sep 1;48(9):2509–18.

18.   Holyoak DT, Otero M, Armar NS, Ziemian SN, Otto A, Cullinane D, et al.

Collagen XI mutation lowers susceptibility to load-induced cartilage damage in

mice. J Orthop Res. 2017 Oct 31;36(2):711–20.

19.   McGuirt WT, Prasad SD, Griffith AJ, Kunst HPM, Green GE, Shpargel KB, et

al. Mutations in COL11A2 cause non-syndromic hearing loss (DFNA13). Nat

Genet. 1999 Dec 1;23(4):413–9.

20.   Luo YY, Karsdal MA. Type XI collagen. In: Biochemistry of Collagens,

Laminins and Elastin. Academic Press; 2016. p. 77–80.

21.   Chen Y, Chen M, Gaffney EA, Brown CP. Effect of crosslinking in cartilage-like

collagen microstructures. J Mech Behav Biomed Mater. 2017 Feb 1;66:138–

43.

22.   Radlanski RJ, Renz H. Genes, forces, and forms: Mechanical aspects of

prenatal craniofacial development. Dev Dyn. 2006 May;235(5):1219–29.

23.   Alexopoulos LG, Haider MA, Vail TP, Guilak F. Alterations in the mechanical

properties of the human chondrocyte pericellular matrix with osteoarthritis. J

Biomech Eng. 2003 Jun;125(3):323–33.

24. Waarsing JH, Kloppenburg M, Slagboom PE, Kroon HM, Houwing-

Duistermaat JJ, Weinans H, et al. Osteoarthritis susceptibility genes influence

the association between hip morphology and osteoarthritis. Arthritis Rheum.

2011 May;63(5):1349–54.

25. Baird DA, Paternoster L, Gregory JS, Faber BG, Saunders FR, Giuraniuc C V.,

et al. Investigation of the relationship between susceptibility loci for hip

osteoarthritis and DXA-derived hip shape in a population based cohort of peri-

menopausal women. Arthritis Rheumatol. 2018 Jun 8;

26. Verbruggen SW, Kainz B, Shelmerdine SC, Hajnal J V., Rutherford MA,

Arthurs OJ, et al. Stresses and strains on the human fetal skeleton during

development. J R Soc Interface. 2018 Jan 24;15(138):20170593.

27. Hammond CL, Moro E. Using transgenic reporters to visualize bone and

cartilage signaling during development in vivo. Front Endocrinol (Lausanne).

2012;3:91.

28. Witten PE, Harris MP, Huysseune A, Winkler C. Small teleost fish provide new

insights into human skeletal diseases. In: Methods in cell biology. 2017. p.

321–46.

29. Eames B, DeLaurier A, Ullmann B, Huycke TR, Nichols JT, Dowd J, et al.

FishFace: interactive atlas of zebrafish craniofacial development at cellular

resolution. BMC Dev Biol. 2013 May 28;13(1):23.

30. Schilling TF, Le Pabic P. Fishing for the signals that pattern the face. J Biol.

2009;8(11):101.

31.   Askary A, Xu P, Barske L, Bay M, Bump P, Balczerski B, et al. Genome-wide

784   analysis of facial skeletal regionalization in zebrafish. Development. 2017 Aug

785   15;144(16):2994–3005.

786   32.   Askary A, Smeeton J, Paul S, Schindler S, Braasch I, Ellis NA, et al. Ancient

787   origin of lubricated joints in bony vertebrates. Elife. 2016;

788   33.   Brunt LH, Skinner REH, Roddy KA, Araujo NM, Rayfield EJ, Hammond CL.

789   Differential effects of altered patterns of movement and strain on joint cell

790   behaviour and skeletal morphogenesis. Osteoarthr Cartil. 2016 Nov

791   1;24(11):1940–50.

792   34.   Brunt LH, Norton JL, Bright JA, Rayfield EJ, Hammond CL. Finite element

793   modelling predicts changes in joint shape and cell behaviour due to loss of

794   muscle strain in jaw development. J Biomech. 2015 Sep 18;48(12):3112–22.

795   35.   Baas D, Malbouyres M, Haftek-Terreau Z, Le Guellec D, Ruggiero F.

796   Craniofacial cartilage morphogenesis requires zebrafish col11a1 activity.

797   Matrix Biol. 2009 Oct 1;28(8):490–502.

798   36.   Westerfield M. The Zebrafish Book: A Guide for the Laboratory Use of

799   Zebrafish (danio rerio). 4th ed. University of Oregon Press; 2000.

800   37.   Hammond CL, Schulte-Merker S. Two populations of endochondral

801   osteoblasts with differential sensitivity to Hedgehog signalling. Development.

802   2009 Dec 1;136(23):3991–4000.

803   38.   Elworthy S, Hargrave M, Knight R, Mebus K, Ingham PW. Expression of

804   multiple slow myosin heavy chain genes reveals a diversity of zebrafish slow

805   twitch muscle fibres with differing requirements for Hedgehog and Prdm1

activity. Development. 2008 Jun 15;135(12):2115–26.

39.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et

al. Fiji: an open-source platform for biological-image analysis. Nat Methods.

2012 Jun 28;9(7):676–82.

40.   Walker M, Kimmel C. A two-color acid-free cartilage and bone stain for

zebrafish larvae. Biotech Histochem. 2007 Jan 12;82(1):23–8.

41.   Thisse C, Thisse B. High-resolution in situ hybridization to whole-mount

zebrafish embryos. Nat Protoc. 2008 Jan 1;3(1):59–69.

42.   Fang M, Adams JS, McMahan BL, Brown RJ, Oxford JT. The expression

patterns of minor fibrillar collagens during development in zebrafish. Gene

Expr Patterns. 2010;

43.   Brunt LH, Roddy KA, Rayfield EJ, Hammond CL. Building Finite Element

Models to Investigate Zebrafish Jaw Biomechanics Video Link. J Vis Exp.

2016;54811(11810).

44.   Glasson SS, Chambers MG, Van Den Berg WB, Little CB. The OARSI

histopathology initiative – recommendations for histological assessments of

osteoarthritis in the mouse. Osteoarthr Cartil. 2010 Oct;18:S17–23.

45.   Chen X, Nadiarynkh O, Plotnikov S, Campagnola PJ. Second harmonic

generation microscopy for quantitative analysis of collagen fibrillar structure.

Nat Protoc. 2012 Mar 8;7(4):654–69.

46.   Blaschke UK, Eikenberry EF, Hulmes DJ, Galla HJ, Bruckner P. Collagen XI

nucleates self-assembly and limits lateral growth of cartilage fibrils. J Biol

Chem. 2000 Apr 7;275(14):10370–8.

47.   Robin NH, Moran RT, Ala-Kokko L. Stickler Syndrome. GeneReviews®.

830   University of Washington, Seattle; 1993.

831   48.   Hunt NCA, Vujani GM. Fibrochondrogenesis in a 17-week fetus: A case

832   expanding the phenotype. Am J Med Genet. 1998 Jan 23;75(3):326–9.

833   49.   Randrianaivo H, Haddad G, Roman H, Lise A, Toutain A, Le Merrer M, et al.

834   Fetal fibrochondrogenesis at 26 weeks' gestation. Prenat Diagn. 2002 Sep

835   1;22(9):806–10.

836   50.   Clément A, Wiweger M, von der Hardt S, Rusch MA, Selleck SB, Chien C-B, et

837   al. Regulation of Zebrafish Skeletogenesis by ext2/dackel and

838   papst1/pinscher. Mullins M, editor. PLoS Genet. 2008 Jul 25;4(7):e1000136.

839   51.   Schmid TM, Bonen DK, Luchene L, Linsenmayer TF. Late events in

840   chondrocyte differentiation: hypertrophy, type X collagen synthesis and matrix

841   calcification. In Vivo. 5(5):533–40.

842   52.   Eames BF, Amores A, Yan Y-L, Postlethwait JH. Evolution of the osteoblast:

843   skeletogenesis in gar and zebrafish. BMC Evol Biol. 2012 Mar 5;12(1):27.

844   53.   Mitchell RE, Huitema LFA, Skinner REH, Brunt LH, Severn C, Schulte-Merker

845   S, et al. New tools for studying osteoarthritis genetics in zebrafish. Osteoarthr

846   Cartil. 2013 Feb;21(2):269–78.

847   54.   Miosge N, Hartmann M, Maelicke C, Herken R. Expression of collagen type I

848   and type II in consecutive stages of human osteoarthritis. Histochem Cell Biol.

849   2004 Sep 17;122(3):229–36.

850   55.   Roddy KA, Skinner RE, Brunt LH, Kague E, Cross S, Rayfield EJ, et al. A

851   zebrafish model of developmental joint dysplasia: Manipulating the larval 1

mechanical environment to drive the malformation and recovery of joint shape.

56. Stickler GB, Belau PG, Farrell FJ, Jones JD, Pugh DG, Steinberg AG, et al.

Hereditary progressive arthro-ophthalmopathy. Mayo Clin Proc. 1965;40.

57. Hernandez LP, Barresi MJF, Devoto SH. Functional Morphology and

Developmental Biology of Zebrafish: Reciprocal Illumination from an Unlikely

Couple. Integr Comp Biol. 2002 Apr 1;42(2):222–31.

58. Buckwalter JA, Anderson DD, Brown TD, Tochigi Y, Martin JA. The Roles of

Mechanical Stresses in the Pathogenesis of Osteoarthritis: Implications for

Treatment of Joint Injuries. Cartilage. 2013 Oct 1;4(4):286–94.

59. Blaschke UK, Eikenberry EF, Hulmes DJ, Galla HJ, Bruckner P. Collagen XI

nucleates self-assembly and limits lateral growth of cartilage fibrils. J Biol

Chem. 2000 Apr;275(14):10370–8.

60. Scanzello CR, Goldring SR. The role of synovitis in osteoarthritis

pathogenesis. Bone. 2012 Aug;51(2):249–57.

61. Sokolove J, Lepus CM. Role of inflammation in the pathogenesis of

osteoarthritis: latest findings and interpretations. Ther Adv Musculoskelet Dis.

2013 Apr;5(2):77–94.

62. Mansour J. Biomechanics of cartilage. In: Kinesiology; the mechanics and

pathomechanics of human movement. Oatis CA, editor. Baltimore: Lippincott

Williams & Wilkins; 2003. 1992-6 p.

63. Hall BK (Brian K. Bones and cartilage : developmental and evolutionary

skeletal biology. 892 p.

64. Prein C, Warmbold N, Farkas Z, Schieker M, Aszodi A, Clausen-Schaumann

H. Structural and mechanical properties of the proliferative zone of the

developing murine growth plate cartilage assessed by atomic force

microscopy. Matrix Biol. 2016 Mar;50:1–15.

65. Radhakrishnan P, Lewis NT, Mao JJ. Zone-specific micromechanical

properties of the extracellular matrices of growth plate cartilage. Ann Biomed

Eng. 2004;(32):284–91.

66. Franz T, Hasler EM, Hagg R, Weiler C, Jakob RP, Mainil-Varlet P. In situ

compressive stiffness, biochemical composition, and structural integrity of

articular cartilage of the human knee joint. Osteoarthr Cartil. 2001 Aug

1;9(6):582–92.

67. Geraghty B, Jones SW, Rama P, Akhtar R, Elsheikh A. Age-related variations

in the biomechanical properties of human sclera. J Mech Behav Biomed Mater.

2012 Dec;16:181–91.

68. Hayes AJ, Reynolds S, Nowell MA, Meakin LB, Habicher J, Ledin J, et al.

Spinal Deformity in Aged Zebrafish Is Accompanied by Degenerative Changes

to Their Vertebrae that Resemble Osteoarthritis. Heymann D, editor. PLoS

One. 2013 Sep 24;8(9):e75787.

69. Askary A, Smeeton J, Paul S, Schindler S, Braasch I, Ellis NA, et al. Ancient

origin of lubricated joints in bony vertebrates. Elife. 2016 Jul 19;5.

70. Naveen SV, Ahmad RE, Hui WJ, Suhaeb AM, Murali MR, Shanmugam R, et

al. Histology, glycosaminoglycan level and cartilage stiffness in

monoiodoacetate-induced osteoarthritis: comparative analysis with anterior

cruciate ligament transection in rat model and human osteoarthritis. Int J Med

Sci. 2014;11(1):97–105.

71.   Sharma AR, Jagga S, Lee S-S, Nam J-S. Interplay between cartilage and

900   subchondral bone contributing to pathogenesis of osteoarthritis. Int J Mol Sci.

901   2013 Sep 30;14(10):19805–30.

902   72.   Pollard AS, Boyd S, McGonnell IM, Pitsillides AA. The role of embryo

903   movement in the development of the furcula. J Anat. 2017 Mar;230(3):435–43.

904   73.   Pitsillides AA. Early effects of embryonic movement: "a shot out of the dark." J

905   Anat. 2006 Apr;208(4):417–31.

906   74.   Rolfe RA, Bezer JH, Kim T, Zaidon AZ, Oyen ML, Iatridis JC, et al. Abnormal

907   fetal muscle forces result in defects in spinal curvature and alterations in

908   vertebral segmentation and shape. J Orthop Res. 2017 Oct;35(10):2135–44.

909   75.   Nowlan NC, Chandaria V, Sharpe J. Immobilized chicks as a model system for

910   early-onset developmental dysplasia of the hip. J Orthop Res. 2014

911   Jun;32(6):777–85.

912   76.   Roddy KA, Prendergast PJ, Murphy P. Mechanical Influences on

913   Morphogenesis of the Knee Joint Revealed through Morphological, Molecular

914   and Computational Analysis of Immobilised Embryos. Agarwal S, editor. PLoS

915   One. 2011 Feb 28;6(2):e17526.

916   77.   Kahn J, Shwartz Y, Blitz E, Krief S, Sharir A, Breitel DA, et al. Muscle

917   Contraction Is Necessary to Maintain Joint Progenitor Cell Fate. Dev Cell.

918   2009 May;16(5):734–43.

919   78.   Shwartz Y, Viukov S, Krief S, Zelzer E. Joint Development Involves a

920   Continuous Influx of Gdf5-Positive Cells. Cell Rep. 2016 Jun 21;15(12):2577–

921   87.

79. Lee W, Leddy HA, Chen Y, Lee SH, Zelenski NA, McNulty AL, et al. Synergy between Piezo1 and Piezo2 channels confers high-strain mechanosensitivity to articular cartilage. Proc Natl Acad Sci U S A. 2014 Nov 25;111(47):E5114-22.

80. Karystinou A, Roelofs AJ, Neve A, Cantatore FP, Wackerhage H, De Bari C. Yes-associated protein (YAP) is a negative regulator of chondrogenesis in mesenchymal stem cells. Arthritis Res Ther. 2015 Dec 30;17(1):147.

81. Deng Y, Wu A, Li P, Li G, Qin L, Song H, et al. Yap1 Regulates Multiple Steps of Chondrocyte Differentiation during Skeletal Development and Bone Repair. Cell Rep. 2016 Mar 8;14(9):2224–37.

82. Rolfe RA, Nowlan NC, Kenny EM, Cormican P, Morris DW, Prendergast PJ, et al. Identification of mechanosensitive genes during skeletal development: alteration of genes associated with cytoskeletal rearrangement and cell signalling pathways. BMC Genomics. 2014 Jan 20;15(1):48.

83. Brunt LH, Begg K, Kague E, Cross S, Hammond CL, Kague E, et al. Wnt Signalling Controls the Response to Mechanical Loading during Zebrafish Joint Development. Development. 2017;144(15):2798–809.

84. Singh PNP, Shea CA, Sonker SK, Rolfe RA, Ray A, Kumar S, et al. Precise spatial restriction of BMP signaling in developing joints is perturbed upon loss of embryo movement. Development. 2018 Mar 1;145(5):dev153460.

85. García-Ibarbia C, Pérez-Núñez MI, Olmos JM, Valero C, Pérez-Aguilar MD, Hernández JL, et al. Missense polymorphisms of the WNT16 gene are associated with bone mass, hip geometry and fractures. Osteoporos Int. 2013 Sep 16;24(9):2449–54.

86. Movérare-Skrtic S, Henning P, Liu X, Nagano K, Saito H, Börjesson AE, et al. Osteoblast-derived WNT16 represses osteoclastogenesis and prevents cortical bone fragility fractures. Nat Med. 2014 Nov 12;20(11):1279–88.

87. Ohlsson C, Henning P, Nilsson KH, Wu J, Gustafsson KL, Sjögren K, et al. Inducible *Wnt16* inactivation: WNT16 regulates cortical bone thickness in adult mice. J Endocrinol. 2018 May;237(2):113–22.

88. Nalesso G, Thomas BL, Sherwood JC, Yu J, Addimanda O, Eldridge SE, et al. WNT16 antagonises excessive canonical WNT activation and protects cartilage in osteoarthritis. Ann Rheum Dis. 2017 Jan;76(1):218–26.

89. Dell'Accio F, De Bari C, Eltawil NM, Vanhummelen P, Pitzalis C. Identification of the molecular response of articular cartilage to injury, by microarray screening: Wnt-16 expression and signaling after injury and in osteoarthritis. Arthritis Rheum. 2008 Apr 25;58(5):1410–21.

90. Sirko-Osadsa DA, Murray MA, Scott JA, Lavery MA, Warman ML, Robin NH. Stickler syndrome without eye involvement is caused by mutations in COL11A2, the gene encoding the α2(XI) chain of type XI collagen. J Pediatr. 1998 Feb 1;132(2):368–71.

91. Shwartz Y, Farkas Z, Stern T, Aszódi A, Zelzer E. Muscle contraction controls skeletal morphogenesis through regulation of chondrocyte convergent extension. Dev Biol. 2012 Oct 1;370(1):154–63.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6