Comparative architectures of direct and social genetic effects from the genome-wide association study of 170 phenotypes in outbred laboratory mice

Amelie Baud[1*], Francesco Paolo Casale[1,2], Jerome Nicod[3], Oliver Stegle[1,4*]

[1] European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, CB10 1SD Hinxton, Cambridge, UK

[2] Microsoft Research New England, Cambridge, Massachusetts, USA

[3] Wellcome Trust Centre for Human Genetics, Roosevelt Drive, Oxford OX3 7BN, UK

[4] European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, DE

* Corresponding authors: abaud@ebi.ac.uk and oliver.stegle@embl.de

Abstract

Social genetic effects (SGE, also called indirect genetic effects) are associations between genotypes of one individual and phenotype of another. SGE can arise when two individuals interact and heritable traits of one individual influence the phenotype of the other. To better understand the architecture of SGE, we re-analysed an existing dataset comprising 170 behavioural, physiological and morphological phenotypes measured in outbred laboratory mice. For all phenotypes and in order to compare SGE with better-known direct genetic effects (DGE, associations between an individual's genotypes and their own phenotype), we analysed polygenic models with random terms for SGE and DGE and performed the genome-wide association study of both SGE (sgeGWAS) and DGE (dgeGWAS). Our analyses yielded two main insights: first, SGE and DGE *acting on the same phenotype* generally arise from partially different loci and/or loci with different effect sizes; secondly, individual SGE associations typically explain less phenotypic variance than DGE associations. Our results shed light on the architecture of SGE and have important implications for the design of future studies. Importantly, we detail and validate methods that can be used for sgeGWAS

33 in outbred populations with any levels of relatedness and group sizes, and provide
34 software to perform these analyses.

Main text

Introduction

Social interactions between two individuals can result in the phenotype of one
individual being affected by genotypes of their social partner. Such effects arise when
heritable traits of the social partner influence the phenotype of interest (Figure 1a),
and are called indirect genetic effects or social genetic effects[1-4] (SGE).

SGE have been shown to contribute significantly and substantially to
phenotypic variation in livestock, wild animals, plants and laboratory model organisms
(see review by Bijma[5] and subsequent references). In laboratory mice, SGE have
been found to affect a broad range of phenotypes including behavioural, physiological,
and morphological traits[6-9], and in humans effects of non-transmitted parental alleles
have been detected on offspring's educational attainment[10-12]. Thus, SGE are an
important component of the genotype to phenotype path, and understanding their
architecture is important.

SGE can be used as a tool to identify traits of social partners affecting a
phenotype of interest. For example, a candidate gene study of SGE on plumage
condition in laying hens[13] found an indirect association with the serotonin receptor 2C
gene. As the serotoninergic system is known to control behaviour, this SGE
association is consistent with observations of cage mates influencing the plumage of
a focal hen through feather pecking. When traits of social partners affecting the
phenotype of interest are unknown, the genome-wide association study of SGE
(sgeGWAS) may be a promising avenue. Indeed, similarly to how GWAS of
"traditional" direct genetic effects (DGE, effects of an individual's genotypes on its own
phenotype) has provided valuable insights into the "within-body" pathways affecting
disease and quantitative phenotypes[14], sgeGWAS could help dissect the "between-
bodies" pathways of social effects and provide clues on the traits of social partners

that mediate social effects. Deploying sgeGWAS reliably and efficiently will however
require a good understanding of the architecture of SGE.

Information on the architecture of SGE is relatively sparse, as few studies have
identified genomic loci that influence phenotypes of social partners. Quasi-Mendelian
SGE exist[15,16] but candidate gene studies and GWAS of SGE have revealed oligo- or
polygenic architectures for a larger number of phenotypes[7,8,13,17,18]. Two key features
of such complex architectures are the degree of overlap between SGE and DGE loci,
and the proportion of variance explained by SGE loci. The degree of overlap between
SGE and DGE has most often been studied in terms of the genome-wide correlation
between SGE and DGE effect sizes, and it has been shown to be an important
determinant of the response of a phenotype to selection[19-24]. In addition, whether the
strongest SGE and DGE for any given phenotype arise from the same loci is of
practical interest: it determines how redundant sgeGWAS may be with dgeGWAS of
the same phenotype and whether loci identified in dgeGWAS may also have effects
on the same phenotype of partners.

Similarly, the variance explained by individual SGE loci is both of fundamental
and practical interest, as it provides insights into the evolutionary process and
determines the power of sgeGWAS. Estimates of the variance explained by individual
SGE loci are most informative when similar estimates for DGE loci acting on the same
phenotype and in the same population are available for comparison. Of the few studies
that have mapped SGE[7,8,13,17,18,25], only two have reported SGE and DGE loci *acting*
on the same phenotype. One of these reported that DGE loci explain a much greater
proportion of phenotypic variance than SGE loci[17] while the other found similar effect
sizes for individual SGE and DGE loci[8]. Given the sparsity of information, expectations
on the effect sizes of SGE are difficult to build.

In order to improve our understanding of the architecture of SGE, we leveraged
an existing dataset of 170 behavioural, physiological and morphological phenotypes
measured in outbred "CFW" laboratory mice[26,27] (Figure 1c). We studied both SGE
and DGE acting on each of these 170 phenotypes (Figure 1b) using polygenic models
and GWAS. Specifically, we investigated whether SGE and DGE arise from similar
loci, and compared the effect sizes of SGE loci to those of DGE loci.

Figure 1 *Illustration of social genetic effects (SGE), definition of specific terms used*

throughout the manuscript, and experimental design. (a) SGE arise when two

individuals interact and heritable traits of one influence the phenotype of the other. In

other words, genotypes of the "social partner" influence - through "mediating traits" -

the "phenotype of interest", measured in the "focal individual". Importantly, these

mediating traits are not measured. (b) 1,812 "CFW" outbred mice were housed in

groups of 3 mice per cage. SGE and DGE contributed by each mouse were modelled,

such that each mouse served as both focal individual and cage mate in our analyses.

(c) Housing conditions and phenotyping.

108
Results

Genome-wide genotypes and 200 behavioural, physiological and morphological

phenotypes for 2,073 commercially-available, outbred CFW mice were available from

Nicod et al.[26] and Davies et al.[27]. Males were always housed with males and females

with females, and mice were left undisturbed in their cages for at least nine weeks

before phenotyping started (Figure 1c). We only kept mice that had the same two cage

mates over the course of the experiment (1,869 total). Furthermore, we excluded 57

mice that formed genetic substructures (see Methods) so that the remaining 1,812

mice were as equally related as possible while retaining as large a sample size as

possible. We analysed a subset of 170 phenotypes that could be satisfactorily
normalised (see Methods). The exact number of mice used for each phenotype, is
shown in Supplementary Table 1.
122

Aggregate contribution of SGE

We first estimated the aggregate contribution of SGE to each phenotype (i.e. the sum
of SGE across the genome). To do so, we used the variance decomposition method
detailed in Baud et al.[9], which features random effects for DGE, SGE, direct and social
environmental effects, and "cage effects" (see Methods). SGE, in aggregate,
explained up to 22% (+/- 6%) of variation in serum LDL levels and an average of 11%
across 9 phenotypes with significant aggregate SGE (FDR < 10%, Supplementary
Table 1). Those 9 phenotypes included behavioural (helplessness, a murine model for
depression), physiological (serum LDL cholesterol, wound healing, blood eosinophils,
serum alpha-amylase concentration, blood platelets, acute hypoxic response), and
morphological (weight of adrenal glands) traits. For many of these phenotypes, the
pathways underlying social effects, i.e. the traits of cage mates that mediate social
effects, are unknown.
136

Overlap between SGE and DGE loci acting on the same phenotype

The polygenic model used for variance decomposition fits a correlation coefficient ρ
that measures the correlation between the SGE and DGE random effects of the model
(see Methods). Thus, ρ quantifies the extent to which SGE and DGE acting on the
same phenotype arise from similar loci, with similar effect sizes. We stress that, in this
section and throughout the manuscript, we compare SGE and DGE *acting on the*
same phenotype (but do this for all 170 phenotypes). Simulations showed that the
precision with which ρ can be estimated depends on the aggregate contribution of both
SGE and DGE (Supplementary Figure 1), so we limited the analysis of ρ to 28
phenotypes for which both aggregate SGE and aggregate DGE explained more than
5% of phenotypic variation. The value of ρ varied between -0.28 (+/-0.35) and 1(+/-
0.09) across these traits, with an average of 0.42 (Figure 2 and Supplementary Table
1). For 10 out of the 28 phenotypes where ρ could be more precisely estimated, ρ was
significantly different from zero (nominal P < 0.05), suggesting that loci affecting a
phenotype directly also sometimes influence the same phenotype of cage mates. The

strongest evidence for shared SGE and DGE loci ($\rho \neq 0$ at Bonferroni-corrected P <

0.05) was for healing from an ear punch, weight of the adrenal glands, serum LDL

cholesterol levels, and mean platelet volume.

We also evaluated evidence that $|\rho|$ was different from one (i.e. ρ different from

one and minus one) in order to empirically evaluate the widely-influential model of

"phenotypic contagion". Phenotypic contagion or "spread" is a model for social effects

whereby the phenotype of interest of a focal individual is affected by the same

phenotype of their social partners. In humans, cognitive susceptibility to depression,

alcohol consumption, stress, obesity and educational attainment, for example, have

been shown to be "contagious" or "spread" across college roommates, spouses,

friends, or parent/offspring[10-12,28-32]. As a result, phenotypic contagion has shaped the

way we think about social effects: for example, phenotypes unlikely to spread have

been used to cast doubt on social network effects[33]. Here we leveraged the parameter

ρ to test whether phenotypic contagion was sufficient to account for SGE: under a

model of pure phenotypic contagion, $|\rho|$ is expected to be equal to one; on the contrary,

if traits of social partners other than the phenotype of interest mediate social effects,

$|\rho|$ is expected to be different from one. We found that $|\rho|$ was significantly different

from one (nominal P < 0.05) for 10 out of 28 phenotypes. The most significant P value

(0.00066, significant after Bonferroni correction) was found for immobility in the first

two minutes of the Porsolt swim test, a measure of helplessness that is relevant to

depression. This latter result suggests that phenotypes that spread may additionally

be affected by other traits of social partners. These results motivate the use of

sgeGWAS as a tool to more broadly understand social effects.

Figure 2 Correlation ρ between SGE and DGE random effects (see Methods). The 28
phenotypes included in this table are those for which the correlation ρ could be more
precisely estimated, i.e. phenotypes with aggregate SGE and aggregate DGE > 5%.
The bars show the standard errors. The stars represent the P value for rejecting H_0: ρ
= 0 (bottom) and H_0: $|\rho|$ = 1 (i.e. pure phenotypic contagion, top). * denotes nominal
P value < 0.05, * denotes Bonferroni-corrected P value < 0.05.

sgeGWAS and dgeGWAS of 170 phenotypes

To map SGE and following Biscarini et al.[13] and Brinker et al.[25], we calculated the
"social genotype" of a mouse at a variant as the sum of the reference allele dosages
of its cage mates at the variant, and tested for association between social genotype
and phenotype. In order to avoid spurious associations, we accounted for background
SGE, DGE and non-genetic effects using an extension of the variance components
model used for variance decomposition. In the sgeGWAS we also accounted for DGE
of the variant tested for SGE, by including direct genotypes at the locus as a covariate
(See Methods). Similarly, in the dgeGWAS we included social genotypes at the locus
as a covariate. We hereafter refer to this strategy as "conditioning". We found that
conditioning was necessary to avoid spurious associations in the sgeGWAS due to
co-localised DGE. As we show in the Supplementary Note, this problem originates
from the use of each mouse as both focal individual and cage mate in the analysis, a

strategy that has been used before to maximise sample size when all individuals are
phenotyped and genotyped[13,25]. Importantly, spurious associations may arise even if
all individuals are strictly unrelated (Supplementary Note). Using each mouse as both
focal individual and cage mate in the analysis results in direct and social genotypes at
a locus being correlated (Supplementary Figures 2a and 2b), which leads to
sgeGWAS P values being inflated under the null in the presence of a simulated, co-
localised large-effect DGE (Supplementary Figure 2c). This issue has not previously
been reported and may have resulted in spurious SGE associations when conditioning
was not used[13].

We show that conditioning on direct genotypes at the locus yielded calibrated
sgeGWAS P values for null phenotypes (Supplementary Figure 2d), indicating that
genome-wide significance thresholds may be derived for sgeGWAS by permuting
social genotypes (see Methods), as long as conditioning is used in the analysis. A
power analysis suggested that conditioning may slightly decrease power to detect
SGE in the absence of co-localised DGE, particularly when direct and social
genotypes are highly correlated (Supplementary Figure 3a and 3c) but would increase
power if the locus also gave rise to DGE (Supplementary Figures 3b and 3d).

In order to compare, for each phenotype, the results of sgeGWAS and
dgeGWAS, we defined loci based on the average size of the 95% confidence interval
in this population, namely 1.5Mb[26], and, following Nicod et al.[26], used a per-phenotype
FDR approach (see Methods). At a 10% FDR threshold, sgeGWAS identified 24
genome-wide significant loci for 17 of the 170 phenotypes (Figure 3 and
Supplementary Table 2). In comparison, dgeGWAS identified 121 genome-wide
significant loci for 63 phenotypes at the same threshold (Figure 3 and Supplementary
Table 3).

There was no overlap between genome-wide significant SGE and DGE loci
acting on the same phenotype. However, variants at genome-wide significant SGE
loci were enriched in small P values in the corresponding dgeGWAS (Supplementary
Figure 4). Together these results suggest a partially distinct basis for SGE and DGE
acting on the same phenotype (i.e. partially different loci and/or effect sizes), which is
consistent with the results from the analysis of the correlation parameter ρ.

Figure 3 *Superimposed manhattan plots corresponding to 170 sgeGWAS (top panel)*
228

and 170 dgeGWAS (bottom panel) of the same phenotypes. DGE associations with a
229

–logP greater than 10 were plotted at –logP 10 (as indicated by 10+). Data points with
230

negative log P < 2 are not shown. Lead variants for all genome-wide significant SGE
231

and DGE loci are represented with a larger dot. In the SGE panel, each color
232

corresponds to a class of phenotypes: behavioural (red, includes 7 behavioural
233

phenotypes with a detected SGE locus), adult neurogenesis (black, 2 phenotypes),
234

immune (orange, 1 phenotype), haematological (yellow, 1 phenotype), blood
235

biochemistry (blue, 2 phenotypes), bone phenotypes (green, 2 phenotypes), heart
236

function (brown, 1 phenotype), and lung function (purple, 1 phenotype). In the DGE
237

panel, a genome-wide significant locus is colored grey when the corresponding
238

phenotype does not have a genome-wide significant SGE association; when the
239

corresponding phenotype does have an SGE association, the same color is used as
240

in the SGE panel.
241

Compared to many other mouse populations used for mapping, linkage
243

disequilibrium decays rapidly in the CFW population[26,34]. At each genome-wide
244

significant SGE locus we identified candidate genes, prioritising well-annotated genes

(see Methods). At five genome-wide significant SGE loci we identified a single

candidate gene (Supplementary Table 2, locus zoom plots in Supplementary Figure

5): *Abca12*, a gene known for its involvement in lipid transport and homeostasis in the

skin[35], at an SGE locus for adult neurogenesis in the hippocampus; *Epha4*, a

signalling genes involved in neural system function, at an SGE locus for helplessness;

H60c, a poorly characterised gene potentially involved in skin immunity[36], at an SGE

locus for locomotor activity; *Pgk1-rs7*, a pseudogene of phosphoglycerate kinase-1, at

and SGE locus for sleep; and *Ighv5-9-1*, a variable region of the T cell receptor, at an

SGE locus for response to hypoxia. None of these genes have known direct effects

that can easily explain the observed SGE, nor did they seem to have DGE on the

phenotype in this dataset (Supplementary Figure 5), so the results of our sgeGWAS

point to yet unknown traits of cage mates that influence the five phenotypes above.

Of these candidate genes, one, *Epha4*, has previously been associated with

the phenotype of interest. *Epha4* expression in the hippocampus was found to be

affected by chronic mild stress in mice and responsive to antidepressant treatment[37].

We also found suggestive DGE of *Epha4* on helplessness (Supplementary Figure 5),

confirming that some level of phenotypic contagion was likely for that phenotype. The

other candidate genes did not immediately permit to generate hypotheses on the traits

of cage mates mediating the social effects. To gain such insights from the results of

sgeGWAS, it is likely that other data types (e.g. gene expression) will need to be

integrated. Alternatively, larger sample sizes would permit identification of additional

SGE loci, some of which might immediately provide insights into the traits of partners

that mediate social effects. SgeGWAS, in that respect, is similar to dgeGWAS[14,38,39].

Architecture of SGE and comparison with that of DGE

Despite being carried out on the same individuals and phenotypes, and in a perfectly

analogous manner, sgeGWAS identified fewer genome-wide significant associations

than dgeGWAS (24 associations for 17 phenotypes and 121 associations for 63

phenotypes respectively). As the determinants of power for SGE have not been

investigated, it is not clear whether we had more or less power to detect SGE

associations compared to DGE associations. In order to get a better understanding of

this issue, we simulated local SGE or DGE arising from a single causal variant and

calculated power to detect these associations. Briefly (see Methods), we considered

random groups of two or three mice per cage, and simulated phenotypes arising from
the sum of local genetic effects (DGE or SGE), polygenic effects (DGE and SGE), and
non-genetic effects. We simulated local SGE according to two alternative generative
models, both consistent with the analysis model used for sgeGWAS: an "additive"
model according to which social effects add up across cage mates, and a
"proportional" model corresponding to a scenario where the focal mouse interacts with
only one cage mate at a time, spending equal time with each cage mate. Note that the
additive and proportional models are equivalent when there is a single cage mate (i.e.
two mice per cage). For all three types of local effects (DGE, additive SGE and
proportional SGE) we simulated the same allelic effect. Finally, we considered variants
with low, medium or high minor (direct) minor allele frequencies (MAF, see Methods).

Our simulations showed that power always increased with MAF (Figure 3a). At
a given MAF, simulating SGE from a single cage mate led to the same power for SGE
and DGE. Simulating SGE arising from two cage mates additively led to greater power
to detect SGE associations compared to DGE associations. In contrast, simulating
SGE arising from two cage mates under the proportional model led to lower power for
SGE compared to DGE. These results are consistent with the fact that, for a given
sample size, power to detect a local effect (DGE or SGE) is determined by the sample
variance of the simulated effect. Noting MAF as p, number of cage mates as N, and
allelic effect as b, that variance is expected to be $2p(1-p)b^2$ for DGE, $2Np(1-p)\ b^2$ for
SGE simulated under the additive model, and $2Np(1-p)/N^2\ b^2$ for SGE simulated under
the proportional model (see Methods). In conclusion, our simulations showed that
power to detect individual SGE associations is determined not only by allelic effect
and MAF of the causal variants, but also by the way SGE arise across cage mates
(additively or not) and the number of cage mates. In the real data, the way SGE arose
across cage mates is not known, so it is not possible to determine the primary cause
for the smaller number of genome-wide significant SGE associations compared to
DGE associations.

Comparing genome-wide significant SGE and DGE associations in terms of
proportion of phenotypic variance explained yielded two main results: firstly, individual
genome-wide significant SGE associations explained a maximum of 2.5% of
phenotypic variance, while eleven genome-wide significant DGE associations
explained more than 5% of phenotypic variance and up to 40% (Figure 3b,
Supplementary Table 2 and Supplementary Table 3). Average values were 1.8% for

SGE and 2.7% for DGE associations. As these results are born from the analysis of 170 phenotypes, it suggests that SGE associations will generally be more difficult to detect than DGE associations. Secondly, for each phenotype we compared the variance explained jointly by all genome-wide SGE (respectively DGE) associations to the variance explained by SGE (respectively DGE) in aggregate. We found that genome-wide significant associations explained a large proportion of the corresponding genetic variance for both SGE and DGE (Figure 3c). More precisely, across 5 phenotypes with aggregate contribution of SGE greater than 5% and at least one genome-wide significant SGE association, we found that an average of 32.5% of the aggregate variance was explained by genome-wide significant associations. For DGE, that figure was calculated across 55 phenotypes and was equal to 32.1%. The proportion of aggregate variance explained by genome-wide significant associations may seem large given the relatively small number of genome-wide significant associations per phenotype (e.g. compared to humans[40]), but is consistent with studies of DGE in other outbred laboratory rodent populations[41,42] and are the result of a relatively small number of variants segregating in the CFW population with relatively high MAFs[26]. In conclusion, our results are consistent with oligo- or polygenic architectures for SGE. A more precise estimation of the number of loci involved will only be possible when more SGE associations are discovered in other datasets.

Figure 3 *Power to detect local SGE and DGE, and characterisation of the architecture of SGE and DGE. (a) Power to local genetic effects in simulations. Three types of local genetic effects were simulated: DGE (or, equivalently, SGE arising from a single cage mate), SGE arising from two cage mates under an additive model, and SGE arising from two cage mates under a proportional model (see Main Text). For each type of effect, results are shown (left to right) for variants with low MAF (MAF < 0.05), medium MAF (0.225<MAF<0.275) and high MAF (MAF>0.45) (MAF: minor allele frequency,*

defined based on direct genotypes). (b) Histogram of the proportion of phenotypic

variance explained by individual genome-wide significant SGE (red) and DGE (black)

associations. (c) Comparison, for each phenotype, of the variance explained by social

(red) and direct (black) genetic effects in aggregate (x axis) and the total variance

explained jointly by all genome-wide significant SGE or DGE associations for a

phenotype (y axis). Each dot corresponds to a phenotype with at least one genome-

wide significant association.

Discussion

In this study we performed the comparative analysis of both SGE and DGE acting on

each one of 170 behavioural, physiological and morphological phenotypes measured

in outbred laboratory mice, using polygenic models and GWAS. Our results provided

two key insights into the architecture of these complex traits: first, SGE and DGE

acting on the same phenotype typically arise from partially different loci and/or loci with

different effect sizes; secondly, SGE associations tend to explain less phenotypic

variation than DGE associations. As we analysed a broad range of phenotypes, the

insights we gained are likely to generalize to other populations and phenotypes.

For 10 phenotypes we uncovered evidence that SGE and DGE were

significantly correlated. For example, ρ was significantly different from zero for the two

measures of helplessness included in this dataset. This result is consistent with prior

evidence that mood spreads across social partners[28,43]. It is also consistent with the

observation that, in this study, two out of the three genome-wide significant SGE loci

for helplessness have suggestive direct effects on helplessness - direct effect that are

further supported by prior reports that *Epha4*, the candidate gene at one of the loci, is

associated with depression and responds to antidepressant treatment[37]. The

pathways that mediate non-zero correlations between SGE and DGE for other

phenotypes were not always obvious (e.g. healing from an ear punch, serum LDL

cholesterol levels) but warrant further investigation of SGE and DGE.

A key result from our study is empirical evidence that phenotypic contagion is

often not sufficient to account for social effects, even when it does play a role. Indeed,

for 10 out of 28 tested phenotypes we found that $|\rho|$ was significantly different from

one, including the two aforementioned measures of helplessness. This result supports

efforts to discover other traits of social partners that mediate social effects, and points to sgeGWAS as a way to do so. It is important to bear in mind, however, that SGE only capture the genetic component of the traits of partners that mediate social effects. Hence, traits that are mostly non-genetically determined will be missed by SGE studies.

Our results on the variance explained by individual SGE loci are an important contribution towards understanding the architecture of SGE and will help design future experiments such as sgeGWAS. In particular, the fact that SGE loci never explained a large fraction of phenotypic variance (max 2.5%), while in comparison 11 DGE loci explained more than 5% of phenotypic variation, shows that sgeGWAS will require larger sample sizes than dgeGWAS to be equally powered.

Finally, our study made several important methodological contributions that will help design, perform and interpret sgeGWAS, particularly in outbred populations where both DGE and SGE contribute to phenotypic variation. Specifically, our study improved our understanding of the determinants of power for SGE and we showed that correlations between direct and social genotypes at a locus need to be accounted for to avoid spurious associations. These correlations arise when the same individuals serve as both focal individuals and social partners in the analysis, even if all individuals are unrelated. Importantly, similar correlations between direct and social genotypes, but potentially much stronger, may arise for different reasons in other datasets, notably when focal individuals and social partners are related, or as a result of direct assortments (e.g. assortative mating[44,45], homophily between friends[10]). The methods we presented here will help avoid spurious associations in such situations. Importantly, we contribute software and code to reproduce our analyses or analyse other datasets.

Methods

Phenotypes and experimental variables

Phenotypes and experimental variables (covariates) for 1,934 Crl:CFW(SW)-US_P08 (CFW) mice were retrieved from http://wp.cs.ucl.ac.uk/outbredmice/. We normalized

each phenotype using the boxcox function (MASS package[46]) in R, and excluded

phenotypes that could not be normalised satisfactorily (lambda outside of -2 to 2

interval). The subset of covariates used for each phenotype is indicated in

Supplementary Table 1. Because data for some phenotypes were missing for some

mice, the sample size varied. The sample size for each phenotype after all filtering

(see below) is indicated in Supplementary Table 1.

Caging information

Mice were four to seven weeks old when they arrived at the phenotyping facility. They

were grouped with their cage mates and then spent nine to twelve weeks undisturbed

in quarantine. They spent a further four weeks together during phenotyping. Males

were always housed with males and females with females.

Cage assignments were not included in the publicly available dataset but were

provided by the authors upon request and are now provided in Supplementary Table

4. Cage assignments were recorded at eleven time points throughout the study and

showed that a few mice were taken out of their original cages and singly housed,

presumably because they were too aggressive to their cage mates. When this

happened, we excluded all the mice in that cage from the analysis. We also excluded

cages where some of the mice were "genetically close" (as defined below) to many

other mice. Finally, we only retained cages with exactly three mice per cage. Although

from the sleep test on all mice were singly housed, we still investigated "persistent"

SGE on sleep and tissue phenotypes (persistence over one day for sleep phenotypes

and over a few days for tissue measures).

Genome-wide genotypes

From http://wp.cs.ucl.ac.uk/outbredmice/ we retrieved both allele dosages for 7 million

variants and allele dosages for a subset of 353,697 high quality, LD-pruned variants

(as described in Nicod et al.[26]). We used high quality, LD-pruned variants for all

analyses but the identification of candidate genes at SGE loci (see below), for which

we used the full set of variants.

Genetic relatedness matrix (GRM) and exclusion of "genetically close" mice

The genetic relatedness matrix was calculated as the cross-product of the dosage

matrix after standardizing the dosages for each variant to mean 0 and variance 1.

We excluded whole cages of mice based on GRM values as follows: we defined

a "close pair" of mice as having a GRM value greater than 0.3 (based on the histogram

of all GRM values). 199 mice in 145 cages were involved in such close pairs. Excluding

all 145 cages would have resulted in excluding 435 mice out of a total of 1,812, which

would have led to substantially reduced power for sgeGWAS and dgeGWAS. Thus,

we made a compromise and only excluded the 19 cages that were involved in 4 or

more close pairs (57 mice excluded).

Variance decomposition

The same method as described in details in Baud et al.[9] was used. Briefly, the model

used was:

$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c}$ (0)

y_f is the phenotypic value of the focal mouse f, X_f is a row of the matrix X of covariate

values and b a column vector of corresponding estimated coefficients. $\underline{a_{D,f}}$ is the

additive direct genetic effects (DGE) of f. Z_f is a row of the matrix Z that indicates

cage mates (importantly $Z_{i,i} = 0$) and $\underline{a_S}$ the column vector of additive social genetic

effects (SGE). $\underline{e_D}$ refers to direct environmental effects and $\underline{e_S}$ to social environmental

effects. W_f is a row of the matrix W that indicates cage assignment and c the column

vector of cage effects.

The joint distribution of all random effects is defined as:

466
$$
\begin{bmatrix} \underline{a_D} \\ \underline{a_S} \\ \underline{e_D} \\ \underline{e_S} \\ \underline{c} \end{bmatrix} \sim \text{MVN} \left(0, \begin{bmatrix} \sigma_{A_D}^2 A & \sigma_{A_{DS}} A & 0 & 0 & 0 \\ \sigma_{A_{DS}} A^T & \sigma_{A_S}^2 A & 0 & 0 & 0 \\ 0 & 0 & \sigma_{E_D}^2 I & \sigma_{E_{DS}} I & 0 \\ 0 & 0 & \sigma_{E_{DS}} I^T & \sigma_{E_S}^2 I & 0 \\ 0 & 0 & 0 & 0 & \sigma_C^2 I \end{bmatrix} \right)
$$

468 where A is the GRM and I the identity matrix.

469 The phenotypic covariance is:

470     $C_{i,j} = cov(y_i, y_j)$

471             $= \sigma_{A_D}^2 A_{i,j} + \sigma_{A_{DS}} + \sigma_{A_S}^2 (ZAZ^T)_{i,j} + \sigma_{E_D}^2 I_{i,j} + \sigma_{E_{DS}} \{ (IZ^T)_{i,j}$

472             $+ (ZI^T)_{i,j} \} + \sigma_{E_S}^2 (ZIZ^T)_{i,j} + \sigma_C^2 (WIW^T)_{i,j}$

The variances explained by DGE and SGE were calculated respectively as

$sampleVar(\sigma_{A_D}^2 A) / sampleVar(C)$ and $sampleVar(\sigma_{A_S}^2 (ZAZ^T)) / sampleVar(C)$

where $sampleVar$ is the sample variance of the corresponding covariance matrix:

suppose that we have a vector $\underline{x}$ of random variables with covariance matrix M, the

sample variance of M is calculated as

$sampleVar(M) = \frac{Tr(PMP)}{n-1}$

Tr denotes the trace, n is the sample size, and $P = I - \frac{11'}{n}$ [47,48].

For those phenotypes where body weight was included as a covariate, we checked

that this did not lead to systematically increased (or decreased) estimates of the

aggregate contribution of SGE (collider bias).

Significance of variance components was assessed using a two-degree of freedom

log likelihood ratio (LLR) test (i.e., the test statistics was assumed to follow a two-

degree of freedom chi2 distribution under the null). Note that this testing procedure is

conservative.

The Q value for the aggregate contribution of SGE was calculated for each phenotype

using the R package qvalue[49]. Significant contributions at FDR < 10% were those with

Q value < 0.1.

Correlation between DGE and SGE

The correlation ρ between $\underline{a_D}$ and $\underline{a_S}$ was calculated as:

$\rho = \dfrac{\sigma_{A_{DS}}}{\sigma_{A_D} \times \sigma_{A_S}}$

ρ reflects the correlation between SGE and DGE *acting on the same phenotype,*

similarly to how "traditional" genetic correlations measure the correlation between

DGE on two traits; ρ can actually be interpreted as the correlation between DGE on

the traits of cage mates mediating social effects and DGE on the phenotype of interest

itself.

We tested whether ρ was significantly different from 0 and whether |ρ| was

significantly different from 1 using a one-degree of freedom LLR test.

Simulations 1: for Supplementary_Figure1.

Phenotypes were simulated based on the genotypes and cage relationships of the full

set of 1,812 mice. Phenotypes were drawn from model (0) with the following variances:

$\sigma_{A_D}^2 = 15$, $\sigma_{A_S}^2 = 8$, $\rho_{A_{DS}} = 0.47$, $\sigma_{E_D}^2 = 22$, $\sigma_{E_S}^2 = 16$, $\rho_{E_{DS}} = -0.97$, $\sigma_C^2 = 26$. These variances

correspond to the median value of estimates across traits with aggregate SGE and

DGE > 5%. After building the phenotypic covariance matrix, the sample variance of

the simulations was calculated and used to calculate "realised" simulation parameters

from the "target" parameters above. The realised parameters were used for

comparison with the parameters estimated from the simulations.

Definition of "social genotype" for sgeGWAS

In the sgeGWAS, we assumed additive effects across cage mates and calculated the

"social genotype" of a mouse as the sum of the reference allele dosages of its cage

mates. The same assumptions were made by Biscarini *et al.*[13] and Brinker *et al.*[25].

Correlation between direct and social genotypes at a variant

Spearman's rank correlation coefficient was used. We tested whether the correlation

was different from 0 using the function cor.test in the R package stats[50].

Models used for sgeGWAS and dgeGWAS

To test SGE of a particular variant, we compared the following two models:

$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + G_f b_D$ (1, null)

$$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + G_f b_D + Z_f G b_S \qquad \text{(2, alternative)}$$

Here, G is the vector of direct genotypes at the tested variant, b_D the estimated

coefficient for local DGE and b_S the estimated coefficient for local SGE.

The models were fitted using LIMIX[51,52] with the covariance of the model estimated

only once per phenotype, in the model with no local genetic effect (model 0).

The significance of local SGE was calculated by comparing models (1) and (2) with a

1-degree of freedom LLR test.

We refer to the inclusion of $G_f b_D$ in model (1, null) as "conditioning".

dgeGWAS was carried out similarly, by comparing the null model (3) below and model

(2) above:

$$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + Z_f G b_S \qquad \text{(3, null)}$$

We refer to the inclusion of $Z_f G b_S$ in model (3, null) as "conditioning".

Identification of genome-wide significant associations

Because we wanted to compare the architecture of DGE and SGE *for each phenotype*

independently, we adopted the per-phenotype FDR approach used by Nicod et al.[26].

Had we used a study-wide FDR approach instead, the comparison of SGE and DGE

loci for a given phenotype would have depended on the SGE and DGE loci identified

for the other phenotypes in the dataset.

The procedure we used to control the FDR accounts for the fact that we report

loci rather than individual variants[53], where a locus is defined as the 1.5 Mb-wide

window around a SNP (this window size is the average 95% confidence interval for

DGE QTLs in [26]). More precisely, for each phenotype and for each type of genetic

effect (social and direct), we ran 100 "permuted GWAS" by permuting the rows of the

matrix of social (respectively direct) genotypes, and testing each variant at a time using

the permuted genotypes together with the un-permuted phenotypes, covariates, GRM

and matrix of direct (respectively social) genotypes (for conditioning). See [52,54] for

references on this permutation approach. For each permutation we then compiled a

list of loci that would be significant at a nominal P value of 0.01. Using the un-permuted

data, we similarly compiled a list of loci that would be significantly associated at a

nominal P value of 0.01. Ordering the latter in order of decreasing significance and

going down the list, we calculated for each locus an associated FDR until the FDR

was above 10%. For a given P value x, the FDR was calculated as:

$$FDR(x) = \frac{\# \ loci \ with \ P < x \ in \ permuted \ data}{100 \times \# \ loci \ with \ P < x \ in \ unpermuted \ data}$$

We report only those loci whose P value corresponds to an FDR < 10%.

Definition of candidate genes at associated loci (Table 2)

At each significantly associated locus we defined a 1.5Mb window centred on the lead

variant, identified all the variants that segregate in this window based on the full set of

7M variants, and reran the sgeGWAS locally with all the variants at the locus. We

highlighted those genes that are located within the most significantly associated

segments and whose MGI symbol does not start by 'Gm', 'Rik', 'Mir', 'Fam', or 'Tmem'

in order to enrich the reported sets in genes with known function.

Variance explained by a genome-wide significant association

The variance explained by a genome-wide significant SGE association was estimated

in an extension of model (0) with additional fixed effects for both direct and social

effects of lead SNPs at all genome-wide significant SGE loci (the lead SNP being the

SNP with the most significant P value at the locus in the sgeGWAS). After fitting the

model, the variance was calculated as:

$$\frac{var(ZGb_S)}{\sum var(X_c b_c) + \sum var(Gb_D) + \sum var(ZGb_S) + sampleVar(C)}$$

where $sampleVar(C)$ is the sample variance of the covariance matrix in this model.

The variance explained by a genome-wide significant DGE association was estimated

in a similar model but considering all genome-wide significant DGE associations and

calculated as:

$$\frac{var(Gb_D)}{\sum var(X_c b_c) + \sum var(Gb_D) + \sum var(ZGb_S) + sampleVar(C)}$$

Variance explained jointly by all genome-wide significant SGE or DGE associations

for a phenotype

The variance explained jointly by all significant SGE associations was estimated using the same model as above with all genome-wide significant SGE associations and calculated as:

$$\frac{\sum var(ZGb_S)}{\sum var(X_c b_c) + \sum var(Gb_D) + \sum var(ZGb_S) + sampleVar(C)}$$

The variance explained jointly by all significant DGE associations was estimated using the same model as above with all genome-wide significant DGE associations and calculated as:

$$\frac{\sum var(Gb_D)}{\sum var(X_c b_c) + \sum var(Gb_D) + \sum var(ZGb_S) + sampleVar(C)}$$

Simulations 2: for Supplementary Figure 2d.

Phenotypes were simulated based on the genotypes and cage relationships of the full set of 1,812 mice. Phenotypes were simulated as the sum of random effects and local DGE (from model (1)), with the following parameters: $\sigma^2_{A_D}$ = 5 or 20, $\sigma^2_{A_S}$ = 5 or 20, $\rho_{A_{DS}}$= 0.5, $\sigma^2_{E_D}$= 30, $\sigma^2_{E_S}$= 30, $\rho_{E_{DS}}$ = -0.97, σ^2_C = 25. The values for $\rho_{A_{DS}}$, $\sigma^2_{E_D}$, $\sigma^2_{E_S}$, $\rho_{E_{DS}}$, and σ^2_C were close to the median of the corresponding estimates from the real data. $\sigma^2_{A_D}$ = 5 and $\sigma^2_{A_S}$ = 5 correspond to low polygenic effects in the real data, and $\sigma^2_{A_D}$ = 20 and $\sigma^2_{A_S}$ = 20 correspond to high polygenic effects in the real data. We simulated local DGE at random variants in the genome, and simulated variances of 0, 5, 20 or 50.

 The results we show in Supplementary Figure 2d are based on a subset of simulations: $\sigma^2_{A_D}$ = 20 and $\sigma^2_{A_S}$ = 20 and local DGE variance of 20.

Simulations 3: for Supplementary Figure 3a-d, and Figure 3a.

Phenotypes were simulated based on the real genotypes but random cages for a
random subset of 1,800 mice (in order to be able to draw full cages of 2 or 3 mice).
Phenotypes were simulated as the sum of random effects, local DGE and local SGE
(model (2) except for Z) with the following parameters: $\sigma_{A_D}^2 = 17$, $\sigma_{A_S}^2 = 17$, $\rho_{A_{DS}} = 0.65$,
$\sigma_{E_D}^2 = 19$, $\sigma_{E_S}^2 = 15$, $\rho_{E_{DS}} = -0.8$, $\sigma_C^2 = 25$. Those values correspond to the median
estimates for phenotypes with aggregate SGE and DGE > 0.1.

We simulated local SGE and DGE at variants where direct and social
genotypes were either lowly correlated (Spearman correlation negative log P value <
0.05) or more highly correlated (Spearman correlation negative log P value > 0.2), and
had with low MAF (MAF < 0.05), medium MAF (0.225<MAF<0.275) or high MAF
(MAF>0.45). We simulated local DGE with an allelic effect of 0 or 1 (1 corresponds to
a large effect in the real data). We simulated local SGE under two alternative
generative models: an "additive" model by using Z as in model (2) (i.e. filled with 0s
and 1s) or a "proportional" model by using $Z' = \frac{Z}{N}$. In all cases we simulated an
allelic effect of 0.2 (similar to the average allelic effect estimated in the SGE GWAS).
The sample variance of the simulated local DGE term is $var(Gb_D) = 2p(1-p)\,b_D^2$; it
is $var(ZGb_S) = 2Np(1-p)\,b_S^2$ for the local SGE term simulated under the additive
model, and $var\left(\frac{Z}{N}Gb_S\right) = 2Np(1-p)/N^2\,b_S^2$ for the local SGE component simulated
under the proportional model.

The results we show in Supplementary Figure 3a-d are based on a subset of
simulations with group size 3 and are averaged across low, medium and high MAF.
Power was calculated at a genome-wide significance threshold of negative log P 5,
which is similar to the significance of associations detected at FDR < 10%.
The results we show in Figure 3a are based on a subset of simulations with
group size 2 and 3, no local DGE, and averaged across high and low genotypic
correlations. Power was also calculated at a genome-wide significance threshold of
negative log P 5.

Scripts used in this study

All the scripts used in this study are available from http://github.com/limix/SGE.
LIMIX can be downloaded from http://github.com/limix/limix.

Acknowledgements

665 We would like to thank Na Cai and Robert W. Davis for helpful discussion of the data

666 from Nicod et al.[26] and Davies et al.[27], as well as Ewan Birney, Emma Huang, Richard

667 Mott, Abraham A. Palmer, and Sandra Sanchez-Roige for comments on a previous

668 version of this manuscript. AB was supported by a fellowship from the Wellcome Trust

669 (105941/Z/14/Z).

References

1. Griffing, B. Selection in reference to biological groups I. Individual and group selection applied to populations of unordered groups. *Australian Journal of Biological Sciences* **20**, 127–140 (1967).

2. Moore, A.J., Brodie III, E.D. & Wolf, J.B. Interacting phenotypes and the evolutionary process: I. Direct and indirect genetic effects of social interactions. *Evolution* **51**, 1352–1362 (1997).

3. Wolf, J.B., Brodie III, E.D., Cheverud, J.M., Moore, A.J. & Wade, M.J. Evolutionary consequences of indirect genetic effects. *Trends in ecology & evolution* **13**, 64–69 (1998).

4. Bergsma, R., Kanis, E., Knol, E.F. & Bijma, P. The contribution of social effects to heritable variation in finishing traits of domestic pigs (Sus scrofa). *Genetics* **178**, 1559–1570 (2008).

5. Bijma, P. The quantitative genetics of indirect genetic effects: a selective review of modelling issues. *Heredity* **112**, 61 (2014).

6. Wilson, A.J., Gelin, U., Perron, M.–C. & Réale, D. Indirect genetic effects and the evolution of aggression in a vertebrate system. *Proceedings of the Royal Society of London B: Biological Sciences* **276**, 533–541 (2009).

7. Ashbrook, D.G., Gini, B. & Hager, R. Genetic variation in offspring indirectly influences the quality of maternal behaviour in mice. *Elife* **4**, 11814 (2015).

8. Ashbrook, D.G., Sharmin, N. & Hager, R. Offspring genes indirectly influence sibling and maternal behavioural strategies over resource share. in *Proc. R. Soc. B* Vol. 284 20171059 (The Royal Society, 2017).

9. Baud, A. *et al.* Genetic variation in the social environment contributes to health and disease. *PLoS genetics* **13**, e1006498 (2017).

10. Domingue, B.W. *et al.* The social genome of friends and schoolmates in the National Longitudinal Study of Adolescent to Adult Health. *Proceedings of the National Academy of Sciences*, 201711803 (2018).

11. Kong, A. *et al.* The nature of nurture: Effects of parental genotypes. *Science* **359**, 424–428 (2018).

12. Bates, T.C. *et al.* The Nature of Nurture: Using a Virtual–Parent Design to Test Parenting Effects on Childrens Educational Attainment in Genotyped Families. *Twin Research and Human Genetics*, 1–11 (2018).

13. Biscarini, F. *et al.* Across–line SNP association study for direct and associative effects on feather damage in laying hens. *Behavior genetics* **40**, 715–727 (2010).

14. Visscher, P.M., Brown, M.A., McCarthy, M.I. & Yang, J. Five years of GWAS discovery. *The American Journal of Human Genetics* **90**, 7–24 (2012).

15. Ferrero, D.M. *et al.* A juvenile mouse pheromone inhibits sexual behaviour through
the vomeronasal system. *Nature* **502**, 368 (2013).

16. Ross, K.G. & Keller, L. Genetic control of social organization in an ant. *Proceedings of*
the National Academy of Sciences **95**, 14232–14237 (1998).

17. Mutic, J.J. & Wolf, J.B. Indirect genetic effects from ecological interactions in
Arabidopsis thaliana. *Molecular ecology* **16**, 2371–2381 (2007).

18. Bailey, N.W. & Hoskins, J.L. Detecting cryptic indirect genetic effects. *Evolution* **68**,
1871–1882 (2014).

19. Wolf, J.B. Genetic architecture and evolutionary constraint when the environment
contains genes. *Proceedings of the National Academy of Sciences* **100**, 4655–4660
(2003).

20. Muir, W.M. Incorporation of Competitive Effects in Plant or Animal Breeding
Programs. *Genetics* (2005).

21. Bijma, P., Muir, W.M., Ellen, E.D., Wolf, J.B. & Van Arendonk, J.A. Multilevel selection
2: estimating the genetic parameters determining inheritance and response to
selection. *Genetics* **175**, 289–299 (2007).

22. Bijma, P., Muir, W.M. & Van Arendonk, J.A. Multilevel selection 1: quantitative
genetics of inheritance and response to selection. *Genetics* **175**, 277–288 (2007).

23. McGlothlin, J.W., Moore, A.J., Wolf, J.B. & Brodie III, E.D. Interacting phenotypes and
the evolutionary process. III. Social evolution. *Evolution* **64**, 2558–2574 (2010).

24. Bijma, P. & Wade, M.J. The joint effects of kin, multilevel selection and indirect genetic
effects on response to genetic selection. *Journal of evolutionary biology* **21**, 1175–
1188 (2008).

25. Brinker, T., Bijma, P., Vereijken, A. & Ellen, E.D. The genetic architecture of socially–
affected traits: a GWAS for direct and indirect genetic effects on survival time in laying
hens showing cannibalism. *Genetics Selection Evolution* **50**, 38 (2018).

26. Nicod, J. *et al.* Genome–wide association of multiple complex traits in outbred mice by
ultra–low–coverage sequencing. *Nature genetics* **48**, 912 (2016).

27. Davies, R.W., Flint, J., Myers, S. & Mott, R. Rapid genotype imputation from sequence
without reference panels. *Nature genetics* **48**, 965 (2016).

28. Haeffel, G.J. & Hames, J.L. Cognitive Vulnerability to Depression Can Be Contagious.
Clinical Psychological Science **2**, 75–85 (2014).

29. Rosenquist, J.N., Murabito, J., Fowler, J.H. & Christakis, N.A. The spread of alcohol
consumption behavior in a large social network. *Annals of internal medicine* **152**, 426–
433 (2010).

30. Grant, J.D. *et al.* Spousal concordance for alcohol dependence: evidence for
assortative mating or spousal interaction effects? *Alcoholism: Clinical and*
Experimental Research **31**, 717–728 (2007).

31.   Bolger, N., DeLongis, A., Kessler, R.C. & Wethington, E. The contagion of stress across
749   multiple roles. *Journal of Marriage and the Family*, 175–183 (1989).
750   32.   Christakis, N.A. & Fowler, J.H. The spread of obesity in a large social network over 32
751   years. *New England journal of medicine* **357**, 370–379 (2007).
752   33.   Cohen–Cole, E. & Fletcher, J.M. Detecting implausible social network effects in acne,
753   height, and headaches: longitudinal analysis. *Bmj* **337**, a2533 (2008).
754   34.   Yalcin, B. *et al.* Commercially available outbred mice for genome–wide association
755   studies. *PLoS genetics* **6**, e1001085 (2010).
756   35.   Smyth, I. *et al.* A mouse model of harlequin ichthyosis delineates a key role for Abca12
757   in lipid homeostasis. *PLoS genetics* **4**, e1000192 (2008).
758   36.   Yoshida, S. *et al.* Involvement of an NKG2D ligand H60c in epidermal dendritic T cell–
759   mediated wound repair. *The Journal of Immunology* **188**, 3972–3979 (2012).
760   37.   Li, Y. *et al.* Differential expression of hippocampal EphA4 and ephrinA3 in anhedonic–
761   like behavior, stress resilience, and antidepressant drug treatment after chronic
762   unpredicted mild stress. *Neuroscience letters* **566**, 292–297 (2014).
763   38.   Visscher, P.M. *et al.* 10 years of GWAS discovery: biology, function, and translation.
764   *The American Journal of Human Genetics* **101**, 5–22 (2017).
765   39.   Baud, A. & Flint, J. Identifying genes for neurobehavioural traits in rodents: progress
766   and pitfalls. *Disease models & mechanisms* **10**, 373–383 (2017).
767   40.   Manolio, T.A. *et al.* Finding the missing heritability of complex diseases. *Nature* **461**,
768   747 (2009).
769   41.   Valdar, W. *et al.* Genome–wide genetic association of complex traits in heterogeneous
770   stock mice. *Nature genetics* **38**, 879 (2006).
771   42.   Sequencing, R.G. *et al.* Combined sequence–based and genetic mapping analysis of
772   complex traits in outbred rats. *Nature genetics* **45**, 767 (2013).
773   43.   Bastiampillai, T., Allison, S. & Chan, S. Is depression contagious? The importance of
774   social networks and the implications of contagion theory. *Australian & New Zealand*
775   *Journal of Psychiatry* **47**, 299–303 (2013).
776   44.   Domingue, B.W., Fletcher, J., Conley, D. & Boardman, J.D. Genetic and educational
777   assortative mating among US adults. *Proceedings of the National Academy of Sciences*
778   **111**, 7996–8000 (2014).
779   45.   Robinson, M.R. *et al.* Genetic evidence of assortative mating in humans. *Nature*
780   *Human Behaviour* **1**, 0016 (2017).
781   46.   Venerables, W. & Ripley, B. Modern applied statistics with S. (new york: Springer,
782   2002).
783   47.   Kang, H.M. *et al.* Variance component model to account for sample structure in
784   genome–wide association studies. *Nature genetics* **42**, 348–354 (2010).
785   48.   Searle, S.R. *Matrix algebra useful for statistics*, (Wiley, 1982).

786   49.   Dabney, A., Storey, J.D. & Warnes, G. qvalue: Q-value estimation for false discovery
787   rate control. *R package version* **1** (2010).

788   50.   Best, D. & Roberts, D. Algorithm AS 89: the upper tail probabilities of Spearmans rho.
789   *Journal of the Royal Statistical Society. Series C (Applied Statistics)* **24**, 377–379
790   (1975).

791   51.   Lippert, C., Casale, F., Rakitsch, B. & Stegle, O. LIMIX: genetic analysis of multiple traits.
792   (2014).

793   52.   Casale, F.P., Rakitsch, B., Lippert, C. & Stegle, O. Efficient set tests for the genetic
794   analysis of correlated traits. *Nature methods* **12**, 755 (2015).

795   53.   Brzyski, D. *et al.* Controlling the rate of GWAS false discoveries. *Genetics* **205**, 61–75
796   (2017).

797   54.   Sudmant, P.H. *et al.* An integrated map of structural variation in 2,504 human
798   genomes. *Nature* **526**, 75 (2015).