Dissecting the mechanisms underlying indirect genetic effects on biomedical phenotypes: a study of 170 behavioural, physiological and morphological phenotypes measured in adult laboratory mice.

Short title:

Mechanisms underlying indirect genetic effects on biomedical phenotypes

Amelie Baud[1*], Francesco Paolo Casale[1,2], Jerome Nicod[3], Oliver Stegle[1,4*]

[1] European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, CB10 1SD Hinxton, Cambridge, UK

[2] Microsoft Research New England, Cambridge, Massachusetts, USA

[3] Wellcome Trust Centre for Human Genetics, Roosevelt Drive, Oxford OX3 7BN, UK

[4] European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, DE

* Corresponding authors: abaud@ebi.ac.uk and oliver.stegle@embl.de

Abstract

The phenotype of an individual can be affected not only by the individual's own genotypes (direct genetic effects, DGE) but also by genotypes of other, interacting individuals (indirect genetic effects, IGE). IGE have been detected, using polygenic models, for a broad range of biomedical phenotypes. However, little is known about the loci, genes and traits of interacting partners mediating the effects, especially where non-familial IGE are concerned. To address this question, we studied IGE arising in cages of unrelated, adult laboratory mice. We leveraged a dataset of 170 behavioural, physiological and morphological phenotypes measured in 1,812 genetically heterogeneous mice and developed two approaches. First, we used variance components models to estimate, for each phenotype, the correlation between DGE and IGE. Our results demonstrate some overlap but also differences between the mechanisms of DGE and IGE for a given phenotype. Second, we developed and applied methods for the genome-wide association study of IGE (igeGWAS) in order to identify loci, genes and traits of cage mates mediating IGE. We identified 24 genome-wide significant IGE loci for 17 phenotypes (FDR < 10%), none of which overlapped with genome-wide significant DGE loci for the same phenotype. Using an exhaustive list of single nucleotide polymorphisms at each locus, we fine-mapped each association. At six loci this pointed to a single candidate gene, which we used to formulate specific hypotheses as to the mechanisms of IGE. The empirical insights from our study and the analytical strategies we have developed pave the way for using igeGWAS to unravel mechanisms of phenotypic variation that are expressed only in the context of social interactions.

Introduction

The phenotype of an individual can be affected not only by the individual's own genotypes (direct genetic effects, DGE) but also by genotypes of other, interacting individuals (indirect genetic effects, IGE)(1-3). IGE arise when the phenotype of the "focal" individual is influenced by heritable traits of "partners", which may include behavioural and non-behavioural traits as well as modifications of the non-social environment by partners (4) (**Figure 1a**). IGE have been detected in many experimental systems(5-15), livestock(16-18), crops(19), wild animals(20-22), and humans(23-25), demonstrating that they are an important component of the genotype to phenotype path.

Most studies of IGE have used polygenic models to study aggregate effects: trait-based models(2, 26) and polygenic risk scores(24, 27) have been used to study IGE mediated by *specific* traits of partners, whereas variance components models have been used to detect IGE mediated by *any* heritable trait of partners (10, 16, 28, 29). More recently, the genome-wide association study of IGE (igeGWAS) has been proposed as a strategy to identify individual genetic loci causing IGE associations(5, 8, 9, 25, 30, 31). The major advantage of this strategy is the ability to leverage associated loci to uncover novel traits of partners influencing a phenotype of interest.

However, so far igeGWAS has been applied only in limited settings. In particular, it has not been used to study non-familial IGE affecting biomedical phenotypes (IGE from "peers"), despite evidence from polygenic models suggesting such IGE are common in laboratory mice(10) and occur in humans too(27).

In that context, it is unknown whether igeGWAS of a phenotype will identify loci other than those identified by dgeGWAS of the same phenotype. This will depend,

first, on the correlation between DGE and IGE for that phenotype (**Figure 1b**). This correlation is expected to be different from zero based on reports of "phenotypic contagion" between mouse and human peers, namely instances where an intermediate phenotype such as stress levels(32-34), infection status, or a component of the gut microbiome(35) is "transferred" to peers and contributes to the phenotype of interest both in focal individuals and their peers. In the extreme, if the correlation between DGE and IGE is close to ±1, there will be little scope for igeGWAS to identify loci not identified by dgeGWAS. Second, whether igeGWAS identifies loci not identified by dgeGWAS will depend on the relative power of igeGWAS and dgeGWAS, which has not been thoroughly investigated.

Finally, when igeGWAS is applied to peer effects, each individual in the study gives rise to DGE and IGE (**Figure 1b**), hence serves as both focal individual and peer in the analysis. Whether this leads to confounding between DGE and IGE is unclear.

To address these questions, we leveraged a published dataset of 170 behavioural, physiological and morphological phenotypes measured in 1,812 genetically heterogeneous laboratory mice (**Figure 1c**), which we supplemented with previously unreported cage information (**Supplementary Table 1**), and used two approaches: first, we used polygenic models to estimate, for each phenotype, the correlation between DGE and IGE (**blue arrows in Figure 1b**; see Methods). This correlation is equivalent to the correlation between DGE on the traits of cage mates influencing the phenotype of interest and DGE on the phenotype of interest itself (**red arrows in Figure 1b**) and can be estimated even when the traits of cage mates mediating IGE are unknown or unmeasured. Second, we developed methods for igeGWAS and applied igeGWAS and dgeGWAS to the 170 phenotypes to compare

the relative power of dgeGWAS and igeGWAS and identify novel genes and traits of

cage mates mediating IGE.

Figure 1 *Illustration of indirect genetic effects (IGE) and experimental design. **(a)** IGE*

arise when two individuals interact and heritable traits of one influence the phenotype

of the other. Importantly, the traits of cage mates that mediate social effects do not

need to be known or measured for many SGE analyses. This figure defines several

terms used throughout this manuscript. We further refer to "peers" as non-familial

"partners". (b) DGE and IGE when all study individuals are genetically heterogeneous.

The correlation ρ between DGE and IGE for a given phenotype is most directly

understood by looking at the blue arrows (mice "projecting" IGE). It is, however,

equivalent to the correlation between DGE on phenotype of interest and DGE on traits

of partners mediating IGE (red arrows). (c) Experimental design. A list of the 170

phenotypes collected on each mouse is presented in Supplementary Table 2.

Results

We used the genome-wide genotypes (both LD-pruned and non-pruned, see Methods) and 200 phenotypes for 2,073 CFW mice reported in Nicod et al.(36) and Davies et al.(37). In addition, we used previously unreported cage information provided by the authors of the original study upon request (**Supplementary Table 1**). Mice were housed in same-sex groups and interacted for at least nine weeks before phenotyping started. For our analyses we only kept mice that had the same two cage mates over the course of the experiment. We also excluded a subset of mice suspected to be siblings in order to mitigate confounding from parental and litter effects, yielding a final sample size of 1,812 mice. We normalised each phenotype and excluded 30 phenotypes that could not be satisfactorily normalised, yielding a total of 170 phenotypes. Due to missing phenotype data, the average sample size for each phenotype was 1,256.

Correlation between DGE and IGE for a given phenotype

Initially, to assess the role of phenotypic contagion in mediating IGE and quantify the scope for igeGWAS to identify loci not identified by dgeGWAS, we used polygenic models to estimate the correlation ρ between DGE and IGE (**blue arrows in Figure 1b**). We limited our analysis to 28 traits with marginal DGE and IGE that exceeded 5%, for which ρ can be accurately estimated (**Supplementary Figure 1**), including behavioural, physiological and morphological phenotypes (**Figure 2 and Supplementary Table 2**). A correlation coefficient of 0 for a phenotype would indicate different intermediate traits (gene expression traits, pathways) mediate DGE and IGE on the phenotype of interest and no role of phenotypic contagion in mediating IGE. In

contrast, a correlation coefficient of ±1 would indicate that DGE and IGE are mediated

by the exact same intermediate traits, implying that IGE are mediated by the

phenotype of interest itself and that no other trait of cage mates is involved. We tested

both models, finding, first, that ρ was significantly different from zero for 10 out of 28

phenotypes (P < 0.05), which is consistent with a role for phenotypic contagion in

mediating IGE. Second, ρ was significantly different from plus or minus one for the

same number of phenotypes (P < 0.05), indicating that phenotypic contagion is usually

not sufficient to explain IGE and that traits of cage mates other than the phenotype of

interest mediate IGE. To uncover those traits, we turned to igeGWAS.

Figure 2 *Correlation coefficients ρ between DGE and IGE. ρ is shown for 28*

phenotypes with marginal DGE and IGE greater than 5%. Error bars denote standard

errors. Asterisks on the left show phenotypes for which ρ is significantly different from

0 (black: P < 0.05, red: Bonferroni-corrected P < 0.05). Asterisks on the right show

phenotypes for which $|\rho|$ is significantly different from 1 (black: P < 0.05, red:

Bonferroni-corrected P < 0.05). Numerical values are provided in Supplementary

Table 2.

igeGWAS and dgeGWAS of 170 phenotypes

To assess the relative power of igeGWAS and dgeGWAS and to gain insights into the mechanisms of IGE, we performed igeGWAS and dgeGWAS in an analogous manner for each one of the 170 phenotypes.

For igeGWAS, we estimated the "social genotype" of a mouse at a variant as the sum of the reference allele dosages across its cage mates at the variant[31, 38], and tested for association between this social genotype and the phenotype of interest. In order to avoid spurious associations, we accounted for background IGE, background DGE and shared environmental (cage) effects using random effect components in a linear mixed model. Using simulation studies, we identified a previously unrecognised source of spurious associations in igeGWAS: correlations between direct and social genotypes that arise when each individual serves as both focal individual and social partner in the analysis, a strategy that maximises sample size when all the individuals are genotyped and phenotyped[31, 38]. Importantly, we show that these genotypic correlations arise even when all individuals are strictly unrelated (**Supplementary Note**). To account for these correlations in our study, we included, in the null model for igeGWAS, a fixed effect covariate for direct genotypes, a strategy we hereafter refer to as "conditioning". Consistent with the results of a simulation study, we observed that conditioning is required to obtain calibrated P values (**Supplementary Figure 2, 4 and 5b**). We also note that conditioning will result in decreased power to detect IGE in the absence of co-localised DGE (**Supplementary Figure 3a and 3c**) but increased power if the locus also gives rise to DGE (**Supplementary Figures 3b and 3d**). Given the intermediate values observed for the correlation ρ between DGE and IGE across phenotypes (**Figure 2**), we expect conditioning to overall increase power to detect IGE in our dataset. Finally,

following the approach considered in Nicod et al.(36), we employed a genome-wide

permutation strategy to control the FDR for each phenotype, thereby taking into

account the specific patterns of linkage disequilibrium present in the sample.

igeGWAS identified a total of 24 genome-wide significant loci across 17 of the 170

tested phenotypes (FDR < 10%; **Supplementary Table 3**).

To enable a direct comparison between igeGWAS and dgeGWAS, we ran

dgeGWAS for each phenotype analogously to the approach taken for igeGWAS,

including random effects for DGE and IGE polygenic effects and cage effects and

conditioning on the social genotype at each locus. This identified 120 genome-wide

significant DGE loci for 63 phenotypes (10% per-phenotype FDR; **Supplementary**

Table 4).

Relative power of igeGWAS and dgeGWAS

igeGWAS and dgeGWAS were run in analogous manners yet, across all 170

phenotypes, igeGWAS identified fewer genome-wide significant loci than dgeGWAS

(24 and 63 loci respectively). Consistent with this observation, genome-wide

significant IGE loci tended to have lower effect sizes (proportion of phenotypic

variance explained) than DGE loci (**Figure 3a**). This trend is likely to be even stronger

across all associated loci due to the Beavis effect(39, 40).

Was lower power to detect IGE to be expected? This question has been

addressed in the context of polygenic effects(41, 42) but not in the context of GWAS.

To do so, we carried out a power analysis where allelic effects were fixed, three

categories of minor allele frequencies (MAF) were considered (low, medium, or high),

and local DGE or IGE were simulated. We used two generative models for IGE: an

additive model, whereby IGE received by a focal mouse correspond to the sum of the

genetic effects projected by its cage mates, and an average model, whereby IGE

received by one mouse correspond to the average of the genetic effects projected by

its cage mates (see **Figure 1b**). The average model corresponds to a scenario where

a mouse can only interact with one cage mate at a time, resulting in a dilution of social

effects when more than one cage mate is present in the cage (two cage mates were

present in this study). Our results show that power to detect IGE loci always increases

with MAF and is higher than power to detect DGE if IGE arise from the sum of genetic

effects across cage mates (additive model)(**Figure 3b**). With dilution however

(average model), power to detect IGE becomes lower than power to detect DGE.

a and **b** labels as shown in figure.

Figure 3 *Effect sizes of genome-wide significant DGE and IGE associations and*

relationship to power of igeGWAS and dgeGWAS. *(a) Histogram of the proportion of*

phenotypic variance explained by genome-wide significant IGE (red) and DGE (black)

associations (FDR < 10%). *(b) Power to detect IGE and DGE associations in*

simulations. Three types of local genetic effects were simulated: DGE, IGE arising

from the sum of genetic effects across cage mates (additive model) and IGE arising

from the average of genetic effects across cage mates (average model). See main

text for an explanation of the biology behind each model. For each type of effect,

results are shown (left to right) for variants with low MAF (MAF < 0.05), medium MAF

(0.225<MAF<0.275) and high MAF (MAF>0.45) (MAF: minor allele frequency, defined

based on direct genotypes). Power was calculated at a genome-wide significance

threshold of negative log P 5.

Given the relatively lower power of igeGWAS, reducing the multiple testing burden of

igeGWAS by testing, for a given phenotype, only those loci that are genome-wide

significant DGE loci for the same or related phenotype (e.g. DGE loci for any

behavioural phenotype, or any immunological phenotype) may seem like a promising

approach(12). However, we show that doing so would miss the most significant IGE

loci as genome-wide significant DGE and IGE loci for a given category of phenotypes

(e.g. behavioural or immunological) do not overlap (**Figure 4**). This observation is

expected based on the intermediate values observed for the correlation ρ between

DGE and IGE and the limited power of dgeGWAS and igeGWAS.

Figure 4 *Superimposed manhattan plots corresponding to igeGWAS (top panel) and dgeGWAS (bottom panel) of the same 170 phenotypes. DGE associations with a negative log P value greater than 10 were truncated at this threshold (as indicated by 10+); also, data points with negative log P values smaller than 2 are not shown. The larger dots correspond to the most significant variant at each genome-wide significant IGE or DGE locus. In the IGE panel (top), each colour corresponds to a class of phenotypes: behavioural (red, includes 7 behavioural phenotypes with a detected IGE locus), adult neurogenesis (black, 2 phenotypes), immune (orange, 1 phenotype), haematological (yellow, 1 phenotype), blood biochemistry (blue, 2 phenotypes), bone phenotypes (green, 2 phenotypes), heart function (brown, 1 phenotype), and lung function (purple, 1 phenotype). In the DGE panel (bottom), the same colouring scheme*

is used as in the IGE panel except for grey dots, which are for phenotype that do not

have a genome-wide significant IGE locus.

Mechanisms of IGE inferred from genome-wide significant IGE loci

Linkage disequilibrium decays faster in the CFW population than in many other mouse

populations used for mapping, which facilitates identification of putatively causal

genes at associated loci(36, 43, 44). To identify such genes, we fine-mapped genome-

wide significant IGE loci using the full set of variants (rather than the pruned set used

for igeGWAS) in the 1.5Mb window surrounding the most significant variant at the

locus, which corresponds, in this sample, to the 95% confidence interval for the

association(36). We then identified, for each genome-wide significant IGE locus,

putatively causal genes (hereafter "candidate genes") that either overlapped the

associated plateau or were located in direct proximity (**Supplementary Table 3**).

Following Nicod et al.(36), we focused on IGE loci with a single candidate gene to

draw hypotheses on the mechanisms of IGE.

Epha4 was the only candidate gene at a locus on chromosome 1 for immobility

in the forced swim test (FST), a measure of stress coping strategy (**Figure 5a**, top

panel, and **Supplementary Figure 4**). Because direct effects of *Epha4* on FST

immobility have been reported(45, 46), we evaluated the possibility that behavioural

contagion in the weeks prior to or during the FST might mediate IGE of *Epha4* on FST

immobility in CFW mice. To do so, we ran a direct association (DGE) analysis using

the full set of variants in the same 1.5Mb window including *Epha4* as for IGE analysis.

We found no evidence that *Epha4* directly affects FST immobility in CFW mice (**Figure**

5a, bottom panel), making it unlikely that behavioural contagion explains the detected

IGE.

To elucidate the mechanisms of action of *Epha4*, we used independent

expression data from the hippocampus of a separate sample of 79 mice from the same

CFW population(44). We did not find a cis-eQTL for *Epha4* but identified *Dlgap1*, a

candidate gene at another IGE locus for FST immobility (**Supplementary Figure 5**),

as the sixth gene most significantly co-expressed with *Epha4* in the hippocampus of

CFW mice (**Supplementary Figure 6**). The correlation in expression was significant

after accounting for the number of genes expressed in the hippocampus (Spearman r

= 0.868, Bonferroni-corrected P = $2,3.10^{-19}$). *Epha4* and *Dlgap1* encode two synaptic

proteins that are important for synaptic plasticity in the hippocampus(47-49) and the

corresponding knock-out mice show altered social approach(50) and social

memory(51). Hence, our results suggest that differences in the social behaviour of

cage mates is responsible for IGE on stress-coping strategy in CFW mice.

Unfortunately, we were not able to replicate our analyses or directly test our hypothesis

as FST measurements were not available for the additional sample of 79 CFW mice,

expression data were not available for the main sample, and social behaviour was not

tested in either sample.

Among the other genome-wide significant IGE loci with a single candidate gene

was a locus on chromosome 12 for a measure reflecting unstable sleep patterns over

the three days of the sleep test (**Figure 5c**, top panel). Importantly, the mice were

singly housed during the sleep test, implying that any influence of cage mates must

have either persisted or revealed itself in the new environment that the mice

encountered. We found no evidence of DGE on the same measure at this locus

(**Figure 5c**, bottom panel), suggesting unstable sleep patterns during the sleep test

were not caused by similarly unstable sleep patterns of cage mates prior to the sleep

test.

The sole candidate gene at this locus, *Pgk1-rs7*, is a pseudogene of *Pgk1*,

which encodes the glycolytic enzyme phosphoglycerate kinase 1. Because *Pgk1* and

Pgk1-rs7 have similar sequences and that could have led to genotyping errors in the

form of similar genotypes at *Pgk1 and Pgk1-rs7*, we first confirmed that variants in the

vicinity of *Pgk1-rs7* and variants in the vicinity of *Pgk1* were not linked, suggestsing

Pgk1-rs7, but not *Pgk1*, is involved. Whether *Pgk1-rs7* is functional or not has not

been reported before. However, the perfect sequence match between the proteins

encoded by *Pgk1-rs7* and *Pgk1,* evolutionary theory related to the position of *Pgk1* on

the X chromosome(52), and the existence of a functional pseudogene for another

glycolytic enzyme(53) suggest that *Pgk1-rs7* is likely to be functional.

A previous report found that *Pgk1-rs7* was expressed specifically in brown

adipose tissue - the tissue responsible for non-shivering thermogenesis – and that its

expression was activated by cold temperatures(54). Therefore, our hypothesis is that,

when mice were group-housed and slept huddled before the sleep test, heat produced

by cage mates raised the skin temperature of focal mice(55), which is permissive for

sleep(56-58); when mice became singly housed for the sleep test and as laboratory

conditions are well below thermoneutrality for mice(59), focal mice that used to have

warmer cage mates struggled more than others to adapt to the new, cold conditions,

resulting in unstable sleep patterns during the sleep test.

Figure 5 *Locus zoom plots for two genome-wide significant IGE loci (FDR < 10%).*

The top and bottom panels show, respectively, IGE and DGE associations for <u>all</u> the

variants at the locus (rather that the set of pruned variants used for GWAS). (a) Locus

on chromosome 1 for immobility during the first two minutes of the forced swim test

(FST), a measure of stress-coping strategy in mice. Epha4 is the only candidate gene

(see Methods) for IGE at this locus. (b) Locus on chromosome 12 for a measure

reflecting unstable sleep patterns during the three days of the sleep test. Pgk1-rs7 is

the only candidate gene for IGE at this locus.

Discussion

To dissect the mechanisms underlying non-familial IGE on behavioural, physiological and morphological phenotypes of adult laboratory mice(10), we applied polygenic models and igeGWAS to 170 such phenotypes.

Using polygenic models, first, we gained insights into the role of phenotypic contagion in mediating IGE and the scope for igeGWAS to identify loci not already identified by dgeGWAS. To do so, we estimated, for each phenotype, the correlation ρ between DGE and IGE, which is equivalent to the correlation between DGE on the phenotype of interest and DGE on the (unknown, unmeasured) traits of cage mates influencing the phenotype of interest. We tested two extreme scenarios: one in which phenotypic contagion plays no role in mediating IGE (ρ=0) and one in which phenotypic contagion is sufficient to explain IGE (ρ=1). Focusing on 28 phenotypes strongly affected by DGE and IGE, we first found that ρ was significantly different from 0 for 10 out of 28 phenotypes (P < 0.05). This result is consistent with a role for phenotypic contagion in mediating IGE, which is expected from previous reports of phenotypic contagion following non-genetic manipulations of the phenotype of laboratory mice(32, 35). We also found that ρ was significantly different from |1| for 10 out of 28 phenotypes (P < 0.05), which demonstrates that phenotypic contagion is not sufficient to account for IGE and that traits of cage mates other than the phenotype of interest are involved.

Using igeGWAS, second, we identified 24 genome-wide significant IGE loci for 17 of the 170 phenotypes considered, none of which were genome-wide significant in the corresponding dgeGWAS. We highlighted three of those loci to illustrate how IGE loci may shed light into the mechanisms of IGE. Briefly, we fine-mapped each locus,

identified candidate genes based on their proximity to the most significantly associated

variants and focused, to begin with, on loci with a single candidate gene. We reviewed

the literature on each single candidate gene to identify traits of cage mates the gene

might contribute to and that could plausibly influence the phenotype of focal mice.

Doing so, we hypothesized a role for *Epha4* and *Dlgap1* in giving rise to IGE on

immobility in the FST, a measure of stress-coping strategy(60). The selective

improvements in social memory observed in *Epha4* knock-out mice(50) and the

selective reduction in social approach observed in *Dlgap1* knock-out mice(51) together

led us to hypothesize that variation in the social behaviour of cage mates mediates

IGE of *Epha4* and *Dlgap1* on stress-coping strategy. Exactly what kind of social

behaviour of cage mates is involved cannot be predicted from the knock-out mice, as

those mice were confronted to an immobilized, stranger mouse outside their home

cage. Similarly, evidence that both aggression from a stranger mouse and social

isolation affect FST immobility(61-65) supports our hypothesis but it does not

necessarily mean aggression or affiliative behaviour cause IGE in CFW mice.

Experimental work will permit testing the hypothesis that *Epha4* and *Dlgap1* can give

rise to IGE on stress coping strategy and identify the specific social behaviour(s)

involved. As a second example we highlighted an IGE locus for unstable sleep

patterns where *Pgk1-rs7* was the single candidate gene. Based on evidence

suggesting *Pgk1-rs7* is involved in non-shivering thermogenesis(54), we proposed

that thermogenesis in cage mates influences the sleep of focal mice by keeping the

skin of the latter warm as all mice sleep huddled.

Understanding the mechanisms underlying IGE on biomedical phenotypes in

laboratory mice serves two goals: first, it may uncover mechanisms of social influence

that are at play in humans too. For example, identifying specific aspects of the social

behaviour of cage mates that influence stress-coping strategy in CFW mice may

provide insights into specific traits of peers influencing stress-coping strategies in

humans. Second, improving our understanding of how mice in a cage influence each

other will permit measuring and accounting for relevant traits of cage mates (e.g. body

temperature or thermogenesis, for studies of sleep) when parameters such as DGE

or a drug treatment are of primary interest. In turn, this will increase the robustness

of mouse studies and improve translation to humans.

Finally, we described specific challenges for igeGWAS and proposed solutions.

First, using simulations we demonstrated that spurious IGE associations can arise in

the presence of co-localized DGE even when all mice are strictly unrelated, as a result

of using each mouse as a focal individual and a cage mate in the analysis. We showed

that conditioning on direct genotypes at the tested locus yields calibrated igeGWAS P

values and avoids spurious IGE associations. Second, by comparing the number of

genome-wide significant loci from the igeGWAS and dgeGWAS of the same 170

phenotypes, we showed that power is typically lower for igeGWAS compared to

dgeGWAS. We demonstrated that dilution of IGE across cage mates, which occurs,

for example, when a focal mouse can interact only with one cage mate at a time, can

contribute to lower power of igeGWAS.

The methods we have developed to estimate the correlation ρ between DGE

and IGE and to carry out igeGWAS are very general and appropriate for many different

types of experimental designs. Together with the insights gained from our analysis of

170 phenotypes, they pave the way for future igeGWAS of biomedical phenotypes in

laboratory mice.

Materials and methods

Phenotypes and experimental variables

Phenotypes and experimental variables (covariates) for 1,934 Crl:CFW(SW)-US_P08 (CFW) mice were retrieved from http://wp.cs.ucl.ac.uk/outbredmice/. Phenotypes were normalized using the boxcox function (MASS package[64]) in R; phenotypes that could not be normalised satisfactorily (transformation parameter lambda outside of -2 to 2 interval) were excluded. The subset of covariates used for each phenotype is indicated in Supplementary Table 2. Because data for some phenotypes were missing for some mice, the sample size varied. The sample size for each phenotype after all filtering (see below) is indicated in Supplementary Table 2.

Cage information

Mice were four to seven weeks old when they arrived at the phenotyping facility and were housed in same-sex groups of three mice. They were left undisturbed for nine to twelve weeks during their time in quarantine and spent another four weeks together during phenotyping.

Cage assignments were not included in the publicly available dataset but were provided by the authors upon request and are now provided in **Supplementary Table 1**. Cage assignments were recorded at eleven time points throughout the study and showed that a few mice were taken out of their original cages and singly housed, presumably because they were too aggressive. We only included in our analyses mice that had the same two cage mates throughout the experiment. We further excluded a subset of mice based on their genotype-based genetic similarity, as described below.

Finally, all mice were singly housed during the sleep test and until sacrifice a few days

later. Hence, we investigated "persistent" IGE on sleep and tissue phenotypes.

Genome-wide genotypes

From http://wp.cs.ucl.ac.uk/outbredmice/ we retrieved both allele dosages for 7 million

variants and allele dosages for a subset of 353,697 high quality, LD-pruned variants

(as described in Nicod et al.[38]; genotyping based on sparse sequencing data). We

used high quality, LD-pruned variants for all analyses but the identification of candidate

genes at IGE loci (see below), for which we used the full set of variants.

Genetic relatedness matrix (GRM) and exclusion of presumed siblings

The genetic relatedness matrix was calculated as the cross-product of the LD-pruned

dosage matrix after standardizing the dosages for each variant to mean 0 and variance

1. A few pairs of mice were outliers in the distribution of GRM values, which made us

suspect that siblings had been included in the sample even though they were not

supposed to be (siblings were excluded by design). To mitigate confounding of DGE

and IGE analyses by litter effects, we excluded 19 cages (57 mice) from all analyses.

Variance components model

The same model as described in detail in Baud et al.[11] was used. Briefly, the model

used is the following:

$$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} \qquad (0)$$

y_f is the phenotypic value of the focal mouse f, X_f is a row of the matrix X of covariate

values and b a column vector of corresponding estimated coefficients. $\underline{a_{D,f}}$ is the

additive direct genetic effects (DGE) of f. Z_f is a row of the matrix Z that indicates

cage mates (importantly $Z_{i,i} = 0$) and $\underline{a_S}$ the column vector of additive indirect (social)

genetic effects (IGE). $\underline{e_D}$ refers to direct environmental effects (DEE) and $\underline{e_S}$ to indirect

(social) environmental effects (IEE). W_f is a row of the matrix W that indicates cage

assignment and c the column vector of cage effects.

The joint distribution of all random effects is defined as:

$$\begin{bmatrix} \underline{a_D} \\ \underline{a_S} \\ \underline{e_D} \\ \underline{e_S} \\ \underline{c} \end{bmatrix} \sim \text{MVN}\left(0, \begin{bmatrix} \sigma_{A_D}^2 A & \sigma_{A_{DS}} A & 0 & 0 & 0 \\ \sigma_{A_{DS}} A^T & \sigma_{A_S}^2 A & 0 & 0 & 0 \\ 0 & 0 & \sigma_{E_D}^2 I & \sigma_{E_{DS}} I & 0 \\ 0 & 0 & \sigma_{E_{DS}} I^T & \sigma_{E_S}^2 I & 0 \\ 0 & 0 & 0 & 0 & \sigma_C^2 I \end{bmatrix}\right)$$

where A is the GRM matrix and I the identity matrix.

The phenotypic covariance is:

$$C_{i,j} = cov(y_i, y_j) = \sigma_{A_D}^2 A_{i,j} + \sigma_{A_{DS}}\left\{(AZ^T)_{i,j} + (ZA^T)_{i,j}\right\} + \sigma_{A_S}^2 (ZAZ^T)_{i,j}$$

$$+ \sigma_{E_D}^2 I_{i,j} + \sigma_{E_{DS}}\left\{(IZ^T)_{i,j} + (ZI^T)_{i,j}\right\} + \sigma_{E_S}^2 (ZIZ^T)_{i,j}$$

$$+ \sigma_C^2 (WIW^T)_{i,j}$$

The variances explained by DGE and IGE were calculated respectively as

$sampleVar(\sigma_{A_D}^2 A) / sampleVar(C)$ and $sampleVar(\sigma_{A_S}^2 (ZAZ^T)) / sampleVar(C)$

where $sampleVar$ is the sample variance of the corresponding covariance matrix:

suppose that we have a vector $\underline{x}$ of random variables with covariance matrix M, the

sample variance of M is calculated as

$sampleVar(M) = \frac{Tr(PMP)}{n-1}$

Tr denotes the trace, n is the sample size, and $P = I - \frac{11'}{n}$ [65,66].

For those phenotypes where body weight was included as a covariate, we

checked that this did not lead to systematically increased (or decreased) estimates of

the aggregate contribution of IGE (collider bias).

Significance of the IGE variance component was assessed using a two-degree

of freedom log likelihood ratio (LLR) test (for the variance component and the

covariance with DGE). Note that this testing procedure is conservative. The Q value

for the aggregate contribution of IGE was calculated for each phenotype using the R

package qvalue[67]. Significant IGE contributions were reported at FDR < 10%

(corresponding to Q value < 0.1).

Alternative use of a less general model

When all cages have the same number of mice, as is the case in this study, the non-

genetic random effects are not identifiable[17,68]. An equivalent model can, in that case,

be defined as:

$cov\left(e_i, e_j\right) = \sigma_E^2 = \sigma_{E_D}^2 + \sigma_{E_S}^2 + \sigma_C^2$ if $i = j$

$cov\left(e_i, e_j\right) = \rho\sigma_E^2 = 2\,\sigma_{E_{DS}} + \sigma_C^2$ if $i \neq j$ and i and j share a cage

$cov\left(e_i, e_j\right) = 0$ if i and j are in different cages

We checked that both model (0) and this alternative model yielded the same genetic

estimates and maximum likelihoods. The alternative model was fitted using the

SimplifNonIdableEnvs option in LIMIX.

Correlation between DGE and IGE

The correlation ρ between $\underline{a_D}$ and $\underline{a_S}$ was calculated as:

$$\rho = \frac{\sigma_{A_{DS}}}{\sigma_{A_D} \times \sigma_{A_S}}$$

We tested whether ρ was significantly different from 0 and whether |ρ| was significantly

different from 1 using a one-degree of freedom LLR test.

Simulations for Supplementary Figure 1.

Phenotypes were simulated based on the genotypes and cage relationships of the full set of 1,812 mice. Phenotypes were drawn from model (0) with the following parameters: IGE explaining between 0 and 35.7% of phenotypic variance, DGE explaining 15% of phenotypic variance, $\rho_{A_{DS}} = 0.47$, DEE explaining 22% pf phenotypic variance, IEE explaining 16% of phenotypic variance, $\rho_{E_{DS}} = -0.97$, and cage effects explaining 26% of phenotypic variance. These variances correspond to the median value of estimates across traits with aggregate IGE and DGE > 5%. After building the phenotypic covariance matrix, the sample variance of the simulations was calculated and used to calculate "realised" simulation parameters from the "target" parameters above. The realised parameters were used for comparison with the parameters estimated from the simulations.

Definition of "social genotype" for igeGWAS

We assumed additive effects across cage mates and calculated the "social genotype" of a mouse as the sum of the reference allele dosages of its cage mates. The same assumption was made by Biscarini *et al.*[37] and Brinker *et al.*[28] among others.

Correlation between direct and social genotypes at a variant (Supplementary Figure 2)

Spearman's rank correlation coefficient was used. We tested whether the correlation was different from 0 using the function cor.test in the R package stats[69].

Models used for igeGWAS and dgeGWAS

To test IGE of a particular variant, we compared the following two models:

$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + G_f b_D$ (1, null)

$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + G_f b_D + Z_f G b_S$ (2, alternative)

Here, G is the vector of direct genotypes at the tested variant, b_D the estimated

coefficient for local DGE and b_S the estimated coefficient for local IGE. In igeGWAS,

we refer to the inclusion of $G_f b_D$ in model (1, null) as "conditioning".

The models were fitted using LIMIX[70,71] with the covariance of the model

estimated only once per phenotype, in the null model with no local genetic effect

(model 0).

The significance of local IGE was calculated by comparing models (1) and (2)

with a 1-degree of freedom LLR test.

dgeGWAS was carried out by comparing model (2) above to the null model (3)

below:

$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + Z_f G b_S$ (3, null)

In dgeGWAS, we refer to the inclusion of $Z_f G b_S$ in model (3, null) as "conditioning".

Identification of genome-wide significant associations

We used a genome-wide permutation strategy to control the FDR for each phenotype,

as done by Nicod et al.(36). This strategy takes into account the specific patterns of

linkage disequilibrium present in the sample and identifies genome-wide significant

associations *for each phenotype independently of the results for the other phenotypes*

in the dataset. More precisely, for each phenotype and for each type of genetic effect

(social and direct), we performed 100 "permuted GWAS" by permuting the rows of the

matrix of social (respectively direct) genotypes, and testing each variant at a time using

the permuted genotypes together with the un-permuted phenotypes, un-permuted

covariates, un-permuted GRM and un-permuted matrix of direct (respectively social)

genotypes (for conditioning). See [71,73] for references on this permutation approach.

For a given P value x, the per-phenotype FDR can be calculated as:

$$FDR(x) = \frac{\#\ loci\ with\ P < x\ in\ permuted\ data}{100\ \times\ \#\ loci\ with\ P < x\ in\ unpermuted\ data}$$

We reported those loci with FDR < 10%.

Definition of candidate genes at associated loci

At each significantly associated locus we defined a 1.5Mb window centred on the lead

variant corresponding, in this sample, to the 95% confidence interval for the

association[38]. We identified all the variants that segregate in this window based on the

full set of 7M variants and reran igeGWAS and dgeGWAS locally using all the variants

at the locus. We defined "candidate" genes as those genes that either overlapped the

associated plateau or were located in direct proximity, and whose MGI symbol does

not start by 'Gm', 'Rik', 'Mir', 'Fam', or 'Tmem' in order to focus on genes with known

function and generate more tractable hypotheses on the pathways of social effects.

Gene expression in the hippocampus of an independent sample of CFW mice

Gene expression in the hippocampus of an independent sample of 79 male CFW mice,

initially published in Parker et al.[40], was available from GeneNetwork

(http://gn2.genenetwork.org/). The data are accessible by selecting Mouse as

Species, CFW Outbred GWAS as *Group*, Hippocampus mRNA as *Type*, and UCSD

CFW Hippocampus (Jan17) RNA-Seq Log2 Z-score as *Dataset*. To retrieve the genes

whose expression is most highly correlated with that of *Epha4*, we entered "Epha4" in

the *Get Any* field. Following selection of the Epha4 record (click on

ENSMUSG00000026235), we used *Calculate Correlations* with Sample r as *Method*,

UCSD CFW Hippocampus (Jan17) RNA-Seq Log2 Z-score as *Database*, and

Spearman rank as correlation *Type*. Supplementary Figure 6 was obtained by clicking

on the value of the correlation between Epha4 and Dlgap1 expression levels (column

Sample rho).

Variance explained by a genome-wide significant association

The variance explained by a genome-wide significant IGE association was estimated

in an extension of model (0) with additional fixed effects for both direct and social

effects of lead SNPs at all genome-wide significant IGE loci (the lead SNP being the

SNP with the most significant P value at the locus in the igeGWAS). After fitting the

model, the variance was calculated as:

583
$$\frac{var(ZG\widehat{b_S})}{\sum var(X_c\widehat{b_c}) + \sum var(G\widehat{b_D}) + \sum var(ZG\widehat{b_S}) + sampleVar(C)}$$

582 where $sampleVar(C)$ is the sample variance of the covariance matrix in this model.

584 The variance explained by a genome-wide significant DGE association was

585 estimated in a similar model but considering all genome-wide significant DGE

586 associations and

587 calculated as:

589
$$\frac{var(ZG\widehat{b_D})}{\sum var(X_c\widehat{b_c}) + \sum var(G\widehat{b_D}) + \sum var(ZG\widehat{b_S}) + sampleVar(C)}$$

Simulations for Supplementary Figure 2c and 2d.

Phenotypes were simulated based on the genotypes and cage relationships of the

1,812 mice. Phenotypes were simulated as the sum of random effects and local DGE

(from model (1)), with the following parameters: $\sigma^2_{A_D}$ = 5 or 20, $\sigma^2_{A_S}$ = 5 or 20, $\rho_{A_{DS}}$= 0.5,

$\sigma^2_{E_D}$= 30, $\sigma^2_{E_S}$= 30, $\rho_{E_{DS}}$ = -0.97, σ^2_C = 25. The values for $\rho_{A_{DS}}$, $\sigma^2_{E_D}$, $\sigma^2_{E_S}$, $\rho_{E_{DS}}$, and σ^2_C

were close to the median of the corresponding estimates from the real data. $\sigma_{A_D}^2 = 5$

and $\sigma_{A_S}^2 = 5$ correspond to low polygenic effects in the real data, and $\sigma_{A_D}^2 = 20$ and

$\sigma_{A_S}^2 = 20$ correspond to high polygenic effects in the real data. We simulated local

DGE at random variants in the genome, and simulated variances of 0, 5, 20 or 50.

The results we show in Supplementary Figure 2c and 2d are based on a subset

of simulations: $\sigma_{A_D}^2 = 20$ and $\sigma_{A_S}^2 = 20$ and local DGE variance of 20.

Simulations for Supplementary Figure 3a-d, and Figure 3a.

Phenotypes were simulated based on the real genotypes but random cages.

Phenotypes were simulated as the sum of random effects, local DGE and local IGE

(model (2)) with the following parameters: $\sigma_{A_D}^2 = 17$, $\sigma_{A_S}^2 = 17$, $\rho_{A_{DS}} = 0.65$, $\sigma_{E_D}^2 = 19$,

$\sigma_{E_S}^2 = 15$, $\rho_{E_{DS}} = -0.8$, $\sigma_C^2 = 25$. Those values correspond to the median estimates for

phenotypes with aggregate IGE and DGE > 0.1.

We simulated local IGE and DGE at variants where direct and social genotypes

were either lowly correlated (Spearman correlation negative log P value < 0.05) or

more highly correlated (Spearman correlation negative log P value > 0.2), and had

with low MAF (MAF < 0.05), medium MAF (0.225<MAF<0.275) or high MAF

(MAF>0.45). We simulated local DGE with an allelic effect of 0 or 1 (1 corresponds to

a large effect in the real data). We simulated local IGE under two alternative generative

models: an "additive" model by using Z as in model (2) (i.e. filled with 0s and 1s) or a

"proportional" model by using $Z' = \frac{Z}{N}$. In all cases we simulated an allelic effect of 0.2

(similar to the average allelic effect estimated in the IGE GWAS). The sample variance

of the simulated local DGE term is $var(Gb_D) = 2p(1-p)\,b_D^2$; it is $var(ZGb_S) =$

$2Np(1-p)\,b_S^2$ for the local IGE term simulated under the additive model, and

$var\left(\frac{Z}{N}Gb_S\right) = \frac{2Np(1-p)}{N^2}b_S^2$ for the local IGE component simulated under the

proportional model.

The results we show in Supplementary Figure 3a-d are based on simulations

averaged across low, medium and high MAF. Power was calculated at a genome-wide

significance threshold of negative log P 5, which is similar to the significance of

associations detected at FDR < 10%.

The results we show in Figure 3b are based on a subset of simulations with no

local DGE, averaged across high and low genotypic correlations. Power was also

calculated at a genome-wide significance threshold of negative log P 5.

Scripts used in this study

630 All the scripts used in this study are available from http://github.com/limix/IGE.

631 LIMIX can be downloaded from http://github.com/limix/limix.

Acknowledgements

634 The authors want to thank Drs. Na Cai and Robert W. Davies and Richard Mott for

635 facilitating access to the genotype data from Nicod et al.(36), as well as Dr. Paul

636 Franken for a discussion of the sleep measure reported in this manuscript. AB was

637 supported by a fellowship from the Wellcome Trust (105941/Z/14/Z).

Authors' contributions

640 AB and OS designed the study. AB and FPC performed the analyses. AB, FPC, JN

641 and OS interpreted the results. AB and OS wrote the manuscript with input from FPC

642 and JN.

References

1.   Griffing B. Selection in reference to biological groups I. Individual and group selection applied to populations of unordered groups. Australian Journal of Biological Sciences. 1967;20(1):127-40.

2.   Moore AJ, Brodie III ED, Wolf JB. Interacting phenotypes and the evolutionary process: I. Direct and indirect genetic effects of social interactions. Evolution. 1997;51(5):1352-62.

3.   Wolf JB, Brodie III ED, Cheverud JM, Moore AJ, Wade MJ. Evolutionary consequences of indirect genetic effects. Trends in ecology & evolution. 1998;13(2):64-9.

4.   Wolf JB, Mutic JJ, Kover PX. Functional genetics of intraspecific ecological interactions in Arabidopsis thaliana. Philosophical Transactions of the Royal Society B: Biological Sciences. 2011;366(1569):1358-67.

5.   Bailey NW, Hoskins JL. Detecting cryptic indirect genetic effects. Evolution. 2014;68(7):1871-82.

6.   Beck JA, Lloyd S, Hafezparast M, Lennon-Pierce M, Eppig JT, Festing MF, et al. Genealogies of mouse inbred strains. Nature genetics. 2000;24(1):23-5.

7.   Bleakley BH, Brodie ED. Indirect genetic effects influence antipredator behavior in guppies: estimates of the coefficient of interaction psi and the inheritance of reciprocity. Evolution. 2009;63(7):1796-806.

8.   Ashbrook DG, Gini B, Hager R. Genetic variation in offspring indirectly influences the quality of maternal behaviour in mice. Elife. 2015;4:11814.

9.   Ashbrook DG, Sharmin N, Hager R, editors. Offspring genes indirectly influence sibling and maternal behavioural strategies over resource share. Proc R Soc B; 2017: The Royal Society.

10.   Baud A, Mulligan MK, Casale FP, Ingels JF, Bohl CJ, Callebert J, et al. Genetic

variation in the social environment contributes to health and disease. PLoS genetics.

2017;13(1):e1006498.

11.   Head ML, Berry LK, Royle NJ, Moore AJ. Paternal care: direct and indirect genetic

effects of fathers on offspring performance. Evolution. 2012;66(11):3570-81.

12.   Mutic JJ, Wolf JB. Indirect genetic effects from ecological interactions in Arabidopsis

thaliana. Molecular ecology. 2007;16(11):2371-81.

13.   Petfield D, Chenoweth SF, Rundle HD, Blows MW. Genetic variance in female

condition predicts indirect genetic variance in male sexual display traits. Proceedings of the

National Academy of Sciences. 2005;102(17):6045-50.

14.   Wilson AJ, Gelin U, Perron M-C, Réale D. Indirect genetic effects and the evolution of

aggression in a vertebrate system. Proceedings of the Royal Society of London B: Biological

Sciences. 2009;276(1656):533-41.

15.   Moore AJ, Haynes KF, Preziosi RF, Moore PJ. The evolution of interacting

phenotypes: genetics and evolution of social dominance. the american naturalist.

2002;160(S6):S186-S97.

16.   Bergsma R, Kanis E, Knol EF, Bijma P. The contribution of social effects to heritable

variation in finishing traits of domestic pigs (Sus scrofa). Genetics. 2008;178(3):1559-70.

17.   Ellen ED, Visscher J, van Arendonk JA, Bijma P. Survival of laying hens: genetic

parameters for direct and associative effects in three purebred layer lines. Poultry science.

2008;87(2):233-9.

18.   Alemu SW, Bijma P, Møller SH, Janss L, Berg P. Indirect genetic effects contribute

substantially to heritable variation in aggression-related traits in group-housed mink (Neovison

vison). Genetics Selection Evolution. 2014;46(1):30.

19.   e Silva JC, Potts B, Gilmour A, Kerr R. Genetic-based interactions among tree

neighbors: identification of the most influential neighbors, and estimation of correlations

among direct and indirect genetic effects for leaf disease and growth in Eucalyptus globulus.

Heredity. 2017;119(3):125.

20.   Wilson A, Coltman D, Pemberton J, Overall A, Byrne K, Kruuk L. Maternal genetic

effects set the potential for evolution in a free-living vertebrate population. Journal of

evolutionary biology. 2005;18(2):405-14.

21.   McAdam AG, Boutin S, Réale D, Berteaux D. Maternal effects and the potential for

evolution in a natural population of animals. Evolution. 2002;56(4):846-51.

22.   Hadfield JD, Burgess MD, Lord A, Phillimore AB, Clegg SM, Owens IP. Direct versus

indirect sexual selection: genetic basis of colour, size and recruitment in a wild bird.

Proceedings of the Royal Society B: Biological Sciences. 2006;273(1592):1347-53.

23.   Bates TC, Maher BS, Medland SE, McAloney K, Wright MJ, Hansell NK, et al. The

Nature of Nurture: Using a Virtual-Parent Design to Test Parenting Effects on Children's

Educational Attainment in Genotyped Families. Twin Research and Human Genetics. 2018:1-

11.

24.   Kong A, Thorleifsson G, Frigge ML, Vilhjalmsson BJ, Young AI, Thorgeirsson TE, et

al. The nature of nurture: Effects of parental genotypes. Science. 2018;359(6374):424-8.

25.   Warrington NM, Beaumont RN, Horikoshi M, Day FR, Helgeland Ø, Laurin C, et al.

Maternal and fetal genetic effects on birth weight and their relevance to cardio-metabolic risk

factors. Nature genetics. 2019:1.

26.   McGlothlin JW, Brodie III ED. HOW TO MEASURE INDIRECT GENETIC

EFFECTS: THE CONGRUENCE OF TRAIT-BASED AND VARIANCE-PARTITIONING

APPROACHES. Evolution. 2009;63(7):1785-95.

27.   Sotoudeh R, Harris KM, Conley D. Effects of the peer metagenomic environment on

smoking behavior. Proceedings of the National Academy of Sciences. 2019;116(33):16302-7.

28.   Bijma P. The quantitative genetics of indirect genetic effects: a selective review of

modelling issues. Heredity. 2014;112(1):61.

29.   Wilson AJ, Morrissey M, Adams M, Walling CA, Guinness FE, Pemberton JM, et al.

Indirect genetics effects and evolutionary constraint: an analysis of social dominance in red

deer, Cervus elaphus. Journal of evolutionary biology. 2011;24(4):772-83.

30.   McGinnis R, Steinthorsdottir V, Williams NO, Thorleifsson G, Shooter S, Hjartardottir

S, et al. Variants in the fetal genome near FLT1 are associated with risk of preeclampsia. Nature

genetics. 2017;49(8):1255.

31.   Brinker T, Bijma P, Vereijken A, Ellen ED. The genetic architecture of socially-

affected traits: a GWAS for direct and indirect genetic effects on survival time in laying hens

showing cannibalism. Genetics Selection Evolution. 2018;50(1):38.

32.   de Oliveira PC, Zaniboni CR, Carmona IM, Fonseca AR, Canto-de-Souza A.

Preliminary behavioral assessment of cagemates living with conspecifics submitted to chronic

restraint stress in mice. Neuroscience letters. 2017;657:204-10.

33.   Bolger N, DeLongis A, Kessler RC, Wethington E. The contagion of stress across

multiple roles. Journal of Marriage and the Family. 1989:175-83.

34.   Haeffel GJ, Hames JL. Cognitive Vulnerability to Depression Can Be Contagious.

Clinical Psychological Science. 2014;2(1):75-85.

35.   Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, et al. Gut microbiota

from twins discordant for obesity modulate metabolism in mice. Science.

2013;341(6150):1241214.

36.   Nicod J, Davies RW, Cai N, Hassett C, Goodstadt L, Cosgrove C, et al. Genome-wide

association of multiple complex traits in outbred mice by ultra-low-coverage sequencing.

Nature genetics. 2016;48(8):912.

37.   Davies RW, Flint J, Myers S, Mott R. Rapid genotype imputation from sequence

without reference panels. Nature genetics. 2016;48(8):965.

38.   Biscarini F, Bovenhuis H, Van Der Poel J, Rodenburg T, Jungerius A, Van Arendonk

J. Across-line SNP association study for direct and associative effects on feather damage in

laying hens. Behavior genetics. 2010;40(5):715-27.

39.   Beavis W. QTL analyses: power, precision and accuracy. Molecular Dissection of

Complex Traits. Edited by: AH P. 1997. CRC Press, New York.

40.   Beavis W, Beavis W. The power and deceit of QTL experiments: lessons from

comparative QTL studies. 1994.

41.   Bijma P. Estimating indirect genetic effects: precision of estimates and optimum

designs. Genetics. 2010;186(3):1013-28.

42.   Bijma P. Multilevel selection 4: modeling the relationship of indirect genetic effects

and group size. Genetics. 2010;186(3):1029-31.

43.   Yalcin B, Nicod J, Bhomra A, Davidson S, Cleak J, Farinelli L, et al. Commercially

available outbred mice for genome-wide association studies. PLoS genetics.

2010;6(9):e1001085.

44.   Parker CC, Gopalakrishnan S, Carbonetto P, Gonzales NM, Leung E, Park YJ, et al.

Genome-wide association study of behavioral, physiological and gene expression traits in

outbred CFW mice. Nature genetics. 2016;48(8):919.

45.   Li Y, Wang H, Wang X, Liu Z, Wan Q, Wang G. Differential expression of

hippocampal EphA4 and ephrinA3 in anhedonic-like behavior, stress resilience, and

antidepressant drug treatment after chronic unpredicted mild stress. Neuroscience letters.

2014;566:292-7.

46. Zhang J-c, Yao W, Qu Y, Nakamura M, Dong C, Yang C, et al. Increased EphA4-

ephexin1 signaling in the medial prefrontal cortex plays a role in depression-like phenotype.

Scientific Reports. 2017;7(1):7133.

47. Bouvier D, Corera AT, Tremblay MÈ, Riad M, Chagnon M, Murai KK, et al. Pre-

synaptic and post-synaptic localization of EphA4 and EphB2 in adult mouse forebrain. Journal

of neurochemistry. 2008;106(2):682-95.

48. Murai KK, Nguyen LN, Irie F, Yamaguchi Y, Pasquale EB. Control of hippocampal

dendritic spine morphology through ephrin-A3/EphA4 signaling. Nature neuroscience.

2003;6(2):153-60.

49. Coba M, Ramaker M, Ho E, Thompson S, Komiyama N, Grant S, et al. Dlgap1

knockout mice exhibit alterations of the postsynaptic density and selective reductions in

sociability. Scientific reports. 2018;8(1):1-12.

50. Poppe L, Rué L, Timmers M, Lenaerts A, Storm A, Callaerts-Vegh Z, et al. EphA4 loss

improves social memory performance and alters dendritic spine morphology without changes

in amyloid pathology in a mouse model of Alzheimer's disease. Alzheimer's Research &

Therapy. 2019;11(1):1-13.

51. Coba M, Ramaker M, Ho E, Thompson S, Komiyama N, Grant S, et al. Dlgap1

knockout mice exhibit alterations of the postsynaptic density and selective reductions in

sociability. Scientific reports. 2018;8(1):2281.

52. Marques AC, Dupanloup I, Vinckenbosch N, Reymond A, Kaessmann H. Emergence

of young human genes after a burst of retroposition in primates. PLoS biology.

2005;3(11):e357.

53.   Betrán E, Wang W, Jin L, Long M. Evolution of the phosphoglycerate mutase

processed gene in human and chimpanzee revealing the origin of a new primate gene.

Molecular biology and evolution. 2002;19(5):654-63.

54.   Cheng Y, Jiang L, Keipert S, Zhang S, Hauser A, Graf E, et al. Prediction of Adipose

Browning Capacity by Systematic Integration of Transcriptional Profiles. Cell reports.

2018;23(10):3112-25.

55.   Sokoloff G, Blumberg MS. Competition and cooperation among huddling infant rats.

Developmental Psychobiology: The Journal of the International Society for Developmental

Psychobiology. 2001;39(2):65-75.

56.   Wisden W. Sleep and thermoregulation Edward C Harding, Nicholas P Franks and

William Wisden. Current Opinion in Physiology. 2020;15:7-13.

57.   Harding EC, Yu X, Miao A, Andrews N, Ma Y, Ye Z, et al. A neuronal hub binding

sleep initiation and body cooling in response to a warm external stimulus. Current Biology.

2018;28(14):2263-73. e4.

58.   Raymann RJ, Swaab DF, Van Someren EJ. Skin deep: enhanced sleep depth by

cutaneous temperature manipulation. Brain. 2008;131(2):500-13.

59.   Gordon C. Thermal physiology of laboratory mice: defining thermoneutrality. Journal

of thermal biology. 2012;37(8):654-85.

60.   Commons KG, Cholanians AB, Babb JA, Ehlinger DG. The rodent forced swim test

measures stress-coping strategy, not depression-like behavior. ACS chemical neuroscience.

2017;8(5):955-60.

61.   Iio W, Matsukawa N, Tsukahara T, Kohari D, Toyoda A. Effects of chronic social

defeat stress on MAP kinase cascade. Neuroscience letters. 2011;504(3):281-4.

62.   Hebert M, Evenson A, Lumley L, Meyerhoff J. Effects of acute social defeat on activity

in the forced swim test: Parametric studies in DBA/2 mice using a novel measurement device.

Aggressive Behavior: Official Journal of the International Society for Research on Aggression.

1998;24(4):257-69.

63.   Iñiguez SD, Riggs LM, Nieto SJ, Dayrit G, Zamora NN, Shawhan KL, et al. Social

defeat stress induces a depression-like phenotype in adolescent male c57BL/6 mice. Stress.

2014;17(3):247-55.

64.   Koike H, Ibi D, Mizoguchi H, Nagai T, Nitta A, Takuma K, et al. Behavioral

abnormality and pharmacologic response in social isolation-reared mice. Behavioural brain

research. 2009;202(1):114-21.

65.   Berry A, Bellisario V, Capoccia S, Tirassa P, Calza A, Alleva E, et al. Social

deprivation stress is a triggering factor for the emergence of anxiety-and depression-like

behaviours and leads to reduced brain BDNF levels in C57BL/6J mice.

Psychoneuroendocrinology. 2012;37(6):762-72.