Comparative architectures of direct and social genetic effects from the genome-wide association study of 170 phenotypes in outbred laboratory mice

Amelie Baud[1*], Francesco Paolo Casale[1,2], Jerome Nicod[3], Oliver Stegle[1,4*]

[1] European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, CB10 1SD Hinxton, Cambridge, UK

[2] Microsoft Research New England, Cambridge, Massachusetts, USA

[3] Wellcome Trust Centre for Human Genetics, Roosevelt Drive, Oxford OX3 7BN, UK

[4] European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, DE

* Corresponding authors: abaud@ebi.ac.uk and oliver.stegle@embl.de

Abstract

Social genetic effects (SGE, also called indirect genetic effects) are associations between genotypes of one individual and phenotype of another. SGE can arise when two individuals interact and heritable traits of one influence the phenotype of the other. To better understand how SGE arise from individual genetic loci, we have derived statistical strategies for the joint analysis of SGE and traditional direct genetic effects (DGE, effects of an individual's genotypes on its own phenotype), which we applied to a dataset of 170 behavioural, physiological and morphological phenotypes measured in 1,812 outbred laboratory mice. Genome-wide association study of SGE and DGE identified 24 and 120 genome-wide significant loci respectively (FDR < 10%). We found no overlap between genome-wide significant SGE and DGE loci acting on the same phenotype, and a moderate correlation between SGE and DGE genome-wide, demonstrating that sgeGWAS has the potential to identify novel genetic loci contributing to inter-individual differences. We focused on two such loci to illustrate how individual SGE loci may help identify traits of social partners mediating social effects. Finally, we contrasted the effect sizes of genome-wide significant SGE and

33 DGE loci and found smaller effect sizes for the former: SGE associations explained a
34 maximum of 2.5% of phenotypic variance when eleven genome-wide significant DGE
35 associations explained more than 5% of phenotypic variance. Our results and the
36 presented methods will guide the design and analysis of future sgeGWAS.
37
38

Main text

Introduction

Social interactions between two individuals can result in the phenotype of one
individual being affected by genotypes of the other. Such effects arise when the
phenotype of a focal individual is influenced by heritable traits of their social partner
(Figure 1a) and are called social genetic effects (SGE) or indirect genetic effects[1-4].
SGE have been shown to contribute significantly and substantially to
phenotypic variation in livestock, wild animals, plants and laboratory model
organisms[5]. In laboratory mice, SGE have been found to affect behavioural,
physiological, and morphological traits, including phenotypes not *a priori* expected to
be socially affected such as wound healing[6-9]. In humans, effects of non-transmitted
parental alleles have been detected on offspring's educational attainment[10-12]. Thus,
SGE are an important component of the genotype to phenotype path.
In addition, SGE may be used as a tool to investigate social effects, in particular
identify traits of social partners affecting a phenotype of interest, measured in focal
individuals. Indeed, identifying individual variants of social partners associated with
the phenotype of interest, through genome-wide association study of SGE
(sgeGWAS), could provide new insights into the pathways of social effects in a similar
manner as GWAS of "traditional" direct genetic effects (DGE, effects of an individual's
genotypes on its own phenotype) has provided insights into the "within-body"
pathways of phenotypic variation[13]. However, robust methods to perform sgeGWAS
are not established and such experiments will require information on the architecture
of SGE[14].
The architecture of SGE is largely unknown, as few studies have identified
genomic loci that influence phenotypes of social partners. Quasi-Mendelian SGE

exist[15,16] but candidate gene studies and GWAS of SGE have revealed oligo- or

polygenic architectures for a larger number of phenotypes[7,8,15-18]. Furthermore, it is

currently unknown to what extent SGE loci overlap with DGE loci and how much

phenotypic variance individual SGE loci explain. To address these questions and

identify loci underlying SGE, we leveraged an existing dataset of 170 behavioural,

physiological and morphological phenotypes measured in outbred "CFW" laboratory

mice[19,20] and studied both SGE and DGE acting on each phenotype (Figure 1b and

1c). Our results and the methods we detail and validate will inform the design and

analysis of future sgeGWAS carried out in outbred populations with any levels of

relatedness and group sizes.

***Figure 1** Illustration of social genetic effects (SGE), definition of specific terms used*

throughout the manuscript, and experimental design. (a) SGE arise when two

individuals interact and heritable traits of one influence the phenotype of the other. In

other words, genotypes of the "social partner" influence - through "mediating traits" -

the "phenotype of interest", measured in the "focal individual". Importantly, these

mediating traits do not need to be measured or known. (b) 1,812 "CFW" outbred mice

were housed in groups of 3 mice per cage. SGE and DGE contributed by each mouse

were modelled, such that each mouse served as both focal individual and cage mate

in our analyses. (c) Experimental design. The experimental paradigms used to collect
the 170 phenotypes are listed.

90
Results

Genome-wide genotypes (both LD-pruned and non pruned, see Methods) and
phenotypes for 2,073 CFW mice were available from Nicod et al.[19] and Davies et al.[20].
Males were always housed with males and females with females, and mice interacted
in their cages for a period of at least nine weeks before phenotyping (Figure 1c). We
only kept mice that had the same two cage mates over the course of the experiment
(1,869 total). Furthermore, we excluded 57 mice that formed genetic substructures so
that the remaining 1,812 mice were as equally related as possible while retaining as
large a sample size as possible (see Methods).

Aggregate contribution of SGE
Initially we estimated the aggregate contribution of SGE (i.e. the sum of SGE across
the genome) to all 170 phenotypes in the dataset, which include phenotypes that have
never before been considered in a study of SGE (muscles and bones characteristics,
response to hypoxia, electrocardiogram, and sleep). We used a random effect model
for SGE and DGE as in Baud et al.[9] and found that SGE, in aggregate, explained up
to 22% (+/- 6%) of variation in serum LDL levels and an average of 11% across 9
phenotypes with significant aggregate SGE (FDR < 10%, Supplementary Table 1).
Phenotypes significantly affected by SGE included behavioural (helplessness),
physiological (serum LDL cholesterol, healing from an ear punch, blood eosinophils,
serum alpha-amylase concentration, blood platelets, acute hypoxic response), and
morphological (weight of adrenal glands) traits. Importantly, among the phenotypes
significantly affected by SGE were both phenotypes known or expected to be affected
by the social environment, namely helplessness (a murine model for depression),
blood eosinophils (a measure of immunity) and weight of adrenal glands (a measure
associated with stress), and phenotypes less expected to be socially affected, such
as LDL levels, response to hypoxia, and healing from an ear punch. Notably, the
significant and substantial SGE detected in this population for healing from an ear
punch (P = 0.00018, variance explained by SGE in aggregate = 17 +/-4%) constitute

the third independent line of evidence that SGE affect wound healing in laboratory

mice[9]. Altogether, our results demonstrate that SGE may affect a broad range of

phenotypes including traits not *a priori* expected to be socially affected, which is

consistent with prior evaluation of SGE in other populations[9].

Correlation between SGE and DGE loci acting on the same phenotype

The polygenic model used for variance decomposition fits a correlation coefficient ρ

(See Methods) that quantifies the extent to which SGE and DGE acting on the same

phenotype arise from similar loci. We stress that, in this section and throughout the

manuscript, we compare SGE and DGE *acting on the same phenotype* (but do this for

all phenotypes). Using simulations, we observed that ρ can be accurately estimated,

provided the aggregate contribution of SGE and DGE is sufficiently large

(Supplementary Figure 1). Consequently, we limited the analysis of ρ to 28

phenotypes for which both aggregate SGE and aggregate DGE explained at least 5%

of phenotypic variation. The SGE-DGE correlation coefficient ρ varied between -0.28

and 1 across these traits, with an average of 0.42 (Figure 2 and Supplementary Table

1). For 10 out of the 28 phenotypes ρ was significantly different from zero (nominal P

< 0.05, Figure 2 and Supplementary Table 1), indicating that loci exist that influence

both the phenotype of the focal animal (DGE) and the same phenotype of cage mates

(SGE). Strong evidence for shared SGE and DGE loci ($\rho \neq 0$ at Bonferroni-corrected

P < 0.05) was observed for healing from an ear punch, weight of adrenal glands, serum

LDL cholesterol levels, and mean platelet volume.

We also evaluated evidence that $|\rho|$ was different from one (i.e. ρ different from

one and minus one) in order to empirically evaluate the widely-influential model of

"phenotypic contagion". Phenotypic contagion or "spread" is a model for social effects

whereby the phenotype of interest of a focal individual is affected by the same

phenotype of their social partners. In humans, cognitive susceptibility to depression,

alcohol consumption, stress, obesity and educational attainment, for example, have

been suggested to be "contagious" or "spread" across college roommates, spouses,

friends, or parent/offspring[10-12,21-25]. As a result, phenotypic contagion has shaped the

way we think about social effects: for example, phenotypes unlikely to spread have

been used to cast doubt on social network effects[26]. Here we leveraged the parameter

ρ to test whether phenotypic contagion was sufficient to account for SGE: under a

model of pure phenotypic contagion, |ρ| is expected to be equal to one; on the contrary, if traits of social partners other than the phenotype of interest mediate social effects, |ρ| is expected to be different from one. We found that |ρ| was significantly different from one (nominal P < 0.05) for 10 out of 28 phenotypes (Figure 2 and Supplementary Table 1), with strong evidence (P < 0.05 after Bonferroni correction) for helplessness and healing from an ear punch. Our results empirically demonstrate that traits of social partners other than the phenotype of interest may be important, even when some level of contagion is likely, and motivate the use of sgeGWAS to uncover such traits.

Figure 2 *Correlation coefficients ρ between SGE and DGE random effects (see Methods). Shown are correlation coefficients for 28 phenotypes with error bars denoting standard errors. The stars denote significance for rejecting H_0: $\rho = 0$ (i.e. SGE and DGE arise from different loci, bottom row) and H_0: $|\rho| = 1$ (i.e. pure phenotypic contagion, top row). Black stars (*) denote nominal P value < 0.05, red stars (*) denote Bonferroni-corrected P value < 0.05.*

sgeGWAS and dgeGWAS of 170 phenotypes

To map SGE, we calculated the "social genotype" of a mouse at a variant as the sum of the reference allele dosages across its cage mates at the variant[17,18], and tested for association between social genotype and phenotype. In order to avoid spurious

associations, we accounted for background SGE, DGE and non-genetic effects using
random effects as in the model used for variance decomposition. In the sgeGWAS we
also accounted for DGE of the tested variant, by including direct genotypes at the
variant as a covariate (See Methods). Similarly, in the dgeGWAS we included social
genotypes at the variant as a covariate. We hereafter refer to this strategy as
"conditioning". We found that conditioning was necessary to avoid spurious
associations in the sgeGWAS due to co-localised DGE (Supplementary Figure 2). As
we show in the Supplementary Note, this problem originates from the use of each
mouse as both focal individual and cage mate in the analysis, a strategy that has been
used before to maximise sample size when all individuals are phenotyped and
genotyped[17,18]. Importantly, spurious associations may arise even if all individuals are
strictly unrelated (Supplementary Note). A power analysis suggested that conditioning
may slightly decrease power to detect SGE in the absence of co-localised DGE,
particularly when direct and social genotypes are highly correlated (Supplementary
Figure 3a and 3c) but would increase power if the locus also gave rise to DGE
(Supplementary Figures 3b and 3d).

In order to compare, for each phenotype, the results of sgeGWAS and
dgeGWAS, we defined a locus as the average size of the 95% confidence interval in
this population, namely 1.5Mb[19], and, following Nicod et al.[19], used a per-phenotype
FDR approach to account for multiple testing (see Methods). sgeGWAS identified 24
genome-wide significant loci for 17 of the 170 phenotypes (FDR < 10%; Figure 3 and
Supplementary Table 2). In comparison, dgeGWAS identified 120 genome-wide
significant loci for 63 phenotypes at the same threshold (Figure 3 and Supplementary
Table 3).

There was no overlap between genome-wide significant SGE and DGE loci
acting on the same phenotype. However, variants at genome-wide significant SGE
loci were enriched in small P values in the corresponding dgeGWAS (Supplementary
Figure 4). Together these results suggest a partially distinct basis for SGE and DGE
acting on the same phenotype (i.e. partially different loci and/or effect sizes), which is
consistent with the results from the analysis of the correlation parameter ρ.

Figure 3 *Superimposed manhattan plots corresponding to to sgeGWAS (top panel) and dgeGWAS (bottom panel) for 170 phenotypes. DGE associations with a negative log P value greater than 10 were truncated at this threshold (as indicated by 10+); data points with negative log P values smaller than 2 are not shown. Lead variants for all genome-wide significant SGE and DGE loci are represented with a larger dot. In the SGE panel, each color corresponds to a class of phenotypes: behavioural (red, includes 7 behavioural phenotypes with a detected SGE locus), adult neurogenesis (black, 2 phenotypes), immune (orange, 1 phenotype), haematological (yellow, 1 phenotype), blood biochemistry (blue, 2 phenotypes), bone phenotypes (green, 2 phenotypes), heart function (brown, 1 phenotype), and lung function (purple, 1 phenotype). In the DGE panel, a genome-wide significant locus is colored grey when the corresponding phenotype does not have a genome-wide significant SGE association; when the corresponding phenotype does have a genome-wide significant SGE association, the same color is used as in the SGE panel.*

Compared to many other mouse populations used for mapping, linkage disequilibrium decays rapidly in the CFW population, which facilitates identification of

candidate genes at associated loci[19,27]. At each genome-wide significant SGE locus
we reran the association analysis using the full set of variants (rather than the pruned
set used for GWAS) and prioritized candidate genes that were underneath or closest
to the most significantly associated genomic segment, excluding certain categories of
genes of unknown function (see Methods). This resulted in five genome-wide
significant SGE loci with a single candidate gene (Supplementary Table 2).

An example is an SGE locus on chromosome 1 (Figure 4a, top panel) for
immobility in the first two minutes of the Porsolt swim test, a measure of helplessness
in CFW mice[28]. *Epha4*, the candidate gene at this locus, is critical for the development
of the central nervous system (CNS) and synaptic plasticity in the adult brain[29]. *Epha4*
mutant mice show a range of altered phenotypes including abnormal CNS
morphology, altered limb coordination and lower locomotor activity, lower
reproductivity, and poor maternal nurturing[30]. *Epha4* has also been associated with
helplessness[31,32], but we found little evidence of DGE from *Epha4* on helplessness in
our study (Figure 4a, bottom panel), in contrast with strong evidence of SGE.
Expression data from the hippocampus of an independent sample of CFW mice[33],
mined using Genenetwork[34] (see Methods), and a second genome-wide significant
locus for helplessness in our study, located on chromosome 17, helped decipher the
pathways mediating the social effects of *Epha4*: of the eight genes prioritized as
candidate genes at the chromosome 17 locus (Figure 4b, top panel), one, *Dlgap1*, is
among the ten genes whose expression in the hippocampus is most strongly
correlated with that of *Epha4*. Thus, *Epha4* and *Dlgap1* of cage mates are likely
involved in a common pathway affecting helplessness of focal mice. *Dlgap1* is a core
component of the post-synaptic density and *Dlgap1* knock-out mice show selective
deficits in sociability[35]. Hence, low sociability of cage mates most likely pathway
explaining social effects of *Epha4* and *Dlgap1* on helplessness.

250
251

c

Figure 4 *Locus zoom plots for three genome-wide significant (FDR < 10%) SGE loci.*
In (a), (b), and (c) the top and bottom panels respectively show SGE and DGE of all
variants at the locus. (a) Locus on chromosome 1 for immobility during the first two
minutes of the Porsolt swim test, a measure of helplessness and model of depression
in mice. Epha4 is the only candidate gene (see Methods) for SGE at this locus. (b)
Locus on chromosome 17 for immobility during the last four minutes of the Porsolt
swim test. Candidate genes for SGE include Dlgap1, whose expression is highly
correlated with that of Epha4 in the hippocampus of CFW mice. (c) Locus on
chromosome 12 for a measure of disrupted sleep, where Pgk1-rs7 is the only
candidate gene for SGE.

Among the other genome-wide significant SGE loci with a single candidate
gene was a locus on chromosome 12 for a measure of disrupted sleep (Figure 4c, top
panel). More precisely, after sharing a cage for 15 weeks mice were singly housed for
three days so that their sleep could be recorded; the measure reported here was
defined as the sum of differences in hourly sleep times between days 2 and 3[19]. Thus,

this measure captures unpredictable, disrupted sleep most likely resulting from mice
struggling to adapt to their new housing conditions. The candidate gene at this locus,
Pgk1-rs7, is a pseudogene of phosphoglycerate kinase 1 (*Pgk1*, located on
chromosome X). Lack of linkage disequilibrium (see Methods) between variants in the
vicinity of *Pgk1-rs7* and variants in the vicinity of *Pgk1* suggests our finding is not the
result of sequencing read misalignment during genotyping. The perfect sequence
match between the proteins encoded by *Pgk1-rs7* and *Pgk1,* the position of *Pgk1* on
the X chromosome, and the existence of a functional pseudogene for the human
phosphoglycerate mutase[36] (the enzyme acting on the product of PGK in glycolysis),
together suggest that *Pgk1-rs7* might be a functional autosomal substitute for *Pgk1*, a
process suggested to have enhanced male germline functions in evolution[37]. Neither
Pgk1-rs7 nor *Pgk1* have been associated with sleep and we found no evidence of
DGE on sleep at this locus (Figure 4c, bottom panel), suggesting that other traits of
cage mates are involved. Given that *Pgk1-rs7* is a marker of brown adipose tissue -
the tissue responsible for non-shivering thermogenesis - and that its expression is
activated by cold temperatures[38], our hypothesis regarding the pathways mediating
social effects of *Pgk1-rs7* on sleep is as follows: first, *Pgk1-rs7* controls the heat
produced by a mouse, which in turn determines the need for its cage mates to develop
their own thermogenic capacity[39]. Indeed mice often huddle in the laboratory because
ambient temperature (20° in the initial study) is well below their thermoneutral zone
(30° [40]), and warm cage mates reduce the need for one mouse to produce its own
heat[39]. Secondly, when the mice were separated for the sleep test after several weeks
of sharing a cage, those with low thermogenic capacity likely got awaken by the cold
conditions, resulting in disrupted sleep. Our results illustrate how the results of
sgeGWAS may be used to generate hypotheses as to the traits of social partners
mediating social effects.
295
Architecture of SGE and comparison with that of DGE
Despite being carried out on the same individuals and phenotypes, and in a perfectly
analogous manner, sgeGWAS identified fewer genome-wide significant loci than
dgeGWAS (24 associations for 17 phenotypes and 120 associations for 63
phenotypes respectively). One possible reason for this is dilution of social genetic
effects occurring when the focal mouse can only interact with one cage mate at a time.

We explored this possibility in a power analysis where allelic effects were fixed, three

categories of minor allele frequencies (MAF) were considered (low, medium, or high;

see Methods), and local DGE or SGE were simulated (see Methods). Briefly, we used

two generative models for SGE : an additive model whereby SGE received by a focal

mouse correspond to the sum of the genetic effects projected by its cage mates, and

an average model, reflecting dilution, whereby SGE received by one mouse

correspond to the average of the genetic effects projected by its cage mates. Our

results show that power to detect SGE loci always increases with MAF and is higher

than power to detect DGE when SGE were simulated under the additive model (Figure

5a). When SGE were simulated under the average model, however, power to detect

SGE was lower than power to detect DGE. Thus, smaller effect sizes or dilution of

social effects may have led to the detection of fewer genome-wide significant SGE

associations compared to DGE associations.

Genome-wide significant SGE associations generally explained less

phenotypic variance than genome-wide significant DGE associations: SGE

associations explained a maximum of 2.5% of phenotypic variance, while eleven

genome-wide significant DGE associations explained more than 5% of phenotypic

variance and up to 40% (Figure 5b, Supplementary Table 2 and Supplementary Table

3); average values across all genome-wide significant associations were 1.8% for

SGE and 2.7% for DGE.

Contrastingly, we found that genome-wide significant associations explained a

similar proportion of the corresponding genetic variance for SGE and DGE (Figure 5c).

More precisely, across 5 phenotypes with aggregate contribution of SGE greater than

5% and at least one genome-wide significant SGE association, we found that an

average of 32.5% of the aggregate variance was explained by genome-wide

significant associations. For DGE, that figure was calculated across 55 phenotypes

and was equal to 32.1%. The proportion of aggregate variance explained by genome-

wide significant associations may seem large given the relatively small number of

genome-wide significant associations per phenotype (e.g. compared to humans[41]), but

is consistent with studies of DGE in other outbred laboratory rodent populations[42,43]

and are the result of a relatively small number of variants segregating in the CFW

population and relatively high MAFs[19]. Our quantification of the proportion of genetic

variance explained by genome-wide significant associations suggests SGE and DGE

likely have similar levels of polygenicity. However, given the small number of

phenotypes where this fraction could be calculated, this conclusion will need to be re-evaluated when more SGE associations are discovered in other datasets.

Figure 5 *Power to detect local SGE and DGE, and characterisation of the architecture of SGE and DGE. (a) Power to detect local genetic effects in simulations. Three types of local genetic effects were simulated: DGE, SGE arising from additive genetic effects across cage mates, and SGE arising from average genetic effects across cage mates (see main text and Methods). For each type of effect, results are shown (left to right) for variants with low MAF (MAF < 0.05), medium MAF (0.225<MAF<0.275) and high MAF (MAF>0.45) (MAF: minor allele frequency, defined based on direct genotypes). Power was calculated at a genome-wide significance threshold of negative log P 5. (b) Histogram of the proportion of phenotypic variance explained by individual genome-wide significant SGE (red) and DGE (black) associations (FDR < 10%). (c) Comparison, for each phenotype, of the variance explained by social (red) and direct (black) genetic effects in aggregate (x axis) and the total variance explained jointly by all genome-wide significant associations (y axis). Each dot corresponds to a phenotype with at least one genome-wide significant association. Only those phenotypes with aggregate SGE (or DGE) > 5% were considered to calculate the average values reported in the main text.*

Discussion

In this study we performed the comparative analysis of SGE and DGE acting on a broad range of phenotypes measured in outbred laboratory mice. For each phenotype we applied GWAS as well as variance decomposition techniques to identify individual loci giving rise to SGE in the CFW mouse population and compare the architecture of SGE to that of DGE. Our results demonstrate that sgeGWAS may be used to identify

novel genetic loci that contribute to inter-individual differences, illustrate how such loci
can help infer the traits of social partners that affect the phenotype of interest, and
suggest comparatively larger sample sizes will be necessary required for sgeGWAS
compared to dgeGWAS.

For this study we chose to investigate not only phenotypes expected to be
affected by social partners, such as behaviours and immune traits, but also
phenotypes that were not *a priori* known or suspected to be socially affected, such as
lung response to hypoxia, and muscle and bones characteristics. Indeed, our previous
work, based on two different populations of mice[9], had suggested that SGE could
affect a broad range of behavioural, physiological and morphological phenotypes. In
this study we found that serum LDL cholesterol and alpha-amylase (a marker of
pancreatic function), blood platelets, acute hypoxic response and healing from an ear
punch, in addition to a number of behavioural and immune phenotypes, were
significantly affected by SGE in aggregate (FDR < 10%, Supplementary Table 1).
Thus, our study identified several phenotypes not previously known to be affected by
the social environment and confirmed that SGE can affect a broad range of
phenotypes. While the exact phenotypes affected by SGE in different populations
vary, presumably due to differences in the pool of genetic variants segregating in the
populations and other experimental variables[9], healing from an ear punch shows
remarkable consistency across populations: it is indeed significantly and strongly
affected by SGE, in aggregate, in all three mouse populations we have studied to
date[9]. Unfortunately, there were no genome-wide significant SGE loci for this
phenotype in this study, which prevented identification of the underlying mechanisms.

Our analysis of a broad range of phenotypes, together with testing of genome-
wide variants, was also meant to provide a view of the architecture of SGE that is as
unbiased as possible. We identified far fewer genome-wide significant SGE loci
compared to DGE loci (24 and 120 respectively, Figure 3) and found that SGE loci
tended to explain less phenotypic variation than DGE loci for the same phenotype
(Figure 5b). Our results contrast with findings from two early mapping studies of SGE,
which identified more SGE loci and found greater effect sizes for SGE loci as
compared to DGE loci[15,17]. This discrepancy could be due to differences in the
phenotypes studied, but more likely arises from the fact that one of those studies
tested loci with genome-wide significant DGE only and the other one failed to account
for DGE of the tested variant, thus not controlling the type 1 error rate if direct and

social genotypes were correlated (as we show in Supplementary Figure 2). Notably,

our results are consistent with two other studies that mapped genome-wide SGE and

DGE acting on a small set of traits: one that focused on mouse pup behaviour and

found similar effect sizes for SGE and DGE loci[8], and one that focused on pecking

behaviour in hens and identified DGE loci but no SGE loci[18]. Given the breadth of

phenotypes considered in our study, lower effect sizes is likely to be a general feature

of SGE loci, which implies future sgeGWAS will require larger sample sizes than

dgeGWAS to be equally powered.

Given the effort that we predict will be necessary to carry out well-powered

sgeGWAS, one goal of our study was to empirically evaluate the promise of sgeGWAS

to identify genetic loci not identified by dgeGWAS and provide clues on yet unknown

traits of social partners affecting the phenotype of interest. To do so, we calculated the

correlation ρ between SGE and DGE acting on the same phenotype and tested

whether it was different from $|1|$. We found that for 10 out of 28 phenotypes ρ was

statistically different from $|1|$ (nominal $P < 0.05$, Figure 2 and Supplementary Table 1),

suggesting that different loci and/or different effect sizes give rise to SGE and DGE.

Consistent with this observation, genome-wide significant sgeGWAS and dgeGWAS

loci never overlapped. SGE and DGE loci were largely different even for phenotypes

with prior evidence of "contagion", such as helplessness[21,44], a murine model for

depression. Hence, our results motivate the use of sgeGWAS to discover more traits

of social partners that influence depression and other phenotypes.

We focused on two genome-wide significant loci with a single candidate

gene and no evidence of DGE to illustrate how sgeGWAS may help identify novel

traits of cage mates affecting the phenotype of interest. A locus for helplessness on

chromosome 1 featured *Epha4* as single candidate gene (Figure 4a), and we found

that *Epha4* expression in the hippocampus of CFW mice was highly correlated with

expression of *Dlgap1*, a candidate gene at another genome-wide significant SGE

locus for helplessness in our study (Figure 4b). *Epha4* and *Dlgap1* both point to

neurobehavioral traits of cage mates affecting helplessness of focal mice, and a report

of selective deficits in sociability of *Dlgap1* knock-out mice[35] further indicates

sociability of cage mates likely contributes to social effects on helplessness.

Consistent with this hypothesis, single housing decreases the effectiveness of

antidepressants in Swiss mice[45], which are genetically similar to CFW mice, and social

C57BL/6NHsd show increased anhedonia (another model of depression) when
housed with less social BALB/cOlaHsd cage mates[46]. The other locus we investigated
suggested an association between *Pgk1-rs7*, a gene associated with
thermogenesis[38], and sleep disruption in cage mates (Figure 4c). We hypothesized
that *Pgk1-rs7* controls the heat produced by a mouse, which in turn determines the
need for its cage mates to develop their own thermogenic capacity[39,40] given that mice
huddle to stay warm when sleeping; when the mice were singly housed for the sleep
test, those with low thermogenic capacity got awaken by the cold conditions, resulting
in the observed disrupted sleep. Consistent with this hypothesis, previous studies
have described how increased cage density reduces thermogenic capacity even at
ambient temperature very close to thermoneutrality (28° in Himms-Hagen et al.[47]) and
how sleep quality is impacted by ambient temperatures[48].

One limitation of using SGE to study social effects is that it is blind to traits of
social partners that are mostly non-genetically determined. Additionally, identifying
traits of partners mediating social effect from SGE associations will likely be
challenging, for the same reasons as it is challenging to infer "within-body" pathways
of phenotypic variation from dgeGWAS results[13,49]. Additional data such as expression
data collected in the cage mates (where SGE originate) may help improve the
prioritization of candidate genes at SGE loci and identify the most relevant tissues.

Finally, our study makes several important methodological contributions that
will help perform sgeGWAS, particularly in outbred populations where both DGE and
SGE contribute to phenotypic variation. Specifically, we described the determinants of
power for SGE and highlighted how dilution can reduce power to detect SGE (Figure
5a). We also showed how using the same individuals as both focal individuals and
cage mates in the analysis leads to correlations between direct and social genotypes
at a locus, which need to be accounted for to avoid spurious associations
(Supplementary Note and Supplementary Figure 2). Similar correlations between
direct and social genotypes, but potentially much stronger, may arise for different
reasons in other datasets, notably when focal individuals and social partners are
related, or as a result of direct assortments (e.g. assortative mating[50,51], homophily
between friends[10]). Our analysis approach is robust to spurious associations that arise
in such situations.

Our study vastly improves our understanding of the architecture of SGE and
will enable future sgeGWAS by guiding their design, analysis, and interpretation.
465

Methods

Phenotypes and experimental variables

Phenotypes and experimental variables (covariates) for 1,934 Crl:CFW(SW)-US_P08
(CFW) mice were retrieved from http://wp.cs.ucl.ac.uk/outbredmice/. Phenotypes
were normalized using the boxcox function (MASS package[52]) in R; phenotypes that
could not be normalised satisfactorily (transformation parameter lambda outside of -2
to 2 interval) were excluded. The subset of covariates used for each phenotype is
indicated in Supplementary Table 1. Because data for some phenotypes were missing
for some mice, the sample size varied. The sample size for each phenotype after all
filtering (see below) is indicated in Supplementary Table 1.

Caging information

Mice were four to seven weeks old when they arrived at the phenotyping facility. They
were grouped with their cage mates and then spent nine to twelve weeks undisturbed
in quarantine. They spent a further four weeks together during phenotyping. Males
were housed with males and females with females.

Cage assignments were not included in the publicly available dataset but were
provided by the authors upon request and are now provided in Supplementary Table
4. Cage assignments were recorded at eleven time points throughout the study and
showed that a few mice were taken out of their original cages and singly housed,
presumably because they were too aggressive to their cage mates. When this
happened, we excluded all the mice in that cage from the analysis. We also excluded
cages where some of the mice were "genetically close" (as defined below) to many
other mice. Finally, we only retained cages with exactly three mice per cage. Although
from the sleep test on all mice were singly housed, we still investigated "persistent"
SGE on sleep and tissue phenotypes (persistence over one day for sleep phenotypes
and over a few days for tissue measures).

Genome-wide genotypes

From http://wp.cs.ucl.ac.uk/outbredmice/ we retrieved both allele dosages for 7 million variants and allele dosages for a subset of 353,697 high quality, LD-pruned variants (as described in Nicod et al.[19]). We used high quality, LD-pruned variants for all analyses but the identification of candidate genes at SGE loci (see below), for which we used the full set of variants.

Genetic relatedness matrix (GRM) and exclusion of "genetically close" mice

The genetic relatedness matrix was calculated as the cross-product of the LD-pruned dosage matrix after standardizing the dosages for each variant to mean 0 and variance 1.

We excluded whole cages of mice based on GRM values as follows: we defined a "close pair" of mice as having a GRM value greater than 0.3 (based on the histogram of all GRM values). 199 mice in 145 cages were involved in such close pairs. Excluding all 145 cages would have resulted in excluding 435 mice out of a total of 1,812, which would have led to substantially reduced power for sgeGWAS and dgeGWAS. Thus, we made a compromise and only excluded the 19 cages that were involved in 4 or more close pairs (57 mice excluded).

Variance decomposition

The same method as described in details in Baud et al.[9] was used. Briefly, the model used was:

$$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} \qquad (0)$$

y_f is the phenotypic value of the focal mouse f, X_f is a row of the matrix X of covariate values and b a column vector of corresponding estimated coefficients. $\underline{a_{D,f}}$ is the additive direct genetic effects (DGE) of f. Z_f is a row of the matrix Z that indicates cage mates (importantly $Z_{i,i} = 0$) and $\underline{a_S}$ the column vector of additive social genetic effects (SGE). $\underline{e_D}$ refers to direct environmental effects and $\underline{e_S}$ to social environmental

528   effects. W_f is a row of the matrix W that indicates cage assignment and c the column

529   vector of cage effects.

530   The joint distribution of all random effects is defined as:

532
$$\begin{bmatrix} a_D \\ \hline a_S \\ \hline e_D \\ \hline e_S \\ \hline c \end{bmatrix} \sim \text{MVN} \left(0, \begin{bmatrix} \sigma_{A_D}^2 A & \sigma_{A_{DS}} A & 0 & 0 & 0 \\ \sigma_{A_{DS}} A^T & \sigma_{A_S}^2 A & 0 & 0 & 0 \\ 0 & 0 & \sigma_{E_D}^2 I & \sigma_{E_{DS}} I & 0 \\ 0 & 0 & \sigma_{E_{DS}} I^T & \sigma_{E_S}^2 I & 0 \\ 0 & 0 & 0 & 0 & \sigma_C^2 I \end{bmatrix} \right)$$

534   where A is the GRM and I the identity matrix.

535   The phenotypic covariance is:

536      $C_{i,j} = cov\,(y_i, y_j)$

537               $= \sigma_{A_D}^2 A_{i,j} + \sigma_{A_{DS}} + \sigma_{A_S}^2 (ZAZ^T)_{i,j} + \sigma_{E_D}^2 I_{i,j} + \sigma_{E_{DS}} \{ (IZ^T)_{i,j}$

538                 $+ (ZI^T)_{i,j} \} + \sigma_{E_S}^2 (ZIZ^T)_{i,j} + \sigma_C^2 (WIW^T)_{i,j}$

The variances explained by DGE and SGE were calculated respectively as

$sampleVar(\sigma_{A_D}^2 A) \,/\, sampleVar(C)$ and $sampleVar(\sigma_{A_S}^2 (ZAZ^T)) \,/\, sampleVar(C)$

where $sampleVar$ is the sample variance of the corresponding covariance matrix:

suppose that we have a vector x of random variables with covariance matrix M, the

sample variance of M is calculated as

$sampleVar(M) = \dfrac{Tr(PMP)}{n-1}$

Tr denotes the trace, n is the sample size, and $P = I - \dfrac{11'}{n}$ [53,54].

For those phenotypes where body weight was included as a covariate, we checked

that this did not lead to systematically increased (or decreased) estimates of the

aggregate contribution of SGE (collider bias).

Significance of variance components was assessed using a two-degree of freedom

log likelihood ratio (LLR) test (i.e., the test statistics was assumed to follow a two-

degree of freedom chi2 distribution under the null). Note that this testing procedure is

conservative.

The Q value for the aggregate contribution of SGE was calculated for each phenotype

using the R package qvalue[55]. Significant SGE contributions were reported at FDR <

10% (corresponding to Q value < 0.1).

Correlation between DGE and SGE

The correlation ρ between $\underline{a_D}$ and $\underline{a_S}$ was calculated as:

$$\rho = \frac{\sigma_{A_{DS}}}{\sigma_{A_D} \times \sigma_{A_S}}$$

with ρ reflecting the correlation between SGE and DGE *acting on the same phenotype,*

similarly to how "traditional" genetic correlations measure the correlation between

DGE on two traits. The parameter ρ can actually be interpreted as the correlation

between DGE on the traits of cage mates mediating social effects and DGE on the

phenotype of interest itself.

We tested whether ρ was significantly different from 0 and whether $|\rho|$ was significantly

different from 1 using a one-degree of freedom LLR test.

Simulations for Supplementary Figure 1.

Phenotypes were simulated based on the genotypes and cage relationships of the full

set of 1,812 mice. Phenotypes were drawn from model (0) with the following variances:

$\sigma_{A_D}^2 = 15$, $\sigma_{A_S}^2 = 8$, $\rho_{A_{DS}} = 0.47$, $\sigma_{E_D}^2 = 22$, $\sigma_{E_S}^2 = 16$, $\rho_{E_{DS}} = -0.97$, $\sigma_C^2 = 26$. These variances

correspond to the median value of estimates across traits with aggregate SGE and

DGE > 5%. After building the phenotypic covariance matrix, the sample variance of

the simulations was calculated and used to calculate "realised" simulation parameters

from the "target" parameters above. The realised parameters were used for

comparison with the parameters estimated from the simulations.

Definition of "social genotype" for sgeGWAS

In the sgeGWAS, we assumed additive effects across cage mates and calculated the

"social genotype" of a mouse as the sum of the reference allele dosages of its cage

mates. The same assumptions were made by Biscarini *et al.*[17] and Brinker *et al.*[18].

Correlation between direct and social genotypes at a variant

Spearman's rank correlation coefficient was used. We tested whether the correlation

was different from 0 using the function cor.test in the R package stats[56].

Models used for sgeGWAS and dgeGWAS

To test SGE of a particular variant, we compared the following two models:

$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + G_f b_D$ (1, null)

$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + G_f b_D + Z_f G b_S$ (2, alternative)

Here, G is the vector of direct genotypes at the tested variant, b_D the estimated

coefficient for local DGE and b_S the estimated coefficient for local SGE.

The models were fitted using LIMIX[57,58] with the covariance of the model estimated

only once per phenotype, in the null model with no local genetic effect (model 0).

The significance of local SGE was calculated by comparing models (1) and (2) with a

1-degree of freedom LLR test.

We refer to the inclusion of $G_f b_D$ in model (1, null) as "conditioning".

dgeGWAS was carried out by comparing model (2) above to the null model (3) below:

$y_f = X_f \underline{b} + a_{D,f} + e_{D,f} + Z_f \underline{a_S} + Z_f \underline{e_S} + W_f \underline{c} + Z_f G b_S$ (3, null)

We refer to the inclusion of $Z_f G b_S$ in model (3, null) as "conditioning".

Identification of genome-wide significant associations

Because we wanted to compare the architecture of DGE and SGE *for each phenotype*

independently, we adopted the per-phenotype FDR approach used by Nicod et al.[19].

In contrast to a study-wise FDR, this approach enables the direct comparison of paired

SGEwas and DGEwas for the same traits.

The procedure we used to control the FDR accounts for the fact that we report

loci rather than individual variants[59], where a locus is defined as the 1.5 Mb-wide

window around a SNP (this window size is the average 95% confidence interval for

DGE QTLs in [19]). More precisely, for each phenotype and for each type of genetic

effect (social and direct), we performed 100 "permuted GWAS" by permuting the rows

of the matrix of social (respectively direct) genotypes, and testing each variant at a

time using the permuted genotypes together with the un-permuted phenotypes,

covariates, GRM and matrix of direct (respectively social) genotypes (for conditioning).

See [58,60] for references on this permutation approach. For a given P value x, the per-

phenotype FDR can be calculated calculated as:

$$FDR(x) = \frac{\# \ loci \ with \ P < x \ in \ permuted \ data}{100 \ \times \ \# \ loci \ with \ P < x \ in \ unpermuted \ data}$$

We report only those loci whose P value corresponds to an FDR < 10%.

Definition of candidate genes at associated loci

At each significantly associated locus we defined a 1.5Mb window centred on the lead

variant, identified all the variants that segregate in this window based on the full set of

7M variants, and reran the sgeGWAS and dgeGWAS locally with all the variants at

the locus. We highlighted those genes that are located within the most significantly

associated segments and whose MGI symbol does not start by 'Gm', 'Rik', 'Mir', 'Fam',

or 'Tmem' in order to focus on genes with known function and generate more tractable

hypotheses on the pathways of social effects.

Calculation of linkage disequilibrium between variants in the vicinity of Pgk1-rs7 and

variants in the vicinity of Pgk1

Dosages at LD-pruned variants in the regions chr12:10,091,326-11,591,326 and chrX:

105,500,000- 106,500,000 were used as input for PLINK2 to generate binary PLINK

files, and linkage (R^2) was subsequently calculated between all pairs of variants.

Variance explained by a genome-wide significant association

The variance explained by a genome-wide significant SGE association was estimated in an extension of model (0) with additional fixed effects for both direct and social effects of lead SNPs at all genome-wide significant SGE loci (the lead SNP being the SNP with the most significant P value at the locus in the sgeGWAS). After fitting the model, the variance was calculated as:

$$\frac{var(ZGb_S)}{\sum var(X_c b_c) + \sum var(Gb_D) + \sum var(ZGb_S) + sampleVar(C)}$$

where $sampleVar(C)$ is the sample variance of the covariance matrix in this model.

The variance explained by a genome-wide significant DGE association was estimated in a similar model but considering all genome-wide significant DGE associations and calculated as:

$$\frac{var(Gb_D)}{\sum var(X_c b_c) + \sum var(Gb_D) + \sum var(ZGb_S) + sampleVar(C)}$$

Variance explained jointly by all genome-wide significant SGE or DGE associations for a phenotype

The variance explained jointly by all significant SGE associations was estimated using the same model as above with all genome-wide significant SGE associations and calculated as:

$$\frac{\sum var(ZGb_S)}{\sum var(X_c b_c) + \sum var(Gb_D) + \sum var(ZGb_S) + sampleVar(C)}$$

The variance explained jointly by all significant DGE associations was estimated using the same model as above with all genome-wide significant DGE associations and calculated as:

$$\frac{\sum var(Gb_D)}{\sum var(X_c b_c) + \sum var(Gb_D) + \sum var(ZGb_S) + sampleVar(C)}$$

Simulations for Supplementary Figure 2c and 2d.

Phenotypes were simulated based on the genotypes and cage relationships of the
1,812 mice. Phenotypes were simulated as the sum of random effects and local DGE
(from model (1)), with the following parameters: $\sigma^2_{A_D} = 5$ or 20, $\sigma^2_{A_S} = 5$ or 20, $\rho_{A_{DS}} = 0.5$,
$\sigma^2_{E_D} = 30$, $\sigma^2_{E_S} = 30$, $\rho_{E_{DS}} = -0.97$, $\sigma^2_C = 25$. The values for $\rho_{A_{DS}}$, $\sigma^2_{E_D}$, $\sigma^2_{E_S}$, $\rho_{E_{DS}}$, and σ^2_C
were close to the median of the corresponding estimates from the real data. $\sigma^2_{A_D} = 5$
and $\sigma^2_{A_S} = 5$ correspond to low polygenic effects in the real data, and $\sigma^2_{A_D} = 20$ and
$\sigma^2_{A_S} = 20$ correspond to high polygenic effects in the real data. We simulated local
DGE at random variants in the genome, and simulated variances of 0, 5, 20 or 50.

The results we show in Supplementary Figure 2c and 2d are based on a subset
of simulations: $\sigma^2_{A_D} = 20$ and $\sigma^2_{A_S} = 20$ and local DGE variance of 20.

Simulations for Supplementary Figure 3a-d, and Figure 3a.

Phenotypes were simulated based on the real genotypes but random cages.
Phenotypes were simulated as the sum of random effects, local DGE and local SGE
(model (2)) with the following parameters: $\sigma^2_{A_D} = 17$, $\sigma^2_{A_S} = 17$, $\rho_{A_{DS}} = 0.65$, $\sigma^2_{E_D} = 19$,
$\sigma^2_{E_S} = 15$, $\rho_{E_{DS}} = -0.8$, $\sigma^2_C = 25$. Those values correspond to the median estimates for
phenotypes with aggregate SGE and DGE > 0.1.

We simulated local SGE and DGE at variants where direct and social
genotypes were either lowly correlated (Spearman correlation negative log P value <
0.05) or more highly correlated (Spearman correlation negative log P value > 0.2), and
had with low MAF (MAF < 0.05), medium MAF (0.225<MAF<0.275) or high MAF
(MAF>0.45). We simulated local DGE with an allelic effect of 0 or 1 (1 corresponds to
a large effect in the real data). We simulated local SGE under two alternative
generative models: an "additive" model by using Z as in model (2) (i.e. filled with 0s
and 1s) or a "proportional" model by using $Z' = {}^Z/_N$. In all cases we simulated an
allelic effect of 0.2 (similar to the average allelic effect estimated in the SGE GWAS).
The sample variance of the simulated local DGE term is $var(Gb_D) = 2p(1-p)\,b^2_D$; it
is $var(ZGb_S) = 2\mathrm{N}p(1-p)\,b^2_S$ for the local SGE term simulated under the additive

model, and $var\left(\frac{Z}{N}Gb_S\right) = 2Np(1-p)/N^2\,b_S^2$ for the local SGE component simulated

under the proportional model.

The results we show in Supplementary Figure 3a-d are based on simulations

averaged across low, medium and high MAF. Power was calculated at a genome-wide

significance threshold of negative log P 5, which is similar to the significance of

associations detected at FDR < 10%.

The results we show in Figure 3a are based on a subset of simulations with no

local DGE, averaged across high and low genotypic correlations. Power was also

calculated at a genome-wide significance threshold of negative log P 5.

Scripts used in this study

All the scripts used in this study are available from http://github.com/limix/SGE.

LIMIX can be downloaded from http://github.com/limix/limix.

Acknowledgements

731 AB was supported by a fellowship from the Wellcome Trust (105941/Z/14/Z).

References

1. Griffing, B. Selection in reference to biological groups I. Individual and group selection applied to populations of unordered groups. *Australian Journal of Biological Sciences* **20**, 127–140 (1967).

2. Moore, A.J., Brodie III, E.D. & Wolf, J.B. Interacting phenotypes and the evolutionary process: I. Direct and indirect genetic effects of social interactions. *Evolution* **51**, 1352–1362 (1997).

3. Wolf, J.B., Brodie III, E.D., Cheverud, J.M., Moore, A.J. & Wade, M.J. Evolutionary consequences of indirect genetic effects. *Trends in ecology & evolution* **13**, 64–69 (1998).

4. Bergsma, R., Kanis, E., Knol, E.F. & Bijma, P. The contribution of social effects to heritable variation in finishing traits of domestic pigs (Sus scrofa). *Genetics* **178**, 1559–1570 (2008).

5. Bijma, P. The quantitative genetics of indirect genetic effects: a selective review of modelling issues. *Heredity* **112**, 61 (2014).

6. Wilson, A.J., Gelin, U., Perron, M.-C. & Réale, D. Indirect genetic effects and the evolution of aggression in a vertebrate system. *Proceedings of the Royal Society of London B: Biological Sciences* **276**, 533–541 (2009).

7. Ashbrook, D.G., Gini, B. & Hager, R. Genetic variation in offspring indirectly influences the quality of maternal behaviour in mice. *Elife* **4**, 11814 (2015).

8. Ashbrook, D.G., Sharmin, N. & Hager, R. Offspring genes indirectly influence sibling and maternal behavioural strategies over resource share. in *Proc. R. Soc. B* Vol. 284 20171059 (The Royal Society, 2017).

9. Baud, A. *et al.* Genetic variation in the social environment contributes to health and disease. *PLoS genetics* **13**, e1006498 (2017).

10. Domingue, B.W. *et al.* The social genome of friends and schoolmates in the National Longitudinal Study of Adolescent to Adult Health. *Proceedings of the National Academy of Sciences*, 201711803 (2018).

11. Kong, A. *et al.* The nature of nurture: Effects of parental genotypes. *Science* **359**, 424–428 (2018).

12. Bates, T.C. *et al.* The Nature of Nurture: Using a Virtual-Parent Design to Test Parenting Effects on Childrens Educational Attainment in Genotyped Families. *Twin Research and Human Genetics*, 1–11 (2018).

13. Visscher, P.M. *et al.* 10 years of GWAS discovery: biology, function, and translation. *The American Journal of Human Genetics* **101**, 5–22 (2017).

14. Timpson, N. J., Greenwood, C. M., Soranzo, N., Lawson, D. J. & Richards, J. B. Genetic
architecture: the shape of the genetic contribution to human traits and disease.
Nature Reviews Genetics **19**, 110 (2018).

15. Mutic, J. J. & Wolf, J. B. Indirect genetic effects from ecological interactions in
Arabidopsis thaliana. *Molecular ecology* **16**, 2371–2381 (2007).

16. Bailey, N. W. & Hoskins, J. L. Detecting cryptic indirect genetic effects. *Evolution* **68**,
1871–1882 (2014).

17. Biscarini, F. *et al.* Across-line SNP association study for direct and associative
effects on feather damage in laying hens. *Behavior genetics* **40**, 715–727 (2010).

18. Brinker, T., Bijma, P., Vereijken, A. & Ellen, E. D. The genetic architecture of
socially-affected traits: a GWAS for direct and indirect genetic effects on survival
time in laying hens showing cannibalism. *Genetics Selection Evolution* **50**, 38
(2018).

19. Nicod, J. *et al.* Genome-wide association of multiple complex traits in outbred mice
by ultra-low-coverage sequencing. *Nature genetics* **48**, 912 (2016).

20. Davies, R. W., Flint, J., Myers, S. & Mott, R. Rapid genotype imputation from
sequence without reference panels. *Nature genetics* **48**, 965 (2016).

21. Haeffel, G. J. & Hames, J. L. Cognitive Vulnerability to Depression Can Be
Contagious. *Clinical Psychological Science* **2**, 75–85 (2014).

22. Rosenquist, J. N., Murabito, J., Fowler, J. H. & Christakis, N. A. The spread of alcohol
consumption behavior in a large social network. *Annals of internal medicine* **152**,
426–433 (2010).

23. Grant, J. D. *et al.* Spousal concordance for alcohol dependence: evidence for
assortative mating or spousal interaction effects? *Alcoholism: Clinical and
Experimental Research* **31**, 717–728 (2007).

24. Bolger, N., DeLongis, A., Kessler, R. C. & Wethington, E. The contagion of stress
across multiple roles. *Journal of Marriage and the Family*, 175–183 (1989).

25. Christakis, N. A. & Fowler, J. H. The spread of obesity in a large social network over
32 years. *New England journal of medicine* **357**, 370–379 (2007).

26. Cohen-Cole, E. & Fletcher, J. M. Detecting implausible social network effects in
acne, height, and headaches: longitudinal analysis. *Bmj* **337**, a2533 (2008).

27. Yalcin, B. *et al.* Commercially available outbred mice for genome-wide association
studies. *PLoS genetics* **6**, e1001085 (2010).

28. Costa, A. P. R. *et al.* A proposal for refining the forced swim test in Swiss mice.
Progress in Neuro-Psychopharmacology and Biological Psychiatry **45**, 150–155
(2013).

29. Yamaguchi, Y. & Pasquale, E. B. Eph receptors in the adult brain. *Current opinion in
neurobiology* **14**, 288–296 (2004).

30.   Smith, C. L. *et al.* Mouse Genome Database (MGD) 2018: knowledgebase for the
laboratory mouse. *Nucleic acids research* **46**, D836–D842 (2017).

31.   Li, Y. *et al.* Differential expression of hippocampal EphA4 and ephrinA3 in
anhedonic-like behavior, stress resilience, and antidepressant drug treatment
after chronic unpredicted mild stress. *Neuroscience letters* **566**, 292–297 (2014).

32.   Zhang, J.-c. *et al.* Increased EphA4-ephexin1 signaling in the medial prefrontal
cortex plays a role in depression-like phenotype. *Scientific Reports* **7**, 7133
(2017).

33.   Parker, C. C. *et al.* Genome-wide association study of behavioral, physiological and
gene expression traits in outbred CFW mice. *Nature genetics* **48**, 919 (2016).

34.   Sloan, Z. *et al.* GeneNetwork: framework for web-based genetics. *The Journal of*
Open Source Software **1**, – (2016).

35.   Coba, M. *et al.* Dlgap1 knockout mice exhibit alterations of the postsynaptic
density and selective reductions in sociability. *Scientific reports* **8**, 2281 (2018).

36.   Betrán, E., Wang, W., Jin, L. & Long, M. Evolution of the phosphoglycerate mutase
processed gene in human and chimpanzee revealing the origin of a new primate
gene. *Molecular biology and evolution* **19**, 654–663 (2002).

37.   Marques, A. C., Dupanloup, I., Vinckenbosch, N., Reymond, A. & Kaessmann, H.
Emergence of young human genes after a burst of retroposition in primates. *PLoS*
biology **3**, e357 (2005).

38.   Cheng, Y. *et al.* Prediction of Adipose Browning Capacity by Systematic Integration
of Transcriptional Profiles. *Cell reports* **23**, 3112–3125 (2018).

39.   Sokoloff, G. & Blumberg, M. S. Competition and cooperation among huddling infant
rats. *Developmental Psychobiology: The Journal of the International Society for*
Developmental Psychobiology **39**, 65–75 (2001).

40.   Gordon, C. Thermal physiology of laboratory mice: defining thermoneutrality.
Journal of thermal biology **37**, 654–685 (2012).

41.   Manolio, T. A. *et al.* Finding the missing heritability of complex diseases. *Nature*
461, 747 (2009).

42.   Valdar, W. *et al.* Genome-wide genetic association of complex traits in
heterogeneous stock mice. *Nature genetics* **38**, 879 (2006).

43.   Sequencing, R. G. *et al.* Combined sequence-based and genetic mapping analysis
of complex traits in outbred rats. *Nature genetics* **45**, 767 (2013).

44.   Bastiampillai, T., Allison, S. & Chan, S. Is depression contagious? The importance
of social networks and the implications of contagion theory. *Australian & New*
Zealand Journal of Psychiatry **47**, 299–303 (2013).

45.　Karolewicz, B. & Paul, I.A. Group housing of mice increases immobility and
antidepressant sensitivity in the forced swim and tail suspension tests. *European*
journal of pharmacology **415**, 197–201 (2001).

46.　Heinla, I., Åhlgren, J., Vasar, E. & Voikar, V. Behavioural characterization of
C57BL/6N and BALB/c female mice in social home cage–Effect of mixed housing
in complex environment. *Physiology & behavior* **188**, 32–41 (2018).

47.　Himms–Hagen, J. & Villemure, C. Number of mice per cage influences uncoupling
protein content of brown adipose tissue. *Proceedings of the Society for*
Experimental Biology and Medicine **200**, 502–506 (1992).

48.　Martire, V.L., Silvani, A., Bastianini, S., Berteotti, C. & Zoccoli, G. Effects of ambient
temperature on sleep and cardiovascular regulation in mice: the role of
hypocretin/orexin neurons. *PLoS One* **7**, e47032 (2012).

49.　Visscher, P.M., Brown, M.A., McCarthy, M.I. & Yang, J. Five years of GWAS
discovery. *The American Journal of Human Genetics* **90**, 7–24 (2012).

50.　Domingue, B.W., Fletcher, J., Conley, D. & Boardman, J.D. Genetic and educational
assortative mating among US adults. *Proceedings of the National Academy of*
Sciences **111**, 7996–8000 (2014).

51.　Robinson, M.R. *et al.* Genetic evidence of assortative mating in humans. *Nature*
Human Behaviour **1**, 0016 (2017).

52.　Venerables, W. & Ripley, B. Modern applied statistics with S. (new york: Springer,
2002).

53.　Kang, H.M. *et al.* Variance component model to account for sample structure in
genome–wide association studies. *Nature genetics* **42**, 348–354 (2010).

54.　Searle, S.R. *Matrix algebra useful for statistics*, (Wiley, 1982).

55.　Dabney, A., Storey, J.D. & Warnes, G. qvalue: Q–value estimation for false
discovery rate control. *R package version* **1** (2010).

56.　Best, D. & Roberts, D. Algorithm AS 89: the upper tail probabilities of Spearmans
rho. *Journal of the Royal Statistical Society. Series C (Applied Statistics)* **24**, 377–
379 (1975).

57.　Lippert, C., Casale, F., Rakitsch, B. & Stegle, O. LIMIX: genetic analysis of multiple
traits. (2014).

58.　Casale, F.P., Rakitsch, B., Lippert, C. & Stegle, O. Efficient set tests for the genetic
analysis of correlated traits. *Nature methods* **12**, 755 (2015).

59.　Brzyski, D. *et al.* Controlling the rate of GWAS false discoveries. *Genetics* **205**, 61–
75 (2017).

60.　Sudmant, P.H. *et al.* An integrated map of structural variation in 2,504 human
genomes. *Nature* **526**, 75 (2015).