Representation of auditory motion directions and sound source locations in the human planum temporale

Ceren Battal [*,1,2], Mohamed Rezk[1,2], Stefania Mattioni[1,2], Jyothirmayi Vadlamudi[2] & Olivier Collignon [*,1,2]

[1]Center of Mind/Brain Sciences, University of Trento, Italy

[2]Institut de recherche en Psychologie (IPSY), Institute of Neuroscience (IoNS), Université catholique de Louvain (UcL)

*Corresponding authors: Ceren Battal; ceren.battal@uclouvain.be; Olivier Collignon; Institut de recherche en Psychologie (IPSY), 10, place du cardinal Mercier, 1348 Louvain-la-Neuve ; olivier.collignon@uclouvain.be.

Key words: spatial hearing, planum temporale, auditory motion, auditory space, direction selectivity, fMRI, multivariate analyses.

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: We would also like to express our gratitude to Giorgia Bertonati, Marco Barilari, Stefania Benetti, Valeria Occelli, Stephanie Cattoir who helped with the data acquisition; Jorge Jovicich for helping setting-up the fMRI acquisition parameters and Pietro Chiesa for continuing support with the auditory hardware. The project was funded by the ERC starting grant MADVIS (Project: 337573) awarded to OC. MR and JV are PhD fellows and OC is research associate at the Fond National de la Recherche Scientifique de Belgique (FRS-FNRS).

ABSTRACT

The ability to compute the location and direction of sounds is a crucial perceptual skill to efficiently interact with dynamic environments. How the human brain implements spatial hearing is however poorly understood. In our study, we used fMRI to characterize the brain activity of male and female humans listening to left, right, up and down moving as well as static sounds. Whole brain univariate results contrasting moving and static sounds varying in their location revealed a robust functional preference for auditory motion in bilateral human Planum Temporale (hPT). Using independently localized hPT, we show that this region contains information about auditory motion directions and, to a lesser extent, sound source locations. Moreover, hPT showed an axis of motion organization reminiscent of the functional organization of the middle-temporal cortex (hMT+/V5) for vision. Importantly, whereas motion direction and location rely on partially shared pattern geometries in hPT, as demonstrated by successful cross-condition decoding, the responses elicited by static and moving sounds were however significantly distinct. Altogether our results demonstrate that the hPT codes for auditory motion and location but that the underlying neural computation linked to motion processing is more reliable and partially distinct from the one supporting sound source location.

SIGNIFICANCE STATEMENT

In comparison to what we know about visual motion, little is known about how the brain implements spatial hearing. Our study reveals that motion directions and sound source locations can be reliably decoded in the human Planum Temporale

(hPT) and that they rely on partially shared pattern geometries. Our study therefore

sheds important new lights on how computing the location or direction of sounds are

implemented in the human auditory cortex by showing that those two computations

rely on partially shared neural codes. Furthermore, our results show that the neural

representation of moving sounds in hPT follows a "preferred axis of motion"

organization, reminiscent of the coding mechanisms typically observed in the occipital

hMT+/V5 region for computing visual motion.

INTRODUCTION

While the brain mechanisms underlying the processing of visual localization and visual motion have received considerable attention (Braddick et al., 2001; Movshon and Newsome, 1996; Newsome and Park, 1988), much less is known about how the brain implements spatial hearing. The representation of auditory space relies on the computations and comparison of intensity, temporal and spectral cues that arise at each ear (Blauert, 1982; Searle et al., 1976). In the auditory pathway, these cues are both processed and integrated in the brainstem, thalamus, and cortex in order to create an integrated neural representation of auditory space (Boudreau and Tsuchitani, 1968; Goldberg and Brown, 1969; Knudsen & Konishi 1978; Ingham et al., 2001; see Grothe et al., 2010 for review).

Similar to the dual-stream processing model in vision (Ungerleider and Mishkin, 1982; Goodale and Milner, 1992), partially distinct ventral "what" and dorsal "where" auditory processing streams have been proposed for auditory processing (Barrett and Hall, 2006; Lomber and Malhotra, 2008; Rauschecker and Tian, 2000; Recanzone, 2000; Romanski et al., 1999; Tian et al., 2001; Warren and Griffiths, 2003). In particular, the dorsal stream is thought to process sound source location and motion both in animals and humans (Arnott et al., 2004; Alain et al., 2001; Maeder et al., 2001; Rauschecker and Tian 2000; Tian et al., 2001). However, it remains poorly understood whether the human brain implements the processing of auditory motion and location using distinct, similar, or partially shared neural substrates (Poirier et al., 2017; Smith et al., 2010; Zatorre et al., 2002).

One candidate region that might integrate spatial cues to compute motion

and location information is the planum temporale (hPT) (Barrett and Hall, 2006;

Baumgart and Gaschler-Markefski, 1999; Warren et al., 2002). hPT is located in the

superior temporal gyrus, posterior to Helsch' gyrus, and is typically considered part

of the dorsal auditory stream (Derey et al., 2016; Poirier et al., 2017; Rauschecker and

Tian, 2000; Warren et al., 2002). Some authors have suggested that hPT engages in

both the processing of moving sounds and the location of static sound-sources

(Barrett and Hall, 2006; Derey et al., 2016; Krumbholz et al., 2005; Smith et al., 2004,

2007, 2010; Zatorre et al., 2002). This proposition is supported by early animal

electrophysiological studies demonstrating the neurons in the auditory cortex that are

selective to sound source location and motion directions (Altman, 1968, 1994; Benson

et al., 1981; Doan et al., 1999; Imig et al., 1990; Middlebrooks and Pettigrew, 1981;

Poirier et al., 1997; Rajan et al., 1990). In contrast, other studies in animals (Poirier et

al., 2017) and humans (Baumgart and Gaschler-Markefski, 1999; Bremmer et al., 2001;

Hall and Moore, 2003; Krumbholz et al., 2005; Lewis et al., 2000; Pavani et al., 2002;

Poirier et al., 2005) pointed toward a more specific role of hPT for auditory motion

processing.

The hPT, selective to process auditory motion/location in a dorsal "where"

pathway is reminiscent of the dorsal region in the middle temporal cortex (hMT+/V5)

in the visual system. hMT+/V5 is dedicated to process visual motion (Movshon and

Newsome, 1996; Watson et al., 1993; Tootell et al., 1995) and displays a columnar

organization in particular of a preferred axis of motion directions (Albright et al., 1984;

Zimmermann et al., 2011). Whether hPT disclose similar characteristic tuning remains

however unknown.

The main goals of the present study were threefold. First, using MVPA, we

investigated whether information about auditory motion direction and sound-source

location can be retrieved from the pattern of activity in hPT. Further, we asked whether

the spatial distribution of the neural representation is in the format of "preferred axis

of motion/location" as observed in the visual motion selective regions (Albright et al.,

1984; Zimmermann et al., 2011). Finally, we aimed at characterizing whether the

processing of motion direction (e.g. going to the left) and sound-source location (e.g.

being in the left) rely on partially common neural representations in the hPT.

MATERIALS AND METHODS

Participants

Eighteen participants with no reported auditory problems were recruited for the

study. Two participants were excluded due to poor spatial hearing performance in the

task, as it was lower by more than 2.5 standard deviations than the average of the

participants. The final sample included 16 right-handed participants (8 females, age

range: 20 to 42, mean ± SD = 32 ± 5.7 years). Participants were blindfolded and

instructed to keep their eyes closed throughout the experiments and practice runs.

All the procedures were approved by the research ethics boards of the Centre for

Mind/Brain Sciences (CIMeC) and University of Trento. Experiments were undertaken

with the understanding and written consent of each participant.

Auditory stimuli

Our limited knowledge of the auditory space processing in the human brain might be a consequence of the technical challenge of evoking vivid perceptual experience of auditory space while using neuroimaging tools such as fMRI, EEG, or MEG. In this experiment, to create an externalized ecological sensation of sound location and motion inside the MRI scanner, we relied on individual in-ear stereo recordings that were recorded in a semi-anechoic room and from 30 loudspeakers on horizontal and vertical planes, mounted on two semi-circular wooden structures with a radius of 1.1m (see Figure 1A). Participants were seated in the center of the apparatus with their head on a chin-rest, such that the speakers on the horizontal and vertical planes were equally distant from participants' ears. Then, these recordings were re-played to the participants when they were inside the MRI scanner. By using such sound system with in-ear recordings, auditory stimuli automatically convolved with each individuals' own pinna and head related transfer function to produce a salient auditory perception in external space.

--- Figure 1 about here ---

The auditory stimuli were prepared using custom MATLAB scripts (r2013b; Matworks). Auditory stimuli were recorded using binaural in-ear omni-directional microphones (Sound Professionals-TFB-2; 'flat' frequency range 20–20,000 Hz) connected to a portable Zoom H4n digital wave recorder (16-bit, stereo, 44.1 kHz sampling rate). Microphones were positioned at the opening of participant's left and right auditory ear canals. While auditory stimuli were played, participants were listening without

performing any task with their head fixed to the chin-rest in front of them. Binaural in-

ear recordings allowed combining binaural properties such as interaural time and

intensity differences, and participant specific monaural filtering cues to create reliable

and ecological auditory space sensation (Pavani et al., 2002).

Stimuli recordings

Sound stimuli consisted of 1250 ms pink noise (50 ms rise/fall time). In the

motion condition, the pink noise was presented moving in 4 directions: leftward,

rightward, upward and downward. Moving stimuli covered 120° of space/visual field

in horizontal and vertical axes. To create the perception of smooth motion, the 1250

ms of pink noise was fragmented into 15 equal length pieces with each 83.333 ms

fragment being played every two speakers, and moved one speaker at a time, from

outer left to outer right (rightward motion), or vice-versa for the leftward motion. For

example, for the rightward sweep, sound was played through speakers located at -

60° and -52° consecutively, followed by -44°, and so on. A similar design was used for

the vertical axis. This resulted in participants perceiving moving sweeps covering an

arc of 120° in 1250 ms (speed = 96°/s; 50 ms fade in/out) containing the same sounds

for all four directions. The choice of the movement speed of the motion stimuli aimed

to create listening experience relevant to everyday-life conditions. Moreover, at such

velocity it has been demonstrated that human listeners are not able to make the

differences between concatenated static stimuli from motion stimuli elicited by a

single moving object (Poirier et al., 2017), supporting the participant's report that our

stimuli were perceived as smoothly moving (no perception of successive snapshots).

In the static condition, the same pink noise was presented separately at one of 4

locations: left, right, up, and down. Static sounds were presented at the second most

outer speakers (-56° and +56° in the horizontal axis, and +56° and -56° in the vertical

axis) in order to avoid possible reverberation difference at the outermost speakers.

The static sounds were fixed at one location at a time instead of presented in multiple

locations (Smith et al., 2008; 2010; Poirier et al., 2017; Krumbholz et al., 2005). This

strategy was purposely adopted for two main reasons. First, randomly presented static

sounds can evoke robust sensation of auditory apparent motion (Strybel & Neale

1994; Lakatos et al., 1997; see Carlile 2016 for review). Second, and crucially for the

purpose of the present experiment, presenting static sounds located in a given

position and moving sounds directed toward the same position allowed us to

investigate whether moving and static sounds share a common representational

format using cross-condition decoding (see below), which would have been

impossible if the static sounds where randomly moving in space.

Before the recordings, the sound pressure levels (SPL) were measured from

the participant's head position and ensured that each speaker conveys 65dB-A SPL.

All participants reported strong sensation of auditory motion and were able to detect

locations with high accuracy (see Fig 1C). Throughout the experiment, participants

were blindfolded. Stimuli recordings were conducted in a session that lasted

approximately 10 minutes, requiring the participant to remain still during this period.

Auditory experiment

Auditory stimuli were presented via MR-compatible closed-ear headphones (Serene

Sound, Resonance Technology; 500-10KHz frequency response) that provided

average ambient noise cancellation of about 30 dB-A. Sound amplitude was adjusted

according to each participant's comfort level. To familiarize the participants with the

task, they completed a practice session outside of the scanner until they reached

above 80% accuracy.

Each run consisted of the 8 auditory categories (4 motion and 4 static)

randomly presented using a block-design. Each category of sound was presented for

15 s (12 repetition of 1250 ms sound, no ISI) and followed by 7 s gap for indicating

the corresponding direction/location in space and 8 s of silence (total inter-block

interval was 15 s). The ramp (50 ms fade in/out) applied at the beginning and at the

end of each sound creates static bursts and minimized adaptation to the static sounds.

During the response gap, participants heard a voice saying "left", "right", "up", and

"down" in pseudo-randomized order. Participants were asked to press a button with

their right index finger when the auditory block's direction or location was matching

with the auditory cue (Figure 1B). The number of targets and the order (position 1-4)

of the correct button press were balanced across category. This procedure was

adopted to ensure that the participants gave their response using equal motor

command for each category and to ensure the response is produced after the end of

the stimulation period for each category. Participants were instructed to emphasize

accuracy of response but not reaction times.

Each scan consisted of one block of each category, resulting in a total of 8

blocks per run, with each run lasting 4 m 10 s. Participants completed a total of 12

runs. The order of the blocks was randomized within each run, and across participants.

Based on pilot experiments, we decided to not rely on a sparse-sampling

design as sometimes done in the auditory literature in order to present the sounds

without the scanner background noise (Hall et al., 1999). These pilot experiments

showed that the increase in the signal to noise ratio potentially provided by sparse

sampling did not compensate for the loss in the number of volume acquisitions.

Indeed, pilot recordings on participants not included in the current sample showed

that, given a similar acquisition time between sparse-sampling designs (several

options tested) and continuous acquisition, the activity maps elicited by our spatial

sounds contained higher and more reliable beta values using continuous acquisition.

fMRI data acquisition and analyses

Imaging parameters

Functional and structural data were acquired with a 4T Bruker MedSpec

Biospin MR scanner, equipped with an 8-channel head coil. Functional images were

acquired with T2*-weighted gradient echo-planar sequence. Acquisition parameters

were: repetition time of 2500 ms, echo time of 26 ms, flip angle of 73°, a field of view

of 192 mm, a matrix size of 64 x 64, and voxel size of 3 x 3 x 3 mm^3. A total of 39

slices were acquired in ascending feet-to-head interleaved order with no gap. The

three initial scans of each acquisition run were discarded to allow for steady-state

magnetization. Before every two EPI runs, we performed an additional scan to

measure the point-spread function (PSF) of the acquired sequence, including fat

saturation, which served for distortion correction that is expected with high-field

imaging (Zeng and Constable, 2002).

High-resolution anatomical scan was acquired for each participant using a T1-

weighted 3D MP-RAGE sequence (176 sagittal slices, voxel size of 1×1×1 mm^3; field

of view 256 x 224 mm; repetition time = 2700 ms; TE = 4.18 ms; FA: 7°; inversion

time: 1020 ms). Participants were blindfolded and instructed to lie still during

acquisition and foam padding was used to minimize scanner noise and head

movement.

Univariate fMRI analysis

Whole brain

Raw functional images were pre-processed and analysed with SPM8 (Welcome

Trust   Centre   for   Neuroimaging   London,   UK;

http://www.fil.ion.ucl.ac.uk/spm/software/spm/) implemented in MATLAB R2014b

(MathWorks). Before the statistical analysis, our preprocessing steps included slice

time correction with reference to the middle temporal slice, realignment of functional

time series, the coregistration of functional and anatomical data, spatial normalization

to an echo planar imaging template conforming to the Montreal Neurological Institute

space, and spatial smoothing (Gaussian kernel, 6 mm FWHM) were performed.

To obtain blood oxygen level-dependent (BOLD) activity related to auditory

spatial processing, we computed single subject statistical comparisons with fixed-

effect general linear model (GLM). In the GLM, we used eight regressors from each

category (four motion direction, four sound source location). The canonical double-

gamma hemodynamic response function implemented in SPM8 was convolved with

a box-car function to model the above mentioned regressors. Motion parameters

derived from realignment of the functional volumes (3 translational motion and 3

rotational motion parameters), button press, and the four auditory response cue

events were modeled as regressors of no interest. During the model estimation, the

data were high-pass filtered with cut-off 128s to remove the slow drifts/ low-frequency

fluctuations from the time series. To account for serial correlation due to noise in fMRI

signal, autoregressive (AR (1)) was used.

In order to obtain activity related to auditory processing in the whole brain,

the contrasts tested the main effect of each category ([Left Motion], [Right Motion],

[Up Motion], [Down Motion], [Left Static], [Right Static], [Up Static], [Down Static]). To

find brain regions responding preferentially to the moving and static sounds, we

combined all motion conditions [Motion] and all static categories [Static]. The

contrasts tested the main effect of each condition ([Motion], [Static]), and comparison

between the conditions ([Motion > Static], and [Static > Motion]). These linear

contrasts generated statistical parametric maps (SPM[T]) which were further spatially

smoothed (Gaussian kernel 8 mm FWHM) and entered in a second-level analysis,

corresponding to a random effects model, accounting for inter-subject variance. One-

sample t-tests were run to characterize the main effect of each condition ([Motion],

[Static]), and the main effect of motion processing ([Motion > Static]) and static

location processing ([Static > Motion]). Statistical inferences were performed at a

threshold of $p<0.05$ corrected for multiple comparisons (Family-Wise Error corrected;

FWE) either over the entire brain volume or after correction for multiple comparisons

over small spherical volumes (12 mm radius) located in regions of interest. Significant

clusters were anatomically labeled using the xjView Matlab toolbox

(http://www.alivelearn.net/xjview) or structural neuroanatomy information provided in

the Anatomy Toolbox (Eickhoff et al., 2007).

Region of interest analysis

ROI Definition

Due to the hypothesis-driven nature of our study we defined hPT as an *a priori*

region of interest for statistical comparisons and in order to define the volume in which

we performed multivariate pattern classification analyses.

To avoid any form of double dipping that may arise when defining the ROI

based on our own data, we decided to independently define hPT, using a meta-

analysis method of quantitative association test, implemented via the online tool

Neurosynth (Yarkoni et al., 2011) using the term query "Planum Temporale". Rather

than showing which regions are disproportionately reported by studies where a

certain term is dominant (uniformity test; P (activation | term)), this method identifies

regions whose report in a neuroimaging study is diagnostic of a certain term being

dominant in the study (association test; P (term | activation)). As such, the definition of

this ROI was based on a set of 85 neuroimaging studies at the moment of the query

(September 2017). This method provides an independent method to obtain masks for

further region-of-interest analysis. The peak coordinate from the meta-analysis map

was used to create a 6 mm spheres (117 voxels) around the peak z-values of hPT (peak

MNI coordinates [-56 -28 8] and [60 -28 8]; lhPT and rhPT hereafter, respectively).

307   Additionally, hPT regions of interest were individually defined using

anatomical parcellation with FreeSurfer (http://surfer.nmr.mgh.harvard.edu). The

individual anatomical scan was used to perform cortical anatomical segmentation

according to the Destrieux (Destrieux et al., 2010) atlas. We selected planum

temporale label defined as [G_temp_sup-Plan_tempo, 36] bilaterally. We equated the

size of ROI across participants and across hemispheres to 110 voxels (each voxel being

3mm isotropic). For anatomically defined hPT ROIs, all further analyses were carried

out in subject space for enhanced anatomical precision and to avoid spatial

normalization across participants. We replicated our pattern of results in anatomically

defined parcels (left and right hPT) obtained from the single subject brain

segmentation (for further analysis see Battal et al., 2018).

ROI Analyses

Univariate

The beta parameter estimates of the 4 motion directions and 4 sound source locations

were extracted from lhPT and rhPT regions (Fig 2C). In order to investigate the

presence of motion directions/sound source locations selectivity and condition effect

in hPT regions, we performed a 2 Conditions (hereafter refers to the stimuli type:

motion, static) x 4 Orientations (hereafter refers to either direction or location of the

stimuli: left, right, down, and up) repeated measures ANOVA in each hemisphere

separately on these beta parameter estimates. Statistical results were then corrected

for multiple comparisons (number of ROIs x number of tests) using the false discovery

rate (FDR) method (Benjamini and Yekutieli, 2001). A Greenhouse–Geisser correction

was applied to the degrees of freedom and significance levels whenever an

assumption of sphericity was violated.

ROI - Multivariate pattern analyses

Within Condition Classification

Four-class and binary classification analyses were conducted within the hPT

region in order to investigate the presence of auditory motion direction and sound

source location information in this area. To perform multi-voxel pattern classification,

we used a univariate one-way ANOVA to select a subset of voxels (n=110) that are

showing the most significant signal variation between the categories of stimuli (in our

study, between orientations). This feature-selection not only ensures a similar number

of voxels within a given region across participants (dimensionality reduction), but,

more importantly, identifies and selects voxels that are carrying the most relevant

information across categories of stimuli (Cox and Savoy 2003; De Martino et al., 2008),

therefore minimizing the chance to include in our analyses voxels carrying noises

unrelated to our categories of stimuli.

Multivariate pattern analyses (MVPA) were performed in the lhPT and rhPT.

Preprocessing steps were identical to the steps performed for univariate analyses,

except for functional time series that were smoothed with a Gaussian kernel of 2 mm

(FWHM). MVPA was performed using CoSMoMVPA (http://www.cosmomvpa.org/;

(Oosterhof et al., 2016), implemented in MATLAB. Classification analyses were

performed using support vector machine (SVM) classifiers as implemented in LIBSVM (http://www.csie.ntu.edu.tw/~cjlin/libsvm; Chang and Lin, 2011). A general linear model was implemented in SPM8, where each block was defined as a regressor of interest. A beta map was calculated for each block separately. Two multi-class and six binary linear SVM classifiers with a linear kernel with a fixed regularization parameter of C = 1 were trained and tested for each participant separately. The two multi-class classifiers were trained and tested to discriminate between the response patterns of the 4 auditory motion directions and locations, respectively. Four binary classifiers were used to discriminate brain activity patterns for motion and location within axes (left vs. right motion, left vs. right static, up vs. down motion, up vs. down static, hereafter within axis classification). We used 8 additional classifiers to discriminate across axes (Left vs. Up, Left vs. Down, Right vs. Up, and Right vs. Down motion directions, Left vs. Up, Left vs. Down, Right vs. Up, and Right vs. Down sound source locations, hereafter across axes classification).

For each participant, the classifier was trained using a cross-validation leave-one-out procedure where training was performed with n-1 runs and testing was then applied to the remaining one run. In each cross-validation fold, the beta maps in the training set were normalized (z-scored) across conditions, and the estimated parameters were applied to the test set. To evaluate the performance of the classifier and its generalization across all the data, the previous step was repeated 12 times where in each fold a different run was used as the testing data and the classifier was trained on the other 11 runs. For each region per participant, a single classification accuracy was obtained by averaging the accuracies of all cross-validation folds.

Cross-condition classification

To test whether motion directions and sound source locations share a similar neural representation in hPT region, we performed cross-condition classification. We carried out the same steps as for the within-condition classification as described above but trained the classifier on sound source locations and tested on motion directions, and vice versa. The accuracies from the two cross-condition classification analyses were averaged. For interpretability reasons, cross-condition classification was only interpreted on the stimuli categories that the classifiers discriminated reliably (above chance level) for both motion and static conditions (e.g. if discrimination of left vs. right was not successful in one condition, either static or motion, then the left vs. right cross-condition classification analysis was not carried out).

Within-orientation classification

To foreshadow our results, cross-condition classification analyses (see previous section) showed that motion directions and sound source locations share, at least partially, a similar neural representation in hPT region. To further investigate the similarities/differences between the neural patterns evoked by motion directions and sound source locations in the hPT, we performed 4 binary classifications in which the classifiers were trained and tested on the same orientation pairs: leftward motion vs. left static, rightward motion vs. right static, upward motion vs. up static, and downward motion vs. down static. If the same orientation (leftward and left location) across conditions (motion and static) generates similar patterns of activity in hPT

region, the classifier would not be able to differentiate leftward motion direction from

left sound location. However, significant within-orientation classification would

indicate that the evoked patterns within hPT contain differential information for

motion direction and sound source location in the same orientation (e.g. left).

The mean of the four binary classifications was computed to produce one

accuracy score per ROI. Prior to performing the within-orientation and cross-condition

MVPA, each individual pattern was normalised separately across voxels so that any

cross or within-orientation classification could not be due to global univariate

activation differences across the conditions.

Statistical analysis: MVPA

Statistical significance in the multivariate classification analyses was assessed

using non-parametric tests permuting condition labels and bootstrapping (Stelzer et

al., 2013). Each permutation step included shuffling of the condition labels and re-

running the classification 100 times on the single-subject level. Next, we applied

bootstrapping procedure in order to obtain a group-level null distribution that is

representative of whole group. From each individual's null distribution one value was

randomly chosen and averaged across all the participants. This step was repeated

100,000 times resulting in a group level null distribution of 100,000 values. The

classification accuracies across participants was considered as significant if $p < 0.05$

after corrections for multiple comparisons (number of ROIs x number of tests) using

the FDR method (Benjamini and Yekutieli, 2001).

Similar approach was adopted to assess statistical difference between the

classification accuracies of two auditory categories (e.g. four motion direction vs. four

sound source location, left motion vs. left static, left motion vs. up motion, etc.). We

performed additional permutation tests (100,000 iterations) by building a null

distribution for t-stats after randomly shuffling the classification accuracy values across

two auditory categories and re-calculating the two-tail t-test between the classification

accuracies of these two categories. All p-values were corrected for multiple

comparisons (number of ROIs x number of tests) using the FDR method (Benjamini

and Yekutieli, 2001).

Representation Similarity analysis

Neural Dissimilarity matrices

To further explore the differences in the representational format between

sound source locations and motion directions in hPT region, we relied on

representation similarity analysis (RSA; Kriegeskorte et al., 2008). More specifically,

we tested the correlation between the representational dissimilarity matrix (RDM) of

right and left hPT in each participant with different computational models that

included condition-invariant models assuming orientation invariance across

conditions (motion, static), condition-distinct models assuming that sound source

location and motion direction sounds elicit highly dissimilar activity patterns, and a

series of intermediate graded models between them. The RSA was performed using

CosmoMVPA toolbox (Oosterhof et al., 2016) implemented in MATLAB. To perform

this analysis, we first extracted in each participant the activity patterns associated with

each condition (Edelman et al., 1998; Haxby et al., 2001). Then, we averaged

individual subject statistical maps (i.e. activity patterns) in order to have a mean

pattern of activity for each auditory category across runs. Finally, we used Pearson's

linear correlation as the similarity measure to compare each possible pair of the

activity patterns evoked by the four different motion directions and four different

sound source locations. This resulted in an 8 x 8 correlation matrix for each participant

that was then converted into a representational dissimilarity matrix (RDMs) by

computing 1 – correlation. Each element of the RDM contains the dissimilarity index

between the patterns of activity generated by two categories, in other words the RDM

represents how different is the neural representation of each category from the neural

representations of all the other categories in the selected ROI. The 16 neural RDMs

(1 per participant) for each of the 2 ROIs were used as neural input for RSA.

Computational models

To investigate shared representations between auditory motion directions and

sound source locations, we created multiple computational models ranging from a

fully condition-distinct model to a fully condition-invariant model with intermediate

gradients in between (Zabicki et al., 2016) (Figure 4C).

Condition-Distinct model

The condition-distinct models assume that dissimilarities between motion and

static condition is 1 (i.e. highly dissimilar), meaning that neural responses/patterns

generated by motion and static conditions are totally unrelated. For instance, there

would be no similarity between any motion directions with any sound source location.

The dissimilarity values in the diagonal were set to 0, simply reflecting that neural

responses for the same direction/location are identical to themselves.

Condition-Invariant model

The condition-invariant models assume a fully shared representation for

specific/corresponding static and motion conditions. For example, the models

consider the neural representation for the left sound source location and the left

motion direction highly similar. All within-condition (left, right, up and down)

comparisons are set to 0 (i.e. highly similar) regardless of their auditory condition. The

dissimilarities between different directions/locations are set to 1 meaning that each

within condition sound (motion or static) is different from all the other within

conditions.

Intermediate models

To detect the degree of similarity/shared representation between motion

direction and sound source location patterns, we additionally tested 2 classes of 5

different intermediate models. The two classes were used to deepen the

understanding of characteristic tuning of hPT for separate direction/location or axis of

motion/location. The two model classes represent 2 different possibilities. The first

scenario was labeled as Within-Axis Distinct, and these models assume that each of

the 4 directions/locations (i.e. left, right, up, down) would generate a distinctive neural

representation different from all of the other within-condition sounds (e.g. the

patterns of activity produced by the left category are highly different from the patterns

produced by right, up and down categories) (see Figure 4C, upper panel). To

foreshadow our results, we observed preference for axis of motion in MVP-

classification, therefore we created another class of models to further investigate

neural representations of within-axis and across-axes of auditory motion/space. The

second scenario was labeled with Within-Axis Combined, and these models assume

that opposite direction/locations within the same axis would generate similar patterns

of activity (e.g. the pattern of activity of horizontal (left and right) categories are

different from the patterns of activity of vertical categories (up and down) (see Figure

4C, lower panel).

In all intermediate models, the values corresponding to the dissimilarity

between same auditory spaces (e.g. left motion and left location) were gradually

modified from 0.9 (motion and static conditions are mostly distinct) to 0.1 (motion and

static conditions mostly similar). These models were labeled IM9, 7, 5, 3, and 1

respectively.

In all condition-distinct and intermediate models, the dissimilarity of within-

condition sounds was fixed to 0.5 and dissimilarity of within-orientation sounds was

fixed to 1. Across all models, the diagonal values were set to 0.

Performing RSA

We computed Pearson's correlation to compare neural RDMs and

computational model RDMs. The resulting correlation captures which computational

model better explains the neural dissimilarity patterns between motion direction and

sound source location conditions. To compare computational models, we performed

Mann-Whitney-Wilcoxon rank-sum test between for every pair of models. All p-values

were then corrected for multiple comparisons using the FDR method (Benjamini and

Yekutieli, 2001). To test differences between two classes of models (Within-Axis

Combined vs. Within-Axis Distinct), within each class, the correlation coefficient values

were averaged across hemispheres and across models. Next, we performed

permutation tests (100,000 iterations) by building a null distribution for differences

between classes of models after randomly shuffling the correlation coefficient values

across two classes, and re-calculating the subtraction between the correlation

coefficients of Within-Axis Combined and Within-Axis Distinct classes.

521   To visualize the distance between the patterns of the motion directions and

522   sound source locations, we used multi-dimensional scaling (MDS) to project the high-

523   dimensional RDM space onto 2 dimensions with the neural RDMs that were obtained

524   from both lhPT and rhPT.

525   Additionally, the single-subject 8 x 8 correlation matrices were used to

calculate the reliability of the data considering the signal-to-noise ratio of the data

(Kriegeskorte et al., 2007). For each participant and each ROI, the RDM was correlated

with the averaged RDM of the rest of the group. The correlation values were then

averaged across participants. This provided the maximum correlation that can be

expected from the data.

RESULTS

Behavioral results

During the experiment, we collected target direction/location discrimination responses (see Figure 1C). The overall accuracy scores were entered into 2 x 4 (Condition, Orientation) repeated measures ANOVA. No main effect of Condition ($F_{1,15}$ = 2.22; p = 0.16) was observed, indicating that the overall accuracy while detecting direction of motion or sound source location did not differ. There was a significant main effect of orientation ($F_{1.6,23.7}$ = 11.688; p = 0.001), caused by greater accuracy in the horizontal orientations (left and right) as compared to the vertical orientations (up and down). Post-hoc two-tailed t-tests (Bonferroni corrected) revealed that accuracies did not reveal significant difference within horizontal orientations (left vs right; t_{15} = -0.15, p=1), and within vertical orientations (up vs down; t_{15} = 0.89, p = 1). However, left orientation accuracy was greater as compared to down (t_{15} = 3.613, p = 0.005), and up (t_{15} = 4.51, p < 0.001) orientations and right orientation accuracy was greater as compared to the down (t_{15} = 3.76, p = 0.003) and up (t_{15} = 4.66, p < 0.001) orientation accuracies. No interaction between Condition x Orientation was observed, pointing out that differences between orientations in terms of performance expresses both for static and motion. A Greenhouse–Geisser correction was applied to the degrees of freedom and significance levels whenever an assumption of sphericity was violated.

fMRI results – whole-brain univariate analyses

To identify brain regions that are preferentially recruited for auditory motion processing, we performed a univariate RFX- GLM contrast [Motion > Static] (Figure 2B). Consistent with previous studies (Dormal et al., 2016; Getzmann and Lewald,

2012; Pavani et al., 2002; Poirier et al., 2005; Warren et al., 2002), whole-brain

univariate analysis revealed activation in the superior temporal gyri, bilateral hPT,

precentral gyri, and anterior portion of middle temporal gyrus in both hemispheres

(Figure 2B, Table 1). The most robust activation (resisting whole brain FWE correction,

p<0.05) was observed in the bilateral hPT (peak MNI coordinates [-46 -32 10] and [60

-32 12]). We also observed significant activation in occipito-temporal regions (in the

vicinity of anterior hMT+/V5) as suggested by previous studies (Dormal et al., 2016;

Poirier et al., 2005; Warren et al., 2002).

fMRI results – ROI univariate analyses

Beta parameter estimates were extracted from the pre-defined ROIs (see

methods) for the four motion directions and four sound source locations from the

auditory experiment (Figure 2C). We investigated the condition effect and the

presence of direction/location selectivity in lhPT and rhPT regions separately by

performing 2 x 4 (Conditions, Orientations) repeated measures of ANOVA with beta

parameter estimates. In lhPT, main effect of Conditions was significant ($F_{1,15}$ = 37.28,

$p < 0.001$), indicating that auditory motion evoked higher response compared to

static sounds. There was no significant main effect of Orientations ($F_{1.5,22.5}$ = 0.77, p =

0.44), and no interaction ($F_{3,45}$ = 2.21, p = 0.11). Similarly, in rhPT, only main effect of

Conditions was significant ($F_{1,15}$ = 37.02, $p < 0.001$). No main effect of Orientation

($F_{1.5,23.2}$ = 1.43, p = 0.26) or interaction ($F_{3,45}$ = 1.74, p = 0.19) was observed. Overall,

brain activity in the hPT as measured with beta parameter estimate extracted from

univariate analysis did not provide evidence of motion direction or sound source

location selectivity.

--- Figure 2 about here ---

fMRI results – ROI multivariate pattern analyses

To further investigate the presence of information about auditory motion direction

and sound source location in hPT, we ran multi-class and binary multivariate pattern

classification. Figure 3A-C shows the mean classification accuracy across categories in

each ROI.

MVPA – Within Condition

Multi-class across four conditions classification accuracy in the hPT was significantly

above chance (chance level = 25%) in both hemispheres for motion direction (lhPT:

mean ± SD = 38.4 ±7, p < 0.001; rhPT: mean ± SD = 37.1 ± 6.5, p<0.001), and sound

source location (lhPT: mean ± SD = 32.4 ± 6.7, p < 0.001; rhPT: mean ± SD = 31.2 ±

7.5, p < 0.001) (Figure 3A). In addition, we assessed the differences between

classification accuracies for motion and static stimuli by using permutation tests in

lhPT (p = 0.024) and rhPT (p = 0.024), indicating greater accuracies for classifying

motion direction than sound source location across all regions.

MVPA – Binary Within-Axis

Binary horizontal (left vs. right) within-axis classification showed significant results in

both lhPT and rhPT for static sounds (lhPT: mean ± SD = 58.6 ± 14.5, p<0.001; rhPT:

mean ± SD = 56.5 ± 11.9, p = 0.008) (Figure 3C), while motion classification was

significant only in the rhPT (mean ± SD = 55.5 ± 13.9, p = 0.018) (Figure 3B).

Moreover, binary vertical (up vs down) within-axis classification was significant only in

the lhPT for both motion (mean ± SD = 55.7 ± 7.1, p = 0.01), and static (mean ± SD

= 54.9 ± 11.9, p = 0.03) conditions (Figure 3B-C).

MVPA – Binary Across-Axis

We used 8 additional binary classifiers to discriminate across axes moving and static

sounds. Binary across-axes (Left vs. Up, Left vs. Down, Right vs. Up, and Right vs.

Down) classifications were significantly above chance level in the bilateral hPT for

motion directions (Left vs. Up lhPT: mean ± SD = 65.8 ±14.8, p<0.001; rhPT: mean ±

SD = 64.8 ± 9.4, p<0.001; Left vs. Down lhPT: mean ± SD = 74 ±15.9, p<0.001; rhPT:

mean ± SD = 66.9 ± 10.4, p<0.001; Right vs. Up lhPT: mean ± SD = 72.4 ±15.8,

p<0.001; rhPT: mean ± SD = 71.4 ± 13.3, p<0.001; Right vs. Down lhPT: mean ± SD

= 73.4 ±11.8, p<0.001; rhPT: mean ± SD = 68.2 ± 15.9, p<0.001) (Figure 3B).

However, sound source locations across-axes classifications were significant only in

certain conditions (Left vs. Up rhPT: mean ± SD = 56 ± 12.7, p=0.018; Left vs. Down

lhPT: mean ± SD = 55.2 ±11.2, p=0.024; rhPT: mean ± SD = 57.3 ± 17.2, p=0.003;

Right vs. Down lhPT: mean ± SD = 57.6 ±8.4, p=0.005) (Figure 3C).

MVPA – "Axis of Motion" Preference

In order to test whether neural patterns within hPT contain information about

opposite directions/locations within an axis, we performed four binary within-axis

classifications. Similar multivariate analyses were performed to investigate the

presence of information about across-axes directions/locations. The classification

accuracies were plotted in Figure 3B-C.

In motion direction classifications, to assess the statistical difference between

classification accuracies of across axes (left vs. up, left vs. down, and right vs. up, right

vs. down) and within axes (left vs. right, and up vs. down) directions, we performed

pairwise permutation tests and FDR-corrected for multiple comparisons. Across-axes

classification accuracies in lhPT ([left vs. up] vs. [left vs. right]: p=0.006, [left vs. down]

vs. [left vs. right]: p<0.001, [right vs. down] vs. [left vs. right]: p<0.001, [right vs. up]

vs. [left vs. right]: p=0.001), and rhPT ([left vs. up] vs. [left vs. right]: p=0.029, [left vs.

down] vs. [left vs. right]: p=0.014, [right vs. down] vs. [left vs. right]: p=0.02, [right vs.

up] vs. [left vs. right]: p=0.003) were significantly higher compared to the horizontal

within-axis classification accuracies. Similarly, across-axes classification accuracies

were significantly higher when compared with vertical within-axis classification

accuracies in lhPT ([up vs. down] vs. [left vs. up], p=0.02; [up vs. down] vs. [left vs.

down], p=0.001; [up vs. down] vs. [right vs. up], p=0.001; [up vs. down] vs. [right vs.

down], p <0.001) and rhPT ([up vs. down] vs. [left vs. up], p= 0.001; [up vs. down] vs.

[left vs. down], p=0.001; [up vs. down] vs. [right vs. up], p= 0.001; [up vs. down] vs.

[right vs. down], p=0.002). No significant difference was observed between the within-

axis classifications in lhPT ([left vs. right] vs. [up vs. down], p=0.24) and rhPT ([left vs.

right] vs. [up vs. down], p=0.31). Similarly, no significance difference was observed

among the across-axes classification accuracies in the bilateral hPT.

In static sound location classifications, no significant difference was observed

between across-axes and within-axes classification accuracies, indicating that the

classifiers did not perform better when discriminating sound source locations across

axes compared to the opposite locations.

--- Figure 3 about here ---

One may wonder whether the higher classification accuracy for across compared to

within axes relates to the perceptual differences (e.g. difficulty in localizing) in

discriminating sounds within the horizontal and vertical axes. Indeed, because we

opted for an ecological design reflecting daily-life listening condition, we observed,

as expected, that discriminating vertical directions was more difficult than

discriminating horizontal ones (Middlebrooks and Green, 1991). In order to address

this issue, we replicated our MVP-classification analyses after omitting the four

participants showing the lowest performance in discriminating the vertical motion

direction, leading to comparable performance (at the group level) within and across

axes. We replicated our pattern of results by showing preserved higher classification

accuracies across-axes than within-axis. Moreover, while accuracy differences

between across- and within-axes classification was only observed in the motion

condition, behavioral differences were observed in both static and motion conditions.

To assess whether the higher across axes classification accuracies are due to

difference in difficulties between horizontal and vertical sounds, we performed

correlation analysis. For each participant, the behavioral performance difference

between horizontal (left and right) and vertical (up and down) conditions was

calculated. The difference between the classification accuracies within axes (left vs.

right and up vs. down) and across axes (left vs. up, left vs. down, right vs. up, and right

vs. down) was calculated. Spearman correlation between the (Δ) behavioral

performance and (Δ) classification accuracies was not significant in the bilateral hPT

(lhPT $r = 0.18$, $p = 0.49$; rhPT $r = 0.4$, $p = 0.12$). An absence of correlation suggests

that there is no relation between the difference in classification accuracies and the

perceptual difference. These results strengthen the notion that "axis of motion"

organization observed in PT does not simply stem from behavioral performance

differences.

MVPA – Cross-condition

To investigate if motion direction and sound source locations rely on shared

representation in hPT, we trained the classifier to distinguish neural patterns from the

motion directions (e.g. going to the left) and then tested on the patterns elicited by

static conditions (e.g. being in the left), and vice versa.

Cross-condition classification revealed significant results across 4

direction/location (lhPT: mean ± SD = 27.8 ± 5.3, $p = 0.008$, rhPT: mean ± SD = 28.7

± 3.8, $p<0.001$) (Figure 3A). Within- axis categories did not reveal any significant

cross-condition classification. These results suggest that a partial overlap between the

neural patterns of moving and static stimuli in the hPT.

MVPA – Within-orientation

Cross-condition classification results indicated a shared representation

between motion directions and sound source locations. Previous studies argued that

successful cross-condition classification reflects an abstract representation of stimuli

conditions (Fairhall and Caramazza, 2013; Higgins et al., 2017; Hong et al., 2012). To

test this hypothesis, patterns of the same orientation of motion and static conditions

(e.g. leftward motion and left location) were involved in within-orientation MVPA. The

rational was that if the hPT region carries fully abstract representation of space, within-

orientation classification would give results in favor of the null hypothesis (no

differences within the same orientation). In the within-orientation classification

analysis, accuracies from the four within-orientation classification analyses were

averaged and survived FDR corrections in bilateral hPT (lhPT: mean ± SD = 65.6 ± 5,

$p < 0.001$, rhPT: mean ± SD = 61.9 ± 5.6, p<0.001), indicating that the neural patterns

of motion direction can be reliably differentiated from sound-source location within

hPT (Figure 4A).

--- Figure 4 about here ---

RSA

Multi-dimensional Scaling

Visualization of the representational distance between the neural patterns

evoked by motion directions and sound source locations further supported that

within-axis directions show similar geometry compared to the across-axes directions,

therefore, it is more difficult to differentiate the neural patterns of opposite directions

in MVP-classification. MDS also showed that in both lhPT and rhPT, motion directions

and sound source locations are separated into 2 clusters (Figure 4B).

RSA with external models

The correlation between model predictions and neural RDMs for the lhPT and

rhPT is shown in Figure 4D. The cross-condition classification results indicated a

shared representation within the neural patterns of hPT for motion and static sounds.

We examined the correspondence between the response pattern dissimilarities

elicited by our stimuli with 14 different model RDMs that included a fully condition

distinct, fully condition-invariant models, and intermediate models with different

degrees of shared representation.

First class of computational RDMs were modeled with the assumption that the

neural patterns of within-axis sounds are fully distinct. The analysis revealed a negative

correlation with the fully condition-invariant model in the bilateral hPT (lhPT: mean r

± SD = -0.12 ± 0.18, rhPT: mean r ± SD = -0.01 ± 0.2) that increased gradually as the

models progressed in the condition-distinct direction. The model that best fit the data

was the M9 model in the bilateral hPT (lhPT: mean r ± SD = 0.66± 0.3, rhPT: mean r

± SD = 0,65 ± 0.3). A similar trend was observed for the second class of models that

have the assumption of within-axis sounds evoke similar neural patterns. Condition-

invariant model provided the least explanation of the data (lhPT: mean r ± SD = 0.14

± 0.25, rhPT: mean r ± SD = 0.2 ± 0.29), and correlations gradually increased as the

models progressed in the condition-distinct direction. The winner models in this class

were the models M9 in lhPT and M7 in the rhPT (lhPT: mean r ± SD = 0.67 ± 0.22,

rhPT: mean r ± SD = 0.69 ± 0.15).

In addition, we assessed differences between correlation coefficients for

computational models using Mann-Whitney-Wilcoxon rank-sum test for each class of

models and hemispheres separately (Figure 4D-E). In Within-Axis Distinct class in

lhPT, pairwise comparisons of correlation coefficients did not show significant difference for [Condition Distinct vs. IM.9; p = 0.8], [Condition Distinct vs. IM7; p = 0.6; rhPT], [Condition Distinct vs. IM.5; p = 0.09], however as the models progressed in further condition invariant direction, the difference between correlation coefficients for models reached significance ([Condition Distinct vs. IM.3], p = 0.012; [Condition Distinct vs. IM.1], p = 0.007; [Condition Distinct vs. Condition Invariant], p <0.001), indicating Condition Distinct model provided stronger correlation compared to the models representing conditions similarly. In rhPT, the rank-sum tests between each pairs revealed no significant difference for [Condition Distinct vs. IM.9; p = 0.9], [Condition Distinct vs. IM7; p = 0.7], [Condition Distinct vs. IM.5; p = 0.3], and also [Condition Distinct vs. IM.3; p = 0.06], however as the models progressed in further condition invariant direction, the difference between correlation coefficients for models reached significance ([Condition Distinct vs. IM.1], p =0.006; [Condition Distinct vs. Condition Invariant], p <0.001).

The two classes of models were used to deepen our understanding of characteristic tuning of hPT for separate direction/location or axis of motion/location. To reveal the differences between Within-Axis Distinct and Within-Axis Combine classes of models, we relied on 2-sided signed-rank test. Within-Axis Combined models explained our stimuli space better than Within-Axis Distinct models supporting similar pattern representation within planes (p = 0.0023).

DISCUSSION

In line with previous studies, our univariate results demonstrate higher activity

for moving than for static sounds in the superior temporal gyri, bilateral hPT,

precentral gyri, and anterior portion of middle temporal gyrus in both hemispheres

(Baumgart and Gaschler-Markefski, 1999; Krumbholz et al., 2005; Pavani et al., 2002;

Poirier et al., 2005; Warren et al., 2002). The most robust cluster of activity was

observed in the bilateral hPT (Figure 2B, Table 1). Moreover, activity estimates

extracted from independently defined hPT also revealed higher activity for moving

relative to static sounds in this region. Both whole-brain and ROI analyses therefore

clearly indicated a functional preference (expressed here as higher activity level

estimates) for motion processing over sound-source location in bilateral hPT regions

(Figure 2).

Does hPT contain information about specific motion directions and sound

source locations? At the univariate level, our four (left, right, up and down) motion

directions and sound source locations did not evoke differential univariate activity in

hPT region (see Figure 2C). We then carried out multivariate pattern classification and

observed that bilateral hPT contains reliable distributed information about the four

auditory motion directions (Figure 3). Our results are therefore similar to the

observations made with fMRI in the human visual motion area hMT+/V5 showing

reliable direction-selective information despite comparable voxel-wise univariate

activity levels across directions (Kamitani and Tong, 2006; but see Zimmermann et al.,

2011). Within-axis MVP-classification results revealed that even if both horizontal (left

versus right), and vertical (up versus down) motion directions can be classified in the

hPT region (see Figure 3C-D), across axes (e.g. left versus up) direction classifications

revealed much higher decoding accuracies compared to within-axis classifications. Such enhanced classification accuracy across axes versus within axis is reminiscent of observations made in MT+/V5 where a large-scale axis of motion selective organization was observed in non-human primates (Albright et al., 1984), and in human area MT+/V5 (Zimmermann et al., 2011). Further examination with RSA provided additional evidence that within-axis combined models (aggregating the opposite directions/location) explain better the neural representational space of hPT by showing higher correlations values compared to within-axis distinct models (see Figure 4D-E). Our findings suggest that the topographic organization principle of hMT+/V5 and hPT shows similarities in representing motion directions. Animal studies have shown that hMT+/V5 contains motion direction selective columns with specific directions organized side by side with their respective opposing motion direction counterparts (Albright et al., 1984; Born and Bradley, 2005; Diogo et al., 2003; Geesaman et al., 1997), an organization also probed using fMRI in humans (Zimmermann et al., 2011; but see below for alternative accounts). The observed difference for within versus between axis direction decoding may potentially stem from such underlying cortical columnar organization (Bartels et al., 2008; Haynes and Rees, 2006; Kamitani and Tong, 2005). Alternatively, it has been suggested that classifying orientation preference reflects much larger scale (e.g. retinotopy) rather than columnar organization (Op de Beeck, 2010; Freeman et al., 2011, 2013). Interestingly, high-field fMRI studies showed that the fMRI signal carries information related to both large- and fine-scale (columnar level) biases (Gardumi et al., 2016; Pratte et al., 2016; Sengupta et al., 2017). The present study sheds important new

lights on the coding mechanism of motion direction within the hPT and demonstrates that fMRI signal in the hPT contains direction specific information and point toward an "axis of motion" organization. This result highlights intriguing similarities between the visual and auditory systems, arguing for common neural coding strategies of motion directions across the senses. However, further studies are needed to test the similarities between the coding mechanisms implemented in visual and auditory motion selective regions, and more particularly, what drives the directional information captured with fMRI in the auditory cortex.

Supporting univariate motion selectivity results in bilateral hPT, MVPA revealed higher classification accuracies for moving than for static sounds (Figure 3A-B). However, despite minimal univariate activity elicited by sound-source location in hPT, and the absence of reliable univariate differences in the activity elicited by each position (see Figure 2C), within-axis MVP-classification results showed that beside the vertical axis (up versus down), sound source location information can be reliably decoded bilaterally in hPT (Figure 3C). Our results are in line with previous studies showing that posterior regions in auditory cortex exhibit location sensitivity both in animals (Recanzone, 2000; Stecker et al., 2005; Tian et al., 2001) and humans (Ahveninen et al., 2006, 2013; Brunetti et al., 2005; Deouell et al., 2007; Derey et al., 2016; Krumbholz et al., 2005; Warren and Griffiths, 2003; Zatorre et al., 2002; Alain et al., 2001). In contrast to what was observed for motion direction however, within-axis classifications did not reveal lower decoding accuracies when compared to across-axis classifications in hPT for static sounds. This indicates that auditory sound

source locations might not follow similar topographic representations than the one

observed for motion directions.

To which extend the neural representation of motion directions and sound

source locations overlaps has long been debated (Grantham, 1986; Kaas et al., 1999;

Poirier et al., 2017; Romanski et al., 2000; Smith et al., 2004, 2007; Zatorre and Belin,

2001). A number of neurophysiological studies have reported direction-selective

neurons along the auditory pathway (Ahissar et al. 1992; Spitzer and Semple 1993;

Doan et al. 1999; Stumpf et al. 1992; Toronchuk et al. 1992; McAlpine et al. 2000).

However, whether these neurons exhibit selectivity to motion direction specifically, or

also to spatial location remains debated (Poirier et al. 1997; McAlpine et al. 2000;

Ingham et al. 2001; Oliver 2003). In our study, we revisit this question by relying on

cross-condition classification that revealed that auditory motion (e.g. going to the left)

and sound-source location (being on the left) share partial neural representations in

hPT (Figure 3A). This observation suggests that there is a shared level of computation

between sounds located on a given position and sounds moving towards this

position. Low-level features of these two types of auditory stimuli vary in many ways

and produce large difference at the univariate level in the cortex (see Figure 2B).

However, perceiving, for instance, a sound going toward the left side or located on

the left side evoke a sensation of location/direction in the external space that is

common across conditions. Our significant cross-condition classification may

therefore relate to the evoked sensation/perception of an object being/going to a

common external spatial location. Electrophysiological studies in animals

demonstrated that motion-selective neurons in the auditory cortex displayed similar

response profile to sounds located or moving toward the same position in external

space, suggesting that the processing of sound-source locations may contribute to

the perception of moving sounds (Ahissar et al., 1992; Doan et al., 1999; Poirier et al.,

1997). Results from human psychophysiological and auditory evoked potential studies

also strengthen the notion that sound source location contributes to motion

perception (Getzmann and Lewald, 2011; Strybel and Neale, 1994). Our cross-

condition MVPA results therefore extend the notion that motion directions and sound

source locations might have common features that are partially shared for encoding

spatial sounds.

Significant cross-condition classification based on multivariate fMRI signal has

typically been considered as a demonstration that the region implements a partly

common and abstracted representation of the tested conditions (Fairhall and

Caramazza, 2013; Higgins et al., 2017; Hong et al., 2012). For instance, a recent study

elegantly demonstrated that the human auditory cortex at least partly integrates

interaural time and level differences (ITD and ILD) into a higher-order representation

of auditory space based on significant cross-cue classification (training on ITD and

classifying ILD, and reversely) (Higgins et al., 2017). In the present study, we argue

that even if cross-condition MVP-classification indeed demonstrate the presence of

partially shared information across sound source location and direction; successful

cross-MVPA results should however not be taken as evidence that the region

implements a purely abstract representation of auditory space. Our successful across-

condition classification (see Figure 4A) demonstrated that, even though there are

shared representations for moving and static sounds within hPT, classifiers are able to

easily distinguish motion directions from sound source locations (e.g. leftward versus

left location). RSA analyses further supported the idea that moving and static sounds

elicit distinct patterns in hPT (see Figure 4B-D). Altogether, our results suggest that

hPT contains both motion direction and sound-source location information but that

the neural patterns related to these two conditions are only partially overlapping. Our

observation of significant cross-condition classification based on highly distinct

pattern of activity between static and moving sounds may support the notion that

even if location information could serve as a substrate for movement detection,

motion encoding does not solely rely on location information (Ducommun et al., 2002;

Getzmann, 2011; Poirier et al., 2017).

REFERENCES

Ahissar, M., Ahissar, E., Bergman, H., and Vaadia, E. (1992). Encoding of sound-

source location and movement: activity of single neurons and interactions between

adjacent neurons in the monkey auditory cortex. J. Neurophysiol. *67*, 203–215.

Ahveninen, J., Jaaskelainen, I.P., Raij, T., Bonmassar, G., Devore, S.,

Hamalainen, M., et al. (2006). Task-modulated "what" and "where" pathways in

human auditory cortex. Proc. Natl. Acad. Sci. U. S. A. *103*, 14608–14613.

Ahveninen, J., Huang, S., Nummenmaa, A., Belliveau, J.W., Hung, A.-Y.,

Jääskeläinen, et al. (2013). Evidence for distinct human auditory cortex regions for

sound location versus identity processing. Nat. Commun. *4*, 871–882.

Alain, C., Arnott, S.R., Hevenor, S.J., Graham, S., Grady, C.L., 2001. "What" and

"where" in the human auditory system. Proc. Natl. Acad. Sci. U.S.A. *98*, 12301–12306.

Albright, T.D., Desimone, R., and Gross, C.G. (1984). Columnar organization of

directionally selective cells in visual area MT of the macaque. J. Neurophysiol. *51*, 16–

31.

Alink, A., Euler, F., Kriegeskorte, N., Singer, W., and Kohler, A. (2012). Auditory

motion direction encoding in auditory cortex and high-level visual cortex. Hum. Brain

Mapp. *33*, 969–978.

Altman, J.A. (1968). Are there neurons detecting direction of sound source

motion? Exp. Neurol. *22*, 13–25.

Altman, J.A. (1994). Processing of Information Concerning Moving Sound

Sources in the Auditory Centers and its Utilization by Brain Integrative Structures.

Sens. Syst. *8*, 255–261.

Arnott, S. R., Binns, M. A., Grady, C. L., and Alain, C. (2004). Assessing the

auditory dual-pathway model in humans. Neuroimage *22*, 401–408.

Barrett, D.J.K., and Hall, D.A. (2006). Response preferences for "what" and

"where" in human non-primary auditory cortex. Neuroimage *32*, 968–977.

Bartels, A., Logothetis, N.K., and Moutoussis, K. (2008). fMRI and its

interpretations: an illustration on directional selectivity in area V5 / MT. Trends in

Neuro. *31*, 444-453.

Battal, C., Rezk, M., Mattioni, S., Vadlamudi, J., Collignon, O. (2018).

Representation of auditory motion directions and sound source locations in the

human planum temporale. http://biorxiv.org/cgi/content/short/302497v2.

Baumgart, F., and Gaschler-Markefski, B. (1999). A movement-sensitive area in

auditory cortex. Nature *400*, 1997–1999.

Beckett, A., Peirce, J.W., Sanchez-Panchuelo, R.M., Francis, S., and Schluppeck,

D. (2012). Contribution of large-scale biases in decoding of direction-of-motion from

high-resolution fMRI data in human early visual cortex. Neuroimage *63*, 1623–1632.

Op de Beeck, H.P. (2010). Against hyperacuity in brain reading: Spatial

smoothing does not hurt multivariate fMRI analyses? Neuroimage *49*, 1943–1948.

Benjamini, Y., and Yekutieli, D. (2001). The control of the false discovery rate in

multiple testing under dependency. Ann. Stat. *29*, 1165–1188.

Benson, D.A., Hienz, R.D., and Goldstein, M.H. (1981). Single-unit activity in the

auditory cortex of monkeys actively localizing sound sources: Spatial tuning and

behavioral dependency. Brain Res. *219*, 249–267.

Blauert, J. (1982). Binaural localization. Scand.Audiol.Suppl *15*, 7–26.

Born, R.T., and Bradley, D.C. (2005). Structure and Function of Visual Area MT.

Ann. Review of Neuro. *28*, 157–189.

Boudreau, J.C. & Tsuchitani, C., (1968). Binaural interaction in the cat superior

olive S segment. *Journal of neurophysiology*, 31(3), pp.442–454

Braddick, O.J., O'Brien, J.M.D., Wattam-Bell, J., Atkinson, J., Hartley, T., and

Turner, R. (2001). Brain areas sensitive to coherent visual motion. Perception *30*, 61–

72.

Bremmer, F., Schlack, A., Shah, N.J., Zafiris, O., Kubischik, M., Hoffmann, K.,

Zilles, K., and Fink, G.R. (2001). Polymodal motion processing in posterior parietal and

premotor cortex: a human fMRI study strongly implies equivalencies between humans

and monkeys. Neuron *29*, 287–296.

Britten, K.H., Newsome, W.T., Shadlen, M.N., Celebrini, S., and Movshon, J.A.

(1996). A relationship between behavioral choice and the visual responses of neurons

in macaque MT. Vis. Neurosci. *13*, 87–100.

Brunetti, M., Belardinelli, P., Caulo, M., Del Gratta, C., Della Penna, S., Ferretti,

et al. (2005). Human brain activation during passive listening to sounds from different

locations: an fMRI and MEG study. Hum. Brain Mapp. *26*, 251–261.

Carlile, S., and Leung, J. (2016). The Perception of Auditory Motion. Trends

Hear. *20*, 1–19.

Chang, C., and Lin, C., (2011). LIBSVM: a library for support vector machines.

ACM Transactions on Intelligent Systems and Technology *2*, 27:1-27:27. Software

available at http://www.csie.ntu.edu.tw/~cjlin/libsvm

Collignon, O., Vandewalle, G., Voss, P., Albouy, G., Charbonneau, G.,

Lassonde, M., et al. (2011). Functional specialization for auditory-spatial processing in

the occipital cortex of congenitally blind humans. Proc. Natl. Acad. Sci. U. S. A. *108*,

4435–4440.

Cox D.D., and Savoy R.L. (2003) Functional magnetic resonance imaging (fMRI)

"brain reading": detecting and classifying distributed patterns of fMRI activity in

human visual cortex. Neuroimage *19*, 261–270.

De Martino, F., Valente, G., Staeren, N., Ashburner, J., Goebel, R., and Formi-

sano, E. (2008). Combining multivariate voxel selection and support vector machines

for mapping and classification of fMRI spatial patterns. NeuroImage *43*, 44–58.

Deouell, L.Y., Heller, A.S., Malach, R., Esposito, M.D., and Knight, R.T. (2007).

Cerebral Responses to Change in Spatial Location of Unattended Sounds. Neuron *55*,

985–996.

Derey, K., Valente, G., De Gelder, B., and Formisano, E. (2016). Opponent

Coding of Sound Location (Azimuth) in Planum Temporale is Robust to Sound-Level

Variations. Cereb. Cortex *26*, 450–464.

Diogo, A.C.M., Soares, J.G.M., Koulakov, A., Albright, T.D., and Gattass, R.

(2003). Electrophysiological imaging of functional architecture in the cortical middle

temporal visual area of Cebus apella monkey. J. Neurosci. *23*, 3881–3898.

Doan, D.E., Saunders, J.C., Field, F., Ingham, N.J., Hart, H.C., and McAlpine,

D. (1999). Sensitivity to Simulated Directional Sound Motion in the Rat Primary

Auditory Cortex. J. Neurophysi. 81, 2075–2087.

Dormal, G., Rezk, M., Yakobov, E., Lepore, F., and Collignon, O. (2016).

Auditory motion in the sighted and blind: Early visual deprivation triggers a large-

scale imbalance between auditory and "visual" brain regions. Neuroimage *134*, 630–

644.

Ducommun, C.Y., Murray, M.M., Thut, G., Bellmann, A., Viaud-Delmon, I.,

Clarke, S., et al. (2002). Segregated processing of auditory motion and auditory

location: an ERP mapping study. Neuroimage *16*, 76–88.

Edelman, S., Grill-Spector, K., Kushnir, T., and Malach, R. (1998). Toward direct

visualization of the internal shape represetation space by fMRI. Psychobiology *26*,

309–321.

Eickhoff, S.B., Paus, T., Caspers, S., Grosbras, M.H., Evans, A.C., Zilles, K., et al.

(2007). Assignment of functional activations to probabilistic cytoarchitectonic areas

revisited. Neuroimage *36*, 511–521.

Fairhall, S.L., and Caramazza, A. (2013). Brain Regions That Represent Amodal

Conceptual Knowledge. J. Neurosci. *33*, 10552–10558.

Freeman, J., Brouwer, G.J., Heeger, D.J., and Merriam, E.P. (2011). Orientation

decoding depends on maps, not columns. J. Neurosci. *31*, 4792–4804.

Freeman, J., Heeger, D.J., and Merriam, E.P. (2013). Coarse-Scale Biases for

Spirals and Orientation in Human Visual Cortex. J. Neurosci. *33*, 19695–19703.

Gardumi, A., Ivanov, D., Hausfeld, L., Valente, G., Formisano, E., and Uludağ,

K. (2016). The effect of spatial resolution on decoding accuracy in fMRI multivariate

pattern analysis. Neuroimage *132*, 32–42.

Geesaman, B.J., Born, R.T., Andersen, R.A., and Tootell, R.B. (1997). Maps of

complex motion selectivity in the superior temporal cortex of the alert macaque

monkey: a double-label 2-deoxyglucose study. Cereb Cortex *7*, 749–757.

Getzmann, S. (2011). Auditory motion perception: onset position and motion

direction are encoded in discrete processing stages. Eur. J. Neurosci. *33*, 1339–1350.

Getzmann, S., and Lewald, J. (2011). The effect of spatial adaptation on auditory

motion processing. Hear. Res. *272*, 21–29.

Getzmann, S., and Lewald, J. (2012). Cortical processing of change in sound

location: Smooth motion versus discontinuous displacement. Brain Res. *1466*, 119–

127.

Grantham, D.W. (1986). Detection and discrimination of simulated motion of

auditory targets in the horizontal plane. J. Acoust. Soc. Am. *79*, 1939–1949.

Griffiths, T.D., Büchel, C., Frackowiak, R.S., and Patterson, R.D. (1998). Analysis

of temporal structure in sound by the human brain. Nat. Neurosci. *1*, 422–427.

Grothe B., Pecka M., McAlpine D. (2010). Mechanisms of sound localization in mammals. *Physiol Rev* **90**:983–1012.

Hall, D.A., Haggard, M.P., Akeroyd, M.A., Palmer, A.R., Summerfield, A.Q., Elliott, et al. (1999). "Sparse" temporal sampling in auditory fMRI. Hum. Brain Mapp. *7*, 213–223.

Hall, D.A., and Moore, D.R. (2003). Auditory Neuroscience : The Salience of Looming Sounds. Curr. Biology *13*, 91–93.

Haxby, J. V, Gobbini, M.I., Furey, M.L., Ishai, A., Schouten, J.L., and Pietrini, P. (2001). Distributed and overlapping representations of faces and objects in ventral temporal cortex. Science *293*, 2425–2430.

Haynes, J.D., and Rees, G. (2006). Decoding mental states from brain activity in humans. Nat. Rev. Neurosci. *7*, 523–534.

Heeger, D.J., Boynton, G.M., Demb, J.B., Seidemann, E., and Newsome, W.T. (1999). Motion opponency in visual cortex. J. Neurosci. *19*, 7162–7174.

Higgins, N.C., McLaughlin, S.A., Rinne, T., and Stecker, G.C. (2017). Evidence for cue-independent spatial representation in the human auditory cortex during active listening. Proc. Natl. Acad. Sci. *114*, E7602–E7611.

Imig, T.J., Irons, W. a, and Samson, F.R. (1990). Single-unit selectivity to azimuthal direction and sound pressure level of noise bursts in cat high-frequency primary auditory cortex. J. Neurophysiol. *63*, 1448–1466.

Imig, T.J., Bibikov, N.G., Poirier, P., and Samson, F.K. (2000). Directionality derived from pinna-cue spectral notches in cat dorsal cochlear nucleus. J. Neurophysiol. *83*, 907–925.

Ingham, N.J., Hart, H.C., and McAlpine, D. (2001). Spatial receptive fields of

inferior colliculus neurons to auditory apparent motion in free field. J. Neurophysiol.

85, 23–33.

Jiang, F., Stecker, G.C., and Fine, I. (2014). Auditory motion processing after

early blindness. J. Vis. *14*, 4–4.

Jiang, F., Stecker, G.C., Boynton, G.M., and Fine, I. (2016). Early Blindness

Results in Developmental Plasticity for Auditory Motion Processing within Auditory

and Occipital Cortex. Front. Hum. Neurosci. *10*, 324.

Kaas, J.H., Hackett, T.A., and Tramo, M.J. (1999). Auditory processing in

primate cerebral cortex. Curr. Opin. Neurobiol. *9*, 164–170.

Kamitani, Y., and Tong, F. (2005). Decoding the visual and subjective contents

of the human brain. *8*, 679–685.

Kamitani, Y., and Tong, F. (2006). Decoding Seen and Attended Motion

Directions from Activity in the Human Visual Cortex. Curr. Biol. *16*, 1096–1102.

Kamitani, Y., and Sawahata, Y. (2010). Spatial smoothing hurts localization but

not information: Pitfalls for brain mappers. Neuroimage *49*, 1949–1952.

Kriegeskorte N, Mur M, Bandettini P. (2008). Representational similarity

analysis—connecting the branches of systems neuroscience. Front. Syst. Neurosci. *2*,

1-28.

Kriegeskorte, N., Formisano, E., Sorger, B., and Goebel, R. (2007). Individual

faces elicit distinct response patterns in human anterior temporal cortex. Proc. Natl.

Acad. Sci. *104*, 20600–20605.

Krumbholz, K., Schönwiesner, M., Rübsamen, R., Zilles, K., Fink, G.R., and von

1051 Cramon, D.Y. (2005). Hierarchical processing of sound location and motion in the

1052 human brainstem and planum temporale. Eur. J. Neurosci. *21*, 230–238.

1053 Lewis, J.W., Beauchamp, M.S., and DeYoe, E.A. (2000). A comparison of visual

1054 and auditory motion processing in human cerebral cortex. Cereb.Cortex *10*, 873–888.

1055 Lomber, S.G., and Malhotra, S. (2008). Double dissociation of "what" and

1056 "where" processing in auditory cortex. Nat. Neurosci. *11*, 609–616.

1057 Maeder, P. P., Meuli, R. A., Adriani, M., Bellmann, A., Fornari, E., Thiran, J. P.,

1058 et al. (2001). Distinct pathways involved in sound recognition and localization: a

1059 human fMRI study. Neuroimage *14*, 802–816.

1060 McAlpine, D., Jiang, D., Shackleton, T.M., and Palmer, A.R. (2000). Responses

1061 of neurons in the inferior colliculus to dynamic interaural phase cues: evidence for a

1062 mechanism of binaural adaptation. J Neurophysiol *83*, 1356– 1365.

1063 Middlebrooks, J., and Pettigrew, J. (1981). Functional classes of neurons in

1064 primary auditory cortex of the cat distinguished by sensitivity to sound location. J

1065 Neurosci *1*, 107–120.

1066 Middlebrooks, J.C., and Green, D.M. (1991). Sound localization by human

1067 listeners. Annu. Rev. Psychol. *42*, 135–159.

1068 Middlebrooks, J.C. (2002). Auditory space processing: Here, there or

1069 everywhere? Nat. Neurosci. *5*, 824–826.

1070 Middlebrooks, J.C., and Bremen, P. (2013). Spatial Stream Segregation by

1071 Auditory Cortical Neurons. J. Neurosci. *33*, 10986–11001.

1072 Movshon, J.A., and Newsome, W.T. (1996). Visual response properties of striate

1073 cortical neurons projecting to area MT in macaque monkeys. J. Neurosci. *16*, 7733–

7741.

Newsome, W.T., and Park, E.B. (1988). A Selective Impairment of Motion Perception Following Lesions of the Middle Temporal Visual Area (MT). J. Neurosci. *8*, 2201–2211.

Norman, K.A., Polyn, S.M., Detre, G.J., Haxby, J.V. (2006). Beyond mind-reading: multi-voxel pattern analysis of fMRI data. Trends Cogn Sci *10*, 424–430.

Oliver, D. L., Beckius, G. E., Bishop, D. C., Loftus, W. C., & Batra, R. (2003). Topography of interaural temporal disparity coding in projections of medial superior olive to inferior colliculus. *The Journal of neuroscience 23*, 7438-7449.

Oosterhof, N.N., Connolly, A.C., and Haxby, J. V. (2016). CoSMoMVPA: Multi-Modal Multivariate Pattern Analysis of Neuroimaging Data in Matlab/GNU Octave. Front. Neuroinform. *10*, 1–27.

Ortiz-Rios, M., Azevedo, F.A.C., Kuśmierek, P., Balla, D.Z., Munk, M.H., Keliris, G.A., et al. (2017). Widespread and Opponent fMRI Signals Represent Sound Location in Macaque Auditory Cortex. Neuron *93*, 971–983.e4.

Pavani, F., Macaluso, E., Warren, J.D., Driver, J., and Griffiths, T.D. (2002). A common cortical substrate activated by horizontal and vertical sound movement in the human brain. Curr. Biol. *12*, 1584–1590.

Poirier, C., Collignon, O., Devolder, A.G., Renier, L., Vanlierde, A., Tranduy, D., et al. (2005). Specific activation of the V5 brain area by auditory motion processing: an fMRI study. Brain Res.Cogn Brain Res. *25*, 650–658.

Poirier, C., Baumann, S., Dheerendra, P., Joly, O., Hunter, D., Balezeau, et al. (2017). Auditory motion-specific mechanisms in the primate brain. PLoS Biol. *15*, 1–

24.

Poirier, P., Jiang, H., Lepore, F., and Guillemot, J.P. (1997). Positional,

directional and speed selectivities in the primary auditory cortex of the cat. Hear. Res.

113, 1–13.

Pratte, M.S., Sy, J.L., Swisher, J.D., and Tong, F. (2016). Radial bias is not

necessary for orientation decoding. Neuroimage *127*, 23–33.

Rajan, R., Aitkin, L.M., and Irvine, D.R. (1990). Azimuthal sensitivity of neurons in

primary auditory cortex of cats. II. Organization along frequency-band strips. J.

Neurophysiol. *64*, 888–902.

Rauschecker, J.P., and Tian, B. (2000). Mechanisms and streams for processing

of "what" and "where" in auditory cortex. Proc.Natl.Acad.Sci.U.S.A. *97*, 11800–

11806.

Recanzone, G.H. (2000). Spatial processing in the auditory cortex of the

macaque monkey. Proc. Natl. Acad. Sci. U. S. A. *97*, 11829–11835.

Romanski, Tian, Fritz, Mishkin, Goldman-Rakic, and Rauschecker (2000). Reply

to "What', "where" and "how" in auditory cortex'. Nat. Neurosci. *3*, 966.

Romanski, L.M., Tian, B., Fritz, J., Mishkin, M., Goldman-Rakic, P.S., and

Rauschecker, J.P. (1999). Dual streams of auditory afferents target multiple domains

in the primate prefrontal cortex. Nat.Neurosci. *2*, 1131–1136.

Searle, C.L., Braida, L.D., Davis, M.F., and Colburn, H.S. (1976). Model for

auditory localization. J. Acoust. Soc. Am. *60*, 1164–1175.

Sengupta, A., Yakupov, R., Speck, O., Pollmann, S., and Hanke, M. (2017). The

effect of acquisition resolution on orientation decoding from V1 BOLD fMRI at 7 T.

Neuroimage *148*, 64–76.

Smith, K.R., Okada, K., Saberi, K., and Hickok, G. (2004). Human cortical

auditory motion areas are not motion selective. Neuroreport *15*, 1523–1526.

Smith, K.R., Saberi, K., and Hickok, G. (2007). An event-related fMRI study of

auditory motion perception: No evidence for a specialized cortical system. Brain Res.

1150, 94–99.

Smith, K.R., Hsieh, I.H., Saberi, K., and Hickok, G. (2010). Auditory spatial and

object processing in the human planum temporale: no evidence for selectivity. J.

Cogn. Neurosci. *22*, 632–639.

Spitzer, M.W., and Semple, M.N. (1993). Responses of inferior colliculus neurons

to time-varying interaural phase disparity: effects of shifting the locus of virtual

motion. J Neurophysiol *69* 1245–1263.

Stecker, G.C., Harrington, I.A., and Middlebrooks, J.C. (2005). Location coding

by opponent neural populations in the auditory cortex. PLoS Biol. *3*, 0520–0528.

Stelzer, J., Chen, Y., and Turner, R. (2013). Statistical inference and multiple

testing correction in classification-based multi-voxel pattern analysis (MVPA): Random

permutations and cluster size control. Neuroimage *65*, 69–82.

Strybel, T.Z., and Neale, W. (1994). The effect of burst duration, interstimulus

onset interval, and loudspeaker arrangement on auditory apparent motion in the free

field. J. Acoust. Soc. Am. *96*, 3463–3475.

Stumpf, E., Toronchuk, J.M., Cynader, M.S., (1992). Neurons in cat primary

auditory cortex sensitive to correlates of auditory motion in three-dimensional space.

Exp. Brain Res. *88*, 158–168.

Tian, B., Reser, D., Durham, A., Kustov, A., and Rauschecker, J.P. (2001).

Functional Specialization in Rhesus Monkey Auditory Cortex. Science *292*, 290–293.

Tootell, R.B.H., Reppas, J.B., Dale, A.M., Look, R.B., Sereno, M.I., Malach, R., et

al. (1995). Visual motion aftereffect in human cortical area MT revealed by functional

magnetic resonance imaging. Nature *375*, 139–141.

Toronchuk, J.M., Stumpf, E., Cynader, M.S., (1992). Auditory cortex neurons

sensitive to correlates of auditory motion: underlying mechanisms. Exp. Brain Res. *88*,

169–180.

Wang, H.X., Merriam, E.P., Freeman, J., and Heeger, D.J. (2014). Motion

Direction Biases and Decoding in Human Visual Cortex. J. Neurosci. *34*, 12601–

12615.

Warren, J., Zielinski, B., and Green, G. (2002). Perception of sound-source

motion by the human brain. Neuron *34*, 139–148.

Warren, J.D., and Griffiths, T.D. (2003). Distinct mechanisms for processing

spatial sequences and pitch sequences in the human auditory brain. J.Neurosci. *23*,

5799–5804.

Woods, T.M., Lopez, S.E., Long, J.H., Rahman, J.E., and Recanzone, G.H.

(2006). Effects of Stimulus Azimuth and Intensity on the Single-Neuron Activity in the

Auditory Cortex of the Alert Macaque Monkey. J. Neurophysiol. *96*, 3323–3337.

Hong, S., Tong, F., and Seiffert, A.E. (2012). Direction-selective patterns of

activity in human visual cortex suggest common neural substrates for different types

of motion. Neuropsychologia *50*, 514–521.

Yarkoni, T., Poldrack, R.A., Nichols, T.E., Van Essen, D.C., and Wager, T.D.

(2011). Large-scale automated synthesis of human functional neuroimaging data. Nat.

Methods *8*, 665–670.

Young, E.D., Spirou, G.A., Rice, J.J., Voigt, H.F., and Rees, A. (1992). Neural

Organization and Responses to Complex Stimuli in the Dorsal Cochlear Nucleus.

Philos. Trans. R. Soc. B Biol. Sci. *336*, 407–413.

Zabicki, A., De Haas, B., Zentgraf, K., Stark, R., Munzert, J., and Krüger, B.

(2016). Imagined and Executed Actions in the Human Motor System: Testing Neural

Similarity Between Execution and Imagery of Actions with a Multivariate Approach.

Cereb. Cortex *27*, 1–14.

Zatorre, R.J., and Belin, P. (2001). Spectral and Temporal Processing in Human

Auditory Cortex. Cereb. Cortex *11*, 946–953.

Zatorre, R.J., Bouffard, M., Ahad, P., and Belin, P. (2002). Where is "where" in

the human auditory cortex? Nat. Neurosci. *5*, 905–909.

Zeng, H., and Constable, R.T. (2002). Image distortion correction in EPI:

Comparison of field mapping with point spread function mapping. Magn. Reson.

Med. *48*, 137–146.

Zhou, Y., and Wang, X. (2012). Level dependence of spatial processing in the

primate auditory cortex. J. Neurophysiol. *108*, 810–826.

Zimmermann, J., Goebel, R., de Martino, F., van de Moortele, P.F., Feinberg,

D., Adriany, G., et al. (2011). Mapping the Organization of Axis of Motion Selective

Features in Human Area MT Using High-Field fMRI. PLoS One *6*, 1–10.

1187

 FIGURE LEGENDS

 Figure 1. Stimuli and Experimental Design. (A). The acoustic apparatus used to

present auditory moving and static sounds while binaural recordings were carried out

for each participant before the fMRI session. **(B).** Auditory stimuli presented inside the

MRI scanner consisted of 8 conditions: leftward, rightward, downward and upward

moving stimuli and left, right, down and up static stimuli. Each condition was

presented for 15 s (12 repetition of 1250 ms sound, no ISI) and followed by 7 s gap

for indicating the corresponding direction/location in space and 8 s of silence (total

inter-block interval was 15 s). Sound presentation and response button press were

pseudo-randomized. Participants were asked to respond as accurately as possible

during the gap period. **(C).** The behavioural performance inside the scanner.

Figure 2. Univariate whole brain results. (A). Association test map was obtained from

the online tool Neurosynth using the term "Planum Temporale" (FDR corrected p

<0.05). The black spheres are illustration of drawn mask (radius = 6mm, 117 voxels)

around the peak coordinate from Neurosynth (search term "planum temporale",

meta-analysis of 85 studies). **(B).** Auditory motion processing [motion > static]

thresholded at p <0.05 FWE-corrected. **(C).** Mean activity estimates (arbitrary units ±

SEM) associated with the perception of auditory motion direction (red) and sound-

source location (blue). ML: motion left, MR: motion right, MD: motion down, MU:

motion up, SL: static left, SR: static right, SD: static down, and SU: static up.

Figure 3. Within and cross-classification results. (A). Classification results for the 4

conditions in functionally defined hPT region. Within-condition and cross-condition

classification results are shown in the same bar plots. Moving: four motion direction;

Static: four sound source location; and Cross: cross-condition classification accuracies.

(B). Classification results of within (left vs. right, up vs. down) and across axes (left vs.

up, left vs. down, right vs. up, right vs. down) motion directions. **(C).** Classification

results of within (left vs. right, up vs. down) and across axes (left vs. up, left vs. down,

right vs. up, right vs. down) sound source locations. LvsR: Left vs. Right, UvsD: Up vs.

Down, LvsU: Left vs. Up, LvsD: Left vs. Down, RvsU: Right vs. Up, RvsD: Right vs. Down

classifications. FDR corrected p-values: (*) $p<0.05$, (**) $p<0.01$, (***) $p<0.001$ testing

differences against chance level (dotted lines; see methods).

Figure 4. Pattern dissimilarity between motion directions and sound source locations.

(A). Across-condition classification results of across 4 conditions are represented in

each ROI (lhPT and rhPT). 4 binary classifications [leftward motion vs. left location],

[rightward motion vs. right location], [upward motion vs. up location], and [downward

motion vs. down location] were computed and averaged to produce one accuracy

score per ROI. FDR corrected p-values: (***) $p<0.001$. Dotted lines represent chance

level. **(B).** The embedded top panel shows neural RDMs extracted from left and right

hPT, and multi-dimensional scaling (MDS) plot visualizes the similarities of the neural

pattern elicited by 4 motion directions (arrows) and 4 sound source locations (dots).

Color codes for arrow/dots are as follows: green indicates left direction/location, red

indicates right direction/location, orange indicates up direction/location, and blue

indicates down direction/location. ML: motion left, MR: motion right, MD: motion

down, MU: motion up, SL: static left, SR: static right, SD: static down, and SU: static

up. **(C-E).** The results of representational similarity analysis (RSA) in hPT are

represented. **(C).** RDMs of the computational models that assume different similarities

of the neural pattern based on auditory motion and static conditions. **(D-E).** RSA

results for every model and each ROI. For each ROI, the dotted lines represent the

reliability of the data considering the signal-to-noise ratio (see Materials and

Methods), which provides an estimate of the highest correlation we can expect in a

given ROI when correlating computational models and neural RDMs. Error bars

indicate SEM. IM1: Intermediate models with within-axis conditions distinct, IM2:

Intermediate model with within-axis conditions combined. Each right up corner of the

bar plots show visualization of significant differences for each class of models and

hemispheres separately (Mann-Whitney-Wilcoxon rank-sum test, FDR corrected).

1243

 TABLES

Table 1. Results of the univariate analyses for the main effect of auditory motion

processing [motion > static], and auditory localization processing [static > motion].

Coordinates reported in this table are significant (p < 0.05 FWE) after correction over

small spherical volumes (SVC, 12 mm radius) of interest (#) or over the whole brain (*).

Coordinates used for correction over small spherical volumes are as follows (x, y, z, in

MNI space): left middle temporal gyrus (hMT+/V5) [-42 -64 4] (Dormal et al., 2016),

right middle temporal gyrus (hMT +/V5) [42 – 60 4] (Dormal et al., 2016), right superior

frontal sulcus [32 0 48] (Collignon et al., 2011), right middle occipital gyrus [48 -76 6]

(Collignon et al., 2011). K represents the number of voxels when displayed at p(unc)

< 0.001. L: left, R: right, G: gyrus, S: sulcus.

1255

Area	k	x (mm)	y (mm)	z (mm)	Z	p
MOTION > STATIC						
L planum temporale	10506	-46	-32	10	6.63	0.000*
L Middle Temporal G		-56	-38	14	6.10	0.000*
L Precentral G		-46	-4	52	5.25	0.004*
L Putamen		-22	2	2	4.98	0.011*
L Middle Temporal G	43	-50	-52	8	3.79	0.01#
R Superior Temporal G	7074	66	-36	12	6.44	0.000*
R Superior Temporal G		62	-2	-4	5.73	0.000*
R Superior Temporal G		52	-14	0	5.56	0.001*
R Precentral G		50	2	50	4.70	0.032*
R Superior Frontal S	159	46	0	50	4.40	0.001#
R Middle Temporal G	136	42	-60	6	4.31	0.001#
R Middle Occipital G	24	44	-62	6	3.97	0.006#

(a) Recording moving and static sounds

(b) fMRI design

One fMRI run (2 conditions x 4 directions/locations)

MOVING SOUNDS

STATIC SOUNDS

x 12 runs

Motion block

Task: Motion direction discrimination

1.25 sec

MOTION BLOCK
~(15 sec)

1 sec

1.5 sec

0.45   1.05
sec   sec

6 sec

U D R L

(c) Behavioral results inside fMRI

Behavioural Accuracy %

Motion
Static

Left   Right   Down   Up

Figure 1

Figure 2

Left hPT   # Right hPT

A. MVPA 4 Directions/Locations

B. MVPA Within/Across-Axis Directions

C. MVPA Within/Across-Axis Locations

Figure 3

Figure 4