Hemispheric asymmetry of globus pallidus relates to alpha modulation in reward-related attentional tasks

Subcortical modulation of visual alpha oscillations

C. Mazzetti[1], T. Staudigl[2], T. R. Marshall[4], J. M. Zumer[3,5,6], S. J. Fallon[4] & O. Jensen[5]

[1] Radboud University Nijmegen, Donders Institute for Brain, Cognition and Behaviour, Nijmegen 6525EN, The Netherlands

[2] Cedars-Sinai Medical Center, Department of Neurology, CA 90048, Los Angeles, United States

[3] School of Psychology, University of Birmingham, B15 2TT, Birmingham, UK;

[4] Department of Experimental Psychology, University of Oxford, OX2 6HG, Oxford, UK;

[5] Centre for Human Brain Health, School of Psychology, University of Birmingham, B15 2TT Birmingham, UK;

[6] Centre for Computational Neuroscience and Cognitive Robotics, University of Birmingham, B15 2TT, Birmingham, UK

Corresponding author: Cecilia Mazzetti, Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen, P.O. Box 9101, NL-6500 HB Nijmegen, The Netherlands; phone: +31-24-36- 65059; email: c.mazzetti@donders.ru.nl

Number of pages: 50

Word count: abstract (263), **introduction** (711), **discussion** (1654)

Number of figures: 11

Conflict of interest: The authors declare no competing financial interests

Abstract

While subcortical structures such as the basal ganglia (BG) have been widely explored in

relation to motor control, recent evidence suggests that their mechanisms extend to the

domain of attentional switching. We here investigated the subcortical involvement in reward

related top-down control of visual alpha-band oscillations (8 – 13 Hz), which have been

consistently linked to the mechanisms supporting the allocation of visual spatial attention.

Given that items associated with contextual saliency (e.g. monetary reward or loss) attract

attention, it is not surprising that alpha oscillations are further modulated by the modulated by

the acquired salience of visual items. The executive network controlling such reward-

dependent modulations of oscillatory brain activity has yet to be fully elucidated. Although

such network has been explored in terms of cortico-cortical interaction, it likely relies also on

the contribution of subcortical regions. To uncover this, we combined MRI and MEG data

from 28 male and female participants, to investigate whether derived measures of subcortical

structural asymmetries could predict interhemispheric modulation of alpha power during a

spatial attention task. We show that volumetric hemispheric lateralization of globus pallidus

(GP) and thalamus (Th) explains individual hemispheric biases in the ability to modulate

posterior alpha power. Importantly, for the GP, this effect became stronger when the value-

saliency parings in the task increased. Our findings suggest that the Th and GP in humans are

part of a subcortical executive control network, differentially involved in modulating

posterior alpha activity in the presence of saliency. Further investigation aimed at uncovering

the interaction between subcortical and neocortical attentional networks would provide useful

insight in future studies.

Significance statement

While the involvement of subcortical regions into higher level cognitive processing, such as attention and reward attribution, has been already established in previous studies, little is known about its relationship with the functional oscillatory underpinnings of said processes. In particular, interhemispheric modulation of alpha band (8-13Hz) oscillations, as recorded with magnetoencephalography (MEG), have been previously shown to vary as a function of salience (i.e. monetary reward/loss) in spatial attention task. We here provide novel insights into the link between subcortical and cortical control of visual attention, as indexed by volumetric variability of subcortical structures lateralization and individual biases in the modulation of visual alpha activity, in the context of a reward-related attentional paradigm.

Introduction

Functioning in the natural world necessitates the presence of neuronal mechanisms capable of prioritising stimuli according to their relevance (Nobre and Kastner, 2014). Deployment of attentional resources is biased towards stimuli associated with salience (e.g. monetary reward or loss), even when unrelated to the current task (Chelazzi et al., 2013). Posterior neuronal oscillations in the alpha band (8 – 13 Hz) reflect the allocation of covert attention (Worden et al., 2000; Kelly, 2006; Thut, 2006) and they have been shown to be mediated by cortico-cortical interactions (Capotosto et al., 2012a; Ptak, 2012; Vossel et al., 2014; Marshall et al., 2015a, 2015b). On the other hand, these cortical networks are further modulated by subcortical input (van Schouwenburg et al., 2010a, 2010b), whose involvement posterior oscillations remains still unclear. Previous literature has indeed linked subcortical activity to cognitive control (Cummings, 1993; Jahfari et al., 2011; Braunlich and Seger, 2013), but a direct link between these structures and alpha band oscillations has not been established. Electrophysiological activity from subcortical regions are poorly detected with magnetoencephalography (MEG). An alternative is to relate subcortical structures measured by magnetic resonance imaging (MRI) to oscillatory brain activity (Tomer et al., 2008, 2013). For instance, it has been demonstrated that individual hemispheric asymmetries in the volume of the superior longitudinal fasciculus (SLF) (Marshall et al., 2015a) relates to the individual ability to modulate posterior alpha oscillations. Importantly, subjects with greater right than left SLF volume also displayed higher modulation of posterior alpha activity in the left hemisphere, compared to the right (and vice versa). Through an analogous approach, we postulated that volumetric asymmetries of subcortical areas would be reflected by individual interhemispheric biases in the modulation of alpha oscillations during selective attention in a reward context. Basal ganglia (BG), in addition to motor control, have a well-established role in reward processing and salience attribution (Hikosaka et al., 2008, 2014; Shulman et al.,

2010a; Braunlich and Seger, 2013), and recent studies have already pointed to their functions

extending into higher level cognitive processing (Arcizet and Krauzlis, 2018). This notion

has been initially explored in animal recordings (Tremblay et al., 1998; Schultz et al., 2000;

Lauwereyns et al., 2002; Shipp, 2004; Saalmann and Kastner, 2011; Schechtman et al.,

2016), while in humans, it has recently been suggested that the BG play also a specific role in

spatial attention and selection (van Schouwenburg et al., 2010a; Tommasi et al., 2014; Van

Schouwenburg et al., 2015). Another subcortical structure playing a crucial role in cognitive

processing is the thalamus (Fiebelkorn et al., 2019; Jaramillo et al., 2019), whose nuclei are

involved in the regulation of synchronized activity in the visual cortex in relation to visual

attention and largely interact with the BG (Lopes da Silva et al., 1980; Zhou et al., 2016).

Yet, an appropriate model attempting to describe basal ganglia influence on reward-related

alpha modulation, needs to take into account the broader network of subcortical

interconnections with neighboring nuclei, such as the amygdala, in coordination with the

hippocampus (Paton et al., 2006; Zheng et al., 2017).

To this end, we made use of MEG and structural data collected in a previous study which

considered the impact of stimuli paired with value-salience on the modulation of oscillatory

brain activity in a covert attention task (Marshall et al., 2017). The participants performed a

spatial cueing task in which Chinese symbols served as targets and distractors. Prior to the

recordings, stimuli were paired with monetary rewards or losses. The authors demonstrated a

role for the saliency (salient vs neutral) of items, but not valence (positive vs negative), in

modulating posterior alpha oscillations. We here re-analysed these data with the aim to

investigate the putative role of the subcortical brain areas in biasing alpha power modulation

during attentional shifts to stimuli paired with value-saliency. MRI data of the participants

were processed in order to estimate volumetric asymmetries of subcortical areas, consistent

with methods employed in previous studies on clinical and healthy population (Womer et al.,

2014; Guadalupe et al., 2016; Okada et al., 2016). We focused on identifying the link

between individual volumetric asymmetries of subcortical areas and individual

interhemispheric bias in the ability to modulate posterior alpha oscillations. Crucially, we

further examined whether this relationship was affected by the degree of stimulus-value

associations in the task.

Materials and Methods

Participants

In the present study, we re-analysed the previously acquired dataset described in (Marshall et al., 2017), where twenty-eight healthy volunteers participated in the study (mean age: 23±2.7 years; 17 female; all right handed). All participant reported normal or corrected-to-normal vision and no prior knowledge of Chinese language. Of these, datasets from three participants were excluded from the analysis (due to respectively: technical error during acquisition, excessive eye movements during MEG recording and structural MRI data not acquired), leaving 25 participants. The experiment was conducted in compliance with the Declaration of Helsinki and was approved by the local ethics board (CMO region Arnhem-Nijmegen, CMO2001/095).

Experimental design

The experiment consisted of two phases: in the learning phase, participants were trained to memorize associations between 6 Chinese characters and 3 different values (positive, negative, neutral). Conditioning was implemented by means of visual and auditory feedback: two symbols were associated with reward (+80 cents and a 'kaching' sound), two with loss (- 80 cents and a 'buzz' sound) and two with no value (0 cents and a 'beep' sound) (see Figure 1A for an example stimulus-reward association). The stimulus-reward pairing was randomized across participants. Each trial started with the display of three fixation crosses (1000ms), followed by the presentation of a Chinese character (1000ms), together with its matching visual and auditory feedback (Figure 1B). Stimuli were displayed on a grey background, each of them was presented twelve times in a randomized order. The learning phase was conducted in a laboratory with attenuated sound and light and without MEG

recording. With the aim of reducing extinction, upon completion of this phase participants

were informed that the learnt stimulus-feedback associations would be signalling real reward

outcomes throughout the testing phase (i.e. the presentation of a Chinese character,

irrespective of its role as target or distractor, would result in a financial reward, loss or none).

After the learning phase, participants performed a testing phase (Figure 1C), when they were

required to perform a covert spatial attention tasks including the stimuli previously associated

with a monetary outcome, while ongoing electromagnetic activity was recorded with MEG.

In the testing phase, participants performed 8 blocks of 72 trials. Each trial started with the

presentation of three fixation crosses for 1000 ms (pre-trial interval), whose contrast

subsequently decreased, as a preparatory cue indicating imminent stimuli presentation. After

500ms, two symbols were presented to the left and right of the screen (8 degrees visual angle)

respectively, together with a central fixation cross flanked by two arrows, indicating the

target side. Participants were instructed to covertly attend the symbol on the cued side

('target') and to ignore the other one ('distractor'), until one of them changed contrast. The

contrast change either increased or decreased with equal probability, with onset after 750 ms

(13% trials), 1450 ms (47% trials, 'short interval trials') or 2350 ms (40%, 'long interval

trials') from stimulus presentation. Participants were asked to report the direction of the

contrast change at the targeted ('cued') location as quickly as possible by button press, using

the index or middle finger of the right hand to indicate their choice (finger-direction mapping

was randomized across participants). Participants were instructed to refrain from responding

when the distractor changed contrast. Shorter intervals of 750 ms were used to ensure that

participants would start covertly directing their attention rapidly after the cue; these trials

were not included in the analysis. The target changed contrast on 95% of the trials (valid

trials), whereas in the remaining trials the distractor did (invalid trials). The approximate

duration of the full task in the MEG was 50 minutes.

As a result of the conditioning manipulation in the learning phase, targets and distractors in

the task would be associated with either a salient (positive or negative) or a neutral value,

resulting in three categories of trials of interest, as represented by different levels of value-

salience, namely: zero (target and distractor neutral), one (target or distractor salient) or two

(target and distractor salient) value-salience levels.

MEG data acquisition

Electromagnetic brain activity was recorded from participants while seated, using a CTF 275-

channels whole-head MEG system with axial gradiometers (CTF MEG Systems, VSM

MedTech Ltd.). The data were sampled at 1200Hz, following an antialiasing filter set at

300Hz. Head position was constantly monitored throughout the experiment via online head-

localization software. This had access to the position of the three head localization coils

placed at anatomical fiducials (nasion, left and right ear), allowing, if necessary, readjustment

of the participant's position between blocks (Stolk et al., 2013). Horizontal and vertical EOG

and ECG electrodes were recorded with bipolar Ag/AgCl electrodes.

MEG data analysis

MEG data analysis was performed using the *FieldTrip* Toolbox running in MATLAB

(Oostenveld et al., 2011). Continuous data were segmented in epochs, centred at the onset of

the target contrast change, encompassing the preceding 1500 ms and the following 200 ms

(this way covering the full stimulus presentation window for short trials). A notch filter was

applied at 50, 100, 150 Hz to remove line noise, the mean was subtracted and the linear trend

removed. Automatic artifact rejection was implemented for detection and removal of trials

containing eye blinks and horizontal eye movements (detected with EOG), MEG sensor

jumps and muscle artifacts. We produced virtual planar gradiometers by computing spatial

derivatives of the magnetic signal recorded with axial gradiometers (Bastiaansen and

Knösche, 2000). The method has the advantage of improving the interpretation of the

topographic mapping since neural sources would produce a gradient field directly above

them. Time-frequency representations (TFR) of power were then calculated for the resulting

pairs of orthogonal planar gradiometers, before summing the power values at each sensor.

The analysis was performed by sliding a fixed time window of 500 ms in steps of 50 ms. The

resulting data segments were multiplied by a Hanning taper and a fast Fourier transform was

applied in the 2 – 30Hz frequency range, in steps of 2Hz. This procedure was applied only for

correct valid trials, separately for left and right cued conditions.

For each participant, TFRs were averaged across trials and a Modulation Index (MI) was

computed for each sensor k and over all time points t belonging to the time window of

interest -750 – 0 ms, according to the formula:

$$MI(f)_{k,t} = \frac{Power\ (f)_{k,t\ att\ right} - Power\ (f)_{k,t\ att\ left}}{Power\ (f)_{k,t\ att\ right} + Power\ (f)_{k,t\ att\ left}} \tag{1}$$

Where $Power(f)_{k,t\ att\ left}$ represents the power at a given frequency f in the condition

'attend left' and $Power\ (f)_{k,t\ att\ right}$ is the power of the same frequency in the condition

'attend right'. As a result, positive (or negative) MI values, at a given sensor k and given

timepoint t, indicate higher power at a given frequency f when attention was covertly directed

towards the right (or left) hemifield.

Two clusters of sensors were then derived, by selecting the twenty symmetrical occipito-

parietal sensors (i.e. ten pairs of sensors) showing the highest interhemispheric difference in

alpha modulation indices, when considering the grand average over all conditions (see Figure

2A) averaged over the previously defined time window of interest. These clusters constituted

the regions of interests (ROIs) on which subsequent analysis was focused. Subsequently, in

order to quantify individual hemispheric-specific bias with respect to modulation indices in

the alpha range (MI(α)), we calculated the Hemispheric Lateralized Modulation (HLM)

index per participant:

$$HLM(\alpha) = \frac{1}{n_{right}} \sum_{k_{right}=1}^{n_{right}} MI(\alpha)_{k_{right}} + \frac{1}{n_{left}} \sum_{k_{left}=1}^{n_{left}} MI(\alpha)_{k_{left}} \qquad (2)$$

Where k_{left} and k_{right} denote sensors belonging to the aforementioned and previously

defined left and right clusters, respectively. Please note that $MI(\alpha)_k$ indices in Eq.2 (for both

k=1, …, n_{right} and k=1, …, n_{left}) are already a result of an average over timepoints of

interest t. Since MI(α) values were obtained by subtracting alpha power in 'attend left' trials

from 'attend right' trials, and given that, as a result of attentional allocation, alpha power is

suppressed in the hemisphere contralateral to the attended hemifield, a positive HLM(α)

value indicated that a given participant displayed higher modulation of absolute magnitude of

alpha power in the right compared to the left hemisphere, and vice versa (see Figure 2B).

Analogously, Lateralized Indices (LI) of power modulation were computed for the alpha

frequency band and for each subject at the cluster level according to:

$$LI(\alpha) = \frac{1}{n_{right}} \sum_{k_{right}=1}^{n_{right}} MI(\alpha)_{k_{right}} - \frac{1}{n_{left}} \sum_{k_{left}=1}^{n_{left}} MI(\alpha)_{k_{left}} \qquad (3)$$

Also in this formula, k_{left} and k_{right} denote sensors belonging to the aforementioned left

and right clusters, respectively. Since MI(α) values were obtained by contrasting alpha power

in right versus left attention trials (see Eq.1), left hemisphere MI(α) were mostly represented

by negative values, and right hemisphere MI(α) by positive values. Consequently, higher

LI(α) indicated higher alpha lateralization for a given subject (i.e., higher interhemispheric

difference in absolute alpha modulation).

Structural data acquisition

T1-weighted images of three out of twenty-five participants were acquired on a 3 T MRI

scanner (Magnetom TIM Trio, Siemens Healthcare, Erlangen, Germany), acquisition

parameters: TR/TE= 2300/3.03 ms; FA=8°; FoV= 256 × 256 mm; slice thickness= 1 mm;

Acquisition matrix= 0×256×256×0. For the remaining participants, a 1.5T MRI scanner was

used (Magnetom AVANTO, Siemens Healthcare, Erlangen, Germany). Acquisition

parameters: TR/TE= 2250/2.95 ms; FA=15°; FoV= 256 × 256 mm; slice thickness= 1 mm;

Acquisition matrix= 0×256×256×0.

Analysis

Structural analyses were conducted using the Integrated Registration and Segmentation Tool

(FIRST) within FMRIB's Software Library (FSL) v5.0.9 (www.fmrib.ox.ac.uk/fsl/, Oxford

Centre for Functional MRI of the Brain, Oxford, UK). A standard 12 degrees of freedom

affine registration to MNI152 space was applied to individual T1 images, adjusted with

optimal sub-cortical weighting. Bayesian models implemented in the software are derived

from a training based on previous manual segmentation of 336 datasets (provided by the

Center for Morphometric Analysis (CMA, MGH, Boston) and applied to registered images to

extract subcortical volumetric outputs for left and right hemispheres (see Figure 3A).

Given the reward components of the task we then focused on regions of the basal ganglia

identified by the algorithm namely the Globus Pallidus (GP), Nucleus Accumbens (Acb),

Caudate (CN), Putamen (Pu), as well as other regions potentially involved in the

mechanisms, namely Hippocampus (Hpc), Amygdala (Am) and Thalamus (Th). To compute

hemispheric Lateralized Volume indices (LV) for each substructure of interest s, we used the

following formula, which controls for individual differences in specific subcortical volumes

via normalization by total bilateral volume, commonly employed to evaluate structural brain

asymmetries (Guadalupe et al., 2016; Okada et al., 2016):

$$LV(s) \; = \frac{V(s)_{right} - \; V(s)_{left}}{V(s)_{right} + \; V(s)_{left}} \qquad \text{(4)}$$

Where $V_{S_{right}}$ and $V_{S_{left}}$ represent respectively the anatomical right and left volumes (in

voxels) for a given substructure s. Analogously to Eq.(2), a positive (or negative) LV_S index,

in a given participant, indicated a greater right (or left) volume for a given substructure s (see

Figure 3B).

Statistical analysis

Generalized Linear Model

In order to determine the relationship between Basal Ganglia Lateralized Volumes (LV_S) and

electromagnetic indices (HLM(α)) we applied a generalized linear regression model (GLM),

specifying subcortical volumes lateralization (LV_s values) as regressors and individual HLM

values as the response vector, according to the formula:

$$HLM(\alpha) \sim \beta_0 + \beta_1 LV_{GP} + \; \beta_2 LV_{NAcc} + \; \beta_3 LV_{CN} + \beta_4 LV_{Pu} + \; \beta_5 LV_{HPC} +$$
$$\beta_6 LV_{Am} + \; \beta_7 LV_{TH} + \; \varepsilon \qquad \text{(5)}$$

All subsequent analysis on the relationship between volumetric and oscillatory data

specifically focused only on the subcortical structure(s) associated with a significant β

coefficient in the model in Eq.(5), below referred as LV_s.

Cluster based permutation test

To evaluate whether the linear association between LV_s and HLM was effectively limited to

the alpha band, a cluster based permutation approach (Maris and Oostenveld, 2007) was

employed over the full time frequency spectrum of interest. This method effectively allows to

statistically control for multiple comparisons over all time and frequency points of interest.

After selecting the *a-priori* sensors belonging to the formerly specified ROIs, we considered

a permutation distribution of regression coefficients derived from randomly pairing

participants' LV_s value (independent variable) and modulation indices (HLM(f)) 1000 times.

At every time-by-frequency point, the actual regression coefficient was evaluated against the

aforementioned distribution by means of a specified critical α value. Afterwards, a time-

frequency map of the cluster level statistics was derived showing sets of sensors associated

with a significant effect.

An equivalent approach was later applied to investigate possible hemisphere-specific

differences in alpha modulation between participants showing a right or left lateralized

substructure s. Directionality of lateralization was determined by median split of the

distribution of LV_s per participant, producing two subgroups of N=12, representing subjects

with a larger left or right volume of substructure s. After having a-priori averaged across the

time-frequency spectrum of interest ([-750 0] ms, 8-13Hz), MI(α) values at every sensor were

compared between the two subgroups (right vs left lateralized substructure). The actual t-

value was then compared with a permutation distribution of t-statistic derived from randomly

partitioning indices between the two groups 1000 times. As a result, a topography map was

plotted displaying eventual cluster(s) of sensors associated with a significant t-value (i.e. a

significant difference in MI(α) between subgroups).

Comparison between Pearson's correlation coefficients

Finally, we aimed at comparing the association between the derived structural and functional

lateralization indices in different value salience occurrences. To this end, we calculated

HLM(α) values for each participant separately for the three reward-related contingencies and

computed the Pearson's correlations with LV_S indices which displayed a significant β as

arising from the model in Eq.(5). We statistically assessed the difference in correlation

coefficients between the three experimental conditions considered, according to the method

described in (Wilcox, 2016a). The test implements a percentile resampling technique by

generating a bootstrap sample of the difference of the correlation coefficients between the

overlapping variable LV_{GP} (Y) and the two variables representing the HLM(α) for the two

experimental conditions (VO levels) to be compared (X_1, X_2). As suggested in the method,

we used a Winsorized correlation to achieve a robust measure of association between

variables. This transformation has been shown to effectively control for the influence of

outliers on the correlation estimate (Wilcox, 2016b). A confidence interval was then

computed on the resulting bootstrap distribution, to assess the statistical significance of the

actual difference between correlation coefficients describing the different VOs.

Behavioural data analysis

To assess whether subjects displayed a spatial bias during the task, we first averaged across

left and right cued trials separately, averaged across all conditions (i.e., irrespective of value-

saliency occurrences (VO)). We then employed paired t-test on the derived reaction times

(RT) and accuracy (ACC) (expressed as percentage of correct responses) measures for the

left and right cued trials. Secondly, we divided trials according to VO pairings, averaging left

and right cued trials, to determine whether behavioural performance varied as a function of

saliency in both RT and ACC. We here employed one-way repeated measures ANOVA to

assess whether group means in the three conditions significantly differ from each other. We

also considered individual lateralized measures of RT and ACC across different VO

conditions. To this end, behavioural asymmetries in performance (BA) for both measures

were calculated according to:

$$BA_{RT/ACC} = \frac{BA_{RT/ACC_{right}} - BA_{RT/ACC_{left}}}{BA_{RT/ACC_{right}} + BA_{RT/ACC_{left}}} \tag{6}$$

Where $BA_{RT\,right}$ and $BA_{RT\,left}$ represent mean reaction times for 'attend right' and 'attend left'

trials, respectively. A positive BA_{RT} for a given subject indicated faster responses when a

participant was validly cued to the left compared to the right hemisphere, while negative

values indicated the opposite pattern. Consequently, positive BA_{ACC} values indicated higher

accuracy on 'attend right' trials compared to 'left attend' trials, and vice versa.

A one-way repeated measures ANOVA was employed to test the difference across group

means in the three VO conditions examined.

In a next step, we sought to investigate the possible association of behavioural performance

with structural and functional hemispheric lateralization, we used Pearson's correlation to

examine the association of individual asymmetries in accuracy (BA_{ACC}) and reaction times

(BA_{RT}) with individual HLM(α) and LV values of subcortical structures which showed

significant correlation with HLM(α).

In a last step, we employed a general linear model (GLM) in order to assess whether spatial

biases in behavioural performance could be explained by a combination of the other

variables, namely HLM(α) and the LV indices of the subcortical areas considered, according

to the formula:

$$BA_{RT/ACC} \sim \beta_0 + \beta_1 LV_{GP} + \beta_2 LV_{NAcc} + \beta_3 LV_{CN} + \beta_4 LV_{Pu} + \beta_5 LV_{HPC} +$$

$$\beta_6 LV_{Am} + \beta_7 LV_{TH} + \beta_4 HLM(\alpha) + \varepsilon$$

(7)

Results

We acquired structural and electrophysiological data from 25 participants. Participants'

performance was tested during a covert attention paradigm, where Chinese symbols served as

targets and distractors (Figure 1). During a learning phase, prior to the actual task, the stimuli

were associated with different values (positive, negative or neutral). In the testing phase, a

central cue probed an upcoming contrast variation of the target, which appeared either at

1450 or 2350 ms, predicting its position in 95% of the trials. Participants were instructed to

indicate, with button press, the direction of the contrast change, which could either increase

or decrease with equal probability. MEG data, eye-tracking and behavioural responses were

acquired during the testing phase.

Individual variability in interhemispheric alpha power modulation

Time-frequency representations of power were calculated from MEG trials after

preprocessing and artifacts rejection. Power modulation (MI) indices were computed by

contrasting power in trials where participants were validly cued to the right (*attend right*

trials) with trials where participants were validly cued to the left (*attend left* trials) (see

Eq.(1), *Materials and Methods*).

As presented in the previously reported results (Marshall et al., 2017), we confirmed that

participants displayed a clear modulation of alpha band activity in parieto-occipital sensors

$(MI(\alpha))$: when covertly orienting attention to the cued side, alpha power decreased in the

contralateral hemisphere while it increased relatively in the ipsilateral hemisphere (Figure

2A). The magnitude of alpha power modulation, as reflected by $MI(\alpha)$, progressively

increased until the target changed contrast (Figure 2B). To best quantify the modulation, we

focused our analysis on the 750 ms interval immediately preceding the onset of the contrast

change. Next, right and left ROIs were identified as clusters of symmetric pairs of sensors

showing the highest alpha lateralization values (see *Materials and Methods*) (i.e., sensors

displaying highest interhemispheric difference in alpha modulation).

Starting from the assumption that, to a certain extent, an inter-subject variability in the ability

to modulate alpha power – in absolute value – must exist in the right compared to the left

hemisphere (and vice versa), we sought to quantify individual hemispheric biases in the

ability to modulate alpha activity. To this purpose, hemispheric lateralized modulation of

alpha power (HLM(α)) values were then computed for each participant by summing the

average MI(α) in the right and left hemisphere ROIs (see Eq.(2), *Materials and Methods*). As

a result of this computation, positive HLM(α) values would demonstrate that a given subject

was better at modulating their right, compared to left, hemisphere alpha power, while a

negative index would reflect higher ability to modulate alpha power on their left, compared to

right, hemisphere. The histogram in Figure 2B depicts the distribution of hemispheric biases

related to attentional modulation of alpha power. HLM(α) indices ranged from about -0.1 to

0.1 (i.e. a 20% variation) but they were normally distributed around zero across participants

(Shapiro Wilk, $W = .958$, $p = .392$).

Volumetric asymmetry of basal ganglia in relation to hemispheric lateralized alpha

modulation

The next step was to determine whether the biases in the ability to modulate left versus right

hemisphere alpha (HLM(α)) was related to individual hemispheric lateralization of

subcortical structures. A semi-automated segmentation tool implemented in FMRIB's

Software Library (FSL), was used to estimate volumes for the left and right subcortical and

limbic structures, namely: Globus Pallidus (GP), Nucleus Accumbens (Acb), Caudate

Nucleus (CN), Putamen (Pu), Hippocampus (Hpc), Amygdala (Am) and Thalamus (Th). We

then calculated the hemispheric lateralized volumes (LV) for each set of structures (see

Materials and Methods, Eq.(4)). Positive (or negative) LV_s values, for a given participant,

indicated whether a specific substructure *s* was larger in the right compared to the left

hemisphere (and vice versa). Further analysis revealed that, over subjects, the Acb and Th

were significantly left lateralized ($z = -3.78$, $p = 1.56 \times 10^{-4}$ and $z = -3.59$, $p = 3.28 \times 10^{-4}$,

respectively; two-sided Wilcoxon signed rank test), whilst the CN was right lateralized ($z = $

2.97, p $= .003$). For the other substructures, no significant lateralizations were identified.

In order to corroborate that the observed anatomical lateralizations where representative of

the population and not merely a fluke in the dataset, we applied the same analysis to a pool of

612 anonymized anatomical MRI scans internally available at the Donders Institute. We

hence estimated left and right volumes for the same subcortical structures considered in our

study, and derived respective LV indices (see Eq.(4)). Importantly, the same direction of

lateralization was found to be present in the in the Donders dataset as the one reported in our

dataset for all the substructures. Specifically, Th and the Acb were significantly left

lateralized ($z = -14.0$, $p = 1.4 \times 10^{-44}$ and $z= -17.04$, $p = 4.1 \times 10^{-65}$, respectively), whilst the CN

was right lateralized ($z = 13.0$, $p = 1.2 \times 10^{-38}$). In addition, the GP was found to be right

lateralized ($z = 4.0$, $p = 5.9 \times 10^{-5}$, an effect only observed as trend in our dataset), as well as

the Hpc ($z = 4.95$, $p = 7.3 \times 10^{-7}$), while the Pu was left lateralized ($z = -3.27$, p $= .001$).

The finding that subcortical lateralization biases in our sample generalize to bigger datasets is

of potential interest, but will not be further considered in the context of this study.

Given the volumetric variability in the set of substructures considered for the segmentation

protocol, we performed a cross correlation analysis between the different substructures,

including left and right volumes, to query about a potential bias in the segmentation

algorithm. No significant effects were found (positive or negative correlations, all *p-values*

>0.688; highest negative correlation $R=-.084$) indicating that, if a given structure is larger for

a given subject, this doesn't imply a bias in the segmentation protocol (e.g. to the expense of

neighbouring, allegedly smaller areas).

To investigate whether individual subcortical asymmetries (LVs, as defined in Eq. (4))

predicted differences in hemispheric lateralized modulation of alpha power (HLM(α)), we

implemented a GLM, where LV indices were included as multiple explanatory variables for

the response variable (individual HLM(α)). A significant regression was found: when

considering the grand average of all conditions, a linear combination of the subcortical LVs

was able to explain HLM(α) values ($F_{7,17} = 3.37$, $p = .019$, adjusted $R^2 = .409$). When

assessing each predictor individually, only the beta coefficients for LV_{GP} and LV_{Th} were

found to be significantly higher than zero (partial correlation: $p = .004$ and $p = .028$) (Figure

4A). Hence, when controlling for the other explanatory variables in the model, only GP and

TH asymmetry (LV_{GP}, LV_{TH}) significantly contributed to explain biases in hemispheric

lateralized alpha band modulation ($\beta = 1.768$ and $\beta = 1.924$, respectively). The independent

contribution of GP and TH lateralization is visible in Figure 4B,C showing the partial

regression plots for LV_{GP} and LV_{TH} in relation to the HLM(α) values. We conclude that

hemispheric biases in GP and TH volume are predictive of the individual abilities to

modulate left versus right hemisphere alpha. Precisely, subjects presenting a larger GP

volume in the left hemisphere compared to the right, also displayed a higher ability to

modulate alpha power (in absolute value) in the left visual hemisphere compared to the right

(and vice versa); the same association holding for the Th in relation to HLM(α) values.

Hemispheric asymmetry of globus pallidus correlates selectively with power modulation in

the alpha band

In order to better interpret the GLM results, we assessed whether the linear relationship

arising from the model was restricted to the alpha band: to this end, a non-parametric

approach was implemented to further explore the LV_{GP} and LV_{TH} in relation to HLM(α).

This method allows circumvention of the multiple comparison problem over frequency and

time points by evaluating the full low-frequency spectrum (2-30 Hz) from -750 to 0ms (Maris

and Oostenveld, 2007). We therefore conducted a cluster-based permutation test using a

dependent samples regression *t*-statistic to evaluate the effect (linear association between

$LV_{GP/TH}$ indices and HLM over all frequencies) at the sample level. A *p*-value of .05 was

chosen for thresholding the *t*-statistic of the permutation distribution and a critical value

corresponding to alpha = .025 (two tailed) was considered for the cluster-level regression test

statistic. As depicted in Figure 5A, we observed a significant cluster (*p* = .004) extending for

the entire 750 ms window of interest (i.e., when covert attention was deployed to the cued

stimulus before the contrast changed), which confirmed a positive linear association (positive

t-value) between LV_{GP} asymmetry and hemispheric lateralized modulation (HLM) of power

constrained to the alpha frequency range. When applying the same analysis to the Th

asymmetry in relation to HLM, no significant clusters of sensors were identified.

Hemisphere specific relations between alpha modulation and GP asymmetry.

Given the specific association found between LV_{GP} and HLM(α), arising from the previous

analysis, we decided to further investigate the hemisphere-specific influence of GP

volumetric asymmetry on alpha modulation indices. For this purpose, we sought to compare

average left and right hemisphere MI(α)s of participants according to the direction of GP

lateralization. This was done by means of median split of the LV_{GP} distribution, hence

resulting in two subgroups, that either had a bias towards a larger left than right GP volume

or vice versa (see *Materials and Methods*). A topographical representation of MI(α) values

together with a bar plot of MI(α) averaged across right and left hemisphere ROIs per each

subgroup is shown in Figure 6. Consistent with the GLM results, participants with a larger

right than left GP, also displayed a higher modulation of alpha band (in absolute value) in the

right hemisphere compared to the left. Given that the assumption of normality, required to

perform a mixed-effect ANOVA, was not met for the distribution of MI(α) indices in the two

subgroups, we implemented a non-parametric cluster-based permutation test to compare the

MI(α) between the two aforementioned subgroups (averaged across specific time and

frequency band of interest), employing an independent sample t-test score, and then

comparing it with the resulting permutation distribution. This allowed us to explore whether

there was a hemispheric-specific difference in the two subgroups in the extent of absolute

alpha modulation. The test indicated a significant cluster of sensors over right posterior

channels (p= .027), hence including the previously defined right ROI and denoting a

significant difference in the right hemisphere absolute alpha modulation (MI(α)) between the

two subgroups. These results might suggest that the linear association arising from the GLM

(Figure 4A, B) in relation to the association between LV_{GP} and HLM(α), was largely driven

by right hemisphere alpha modulation. We conducted the analogous analysis performing a

median split of the distribution of LV_{TH} indices. In that case we did not find interhemispheric

dominance in alpha modulation indices related to lateralization of the thalamus in the right

compared to left hemisphere.

The involvement of globus pallidus and thalamus in relation to stimulus-value associations

Crucially, we aimed at assessing whether the level of value-saliency occurrences (VO) in a

given trial influenced the association between the structural and functional lateralization

indices arisen from the GLM. We first calculated HLM(α) (see Eq.(2), *Materials and*

Methods) values for each participant, separately for the three VO levels, namely two, one and

zero value saliency occurrences (see *Materials and Methods*). We then examined Pearson

correlations between HLM(α) and LV values for both GP and Th, which showed a positive

significant β in the model, across the three levels considered (Figure 7, Figure 8). LV_{GP}

significantly correlated with HLM(α) only in trials where both target and distractors had

value-salience (two VO) ($r = .68$, $p=1.75\times10^{-4}$; Figure 7A). This denotes that, in trials with

two value-salient items presented, participants exhibiting a right lateralized GP volume, also

displayed a stronger alpha modulation in the right compared to the left hemisphere, and vice

versa. LV_{GP} did not significantly correlate with HLM(α) when only one or none of the

stimuli presented were associated with a salient value ($p = .144$ and $p = .314$, respectively;

Figure 7A).

In order to statistically quantify the influence of the stimulus-value association on the

relationship between LV_{GP} and HLM(α), we compared *robust correlations* in the three

conditions according to the bootstrap method described in (Wilcox, 2016c) for dependent

overlapping correlations (see *Materials and Methods*). The correlation between LV_{GP} and

HLM(α) in trials with two occurrences of value-salience, significantly differed both from the

condition characterized by one (95% CI [.106, .672]) and zero occurrences (95% CI [.125,

.897]). This confirmed that the association between lateralized GP volume and alpha

modulation bias significantly increased as a function of the number of value-salient

occurrences in the task (Figure 7B). We performed the same analysis in order to assess

whether value-saliency occurrences mediated also the association between LV_{TH} and

HLM(α). When considering the correlation indices in the three conditions separately, no

significant linear relationship was found between the two indices (Figure 8A). Also in this

case, when comparing robust correlations between the three conditions, according to the

same method above, no significant difference was found. This suggested that the relationship

between thalamus volumetric lateralization and alpha modulation arising from the model in

Eq.(5), was not driven by the number of value-salient occurrences in the task.

 Behavioural analysis

 Saliency occurrences in the task stimuli didn't affect behavioural responses

At the behavioural level, we sought to investigate whether subjects displayed a spatial bias in

task performance, irrespective of the value-saliency levels. To this end we performed a paired

t-test to assess whether participants' performance differed between left and right cued trials,

in both reaction times (RT) and accuracy measures. No behavioural spatial bias was found

neither in RT (p=.341) nor in accuracy (p=.572) values. Secondly, we examined whether

value-salient occurrences (VO) levels, modulated participants' behavioural performance.

We then compared mean RT and accuracy for the three VOs levels (see *Materials and*

Methods). There were no statistically significant differences between the three groups, as

determined by one-way ANOVA, in RT ($F_{(2,72)}$ =.004, p=.995) (Figure 8A) nor in accuracy

($F_{(2,72)}$=.003, p=.996) (Figure 9C). We then tested whether a behavioural spatial bias occurred

across value saliency occurrences (i.e., whether subjects displayed a difference in RT or

accuracy asymmetry across VOs). We computed measures of behavioural asymmetry in

accuracy (BA_{ACC}) and reaction times (BA_{RT}) (see Eq.(6), *Materials and Methods*).

Analogously to the method used to compute $HLM(\alpha)$, we created asymmetry indices for

every subject by contrasting behavioural measures for *attend right* with *attend left* trials. As

such, a positive BA_{RT} would indicate that subjects were faster when cued to the left compared

to the right hemisphere, and vice versa. Similarly, positive BA_{ACC} indices reflected higher

accuracy when cued to the right compared to the left hemisphere. With the method

aforementioned, we performed a one-way ANOVA to assess whether a significant difference

in behavioural bias occurred across the three VO conditions. Neither BA_{RT} nor BA_{ACC} values

significantly differed across VOs ($F_{(2,72)}$=.191, p=.826 and $F_{(2,72)}$=.669, p=.515, respectively)

(Figure 9B, 9D). With the aim of determining a potential link between lateralized indices of

behavioural performance and the anatomical (LVs) and functional ($HLM(\alpha)$) lateralization

indices of interest, we employed three separate GLMs to assess whether a linear combination

of BA_{RT} and BA_{ACC} values could explain LV_{GP}, LV_{TH} and/or $HLM(\alpha)$ indices. Neither LV_{GP}

nor LV_{TH} could be explained by the behavioural lateralized measures ($F_{1,23}=.18$, $p=.834$,

adjusted $R^2=-.07$ and $F_{1,23}=.16$, $p=.849$, adjusted $R^2=-.07$). The same result held for the

prediction of $HLM(\alpha)$, yielding also in this case no significant regression coefficients

($F_{1,23}=1.17$, $p=.33$, adjusted $R^2=-.01$).

Last, we investigated whether individual behavioural spatial biases could be accounted for by

a combination of the other measures examined. To this end, we considered all subcortical

LV_S and $HLM(\alpha)$ indices and specified them as regressors in a general linear model (see

Eq.(7), *Materials and Methods*), in order to determine whether they could explain biases in

RT and accuracy (BA_{RT} and BA_{ACC}). No significant regression was found which could

account for BA_{RT} indices ($F_{8,16}=.85$, $p=.570$, adjusted $R^2=-.05$) nor for BA_{ACC} indices

($F_{8,16}=1.07$, $p=.429$, adjusted $R^2=-.023$, respectively). (Figure 10A, B).

Stronger alpha lateralization corresponded to faster reaction times in neutral trials

Given that the lack of a relationship between spatial bias in task performance and functional

and structural hemispheric lateralization indices is likely explained by the orthogonalization

of attentional orienting and stimulus-value associations, we further investigated these

potential effects in trials which did not require saliency processing (0 V.O.s).

To this end, we performed a trial-based analysis, by first grouping, for each subject, fast and

slow trials, with respect to the reaction times distribution median. This was done separately

for left-cued and right-cued (valid) trials. For each subject, we then computed alpha

modulation indices ($MI(\alpha)$) of derived fast and slow trials, according to Eq.1 (see Materials

and Methods). Next, we averaged $MI(\alpha)$s in slow and fast trials across subjects. Figure 11A

shows the topographical representation of $MI(\alpha)$s values for the two trial groups (fast versus

slow trials). Figure11B, shows mean values for MI(α), averaged over left and right ROIs in

the two subgroups. To statistically assess the difference in alpha lateralization between the

two subgroups, we compared LI(α)s (according to Eq.(3)) between fast and slow trials, by

means of dependent sample t-test. This revealed that, on average, subjects displayed a

stronger alpha lateralization in fast trials, as compared to slow trials (p=.038), when no

saliency processing was required (0 V.O. trials).

Discussion

The aim of this study was to investigate the involvement of subcortical structures in modulating spatial attention to stimuli associated with contextual salience. We observed that volumetric lateralization of subcortical areas explained individual differences in the ability to modulate interhemispheric alpha power. Specifically, participants exhibiting a right lateralized globus pallidus (GP) also had a better ability to modulate posterior alpha oscillations in the right compared to left hemisphere, and vice versa. The same association held for the relationship between thalamic (Th) hemispheric asymmetry and alpha modulation. Importantly, only the correlation between GP and alpha hemispheric lateralized modulation increased as a function of value-saliency occurrences in the task. To the best of our knowledge, this is the first finding relating individual volumetric differences in BG and thalamus to the modulation of posterior alpha oscillations.

Subcortical areas and alpha synchronization

Our first finding is in line with a growing body of literature demonstrating a subcortical involvement in high level cognitive functions, such as conscious perception (Slagter et al., 2016), working memory performance (Frank et al., 2001), cognitive control (Reilly et al., 2011; Ceaser and Barch, 2015; Piray et al., 2016) and attentional control (Yantis et al., 2012; Tommasi et al., 2015). We showed that volumetric asymmetry of subcortical areas predicts individual biases in the ability to efficiently allocate attention, as indexed by interhemispheric modulation of alpha power. This is strong support in favour of a subcortical involvement in attentional processing, given the well-established role of alpha oscillations in the allocation of spatial visuospatial attention (Jensen and Mazaheri, 2010). In line with the functional association between BG and cognitive control in the context of reward (Fallon and Cools, 2014; Fallon et al., 2017), we provide novel insights into the involvement of subcortical

regions in the modulation of posterior alpha oscillations. We here hypothesized that a larger

subcortical volume would reflect higher control over alpha oscillations in a hemispheric-

specific fashion.

Pulsed inhibition

A well-recognized function of the BG is to inhibit or promote cortical activity via

GABAergic signalling, through the globus pallidus pars interna (GPi), one of its major output

structures (Lanciego et al., 2012; Goldberg et al., 2013). The BG might exercise its influence

by applying control over activity in the prefrontal cortex or it might directly coordinate

posterior regions (as reflected by its relationship to alpha power modulation during reward

processing). Our results suggest that individual differences in GP volume lateralization may

correspond to interhemispheric variability in GABAergic signalling and thus reflect the

subcortical potential to inhibit cortex. This input is likely responsible for producing the

mechanisms of '*pulsed inhibition*' in the visual cortex (Jensen and Mazaheri, 2010), reflected

by interhemispheric modulation of alpha power, allowing the selective processing of stimuli.

Implicitly, we assumed that the volume of the GP indirectly reflects its ability to exert its top-

down control over posterior areas, its size possibly representing a determinant for the number

of GABAergic neurons involved in the control mechanism.

GP in relation to attentional selection and cognitive control

Interestingly, our results emphasize the specific contribution of the GP in supporting

stimulus-driven allocation of attention in a value-based context. The GPi is considered to

mediate the output of the BG (Lanciego et al., 2012; Goldberg et al., 2013), and previous

literature has implicated this structure in voluntary movement regulation: its functions have

indeed been predominantly investigated in clinical and animal models in association with

motor functions and action control (Filion and Tremblay, 1991; Jahfari et al., 2011),

describing, for instance, reduction of hypokinetic and rigidity symptoms following

pallidotomy in humans (Schuurman et al., 1997; Dostrovsky et al., 2002). Nevertheless,

recent results from single unit recordings in humans provided indications that

electrophysiological activity in the GPi reflects processing of stimuli associated with different

reward contingencies (Howell et al., 2016). This is corroborated by evidence of alterations of

cognitive, in addition to motor, abilities, following pallidotomy in Parkinson's disease (PD)

patients (Lombardi et al., 2000); In addition, GPi DBS in the treatment of PD has been

reported to be associated with several cognitive impairments, such as subtle declines in

attention and concentration, although to a lesser extent when compared to subthalamic (STN)

DBS (Combs et al., 2015). This aspect has been further addressed in clinical studies showing

a link between PD, associated with abnormal pallidal activity (Dostrovsky et al., 2002;

Rosenberg-Katz et al., 2016), and altered reward processing as well as updating (Aarts et al.,

2012; Chong et al., 2015). Structural GP abnormalities have also been linked to impaired

suppression of distractors in ADHD (Aylward et al., 1996; Qiu et al., 2009) and psychotic

symptoms in schizophrenia (Hokama et al., 1995; Spinks et al., 2005; Mamah et al., 2007),

which has been related to aberrant salience attribution and reward learning (Early et al., 1987;

Okada et al., 2016). As an important output component of the reward circuit (Haber, 2011),

the GPi might serve to indirectly influence the cortical information flow by biasing selective

processing of value-related stimuli. Our data expands on this notion by suggesting a further

pallidal influence on the modulation of visual alpha oscillations. Importantly, given that the

association between LV_{GP} and $HLM(\alpha)$ increases as a function of the saliency, we here

postulate a specific role for the GP in reward-related shifts of attention. On the other hand, it

is still unclear to what extent this modulation is dependent on reward-related stimuli rather

than covert visual attention: additional studies would be valuable to further disentangle the

role of these two features and generalize the findings.

Right lateralization of the association between GP and alpha modulation

Notably, the association between GP lateralization and interhemispheric alpha power (Figure

6) was largely related to right hemisphere differences in absolute alpha modulation between

subjects exhibiting a right, as compared to left, lateralized GP volume. This finding possibly

reflects the right hemisphere dominance allegedly characterizing spatial attention processes

(Shulman et al., 2010b), corroborated by the right lateralized feature of the ventral attentional

network, which has been described as specifically involved in the processing of behaviourally

salient stimuli (Corbetta and Shulman, 2002)

Differential role of GP and Th in relation to posterior alpha modulation

Our results show that GP and Th lateralizations were related to the interhemispheric bias in

alpha modulation during selective allocation of attention. However, only GP lateralization

was related to the value-saliency pairings in the task. The different contribution from GP and

Th in relation to saliency occurrences might likely reflect different roles of the two areas in

the top-down control of attentional processing. The GP provides a modulatory signal related

to the processing of stimuli that draw attention due to their strong saliency associations. The

perceptual competition resulting from attending to a salient target whilst required to suppress

an equally salient distractor, might be resolved by a network involving the GP. Increased

midbrain activity has indeed been shown to accompany attentional suppression of a highly

rewarding distractor carrying a strong perceptual competition with the target (Gong et al.,

2017), suggesting that dopaminergic networks might flexibly modulate attentional selection

in reward-related contexts.

The correlation between Th lateralization and attention-related alpha modulation, which was

irrespective of the saliency component in the current task, is in line with the evidence that

thalamic activity, particularly arising from its largest nucleus, the pulvinar, modulates the

alpha rhythm in extended visual areas (Lopes da Silva et al., 1980; Wilke et al., 2009;

Saalmann et al., 2012; Zhou et al., 2016; Green et al., 2017). The pulvinar was first shown to

contribute to the generation of the posterior alpha rhythm in dogs (Lopes da Silva et al.,

1980) and also to regulate synchronized activity between visual cortical areas to support the

allocation of attention in human and nonhuman primates (Petersen et al., 1987; Kastner et al.,

2004; Saalmann et al., 2012; Green et al., 2017). Our findings, therefore, add to the growing

body of evidence suggesting that thalamo-cortical interactions play a fundamental role in

shaping cognitive processing (Saalmann and Kastner, 2011; Sherman, 2016; Green et al.,

2017; Halassa and Kastner, 2017; Fiebelkorn and Kastner, 2019). With regard to thalamic

regulation of interhemispheric alpha power, it is important to mention that our interpretation

is limited by the pragmatic difficulty in reliably estimating pulvinar volume by means of the

automatic segmentation algorithm. Yet, an appropriate model attempting to describe basal

ganglia influence on reward-related alpha modulation, needs to take into account the broader

network of basal ganglia interconnections with other subcortical nuclei, such as the thalamus.

Pallido-cortical pathways

Through which route does the GP influence visual alpha oscillations? A possibility is that the

GP modulates prefrontal activity which in turn engages and affects dorsal attentional

networks (Cummings, 1993; Pauls et al., 2014). The dorsal attention network, with the

intraparietal sulcus (IPS) and frontal eye-fields (FEF) as its major hubs, has been suggested to

mediate top-down allocation of attention. Both the IPS and FEF have been indeed causally

implicated in the control over posterior alpha oscillations in relation to attentional shifts

(Corbetta and Shulman, 2002; Capotosto et al., 2012b; Ptak, 2012; Vossel et al., 2014;

Marshall et al., 2015a). With our results, we propose the existence of a network which allows

salience driven signals from the BG to influence the prefrontal cortex in biasing the

competition among posterior regions. The idea of a BG-cortico loop involved in stimulus

driven reorienting of attention has been already introduced (Alexander, 1986; Shulman et al.,

2010a) and is consistent with the notion of a 'salience network', which integrates

behaviourally relevant input in order to bias and guide cognitive control (Seeley et al., 2007;

Metzger, 2010; Chen et al., 2015; Peters et al., 2016). Within this framework, the BG,

through their main output via the GPi, are thought to influence the connectivity between

frontoparietal regions by updating goal-directed behaviour, in order to adapt to changes in the

environment (van Schouwenburg et al., 2010b).

The influence of GP on posterior alpha oscillations could further be mediated through

indirect projections via the thalamus. The major target of GPi projections is the motor

thalamus, including ventrolateral and ventral anterior thalamic nuclei, which innervates

motor and premotor cortex (Herrero et al., 2002; Sommer, 2003; Goldberg et al., 2013).

However, cortical projections from thalamic nuclei receiving input from the BG might be

more diverse and target also prefrontal areas (McFarland and Haber, 2002), which would

enable an indirect modulation of frontoparietal networks by the GPi via the thalamus.

Additionally, intra-thalamic connectivity (Crabtree et al., 1998; Crabtree and Isaac, 2002) as

well as complex interactions between the thalamic reticular nucleus and thalamic nuclei

(Guillery et al., 1998; Halassa and Acsády, 2016) may provide multiple alternative pathways

to convey influence of the GPi on cortical areas and modulate behaviour (Haber and

Calzavara, 2009).

The proposed models provide a theoretical framework in favour of a flexible subcortical

modulation of top-down regulation of attentional allocation, which for the GP appears to be

specifically involved in tasks involving value-saliency processing. Nevertheless, the

aforementioned possible modulatory routes should not be considered as mutually exclusive: a

more comprehensive model of attentional control should instead account for multiple cortical

and subcortical pathways operating in parallel, which would allow optimization of the

organism's interaction with the environment.

Data availability

The preprocessed MEG and MRI anonymised datasets that support the findings of this study

are available as downloadable online data collection in the Donders Data Repository

(https://data.donders.ru.nl), with persistent identifier: 11633/di.dccn.DSC_3016045.01_337,

upon reasonable request to the corresponding author.

Acknowledgements

The authors gratefully acknowledge the support of the Netherlands Organisation for

Scientific Research (NWO, VICI grants 453-09-002 and 453-14-015 and the James S.

McDonnell Foundation (grants 220020328 and 220020448). We also would like to thank

Sebastiaan den Boer for his in contribution in the data collection process and experimental

design.

References

Aarts E, Helmich RC, Janssen MJR, Oyen WJG, Bloem BR, Cools R (2012) Aberrant reward

processing in Parkinson's disease is associated with dopamine cell loss. Neuroimage

59:3339–3346.

Alexander G (1986) Parallel Organization of Functionally Segregated Circuits Linking Basal

Ganglia and Cortex. Annu Rev Neurosci 9:357–381.

Arcizet F, Krauzlis RJ (2018) Covert spatial selection in primate basal ganglia. PLoS Biol

16:e2005930.

Aylward EH, Reiss AL, Reader MJ, Singer HS, Brown JE, Denckla MB (1996) Basal ganglia

volumes in children with attention-deficit hyperactivity disorder. J Child Neurol 11:112–

115.

Bastiaansen MCM, Knösche TR (2000) Tangential derivative mapping of axial MEG applied

to event-related desynchronization research. Clin Neurophysiol 111:1300–1305.

Bischoff-Grethe A, Ozyurt IB, Busa E, Quinn BT, Fennema-Notestine C, Clark CP, Morris

S, Bondi MW, Jernigan TL, Dale AM, Brown GG, Fischl B (2007) A technique for the

deidentification of structural brain MR images. Hum Brain Mapp 28:892–903.

Braunlich K, Seger C (2013) The basal ganglia. Wiley Interdiscip Rev Cogn Sci 4:135–148.

Capotosto P, Corbetta M, Romani GL, Babiloni C (2012a) Electrophysiological correlates of

stimulus-driven reorienting deficits after interference with right parietal cortex during a

spatial attention task: a TMS-EEG study. J Cogn Neurosci 24:10.1162/jocn_a_00287.

Capotosto P, Corbetta M, Romani GL, Babiloni C (2012b) Electrophysiological correlates of

stimulus-driven reorienting deficits after interference with right parietal cortex during a

spatial attention task: a TMS-EEG study. J Cogn Neurosci 24:10.1162/jocn_a_00287.

Ceaser AE, Barch DM (2015) Striatal Activity is Associated with Deficits of Cognitive

Control and Aberrant Salience for Patients with Schizophrenia. Front Hum Neurosci
9:687.

Chelazzi L, Perlato A, Santandrea E, Della Libera C (2013) Rewards teach visual selective
attention. Vision Res 85:58–62.

Chen MC, Ferrari L, Sacchet MD, Foland-Ross LC, Qiu MH, Gotlib IH, Fuller PM, Arrigoni
E, Lu J (2015) Identification of a direct GABAergic pallidocortical pathway in rodents.
Eur J Neurosci 41:748–759.

Chong TTJ, Bonnelle V, Manohar S, Veromann KR, Muhammed K, Tofaris GK, Hu M,
Husain M (2015) Dopamine enhances willingness to exert effort for reward in
Parkinson's disease. Cortex 69:40–46.

Combs HL, Folley BS, Berry DTR, Segerstrom SC, Han DY, Anderson-Mooney AJ, Walls
BD, van Horne C (2015) Cognition and Depression Following Deep Brain Stimulation
of the Subthalamic Nucleus and Globus Pallidus Pars Internus in Parkinson's Disease: A
Meta-Analysis. Neuropsychol Rev 25:439–454.

Corbetta M, Shulman GL (2002) Control of Goal-Directed and Stimulus-Driven Attention in
the Brain. Nat Rev Neurosci 3:215–229.

Crabtree JW, Collingridge GL, Isaac JTR (1998) A new intrathalamic pathway linking
modality-related nuclei in the dorsal thalamus. Nat Neurosci 1:389–394.

Crabtree JW, Isaac JTR (2002) New intrathalamic pathways allowing modality-related and
cross-modality switching in the dorsal thalamus. J Neurosci 22:8754–8761.

Cummings J (1993) Frontal-subcortical circuits and human behavior. Arch Neurol 50:873–
880.

Dostrovsky JO, Hutchison WD, Lozano AM (2002) The Globus Pallidus , Deep Brain
Stimulation , and Parkinson ' s Disease. Neurosci 8:284–290.

Early TS, Reiman EM, Raichle ME, Spitznagel EL (1987) Left globus pallidus abnormality

in never-medicated patients with schizophrenia. Proc Natl Acad Sci U S A 84:561–563.

Fallon SJ, Cools R (2014) Reward acts on the pFC to enhance distractor resistance of

working memory representations. J Cogn Neurosci 26:2812–2826.

Fallon SJ, Zokaei N, Norbury A, Manohar SG, Husain M (2017) Dopamine Alters the

Fidelity of Working Memory Representations according to Attentional Demands. J

Cogn Neurosci 29:728–738.

Fiebelkorn IC, Kastner S (2019) The Puzzling Pulvinar. Neuron 101:201–203.

Fiebelkorn IC, Pinsk MA, Kastner S (2019) The mediodorsal pulvinar coordinates the

macaque fronto-parietal network during rhythmic spatial attention. Nat Commun 10.

Filion M, Tremblay L (1991) Abnormal spontaneous activity of globus pallidus neurons in

monkeys with MPTP-induced parkinsonism. Brain Res 547:140–144.

Frank MJ, Loughry B, O'Reilly RC (2001) Interactions between frontal cortex and basal

ganglia in working memory: A computational model. Cogn Affect Behav Neurosci

1:137–160.

Goldberg JH, Farries MA, Fee MS (2013) Basal ganglia output to the thalamus: still a

paradox. Trends Neurosci 36:10.1016/j.tins.2013.09.001.

Gong M, Jia K, Li S (2017) Perceptual Competition Promotes Suppression of Reward

Salience in Behavioral Selection and Neural Representation. J Neurosci 37:6242–6252.

Green JJ, Boehler CN, Roberts KC, Chen L-C, Krebs RM, Song AW, Woldorff MG (2017)

Cortical and Subcortical Coordination of Visual Spatial Attention Revealed by

Simultaneous EEG–fMRI Recording. J Neurosci 37:7803–7810.

Guadalupe T et al. (2016) Human subcortical brain asymmetries in 15,847 people worldwide

reveal effects of age and sex. Brain Imaging Behav:1–18.

Guillery RW, Feig SL, Lozsádi D a., Lozsadi DA, Lozsádi D a. (1998) Paying attention to the thalamic reticular nucleus. Trends Neurosci 21:28–32.

Haber S (2011) Neuroanatomy of Reward: A View from the Ventral Striatum. In: Neurobiology of Sensation and Reward (Jay A. Gottfried, ed). CRC Press.

Haber SN, Calzavara R (2009) The cortico-basal ganglia integrative network: The role of the thalamus. Brain Res Bull 78:69–74.

Halassa MM, Acsády L (2016) Thalamic Inhibition: Diverse Sources, Diverse Scales. Trends Neurosci 39:680–693.

Halassa MM, Kastner S (2017) Thalamic functions in distributed cognitive control. Nat Neurosci 20:1669–1679.

Herrero MT, Barcia C, Navarro JM (2002) Functional anatomy of thalamus and basal ganglia. Child's Nerv Syst 18:386–404.

Hikosaka O, Bromberg-Martin E, Hong S, Matsumoto M (2008) New insights on the subcortical representation of reward. Curr Opin Neurobiol 18:203–208.

Hikosaka O, Kim HF, Yasuda M, Yamamoto S (2014) Basal Ganglia Circuits for Reward Value–Guided Behavior. Annu Rev Neurosci 37:289–306.

Hokama H, Shenton ME, Nestor PG, Kikinis R, Levitt JJ, Metcalf D, Wible CG, O'Donnella BF, Jolesz FA, McCarley RW (1995) Caudate, putamen, and globus pallidus volume in schizophrenia: A quantitative MRI study. Psychiatry Res Neuroimaging 61:209–229.

Howell NA, Prescott IA, Lozano AM, Hodaie M, Voon V, Hutchison WD (2016) Preliminary evidence for human globus pallidus pars interna neurons signaling reward and sensory stimuli. Neuroscience 328:30–39.

Jahfari S, Waldorp L, van den Wildenberg WPM, Scholte HS, Ridderinkhof KR, Forstmann

BU (2011) Effective Connectivity Reveals Important Roles for Both the Hyperdirect

(Fronto-Subthalamic) and the Indirect (Fronto-Striatal-Pallidal) Fronto-Basal Ganglia

Pathways during Response Inhibition. J Neurosci 31:6891–6899.

Jaramillo J, Mejias JF, Wang XJ (2019) Engagement of Pulvino-cortical Feedforward and

Feedback Pathways in Cognitive Computations. Neuron 101:321-336.e9.

Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity:

gating by inhibition. Front Hum Neurosci 4:186.

Kastner S, O'Connor DH, Fukui MM, Fehd HM, Herwig U, Pinsk MA (2004) Functional

imaging of the human lateral geniculate nucleus and pulvinar. J Neurophysiol 91:438–

448.

Kelly SP (2006) Increases in Alpha Oscillatory Power Reflect an Active Retinotopic

Mechanism for Distracter Suppression During Sustained Visuospatial Attention. J

Neurophysiol 95:3844–3851.

Lanciego JL, Luquin N, Obeso JA (2012) Functional neuroanatomy of the basal ganglia.

Cold Spring Harb Perspect Med 2:1–20.

Lauwereyns J, Takikawa Y, Kawagoe R, Kobayashi S, Koizumi M, Coe B, Sakagami M,

Hikosaka O (2002) Feature-based anticipation of cues that predict reward in monkey

caudate nucleus. Neuron 33:463–473.

Lombardi WJ, Gross RE, Trepanier LL, Lang AE, Lozano AM, Saint-Cyr JA (2000)

Relationship of lesion location to cognitive outcome following microelectrode-guided

pallidotomy for Parkinson's disease: support for the existence of cognitive circuits in the

human pallidum. Brain 123:746–758.

Lopes da Silva FH, Vos JE, Mooibroek J, van Rotterdam A (1980) Relative contributions of

intracortical and thalamo-cortical processes in the generation of alpha rhythms, revealed

by partial coherence analysis. Electroencephalogr Clin Neurophysiol 50:449–456.

Mamah D, Wang L, Barch D, de Erausquin GA, Gado M, Csernansky JG (2007) Structural

analysis of the basal ganglia in schizophrenia. Schizophr Res 89:59–71.

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. J

Neurosci Methods 164:177–190.

Marshall TR, Bergmann TO, Jensen O (2015a) Frontoparietal Structural Connectivity

Mediates the Top-Down Control of Neuronal Synchronization Associated with Selective

Attention. PLoS Biol 13:1–17.

Marshall TR, den Boer S, Cools R, Jensen O, Fallon SJ, Zumer JM (2017) Occipital Alpha

and Gamma Oscillations Support Complementary Mechanisms for Processing Stimulus

Value Associations. J Cogn Neurosci:1–11.

Marshall TR, O'Shea J, Jensen O, Bergmann TO (2015b) Frontal Eye Fields Control

Attentional Modulation of Alpha and Gamma Oscillations in Contralateral

Occipitoparietal Cortex. J Neurosci 35:1638–1647.

McFarland NR, Haber SN (2002) Thalamic relay nuclei of the basal ganglia form both

reciprocal and nonreciprocal cortical connections, linking multiple frontal cortical areas.

J Neurosci 22:8117–8132.

Metzger CD (2010) High field fMRI reveals thalamocortical integration of segregated

cognitive and emotional processing in mediodorsal and intralaminar thalamic nuclei.

Front Neuroanat 4:1–17.

Nobre AC (Kia), Kastner S (2014) The Oxford handbook of attention. Oxford: Oxford

University Press.

Okada N et al. (2016) Abnormal asymmetries in subcortical brain volume in schizophrenia.

Mol Psychiatry 21:1–7.

Oostenveld R, Fries P, Maris E, Schoffelen JM (2011) FieldTrip: Open source software for

advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell

Neurosci 2011.

Paton JJ, Belova MA, Morrison SE, Salzman CD (2006) The primate amygdala represents

the positive and negative value of visual stimuli during learning. Nature 439:865–870.

Pauls DL, Abramovitch A, Rauch SL, Geller DA (2014) Obsessive–compulsive disorder: an

integrative genetic and neurobiological perspective. Nat Rev Neurosci 15:410–424.

Peters SK, Dunlop K, Downar J (2016) Cortico-Striatal-Thalamic Loop Circuits of the

Salience Network: A Central Pathway in Psychiatric Disease and Treatment. Front Syst

Neurosci 10:1–23.

Petersen SE, Robinson DL, Morris JD (1987) Contributions of the pulvinar to visual spatial

attention. Neuropsychologia 25:97–105.

Piray P, Toni I, Cools R (2016) Human Choice Strategy Varies with Anatomical Projections

from Ventromedial Prefrontal Cortex to Medial Striatum. J Neurosci 36:2857–2867.

Ptak R (2012) The Frontoparietal Attention Network of the Human Brain. Neurosci 18:502–

515.

Qiu A, Crocetti D, Adler M, Mahone EM, Denckla MB, Miller MI, Mostofsky SH (2009)

Basal ganglia volume and shape in children with attention deficit hyperactivity disorder.

Am J Psychiatry 166:74–82.

Reilly RCO, Herd SA, Pauli WM (2011) Computational Models of Cognitive Control.

Psychology 20:257–261.

Rosenberg-Katz K, Herman T, Jacob Y, Kliper E, Giladi N, Hausdorff JM (2016) Subcortical

Volumes Differ in Parkinson's Disease Motor Subtypes: New Insights into the

Pathophysiology of Disparate Symptoms. Front Hum Neurosci 10:1–9.

Saalmann YB, Kastner S (2011) Cognitive and Perceptual Functions of the Visual Thalamus.

Neuron 71:209–223.

Saalmann YB, Pinsk MA, Wang L, Li X, Kastner S (2012) The Pulvinar Regulates

Information Transmission Between Cortical Areas Based on Attention Demands.

Science (80-) 337:753–756.

Schechtman E, Noblejas MI, Mizrahi AD, Dauber O, Bergman H (2016) Pallidal spiking

activity reflects learning dynamics and predicts performance. Proc Natl Acad Sci

113:E6281–E6289.

Schultz W, Tremblay L, Hollerman JR, Schultz (2000) Reward processing in primate

orbitofrontal cortex and basal ganglia. Cereb Cortex 10:272–284.

Schuurman PR, de Bie RMA, Speelman JD, Bosch DA (1997) Posteroventral Pallidotomy in

Movement Disorders. In: Advances in Stereotactic and Functional Neurosurgery 12:

Proceedings of the 12th Meeting of the European Society for Stereotactic and Functional

Neurosurgery, Milan 1996 (Ostertag CB, Thomas DGT, Bosch A, Linderoth B, Broggi

G, eds), pp 14–17. Vienna: Springer Vienna.

Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, Reiss AL, Greicius

MD (2007) Dissociable Intrinsic Connectivity Networks for Salience Processing and

Executive Control. J Neurosci 27:2349–2356.

Sherman SM (2016) Thalamus plays a central role in ongoing cortical functioning. Nat

Neurosci 19:533–541.

Shipp S (2004) The brain circuitry of attention. Trends Cogn Sci 8:223–230.

Shulman GL, Astafiev S V, Franke D, Pope DLW, Abraham Z, Mcavoy MP, Corbetta M

(2010a) Interaction of stimulus-driven reorienting and expectation in ventral and dorsal

fronto-parietal and basal ganglia-cortical networks. 29:4392–4407.

Shulman GL, Pope DLW, Astafiev S V., McAvoy MP, Snyder AZ, Corbetta M (2010b)

Right Hemisphere Dominance during Spatial Selective Attention and Target Detection

Occurs Outside the Dorsal Frontoparietal Network. J Neurosci 30:3640–3651.

Slagter HA, Mazaheri A, Reteig LC, Smolders R, Figee M, Mantione M, Schuurman PR,

Denys D (2016) Contributions of the ventral striatum to conscious perception: An

intracranial EEG study of the attentional blink. J Neurosci 37:1081–1089.

Sommer MA (2003) The role of the thalamus in motor control. Curr Opin Neurobiol 13:663–

670.

Spinks R, Nopoulos P, Ward J, Fuller R, Magnotta VA, Andreasen NC (2005) Globus

pallidus volume is related to symptom severity in neuroleptic naive patients with

schizophrenia. Schizophr Res 73:229–233.

Stolk A, Todorovic A, Schoffelen JM, Oostenveld R (2013) Online and offline tools for head

movement compensation in MEG. Neuroimage 68:39–48.

Thut G (2006) -Band Electroencephalographic Activity over Occipital Cortex Indexes

Visuospatial Attention Bias and Predicts Visual Target Detection. J Neurosci 26:9494–

9502.

Tomer R, Goldstein RZ, Wang G-J, Wong C, Volkow ND (2008) Incentive motivation is

associated with striatal dopamine asymmetry. Biol Psychol 77:98–101.

Tomer R, Slagter HA, Christian BT, Fox AS, King CR, Murali D, Davidson RJ (2013)

Dopamine Asymmetries Predict Orienting Bias in Healthy Individuals. Cereb Cortex

23:2899–2904.

Tommasi G, Fiorio M, Yelnik J, Krack P, Sala F, Schmitt E, Fraix V, Bertolasi L, Le Bas J-F,

Ricciardi GK, Fiaschi A, Theeuwes J, Pollak P, Chelazzi L (2014) Disentangling the

953   Role of Cortico-Basal Ganglia Loops in Top–Down and Bottom–Up Visual Attention:

954   An Investigation of Attention Deficits in Parkinson Disease. J Cogn Neurosci 27:1215–

955   1237.

956   Tommasi G, Fiorio M, Yelnik J, Krack P, Sala F, Schmitt E, Fraix V, Bertolasi L, Le Bas J-F,

957   Ricciardi GK, Fiaschi A, Theeuwes J, Pollak P, Chelazzi L (2015) Disentangling the

958   Role of Cortico-Basal Ganglia Loops in Top–Down and Bottom–Up Visual Attention:

959   An Investigation of Attention Deficits in Parkinson Disease. J Cogn Neurosci 27:1215–

960   1237.

961   Tremblay L, Hollerman JR, Schultz W (1998) Modifications of reward expectation-related

962   neuronal activity during learning in primate striatum. J Neurophysiol 80:964–977.

963   van Schouwenburg M, Aarts E, Cools R (2010a) Dopaminergic modulation of cognitive

964   control: distinct roles for the prefrontal cortex and the basal ganglia. Curr Pharm Des

965   16:2026–2032.

966   van Schouwenburg MR, den Ouden HEM, Cools R (2010b) The Human Basal Ganglia

967   Modulate Frontal-Posterior Connectivity during Attention Shifting. J Neurosci 30:9910–

968   9918.

969   Van Schouwenburg MR, Den Ouden HEM, Cools R (2015) Selective attentional

970   enhancement and inhibition of fronto-posterior connectivity by the basal ganglia during

971   attention switching. Cereb Cortex 25:1527–1534.

972   Vossel S, Geng JJ, Fink GR (2014) Dorsal and Ventral Attention Systems. Neurosci 20:150–

973   159.

974   Wilcox RR (2016a) Comparing dependent robust correlations. Br J Math Stat Psychol

975   69:215–224.

Wilcox RR (2016b) Introduction to Robust Estimation and Hypothesis Testing. Academic

Press.

Wilcox RR (2016c) Comparing dependent robust correlations. Br J Math Stat Psychol

69:215–224.

Wilke M, Mueller K-M, Leopold DA (2009) Neural activity in the visual thalamus reflects

perceptual suppression. Proc Natl Acad Sci 106:9465–9470.

Womer FY, Wang L, Alpert KI, Smith MJ, Csernansky JG, Barch DM, Mamah D (2014)

Basal ganglia and thalamic morphology in schizophrenia and bipolar disorder.

Psychiatry Res - Neuroimaging 223:75–83.

Worden MS, Foxe JJ, Wang N, Simpson G V (2000) Anticipatory biasing of visuospatial

attention indexed by retinotopically specific alpha-band electroencephalography

increases over occipital cortex. J Neurosci 20:RC63.

Yantis S, Anderson BA, Wampler EK, Laurent PA (2012) Reward and attentional control in

visual search. Nebr Symp Motiv 59:91–116.

Zheng J, Anderson KL, Leal SL, Shestyuk A, Gulsen G, Mnatsakanyan L, Vadera S, Hsu

FPK, Yassa MA, Knight RT, Lin JJ (2017) Amygdala-hippocampal dynamics during

salient information processing. Nat Commun 8:1–11.

Zhou H, Schafer RJ, Desimone R (2016) Pulvinar-Cortex Interactions in Vision and

Attention. Neuron 89:209–220.

Figure 1. Illustration of selective attention task: stimuli and reward manipulation.

(A) Six Chinese symbols served as stimuli for the task and were associated with three values: two paired with reward, two with loss and two with no financial change (neutral). (B) Representative trial of the learning phase. Symbols were displayed for 1000ms, systematically paired with the corresponding (positive, negative or neutral) value, via visual and auditory feedback. Characters presentation was alternated with a 1000ms fixation period. During the training phase, participants learned associations between the stimuli and their reward value. **C.** Representative trial of the testing phase. After a 1000ms pretrial interval, participants were primed with a 500ms preparatory cue signalling the upcoming stimuli. Two characters were then presented to the left and right hemifield, together with a spatial cue, instructing participants to covertly attend the symbol on the cued side (target) and ignore the other one (distractor). Participants' task was to report when the target stimulus changed contrast. Contrast change could either occur after 750ms (13% of trials), 1450ms (47% of trials) or 2350ms (40% of the trials). In 95% of the trials, the target changed contrast (valid trials), whilst in 5% of the trials, the distractor changed contrast (invalid trials). Figure adapted from (Marshall et al., 2017).

Figure 2. Grand average MI and HLM distribution across participants.

(A) Time-frequency representations of power (TFRs) and topographical plot showing contrast between the 'attention right' – 'attention left' trials. A clear modulation is visible at posterior sensors in the alpha band (8 – 13Hz) in the –750 – 0ms interval (this time window being considered for the computation of HLM(α) indices in (B)). Sensors included in the left and right ROIs are marked as dots. Trials are locked to the onset of the contrast change (t = 0). (B) Side panels show the temporal evolution of modulation indices in the alpha range (MI(α)), averaged over sensors within left and right hemisphere ROIs. The magnitude (absolute value) of MI(α) progressively increased in the stimulus interval until the onset of the contrast change. Middle: distribution of HLM(α) indices across participants, computed over the ROIs and 8 – 13 Hz frequency band (see Materials and Methods). A normal density function is superimposed, denoting no hemispheric bias in lateralized modulation values across participants (Shapiro Wilk, $W = .958$, $p = .392$).

Figure 3. Basal Ganglia volumes resulting from semiautomated subcortical segmentation implemented.

(A) Orthogonal view and 3D rendering. Subcortical volumes are overlaid as meshes on the anatomical MRI of one of the participants (following defacing procedure in Freesurfer, where voxels outside the brain mask with identifiable facial features were excluded (Bischoff-Grethe et al., 2007)). (B) Histograms with superimposition of normal density function, showing the distribution of subcortical lateralization indices for each substructure. In our sample, NAcb and Th volumes were left lateralized ($p = .0001$ and $p=.0003$, respectively) while CN showed a right lateralization ($p = .0029$).

Figure 4. Lateralization of individual subcortical structures in relation to alpha hemispheric lateralized modulation (HLM) in the task. (A) Bar plot displays the Beta coefficients associated with a general linear model where LV values were defined as explanatory variables for HLM(α). Error bars indicate standard error of the mean. Asterisks denote statistical significance; **p < .01. (B) Partial regression plot showing the association between LV_{GP} and HLM(α), while controlling for the other regressors in the model in (A). (C) Partial regression plot showing the association between LV_{Th} and HLM(α), while controlling for the other regressors in the model in (A). Given Eqs.(1) and (2) (see *Materials and Methods*), positive HLM(α) values indicate stronger modulation of alpha power in the right compared to the left hemisphere, and vice versa; similarly, positive (or negative) LV_S indices denote greater right(or left) volume for a given substructure s. The dotted curves in (B) and (C) indicate 95% confidence bounds for the regression line, fitted on the plot in black.

Figure 7. Linear association between GP volumetric asymmetry and alpha modulation asymmetry as a function of value-saliency occurrences in the task. (A) Correlation between GP volume lateralization and HLM(α), grouped accordingly to the number of value-salient stimuli in the trials (see *Materials and Methods*). From left to right, respectively, two, one and zero value-saliency occurrences are displayed. GP asymmetry significantly explained HLM(α) only when value-salient stimuli featured as both target and distractors, irrespective of their valence ($r = .68$, significant at the $p < .001$ level after Bonferroni correction for three comparisons). (B) The association between HLM(α) and GP volume lateralization increased as a function of value saliency in the task: the linear relationship was stronger when two value-salient stimuli were presented, when compared to conditions characterized by either one or value-salience pairings (95% CI [.106, .672] and [.125, .897], respectively for the two comparisons). This suggests that, when both target and distractor were associated with a salient value, participants exhibiting bigger GP volume in the left hemisphere than in the right hemisphere, were also better at modulating alpha oscillations in the left compared to the right hemisphere. Asterisks denote statistical significance; **$p < .01$.

Figure 8. Linear association between Th volumetric asymmetry and alpha modulation asymmetry as a function of value-saliency occurrences in the task. (A) Correlation between TH volume lateralization and HLM(α), grouped accordingly to the number of value-salient stimuli in the trials (see *Materials and Methods*). From left to right, respectively, two, one and zero value-saliency occurrences are displayed. When considering individual correlations between Th asymmetry and HLM(α), no significant linear relationship was found. (B) The association between the two measures also didn't significantly differ as a function of saliency in the trials.

Figure 9. Mean and lateralized reaction times (RT) and accuracy values across the three value-saliency occurrences in the task.

Mean RT (A) and accuracy (C) values averaged across participants in the three value-salient occurrences conditions in the task. Error bars indicate standard error of the means. No significant difference was found between groups by means of one-way repeated measures ANOVA, indicating that different levels of value-saliency pairings didn't influence behavioural performance. No significant difference emerged also when comparing average lateralized values of RT (B) and accuracy (D) across the same conditions, and by means of same statistical analysis, indicating that the behavioural spatial bias was not affected by the different levels of value-saliency pairings.

A

B

Figure 10. General linear model displaying combined lateralized subcortical volumes and hemispheric lateralized

modulation as multiple regressors for the prediction of spatial behavioural bias in accuracy (A) and RT (B). No significant

regression was found which could account for either the lateralized accuracy or RTs (p=.429 and p=.570, respectively).

Figure 11. Alpha modulation indices for left and right hemispheres associated with fast versus slow trials, neutral

condition only. (A) Topographical plot of MI(α) values for the two trial groups, clustered according to median split of

reaction times (fast versus slow trials). Left and right sensors of interest are marked as dots and correspond to the same ROIs

as in Figure 2. (B) Bar graph showing mean MI(α) averaged over ROIs in the two subgroups. Error bars represent standard

error of the mean. Horizontal red lines in each bar plot indicate the average LI (α) for the corresponding subgroup.