Genetic divergence and phenotypic plasticity contribute to variation in cuticular hydrocarbons in the seaweed fly *C. frigida*

Emma Berdan[1*], Swantje Enge[2], Göran Nylund[1], Maren Wellenreuther[3,4], Gerrit A. Martens[5], and Henrik Pavia[1]

[1]Department of Marine Sciences University of Gothenburg, Göteborg, Sweden

[2]Institute for Chemistry and Biology of the Marine Environment, Carl-von-Ossietzky University Oldenburg, Wilhelmshaven, Germany

[3]The New Zealand Institute for Plant & Food Research Limited, Nelson, New Zealand

[4]School of Biological Sciences, the University of Auckland, New Zealand

[5]Hochschule Bremen, City University of Applied Sciences, Bremen, Germany

* - Corresponding author: Emma Berdan, Department of Marine Sciences, Box 461, 405 30 Göteborg, Sweden. Email: emma.berdan@gmail.com

ABSTRACT

Sexual signals facilitate female choice, reducing mating costs. The seaweed fly, *Coelopa*

frigida is a species characterized by intense sexual conflict yet it is unclear which traits

are under sexual selection. In this study we investigated phenotypic and genetic variation

in cuticular hydrocarbons (CHCs), which act as sex pheromones in most dipterans. We

characterized CHCs in *C. frigida* for the first time and find that CHCs likely act as sexual

signals and that CHC composition is highly varied. Variation in CHC composition was

caused by both genetic (sex and population of origin) and environmental (larval diet)

effects. We identified individual compounds that contribute to these differences as well

as general shifts in chemistry. Changes in proportion of methylated compounds, mean

chain length, proportion of alkenes, and normalized total CHCs were found to differ

between these groups. We combined this data with whole genome resequencing data to

examine the genetic underpinnings of these changes. We identified 11 genes related to

CHC synthesis and find putative F_{ST} outlier SNPs in 5 of them that are concordant with

the phenotypic differences we observe. This reveals that CHC composition of *C. frigida*

is dynamic, strongly affected by the larval environment, and likely under natural and/or

sexual selection.

Keywords: cuticular hydrocarbons, *Coelopa frigida*, sexual conflict

INTRODUCTION

Females and males often differ in optimal reproductive strategies, particularly regarding

the mode and frequency of matings. Males can increase their fitness by mating with

multiple females, while one or a few matings are generally sufficient for females to

maximize their reproductive success (Bonduriansky, 2010). This sexual conflict over

mating rates commonly leads to males harassing females to increase the likelihood that

their genes get passed on to the next generation (Sánchez‑Guillén *et al.*, 2017). Such

male mating harassment can carry considerable costs for females, in some cases reducing

reproductive success and increasing mortality (Gosden & Svensson, 2007). For example,

repeated matings are the norm in the blue-tailed damselfly *Ischnura elegans*, particularly

when population densities are high (Corbet, 1999), with females commonly suffering

injuries (Chauhan *et al.*, 2016). Likewise, in the guppy *Poecilia reticulata,* males attempt

to copulate with females by thrusting their gonopodium towards the female's genital pore

as often as once a minute, disrupting normal activities like foraging (Magurran & Nowak,

1991).

Females can minimize the cost of unwanted copulations by making it hard to be detected

by males (Wellenreuther *et al.*, 2014) or evolving mate rejection responses (Andersson,

1994); however, these responses often carry associated costs (Watson *et al.*, 1998).

Consequently, females frequently endure costly matings with males when the costs of

rejection outweigh the costs of mating. Females can further modulate mating interactions

by evolving defined mate preferences. Mate preferences allow females to obtain high

quality or compatible genes from a male, with indirect benefits that include improved

offspring fecundity, survival and immune response (Neff & Pitcher, 2005). In many

solitary and social insects, cuticular hydrocarbons (CHCs) are used as one of the primary

recognition cues to recognize, and possibly discriminate, species, sex, or kin (Bagneres *et*

al., 1996; Ferveur, 2005; Blomquist & Bagnères, 2010; Van Oystaeyen *et al.*, 2014).

Cuticular hydrocarbons are thought to be a primary adaptation to life on land because

they protect insects against desiccation (Wigglesworth, 1945; Blomquist & Bagnères,

2010). *De novo* synthesis of CHCs is conserved across insects (Howard & Blomquist,

2005; Blomquist & Bagnères, 2010): Synthesis takes place in oenocytes and starts with

acetyl-CoA. A fatty acid synthase (FAS) then elongates the acetyl-CoA, forming a long-

chained fatty acyl-CoA. FASs may also add methyl groups to these fatty acids (de

Renobales *et al.*, 1986; Blomquist *et al.*, 1994; Juarez *et al.*, 1996; Chung *et al.*, 2014).

Elongases then further lengthen these fatty acyl-CoAs and double bonds or triple bonds

are added by desaturases (Howard & Blomquist, 2005; Blomquist & Bagnères, 2010).

These are reduced to aldehydes by reductases and then these aldehydes are converted to

hydrocarbons by a decarboxylation reaction, which is mediated by a cytochrome P450

(Qiu *et al.*, 2012). Coding or expression variation in any number of these enzymes can

lead to significant changes in CHC composition (ex: de Renobales *et al.*, 1986; Reed *et*

al., 1995; Qiu *et al.*, 2012; Chung *et al.*, 2014; Chen *et al.*, 2016).

CHC composition is affected by both genetic and environmental factors. Changes in gene

expression (Dallerac *et al.*, 2000a; Chertemps *et al.*, 2007; Shirangi *et al.*, 2009;

Feldmeyer *et al.*, 2014) and/or coding sequences (Keays *et al.*, 2011; Badouin *et al.*,

2013; Kulmuni *et al.*, 2013) can affect CHC composition and length. Multiple

environmental factors have also been shown to influence CHC composition. For

example, CHCs can vary under different temperature regimes (Toolson & Kupersimbron,

1989; Savarit & Ferveur, 2002; Rouault *et al.*, 2004), diets (Stennett & Etges, 1997;

Liang & Silverman, 2000; Stojkovic *et al.*, 2014) or climatic variables (Howard *et al.*,

1995; Kwan & Rundle, 2010). Moreover, multiple studies have found that CHCs vary

between populations (ex: Haverty *et al.*, 1990; Etges & Ahrens, 2001; Frentiu &

Chenoweth, 2010; Perdereau *et al.*, 2010) but it is unknown how much of this variation is

due to plasticity vs genetic divergence. These can be teased apart using laboratory

studies: genetic divergence can be assessed by raising different populations in the same

environment and plasticity can be assessed by raising the same population in multiple

environments.

The seaweed fly *Coelopa frigida* is a common inhabitant of "wrackbeds" (accumulations

of decomposing seaweed) on shorelines. These wrackbeds act as both habitat and food

source for larvae and adults of this species. Fitness trade-offs and local adaptation in

response to wrackbed composition have been demonstrated in Swedish *C. frigida*

(Wellenreuther *et al.*, 2017). The mating system in *C. frigida* is characterized by sexual

conflict. Males will approach and mount females. Females are reluctant to mate and use a

variety of responses to dislodge the male, such as downward curling of the abdomen (to

prevent contact), kicking of the legs, and shaking from side to side (Day *et al.*, 1990).

Although there is no male courtship of any kind it has been consistently reported that

mating attempts by large males are more likely to result in copulation (Butlin *et al.*, 1982;

Gilburn *et al.*, 1992; 1993). Two lines of evidence show that females are actively

choosing larger males rather than "giving up" due to exhaustion. First, under "giving up"

there should be a negative correlation between struggle time and the size differential

between the male and female. However, there is absolutely no correlation between these

factors (Edward, 2008). Second, if females only allow copulation when they have been

exhausted then struggle times should differ between mountings in which copulation

occurs and mountings where copulation does not occur. However, there is no significant

difference and a slight trend towards longer struggle times for interactions that do not end

in copulation (Edward, 2008). Currently, the mechanism by which female choice occurs

is unknown in this system.

Here we investigate cuticular hydrocarbons, a possible male signal that could enable

female choice in *C. frigida.* Specifically, we investigate if the CHC signatures vary

between sexes, populations, and wrackbed diet. We also explore putative genes involved

in CHC synthesis and look for correlations between genetic and phenotypic variation. We

discuss out findings in light of sexual selection in this species and others and outline

future research areas that deserve attention.

METHODS

Study System

Coelopa frigida belongs to the group of acalyptrate flies which exclusively forage on

decomposing seaweed (Cullen *et al.*, 1987) and is found all along the seashores of

Northern Europe (Mcalpine, 1991). This species plays a vital role in coastal

environmental health by accelerating the decomposition of algae, allowing for faster

release of nutrients (Cullen *et al.*, 1987), and larvae are an important food source for

predatory coastal invertebrates and seabirds (Summers *et al.*, 1990; Fuller, 2003; Griffin

et al., 2018).

Sampling

Fly larvae were collected in April and May 2017 from two Norwegian populations,

namely Skeie (58.69733, 5.54083) and Østhassel (58.07068, 6.64346), and two Swedish

populations, namely Stavder (57.28153, 12.13746) and Ystad (55.425, 13.77254). These

populations are situated along an environmental cline in salinity, wrackbed composition,

and wrackbed microbiome (Berdan et al., in prep; Day *et al.*, 1983)). Larvae were

brought to Tjärnö Marine Biological Laboratory of the University of Gothenburg where

they completed development in an aerated plastic pot filled with their own field wrack in

a temperature controlled room at 25° with a 12h/12h light-dark cycle. After adults

eclosed in the lab they were transferred to a new pot filled with standard wrack consisting

of 50% *Fucus spp*. and 50% *Saccharina latissima* which had been chopped, frozen, and

then defrosted. The exception to this is a subset of Ystad adults that were allowed to mate

on the same material they had been collected in (field wrack). Thus we had 5 treatment

groups: Skeie on standard wrack, Østhassel on standard wrack, Stavder on standard

wrack, Ystad on standard wrack, and Ystad on field wrack. Adult flies were allowed to

lay eggs on the provided wrack and this second generation was raised entirely in a

temperature controlled room at 25° with a 12h/12h light dark cycle. As larvae pupated

they were transferred to individual 2 mL tubes with a small amount of cotton soaked in

5% glucose to provide moisture and food for the eclosing adult. This ensured virginity in

all eclosing flies. The pots were checked every day for eclosing adults and the date of

eclosure was noted. Two days after eclosure adult flies were frozen at -80° and kept there

until CHC extraction.

<u>CHC Analysis</u>

Frozen flies were placed in 12-well porcelain plates to defrost and dry, as moisture can

affect lipid dissolution. Each fly was then placed in a 1.5 ml high recovery vial

containing 300 µl of *n*-hexane, vortexed at a low speed for 5 second, and extracted for 5

minutes. Afterwards flies were removed from the vial and allowed to air dry on the

porcelain plates before they were weighed (Sartorius Quintix 124-1S microbalance) to

the nearest 0.0001 grams and sexed. Extracts were evaporated to dryness under nitrogen

gas and then stored at -20°. Before analysis 20 µl of *n*-hexane containing 1 µg/ml *n*-

nonane as an internal standard was added to each vial, which was then vortexed at

maximum speed for 10 seconds.

The extracts from 20 male and 20 female flies from every treatment group were analyzed

on GC (Agilent GC 6890) coupled to MS (Agilent 5973 MSD) using a HP-5MS capillary

column (Agilent) (see Table S1 in supplementary materials for instrument settings).

The total ion chromatograms were quality checked, de-noised and Savitzky-Golay

filtered before peak detection and peak area integration in OpenChrom®. Peaks were

then aligned using the R-package 'GCalignR'(Ottensmann, Stoffel, & Hoffman, 2017)

prior to statistical analysis. Peaks were tentatively identified by its mass spectra and

retention time index based on a C21-C40 *n*-alkane standard solution (Sigma-Aldrich).

<u>Statistical Analysis</u>

The effects of sex and population on CHC profile were assessed using flies raised on

standard wrack using a balanced sampling design (N=13 for each combination) and 127

peaks identified by R package "GCalignR". The effects of sex and diet (i.e. wrack type)

during the larval stage on CHC profiles were analyzed using a balanced design of flies

from the Ystad population (N=12 for each combination) and 111 peaks identified by R

package "GCalignR". Peak areas were normalized on the peak area of the internal

standard and on the weight of the fly before being auto-scaled prior to statistical analysis.

Clustering of samples was visualized with a PCA, and group differences were analyzed

using a PERMANOVA followed by multiple group comparisons using the PRIMER-E 7

software. We also analyzed group differences via (O)PLS-DA using the "ropls"-packages

in R (Thevenot, Roux, Xu, Ezan, & Junot, 2015). Candidate compounds for group

differentiation were determined from variables of importance for OPLS-DA projection

(VIP scores), which were further assessed by univariate analysis using a false discovery

adjusted significance level of $\alpha = 0.05$.

We examined sex and country effects in more depth to look for shifts in chemistry that

could be linked to our genetic data (see below). Our results above indicated that the

differences in CHC profiles between sexes and populations of *C. frigida* were due to

quantitative rather than qualitative differences (i.e. changes in relative amounts rather

than presence/absence of certain compounds). Thus, we looked for general shifts in CHC

chemistry by examining 1. The total normalized peak area (i.e. the sum of all peaks used

in analysis), 2. The proportion of alkenes, and 3. The proportion of methylated

compounds. For #1 we used all 127 peaks used in the previous analysis and for #2 and #3

we used the subset of these peaks that could be accurately categorized (116 peaks). For

the total peak area we used a log transformation and then used the 'dredge' function from

the MuMIn package (Barton, 2017) to compare AIC values for nested glm (gaussian

distribution link=identity) models with the terms sex, country, and country*sex. If there

were two models that differed in AIC by less than 1, we chose the simpler model. For

alkenes and methylated compounds our data was proportional and did not include 1 or 0

so we used the 'betareg' function from the betareg package (Grun *et al.*, 2012). We again

used the 'dredge' function to compare AIC values for nested models with the terms sex,

country, and country*sex and took the simpler model in case of an AIC difference of less

than 1.

We also analysed the distribution of chain lengths in our samples. We subset all peaks

where we had accurate length information (107 peaks). We calculated weighted mean

chain length as

$$N = \Sigma \, l_i p_i$$

where l_i is the length of a carbon chain and p_i is the proportional concentration of all

chains of that length. We also calculated dispersion around this mean as

198                              $$D = \Sigma\, p_i(l_i - N)^2$$

We analyzed these in the same manner as described above, using AIC values to conduct

model choice from glm models (Gaussian distribution link=identity).

<u>Candidate gene analysis</u>

Based on a literature search for genes involved in CHC synthesis, we identified 37

reference protein sequences, which were downloaded from Flybase (flybase.org) in

October 2017. We used tblastx with the default settings to identify orthologs in the *C.*

frigida transcriptome (Berdan & Wellenreuther, unpublished). We then used gmap (Wu

& Watanabe, 2005) to find areas of the *C. frigida* genome (Wellenreuther, *et al.*

unpublished) that might contain these genes. We output all alignments as a gff3 file. We

have a previously made a VCF file created from 30x whole genome re-sequencing of 46

C. frigida across five populations in Scandinavia (Berdan *et. al*, unpublished). We used

this VCF file and the gff3 output from gmap in SNPeff (Cingolani *et al.*, 2012) to

annotate the SNPs in these genes.

To find which SNPs are diverged between populations we used vcftools (Danecek *et al.*,

2011) to calculate Cockerham and Weir's F_{ST} estimate (Weir & Cockerham, 1984) for all

SNPs in our reference genome. We retained the top 5% of this distribution ($F_{ST} \geq 0.142$,

mean F_{ST} 0.024) as SNPs that were potentially divergent. We then subset the VCF file to

include only outlier SNPs located in our previously identified genomic regions.

RESULTS

<u>CHC analysis</u>

A rich blend of hydrocarbons with more than 100 different compounds were found in the

cuticular extract of *Coelopa frigida*. The majority of CHCs ranged in chain length from

23 to 33 carbon atoms. They contained odd numbered *n*-alkanes, methyl-, dimethyl- and

trimethyl-alkanes as well as odd numbered alkenes. However, some even numbered

alkanes were present in lower amounts.

<u>Sex and population effects</u>

Results of the PERMANOVA indicated significant differences between females and

males and between populations but also a significant interaction between the two factors

(Table 1A). Pair-wise tests on the interaction showed a significant difference between

males and females for all populations except Ystad (Table S2). Multiple comparisons of

populations showed that most of our populations differed from one another and a PCA

and PLS-DA indicated that our Swedish (Ystad and Stavder) and Norwegian (Skeie and

Østhassel) populations seemed to group together by country (Figure 2A, C, Table S3).

After combining the Swedish and Norwegian populations, the PERMANOVA indicated

highly significant differences between females and males as well as between the Sweden

and Norway (Table 1B). OPLS-DA models for either sex or country were significant and

lead to separate clusters of females and males and Sweden and Norway, respectively

(Figure 2B, D, Table S3). According to the OPLS-DA model there were 41 peaks that

contributed with a VIP score >1 to the differentiation between females and males. Of

these 31 turned out to be also significant in t-tests with a FDR adjusted p-values (Table

2). These were highly male biased with all except 3 peaks more abundant in males.

Looking at differentiation between Swedish and Norwegian populations we found 43

compounds with a VIP score >1, 29 of which were significant in t-tests with FDR

adjusted p-values and consisted of an equal mix of compounds more abundant in either of

the countries (Table 2).

Sex and diet effects

PERMANOVA demonstrated a significant difference between females and males in the

Ystad population and also a significant effect of the larval diet (wrack type) (Table 1C).

Both OPLS-DA models for either sex or wrack type were significant and lead to separate

clusters of females and males and of flies raised on standard or field wrack (Figure 3B,C,

Table S3). According to the OPLS-DA model there were 38 peaks that contributed to the

differentiation between females and males with a VIP score >1. Of these, 24 compounds

were also significant in a t-test with FDR adjusted p-values (Table 2). These were highly

male biased with all except 3 peaks more abundant in males. Overall 65% (26

compounds) were shared between this analysis and the previous analysis of sex and

population effects. Looking at differentiation between flies raised on diet of standard or

field wrack we found 33 compounds with a VIP score >1. Of these, 27 compounds were

significant using a t-test with FDR adjusted p-values, all with higher amounts in flies

reared on field wrack (Table 2). Flies reared on field wrack had more CHCs per gram

body weight than flies reared on standard wrack (normalized totals: Females field wrack

7.96 ± 0.94, females standard wrack 4.09 ± 0.47, males field wrack 11.24 ± 1.03, males

standard wrack 7.09 ± 1.21).

Overall, the results indicated that the differences in CHC profiles between sexes,

populations and diet type of *C. frigida* were due to quantitative differences, affecting a

wide range of compounds instead of qualitative differences, i.e. the presence/ absence of

a few distinct compounds.

<u>Shifts in Chemistry</u>

As our results above indicated an overall effect of country rather than population we used

country and sex for our terms in these models. For total normalized peak area, the best

model contained only sex as a main effect (for full AIC comparison for all models see

Table S4A-E). In general, males had more CHCs per gram body weight than females

(Figure 4A). For proportion of alkenes, the best model contained country and sex as main

effects (for type II analysis of deviance tables for all best models see Table S5A-E).

Females tended to have more alkenes than males and individuals from Norway tended to

have more alkenes than individuals from Sweden (Figure 4B). The best model for the

proportion of methylated compounds contained sex and country as the main effects. As

with alkenes, females tended to have more methylated compounds than males and

individuals from Norway tended to have more methylated compounds than individuals

from Sweden (Figure 4C).

The best model for weighted mean chain length was country + sex. Overall, males had

longer mean chain lengths than females and individuals from Sweden had longer mean

chain lengths than individuals from Norway (Figure 5A). The best model for dispersion

around mean chain length was also country + sex. Males had higher dispersion than

females and individuals from Norway had higher dispersion than individuals from

Sweden (Figure 5B).

<u>Genetic analysis</u>

We identified 15 isoforms from 13 transcripts in our transcriptome with matches to our

query proteins. These transcripts mapped to 16 unique areas of the genome assembly. We

blasted these areas against the NCBI nr database (accessed in April 2018) to confirm our

annotation. Five of these areas had either no match or matched an unrelated protein and

were removed. The remaining genes include 2 putative fatty acid synthases, 2 putative

desaturases, 5 putative elongases, 1 putative cytochrome P450 reductase, and 1 putative

cytochrome P450-4G1 (Table 3). We found 1042 SNPs within these genes (including 5K

upstream and 5K downstream), of these 11 had an F_{ST} value > 0.142 (i.e. in the top 5% of

the F_{ST} distribution, Table 4).

DISCUSSION

In a sexual conflict system females may minimize the cost of unwanted copulations via

mate rejection responses; however, these responses typically carry associated costs

(Watson *et al.*, 1998). The ability to modulate mating responses and the level of

harassment based on male traits is especially beneficial in systems where no traditional

"courtship" exists. Here we explored a potential male trait, cuticular hydrocarbons

(CHCs) and assessed variation between sexes, populations, and different diets. We

supplemented this with a scan of the *C. frigida* genome to search for candidate genes for

CHC synthesis and to determine if genetic variation could be tied to the phenotypic

patterns we saw. We found phenotypic variation attributable to sex, population, and diet

and genetic variation attributable to population. We discuss these results and their

consequences below.

Males and females differed in their CHC composition. Not only did the composition

between males and females differ (Table 2, Figure 2-5) but, in addition, males

consistently had more CHC per gram body weight than females (Figure 4A). This is a

strong indication that CHCs play a role in sexual communication in *C. frigida* under a

female choice scenario, as it is unlikely that desiccation risk would differ between males

and females. A role for CHCs in sexual selection in *C. frigida* would be in line with the

general finding that CHCs play a major role in communication in insects, specifically in

Diptera (Howard & Blomquist, 2005; Blomquist & Bagnères, 2010; Ferveur & Cobb,

2010). Hydrocarbons may be involved in a wide variety of behaviours in Diptera such as

courtship, aggregation, and dominance (Ferveur & Cobb, 2010). For instance, 11-*cis*-

vaccenyl acetate, a male specific hydrocarbon in *Drosophila melanogaster*, increases

male-male aggression as well as influencing male mating behavior (Wang & Anderson,

2010). Future behavioural tests will be necessary to confirm that CHCs are used for mate

selection but attention should also be paid to behaviours such as aggression and

aggregation. Aggregation hydrocarbons, such as (Z)-10-heneicosene in *D. virilis* can

attract males and females of certain ages (Bartelt & Jackson, 1984). *Coelopa frigida* is

famous for forming large aggregations that often disrupt human activity (Mather, 2011).

The majority and most abundant CHCs found in *C. frigida* had chain lengths from 23 to

33 carbon atoms. The majority consisted of odd numbered *n*-alkanes, methyl-, dimethyl-

and trimethyl-alkanes as well as odd numbered alkenes, which is consistent with the

general biosynthetic pathway of long-chain hydrocarbons derived from fatty acids

(Blomquist, 2010). However, some even numbered alkanes (mainly C30 and C26) were

present in lower amounts.

We found shifts in CHC composition between sexes, populations, and diet. As we only

examined coding variation, only one of these factors, population, could be tied to genetic

differences. Rather than population specific variation we observed country level variation

with strong differences between Norway and Sweden (Figure 2D, Figure 4, Figure 5).

This pattern mirrors a strong genetic split between the two countries (representing

approximately 14% of genetic variation, Berdan *et al*, unpublished). We found potential

outlier SNPs in desaturases, elongases, a cytochrome P450-4G1, and a cytochrome P450

reductase. Desaturases add double bonds or triple bonds to alkanes (Howard &

Blomquist, 2005; Blomquist & Bagnères, 2010). We found an outlier missense variant in

a putative Desaturase 1 as well as a missense and two downstream outlier variants in a

putative Desaturase (Table 4). This aligns with our phenotypic data showing differences

in proportion of alkanes vs. alkenes between Norway and Sweden. Elongases lengthen

fatty acyl-CoAs and are necessary for chains longer than 16 or so (Blomquist &

Bagnères, 2010). We found both synonymous and downstream outlier variants in putative

C. frigida elongases (Table 4) and corresponding differences in both mean chain length

and variance in chain length (dispersion around the mean) due to country of origin. P450

reductases are responsible for reducing fatty acids to aldehydes (Blomquist & Bagnères,

2010). Knockdown of this in *D. melanogaster* leads to a striking reduction of cuticular

hydrocarbons (Qiu *et al.*, 2012). P450-4G genes, which are unique to insects, encode an

oxidative decarboxylase that catalyzes the aldehyde to hydrocarbon reaction. Work in

other insect species has found that a knockdown or knockout of this gene leads to a

decrease in overall hydrocarbon levels (Reed *et al.*, 1995; Qiu *et al.*, 2012; Chen *et al.*,

2016). We found a downstream outlier variant in our cytochrome P450-4G1 and multiple

kinds of outlier variants in a cytochrome P450 reductase. However, total amount of

CHCs did not differ as a result of anything except for sex. As males and females share a

genome it is unlikely that these SNPs affect this difference unless they are located on the

sex chromosome. Finally, we also found variation between Sweden and Norway in the

proportion of methylated compounds. However, research suggests that these differences

are often caused by either variation in Fatty Acid Synthase (FAS, de Renobales *et al.*,

1986; Blomquist *et al.*, 1994; Juarez *et al.*, 1996) or multiple copies of FAS with different

functions (Chung *et al.*, 2014). We found two different FAS' in our genome (Table 3) but

neither had divergent SNPs. Future studies are needed to examine if these genetic and

phenotypic changes are related. Specifically, investigation into both RNA expression and

tests of function of these loci will be necessary to provide a direct link. However, the

association between the known function of several of these genes (elongases, desaturases)

and the corresponding differences in populations are marked. Given that we see shifts in

multiple chemical aspects (length, methylation, and alkene/alkane ratio) it is likely that

multiple loci are responsible for these patterns.

We found a shift in the composition rather than differences in the presence absence of

specific compounds. This type of pattern differs from some Dipterans where sex and

population differences may be mostly qualitative (Carlson & Schlein, 1991; Ferveur &

Sureau, 1996; Dallerac *et al.*, 2000b; Gomes *et al.*, 2008; Everaerts *et al.*, 2010) although

plenty of Dipterans show quantitative differences as well (ex: Jallon & David, 1987;

Byrne *et al.*, 1995). At least part of this discrepancy is likely explained by the number of

compounds that make up the CHCs in *C. frigida* versus other Dipterans. Many Dipterans

have principal CHC components that make up a large proportion of the CHC composition

(Ferveur & Sureau, 1996; Etges & Jackson, 2001; Gomes *et al.*, 2008). Cuticular extracts

from *C. frigida*, have a large number of compounds but do not appear to have compounds

that are dominant in any one sex or population. When averaging across all samples, the

most abundant compound was the C25 alkane, which made up on average 15% of the

composition.

The wrackbed itself also had a strong influence on the CHC composition (Figure 3C).

Wrackbed composition and microbiome (i.e. the food source for *C. frigida* larvae) varies

across *C. frigida* populations in Europe (Berdan, et al., in prep, Day *et al.*, 1983; Butlin &

Day, 1989; Wellenreuther *et al.*, 2017), indicating that natural populations may be

composed of different CHC composition. Adding to this, we found strong population

(country) level differences in CHC composition even when larvae were reared on the

same substrate. We also found several outlier SNPs in a variety of genes involved in

CHC synthesis. Together these results indicate several possibilities: 1. Natural selection

may select for different CHC composition in different environments and natural

polymorphism in CHC genes may be under direct selection due to this. 2. Natural

populations likely differ strongly in their CHC composition. If assortative mating or

selection on female preferences is present that could lead to locally adapted male traits

and female preferences. This would reduce effective migration between populations, as

foreign males would be at a disadvantage.

In conclusion, the analysis of cuticular hydrocarbon extracts from male and female *C.*

frigida from multiple populations revealed a complex and variable mix of more than 100

different compounds. We found extensive phenotypic variation attributable to diet, sex,

and population and potentially underlying genetic variation attributable to population.

This reveals that CHC composition is dynamic, strongly affected by the larval

environment, and most likely under natural and/or sexual selection. Further work is

needed to test the exact role of CHCs in this system.

REFERENCES

Andersson, M.B. 1994. *Sexual selection*. Princeton University Press.

Badouin, H., Belkhir, K., Gregson, E., Galindo, J., Sundström, L., Martin, S.J., Butlin, R.K. & Smadja, C.M. 2013. Transcriptome characterisation of the ant Formica exsecta with new insights into the evolution of desaturase genes in social hymenoptera. *PloS one* **8:** e68200.

Bagneres, A.G., Lorenzi, M.C., Dusticier, G., Turillazzi, S. & Clement, J.L. 1996. Chemical usurpation of a nest by paper wasp parasites. *Science* **272:** 889-892.

Bartelt, R. & Jackson, L. 1984. Hydrocarbon component of the Drosophila virilis (Diptera: Drosophilidae) aggregation pheromone:(Z)-10-heneicosene. *Annals of the Entomological Society of America* **77:** 364-371.

Barton, K. 2017. MuMIn.

Blomquist, G.J., Guo, L., Gu, P., Blomquist, C., Reitz, R.C. & Reed, J.R. 1994. Methyl-branched fatty acids and their biosynthesis in the housefly, Musca domestica L.(Diptera: Muscidae). *Insect biochemistry and molecular biology* **24:** 803-810.

Blomquist, G.J. 2010. Biosynthesis of cuticular hydrocarbons. *Insect hydrocarbons: biology, biochemistry and chemical ecology* **35:** 52.

Blomquist, G.J. & Bagnères, A.-G. 2010. *Insect hydrocarbons: biology, biochemistry, and chemical ecology*. Cambridge University Press.

Bonduriansky, R. 2010. Sexual conflict, John Wiley & Sons, Ltd. Chichester.

Butlin, R. & Day, T. 1989. Environmental correlates of inversion frequencies in natural populations of seaweed flies (Coelopa frigida). *Heredity* **62:** 223.

Butlin, R.K., Read, I.L. & Day, T.H. 1982. The effects of a chromosomal inversion on adult size and male mating success in the seaweed fly, *Coelopa frigida*. *Heredity* **49:** 51-62.

Byrne, A., Camann, M., Cyr, T., Catts, E. & Espelie, K. 1995. Forensic implications of biochemical differences among geographic populations of the black blow fly, Phormia regina (Meigen). *Journal of Forensic Science* **40:** 372-377.

Carlson, D. & Schlein, Y. 1991. Unusual polymethyl alkenes in tsetse flies acting as abstinon inGlossina morsitans. *Journal of chemical ecology* **17:** 267-284.

Chauhan, P., Wellenreuther, M. & Hansson, B. 2016. Transcriptome profiling in the damselfly *Ischnura elegans* identifies genes with sex-biased expression. *BMC genomics* **17:** 985.

Chen, N., Fan, Y.L., Bai, Y., Li, X.D., Zhang, Z.F. & Liu, T.X. 2016. Cytochrome P450 gene, CYP4G51, modulates hydrocarbon production in the pea aphid, Acyrthosiphon pisum. *Insect Biochemistry and Molecular Biology* **76:** 84-94.

Chertemps, T., Duportets, L., Labeur, C., Ueda, R., Takahashi, K., Saigo, K. & Wicker-Thomas, C. 2007. A female-biased expressed elongase involved in long-chain hydrocarbon biosynthesis and courtship behavior in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America* **104:** 4273-4278.

Chung, H., Loehlin, D.W., Dufour, H.D., Vaccarro, K., Millar, J.G. & Carroll, S.B. 2014. A single gene affects both ecological divergence and mate choice in Drosophila. *Science* **343:** 1148-1151.

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X.Y. & Ruden, D.M. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w(1118); iso-2; iso-3. *Fly* **6:** 80-92.

Corbet, P. 1999. *Dragonflies: Behavior and Ecology of Odonata*. Essex: Harley Books.

Cullen, S.J., Young, A.M. & Day, T.H. 1987. Dietary requirements of seaweed flies (Coelopa frigida). *Estuarine, Coastal and Shelf Science* **24:** 701-710.

Dallerac, R., Labeur, C., Jallon, J.-M., Knipple, D.C., Roelofs, W.L. & Wicker-Thomas, C. 2000a. A Δ9 desaturase gene with a different substrate specificity is responsible for the cuticular diene hydrocarbon polymorphism in Drosophila melanogaster. *Proceedings of the National Academy of Sciences* **97:** 9449-9454.

Dallerac, R., Labeur, C., Jallon, J.M., Knippie, D.C., Roelofs, W.L. & Wicker-Thomas, C. 2000b. A Delta 9 desaturase gene with a different substrate specificity is responsible for the cuticular diene hydrocarbon polymorphism in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America* **97:** 9449-9454.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R. & Grp, G.P.A. 2011. The variant call format and VCFtools. *Bioinformatics* **27:** 2156-2158.

Day, T.H., Dawe, C., Dobson, T. & Hillier, P.C. 1983. A chromosomal inversion polymorphism in Scandinavian populations of the seaweed fly, Coelopa frigida. *Hereditas* **99:** 135-145.

Day, T.H., Foster, S.P. & Engelhard, G. 1990. Mating behavior in seaweed flies (Coelopa frigida). *Journal of Insect Behavior* **3:** 105-120.

de Renobales, M., Woodin, T.S. & Blomquist, G.J. 1986. Drosophila melanogaster fatty acid synthetase: characteristics and effect of protease inhibitors. *Insect biochemistry* **16:** 887-894.

Edward, D.A. 2008. Habitat composition, sexual conflict and life history evolution in Coelopa frigida.

Etges, W.J. & Ahrens, M.A. 2001. Premating isolation is determined by larval-rearing substrates in cactophilic Drosophila mojavensis. V. Deep geographic variation in epicuticular hydrocarbons among isolated populations. *The American Naturalist* **158:** 585-598.

Etges, W.J. & Jackson, L.L. 2001. Epicuticular hydrocarbon variation in Drosophila mojavensis cluster species. *Journal of chemical ecology* **27:** 2125-2149.

Everaerts, C., Farine, J.P., Cobb, M. & Ferveur, J.F. 2010. Drosophila Cuticular Hydrocarbons Revisited: Mating Status Alters Cuticular Profiles. *Plos One* **5**.

Feldmeyer, B., Elsner, D. & Foitzik, S. 2014. Gene expression patterns associated with caste and reproductive status in ants: worker‐specific genes are more derived than queen‐specific ones. *Molecular ecology* **23:** 151-161.

Ferveur, J.-F. 2005. Cuticular hydrocarbons: their evolution and roles in Drosophila pheromonal communication. *Behavior genetics* **35:** 279.

Ferveur, J.-F. & Cobb, M. 2010. Behavioral and evolutionary roles of cuticular hydrocarbons in Diptera. *Insect hydrocarbons: biology, biochemistry, and chemical ecology***:** 325-343.

Ferveur, J.F. & Sureau, G. 1996. Simultaneous influence on male courtship of stimulatory and inhibitory pheromones produced by live sex-mosaic Drosophila melanogaster. *Proceedings of the Royal Society B-Biological Sciences* **263:** 967-973.

Frentiu, F.D. & Chenoweth, S.F. 2010. Clines in cuticular hydrocarbons in two Drosophila species with independent population histories. *Evolution* **64:** 1784-1794.

Fuller, R.A. 2003. Factors influencing foraging decisions in ruddy turnstones Arenaria interpres (L.), Durham University.

Gilburn, A., Foster, S. & Day, T. 1992. Female mating preference for large size in Coelopa frigida(seaweed fly). *Heredity* **69:** 209-216.

Gilburn, A., Foster, S. & Day, T. 1993. Genetic correlation between a female mating preference and the preferred male character in seaweed flies (Coelopa frigida). *Evolution* **47:** 1788-1795.

Gomes, C.C.G., Trigo, J.R. & Eiras, Á.E. 2008. Sex pheromone of the American warble fly, Dermatobia hominis: the role of cuticular hydrocarbons. *Journal of chemical ecology* **34:** 636.

Gosden, T. & Svensson, E.I. 2007. Female sexual polymorphism and fecundity consequences of male mating harassment in the wild. *PLoS ONE***:** 2(6):e580 doi:510.1371/journal.pone.0000580.

Griffin, C., Day, N., Rosenquist, H., Wellenreuther, M., Bunnefeld, N. & Gilburn, A. 2018. Tidal range and recovery from the impacts mechanical beach grooming. *Ocean and Coastal Management* **154:** 66-71.

Grun, B., Kosmidis, I. & Zeileis, A. 2012. Extended Beta Regression in R: Shaken, Stirred, Mixed, and Partitioned. *Journal of Statistical Software* **48:** 1-25.

Haverty, M.I., Nelson, L.J. & Page, M. 1990. Cuticular hydrocarbons of four populations of Coptotermes formosanus Shiraki in the united states. *Journal of chemical ecology* **16:** 1635-1647.

Howard, R.W., Howard, C.D. & Colquhoun, S. 1995. Ontogenetic and environmentally induced changes in cuticular hydrocarbons of Oryzaephilus surinamensis (Coleoptera: Cucujidae). *Annals of the Entomological Society of America* **88:** 485-495.

Howard, R.W. & Blomquist, G.J. 2005. Ecological, behavioral, and biochemical aspects of insect hydrocarbons. *Annu. Rev. Entomol.* **50:** 371-393.

Jallon, J.M. & David, J.R. 1987. Variations in cuticular hydrocarbons among the eight species of the Drosophila melanogaster subgroup. *Evolution* **41:** 294-302.

Juarez, M.P., Ayala, S. & Brenner, R.R. 1996. Methyl-branched fatty acid biosynthesis in Triatoma infestans. *Insect Biochemistry and Molecular Biology* **26:** 599-605.

Keays, M.C., Barker, D., WICKER‐THOMAS, C. & Ritchie, M.G. 2011. Signatures of selection and sex‐specific expression variation of a novel duplicate during the evolution of the Drosophila desaturase gene family. *Molecular ecology* **20:** 3617-3630.

Kulmuni, J., Wurm, Y. & Pamilo, P. 2013. Comparative genomics of chemosensory protein genes reveals rapid evolution and positive selection in ant-specific duplicates. *Heredity* **110:** 538-547.

Kwan, L. & Rundle, H.D. 2010. Adaptation to Desiccation Fails to Generate Pre- and Postmating Isolation in Replicate Drosophila Melanogaster Laboratory Populations. *Evolution* **64:** 710-723.

Liang, D. & Silverman, J. 2000. "You are what you eat": diet modifies cuticular hydrocarbons and nestmate recognition in the Argentine ant, Linepithema humile. *Naturwissenschaften* **87:** 412-416.

Magurran, A.E. & Nowak, M.A. 1991. Another battle of the sexes: the consequences of sexual asymmetry in mating costs and predation risk in the guppy, Poecilia reticulata. *Proc. R. Soc. Lond. B* **246:** 31-38.

Mather, K. 2011. South Bay beaches hit my swarms of kelp flies. *Los Angeles Times.* Los Angeles.

Mcalpine, D.K. 1991. Review of the Australian Kelp Flies (Diptera, Coelopidae). *Systematic Entomology* **16:** 29-84.

Neff, B.D. & Pitcher, T.E. 2005. Genetic quality and sexual selection: an integrated framework for good genes and compatible genes. *Mol Ecol* **14:** 19-38.

Perdereau, E., Dedeine, F., Christidès, J.-P. & Bagnères, A.-G. 2010. Variations in worker cuticular hydrocarbons and soldier isoprenoid defensive secretions within and among introduced and native populations of the subterranean termite, Reticulitermes flavipes. *Journal of chemical ecology* **36:** 1189-1198.

Qiu, Y., Tittiger, C., Wicker-Thomas, C., Le Goff, G., Young, S., Wajnberg, E., Fricaux, T., Taquet, N., Blomquist, G.J. & Feyereisen, R. 2012. An insect-specific P450 oxidative decarbonylase for cuticular hydrocarbon biosynthesis. *Proceedings of the National Academy of Sciences of the United States of America* **109:** 14858-14863.

Reed, J.R., Quilici, D.R., Blomquist, G.J. & Reitz, R.C. 1995. Proposed mechanism for the cytochrome P 450-catalyzed conversion of aldehydes to hydrocarbons in the house fly, Musca domestica. *Biochemistry* **34:** 16221-16227.

Rouault, J.D., Marican, C., Wicker-Thomas, C. & Jallon, J.M. 2004. Relations between cuticular hydrocarbon (HC) polymorphism, resistance against desiccation and breeding temperature; a model for HC evolution in D-melanogaster and D-simulans. *Genetica* **120:** 195-212.

Sánchez‐Guillén, R.A., Wellenreuther, M., Chávez‐Ríos, J.R., Beatty, C.D., Rivas‐Torres, A., Velasquez‐Velez, M. & Cordero‐Rivera, A. 2017. Alternative reproductive strategies and the maintenance of female color polymorphism in damselflies. *Ecology and Evolution* **7:** 5592-5602.

Savarit, F. & Ferveur, J.F. 2002. Temperature affects the ontogeny of sexually dimorphic cuticular hydrocarbons in Drosophila melanogaster. *Journal of Experimental Biology* **205:** 3241-3249.

Shirangi, T.R., Dufour, H.D., Williams, T.M. & Carroll, S.B. 2009. Rapid evolution of sex pheromone-producing enzyme expression in Drosophila. *PLoS biology* **7:** e1000168.

Stennett, M.D. & Etges, W.J. 1997. Premating Isolation Is Determined by Larval Rearing Substrates in CactophilicDrosophila mojavensis. III. Epicuticular Hydrocarbon Variation Is Determined by Use of Different Host Plants inDrosophila mojavensis andDrosophila arizonae. *Journal of Chemical Ecology* **23:** 2803-2824.

Stojkovic, B., Savkovic, U., Dordevic, M. & Tucic, N. 2014. Host-shift effects on mating behavior and incipient pre-mating isolation in seed beetle. *Behavioral Ecology* **25:** 553-564.

Summers, R.W., Smith, S., Nicoll, M. & Atkinson, N.K. 1990. Tidal and Sexual Differences in the Diet of Purple Sandpipers Calidris-Maritima in Scotland. *Bird Study* **37:** 187-194.

Toolson, E.C. & Kupersimbron, R. 1989. Laboratory Evolution of Epicuticular Hydrocarbon Composition and Cuticular Permeability in Drosophila-

Pseudoobscura - Effects on Sexual Dimorphism and Thermal-Acclimation Ability. *Evolution* **43:** 468-473.

Van Oystaeyen, A., Oliveira, R.C., Holman, L., van Zweden, J.S., Romero, C., Oi, C.A., d'Ettorre, P., Khalesi, M., Billen, J. & Wäckers, F. 2014. Conserved class of queen pheromones stops social insect workers from reproducing. *Science* **343:** 287-290.

Wang, L.M. & Anderson, D.J. 2010. Identification of an aggression-promoting pheromone and its receptor neurons in Drosophila. *Nature* **463:** 227-U114.

Watson, P.J., Stallmann, R.R. & Arnqvist, G. 1998. Sexual conflict and the energetic costs of mating and mate choice in water striders. *The American Naturalist* **151:** 46-58.

Weir, B.S. & Cockerham, C.C. 1984. Estimating F-Statistics for the Analysis of Population-Structure. *Evolution* **38:** 1358-1370.

Wellenreuther, M., Svensson, E.I. & Hansson, B. 2014. Sexual selection and genetic colour polymorphisms in animals. *Molecular Ecology* **23:** 5398-5414.

Wellenreuther, M., Rosenquist, H., Jaksons, P. & Larson, W. 2017. Local adaptation along an environmental cline in a species with an inversion polymorphism. *Journal of Evolutionary Biology* **30:** 1068-1077.

Wigglesworth, V. 1945. Transpiration through the cuticle of insects. *Journal of Experimental Biology* **21:** 97-114.

Wu, T.D. & Watanabe, C.K. 2005. GMAP: a genomic mapping and alignment program for mRNA and EST sequences. *Bioinformatics* **21:** 1859-1875.

Tables

Table 1– PERMANOVA and PERMDISP results for A. Sex and Population, B. Sex and Country, and C. Sex and Wrack type. $* < 0.05$ $** < 0.01$ $*** < 0.001$

A.

		PERMANOVA: Sex and Population					PERMDISP
	df	SS	MS	F	p-value		
Sex	1	658.1	658.1	5.86	<0.001	***	see S2
Population	3	1150.3	383.4	3.41	<0.001	***	
Sex:Population	3	486.9	162.3	1.44	0.015	*	
Residuals	96	10786.0	112.4				
Total	103	13081					

B.

		PERMANOVA: Sex and Country					PERMDISP	
	df	SS	MS	F	p-value		F	p-value
Sex	1	658.0	658.1	5.60	<0.001	***	8.94	0.003 **
Country	1	517.8	517.8	4.41	<0.001	***	1.19	0.279
Sex:Country	1	154.1	154.1	1.31	0.128			
Residuals	100	11751.0	117.5					
Total	103	13081						

C.

		PERMANOVA: Sex and Wrack type					PERMDISP	
	df	SS	MS	F	p-value		F	p-value
Sex	1	360.9	360.9	3.73	<0.001	***	2.36	0.129
Wrack	1	481.3	481.3	4.98	<0.001	***	2.29	0.139
Sex:Wrack	1	122.3	122.3	1.27	0.1629			
Residuals	44	4252.5	96.7					
Total	47	5217						

Table 2 – List of compounds that are significantly important for differentiation of at least factor (sex, diet, or country). If compounds are significant for differentiation of any category in any comparison a * appears under that factor. Color indicates the direction of the differentiation (red- higher in females, blue – higher in males, grey – higher in Norwegian populations, green – higher in Swedish populations, yellow – higher in flies raised on standard wrack, purple – higher in flies raised on field wrack).

Colors in cells are indicated in parentheses: (red) higher in females, (blue) higher in males, (grey) higher in Norwegian populations, (green) higher in Swedish populations, (yellow) higher in flies raised on standard wrack, (purple) higher in flies raised on field wrack.

Compound number	RI	Identified as	Methyl group position	Sex in Sex & Country	Country in Sex & Country	Sex in Sex & Diet	Wrack type in Sex & Diet
1	2096	unknown alkane					* (purple)
2	2101	C21		(blue)	* (green)		(purple)
3	2176	unknown alkane			* (green)		* (yellow)
5	2202	C22		* (blue)			
7	2300	C23		* (blue)	* (green)	* (blue)	
8	2344	unknown alkane		* (blue)	* (green)	* (blue)	
9	2365	2-ME-C23			* (green)		
11	2400	C24		* (blue)		* (blue)	* (purple)
14b	2478	25:x-C25			* (green)		
16	2500	C25					* (purple)
17	2524	unknown alkane					* (purple)
20	2564	2-ME-C25					* (purple)
21	2576	3-ME-C25					* (purple)
22	2584	5,X-Me-C25	X=9,11,13,15	* (blue)		* (blue)	* (purple)
23	2600	C26					* (purple)
24	2638	unknown alkane		* (red)			
26	2665	2-Me-C26					* (purple)
27	2680	27:x-C27		* (red)		* (red)	
29	2701	C27			* (grey)		* (purple)
30	2712	unknown alkane					* (purple)
31	2735	11+13-Me-C27			* (grey)		* (purple)
33	2765	2-Me-C27 + 11,15-Me-C27			* (grey)		* (purple)
34	2775	3-ME-C27					* (purple)
37a	2807	3,X-Me-C27	X=15,13,11,7				* (purple)
40	2850	29:x,x-C29				* (red)	* (purple)
41	2857	29:x,x-C29			* (green)		
42	2865	2-ME-C28 or x:C29			* (grey)		* (purple)
44	2880	29:x-C29				* (red)	* (purple)
45	2882	29:x-C29		* (red)	* (grey)		
46	2891	4,12-Me-C28					* (purple)
47	2902	C29			* (grey)		* (purple)
48	2933	15+13+11-Me-C29			* (grey)		* (purple)
50	2961	11,X-ME-C29	X=15,17	* (blue)	* (grey)	* (blue)	
51	2972	7,X-ME-C29	X=11,13,15	* (blue)	* (grey)	* (blue)	
52	2982	5,13-Me-C29		* (blue)	* (grey)		* (purple)
53	3007	3,13-ME-C29			* (grey)		
54	3033	3,X,15-Me-C29	X= 11,13	* (blue)		* (blue)	
55	3042	3,7,15-Me-C29			* (green)		
56	3054	31:x,x-C31		* (red)			
57	3058	12,16-Me-C30			* (green)		
60	3074	31:x-C31 or 6,16-Me-C30			* (grey)		
61	3082	31:x-C31		* (red)	* (grey)	* (red)	
62	3090	31:x-C31 or 4,16-Me-C30		* (blue)			
63	3101	C31		* (blue)	* (grey)	* (blue)	* (purple)
64	3118	unknown alkane		* (blue)	* (grey)		* (purple)
65	3131	15+13+11+9-Me-C31		* (blue)		* (blue)	* (purple)
66	3157	15,19-Me-C31		* (blue)		* (blue)	
67	3163	9,X-Me-C31	X=13,15	* (blue)	* (grey)	* (blue)	
68	3170	7,X-Me-C31	X=15,17	* (blue)		* (blue)	
69	3180	5,15-Me-C31		* (blue)		* (blue)	
71	3194	unknown alkane		* (blue)		* (blue)	
72	3205	3,X-Me-C31	X=9,11,13	* (blue)		* (blue)	* (purple)
73a	3231	3,X,Y-Me-C31	X=9,11 ; Y=13,15	* (blue)		* (blue)	
74	3246	33:x,x,x-C33		* (blue)			
75	3256	33:x,x-C33			* (green)		
76	3272	33:x-C33			* (green)		
77	3284	33:x-C33		* (blue)	* (grey)		
78	3300	unknown alkane		* (blue)	* (grey)	* (blue)	* (purple)
79	3315	unknown alkene					* (purple)
80	3330	17+15+13+11-Me-C33		* (blue)		* (blue)	* (purple)
81	3353	11,X-Me-C33	X=13,15,17	* (blue)			
82	3359	9,X-Me-C33	X=11,13,15		* (grey)		
83a	3365	7,X-Me-C33	X=11,13,15,17			* (blue)	
83b	3368	7,X-Me-C33	X=11,13,15,17	* (blue)		* (blue)	
84	3377	5,X-Me-C33	X=15,17	* (blue)		* (blue)	
85a	3391	7,13,17-Me-C33		* (blue)		* (blue)	
85b	3393	7,13,17-Me-C33		* (blue)			

Importantance in differentiation for factor

Legend:
- higher in females (red)
- higher in males (blue)
- higher in Swedish populations (grey)
- higher in Norwegian populations (green)
- higher in flies raised on standard wrack (yellow)
- higher in flies raised on field wrack (purple)
- * significant t-test with FDA α=0.05

Table 3 – Putative genes involved in CHC synthesis in *C. frigida*

Scaffold[1]	Gene Position[1]	Best Blast Hit[2]
scaffold_129	185034-197861	Desaturase 1
scaffold_155	109640-112144	Cytochrome P450-4G1
scaffold_206	225893-231312	Cytochrome P450 reductase
scaffold_480	98147-106166	CG5326 (elongase family)
scaffold_480	2453-12223	CG31522 (elongase family)
scaffold_545	1553-3217	CG5326 (elongase family)
scaffold_340	1141-16368	Baldspot (elongase family)
scaffold_149	7122-16098	Fatty Acid Synthase 3
scaffold_51	250672-259608	Fatty Acid Synthase 1
scaffold_716	36957-55451	CG2781 (elongase family)
scaffold_994	14376-17314	CG9743 (contains Fatty acid desaturase domain and Acyl-CoA desaturase)

1. From unpublished genome assembly (Wellenreuther *et. al*)
2. From a BLAST against *D. melanogaster* annotated nucleotides on flybase.org

Table 4 – Outlier SNPs putatively involved in geographic differences

Scaffold[1]	Position[1]	Variant type[2]	Best Blast Hit[3]	F_{ST}[4]
scaffold_129	195074	missense variant	Desaturase 1	0.16
scaffold_155	116876	downstream gene variant	Cytochrome P450-4G1	0.22
scaffold_206	225645	downstream gene variant	Cytochrome P450 reductase	0.17
scaffold_206	226804	missense variant	Cytochrome P450 reductase	0.17
scaffold_206	229444	missense variant	Cytochrome P450 reductase	0.20
scaffold_206	230032	intron variant	Cytochrome P450 reductase	0.30
scaffold_480	105896	synonymous variant	CG5326 (elongase family)	0.22
scaffold_480	110918	downstream gene variant	CG5326 (elongase family)	0.16
scaffold_994	16976	missense variant	CG9743 (contains Fatty acid desaturase domain and Acyl-CoA desaturase)	0.16
scaffold_994	18844	downstream gene variant	CG9743 (contains Fatty acid desaturase domain and Acyl-CoA desaturase)	0.20
scaffold_994	19574	downstream gene variant	CG9743 (contains Fatty acid desaturase domain and Acyl-CoA desaturase)	0.20

1. From unpublished genome assembly (Wellenreuther *et. al*)
2. Annotated in SnpEff (see text for details)
3. From a BLAST against *D. melanogaster* annotated nucleotides on flybase.org
4. Calculated from a separate unpublished data set of WGS from 46 *C. frigida*

Figure Legends

Figure 1 - Map of *C. frigida* populations used in this study. Base map from d-maps (http://d-maps.com/carte.php?num_car=5972)

Figure 2 – A. PCA of all samples from the balanced data set indicating population (grey – Østhassel, orange – Skeie, blue – Stavder, green – Ystad) and sex (triangle – female, star – male). B. OPLS-DA for sex using all samples from the balanced data set, C. PLS-DA for population using all samples from the balanced data set, and D. OPLS-DA for country using all samples from the balanced data set (yellow – Sweden, purple – Norway). Circles indicate 95% confidence regions.

Figure 3 – A. PCA of samples from Ystad indicating diet (blue – field, yellow – standard) and sex (triangle – female, star – male). B. OPLS-DA for sex using only Ystad samples. C. OPLS-DA for diet using only Ystad samples. Circles indicate 95% confidence regions.

Figure 4 – Boxplot of differences in A. Log transformed total peak area, B. proportion of compounds that are alkenes, C. proportion of methylated compounds. Females are in red, males are in blue.

Figure 5 – Boxplot of differences in A. Weighted mean chain length and B. Dispersion around the mean. Females are in red, males are in blue.

Skeie

Østhassel

Stavder

Ystad

200 km

100 mi

A) PCA

B) OPLS-DA for sex

C) PLS-DA for populations

D) OPLS-DA for countries

Sex

F
M

Country and populations

Norway
  Østhassel
  Skeie

Sweden
  Stavder
  Ystad

A) PCA

B) OPLS-DA for sex

C) OPLS-DA for wrack type