Genetic divergence and phenotypic plasticity contribute to variation in cuticular hydrocarbons in the seaweed fly *Coelopa frigida*

Emma Berdan[1*], Swantje Enge[2], Göran M. Nylund[3], Maren Wellenreuther[4,5], Gerrit A. Martens[6], and Henrik Pavia[3]

[1]Department of Marine Sciences, University of Gothenburg, Göteborg, Sweden

[2]Institute for Chemistry and Biology of the Marine Environment, Carl-von-Ossietzky University Oldenburg, Wilhelmshaven, Germany

[3]Department of Marine Sciences – Tjärnö, University of Gothenburg, Strömstad, Sweden

[4]The New Zealand Institute for Plant & Food Research Limited, Nelson, New Zealand

[5]School of Biological Sciences, the University of Auckland, New Zealand

[6]Hochschule Bremen, City University of Applied Sciences, Bremen, Germany

* - Corresponding author: Emma Berdan, Department of Marine Sciences, Box 461, 405 30 Göteborg, Sweden. Email: emma.berdan@gmail.com

ABSTRACT

Male sexual signals facilitate female choice and thereby reduce mating costs. The

seaweed fly *Coelopa frigida* is characterized by intense sexual conflict, yet it is unclear

which traits are under sexual selection. In this study, we investigated phenotypic and

genetic variation in cuticular hydrocarbons (CHCs) in *C. frigida* for the first time. CHCs

are ubiquitous in insects and they act as sex pheromone in many dipterans. We

characterized CHCs and found that composition was affected by both genetic (sex and

population origin) and environmental (larval diet) factors. We identified general shifts in

CHC chemistry as well as individual compounds and found that the methylated

compounds, mean chain length, proportion of alkenes, and normalized total CHCs

differed between sexes and populations. We combined this data with whole genome re-

sequencing data to examine the genetic underpinnings of these differences. We identified

11 genes related to CHC synthesis and found population level outlier SNPs in 5 that are

concordant with phenotypic differences. Together these results reveal that the CHC

composition of *C. frigida* is dynamic, strongly affected by the larval environment, and

likely under natural and/or sexual selection.

Keywords: cuticular hydrocarbons, *Coelopa frigida*, sexual conflict, sexual selection,

population differentiation, diet.

INTRODUCTION

Females and males often differ in optimal reproductive strategies, particularly regarding the mode and frequency of matings. Males can increase their fitness by mating with multiple females, while one or a few matings are generally sufficient for females to maximize their reproductive success (Bonduriansky, 2010). This sexual conflict over mating rates commonly leads to males harassing females to increase the likelihood that their genes get passed on to the next generation (Sánchez‑Guillén *et al.*, 2017). Such male mating harassment can carry considerable costs for females, e.g. by reducing reproductive success and increasing mortality (Gosden & Svensson, 2007). For example, repeated matings are the norm in the blue-tailed damselfly *Ischnura elegans*, particularly when population densities are high (Corbet, 1999), with females showing elevated immune responses, possibly because of mating related injuries or sexually transmitted diseases (Chauhan *et al.*, 2016). Likewise, in the guppy *Poecilia reticulata,* males attempt to copulate with females by thrusting their gonopodium towards the female's genital pore as often as once every minute, disrupting normal activities like foraging (Magurran & Nowak, 1991).

Females can minimize the cost of unwanted copulations by making it difficult for males to detect them (as seen in species with female limited colour polymorphism, Wellenreuther *et al.*, 2014) or evolving mate rejection responses (Andersson, 1994). However, these responses often carry associated costs (Watson *et al.*, 1998) and females therefore frequently endure costly matings with males when the costs of rejection

outweigh the costs of mating. Mating interactions can further be modulated by the

evolution of divergent inter-and intraspecific mate preferences. Mate preferences in

females, for example, can allow them to obtain high quality or compatible genes from a

male, with indirect benefits that include improved offspring fecundity, survival and

immune response (Neff & Pitcher, 2005). In many solitary and social insects, cuticular

hydrocarbons (CHCs) are used as one of the primary cues to recognize, and possibly

discriminate between species, sexes, and among kin (Bagneres *et al.*, 1996; Ferveur,

2005; Blomquist & Bagnères, 2010; Van Oystaeyen *et al.*, 2014).

Cuticular hydrocarbons are thought to be a primary adaptation to life on land because

they protect insects against desiccation (Wigglesworth, 1945; Blomquist & Bagnères,

2010). *De novo* synthesis of CHCs is conserved across insects (Howard & Blomquist,

2005; Blomquist & Bagnères, 2010): Synthesis takes place in oenocytes and starts with

acetyl-CoA. A fatty acid synthase (FAS) then elongates the acetyl-CoA, forming a long-

chained fatty acyl-CoA. FASs may also add methyl groups to these fatty acids (de

Renobales *et al.*, 1986; Blomquist *et al.*, 1994; Juarez *et al.*, 1996; Chung *et al.*, 2014).

Elongases then further lengthen these fatty acyl-CoAs and double bonds or triple bonds

are added by desaturases (Howard & Blomquist, 2005; Blomquist & Bagnères, 2010).

These are reduced to aldehydes by reductases and then these aldehydes are converted to

hydrocarbons by a decarboxylation reaction, which is mediated by a cytochrome P450

(Qiu *et al.*, 2012). Coding or expression variation in any number of these enzymes can

lead to significant changes in CHC composition (ex: de Renobales *et al.*, 1986; Reed *et*

al., 1995; Qiu *et al.*, 2012; Chung *et al.*, 2014; Chen *et al.*, 2016).

CHC composition is affected by both genetic and environmental factors. Changes in gene

expression (Dallerac *et al.*, 2000a; Chertemps *et al.*, 2007; Shirangi *et al.*, 2009;

Feldmeyer *et al.*, 2014) and/or coding sequences (Keays *et al.*, 2011; Badouin *et al.*,

2013; Kulmuni *et al.*, 2013) can affect both the overall composition of CHCs and the

length of specific compounds. Multiple environmental factors have also been shown to

influence CHC composition. For example, CHCs can vary under different temperature

regimes (Toolson & Kupersimbron, 1989; Savarit & Ferveur, 2002; Rouault *et al.*, 2004),

diets (Stennett & Etges, 1997; Liang & Silverman, 2000; Stojkovic *et al.*, 2014) or

climatic variables (Howard *et al.*, 1995; Kwan & Rundle, 2010). Moreover, multiple

studies have found that CHCs vary between natural populations (ex: Haverty *et al.*, 1990;

Etges & Ahrens, 2001; Frentiu & Chenoweth, 2010; Perdereau *et al.*, 2010), but it is

unknown how much of this variation is due to plasticity versus genetic divergence. These

can be teased apart using laboratory studies: genetic divergence can be assessed by

raising different populations in the same environment and plasticity can be assessed by

raising the same population in multiple environments (i.e. common garden and reciprocal

transplant experiments).

The seaweed fly *Coelopa frigida* is a common inhabitant of "wrackbeds" (accumulations

of decomposing seaweed) on shorelines. These wrackbeds act as both habitat and food

source for larvae and adults of this species. Fitness trade-offs and local adaptation in

response to wrackbed composition have been demonstrated in Swedish *C. frigida*

(Wellenreuther *et al.*, 2017). The mating system in *C. frigida* is characterized by intense

sexual conflict. Males will approach and forcefully mount females. Females are reluctant

to mate and use a variety of responses to dislodge the male, such as downward curling of

the abdomen (to prevent contact), kicking of the legs, and shaking from side to side (Day

et al., 1990; Blyth & Gilburn, 2011). Although there is no male courtship of any kind, it

has been consistently reported that mating attempts by large males are more likely to

result in copulation (Butlin *et al.*, 1982; Gilburn *et al.*, 1992; 1993). Two lines of

evidence show that females are actively choosing larger males rather than "giving up"

their struggle due to exhaustion. First, under "giving up" there should be a negative

correlation between struggle time and the size differential between the male and female.

However, there is absolutely no correlation between these factors (Edward, 2008).

Second, if females only allow copulation when they have been exhausted then struggle

times should differ between mountings in which copulation occurs and mountings where

copulation does not occur. However, there is no significant difference and a slight trend

towards longer struggle times for interactions that do not end in copulation (Edward,

2008). To date, the mechanism by which female choice occurs is currently unknown in

this system.

Here we investigate cuticular hydrocarbons as a possible male mating signal to test

whether this may be involved in female choice in *C. frigida.* Specifically, we examine if

CHC signatures vary between sexes, populations, and larval environment/diet

(wrackbed). We also explore putative genes involved in CHC synthesis and look for

correlations between genetic and phenotypic variation. We discuss our findings in light of

sexual selection in this species and others and outline future research areas that deserve

attention.

METHODS

Study Species

Coelopa frigida belongs to the group of acalyptrate flies which exclusively forage on

decomposing seaweed (Cullen *et al.*, 1987). This species is found along the seashores of

Northern Europe (Mcalpine, 1991), plays a vital role in coastal environmental

biodiversity (Griffin *et al.*, 2018) and health by accelerating the decomposition of algae,

allowing for faster release of nutrients (Cullen *et al.*, 1987). Additionally, larvae are an

important food source for predatory coastal invertebrates and seabirds (Summers *et al.*,

1990; Fuller, 2003).

Sampling

Fly larvae were collected in April and May 2017 from two Norwegian populations,

namely Skeie (58.69733, 5.54083) and Østhassel (58.07068, 6.64346), and two Swedish

populations, namely Stavder (57.28153, 12.13746) and Ystad (55.425, 13.77254). These

populations are situated along an environmental cline in salinity, wrackbed composition,

and wrackbed microbiome (Berdan et al., in prep; Day *et al.*, 1983) from the North Sea to

the Baltic Sea. Larvae were transported to Tjärnö Marine Laboratory of the University of

Gothenburg where they completed development in an aerated plastic pot filled with their

own field wrack in a temperature controlled room at 25 °C with a 12h/12h light-dark

cycle. After adults eclosed in the lab, they were transferred to a new pot filled with

standard wrack consisting of 50 % *Fucus spp*. and 50 % *Saccharina latissima* which had

been chopped, frozen, and then defrosted. The exception to this is a subset of Ystad

adults that were allowed to mate on the same material they had been collected in (field

wrack). Thus we had 5 treatment groups: All four populations (Skeie, Østhassel, Stavder

,Ystad) on standard wrack, and Ystad also on field wrack. Adult flies were allowed to lay

eggs on the provided wrack and this second generation was raised entirely in a

temperature controlled room at 25 °C with a 12 h/12 h light dark cycle. As larvae pupated

they were transferred to individual 2 mL tubes with a small amount of cotton soaked in

5 % (w/v) glucose to provide moisture and food for the eclosing adult. This ensured

virginity in all eclosing flies. The tubes were checked every day and the date of eclosure

of each fly was noted. Two days after eclosure adult flies were frozen at -80 °C and kept

there until CHC extraction.

<u>CHC Analysis</u>

Frozen flies were placed in 12-well porcelain plates to defrost and dry, as moisture can

affect lipid dissolution. Each fly was then placed in a 1.5 ml high recovery vial

containing 300 µl of *n*-hexane, vortexed at a low speed for 5 second, and extracted for 5

minutes. Afterwards flies were removed from the vial and allowed to air dry on the

porcelain plates before they were weighed (Sartorius Quintix 124-1S microbalance) to

the nearest 0.0001 grams and sexed. Extracts were evaporated until dry under nitrogen

gas and then stored at -20°. Before analysis 20 µl of *n*-hexane containing 1 µg/ml *n*-

nonane was added as an internal standard to each vial, which was then vortexed at

maximum speed for 10 seconds.

The extracts from 20 male and 20 female flies from every treatment group were analyzed

on a GC (Agilent GC 6890) coupled to a MS (Agilent 5973 MSD) using a HP-5MS

capillary column (Agilent) (see Table S1 in supplementary materials for instrument

settings).

The total ion chromatograms were quality checked, de-noised and Savitzky-Golay

filtered before peak detection and peak area integration in OpenChrom®. Peaks were

then aligned using the R-package 'GCalignR'(Ottensmann, Stoffel, & Hoffman, 2017)

prior to statistical analysis. Peaks were tentatively identified by their mass spectra and

retention time index based on a C21-C40 *n*-alkane standard solution (Sigma-Aldrich).

Statistical Analysis

The effects of sex and population on CHC profile were assessed using flies raised on

standard wrack using a balanced sampling design (N=13 for each combination) and 127

peaks were identified by the R package "GCalignR". The effects of sex and diet (i.e.

wrack type) during the larval stage on CHC profiles were analyzed using a balanced

design of flies from the Ystad population (N=12 for each combination) and 111 peaks

were identified by R package "GCalignR". Peak areas were normalized on the peak area

of the internal standard and based on the weight of the fly before being auto-scaled prior

to statistical analysis. Clustering of samples was visualized with a PCA, and group

differences were analyzed using a PERMANOVA followed by multiple group

comparisons using the PRIMER-E 7 software. We also analyzed group differences via

(O)PLS-DA using the "ropls"-packages in R (Thevenot, Roux, Xu, Ezan, & Junot, 2015).

Candidate compounds for group differentiation were determined from variables of

importance for OPLS-DA projection (VIP scores) (Galindo‑Prieto *et al.*, 2014), which

were further assessed by univariate analysis using a false discovery adjusted significance

level of $\alpha = 0.05$ using the Benjamini and Hochberg's FDR-controlling procedure

(Benjamini & Hochberg, 1995).

We examined sex and country effects in more depth to examine shifts in chemistry that

could be linked to our genetic data (see below). Our results above indicated that the

differences in CHC profiles between sexes and populations of *C. frigida* were due to

quantitative rather than qualitative differences (i.e. changes in relative amounts rather

than presence/absence of certain compounds). Thus, we looked for general shifts in CHC

chemistry by examining 1. The total normalized peak area (i.e. the sum of all peaks used

in analysis), 2. The proportion of alkenes, and 3. The proportion of methylated

compounds. For #1 we used all 127 peaks used in the previous analysis and for #2 and #3

we used the subset of these peaks that could be accurately categorized (116 peaks). For

the total peak area we transformed the data with a log transformation and then used the

'dredge' function from the MuMIn package (Barton, 2017) to compare AIC values for

nested glm (Gaussian distribution, link=identity) models with the terms sex, country, and

country*sex. If there were two models that differed in AIC by less than 1, we chose the

simpler model. For alkenes and methylated compounds our data was proportional and did

not include 1 or 0 so we used the 'betareg' function from the betareg package (Grun *et*

al., 2012). We again used the 'dredge' function to compare AIC values for nested models

with the terms sex, country, and country*sex and took the simpler model in case of an

AIC difference of less than 1.

We also analysed the distribution of chain lengths in our samples. To do this, we

subsetted all peaks where we had accurate length information (107 peaks) and then

calculated the weighted mean chain length as

$$N = \Sigma\ l_i p_i$$

where l_i is the length of a carbon chain and p_i is the proportional concentration of all

chains of that length. We also calculated dispersion around this mean as

$$D = \Sigma\ p_i(l_i - N)^2$$

We analyzed these in the same manner as described above, using AIC values to conduct

model choice from glm models (Gaussian distribution, link=identity).

208   Candidate gene analysis

Based on a literature search for genes involved in CHC synthesis, we identified 37

reference protein sequences, which were downloaded from Flybase (flybase.org) in

October 2017. We used tblastx with the default settings to identify orthologs in the *C.*

frigida transcriptome (Berdan & Wellenreuther, unpublished). We then used gmap (Wu

& Watanabe, 2005) to find areas of the *C. frigida* genome (Wellenreuther, *et al*.,

unpublished) that might contain these genes. We output all alignments as a gff3 file. We

have a previously made a VCF file created from 30x whole genome re-sequencing of 46

C. frigida across five populations in Scandinavia (Berdan *et. al*, unpublished). This VCF

file was made by aligning raw reads to the *C. frigida* genome using bwa-mem (Li &

Durbin, 2010), duplicate reads were marked using 'picard' (http://picard.sourceforge.net).

SNPs were called with the Genome Analysis Toolkit (GATK; DePristo *et al*. 2011; Van

der Auwera *et al.* 2013) specifically the GATK-module 'UnifiedGenotyper' (Van der

Auwera *et al.* 2013). VCF filtering was done as described in Berdan *et al.* 2015. We used

this VCF file and the gff3 output from gmap in SNPeff (Cingolani *et al.*, 2012) to

annotate the SNPs in these genes.

To find which SNPs are diverged between populations we used vcftools (Danecek *et al.*,

2011) to calculate Cockerham and Weir's F_{ST} estimate (Weir & Cockerham, 1984) for all

SNPs in our reference genome. We retained the top 5% of this distribution ($F_{ST} \geq 0.142$,

mean F_{ST} 0.024) as SNPs that were potentially divergent. We then subset the VCF file to

include only outlier SNPs located in our previously identified genomic regions.

RESULTS

CHC analysis

232 A diverse blend of hydrocarbons with more than 100 different compounds were found in

233 the cuticular extract of *C. frigida*. The majority of CHCs ranged in chain length from 23

234 to 33 carbon atoms. They contained odd numbered *n*-alkanes, methyl-, dimethyl- and

235 trimethyl-alkanes, as well as odd numbered alkenes. However, some even numbered

236 alkanes were also present but in lower quantities.

Sex and population effects

Results of the PERMANOVA indicated significant differences between females and

males and between populations but also a significant interaction between sex and

population (Table 1A). Pairwise tests on the interaction showed a significant difference

between males and females for all populations except Ystad (Table S2). Multiple

comparisons of populations showed that the Norwegian populations (Skeie and

Østhassel) were similar while the Swedish populations (Ystad and Stavder) differed for

males but not for females. In general the Norwegian populations tended to differ from the

Swedish and a similar pattern was also shown in the results of the PCA and PLS-DA

(Figure 2A, C, Table S3). After combining the Swedish and Norwegian populations, the

PERMANOVA indicated highly significant differences between females and males as

well as between Sweden and Norway (Table 1B). OPLS-DA models for both sex and

country were significant and lead to separate clusters of females and males, and Sweden

and Norway, respectively (Figure 2B, D, Table S3). The OPLS-DA model revealed 41

peaks that contributed with a VIP score >1 to the differentiation between females and

males. Of these, 31 were also significant in t-tests with a FDR adjusted p-values (Table

2). Generally compounds were more abundant in males with the exception of only 3

peaks that were more abundant in females. Looking at country differentiation between

Swedish and Norwegian populations we found 43 compounds with a VIP score >1, of

which 29 were significant in t-tests with FDR adjusted p-values and consisted of an equal

mix of compounds more abundant in either of the countries (Table 2). Differences in

CHC profiles between sexes and populations were due to quantitative differences,

affecting a wide range of compounds instead of qualitative differences, i.e. the presence/

absence of a few distinct compounds

<u>Sex and diet effects</u>

PERMANOVA demonstrated a significant difference between females and males in the

Ystad population and also a significant effect of the larval diet (wrack type) (Table 1C).

Both OPLS-DA models for either sex or wrack type were significant and lead to separate

clusters of females and males and of flies raised on standard or field wrack (Figure 3B,C,

Table S3). The OPLS-DA model revealed 38 peaks that contributed to the differentiation

between females and males with a VIP score >1. Of these, 24 compounds were also

significant in a t-test with FDR adjusted p-values (Table 2). All of these, except for 3

peaks, were more abundant in males. Overall 65% (26 compounds) were shared between

this analysis and the previous analysis investigating sex and population effects. Looking

at differentiation between flies raised on a diet consisting of standard or field wrack we

found 33 compounds with a VIP score >1. Of these, 27 compounds were significant

using a t-test with FDR adjusted p-values, all with higher amounts in flies reared on field

wrack (Table 2). Flies reared on field wrack had more CHCs per gram body weight than

flies reared on standard wrack (normalized totals: Females field wrack 7.96 ± 0.94,

females standard wrack 4.09 ± 0.47, males field wrack 11.24 ± 1.03, males standard

wrack 7.09 ± 1.21). Differences in CHC profiles between sexes and diets were

exclusively due to quantitative differences, affecting a wide range of compounds instead

of qualitative differences, i.e. the presence/ absence of a few distinct compounds

<u>Shifts in Chemistry</u>

As our results above indicated an overall effect of country rather than population we used

country and sex for our terms in the glm models. For total normalized peak area, the best

model contained only sex as a main effect (for a full AIC comparison for all models see

Table S4A-E). In general, males had more CHCs per gram body weight than females

(Figure 4A). For the proportion of alkenes, the best model contained country and sex as

main effects (for type II analysis of deviance tables for all best models see Table S5A-E).

Females tended to have more alkenes than males and individuals from Norway had more

alkenes than individuals from Sweden (Figure 4B). The best model for the proportion of

methylated compounds contained sex and country as the main effects. As with alkenes,

females tended to have more methylated compounds than males and individuals from

Norway tended to have more methylated compounds than individuals from Sweden

(Figure 4C).

The best model for weighted mean chain length was country and sex. Overall, males had

longer mean chain lengths than females and individuals from Sweden had longer mean

chain lengths than individuals from Norway (Figure 5A). The best model for dispersion

around mean chain length was also country and sex. Males had higher dispersion than

females and individuals from Norway had higher dispersion than individuals from

Sweden (Figure 5B).

Genetic analysis

We identified 15 isoforms from 13 transcripts in our transcriptome with matches to our

query proteins. These transcripts mapped to 16 unique areas of the genome assembly. We

blasted these areas against the NCBI nr database (accessed in April 2018) to confirm our

annotation. Five of these areas had either no match or matched to an unrelated protein

and were thus removed. The remaining genes include 2 putative fatty acid synthases, 2

putative desaturases, 5 putative elongases, 1 putative cytochrome P450 reductase, and 1

putative cytochrome P450-4G1 (Table 3). We found 1042 SNPs within these genes

(including 5K upstream and 5K downstream), of these 11 had an F_{ST} value > 0.142 (i.e.

in the top 5% of the F_{ST} distribution, Table 4).

DISCUSSION

In species with sexual conflict females may minimize unwanted copulations via mate

rejection responses; however, these responses typically carry associated costs (Watson *et*

al., 1998). The ability to modulate mating outcomes and thus the level of male

harassment experienced by the female based on male traits is especially beneficial in

systems where no "courtship" exists. Here we explored a potential male trait, cuticular

hydrocarbons (CHCs), and detected considerable phenotypic CHC variation attributable

to sex, population, and diet and genetic variation in candidate genes involved in CHC

synthesis that was attributable to population. We discuss these results and their

consequences below.

Males and females differed in their CHC composition. Not only did the composition

between males and females differ (Table 2, Figure 2-5) but, in addition to that, males

consistently had more CHC per gram body weight than females (Figure 4A). This is a

strong indication that CHCs play a role in sexual communication in *C. frigida* under a

female choice scenario, as it is unlikely that desiccation risk would differ between males

and females. A role for CHCs in sexual selection in *C. frigida* would be in line with the

general finding that CHCs play a major role in communication in insects, specifically in

Diptera (Howard & Blomquist, 2005; Blomquist & Bagnères, 2010; Ferveur & Cobb,

2010). Differences in CHCs have been shown to be involved in a wide variety of

behaviours in diptera such as courtship, aggregation, and dominance (Ferveur & Cobb,

2010). For instance, 11-*cis*-vaccenyl acetate, a male specific hydrocarbon in *Drosophila*

melanogaster, increases male-male aggression while suppressing male mating (Wang &

Anderson, 2010). The same compound is also involved in aggregation behavior in both

D. simulans and *D. melanogaster* (Bartelt *et al.*, 1985; Schaner *et al.*, 1987). Other

aggregation hydrocarbons, such as (Z)-10-heneicosene, in *D. virilis* have been shown to

attract males and females of certain ages (Bartelt & Jackson, 1984). Behavioural tests

will be necessary to confirm that CHCs are used for mate selection in *C. frigida* but

attention should also be paid to their possible role in behaviours such as aggression and

aggregation. This may be particularly relevant to *Coelopa frigida* as this species has been

described as forming large aggregations (Mather, 2011).

We found a shift in the composition between sexes, populations, and diets rather than

differences in the presence/absence of specific compounds. This type of pattern differs

from some Dipterans where sex and population differences are mostly qualitative

(Carlson & Schlein, 1991; Ferveur & Sureau, 1996; Dallerac *et al.*, 2000b; Gomes *et al.*,

2008; Everaerts *et al.*, 2010) although many dipterans show quantitative differences (ex:

Jallon & David, 1987; Byrne *et al.*, 1995). This discrepancy is likely to be partially

explained by the distribution of compounds that make up the CHCs in *C. frigida*. While

C. frigida chain lengths and compound classes are comparable with other dipteran

species (Ferveur & Cobb, 2010) the distribution of these compounds is somewhat

different. Many dipterans have principal CHC components that make up a large

proportion of the CHC composition (Ferveur & Sureau, 1996; Etges & Jackson, 2001;

Gomes *et al.*, 2008). Cuticular extracts from *C. frigida* have a large number of

compounds, but do not appear to have specific compounds that are dominant in either sex

or any of the populations. When averaging across all samples, the most abundant

compound was the C25 alkane, which made up on average 15% of the CHCs.

We examined coding variation to determine if we could tie population level phenotypic

differences to genetic differences. We observed country level variation with strong

differences between Norway and Sweden (Figure 2D, Figure 4, Figure 5). This pattern

mirrors a strong genetic split between the two countries (representing approximately 14%

of genetic variation, Berdan *et al*, unpublished). We found potential outlier SNPs in

desaturases, elongases, a cytochrome P450-4G1, and a cytochrome P450 reductase.

Desaturases add double bonds or triple bonds to alkanes (Howard & Blomquist, 2005;

Blomquist & Bagnères, 2010). We found an outlier missense variant in a putative

Desaturase 1 as well as a missense and two downstream outlier variants in a putative

Desaturase (Table 4). This aligns with our phenotypic data showing differences in the

proportion of alkanes vs. alkenes between Norway and Sweden. Elongases lengthen fatty

acyl-CoAs and are necessary for chains longer than 16 carbon atoms (Blomquist &

Bagnères, 2010). We found both synonymous and downstream outlier variants in putative

C. frigida elongases (Table 4) and corresponding differences in both the mean chain

length and variance of the chain length (dispersion around the mean) due to country of

origin. P450 reductases are responsible for reducing fatty acids to aldehydes (Blomquist

& Bagnères, 2010). Knockdown of P450 reductases in *D. melanogaster* leads to a

striking reduction of cuticular hydrocarbons (Qiu *et al.*, 2012). P450-4G genes, which are

unique to insects, encode an oxidative decarboxylase that catalyzes the aldehyde to

hydrocarbon reaction. Work in other insect species has found that a knockdown or

knockout of this gene leads to a decrease in overall hydrocarbon levels (Reed *et al.*, 1995;

Qiu *et al.*, 2012; Chen *et al.*, 2016). We found a downstream outlier variant in our

cytochrome P450-4G1 and multiple kinds of outlier variants in a cytochrome P450

reductase. However, the total amount of CHCs did not differ between populations, only

sexes. As males and females share a genome (with the exception of the sex chromosome)

it is unlikely that these SNPs affect this difference. Finally, we also found variation

between Sweden and Norway in the proportion of methylated compounds. Other studies

suggest that differences in the proportion of methylated compounds are often caused by

either variation in Fatty Acid Synthase (FAS, de Renobales *et al.*, 1986; Blomquist *et al.*,

1994; Juarez *et al.*, 1996) or multiple copies of FAS with different functions (Chung *et*

al., 2014). We found two different FAS' in our genome (Table 3) but neither of them

contained divergent SNPs. Future studies are needed to examine if these genetic and

phenotypic changes are related. Specifically, investigations into both RNA expression

and tests of function of these loci will be necessary to provide a direct link. However, the

association between the known function of several of these genes (elongases, desaturases)

and the corresponding differences in populations are marked. Given that we see shifts in

multiple chemical aspects (length, methylation, and alkene/alkane ratio) it is likely that

multiple loci are responsible for these patterns.

The wrackbed itself also had a strong influence on the CHC composition (Figure 3C).

Wrackbed composition and microbiome (i.e. the food source for *C. frigida* larvae) varies

across *C. frigida* populations in Europe (Berdan, et al., in prep, Day *et al.*, 1983; Butlin &

Day, 1989; Wellenreuther *et al.*, 2017), indicating that natural populations may be have

different CHC composition. This is concordant with other research showing that larval

diet impacts CHC composition (Liang & Silverman, 2000; Rundle *et al.*, 2005; Etges &

de Oliveira, 2014; Stojkovic *et al.*, 2014). Adding to this, we found strong population

(country) level differences in CHC composition even when larvae were reared on the

same substrate. We also found several outlier SNPs in a variety of genes involved in

CHC synthesis. Together these results indicate several possibilities: 1. Natural selection

may select for a different CHC composition in different environments and natural

polymorphism in CHC genes may be under direct selection due to this. For example,

Drosophila serrata and *D. birchii* differ in the environment they inhabit (habitat

generalist vs. habitat specialist for humid rainforests). These two species also differ in the

concentration of methyl branched CHCs (especially important in desiccation resistance)

caused, in part, by changes in the cis-regulatory sequence likely under natural selection

(Chung *et al.*, 2014). 2. Natural populations differ strongly in their CHC composition. If

assortative mating or selection on female preferences is present that could lead to locally

adapted male traits and female preferences (Chung & Carroll, 2015). This would reduce

effective migration between populations, as foreign males would be at a disadvantage.

Differences in CHC composition and corresponding preferences have been shown to

cause behavioral isolation between many different *Drosophila* species or populations

(Coyne *et al.*, 1994; Rundle *et al.*, 2005; Etges & de Oliveira, 2014).

In conclusion, the analysis of cuticular hydrocarbon extracts from male and female *C.*

frigida from multiple populations revealed a complex and variable mix of more than 100

different compounds. We found extensive phenotypic variation attributable to diet, sex,

and country as well as associated genetic variation attributable to the latter. This reveals

that CHC composition is dynamic, strongly affected by the larval environment, and most

likely under natural and/or sexual selection. Further work is needed to confirm that CHCs

are used as sexual signals in this system and to explore the evolutionary consequences of

these differences.

434 ACKNOWLEDGEMENTS

We thank the members of the Marine Chemical Ecology group at Gothenburg University
for helpful comments on experimental design. E.B. was supported by a Marie
Skłodowska-Curie fellowship 704920 — ADAPTIVE INVERSIONS — H2020-MSCA-
IF-2015. MW and the research were supported by grants from Vetenskapsrådet to MW
(2012-3996), from the Mistra Foundation to HP (AquaAgriKelp) and from the Swedish
Foundation for Strategic Research to HP (Sweaweed).

REFERENCES

Andersson, M.B. 1994. *Sexual selection*. Princeton University Press.

Badouin, H., Belkhir, K., Gregson, E., Galindo, J., Sundström, L., Martin, S.J., Butlin, R.K. & Smadja, C.M. 2013. Transcriptome characterisation of the ant Formica exsecta with new insights into the evolution of desaturase genes in social hymenoptera. *PloS one* **8:** e68200.

Bagneres, A.G., Lorenzi, M.C., Dusticier, G., Turillazzi, S. & Clement, J.L. 1996. Chemical usurpation of a nest by paper wasp parasites. *Science* **272:** 889-892.

Bartelt, R. & Jackson, L. 1984. Hydrocarbon component of the Drosophila virilis (Diptera: Drosophilidae) aggregation pheromone:(Z)-10-heneicosene. *Annals of the Entomological Society of America* **77:** 364-371.

Bartelt, R.J., Schaner, A.M. & Jackson, L.L. 1985. cis-Vaccenyl acetate as an aggregation pheromone inDrosophila melanogaster. *Journal of chemical ecology* **11:** 1747-1756.

Barton, K. 2017. MuMIn.

Berdan, E.L., Mazzoni, C.J., Waurick, I., Roehr, J.T. & Mayer, F. 2015. A population genomic scan in *Chorthippus* grasshoppers unveils previously unknown phenotypic divergence. *Molecular Ecology* **24:** 3918-3930.

Blomquist, G.J., Guo, L., Gu, P., Blomquist, C., Reitz, R.C. & Reed, J.R. 1994. Methyl-branched fatty acids and their biosynthesis in the housefly, Musca domestica L.(Diptera: Muscidae). *Insect biochemistry and molecular biology* **24:** 803-810.

Blomquist, G.J. & Bagnères, A.-G. 2010. *Insect hydrocarbons: biology, biochemistry, and chemical ecology*. Cambridge University Press.

Blyth, J.E. & Gilburn, A.S. 2011. The function of female behaviours adopted during premating struggles in the seaweed fly, Coelopa frigida. *Animal Behaviour* **81:** 77-82.

Bonduriansky, R. 2010. Sexual conflict, John Wiley & Sons, Ltd. Chichester.

Butlin, R. & Day, T. 1989. Environmental correlates of inversion frequencies in natural populations of seaweed flies (Coelopa frigida). *Heredity* **62:** 223.

Butlin, R.K., Read, I.L. & Day, T.H. 1982. The effects of a chromosomal inversion on adult size and male mating success in the seaweed fly, *Coelopa frigida*. *Heredity* **49:** 51-62.

Byrne, A., Camann, M., Cyr, T., Catts, E. & Espelie, K. 1995. Forensic implications of biochemical differences among geographic populations of the black blow fly, Phormia regina (Meigen). *Journal of Forensic Science* **40:** 372-377.

Carlson, D. & Schlein, Y. 1991. Unusual polymethyl alkenes in tsetse flies acting as abstinon inGlossina morsitans. *Journal of chemical ecology* **17:** 267-284.

Chauhan, P., Wellenreuther, M. & Hansson, B. 2016. Transcriptome profiling in the damselfly *Ischnura elegans* identifies genes with sex-biased expression. *BMC genomics* **17:** 985.

Chen, N., Fan, Y.L., Bai, Y., Li, X.D., Zhang, Z.F. & Liu, T.X. 2016. Cytochrome P450 gene, CYP4G51, modulates hydrocarbon production in the pea aphid, Acyrthosiphon pisum. *Insect Biochemistry and Molecular Biology* **76:** 84-94.

Chertemps, T., Duportets, L., Labeur, C., Ueda, R., Takahashi, K., Saigo, K. & Wicker-Thomas, C. 2007. A female-biased expressed elongase involved in long-chain hydrocarbon biosynthesis and courtship behavior in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America* **104:** 4273-4278.

Chung, H., Loehlin, D.W., Dufour, H.D., Vaccarro, K., Millar, J.G. & Carroll, S.B. 2014. A Single Gene Affects Both Ecological Divergence and Mate Choice in Drosophila. *Science* **343:** 1148-1151.

Chung, H. & Carroll, S.B. 2015. Wax, sex and the origin of species: Dual roles of insect cuticular hydrocarbons in adaptation and mating. *Bioessays* **37:** 822-830.

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X.Y. & Ruden, D.M. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w(1118); iso-2; iso-3. *Fly* **6:** 80-92.

Corbet, P. 1999. *Dragonflies: Behavior and Ecology of Odonata*. Essex: Harley Books.

Coyne, J.A., Crittenden, A.P. & Mah, K. 1994. Genetics of a pheromonal difference contributing to reproductive isolation in Drosophila. *Science* **265:** 1461-1464.

Cullen, S.J., Young, A.M. & Day, T.H. 1987. Dietary requirements of seaweed flies (Coelopa frigida). *Estuarine, Coastal and Shelf Science* **24:** 701-710.

Dallerac, R., Labeur, C., Jallon, J.-M., Knipple, D.C., Roelofs, W.L. & Wicker-Thomas, C. 2000a. A Δ9 desaturase gene with a different substrate specificity is responsible for the cuticular diene hydrocarbon polymorphism in Drosophila melanogaster. *Proceedings of the National Academy of Sciences* **97:** 9449-9454.

Dallerac, R., Labeur, C., Jallon, J.M., Knippie, D.C., Roelofs, W.L. & Wicker-Thomas, C. 2000b. A Delta 9 desaturase gene with a different substrate specificity is responsible for the cuticular diene hydrocarbon polymorphism in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America* **97:** 9449-9454.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R. & Grp, G.P.A. 2011. The variant call format and VCFtools. *Bioinformatics* **27:** 2156-2158.

Day, T.H., Dawe, C., Dobson, T. & Hillier, P.C. 1983. A chromosomal inversion polymorphism in Scandinavian populations of the seaweed fly, Coelopa frigida. *Hereditas* **99:** 135-145.

Day, T.H., Foster, S.P. & Engelhard, G. 1990. Mating behavior in seaweed flies (Coelopa frigida). *Journal of Insect Behavior* **3:** 105-120.

de Renobales, M., Woodin, T.S. & Blomquist, G.J. 1986. Drosophila melanogaster fatty acid synthetase: characteristics and effect of protease inhibitors. *Insect biochemistry* **16:** 887-894.

Edward, D.A. 2008. Habitat composition, sexual conflict and life history evolution in Coelopa frigida.

Etges, W.J. & Ahrens, M.A. 2001. Premating isolation is determined by larval-rearing substrates in cactophilic Drosophila mojavensis. V. Deep geographic variation in epicuticular hydrocarbons among isolated populations. *The American Naturalist* **158:** 585-598.

Etges, W.J. & Jackson, L.L. 2001. Epicuticular hydrocarbon variation in Drosophila mojavensis cluster species. *Journal of chemical ecology* **27:** 2125-2149.

Etges, W.J. & de Oliveira, C.C. 2014. Premating isolation is determined by larval rearing substrates in cactophilic Drosophila mojavensis. X. Age-specific dynamics of adult epicuticular hydrocarbon expression in response to different host plants. *Ecology and Evolution* **4:** 2033-2045.

Everaerts, C., Farine, J.P., Cobb, M. & Ferveur, J.F. 2010. Drosophila Cuticular Hydrocarbons Revisited: Mating Status Alters Cuticular Profiles. *Plos One* **5**.

Feldmeyer, B., Elsner, D. & Foitzik, S. 2014. Gene expression patterns associated with caste and reproductive status in ants: worker‐specific genes are more derived than queen‐specific ones. *Molecular ecology* **23:** 151-161.

Ferveur, J.-F. 2005. Cuticular hydrocarbons: their evolution and roles in Drosophila pheromonal communication. *Behavior genetics* **35:** 279.

Ferveur, J.-F. & Cobb, M. 2010. Behavioral and evolutionary roles of cuticular hydrocarbons in Diptera. *Insect hydrocarbons: biology, biochemistry, and chemical ecology***:** 325-343.

Ferveur, J.F. & Sureau, G. 1996. Simultaneous influence on male courtship of stimulatory and inhibitory pheromones produced by live sex-mosaic Drosophila melanogaster. *Proceedings of the Royal Society B-Biological Sciences* **263:** 967-973.

Frentiu, F.D. & Chenoweth, S.F. 2010. Clines in cuticular hydrocarbons in two Drosophila species with independent population histories. *Evolution* **64:** 1784-1794.

Fuller, R.A. 2003. Factors influencing foraging decisions in ruddy turnstones Arenaria interpres (L.), Durham University.

Galindo‐Prieto, B., Eriksson, L. & Trygg, J. 2014. Variable influence on projection (VIP) for orthogonal projections to latent structures (OPLS). *Journal of Chemometrics* **28:** 623-632.

Gilburn, A., Foster, S. & Day, T. 1992. Female mating preference for large size in Coelopa frigida(seaweed fly). *Heredity* **69:** 209-216.

Gilburn, A., Foster, S. & Day, T. 1993. Genetic correlation between a female mating preference and the preferred male character in seaweed flies (Coelopa frigida). *Evolution* **47:** 1788-1795.

Gomes, C.C.G., Trigo, J.R. & Eiras, Á.E. 2008. Sex pheromone of the American warble fly, Dermatobia hominis: the role of cuticular hydrocarbons. *Journal of chemical ecology* **34:** 636.

Gosden, T. & Svensson, E.I. 2007. Female sexual polymorphism and fecundity consequences of male mating harassment in the wild. *PLoS ONE***:** 2(6):e580 doi:510.1371/journal.pone.0000580.

Griffin, C., Day, N., Rosenquist, H., Wellenreuther, M., Bunnefeld, N. & Gilburn, A. 2018. Tidal range and recovery from the impacts mechanical beach grooming. *Ocean and Coastal Management* **154:** 66-71.

Grun, B., Kosmidis, I. & Zeileis, A. 2012. Extended Beta Regression in R: Shaken, Stirred, Mixed, and Partitioned. *Journal of Statistical Software* **48:** 1-25.

Haverty, M.I., Nelson, L.J. & Page, M. 1990. Cuticular hydrocarbons of four populations of Coptotermes formosanus Shiraki in the united states. *Journal of chemical ecology* **16:** 1635-1647.

Howard, R.W., Howard, C.D. & Colquhoun, S. 1995. Ontogenetic and environmentally induced changes in cuticular hydrocarbons of Oryzaephilus surinamensis (Coleoptera: Cucujidae). *Annals of the Entomological Society of America* **88:** 485-495.

Howard, R.W. & Blomquist, G.J. 2005. Ecological, behavioral, and biochemical aspects of insect hydrocarbons. *Annu. Rev. Entomol.* **50:** 371-393.

Jallon, J.M. & David, J.R. 1987. Variations in cuticular hydrocarbons among the eight species of the Drosophila melanogaster subgroup. *Evolution* **41:** 294-302.

Juarez, M.P., Ayala, S. & Brenner, R.R. 1996. Methyl-branched fatty acid biosynthesis in Triatoma infestans. *Insect Biochemistry and Molecular Biology* **26:** 599-605.

Keays, M.C., Barker, D., WICKER‐THOMAS, C. & Ritchie, M.G. 2011. Signatures of selection and sex‐specific expression variation of a novel duplicate during the evolution of the Drosophila desaturase gene family. *Molecular ecology* **20:** 3617-3630.

Kulmuni, J., Wurm, Y. & Pamilo, P. 2013. Comparative genomics of chemosensory protein genes reveals rapid evolution and positive selection in ant-specific duplicates. *Heredity* **110:** 538-547.

Kwan, L. & Rundle, H.D. 2010. Adaptation to Desiccation Fails to Generate Pre- and Postmating Isolation in Replicate Drosophila Melanogaster Laboratory Populations. *Evolution* **64:** 710-723.

Li, H. & Durbin, R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26:** 589-595.

Liang, D. & Silverman, J. 2000. "You are what you eat": diet modifies cuticular hydrocarbons and nestmate recognition in the Argentine ant, Linepithema humile. *Naturwissenschaften* **87:** 412-416.

Magurran, A.E. & Nowak, M.A. 1991. Another battle of the sexes: the consequences of sexual asymmetry in mating costs and predation risk in the guppy, Poecilia reticulata. *Proc. R. Soc. Lond. B* **246:** 31-38.

Mather, K. 2011. South Bay beaches hit my swarms of kelp flies. *Los Angeles Times*. Los Angeles.

Mcalpine, D.K. 1991. Review of the Australian Kelp Flies (Diptera, Coelopidae). *Systematic Entomology* **16:** 29-84.

Neff, B.D. & Pitcher, T.E. 2005. Genetic quality and sexual selection: an integrated framework for good genes and compatible genes. *Mol Ecol* **14:** 19-38.

Perdereau, E., Dedeine, F., Christidès, J.-P. & Bagnères, A.-G. 2010. Variations in worker cuticular hydrocarbons and soldier isoprenoid defensive secretions within and among introduced and native populations of the subterranean termite, Reticulitermes flavipes. *Journal of chemical ecology* **36:** 1189-1198.

Qiu, Y., Tittiger, C., Wicker-Thomas, C., Le Goff, G., Young, S., Wajnberg, E., Fricaux, T., Taquet, N., Blomquist, G.J. & Feyereisen, R. 2012. An insect-specific P450 oxidative decarbonylase for cuticular hydrocarbon biosynthesis. *Proceedings of the National Academy of Sciences of the United States of America* **109:** 14858-14863.

Reed, J.R., Quilici, D.R., Blomquist, G.J. & Reitz, R.C. 1995. Proposed mechanism for the cytochrome P 450-catalyzed conversion of aldehydes to hydrocarbons in the house fly, Musca domestica. *Biochemistry* **34:** 16221-16227.

Rouault, J.D., Marican, C., Wicker-Thomas, C. & Jallon, J.M. 2004. Relations between cuticular hydrocarbon (HC) polymorphism, resistance against desiccation and breeding temperature; a model for HC evolution in D-melanogaster and D-simulans. *Genetica* **120:** 195-212.

Rundle, H.D., Chenoweth, S.F., Doughty, P. & Blows, M.W. 2005. Divergent selection and the evolution of signal traits and mating preferences. *Plos Biology* **3:** 1988-1995.

Sánchez‑Guillén, R.A., Wellenreuther, M., Chávez‑Ríos, J.R., Beatty, C.D., Rivas‑Torres, A., Velasquez‑Velez, M. & Cordero‑Rivera, A. 2017. Alternative reproductive strategies and the maintenance of female color polymorphism in damselflies. *Ecology and Evolution* **7:** 5592-5602.

Savarit, F. & Ferveur, J.F. 2002. Temperature affects the ontogeny of sexually dimorphic cuticular hydrocarbons in Drosophila melanogaster. *Journal of Experimental Biology* **205:** 3241-3249.

Schaner, A.M., Bartelt, R.J. & Jackson, L.L. 1987. (Z)-11-octadecenyl acetate, an aggregation pheromone inDrosophila simulans. *Journal of chemical ecology* **13:** 1777-1786.

Shirangi, T.R., Dufour, H.D., Williams, T.M. & Carroll, S.B. 2009. Rapid evolution of sex pheromone-producing enzyme expression in Drosophila. *PLoS biology* **7:** e1000168.

Stennett, M.D. & Etges, W.J. 1997. Premating Isolation Is Determined by Larval Rearing Substrates in CactophilicDrosophila mojavensis. III. Epicuticular Hydrocarbon Variation Is Determined by Use of Different Host Plants inDrosophila mojavensis andDrosophila arizonae. *Journal of Chemical Ecology* **23:** 2803-2824.

Stojkovic, B., Savkovic, U., Dordevic, M. & Tucic, N. 2014. Host-shift effects on mating behavior and incipient pre-mating isolation in seed beetle. *Behavioral Ecology* **25:** 553-564.

Summers, R.W., Smith, S., Nicoll, M. & Atkinson, N.K. 1990. Tidal and Sexual Differences in the Diet of Purple Sandpipers Calidris-Maritima in Scotland. *Bird Study* **37:** 187-194.

Toolson, E.C. & Kupersimbron, R. 1989. Laboratory Evolution of Epicuticular Hydrocarbon Composition and Cuticular Permeability in Drosophila-Pseudoobscura - Effects on Sexual Dimorphism and Thermal-Acclimation Ability. *Evolution* **43:** 468-473.

Van Oystaeyen, A., Oliveira, R.C., Holman, L., van Zweden, J.S., Romero, C., Oi, C.A., d'Ettorre, P., Khalesi, M., Billen, J. & Wäckers, F. 2014. Conserved class of queen pheromones stops social insect workers from reproducing. *Science* **343:** 287-290.

Wang, L.M. & Anderson, D.J. 2010. Identification of an aggression-promoting pheromone and its receptor neurons in Drosophila. *Nature* **463:** 227-U114.

Watson, P.J., Stallmann, R.R. & Arnqvist, G. 1998. Sexual conflict and the energetic costs of mating and mate choice in water striders. *The American Naturalist* **151:** 46-58.

Weir, B.S. & Cockerham, C.C. 1984. Estimating F-Statistics for the Analysis of Population-Structure. *Evolution* **38:** 1358-1370.

Wellenreuther, M., Svensson, E.I. & Hansson, B. 2014. Sexual selection and genetic colour polymorphisms in animals. *Molecular Ecology* **23:** 5398-5414.

Wellenreuther, M., Rosenquist, H., Jaksons, P. & Larson, W. 2017. Local adaptation along an environmental cline in a species with an inversion polymorphism. *Journal of Evolutionary Biology* **30:** 1068-1077.

Wigglesworth, V. 1945. Transpiration through the cuticle of insects. *Journal of Experimental Biology* **21:** 97-114.

Wu, T.D. & Watanabe, C.K. 2005. GMAP: a genomic mapping and alignment program for mRNA and EST sequences. *Bioinformatics* **21:** 1859-1875.

Tables

Table 1– PERMANOVA and PERMDISP results for A. Sex and Population, B. Sex and Country, and C. Sex and Wrack type. * < 0.05 ** < 0.01 *** < 0.001

A.

		PERMANOVA: Sex and Population					PERMDISP
	df	SS	MS	F	p-value		
Sex	1	658.1	658.1	5.86	<0.001	***	see S2
Population	3	1150.3	383.4	3.41	<0.001	***	
Sex:Population	3	486.9	162.3	1.44	0.015	*	
Residuals	96	10786.0	112.4				
Total	103	13081					

B.

		PERMANOVA: Sex and Country					PERMDISP		
	df	SS	MS	F	p-value		F	p-value	
Sex	1	658.0	658.1	5.60	<0.001	***	8.94	0.003	**
Country	1	517.8	517.8	4.41	<0.001	***	1.19	0.279	
Sex:Country	1	154.1	154.1	1.31	0.128				
Residuals	100	11751.0	117.5						
Total	103	13081							

C.

		PERMANOVA: Sex and Wrack type					PERMDISP	
	df	SS	MS	F	p-value		F	p-value
Sex	1	360.9	360.9	3.73	<0.001	***	2.36	0.129
Wrack	1	481.3	481.3	4.98	<0.001	***	2.29	0.139
Sex:Wrack	1	122.3	122.3	1.27	0.1629			
Residuals	44	4252.5	96.7					
Total	47	5217						

Table 2 – List of compounds that are significantly important for differentiation of at least factor (sex, diet, or country). If compounds are significant for differentiation of any category in any comparison a * appears under that factor. Color indicates the direction of the differentiation (red- higher in females, blue – higher in males, grey – higher in Norwegian populations, green – higher in Swedish populations, yellow – higher in flies raised on standard wrack, purple – higher in flies raised on field wrack).

Importantance in differentiation for factor

Compound number	RI	Identified as	Methyl group position	Sex in Sex & Country	Country in Sex & Country	Sex in Sex & Diet	Wrack type in Sex & Diet
1	2096	unknown alkane					*
2	2101	C21		(blue)	*		
3	2176	unknown alkane			*		(yellow)
5	2202	C22		*			
7	2300	C23		*	*	*	
8	2344	unknown alkane		*	*	*	
9	2365	2-ME-C23			*		
11	2400	C24		*		*	*
14b	2478	25:x-C25			*		
16	2500	C25					*
17	2524	unknown alkane					*
20	2564	2-ME-C25					*
21	2576	3-ME-C25					*
22	2584	5,X-Me-C25	X=9,11,13,15	*		*	*
23	2600	C26					*
24	2638	unknown alkane		*			
26	2665	2-Me-C26					*
27	2680	27:x-C27		*		*	
29	2701	C27			*		*
30	2712	unknown alkane					*
31	2735	11+13-Me-C27			*		*
33	2765	2-Me-C27 + 11,15-Me-C27			*		*
34	2775	3-ME-C27					*
37a	2807	3,X-Me-C27	X=15,13,11,7				*
40	2850	29:x,x-C29				*	*
41	2857	29:x,x-C29			*		
42	2865	2-ME-C28 or x:C29			*		*
44	2880	29:x-C29				*	*
45	2882	29:x-C29		*	*		
46	2891	4,12-Me-C28					*
47	2902	C29			*		*
48	2933	15+13+11-Me-C29			*		*
50	2961	11,X-ME-C29	X=15,17	*	*	*	
51	2972	7,X-ME-C29	X=11,13,15	*	*	*	
52	2982	5,13-Me-C29		*	*		*
53	3007	3,13-ME-C29			*		*
54	3033	3,X,15-Me-C29	X= 11,13	*		*	
55	3042	3,7,15-Me-C29			*		
56	3054	31:x,x-C31		*			
57	3058	12,16-Me-C30			*		
60	3074	31:x-C31 or 6,16-Me-C30			*		
61	3082	31:x-C31		*	*	*	
62	3090	31:x-C31 or 4,16-Me-C30		*			
63	3101	C31		*	*	*	*
64	3118	unknown alkane		*	*		*
65	3131	15+13+11+9-Me-C31		*		*	*
66	3157	15,19-Me-C31		*		*	
67	3163	9,X-Me-C31	X=13,15	*	*	*	
68	3170	7,X-Me-C31	X=15,17	*		*	
69	3180	5,15-Me-C31		*		*	
71	3194	unknown alkane		*		*	
72	3205	3,X-Me-C31	X=9,11,13	*		*	*
73a	3231	3,X,Y-Me-C31	X=9,11 ; Y=13,15	*		*	
74	3246	33:x,x,x-C33		*			
75	3256	33:x,x-C33			*		
76	3272	33:x-C33			*		
77	3284	33:x-C33		*	*		
78	3300	unknown alkane		*	*	*	*
79	3315	unknown alkene					*
80	3330	17+15+13+11-Me-C33		*		*	*
81	3353	11,X-Me-C33	X=13,15,17	*			
82	3359	9,X-Me-C33	X=11,13,15		*		
83a	3365	7,X-Me-C33	X=11,13,15,17			*	
83b	3368	7,X-Me-C33	X=11,13,15,17	*		*	
84	3377	5,X-Me-C33	X=15,17	*		*	
85a	3391	7,13,17-Me-C33		*		*	
85b	3393	7,13,17-Me-C33		*			

(red)	higher in females
(blue)	higher in males
(grey)	higher in Swedish populations
(green)	higher in Norwegian populations
(yellow)	higher in flies raised on standard wrack
(purple)	higher in flies raised on field wrack
*	significant t-test with FDA α=0.05

Table 3 – Putative genes involved in CHC synthesis in *C. frigida*

Scaffold[1]	Gene Position[1]	Best Blast Hit[2]
scaffold_129	185034-197861	Desaturase 1
scaffold_155	109640-112144	Cytochrome P450-4G1
scaffold_206	225893-231312	Cytochrome P450 reductase
scaffold_480	98147-106166	CG5326 (elongase family)
scaffold_480	2453-12223	CG31522 (elongase family)
scaffold_545	1553-3217	CG5326 (elongase family)
scaffold_340	1141-16368	Baldspot (elongase family)
scaffold_149	7122-16098	Fatty Acid Synthase 3
scaffold_51	250672-259608	Fatty Acid Synthase 1
scaffold_716	36957-55451	CG2781 (elongase family)
scaffold_994	14376-17314	CG9743 (contains Fatty acid desaturase domain and Acyl-CoA desaturase)

1. From unpublished genome assembly (Wellenreuther *et. al*)
2. From a BLAST against *D. melanogaster* annotated nucleotides on flybase.org

Table 4 – Outlier SNPs putatively involved in geographic differences

Scaffold[1]	Position[1]	Variant type[2]	Best Blast Hit[3]	F_{ST}[4]
scaffold_129	195074	missense variant	Desaturase 1	0.16
scaffold_155	116876	downstream gene variant	Cytochrome P450-4G1	0.22
scaffold_206	225645	downstream gene variant	Cytochrome P450 reductase	0.17
scaffold_206	226804	missense variant	Cytochrome P450 reductase	0.17
scaffold_206	229444	missense variant	Cytochrome P450 reductase	0.20
scaffold_206	230032	intron variant	Cytochrome P450 reductase	0.30
scaffold_480	105896	synonymous variant	CG5326 (elongase family)	0.22
scaffold_480	110918	downstream gene variant	CG5326 (elongase family)	0.16
scaffold_994	16976	missense variant	CG9743 (contains Fatty acid desaturase domain and Acyl-CoA desaturase)	0.16
scaffold_994	18844	downstream gene variant	CG9743 (contains Fatty acid desaturase domain and Acyl-CoA desaturase)	0.20
scaffold_994	19574	downstream gene variant	CG9743 (contains Fatty acid desaturase domain and Acyl-CoA desaturase)	0.20

1. From unpublished genome assembly (Wellenreuther *et. al*)
2. Annotated in SnpEff (see text for details)
3. From a BLAST against *D. melanogaster* annotated nucleotides on flybase.org
4. Calculated from a separate unpublished data set of WGS from 46 *C. frigida*

Figure Legends

Figure 1 - Map of *C. frigida* populations used in this study. Base map from d-maps (http://d-maps.com/carte.php?num_car=5972)

Figure 2 – A. PCA of all samples from the balanced data set indicating population (grey – Østhassel, orange – Skeie, blue – Stavder, green – Ystad) and sex (triangle – female, star – male). B. OPLS-DA for sex using all samples from the balanced data set, C. PLS-DA for population using all samples from the balanced data set, and D. OPLS-DA for country using all samples from the balanced data set (yellow – Sweden, purple – Norway). Circles indicate 95% confidence regions.

Figure 3 – A. PCA of samples from Ystad indicating diet (blue – field, yellow – standard) and sex (triangle – female, star – male). B. OPLS-DA for sex using only Ystad samples. C. OPLS-DA for diet using only Ystad samples. Circles indicate 95% confidence regions.

Figure 4 – Boxplot of differences in A. Log transformed total peak area, B. proportion of compounds that are alkenes, C. proportion of methylated compounds. Females are in red, males are in blue.

Figure 5 – Boxplot of differences in A. Weighted mean chain length and B. Dispersion around the mean. Females are in red, males are in blue.

Skeie

Østhassel

Stavder

Ystad

200 km

100 mi

© d-maps.com

A) PCA

B) OPLS-DA for sex

C) OPLS-DA for wrack type