1

IRES-mediated ribosome repositioning directs translation of a +1 overlapping ORF that enhances viral pathogenicity

[†]Craig H Kerr[1,2], [†]Qing S Wang[1], Kyung-Mee Moon[1,2], Kathleen Keatings[3], Douglas W Allan[3], Leonard J Foster[1,2], Eric Jan*[1]

[1]Department of Biochemistry and Molecular Biology, [2]Centre for High-Throughput Biology, [3]Department of Cellular and Physiological Sciences, University of British Columbia, Vancouver BC, V6T 1Z3 Canada

[†]contributed equally

*corresponding author: ej@mail.ubc.ca

keywords: ribosome, frameshift, dicistrovirus, drosophila, virus

ABSTRACT

RNA structures can interact with the ribosome to alter translational reading frame

maintenance and promote recoding that result in alternative protein products. Here, we

show that the internal ribosome entry site (IRES) from the dicistrovirus Cricket paralysis

virus drives translation of the 0-frame viral polyprotein and an overlapping +1 open

reading frame, called ORFx, via a novel mechanism whereby a subset of ribosomes

recruited to the IRES bypasses downstream to resume translation at the +1-frame 13th

non-AUG codon. A mutant of CrPV containing a stop codon in the +1 frame ORFx

sequence, yet synonymous in the 0-frame, is attenuated compared to wild-type virus in

a *Drosophila* infection model, indicating the importance of +1 ORFx expression in

promoting viral pathogenesis. This work demonstrates a novel programmed IRES-

mediated recoding strategy to increase viral coding capacity and impact virus infection,

highlighting the diversity of RNA-driven translation initiation mechanisms in eukaryotes.

INTRODUCTION

The ribosome mediates translation involving decoding the open reading frame codon by codon through delivery of the correct aminoacyl-tRNAs to the ribosomal A site. This fundamental process occurs with high fidelity for proper gene expression in all species. However, mechanisms exist that can alter the translational reading frame, thus producing alternative protein products from a single RNA (Baranov, Atkins, & Yordanova, 2015). In general, these mechanisms termed recoding, involve a specific RNA structure or element that interacts with the ribosome to cause the translating ribosome to shift reading frame by -1/+1/+2 allowing it to read through stop codons or bypass sequences and restart translation downstream (Dinman, 2012; Firth & Brierley, 2012; Jan, Mohr, & Walsh, 2016). Study of these mechanisms has been enlightening; revealing key ribosome:RNA interactions that alter fundamental processes in the mechanics of ribosome decoding and reading frame maintenance. Importantly, recoding mechanisms are now appreciated as important regulatory processes that can impact the fate of protein expression in cells and virus infections (Au & Jan, 2014; Baranov et al., 2015). Unlike these recoding mechanisms that involve an actively translating ribosome, the intergenic internal ribosome entry site (IRES) within a subset of dicistroviruses has the unusual property to directly recruit the ribosome and initiate translation from overlapping 0 and +1-frame codons to produce two distinct proteins (Ren et al., 2012). Here, we report a novel recoding mechanism and translational initiation pathway whereby a related dicistrovirus IRES directs the ribosome to initiate translation downstream.

Most eukaryotic mRNAs utilize a cap-dependent scanning mechanism involving >12 translation initiation factors to recruit the ribosome and initiate translation from an AUG start codon (Jackson, Hellen, & Pestova, 2010). Alternatively, an IRES, in general, is a structured RNA element that facilitates 5' end-independent translation using subset of translation initiation factors thus providing an advantage during viral infection or under times of stress when cap-dependent translation is compromised (Au & Jan, 2014; Firth & Brierley, 2012; Jan et al., 2016).

Of the different classes of viral IRESs based on factor requirement and mechanism, the intergenic IRES of the *dicistroviridae* family stands out as the most streamlined using a unique mechanism where it directly binds 40S and 80S ribosomes without the need for canonical initiation factors or initiator Met-tRNA$_i$ and initiates translation from a non-AUG codon (Jan, Kinzy, & Sarnow, 2003; Jan & Sarnow, 2002; Pestova & Hellen, 2003; Petrov, Grosely, Chen, O'Leary, & Puglisi, 2016; Wilson, Pestova, Hellen, & Sarnow, 2000). The dicistrovirus IRES is composed of three pseudoknots (PKI, II, and III) that separate into distinct domains; PKII and PKIII fold independently to create the ribosome-binding domain while PKI mediates positioning of the ribosome and establishes the translational reading frame (Jan & Sarnow, 2002; Nishiyama et al., 2003; Wilson et al., 2000). Structural studies have indicated that PKII and PKIII form a compact core structure, and the PKI region adopts a conformation that mimics an anti-codon:codon interaction that binds that conserved core of the ribosome in the A site (Costantino, Pfingsten, Rambo, & Kieft, 2008; Fernandez, Bai, Murshudov, Scheres, & Ramakrishnan, 2014; Pfingsten, Costantino, & Kieft, 2006; Schuler et al., 2006). From here, in an elongation factor 2 dependent manner, the IGR IRES pseudo-

translocates to the ribosomal P site, followed by aminoacyl-tRNA delivery to the A site

and a second round of eEF2-dependent pseudo-translocation of the IGR IRES to the E

site of the ribosome (Au et al., 2015; Fernandez et al., 2014; Muhs et al., 2015).

Altogether, the IGR IRES acts as a complete RNA machine that supersedes initiation

factors and commandeers the ribosome, a strategy that is essential for viral protein

synthesis in dicistrovirus-infected cells (Kerr, Ma, Jang, Thompson, & Jan, 2016).

In general, the dicistrovirus IGR IRESs are conserved at the structural, but not

sequence level and are classified into two sub-groups (termed Type I and II) based on

the presence of distinct structural elements; the main distinction comes from a larger

L1.1 loop and an additional stem-loop (SLIII) in Type II IRESs (Jan, 2006; Nakashima &

Uchiumi, 2009). SLIII allows for the PKI domain of Type II IRESs to mimic the global

shape of a tRNA in addition to assisting in reading frame selection and the larger L1.1

region functions to mediate 60S recruitment (Au et al., 2015; Jang, Lo, & Jan, 2009;

Pfingsten et al., 2006). The domains of Type I and II IGR IRESs function similarly to

directly recruit 80S ribosomes and initiate translation (Cevallos & Sarnow, 2005;

Hatakeyama, Shibuya, Nishiyama, & Nakashima, 2004; Hertz & Thompson, 2011; Jang

& Jan, 2010). Recent high resolution cryo-EM structures of the IGR IRES bound to the

80S ribosome have demonstrated that the IRES initially binds in the A site (Abeyrathne,

Koh, Grant, Grigorieff, & Korostelev, 2016; Fernandez et al., 2014; Muhs et al., 2015).

Movement of the IRES involves an eEF2-dependent pseudo-translocation event where

the ribosome rotates up to 10° allowing PKI to move into the P site in an inchworm-like

manner (Abeyrathne et al., 2016). This allows for the non-AUG initiation codon of the

IRES to be presented in the A site for the incoming amino-acyl tRNA. The first pseudo-

translocation event and delivery of the first amino-acyl tRNA are the rate-limiting steps

of initiation on the IGR IRES (Zhang, Ng, Chen, & Cooperman, 2016).

Biochemical, phylogenetic, and bioinformatics analyses have demonstrated that

a subset of Type II IGR IRESs are able to direct synthesis of a hidden +1 open reading

frame (ORF), termed ORFx, within ORF2 of the viral genome (Firth, Wang, Jan, &

Atkins, 2009; Ren et al., 2012). The functional role of ORFx during viral infection

remains elusive. Extensive mutagenesis of the PKI domain of the *Israeli acute paralysis*

virus (IAPV) IGR IRES has revealed that 0 and +1 frame translation can be uncoupled,

suggesting that the IGR IRES may adopt specific conformations that govern the

translational reading frame (Au et al., 2015; Ren, Au, Wang, Lee, & Jan, 2014).

Generally, Type I and II IGR IRESs are thought to operate similarly in mechanism.

Specific domains between the two types are functionally interchangeable (Jang & Jan,

2010). In the present study, we investigate the capacity of other IGR IRESs from

Dicistroviruses to facilitate +1-frame translation. We show that the IGR IRES from

Cricket paralysis virus (CrPV) can synthesize a 43 amino acid long ORFx protein using

an unexpected mechanism that involves IRES-mediated ribosome bypassing.

Furthermore, we provide insight into the role of ORFx during CrPV infection and show

that mutants deficient in ORFx have impaired virulence in adult flies, thus uncovering a

novel viral recoding strategy that is essential for viral pathogenesis.

RESULTS

IGR IRES-dependent +1-frame translation is not conserved throughout

Dicistroviridae

We previously showed that a subset of dicistrovirus IGR IRESs can direct translation in the 0- and +1-frames and that a base pair adjacent to the PKI domain is important for initiation in the +1-frame (Figure 1) (Ren et al., 2014; Ren et al., 2012). The IGR IRESs are classified into two types: Type I, and II, with the main difference being an extra SLIII within the PKI domain of Type II IRES. Since the honeybee and fire ant viruses harbour Type II IGR IRESs that can support +1 frame ORFx translation, we investigated whether Type I IRESs also had the capacity for +1-frame translation. We first surveyed the other dicistrovirus IGR IRESs and their downstream sequences for a potential base pair adjacent to the PKI domain and an overlapping +1 open reading frame both *in silico* (Figure 1A, B). The IAPV and ABPV IGR IRESs contain an adjacent U-G base pair whereas the SINV-1 and KBV IRESs have a C-G base pair (Figure 1A). All but one of the IGR IRESs contains a potential base pair adjacent to the PKI domain, with the majority of them possessing a potential U-G base pair (Figure 1A). As shown previously, the honeybee and fire ant dicistrovirus ORFx proteins are approximately 94-125 amino acids in length (Firth et al., 2009). The predicted +1 ORFx lengths of other dicistroviruses range from 1-53 amino acids in length (Figure 1A). Apart from the honeybee and fire ant dicistrovirus ORFx, the longest putative ORFx sequences are found within the genomes of the CrPV and DCV at 53 amino acids in length.

To determine whether the other IGR IRESs can direct +1-frame translation, we cloned each IGR IRES within the intergenic region of a dual luciferase bicistronic

construct (Figure 1B). The Renilla luciferase (36 kDa; RLuc) monitors scanning-

dependent translation whereas firefly luciferase (FLuc) is translated by the IRES. To

measure reading frame selection, the firefly luciferase gene is fused in-frame either in

the 0 or +1-frame. Translation was measured by incubating the bicistronic constructs in

Sf-21 extracts in the presence of [^{35}S]-methionine/cysteine and the resulting proteins

were detected by SDS-PAGE and quantified via phosophoimager analysis. Additionally,

we generated bicistronic reporter constructs that contain a T2A 'stop-go' peptide, which

allows sensitive and accurate measurement of luciferase activity (Wang et al 2013

Methods) (Figure 1- figure supplement 1). Since we previously showed robust IAPV

IRES 0 and +1-frame translation (Ren et al., 2012), we chose it as a benchmark to

compare the +1-frame activity of the other IGR IRESs. In general, all IGR IRESs can

direct 0-frame translation to varying extents in this *in vitro* system. Normalizing to the

IAPV 0-frame translational activity, the CrPV IGR IRES showed the highest 0-frame

translational activity (~170%) whereas the MCDV IGR IRES had the lowest activity

(~4%; Figure 1B). By contrast, only a few IGR IRESs can support +1-frame translation.

The Type II IGR IRESs from Taura Syndrome Virus (TSV) and Mud Crab Dicistrovirus

(MCDV) cannot support +1 frame translation indicating that only a subset of Type II IGR

IRESs can direct 0 and +1 frame translation. Interestingly, besides the IAPV IGR IRES,

the BQCV and CrPV IRESs mediated +1-frame translation above background levels, 80%

and 5% of 0-frame translation, respectively. In summary, only a subset of IGR IRESs

can facilitate both 0 and +1-frame translation.

CrPV +1-frame translation is IGR IRES-dependent and initiates downstream

To explore +1 frame translation mechanisms further, we focused on the CrPV

IRES (Figure 1C). We first determined whether the structural integrity of the CrPV PKI

domain is important for +1-frame translation. For these assays, we used a bicistronic

reporter construct that contains the CrPV IRES where the firefly luciferase gene was

subcloned into the +1-frame downstream of the IRES with CrPV nucleotides 6217-6387

(Accession KP974707.1), which includes the predicted +1-frame CrPV ORFx (65 kDa;

ORFx-FLuc). A shortened IRES-mediated 0-frame protein (~11 kDa; sORF2) is also

translated, thus scanning-dependent and CrPV IRES-mediated 0/+1-frame translation

can be simultaneously monitored from this dual luciferase reporter construct (Figure 2A).

Synthesis of all three proteins is detected by incubating the bicistronic construct in a

Sf21 translation extract in the presence of $[^{35}S]$-methionine/cysteine (Figure 2B, lane 1)

(Ren et al., 2012). As shown in Figure 1A, CrPV IRES +1-frame ORFx translation is

approximately 5% of 0-frame translation. Mutating CC_{6214-5} to GG, which disrupts PKI

base pairing and abolishes CrPV IRES activity, resulted in negligible or diminished 0-

and +1 frame translation whereas a compensatory mutation that restores PKI base

pairing rescued translation (Jan & Sarnow, 2002) (Figures 2B, lanes 2 and 3, Figure 1-

figure supplement 1), indicating that the integrity of the IRES and the PKI domain is

required for CrPV IRES +1-frame translation.

The adjacent U-G base pairing of the IAPV IRES is important for +1-frame

translation (Ren et al., 2012). CrPV also has the capacity to form a wobble base pair

adjacent to the PKI domain through nucleotides U_{6186} and G_{6217}, potentially directing

translation from the first +1-frame CUA leucine codon (Figure 1A). To determine if this

base pair is necessary to drive CrPV IRES +1-frame translation, we mutated U_{6186} and

G_{6217} to other bases. Mutating U_{6186} to C or G led to an approximate 18%-23% reduction

in +1 frame activity and mutating G_{6217} to C, A, or U resulted in roughly a 39%, 24%,

and 36% reduction in +1 frame activity, respectively. Although each mutation reduced

+1 frame translational activity to some degree, none of the mutations abolished it

(Figures 2B, lanes 4-9). These results suggest that unlike with IAPV, base pairing

between nucleotides 6186 and 6217 is not absolutely required for CrPV IRES +1-frame

translation.

To determine the potential initiation site of CrPV ORFx, we systematically

replaced codons downstream of the IRES with a stop codon and monitored 0 and +1-

frame translation *in vitro* using the bicistronic reporter construct (Figure 2B, C). Overall,

stop codons placed in the +1-frame did not significantly affect 0-frame translation,

indicating that IRES activity was not compromised (Figure 2B, C). Replacing individual

codons between the 1st and the 12th +1-frame codon with a stop codon inhibited to

varying extents (between 36%-71% reduction compared to wild-type) but did not

completely abolish +1 frame translation (Figure 2B, lanes 11-16; 2C). Conversely, +1-

frame translation was completely inhibited when the 13th +1-frame codon and codons

thereafter was replaced with a stop codon (Figure 2B, lanes 15-16; 2C). Replacing both

the 3rd and 5th +1-frame codons with stop codons reduced +1-frame translation by 33%

but did not eliminate it, ruling out ribosome read-through. To address the possibility that

IRES translation initiates in the 0-frame and then a fraction of translating ribosomes shift

into the +1-frame, we inserted a stop codon in the 0-frame downstream of the IRES. As

expected, a stop codon in the 0-frame 1st to 4th codons downstream of the IRES

abolished 0-frame translation (Figure 2B, lanes 18-21). However, the 0-frame stop

codon insertions reduced by 52-77% but did not eliminate +1 frame translation

indicating that a subset of ribosomes do not shift from the 0 to +1 reading frame. Further

supporting this, introducing stop codons in both the 0 and +1 frames did not abolish +1

frame translation, though inhibited +1 frame translation by ~80% (Fig. 2C, 0 S4/+1 S5).

We noted that the adjacent 14th codon is an AUG methionine. Mutating the AUG to a

CGG (M14R) decreased by 63% but did not abolish +1-frame translation (Fig. 2C). In

summary, the mutational analysis indicated that CrPV ORFx translation requires an

intact IRES and the +1-frame AAA lysine 13th codon. Mutating the 13th AAA codon to

UUU slightly reduced but did not abolish +1-frame translation (Fig. 2D), suggesting that

there may be flexibility in the codon identity for CrPV ORFx +1-frame translation.

CrPV +1-frame translation occurs independently of 40S scanning

The current data led us to generate two hypotheses: i) a subset of 40S subunits

recruited to the CrPV IRES scan downstream to start translation at the 13th codon

(scanning hypothesis) or ii) 40S or 80S ribosomes recruited to the IGR IRES bypass the

spacer region to the 13th codon (bypass hypothesis). To address the scanning

hypothesis, we took a three-pronged approach. First, we utilized mutants of the CrPV

IRES in the L1.1 loop that is known to be deficient in recruitment of the 60S subunit

(Jang et al., 2009; Pfingsten et al., 2006). If scanning is occurring, 40S subunits

recruited to the L1.1 IRES may scan downstream to the downstream +1-frame initiation

codon. Reporter constructs harbouring mutations in the L1.1 loop were deficient in 0

and +1-frame translation (Figure 3A), suggesting that 60S recruitment by the IGR IRES

is necessary for translation in the +1-frame.

Secondly, we utilized the translational inhibitor edeine to assess if scanning was

occurring (Figure 3B). Edeine prevents the 40S ribosomal subunit from recognizing an

AUG start codon (Dinos et al., 2004; Kozak & Shatkin, 1978). Both 0- and +1-frame

translation were resistant to edeine relative to that of scanning-dependent translation

(Figure 3B). This result further supports that the +1-frame 14th AUG codon is not

utilized for +1-frame translation.

Finally, we introduced a stable hairpin (HP1 ΔG = -17.0 or HP2 ΔG = -37.0

kcal/mole) into the spacer region between the CrPV IRES and the downstream 13th

codon to prevent 40S scanning (Figure 3C). Introducing HP1 or HP2 into the spacer

region did not affect 0-frame translation showing that these mutations did not affect

IRES activity (Figure 3C). In contrast, +1-frame translation was decreased by ~50%.

However, in these constructs, the hairpins are likely unwound due to 80S ribosomes

translating in the 0-frame, thus confounding the interpretation of the +1-frame translation

result (Figure 3C). To uncouple these effects, we mutated the 0-frame 3rd codon to a

stop codon upstream of the hairpin (0S3), thus blocking translating ribosomes from

reaching the hairpin and maintaining hairpin integrity. Surprisingly, placing a stop codon

upstream of the hairpin abolished +1-frame translation (Figure 3C), suggesting that 80S

ribosomes may scan downstream to the 13th codon. Given the reduction in +1-frame

translation observed when stop codons are placed adjacent to PKI (Figure 2A), it is

possible that these hairpin insertions, in addition to the nucleotide identity, are disrupting

a functional RNA element within the spacer region that mediates translation at the

downstream +1 frame codon.

+2 frame translation mediated by the CrPV IRES

Our data thus far suggests that scanning of 40S ribosomes to the 13th position is

likely not occurring. Therefore, we next asked whether ribosomes bound to the IGR

IRES are repositioned to the downstream codon akin to translational bypass (Huang et

al., 1988; Samatova, Konevega, Wills, Atkins, & Rodnina, 2014). We hypothesized that

if the ribosome is indeed bypassing to a specific 'landing site' then it should be

independent of frame, whereas if the ribosome merely begins translation in the 0-frame

before slipping into the +1-frame, then we expect to see no ORFx translation. To

address this, we inserted a series of nucleotides into our bicistronic construct that shifts

only the ORFx-Fluc into the +2 frame (Figure 4). Specifically, we inserted either a single

nucleotide or up to 7 proceeding the 13th AAA codon. As a control, we inserted either 6

or 9 nucleotides in the same position that does not introduce an additional frameshift.

Insertion of a U creates a stop codon in the +1-frame while an inserted C does not. To

our surprise, we observed ORFx expression with all insertions (Figure 4). These results

suggest that repositioning of the ribosome from the IRES to a downstream 13th +1 or

+2 frame codon is reading frame-independent.

The integrity of the VLR and pseudo-translocation of the IGR IRES through the

ribosome is required for ORFx expression

How do 80S ribosomes recruited to the IRES mediate bypass at the downstream

13th +1-frame codon? Since the IGR IRES, which occupies the ribosomal A site upon

ribosome binding, undergoes pseudotranslocation to the ribosomal P site to vacate the

A site for delivery of the first aminoacyl-tRNA (Fernandez et al., 2014), we reasoned

that the ribosome bound to the CrPV IGR IRES must have an empty ribosomal P or A

site in order to reposition downstream and accommodate the +1 frame start codon. To

address this, we introduced mutations in the VLR, which has been shown to disrupt the

IGR IRES-mediated pseudotranslocation event (Figure 1C) (Ruehle et al., 2015).

Specifically, shortening the length of the VLR by 2 or 3 nucleotides (Δ2 and Δ3,

respectively) inhibits the first pseudo-translocation event from the A site to the P site

whereas altering the identity of nucleotides A_{6204}, and AA_{6208-9} to guanosines (G-rich)

inhibited IRES translocation from the P site to the E site (Ruehle et al., 2015).

Interestingly, all three VLR mutants decreased +1-frame activity (Figure 5); both the G-

rich and Δ3 mutants demonstrated little to no activity, while the Δ2 mutant still exhibited

roughly 50% activity to that of WT. This result is consistent with previous data that the

Δ2 mutant IRESs are still able to accommodate a fraction of aminoacyl-tRNA in the A

site (~25%), allowing translation to occur (Ruehle et al., 2015). Altogether, these results

indicate that the pseudotranslocation event of the IGR IRES through the ribosome

contributes to +1-frame translation downstream.

Given that pseudotranslocation is required for +1 frame translation, we propose

two possible models: After ribosomes that are recruited to the CrPV IRES-mediated

pseudotranslation, i) a fraction of ribosomes start translation downstream at the 13th +1

frame codon or ii) a fraction of ribosomes initiates translation in the 0- or +1-frame (Ala

or Leu, respectively) of CrPV ORF2 before ribosome bypass and resuming translation

downstream at the 13th +1 frame codon. To distinguish these possibilities, we

attempted to identify the N-terminal amino acids of ORFx. To this end, we fused a 6X

His-tag on the C-terminus of ORFx-FLuc in the bicistronic construct lacking the T2A

sequence (Figure 1 – figure supplement 1). Using this construct, we used a large scale

(100X) *in vitro* Sf-21 translation reaction in an effort to isolate the ORFx-FLuc-6XHis to

appreciable levels (Figure 5 – figure supplement 1A). In a parallel reaction, we

incubated radioactive [^{35}S] Met/Cys to reactions in order to follow purification of ORFx.

A clear band corresponding to radioactive ORFx-Fluc-6XHis was observed via

autoradiography and after purification with Ni^{2+}-NTA beads (Figure 5 – figure

supplement 1A, 1B). The purified ORFx-Fluc-6XHis was then subjected to Edman

degradation analysis (Figure 5 – figure supplement 1C, 1D). Subsequently, a likelihood

score was calculated for each residue within a cycle to determine the appropriate amino

acid assignment (see Materials and Methods). From this, we reproducibly observed Lys

and Met residues in the 3rd and 4th positions, respectively (Figure 5 – figure

supplement 1E). Unfortunately, the amino acids in positions 1 and 2 could not be

unambiguously assigned by Edman degradation due to the low yield of +1 frame ORFx

translation. Indeed, subjecting low picomole amounts of protein to N-terminal

sequencing can result in convoluted traces in early cycles often due to contaminants

from sample loading or the sample itself (Speicher, Gorman, & Speicher, 2009). Despite

this, the amino acids with the best likelihood score for position 1 are Ala and Thr (note

that Ser is a common contaminant and Trp is near background levels) while position 2 is

observed as a Leu or Ile residue. Taken together with our biochemical analysis, this

data suggests that a fraction of ribosomes recruited to the CrPV IRES undergoes

pseudotranslocation in either the 0 or +1 frame and then bypasses downstream to

resume translation at the 13th Lys codon.

The spacer region downstream of the IGR IRES is necessary for +1 frame

translation

We next investigated whether the spacer region located between PKI of the IGR

IRES and the downstream AAA codon contributes to IRES-mediated ribosome bypass.

Our results showed that inserting stop codons in the 'spacer region' (Figure 2A)

between the IRES PKI domain and the +1-frame 13th codon of ORFx inhibited but did

not completely abolish +1-frame translation, suggesting that an element within this

spacer region may promote +1-frame translation. We first addressed whether the CrPV

spacer region is sufficient to direct +1-frame translation by generating a chimeric

construct whereby the PSIV IRES is fused with the CrPV spacer region (Figure 6). The

PSIV IRES directs strong 0-frame but no or relatively weak +1-frame translation (Figure

1). The PSIV-CrPV spacer chimeric reporter resulted in 0-frame translation (Figure 6A),

indicating that the spacer region does not affect the activity of the PSIV IRES. In

contrast to the PSIV IRES construct, the chimeric PSIV-CrPV reporter resulted in +1-

frame translation, implying that the CrPV spacer region is sufficient to drive +1-frame

translation in the presence of a functional IGR IRES (Figure 6A).

To examine more closely the molecular determinants of the CrPV spacer region

that direct +1-frame translation, we deleted or inserted a randomized sequence within

the spacer region to test whether the length was important (Figure 6 and Figure 6 -

figure supplement 1A). Insertions within or deletion of the entire spacer region resulted

in loss of +1-frame but not 0-frame translation, suggesting that sequence context is

important for +1-frame translation (Figure 6A - figure supplement 1B). Additionally, we

examined the possibility of the spacer region to form intramolecular interactions. *In silico*

analysis using RNA folding algorithms did not reveal any obvious stable secondary

structures in the spacer region (data not shown), however we noticed that nucleotides

AUUU$_{6222-25}$ and AAAU$_{6240-43}$ have the potential to interact. Mutating UUU$_{6223-25}$ to AAA

(A/U) or AAAA$_{6240-43}$ to UUUU (U/A) drastically reduced +1-frame translation (Figure 6 -

figure supplement 1C). However, compensatory mutations did not restore +1-frame

translation. To delineate a functional element within the spacer region, we

systematically tested deletions from the 3' end of the spacer region (Figure 6B).

Interestingly, deleting 3 to 27 nucleotides did not affect +1 frame translation. However,

deleting 30 nucleotides and leaving seven nucleotides adjacent to the IRES abolished

+1 translation (Figure 6B). Interestingly, Δ3, Δ6 and Δ27 mutants appear to have much

higher Fluc/Rluc activity when compared to WT, an observation that requires further

examination. In summary, these results suggest that the specific sequences within the

spacer region, in particular the sequences immediately downstream of the PKI domain,

are important for mediating IRES-dependent +1 frame translation.

CrPV +1-frame ORFx is expressed yet not required for infection in *Drosophila* S2

cells

Our *in silico* and biochemical data indicate that ribosomes recruited to the CrPV

IRES can bypass downstream to resume +1-frame translation from the 13th codon. We

termed the predicted +1 frame open frame CrPV ORFx, which is predicted to be 43

amino acids in length (Figure 7). To determine whether CrPV ORFx is synthesized

during infection, we used a proteomics approach to identify ORFx peptides in CrPV-

infected S2 cells. Specifically, *Drosophila* S2 cells infected with CrPV (MOI 10) were

harvested at 6 hours post infection and lysed. Proteins were subsequently digested with

trypsin and peptides were analyzed by LC-MS/MS. We identified two peptides that

correspond to CrPV ORFx both of which were located downstream of the +1-frame 13[th]

codon (Figure 7). Importantly, these peptides were not identified in mock-infected S2

cells. Thus, ORFx is expressed in CrPV-infected S2 cells.

We next sought to determine if CrPV ORFx expression influences the outcome of

viral infection. Using a recently developed a CrPV infectious clone, termed CrPV-2, we

introduced mutations that would abolish ORFx expression (Kerr et al., 2015). To this

end, we created two separate mutant clones: the +1-frame 12[th] codon (UUG_{6251-3}) was

altered to an amber stop codon (UAG; CrPV-S12) and the +1-frame 19[th] codon

(UUA_{6272-4}) changed to an ochre stop codon (UAA; CrPV-S19) (Figure 8A). Both

mutations are synonymous in the 0 frame. Based on our translation data (Figure 2), the

+1-frame S19 would inhibit ORFx expression whereas the +1-frame S12 would inhibit

but not abolish ORFx expression.

We first examined whether ORFx influences viral protein synthesis *in vitro.*

Incubation of *in vitro* transcribed CrPV-2 RNA in Sf21 translation extracts led to

synthesis and processing of all viral proteins as reported previously (Figure 8B) (Kerr et

al., 2015). Introduction of stop codons in the 0 frame of ORF1 or ORF2 inhibited

synthesis of either viral non-structural or structural proteins, respectively (Figure 8B)

(Kerr et al., 2015). Both CrPV-S12 and CrPV-S19 RNAs resulted in viral protein

synthesis that was indistinguishable compared to wild-type CrPV -2 *in vitro* (Figure 8B).

Note that CrPV ORFx is predicted to be ~4.9 kDa and is undetectable in this analysis.

These results demonstrate that CrPV ORFx is not necessary for viral protein synthesis

in vitro.

We next assessed the viability of the CrPV-S12 and -S19 viruses in cell culture.

To determine if ORFx plays a role in the first replication cycle of CrPV, we transfected

S2 cells with *in vitro* transcribed RNA from wild-type or mutant S12 or S19 CrPV-2 to

generate virus. After 48 hours, virus was harvested and used to infect naïve S2 cells at

an MOI of 10 to follow the first round of infection. We monitored viral protein synthesis

via metabolic [^{35}S]-Met/Cys labeling as well as viral protein and RNA accumulation by

Western and Northern blotting, respectively. Infection with wild-type CrPV-2, CrPV-S12

or CrPV-S19 all resulted in accumulation of viral proteins and RNA and shutdown of

host translation in a similar manner (Figure 8C). Furthermore, there was no visible

difference in viral RNA and protein levels between wild-type and mutant virus infections.

These results indicated that ORFx does not play a role in the first round of CrPV

replication in S2 cells.

To determine if ORFx influences subsequent rounds of infection, we infected S2

cells at an MOI of 1 and monitored viral titres (Figure 8D). Neither infection at an MOI of

1 produced significantly different titres between wild type CrPV-2 and either stop codon

mutant at any time point, apart from both CrPV-S12 and CrPV-S19 resulting in higher

titres than CrPV-2 after 24 hours post infection at MOI 1. A similar result was observed

in CrPV-infected S2 cells at an MOI 0.1 (data not shown). Taken together, ORFx has no

observable effect on the life cycle of CrPV in S2 cells.

ORFx contributes to CrPV pathogenesis in adult flies and associates with

membranes

We next addressed whether ORFx contributes to CrPV pathogenesis in adult fruit

flies. To test this, we injected adult flies intrathoracically with PBS, CrPV-2, CrPV-S12,

or CrPV-S19 and monitored fly mortality daily. By day 5, flies injected with CrPV-2 or

CrPV-S12 exhibited 50% mortality and were 100% dead by day 11 and 12, respectively

(Figure 9A). By contrast, flies injected with CrPV-S19 did not reach 50% mortality until

day 9 and reached 100% mortality at day 14 (Figure 9A). These results demonstrate

that ORFx contributes to CrPV pathogenesis in adult flies.

To determine if the effect seen on CrPV pathogenesis is a result of defects in

viral replication, we measured viral titres and assessed viral protein levels in adult flies.

At 5 days post infection, flies were homogenized and lysates were analyzed via

Western blot for viral proteins and viral titres were determined. Interestingly, viral titres

showed no significant differences between wild type CrPV-2, CrPV-S12, and CrPV-S19

(Figure 9B). As well, protein levels in infected flies were unaffected by the S19 mutation

compared to wild-type or the S12 mutant (Figure 9C). Finally, to ensure that the S12

and S19 mutations in the mutant CrPV clones are stable, we performed RT-PCR

analysis and sequenced this region. Our results showed that the S12 and S19

mutations are stable during virus propagation in S2 cells (data not shown). In summary,

our results indicate that the defect in viral pathogenesis observed in the S19 mutant

virus is not due to a defect in viral replication.

From our data, CrPV ORFx likely expresses a 43 amino acid protein whereby

ribosomes bypass to resume translation from the 13th +1-frame AAA codon. Comparing

the sequence of ORFx from CrPV and other species shows no appreciable homology to

other proteins. Nevertheless, *in silico* topology predictions suggest that ORFx has the

propensity to form an alpha helical transmembrane segment at its C-terminus (amino

acids 22-39; Figure 10 – figure supplement 1). To examine ORFx function, we

generated Drosophila expression constructs containing either N- or C-terminal HA-

tagged ORFx . We first assessed whether expression of HA-ORFx affected cell viability

by a trypan blue exclusion assay. Transfection for 10 or 24 hours with the ORFx-HA and

HA-ORFx construct did not result in a dramatic decrease in cell viability (Figure 10 –

figure supplement 1). However, by 48 hours, transfection of ORFx-HA and HA-ORFx

led to a statistically significant but relatively minor reduction (13% and 15% decrease,

respectively) in cell viability, suggesting that ORFx expression has a slight toxic effect in

S2 cells. Immunoblotting for HA showed that HA-ORFx is expressed in S2 cells (Figure

10 - figure supplement 2). To determine the subcellular localization in S2 cells, we

transfected the HA-tagged ORFx constructs and monitored ORFx localization (pORFx-

HA) by HA-antibody immunofluorescence staining in comparison with cytoplasmic, ER,

Golgi, and nuclear marker protein antibodies. We also mutated two pairs of amino acids,

LV and LI, to KK and KR respectively, which would disrupt the transmembrane domain

(pORFx-TMmut-HA). Co-staining showed that the wild-type HA-tagged ORFx overlaps

mainly with ER protein marker Calnexin, and partially overlaps with Golgi- and nuclear-

associated proteins, Golgin84 and nuclear Lamin, respectively (Figure 10, pORFx-HA).

The wild-type HA-ORFx displayed little to no overlap with α-Tubulin staining. By

contrast, the mutant transmembrane HA-tagged ORFx staining showed no overlap with

Calnexin, Golgin84, nor Lamin (Figure 10, pORFx-TmMut-HA). Collectively, these

results suggest that ORFx associates with membranous organelles and potentially the

ER specifically (Figure 10). To examine this further using another approach, we

transfected S2 cells with the HA-tagged ORFx expression plasmid and used differential

centrifugation on cells to separate subcellular components. As controls, we monitored

protein markers that should be enriched in different subcellular fractions. As expected,

tubulin is found in the cytoplasmic fraction and the ER-associated KDEL protein and

cytochrome C are enriched in nuclei and ER fractions (Figure 10- figure supplement 2).

Both HA-tagged ORFx are detected within membranous fractions and not within the

cytoplasmic fraction (Figure 10- figure supplement 2), further supporting that ORFx

resides within the membrane of cells that may be important for its function.

DISCUSSION

Recoding mechanisms have illuminated diverse RNA structural elements that interact with the ribosome to affect reading frame maintenance (Dinman, 2012). In this study, we have demonstrated a novel translation recoding mechanism by which an IRES can promote ribosome repositioning at a downstream codon. In contrast to the IAPV IGR IRES, which directs ribosome reading frame selection (Ren et al., 2012), the CrPV IGR IRES can facilitate the expression of a downstream +1 overlapping frame, which we termed ORFx. We provide extensive analysis using mutagenesis that translation of CrPV ORFx occurs 37 nucleotides downstream at the 13[th] AAA (Lys) codon. Moreover, we show that ORFx is expressed in CrPV-infected cells by mass spectrometry and that ORFx is required for promoting CrPV pathogenesis in a *Drosophila* injection model. Our data suggest a model whereby after 40S and 60S ribosomal subunits binding to the CrPV IGR IRES, the majority of the ribosomes translate in the 0-frame by delivery of the incoming Ala-tRNA^{Ala} to the 0-frame GCU codon whereas a fraction of ribosomes undergoes pseudotranslocation in either the 0 or +1 frames and then bypasses or "slides" 37 nucleotides downstream to resume translation at the +1-frame AAA (Lys) codon (Figure 11). From our results, the following rules appear to apply to +1-frame translation in CrPV: i) the PKI must be intact, ii) both 40S and 60S ribosomal subunits are required to bind to the IRES, iii) pseudotranslocation of the IRES through the ribosome is necessary, iv) the spacer region between PKI and the 13[th] AAA codon is essential, and v) the nucleotide identity of the spacer region is crucial for efficient +1 frame translation.

A unique feature of this repositioning mechanism is that an intact IRES is essential for +1-frame translation (Figure 2). The current model is that the PKI domain

of the IRES occupies the ribosomal A site upon ribosome binding to the IRES, followed

by a pseudotranslocation event in order to vacate the A site to allow delivery of the first

aminoacyl-tRNA in the 0-frame (Fernandez et al., 2014). Given this model, it is difficult

to envision how IRES can direct the ribosome to bypass 37 nucleotides to translate at a

+1-frame AAA Lys codon. A potential clue comes from our mutagenesis analysis that

suggests that translocation of the CrPV IRES through the ribosome is a prerequisite for

downstream +1-frame translation (Figure 5). This is intuitive as the PKI domain must

vacate the A site in order to allow translation in both the 0 and +1 frames. Building on

this, N-terminal sequencing of the *in vitro* translated CrPV ORFx clearly showed that the

3rd and 4th N-terminal amino acids are Lys and Met, respectively, which are likely

encoded by the 13th and 14th +1 frame codons from the CrPV PKI domain (Figure 5 –

Figure supplement 1). Although the first two N-terminal residues of ORFx were not

confidently assigned, it is clear that two translocation cycles occur prior to translation of

the 13th and 14th codons. Thus, the ribosome bound to the IRES must undergo two

pseudotranslocation events before ribosome bypass and 'landing' at the 13th +1 frame

codon to resume translation (Figure 11). Stop codon insertions upstream of +1 frame

13th codon do not support +1 frame translation and insertion of a stop codon in the 12th

codon of the CrPV infectious clone does not affect viral pathogenesis (Figures 2 and 9),

thus ribosomes bound to the IRES must initiate translation then bypass to resume

translation at the 13th codon. Given this, how can +1 frame translation still occur in the

mutational analysis with 0- or +1-frame stop codons insertions directly downstream of

the IRES (Figure 2)? A recent *in vitro* study using single molecule fluorescence

spectroscopy by the Puglisi group demonstrated that the CrPV IGR IRES can facilitate

+1-frame translation approximately 5% of the time compared to 0-frame translation

whereby reading frame selection is dictated by the kinetics of tRNA binding in the first 0-

or +1-frame codon (Petrov et al., 2016). Thus, in the context of the stop codons in the 0

or +1 frames, a fraction of ribosomes recruited to the IRES can still initiate translation in

either frame and then undergoes ribosome bypass downstream to the 13th +1 frame

AAA codon (Figure 11). Determination of the aminoacyl-tRNA(s) that are delivered

during the initial CrPV IGR IRES pseudotranslocation event should provide insights

whether the ribosomal A site is vacant prior to ribosome sliding downstream to the 13th

+1 frame codon.

Another key feature is that the downstream spacer region is necessary for

efficient CrPV IRES-mediated +1-frame synthesis (Figures 3C, 6, and Figure 6 - figure

supplement 1). Interestingly, the majority of mutations within the spacer region causes a

reduction ranging from 10-80% in the amount of CrPV IRES-mediated +1-frame

synthesis (Figure 2), suggesting that nucleotide or codon identity is crucial for triggering

the bypass event. There are no obvious RNA secondary structures within the spacer

region, however, we cannot rule out any long distance RNA:RNA interactions that may

contribute to +1-frame translation. Furthermore, sequences immediately downstream of

the IRES PKI domain are important for +1 frame translation (Figure 3B). Investigation

into how the spacer sequence influences ribosomes bound to the IRES to start

translation in the 0 frame or at the downstream +1 frame initiating codon should shed

light onto this ayptical ribosome bypass mechanism.

Our results indicate that the 13th +1 frame AAA Lys codon is not a strict

requirement as appreciable CrPV IRES +1-frame translation still occurs when the AAA

is altered to other codons (Figures 2C and 6B). Although there is an AUG codon at the

14th codon, altering this codon did not eliminate +1-frame translation. In addition, +1-

frame translation is edeine-insensitive and requires an intact IRES PKI domain and 60S

joining, thus ruling out a 40S scanning mechanism per se (Figures 2 and 3). Lastly,

another interesting feature is that once ribosomes 'leave' the IRES, translation of the

downstream codon is frame-independent, similar to that observed in the ribosome

bypass event in *gene 60* of T4 bacteriophage (Huang et al., 1988; Samatova et al.,

2014).

How does the ribosome reposition to the downstream +1-frame initiation codon

after IRES binding? It is possible that the repositioning of the ribosomes occurs via

mechanism similar to that observed with prokaryotic 70S 'scanning' (Yamamoto et al.,

2016) or 70S 'sliding' that occurs in coupled translational reinitiation (Adhin & van Duin,

1990; Nomura, Gourse, & Baughman, 1984) (Figure 11). Indeed, a related dicistrovirus

PSIV IRES can direct translation using a scanning-like mechanism in prokaryotes,

which may suggest a similar property observed with our studies on the CrPV IRES

(Colussi et al., 2015). Moreover, it is has been reported and proposed that energy-

independent scanning or diffusion of the ribosome or ribosomal subunits (i.e. phaseless

wandering) can occur to locate an AUG codon (Abaeva, Pestova, & Hellen, 2016; Adhin

& van Duin, 1990; Sarabhai & Brenner, 1967; Skabkin, Skabkina, Hellen, & Pestova,

2013)(Figure 11). Nevertheless, our study shows that CrPV IRES +1-frame translation

requires 80S ribosome binding to the IRES (Figure 3A) and is edeine-insensitive (Figure

3B), thus we favour a model that 80S ribosomes reposition to the 13th +1-frame codon.

This warrants comparison is the translational bypassing observed in *gene 60* of T4

bacteriophage (Wills et al., 2008). In *gene 60*, translating ribosomes stall in a non-

canonical rotated state at a 'take-off' Gly codon with a peptidyl-tRNA^{Gly}, which

dissociates from the anticodon, and 'lands' at a matching Gly codon 50 nucleotides

downstream and allowing translation to resume. This bypass mechanism is dependent

on a post-translocation step requiring a downstream 5' 'take-off' stem-loop structure and

a nascent translated peptide (Chen et al., 2015; Samatova et al., 2014). Although CrPV

does not have an obvious RNA structure within the spacer region, it is possible that the

highly complex structure of the IGR IRES itself may contribute to downstream +1 frame

translation (Figure 11). Moreover, it is known that the ribosome bound to the IGR IRES

is in a rotated state and that the first pseudotranslocation step is rate limiting, which may

contribute to CrPV IRES reading frame selection (Fernandez et al., 2014; Zhang et al.,

2016). Finally, it is possible is that CrPV IRES bypass could be occurring through an

RNA looping event (Paek et al., 2015); the downstream RNA is brought into close

proximity with the 80S ribosome allowing it to transition to 13th AAA codon potentially by

an unknown protein factor or a long-range RNA:RNA interaction. Taken together, it is

likely that it is a combination of tRNA kinetics and conformational changes of the IRES

and the downstream spacer region that lead to bypassing, of which the contributions of

each element require further investigation.

The biological relevance of CrPV +1-frame translation was initially evidenced by

the detection of ORFx peptides in CrPV-infected S2 cells (Figure 7). To our surprise,

disruption of ORFx synthesis by stop codon insertion in the +1-frame did not perturb

viral infection in tissue culture cells but showed retarded mortality in adult flies even

though viral load remained similar between wild type and mutant viruses (Figures 8 and

9). Naturally, CrPV infection is thought to cause death through paralysis, subsequently

leading to dehydration or starvation of the host (Chtarbanova et al., 2014; Reinganum,

1975). CrPV can infect several tissues in the fly including the trachea, midgut, and

central nervous system although the latter has not been demonstrated directly (Bonning

& Miller, 2010; Chtarbanova et al., 2014; Paek et al., 2015). How ORFx contributes to

CrPV pathogenesis is an outstanding question. Our results indicate that ORFx is

membrane associated (Figure 10) and does not contribute directly to viral replication but

rather to the pathogenesis of CrPV infection in fruit flies (Figure 9). Furthermore,

expression of ORFx is slightly cytotoxic in Drosophila cells, a property that may also

contribute to pathogenesis of CrPV infection. Future studies into the localization of

ORFx, potential interacting partners, and tissue tropism of wild-type versus mutant virus

infection in the fly should provide insights into the role of ORFx.

Viruses continue to surprise us with their ability to manipulate the ribosome in

remarkable ways. Here, in addition to the previous findings on the honey bee

dicistrovirus IRES, we have revealed another recoding mechanism utilizing an IRES,

thus highlighting the strong selection to increase the coding capacity of the

dicistroviridae genome. Furthermore, an IRES that can direct ribosome repositioning to

facilitate the translation of a hidden +1 overlapping ORFx adds to the growing list of

diverse pathways of ribosome translational recoding. The CrPV IGR IRES may serve as

a model for other IRESs such as those from Hepatitis C virus and Theiler's murine

encephalomyelitis virus that can direct translation of alternative reading frames

(Vassilaki & Mavromara, 2009; Yamasaki, Weihl, & Roos, 1999). It will be of

considerable interest to investigate whether other IRESs direct reading frame selection

by a similar CrPV IGR IRES-mediated ribosome repositioning mechanism. Ribosome

repositioning or bypass is not specific to bacteria or mitochondria (Herr, Gesteland, &

Atkins, 2000; Lang et al., 2014) but may be a more general phenomenon in eukaryotes

that initially thought.

MATERIALS AND METHODS

un

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background ()	CrPV-2	PMID: 25810541	Accession: KP974706	Infectious clone of CrPV
strain, strain background ()	CrPV-ORF1-STOP	PMID: 25810541		
strain, strain background ()	CrPV-S12	This paper		
strain, strain background ()	CrPV-S19	This paper		
genetic reagent (D. melanogaster)	w[1118]	Bloomington Drosophila Stock Center	FLYB: FBst0003605	Maintained by the D. Allan Lab at UBC
cell line ()	Schneider 2 cells	Invitrogen	Catalog #: R690-07; FLYB:FBtc0000181; RRID:CVCL_Z992	Cell line maintained in E. Jan lab; FlyBase symbol: S2-DRSC.
antibody	CrPV anti-ORF2	PMID: 19889774	NA	Dilution: 1:1000
antibody	Golgin84	Developmental Studies Hybridoma Bank	Golgin84 12-1	Dilution: 1:1000
antibody	Calnexin99a	Developmental Studies Hybridoma Bank	Cnx99A 6-2-1	Dilution: 1:1000
antibody	Tubulin	Santa Cruz	sc-8035	Dilution: 1:1000
antibody	Lamin	Developmental Studies Hybridoma Bank	ADL84.12	Dilution: 1:1000
antibody	Cytochrome C	abcam	ab13575	Dilution: 1:1000
antibody	KDEL	Enzo Life Sciences	ADI-SPA-827	Dilution: 1:1000
other	Hoeschst Stain	Molecular Probes		Dilution: 1:10000
recombinant DNA reagent	pEJ551 (plasmid)	PMID: 23009811		Bicistronic construct lacking T2A sequence. Used in Figure 1 & 2.
recombinant DNA reagent	pEJ588 (plasmid)	this paper		Bicistronic construct containing T2A sequence allowing for luciferase activity monitoring used in Figure 2 and onwards.
recombinant DNA reagent	pORFx-HA	this paper		C-terminally HA-tagged CrPV ORFx sequence expressed from pAc5.1/V5-HisB (Thermo)
recombinant DNA reagent	pHA-ORFx	this paper		N-terminally HA-tagged CrPV ORFx sequence expressed from pAc5.1/V5-HisB (Thermo)
commercial assay or kit	Sf-21 TnT Extract	Promega	Catalog #: L1102	
chemical compound, drug	Edeine	NA	NA	Limited reagent held in the Jan lab

 Cell culture and virus. *Drosophila* Schneider line 2 (S2; Invitrogen) cells were

 maintained and passaged in Shield's and Sang medium (Sigma) supplemented with 10%

 fetal bovine serum.

 Propagation of CrPV in *Drosophila* S2 cells has been previously described

 (Garrey, Lee, Au, Bushell, & Jan, 2010). CrPV-2 and mutant viruses were generated

 from *Drosophila* S2 cells using an adapted protocol (Krishna, Marshall, & Schneemann,

 2003). Briefly, 5.0×10^7 S2 cells were transfected with *in vitro* transcribed RNA derived

 from pCrPV-3 or mutant plasmids and incubated for 48h. Cells were dislodged into the

 media, treated with 0.5% Igepal CA-630 (Nonidet P-40) and 0.1% 2-mercaptoethanol,

 and incubated on ice for 10 min. Cell debris was cleared by centrifugation at 13,800

 RCF for 15 min at 4°C. Viral particles were then concentrated by ultracentrifugation at

 141,000 RCF for 2.5 h at 4°C. The pellet was resuspended in PBS and sterilized

through a 0.2 µM filter. Viral titres and yield were determined as previously described

(Kerr et al., 2016). All viruses were sequence verified via RT-PCR with primers directed

against the CrPV IGR IRES.

Plasmids and bicistronic reporter constructs. Each IGR IRES with flanking

upstream and downstream sequences was cloned between the EcoRI and the NcoI

sites within the intergenic region of pRΔDEF as described (Johannes, Carter, Eisen,

Brown, & Sarnow, 1999). The Firefly luciferase (FLuc) open reading frame was fused in

frame to either the 0 or +1 frame. The sequences that were cloned are as follows:

nucleotides 6372–6908 of Israeli acute paralysis virus (IAPV; NC_009025.1);

nucleotides 6741-6969 of Taura syndrome virus (TSV; NC_003005.1); nucleotides

6860-7243 of Mud crab dicistrovirus (MCDV; HM777507.1); nucleotides 5974–6372 of

Cricket paralysis virus (CrPV; NC_003924.1); nucleotides 6060-6422 of Drosophila C

Virus (DCV; NC_001834); nucleotides 5982-6324 of Plautia stali intestine virus (PSIV;

NC_003779.1); nucleotides 151-771 of Big Sloux River virus (BSRV; JF423197.1);

nucleotides 5626-5917 of Black queen cell virus (BQCV; NC_003784.1); nucleotides

3453-3712 of Aphid lethal paralysis virus (ALPV) strain brookings non-structural

polyprotein and capsid protein precursor (HQ871932.1); nucleotides 6546-7159 of

Rhopalosiphum padi virus (RhPV; NC_001874.1). The PSIV_CrPV chimera contains

the nucleotides 5981-6192 of PSIV and 6217-6371 of Cricket paralysis virus isolate

CrPV-2 (KP974706.1)(Kerr et al., 2015).

For CrPV IRES-containing bicistronic constructs, the AUG start codon of the

FLuc gene in the +1 frame was removed by PCR-based site-directed mutagenesis

(Stratagene). For the T2A-containing constructs, the Thosea asigna virus (accession:

AF062037) 2A sequence was inserted in frame preceding the FLuc gene into plasmid

pEJ566 between EcoRI and NdeI restriction sites as described previously (Wang, Au, &

Jan, 2013).

Hairpin structure sequences HP1 and HP2 were inserted between the 6th codon

AAC and 7th codon AAG of CrPV (+1) frame. The HP1 sequence is 5'-CAGCGTTATCG

GGAACGTCGAAGGGGTACGCCATTGTTCGACCACCGGCTCGAGGGGCCC-3' (ΔG

= -17.0 kcal/mole). The HP2 sequence is 5'- ACTGCAGCCACAACCGGCCCCAAGCTT

GGGCCGTGGTGGCTGCAG-3' (ΔG = -32.70 kcal/mole).

Other dicistrovirus sequences used for analysis in in Table S1 are: Solenopsis

invicta virus-1 (SINV-1) NC_006559; Kashmir bee virus (KBV) NC_004807; Acute bee

paralysis virus (ABPV) NC_002548; Homalodisca coagulata virus-1 (HoCV-1)

NC_008029; Himetobi P virus (HiPV) NC_003782; Cripavirus NB-1/2011/HUN (Bat

guano dicistrovirus) NC_025219.1; Aphid lethal paralysis virus (ALPV) NC_004365;

Aphid lethal paralysis virus strain brookings (ALPV-brookings) HQ871932; and Triatoma

virus (TrV) NC_003783.

The constructs encoding HA-tagged CrPV ORFx were generated as follows. For

the C-terminally tagged ORFx-HA, a 3X HA tag was inserted between XbaI and ApaI

sites in pAc5.1/V5-HisB (Thermo). CrPV ORFx (nucleotides 6254-6376) was

subsequently cloned in-frame using KpnI and XbaI sites. For the N-terminally tagged

HA-ORFx a synthesized fragment of DNA (gBlock; IDT) containing a 3X HA tag

followed by the CrPV Orfx sequence was cloned into the pAc5.1 vector with KpnI and

ApaI. The transmembrane mutant of ORFx was generated via site-directed

mutagenesis on the C-terminally HA-tagged construct. All constructs were verified via

sequencing.

In vitro transcription and RNA transfection. pCrPV-2 (Kerr et al., 2015) and

derivative plasmids were linearized with Ecl136II. RNA was transcribed using a T7 RNA

polymerase reaction and subsequently purified with a RNeasy kit (Qiagen). The integrity

and purity of the RNA were confirmed on a 1.2% denaturing formaldehyde agarose gel.

Transfection of _in vitro_ synthesized RNA into S2 cells was performed using

Lipofectamine 2000 (Invitrogen) as per the manufacturer's instructions. 3 µg of RNA

derived from either pCrPV-2 or its cognate mutants were used for transfection using 2.5

x 10^6 cells.

In vitro translation assays. Plasmids were linearized with BamHI and then purified

using QiaQuick Purification columns (Qiagen). The Sf-21 TnT Coupled

Transcription/Translation System was utilized for all translation assays (Promega). In

this system, linearized plasmid is added to the extracts and RNA is synthesized by T7

polymerase and subsequently translated. Total reaction volume was 10 µL containing:

6.7 µL of Sf-21 cell extract (Promega), 0.3 µL of L-[^{35}S]- methionine/cysteine

(PerkinElmer, >1000 Ci/mmol) and 1 µg linearized plasmid DNA. Each reaction mixture

was incubated at 30°C for 2 hours and then resolved on a 15% SDS-PAGE. Gels were

dried and radioactive bands were monitored via phosphoimager analysis. For

quantitation of bands, the number of methionines and cysteines were accounted for and

normalized for each protein. For T2A containing constructs (Wang et al., 2013),

luciferase activity was monitored using a Dual-Luciferase reporter assay system

(Promega) and an Infinite M1000 PRO microplate reader (Tecan).

Northern blot analysis. Total RNA was isolated from cells using TRIzol reagent. Equal

amounts of RNA were loaded on a denaturing agarose gel and subsequently

transferred to Zeta-probe blotting membrane (Bio-Rad). DNA probes were radiolabelled

with $\alpha[P^{32}]$-ATP (MP Biomedicals) using a DecaLabel DNA labeling kit (Fermentas).

Probes were incubated with membrane in hybridizatn buffer (7% SDS, 0.5 M Sodium

phosphate, 1 mM EDTA) at 65°C overnight. Membranes were rinsed with H_2O and

radioactive bands were detected via phosphoimager analysis (Storm; GE Healthcare).

The DNA probe to detect CrPV RNA was generated by PCR amplification using primers

5'-TCCTCAAGCCATGTGTATAGGA-3' and 5'-GTGGCTGAAATACTATCTCTGG-3'.

Western blots. Equal amounts of S2 protein lysates were resolved on a 12% SDS-

PAGE gel and then transferred to a polyvinylidene difluoride Immobilon-FL membrane

(Millipore). Membranes were blocked for 30 min at room temperature with 5% skim milk

in TBST (50 mM Tris, 150 mM NaCl, 1% Tween-20, pH 7.4). Blots were incubated for 1

h at room temperature with the following antibodies: CrPV ORF1 (raised against CrPV

RdRp) rabbit polyclonal (1:10,000), CrPV ORF2 (raised against CrPV VP2) rabbit

polyclonal (1:10,000) (Garrey et al., 2010), mouse anti-tubulin (1:1000; Santa Cruz),

mouse anti-KDEL (1:1000;), or mouse anti-cytochrome C (1:1000; AbCam).

Membranes were washed 3 times with TBST and incubated with either goat anti-rabbit

IgG-HRP (1:20,000; GE Healthcare), goat anti-mouse IgG-HRP (1:5000; Santa Cruz),

IRDye 800CW goat anti-mouse (1:10,000; Li-Cor Biosciences), or IRDye 800CW goat

anti-rabbit (1:20,000; Li-Cor Biosciences) for 1 h at room temperature. An Odyssey

imager (Li-Cor Biosciences) or enhanced chemiluminescence (Thermo) were used for

detection.

Protein isolation and N-terminal sequencing. An ORFx-6XHis-tag was introduced

into the bicistronic construct lacking the T2A sequence. The construct was linearized

and protein was synthesized in an *in vitro* translation extract as previously described

(see above). Reactions were pooled and diluted with Buffer A (50 mM NH_4HCO_3, 500

mM NaCl, 15 mM Imidazole, pH 8.0) before incubating with Ni-NTA beads (QIAGEN)

overnight at 4°C. After incubation, beads were then washed 3 times with Buffer A before

eluting with Buffer B (50 mM NH_4HCO_3, 500 mM NaCl, 500 mM Imidazole, pH 8.0).

Eluted protein was concentrated over a 30 kDa MWCO filter, separated via Tris-glycine

SDS-PAGE, and transferred to a PVDF membrane. Membranes were stained with

Coomassie brilliant blue dye and the corresponding band was analyzed by Edman

sequencing (SickKids SPARC Biocentre). Likelihood scores were calculated as follows:

area under the curve (AUC) for each amino acid in each cycle was determined using

ImageJ software. Each AUC was then divided by the AUC from the previous cycle

subtracted by one to obtain a difference ratio (i.e. $(AUC\ X_n / AUC\ X_{n-1}) - 1$; where X = a

given amino acid and n = cycle number).

LC-MS/MS Analysis. Cell pellets harvested from CrPV-infected cells at 6 hpi were solubilized in 1% sodium deoxy cholate and 50 mM NH_4HCO_3. Protein concentrations were determined via BCA assay (Thermo). Proteins (100 µg) were reduced (2 µg DTT, 37°C, 30 min) and alkylated (5 µg iodoacetamide, RT, 20 min). Samples were digested with trypsin overnight at room temperature. Peptides were acidified with 1% TFA to pH <2.5 and the precipitated deoxycholate was remove via centrifugation. Peptides were desalted and concentrated on C18 STAGE-tips, eluted in 80% acetonitrile, 0.5% acetic acid, and dried in a vacuum concentrator (Eppendorf)(Rappsilber, Ishihama, & Mann, 2003). Samples were resuspended in 20% acetonitrile and 0.1% formic acid before loading on an Agilent 6550 mass spectrometer.

Data were searched using MaxQuant (v1.5.3.30)(Cox & Mann, 2008). Parameters included: carbamidomethylated cysteine (fixed), methionine oxidation (variable), glutamine and asparagine deamidation (variable), and protein N-terminal acetylation (variable); trypsin specific; maximum 2 missed cleavages; 10 ppm precursor mass tolerance; 0.05 Da fragment mass tolerance; 1% FDR; +1 to +7 charge states; and common contaminants were included. Both the *Drosophila* and CrPV protein databases used were the most recent annotations downloaded from UniProt (www.uniprot.org).

Subcellular fractionation by differential centrifugation. 3 X 10^7 S2 cells were transfected with either Drosophila expression vectors (pAc5.1) containing ORFx-HA or HA-ORFx (pORFx-HA or pHA-ORFx) for 48 h at 25°C (XtremeGene; Roche). Cells were then collected by centrifugation at 200 x g for 5 min and washed twice with ice

cold PBS. Cell pellets were then resuspended in SF Buffer (20 mM HEPES [pH 7.4], 10

mM KCl, 1.5 mM MgCl$_2$, 1 mM EDTA, 1 mM EGTA) and incubated on ice for 10 min.

Cells were subsequently disrupted by passaging through a 25 gauge needle 20 times.

Whole cells were removed by centrifugation at 200 x g for 5 min and nuclei were then

collected by centrifugation at 600 x g for 10 min. The mitochondria and endoplasmic

reticulum (ER) were pelleted by centrifugation at 3000 x g for 15 min. Finally, remaining

membrane was pelleted at 16,000 x g for 30 min. The remaining supernatant was

collected as the cytosolic fraction.

Immunofluorescence. S2 cells (1.5 x 10^6) were transfected for 48 h with 2 μg of

pAc5.1/V5-HisB plasmid constructs expressing N- or C-terminally HA-tagged ORFx.

Cells were seeded onto coverslips and fixed with 3% paraformaldehyde for 15 minutes.

Subsequently, cells were permeabilized with 0.2% Triton X-100 in 1X PBS for 1 h

before blocking coverslips with 3% BSA in 1X PBS for 1 h. Coverslips were then

incubated with rabbit anti-HA (1:1000; Cell Signalling) and either mouse anti-Lamin

(1:1000; DSHB), mouse anti-Golgin84 (1:1000, DSHB), or mouse anti-Calnexin 99A

(1:1000, DSHB) in blocking solution overnight at 4°C. Cells were then washed three

times with 1X PBS. Finally, slides were analyzed using a Leica SP5 confocal

microscope with a 63x oil objective lens and a 2x digital zoom. Z-stacks of 15 slices

each were taken of for each condition.

Fly stocks and viral injections. Flies (*Isogenic w^{1118}*; Bloomington Drosophila Stock

Center) were maintained on standard cornmeal food at 25°C and 70% humidity with a

12 h light-dark cycle. Freshly eclosed virgin males and females were separated and

collected in groups of 10 each. Flies (10 males and 10 females) were injected with 200

nL of PBS, CrPV-2, CrPV-S12, or CrPV-S19 (5000 FFU) using a PV830 PicoPump

(World Precision Instruments) and transferred to standard food. Mortality was monitored

daily.

ACKNOWLEDGEMENTS

We thank John Atkins for insightful discussions. Kathy Pan and Seonghoon Lee

contributed a subset of data in Figure 1B and 2B, respectively. This study was

supported by a Canadian Institutes of Health Research (PJT-148761) to E.J. and to D.A.

(MOP-130517), an NSERC Discovery Grant to L.F. and an NSERC CGS-D fellowship

to C.K.

COMPETING INTERESTS

The authors declare no competing interests.

 FIGURE LEGENDS

Figure 1. Biochemical analysis of potential +1-frame translation by Dicistrovirus

IGR IRESs. A) *In silico* **analysis for potential +1-frame translation mediated by IGR**

IRESs in Dicistroviruses. List of type I and type II IGR IRESs based on their

secondary structures. Potential +1-frame base pair immediately downstream of their

IGR IRES PKI region as in IAPV IGR IRES, together with the predicted length of ORFxs

and the first amino acids, are shown. **B) IGR IRES mediated +1-frame translation**

using an *in vitro* **translation assay.** The translation of firefly luciferase (FLuc), which is

fused in the 0-frame or +1-frame, is driven by the individual IGR IRES within the

bicistronic reporter construct. Linearized reporter constructs are incubated in Sf21 *in*

vitro translation insect cell extract at 30 °C for 2 hours in the presence of [^{35}S]

methionine/cysteine. Translation of FLuc and RLuc was monitored by phosphorimager

analysis after resolving on a 16% SDS PAGE. On the bottom panel, the ratio of

FLuc/RLuc are quantified and normalized to the IAPV 0-frame translation. Shown are

averages from at least three independent biological experiments (± SD). **C) The**

secondary structure of CrPV IGR IRES. The ΔPKI, ΔPKII, ΔPKIII, L1.1A, and L1.1B

mutants are shown. Comp denotes compensatory mutations combining ΔPKI CC/GG

and ΔPKI GG/CC. DM - double mutation of ΔPKI and ΔPKII. TM - triple mutation of

ΔPKI, ΔPKII, and ΔPKIII.

Figure 1 - figure supplement 1. **Construction of a T2A construct +1-frame reporter**

construct that recover FLuc enzyme activity. A) The schemes of bicistronic

constructs in which firefly luciferase gene (FLuc) is in frame with the CrPV ORFx are

shown. The T2A sequence (dark grey) is inserted between the CrPV ORFx and the

FLuc gene. The arrow indicates the 'self-cleavage' or 'stop-go' site. A mutation within

the T2A peptide (D to E) inactivates T2A 'self-cleavage' activity. T2A-minus (-T2A) and

T2A-containing ((+)T2A) +1-frame bicistronic constructs were incubated in Sf21 extracts

for 2 h in the presence of [^{35}S] methionine and cysteine. The left panel is a

representative SDS-PAGE monitored by autoradiography, whereas the right panel is the

quantitation of relative luciferase activity (RLA) measured by Dual-Luciferase reporter

assay (Promega). RLA are normalized to that observed with the +1-frame T2A-

containing reporter construct. Shown are averages from at least three independent

biological experiments (± SD). **B)** The quantitation of luciferase activities of IGR IRES

mutants compared to WT T2A containing +1-frame constructs are shown. Displayed are

averages from at least three independent biological experiments (± SD). **C)** The

comparison between +1-frame and 0-frame T2A containing constructs. The ratio of

+1/0-frame translation *in vitro* is 7.1 ± 0.5 %, consistent with that monitored by

autoradiography using T2A-minus constructs (Figure 1B). Note that the FLuc activity

levels from both the 0-frame ΔPKI mutant and the +1-frame ΔPKI mutant indicates to

some extend IGR IRES independent translation. Shown are averages from at least

three independent biological experiments (± SD).

Figure 2. Translation of +1 frame CrPV ORFx is IGR IRES-dependent and initiates

downstream of the PKI region. A) Schematic of mutations introduced downstream

of the IGR IRES. The PKI region and downstream 'spacer' sequence of CrPV IGR

IRES are shown. A series of mutations and stop codons were introduced either on the

0-frame or +1-frame. **B) Analysis of the IGR IRES translation *in vitro*.** Linearized

reporter constructs are incubated in Sf21 translation extract at 30°C for 2 hours in the

present of [^{35}S]-methionine/cysteine. Translation of FLuc and RLuc was monitored by

autoradiography after resolving on a 16% SDS PAGE. A representative gel from at least

three independent biological experiments is shown. * = p-value < 0.05. **C) Quantitation**

of IRES-mediated translation. Translation of FLuc and RLuc monitoring FLuc/RLuc

enzymatic activity were quantified and normalized to wild-type (WT) IAPV IRES. RLuc

monitors scanning-dependent translation acting as an internal control. RLA- relative

light units. Shown are averages from at least three independent biological experiments

(± SD). * = p-value < 0.05.

Figure 3. Translation of CrPV ORFx is dependent on 80S binding and scanning-

independent. A) Effect of L1.1 mutation on +1 frame translation. Comparison of

luciferase activities between WT IGR IRES and L1.1 mutants (L1.1A and L1.1B) that do

not bind the 60S subunit. The value of RLuc and FLuc are normalized to WT. **B) +1-**

frame translation is relatively insensitive to edeine. CrPV IRES-mediated 0- and +1-

frame translation is monitored in the absence or presence of 2 μM edeine. The ΔPKI

mutants in both 0- and +1-frame are used as controls for IGR IRES dependent

translation. The luciferase activities of RLuc and FLuc on the left panel are normalized

to the WT of the 0-frame, whereas the luciferase activities of RLuc and FLuc on the right

panel are normalized to the WT of the +1-frame. **C) Effect of RNA hairpin structures**

on +1 frame translation. Translation of 0- and +1-frame translation of bicistronic

reporter constructs that contain RNA hairpins structures (HP1 ΔG = -17.00 kcal/mole

and HP2 ΔG = -32.70 kcal/mole) between the nucleotides 6135 and 6136. To block the

ribosome scanning the downstream sequence driven by the 0-frame initiation, we

introduced a stop codon at the 3rd codon on the 0-frame on both HP1 and HP2

constructs (0S3). Shown are averages from at least three independent biological

experiments (± SD).

Figure 4. CrPV IRES-mediated +2 frame translation. Mutations and insertions that

shift the ORFx into the +2 frame are shown in the schematic above. Translation assays

were monitored and normalized to that of the wild-type CrPV IRES. Shown are

averages from at least three independent biological experiments (± SD).

Figure 5. Pseudotranslocation is necessary for +1-frame translation. Constructs

containing mutations in the variable loop region (VLR) of the IGR IRES known to disrupt

pseudotranslocation were monitored for translation activity. Luciferase activities were

compared between WT IGR IRES and VLR mutants after *in vitro* translation assays.

Shown are averages from at least three independent biological experiments (± SD).

Figure 5 – figure supplement 1. N-terminal sequencing of ORFx-Fluc-6XHis. (A)

Schematic of the predicted ORFx sequence. 0- and +1-frame protein sequence of

ORF2 (green) and ORFx (red) are shown. **(B) *In* vitro translation of ORFx-FLuc-**

6XHis. Constructs were linearized with BamHI, purified, and incubated with Sf-21

extract including [S^{35}]-Met/Cys. Proteins were separated via SDS-PAGE and visualized

by autoradiography. **(C) Pulldown of radiolabeled ORFx-FLuc-6XHis.** Sf-21

translation extracts were incubated with Ni-NTA beads (see Materials and Methods).

Gels were exposed using autoradiography. Shown is an representative gel from two

independent experiments. **(D) Large scale isolation of ORFx-Fluc-6XHis.** Proteins

from elution fractions in (C) were pooled and concentrated before transferring to a

PVDF membrane. Membranes were stained with Coomassie before band excision and

N-terminal sequencing. The band corresponding to ORFx-FLuc-6XHis is marked with

an asterisk. Shown is a representative pulldown result from two independent

experiments. **(E) Elution profiles of N-terminal sequencing (Edman degradation) of**

ORFx-Fluc-6XHis. The N-terminal elution profiles of amino acid residues (1-4) from

Edman degradation sequencing. Amino acids are shown as individual letter codes. dpu

= diphenolurea; dptu = diphenylthiourea. **(F) Calculated likelihood scores for amino**

acids at each position of ORFx-FLuc-6XHis. Scores were calculated as in Materials

and Methods. Amino acid assignments with the highest confidence are encompassed

by a blue box. Arrows denote which amino acids likely reside in the first position.

Figure 6. The downstream sequence of CrPV IGR IRES promotes +1-frame

translation. A) Chimeric IRES displays +1-frame translation activity. The PSIV IGR

IRES, which shows no +1-frame translation activity (Figure 1 and Figure 3A), is

combined with the downstream sequence of CrPV IGR IRES PKI region. The luciferase

activities are normalized to the wild-type CrPV constructs in both +1- and 0-frame. The

PSIV-CrPV chimera in the 0-frame is used as control to demonstrate that PSIV IGR

IRES functions under this context. **B) Truncation of the spacer region impairs +1-**

frame translation from the CrPV IGR IRES. Schematic displaying truncations within

the spacer region between the PKI domain and the +1 frame start site is shown above.

In vitro translation assay of the CrPV IRES bicistronic construct with the respective

mutations were monitored by luciferase activities. Both RLuc and FLuc activities are

normalized to that of WT constructs. Results are shown as a normalized ratio of

FLuc/RLuc activity (left) and as separated RLuc and Fluc activities (right) Shown are

averages from at least three independent biological experiments (± SD). RLA = relative

luciferase activity.

Figure 6 - figure supplement 1. Effects of deletion and insertion mutations in the

CrPV spacer region on +1-frame translation. Schematic of the spacer region

between the PKI domain and the +1 frame start site is shown above. *In vitro* translation

assay of the CrPV IRES bicistronic construct with the indicated mutations or an insertion

mutation (Ins) of 30 random nucleotides between G_{6217} and C_{6218} or between C_{6235} and

A_{6236} are monitored by luciferase activities. Both 0- and +1-frame activities are

normalized to that of WT constructs. Shown are averages from at least three

independent biological experiments (± SD). RLA = relative luciferase activity.

Figure 7. ORFx is expressed in CrPV infected S2 cells. The predicted +1 frame

amino acid sequence is shown above. Residues that are italicized represent the amino

acid sequence if initiation occurred adjacent to the IGR IRES, whereas the bolded K

lysine residue denotes the position of the 13[th] amino acid. *Drosophila* S2 cells were

infected with CrPV (MOI 10). Proteins were extracted, digested with trypsin and

subjected to LC-MS/MS analysis. Peptide fragment spectra were searched against a

Drosophila uniprot database plus CrPV proteins. Two peptides (highlighted) were

detected from the trypsin digestion of S2 cell lysate at 6 hours post infection. Individual

fragment ions are annotated in the spectra and in the sequence representation.

Figure 8. Role of CrPV ORFx in CrPV-infected S2 cells. A) Schematic of mutations

in the CrPV infectious clone, CrPV-2. The downstream nucleotide sequence of CrPV-

2 IGR IRES and its potential amino acid sequence of +1-frame ORFx are shown. The

ORF2-Stop mutant depletes the synthesis of 0-frame viral structural proteins. Mutants

S12 and S19 place a stop codon on the +1-frame with a synonymous mutation in the 0-

frame. Note that CrPV-2 (Accession: KP974706) sequence has a $C_{6279}A$ mutation

(denoted with a black arrowhead) compared to the original CrPV (Accession:

AF218039) sequence. Residues that are italicized, underlined and in bold are the

peptides identified by mass spectrometry in Figure 7. **B) CrPV ORFx has no effect on**

viral protein synthesis *in vitro.* The *in vitro* transcribed RNA from CrPV-2 or the

cognate mutants was incubated in Sf21 cell extract at 30°C for 2 hours in the present of

[^{35}S]-methionine/cysteine. Translation of viral proteins were monitored by

autoradiography using 12% SDS PAGE. A representative gel of at least three

independent biological experiments is shown. **C) CrPV ORFx has no effect on viral**

protein synthesis and RNA replication in S2 cells. The *in vitro* transcribed RNA of

CrPV-2, S12, or S19 mutants was transfected into S2 cells. Virus was harvested and

used to infect naïve S2 cells at MOI 10. Cells were metabolically labeled with [^{35}S]-

methionine/cysteine for 30 min at 2, 4, 6 and 8 h.p.i. Lysates were collected and

resolved on a 12% SDS PAGE. Western blot analysis against the CrPV RdRp and VP2

and Northern blot analysis of viral RNA are shown. Shown is a representative gel from

three independent biological experiments. **D) CrPV ORFx has no effect on viral**

replication in S2 cells. S2 cells were infected with CrPV-2, CrPV-S12, or CrPV-S19

viruses at MOI 1.0. Viral titres were measured at 6, 9, 12, 24, and 30 h.p.i for MOI 1.

Shown are average from at least three independent biological experiments (±SD).

Figure 9. Pathogenesis of a CrPV mutant lacking ORFx is attenuated in adult

***Drosophila melanogaster* flies. A)** Adult flies (Iso w^{1118}; 10 male, 10 female) were

injected intrathoracically with 5000 FFU of CrPV-2, CrPV-S12, CrPV-S19, or PBS.

Subsequently, flies were flipped onto standard media and survival was monitored daily.

Shown is a graph representing the average from three separate biological experiments

(±SEM). **B)** Flies were harvested at 5 days post injection (dpi) and homogenized in PBS.

Viral titres were determined as described in materials and methods. Error bars

represent the standard error of the mean. **C)** Western blot against CrPV VP2 on lysates

from flies 5 dpi. Shown is a representative gel from at least three independent biological

experiments.

Figure 10. Subcellular localization of CrPV ORFx. S2 cells were transfected with a

construct expressing C-terminally HA-tagged ORFx or a mutant version of ORFx in the

transmembrane region and incubated for 48 h. Following incubation, cells were fixed,

permeabilized and co-stained with HA antibody and antibodies against nuclear lamina

(Lamin), the Golgi apparatus (Golgin84), the ER (Calnexin 99A), and cytoplasm (alpha-

Tubulin). Shown are representative Z-stack micrographs from three independent

experiments. Scale bars represent 15 µm.

Figure 10 – figure supplement 1. Cytotoxicity of CrPV ORFx expression in

Drosophila S2 cells. A) The transmembrane helix property of CrPV ORFx. The

predicted CrPV ORFx sequence stemming from the 13^{th} codon was analyzed for

transmembrane helices using ProtScale and HMMTOP. The mutated form of ORFx is

shown on the right. Shown above are hydropathy plot outputs from ProtScale and below

are HMMTOP predictions. The red amino acid region indicates the putative

transmembrane helix predicted by HMMTOP. **B) Ectopic expression of CrPV ORFx**

causes slight cytotoxicity in S2 cells. S2 cells (1.5×10^{6}) were transfected with

expression constructs containing N- or C-terminally HA-tagged ORFx or empty vector

and monitored over the course of 48 hours for cell death via trypan blue staining. * = p-

value < 0.05.

1010

Figure 10 – figure supplement 2. CrPV ORFx associates with membrane fractions.

S2 cells were transfected with expression constructs encoding HA-tagged ORFx at the

N- or C-terminus for 48 hours. Cells were then lysed and fractionated by differential

centrifugation (see material and methods). Fractions were then analyzed by Western

blot to detect the presence of CrPV ORFx or cellular markers. Shown is a

representative gel from two independent biological experiments.

1017

Figure 11. Model of CrPV IGR IRES mediated +1-frame ORFx translation. The

majority of ribosomes bounds to the IGR IRES initiate translation in the 0-frame from the

GCU start codon. A subset of ribosomes bound to the CrPV IGR IRES (~5%) initiates

translation in the 0 frame and undergoes two pseudotranslocation events before

repositioning downstream at the +1 frame 13th codon (AAA) to translate ORFx. The

model proposes the AAA codon is vacant to allow delivery of the Lys-tRNA.

1024

REFERENCES

Abaeva, I. S., Pestova, T. V., & Hellen, C. U. (2016). Attachment of ribosomal complexes and retrograde scanning during initiation on the Halastavi arva virus IRES. *Nucleic Acids Res, 44*(5), 2362-2377. doi: 10.1093/nar/gkw016

Abeyrathne, P. D., Koh, C. S., Grant, T., Grigorieff, N., & Korostelev, A. A. (2016). Ensemble cryo-EM uncovers inchworm-like translocation of a viral IRES through the ribosome. *Elife, 5*. doi: 10.7554/eLife.14874

Adhin, M. R., & van Duin, J. (1990). Scanning model for translational reinitiation in eubacteria. *J Mol Biol, 213*(4), 811-818. doi: 10.1016/S0022-2836(05)80265-7

Au, H. H., Cornilescu, G., Mouzakis, K. D., Ren, Q., Burke, J. E., Lee, S., . . . Jan, E. (2015). Global shape mimicry of tRNA within a viral internal ribosome entry site mediates translational reading frame selection. *Proc Natl Acad Sci U S A, 112*(47), E6446-6455. doi: 10.1073/pnas.1512088112

Au, H. H., & Jan, E. (2014). Novel viral translation strategies. *Wiley Interdiscip Rev RNA, 5*(6), 779-801. doi: 10.1002/wrna.1246

Baranov, P. V., Atkins, J. F., & Yordanova, M. M. (2015). Augmented genetic decoding: global, local and temporal alterations of decoding processes and codon meaning. *Nat Rev Genet, 16*(9), 517-529. doi: 10.1038/nrg3963

Bonning, B. C., & Miller, W. A. (2010). Dicistroviruses. *Annu Rev Entomol, 55*, 129-150. doi: 10.1146/annurev-ento-112408-085457

Cevallos, R. C., & Sarnow, P. (2005). Factor-independent assembly of elongation-competent ribosomes by an internal ribosome entry site located in an RNA virus that infects penaeid shrimp. *J Virol, 79*(2), 677-683.

Chen, J., Coakley, A., O'Connor, M., Petrov, A., O'Leary, S. E., Atkins, J. F., & Puglisi, J. D. (2015). Coupling of mRNA Structure Rearrangement to Ribosome Movement during Bypassing of Non-coding Regions. *Cell, 163*(5), 1267-1280. doi: 10.1016/j.cell.2015.10.064

Chtarbanova, S., Lamiable, O., Lee, K. Z., Galiana, D., Troxler, L., Meignin, C., . . . Imler, J. L. (2014). Drosophila C virus systemic infection leads to intestinal obstruction. *J Virol, 88*(24), 14057-14069. doi: 10.1128/JVI.02320-14

Colussi, T. M., Costantino, D. A., Zhu, J., Donohue, J. P., Korostelev, A. A., Jaafar, Z. A., . . . Kieft, J. S. (2015). Initiation of translation in bacteria by a structured eukaryotic IRES RNA. *Nature, 519*(7541), 110-113. doi: 10.1038/nature14219

Costantino, D. A., Pfingsten, J. S., Rambo, R. P., & Kieft, J. S. (2008). tRNA-mRNA mimicry drives translation initiation from a viral IRES. *Nat Struct Mol Biol, 15*(1), 57-64. doi: nsmb1351 [pii]10.1038/nsmb1351

Cox, J., & Mann, M. (2008). MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol, 26*(12), 1367-1372. doi: 10.1038/nbt.1511

Dinman, J. D. (2012). Mechanisms and implications of programmed translational frameshifting. *Wiley Interdiscip Rev RNA, 3*(5), 661-673. doi: 10.1002/wrna.1126

Dinos, G., Wilson, D. N., Teraoka, Y., Szaflarski, W., Fucini, P., Kalpaxis, D., & Nierhaus, K. H. (2004). Dissecting the ribosomal inhibition mechanisms of edeine and pactamycin: the universally conserved residues G693 and C795 regulate P-site RNA binding. *Mol Cell, 13*(1), 113-124.

Fernandez, I. S., Bai, X. C., Murshudov, G., Scheres, S. H., & Ramakrishnan, V. (2014). Initiation of Translation by Cricket Paralysis Virus IRES Requires Its Translocation in the Ribosome. *Cell*. doi: 10.1016/j.cell.2014.04.015

Firth, A. E., & Brierley, I. (2012). Non-canonical translation in RNA viruses. *J Gen Virol, 93*(Pt 7), 1385-1409. doi: 10.1099/vir.0.042499-0

Firth, A. E., Wang, Q. S., Jan, E., & Atkins, J. F. (2009). Bioinformatic evidence for a stem-loop structure
5'-adjacent to the IGR-IRES and for an overlapping gene in the bee paralysis dicistroviruses. *Virol*
J, 6, 193. doi: 1743-422X-6-193 [pii]10.1186/1743-422X-6-193
Garrey, J. L., Lee, Y. Y., Au, H. H., Bushell, M., & Jan, E. (2010). Host and viral translational mechanisms
during cricket paralysis virus infection. *J Virol, 84*(2), 1124-1138. doi: JVI.02006-09
[pii]10.1128/JVI.02006-09
Hatakeyama, Y., Shibuya, N., Nishiyama, T., & Nakashima, N. (2004). Structural variant of the intergenic
internal ribosome entry site elements in dicistroviruses and computational search for their
counterparts. *RNA, 10*(5), 779-786.
Herr, A. J., Gesteland, R. F., & Atkins, J. F. (2000). One protein from two open reading frames:
mechanism of a 50 nt translational bypass. *EMBO J, 19*(11), 2671-2680. doi:
10.1093/emboj/19.11.2671
Hertz, M. I., & Thompson, S. R. (2011). In vivo functional analysis of the Dicistroviridae intergenic region
internal ribosome entry sites. *Nucleic Acids Res, 39*(16), 7276-7288. doi: 10.1093/nar/gkr427
Huang, W. M., Ao, S. Z., Casjens, S., Orlandi, R., Zeikus, R., Weiss, R., . . . Fang, M. (1988). A persistent
untranslated sequence within bacteriophage T4 DNA topoisomerase gene 60. *Science,*
239(4843), 1005-1012.
Jackson, R. J., Hellen, C. U., & Pestova, T. V. (2010). The mechanism of eukaryotic translation initiation
and principles of its regulation. *Nat Rev Mol Cell Biol, 11*(2), 113-127. doi: 10.1038/nrm2838
Jan, E. (2006). Divergent IRES elements in invertebrates. *Virus Res, 119*(1), 16-28.
Jan, E., Kinzy, T. G., & Sarnow, P. (2003). Divergent tRNA-like element supports initiation, elongation,
and termination of protein biosynthesis. *Proc Natl Acad Sci U S A, 100*(26), 15410-15415.
Jan, E., Mohr, I., & Walsh, D. (2016). A Cap-to-Tail Guide to mRNA Translation Strategies in Virus-
Infected Cells. *Annu Rev Virol, 3*(1), 283-307. doi: 10.1146/annurev-virology-100114-055014
Jan, E., & Sarnow, P. (2002). Factorless ribosome assembly on the internal ribosome entry site of cricket
paralysis virus. *J Mol Biol, 324*(5), 889-902.
Jang, C. J., & Jan, E. (2010). Modular domains of the Dicistroviridae intergenic internal ribosome entry
site. *RNA, 16*(6), 1182-1195. doi: rna.2044610 [pii]10.1261/rna.2044610
Jang, C. J., Lo, M. C., & Jan, E. (2009). Conserved element of the dicistrovirus IGR IRES that mimics an E-
site tRNA/ribosome interaction mediates multiple functions. *J Mol Biol, 387*(1), 42-58. doi:
S0022-2836(09)00093-X [pii]10.1016/j.jmb.2009.01.042
Johannes, G., Carter, M. S., Eisen, M. B., Brown, P. O., & Sarnow, P. (1999). Identification of eukaryotic
mRNAs that are translated at reduced cap binding complex eIF4F concentrations using a cDNA
microarray. *Proc Natl Acad Sci U S A, 96*(23), 13118-13123.
Kerr, C. H., Ma, Z. W., Jang, C. J., Thompson, S. R., & Jan, E. (2016). Molecular analysis of the factorless
internal ribosome entry site in Cricket Paralysis virus infection. *Sci Rep, 6*, 37319. doi:
10.1038/srep37319
Kerr, C. H., Wang, Q. S., Keatings, K., Khong, A., Allan, D., Yip, C. K., . . . Jan, E. (2015). The 5' untranslated
region of a novel infectious molecular clone of the dicistrovirus cricket paralysis virus modulates
infection. *J Virol, 89*(11), 5919-5934. doi: 10.1128/JVI.00463-15
Kozak, M., & Shatkin, A. J. (1978). Migration of 40 S ribosomal subunits on messenger RNA in the
presence of edeine. *J Biol Chem, 253*(18), 6568-6577.
Krishna, N. K., Marshall, D., & Schneemann, A. (2003). Analysis of RNA packaging in wild-type and mosaic
protein capsids of flock house virus using recombinant baculovirus vectors. *Virology, 305*(1), 10-
24.
Lang, B. F., Jakubkova, M., Hegedusova, E., Daoud, R., Forget, L., Brejova, B., . . . Nosek, J. (2014).
Massive programmed translational jumping in mitochondria. *Proc Natl Acad Sci U S A, 111*(16),
5926-5931. doi: 10.1073/pnas.1322190111

Muhs, M., Hilal, T., Mielke, T., Skabkin, M. A., Sanbonmatsu, K. Y., Pestova, T. V., & Spahn, C. M. (2015).
Cryo-EM of ribosomal 80S complexes with termination factors reveals the translocated cricket
paralysis virus IRES. *Mol Cell, 57*(3), 422-432. doi: 10.1016/j.molcel.2014.12.016
Nakashima, N., & Uchiumi, T. (2009). Functional analysis of structural motifs in dicistroviruses. *Virus Res,*
139(2), 137-147. doi: S0168-1702(08)00230-X [pii]10.1016/j.virusres.2008.06.006
Nishiyama, T., Yamamoto, H., Shibuya, N., Hatakeyama, Y., Hachimori, A., Uchiumi, T., & Nakashima, N.
(2003). Structural elements in the internal ribosome entry site of Plautia stali intestine virus
responsible for binding with ribosomes. *Nucleic Acids Res, 31*(9), 2434-2442.
Nomura, M., Gourse, R., & Baughman, G. (1984). Regulation of the synthesis of ribosomes and ribosomal
components. *Annu Rev Biochem, 53*, 75-117. doi: 10.1146/annurev.bi.53.070184.000451
Paek, K. Y., Hong, K. Y., Ryu, I., Park, S. M., Keum, S. J., Kwon, O. S., & Jang, S. K. (2015). Translation
initiation mediated by RNA looping. *Proc Natl Acad Sci U S A, 112*(4), 1041-1046. doi:
10.1073/pnas.1416883112
Pestova, T. V., & Hellen, C. U. (2003). Translation elongation after assembly of ribosomes on the Cricket
paralysis virus internal ribosomal entry site without initiation factors or initiator tRNA. *Genes*
Dev, 17(2), 181-186.
Petrov, A., Grosely, R., Chen, J., O'Leary, S. E., & Puglisi, J. D. (2016). Multiple Parallel Pathways of
Translation Initiation on the CrPV IRES. *Mol Cell, 62*(1), 92-103. doi:
10.1016/j.molcel.2016.03.020
Pfingsten, J. S., Costantino, D. A., & Kieft, J. S. (2006). Structural basis for ribosome recruitment and
manipulation by a viral IRES RNA. *Science, 314*(5804), 1450-1454. doi: 1133281
[pii]10.1126/science.1133281
Reinganum, C. (1975). The isolation of cricket paralysis virus from the emperor gum moth, Antheraea
eucalypti Scott, and its infectivity towards a range of insect species. *Intervirology, 5*(1-2), 97-102.
Ren, Q., Au, H. H., Wang, Q. S., Lee, S., & Jan, E. (2014). Structural determinants of an internal ribosome
entry site that direct translational reading frame selection. *Nucleic Acids Res, 42*(14), 9366-9382.
doi: 10.1093/nar/gku622
Ren, Q., Wang, Q. S., Firth, A. E., Chan, M. M., Gouw, J. W., Guarna, M. M., . . . Jan, E. (2012). Alternative
reading frame selection mediated by a tRNA-like domain of an internal ribosome entry site. *Proc*
Natl Acad Sci U S A, 109(11), E630-639. doi: 10.1073/pnas.1111303109
Ruehle, M. D., Zhang, H., Sheridan, R. M., Mitra, S., Chen, Y., Gonzalez, R. L., . . . Kieft, J. S. (2015). A
dynamic RNA loop in an IRES affects multiple steps of elongation factor-mediated translation
initiation. *Elife, 4*. doi: 10.7554/eLife.08146
Samatova, E., Konevega, A. L., Wills, N. M., Atkins, J. F., & Rodnina, M. V. (2014). High-efficiency
translational bypassing of non-coding nucleotides specified by mRNA structure and nascent
peptide. *Nat Commun, 5*, 4459. doi: 10.1038/ncomms5459
Sarabhai, A., & Brenner, S. (1967). A mutant which reinitiates the polypeptide chain after chain
termination. *J Mol Biol, 27*(1), 145-162.
Schuler, M., Connell, S. R., Lescoute, A., Giesebrecht, J., Dabrowski, M., Schroeer, B., . . . Spahn, C. M.
(2006). Structure of the ribosome-bound cricket paralysis virus IRES RNA. *Nat Struct Mol Biol,*
13(12), 1092-1096. doi: nsmb1177 [pii]10.1038/nsmb1177
Skabkin, M. A., Skabkina, O. V., Hellen, C. U., & Pestova, T. V. (2013). Reinitiation and other
unconventional posttermination events during eukaryotic translation. *Mol Cell, 51*(2), 249-264.
doi: 10.1016/j.molcel.2013.05.026
Speicher, K. D., Gorman, N., & Speicher, D. W. (2009). N-terminal sequence analysis of proteins and
peptides. *Curr Protoc Protein Sci, Chapter 11*, Unit11 10. doi: 10.1002/0471140864.ps1110s57
Vassilaki, N., & Mavromara, P. (2009). The HCV ARFP/F/core+1 protein: production and functional
analysis of an unconventional viral product. *IUBMB Life, 61*(7), 739-752. doi: 10.1002/iub.201

Wang, Q. S., Au, H. H., & Jan, E. (2013). Methods for studying IRES-mediated translation of positive-
1170   strand RNA viruses. *Methods, 59*(2), 167-179. doi: 10.1016/j.ymeth.2012.09.004

1171   Wills, N. M., O'Connor, M., Nelson, C. C., Rettberg, C. C., Huang, W. M., Gesteland, R. F., & Atkins, J. F.
1172   (2008). Translational bypassing without peptidyl-tRNA anticodon scanning of coding gap mRNA.
1173   *EMBO J, 27*(19), 2533-2544. doi: 10.1038/emboj.2008.170

1174   Wilson, J. E., Pestova, T. V., Hellen, C. U., & Sarnow, P. (2000). Initiation of protein synthesis from the A
1175   site of the ribosome. *Cell, 102*(4), 511-520.

1176   Yamamoto, H., Wittek, D., Gupta, R., Qin, B., Ueda, T., Krause, R., . . . Nierhaus, K. H. (2016). 70S-
1177   scanning initiation is a novel and frequent initiation mode of ribosomal translation in bacteria.
1178   *Proc Natl Acad Sci U S A, 113*(9), E1180-1189. doi: 10.1073/pnas.1524554113

1179   Yamasaki, K., Weihl, C. C., & Roos, R. P. (1999). Alternative translation initiation of Theiler's murine
1180   encephalomyelitis virus. *J Virol, 73*(10), 8519-8526.

1181   Zhang, H., Ng, M. Y., Chen, Y., & Cooperman, B. S. (2016). Kinetics of initiating polypeptide elongation in
1182   an IRES-dependent system. *Elife, 5*. doi: 10.7554/eLife.13429

1183

1184

A

Dicistrovirus	IGR IRES Type	Potential +1 basepair	Predicted +1 ORFx Length (aa)	First +1 frame amino acid
Solenopsis invicta virus-1 (SINV-1)	II	C-G	125	L
Israeli acute paralysis virus (IAPV)	II	U-G	94	A
Kashmir bee virus (KBV)	II	C-G	93	L
Acute bee paralysis virus (ABPV)	II	U-G	92	P
Cricket paralysis virus (CrPV)	I	U-G	53	L
Drosophila C virus (DCV)	I	U-G	51	L
Plautia stali intestine virus (PSIV)	I	U C (?)	43	K
Big Sioux River virus (BSRV)	I	U-G	40	L
Black queen cell virus (BQCV)	I	U-G	26	L
Homalodisca coagulata virus- 1	I	U-G	17	Q
Taura Syndrome virus (TSV)	II	U-G	16	L
Himetobi P virus (HiPV)	I	U-G	14	Q
Bat guano dicistrovirus	I	U-G	14	Q
Mud crab dicistrovirus (MCDV)	II	U-G	9	L
Aphid lethal paralysis virus (ALPV)	I	U-G	7	L
ALPV like Brookings virus	I	U-G	7	Q
Rhopalosiphum padi virus (RhPV)	I	U-G	2	Q
Triatoma virus (TrV)	II	C-G	1	L

B

C

Cricket paralysis virus IGR IRES

Kerr_Figure 2

A

B

C

A

(+1) frame — RLA axis. Groups: WT, ΔPKI, L1.1A, L1.1B

(0) frame — RLA axis. Groups: WT, ΔPKI, L1.1B

Legend: RLuc (open), FLuc (filled)

B

RLA axis. Edeine +/−. Frame WT (0), ΔPKI (0), WT (+1), ΔPKI (+1)

RLA Normalized to +1 frame WT. Edeine +/−

Legend: RLuc (open), FLuc (filled)

C

```
        PKI      6218          0S3                    HP
                        U  GA
WT  UACCU G CUA  CAU UUC AAG AUA AAC AAG AAA AUU CAC ACA UUG AAA AUG AAG AUA ...
```

(+1) frame — RLA axis. Groups: WT, ΔPKI, HP1, HP1 0S3, HP2, HP2 0S3

(0) frame — RLA axis. Groups: WT, ΔPKI, HP1, HP2

Legend: RLuc (open), FLuc (filled)

Peptide #1 **Peptide #2**

Predicted Sequence: N - *AL* **K**MKIKDLCQNRKK **LYILLAK ELPLVLLR** SLISLIFQLIIW - **C**

A

```
                                0 frame
                                ORF2-stop                    S12 (+1 frame)
                                  uaa                          uag    14
                                 ___                          ___     •
(+1) frame        L  H  F  K  I  N  K  K  I  H  T  L  K  M  K  I  K  D
(0)  frame        A  T  F  Q  D  K  Q  E  N  S  H  I  E  N  E  D  K  R
           UACCU gcuacauuucaagauaaacaagaaaauucacacauugaaaaugaagauaaaaga
           _____              •                      •
           PKI    6218       6232                  6251

           S19 (+1 frame)
             uaa
            ___
(+1) frame L  C  Q  N  R  K  K  L  Y  I  L  L  A  K  E  L  P  L  V  L
(0)  frame L  M  S  E  Q  K  E  I  V  H  F  V  S  E  G  V  T  P  S  T
           cuuagucagaacagaaagaaaauuguacauuuguguuagcgaaggaguuaccccuaguacu
            •                                                     •
           6272    ↑                                            6317

(+1) frame L  R  S  L  I  S  L  I  F  Q  L  I  I  W  T
(0)  frame T  A  L  P  D  I  V  N  L  S  T  N  Y  L  D  M
           acugcgcucccugauaucguuaaucuuucaacuaauuaucuggacauga
           •                              •         •
           6332                          6368      6379
```

B

D

C

Kerr_Figure 9

A

B

C

Kerr_Figure 11

Translational Bypass