1

IRES-mediated ribosome repositioning directs translation of a +1 overlapping

ORF that enhances viral infection

[†]Craig H Kerr[1,2], [†]Qing S Wang[1], Kyung-Mee Moon[1,2], Kathleen Keatings[3], Douglas W Allan[3], Leonard J Foster[1,2], Eric Jan*[1]

[1]Department of Biochemistry and Molecular Biology, [2]Centre for High-Throughput Biology, [3]Department of Cellular and Physiological Sciences, University of British Columbia, Vancouver BC, V6T 1Z3 Canada

[†]contributed equally

*corresponding author: ej@mail.ubc.ca

keywords: ribosome, frameshift, dicistrovirus, drosophila, virus

ABSTRACT

RNA structures can interact with the ribosome to alter translational reading frame

maintenance and promote recoding that result in alternative protein products. Here, we

show that the internal ribosome entry site (IRES) from the dicistrovirus Cricket paralysis

virus drives translation of the 0-frame viral polyprotein and an overlapping +1 open

reading frame, called ORFx, via a novel mechanism whereby a subset of ribosomes

recruited to the IRES bypasses downstream to resume translation at the +1-frame 13th

non-AUG codon. A mutant of CrPV containing a stop codon in the +1 frame ORFx

sequence, yet synonymous in the 0-frame, is attenuated compared to wild-type virus in

a *Drosophila* infection model, indicating the importance of +1 ORFx expression in

promoting viral pathogenesis. This work demonstrates a novel programmed IRES-

mediated recoding strategy to increase viral coding capacity and impact virus infection,

highlighting the diversity of RNA-driven translation initiation mechanisms in eukaryotes.

Significance Statement

Viruses use alternate mechanisms to increase the coding capacity of their viral

genomes. Here, we provide biochemical evidence that ribosomes recruited to the

dicistrovirus cricket paralysis virus IRES undergo a bypass event to direct translation of

a downstream +1 frame overlapping open reading frame, called ORFx. Mutations that

block ORFx expression inhibit +1 frame translation and infection in fruit flies. These

findings highlight the diversity of RNA-driven translation initiation mechanisms in

eukaryotes.

INTRODUCTION

The ribosome mediates translation involving decoding the open reading frame codon by codon through delivery of the correct aminoacyl-tRNAs to the ribosomal A site. This fundamental process occurs with high fidelity for proper gene expression in all species. However, mechanisms exist that can alter the translational reading frame, thus producing alternative protein products from a single RNA (1). In general, these mechanisms termed recoding, involve a specific RNA structure or element that interacts with the ribosome to cause the translating ribosome to shift reading frame by -1/+1/+2 allowing it to read through stop codons or bypass sequences and restart translation downstream (2-4). Study of these mechanisms has been enlightening; revealing key ribosome:RNA interactions that alter fundamental processes in the mechanics of ribosome decoding and reading frame maintenance. Importantly, recoding mechanisms are now appreciated as important regulatory processes that can impact the fate of protein expression in cells and viral infections (1, 5). Unlike these recoding mechanisms that involve an actively translating ribosome, the intergenic internal ribosome entry site (IRES) within a subset of dicistroviruses has the unusual property to directly recruit the ribosome and initiate translation from overlapping 0 and +1-frame codons to produce two distinct proteins (6). Here, we report a novel recoding mechanism and translational initiation pathway whereby a related dicistrovirus IRES directs the ribosome to initiate translation downstream.

Most eukaryotic mRNAs utilize a cap-dependent scanning mechanism involving >12 translation initiation factors to recruit the ribosome and initiate translation from an AUG start codon (7). Alternatively, an IRES, in general, is a structured RNA

element that facilitates 5' end-independent translation using subsets of translation

initiation factors, thus providing an advantage during viral infection or under times of

stress when cap-dependent translation is compromised (3-5).

Of the different classes of viral IRESs based on factor requirement and

mechanism, the intergenic IRES of the *dicistroviridae* family stands out as the most

streamlined using a unique mechanism where it directly binds 40S and 80S ribosomes

without the need for canonical initiation factors or initiator Met-tRNA$_i$ and initiates

translation from a non-AUG codon (8-12). The dicistrovirus IRES is composed of three

pseudoknots (PKI, II, and III) that separate into distinct domains; PKII and PKIII fold

independently to create the ribosome-binding domain while PKI mediates positioning of

the ribosome and establishes the translational reading frame (8, 10, 13). Structural

studies have indicated that PKII and PKIII form a compact core structure, and the PKI

region adopts a conformation that mimics an anti-codon:codon interaction that binds the

conserved core of the ribosome in the A site (14-17). From here, in an elongation factor

2 dependent manner, the IGR IRES pseudo-translocates to the ribosomal P site,

followed by aminoacyl-tRNA delivery to the A site and a second round of eEF2-

dependent pseudo-translocation of the IGR IRES to the E site of the ribosome (17-19).

Altogether, the IGR IRES acts as a complete RNA machine that supersedes initiation

factors and commandeers the ribosome, a strategy that is essential for viral protein

synthesis in dicistrovirus-infected cells (20).

In general, the dicistrovirus IGR IRESs are conserved at the structural, but not

sequence level and are classified into two sub-groups (termed Type I and II) based on

the presence of distinct structural elements; the main distinction comes from a larger

L1.1 loop and an additional stem-loop (SLIII) in Type II IRESs (21, 22). SLIII allows for

the PKI domain of Type II IRESs to mimic the global shape of a tRNA in addition to

assisting in reading frame selection and the larger L1.1 region functions to mediate 60S

recruitment (15, 18, 23). The domains of Type I and II IGR IRESs function similarly to

directly recruit 80S ribosomes and initiate translation (24-27). Recent high resolution

cryo-EM structures of the IGR IRES bound to the 80S ribosome have demonstrated that

the IRES initially binds in the A site (17, 19, 28). Movement of the IRES involves an

eEF2-dependent pseudo-translocation event where the ribosome rotates up to 10°

allowing PKI to move into the P site in an inchworm-like manner (28). This allows for the

non-AUG initiation codon of the IRES to be presented in the A site for the incoming

amino-acyl tRNA. The first pseudo-translocation event and delivery of the first amino-

acyl tRNA are the rate-limiting steps of initiation on the IGR IRES (29).

Biochemical, phylogenetic, and bioinformatics analyses have demonstrated that

a subset of Type II IGR IRESs can direct translation of a hidden +1 open reading frame

(ORF), termed ORFx, within ORF2 of the viral genome (6, 30). The functional role of

ORFx during viral infection remains elusive. Extensive mutagenesis of the PKI domain

of the *Israeli acute paralysis virus* (IAPV) IGR IRES has revealed that 0 and +1 frame

translation can be uncoupled, suggesting that the IGR IRES may adopt specific

conformations that govern the translational reading frame (18, 31). Generally, Type I

and II IGR IRESs are thought to operate similarly in mechanism. Specific domains

between the two types are functionally interchangeable (24). In the present study, we

investigate the capacity of other IGR IRESs from dicistroviruses to facilitate +1-frame

translation. We show that the IGR IRES from Cricket paralysis virus (CrPV) can

synthesize an ORFx protein using an unexpected mechanism that involves IRES-

mediated ribosome bypassing. Furthermore, we provide insight into the role of ORFx

during CrPV infection and show that mutants deficient in ORFx have impaired virulence

in adult flies, thus uncovering a novel viral recoding strategy that is essential for viral

infection.

RESULTS

IGR IRES-dependent +1-frame translation is not conserved throughout

Dicistroviridae

We previously showed that a subset of dicistrovirus IGR IRESs can direct

translation in the 0- and +1-frames and that a base pair adjacent to the PKI domain is

important for initiation in the +1-frame (6, 31). The IGR IRESs are classified into two

types: Type I, and II, with the main difference being an extra SLIII within the PKI domain

of Type II IRES. Since the honeybee and fire ant viruses harbour Type II IGR IRESs

that can support +1-frame ORFx translation, we investigated whether Type I IRESs also

had the capacity for +1-frame translation. We first surveyed the other dicistrovirus IGR

IRESs and their downstream sequences for a potential base pair adjacent to the PKI

domain and an overlapping +1 open reading frame both *in silico* (Fig. S1). The IAPV

and ABPV IGR IRESs contain an adjacent U-G base pair whereas the SINV-1 and KBV

IRESs have a C-G base pair (Fig. S1). All but one of the IGR IRESs contains a potential

base pair adjacent to the PKI domain, with the majority of them possessing a potential

U-G base pair. As shown previously, the honeybee and fire ant dicistrovirus ORFx

proteins are approximately 94-125 amino acids in length (30). The predicted +1 ORFx

lengths of other dicistroviruses range from 1-53 amino acids in length. Apart from the

honeybee and fire ant dicistrovirus ORFx, the longest putative ORFx sequences are

found within the genomes of the CrPV and DCV at 53 amino acids in length.

To determine whether the other IGR IRESs can direct +1-frame translation, we

cloned each IGR IRES within the intergenic region of a dual luciferase bicistronic

construct (Fig. 1A). The Renilla luciferase (36 kDa; RLuc) monitors scanning-dependent

translation whereas firefly luciferase (FLuc) is translated by the IRES. To measure

reading frame selection, the firefly luciferase gene is fused in-frame either in the 0 or +1-

frame and translated proteins were monitored by incorporation of [^{35}S]-

methionine/cysteine. Additionally, we generated bicistronic reporter constructs that

contain a T2A 'stop-go' sequence, which allows quantitation of luciferase activity (32)

(Fig. S2). Since we previously showed robust IAPV IRES 0 and +1-frame translation (6),

we chose it as a benchmark to compare the +1-frame activity of the other IGR IRESs. In

general, all IGR IRESs can direct 0-frame translation to varying extents *in vitro*.

Normalizing to the IAPV 0-frame translational activity, the CrPV IGR IRES showed the

highest 0-frame translational activity (~170%) whereas the Mud crab dicistrovirus

(MCDV) IGR IRES had the lowest activity (~4%; Fig. 1A). By contrast, only a few IGR

IRESs can support +1-frame translation. The Type II IGR IRESs from Taura syndrome

virus (TSV) and MCDV did not support +1 frame translation suggesting that only a

subset of Type II IGR IRESs can direct 0- and +1-frame translation. Interestingly,

besides the IAPV IGR IRES, the BQCV and CrPV IRESs mediated +1-frame translation

above background levels, 80% and 5% of 0-frame translation, respectively. In summary,

only a subset of IGR IRESs can facilitate both 0 and +1-frame translation.

CrPV +1-frame translation is IGR IRES-dependent and initiates downstream

To explore +1-frame translation mechanisms further, we focused on the CrPV

IRES (Fig. 1B). We first determined whether the structural integrity of the CrPV PKI

domain is important for +1-frame translation. For these assays, we used a bicistronic

reporter construct that contains the CrPV IRES where a firefly luciferase gene was

subcloned into the +1-frame downstream of the IRES with CrPV nucleotides 6217-6387

(Accession KP974707.1), which includes the predicted +1-frame CrPV ORFx (65 kDa;

ORFx-FLuc). This dual luciferase reporter construct allows simultaneous monitoring of

scanning-dependent and CrPV IRES-mediated 0/+1-frame translation as a shortened 0-

frame protein (~11 kD) is also translated in addition to the +1-frame (Fig. 2A). Synthesis

of all three proteins is detected by incubating the bicistronic construct in a Sf21

translation extract in the presence of [^{35}S]-met/cys (Fig. 2B, lane 1) (6). As shown in Fig.

1A, CrPV IRES +1-frame ORFx translation is approximately 5% of 0-frame translation.

Mutating CC_{6214-5} to GG, which disrupts PKI base pairing and abolishes CrPV IRES

activity, resulted in negligible or diminished 0- and +1-frame translation whereas a

compensatory mutation that restores PKI base pairing rescued translation (10) (Fig. 2B,

lanes 2 and 3, Fig. S2), indicating that the integrity of the IRES and the PKI domain is

required for CrPV IRES +1-frame translation.

The adjacent U-G base pairing of the IAPV IRES is important for +1-frame

translation (6). CrPV also has the capacity to form a wobble base pair adjacent to the

PKI domain through nucleotides U_{6186} and G_{6217}, potentially directing translation from

the first +1-frame CUA leucine codon (Fig. 1B). To determine if this base pair is

necessary to drive CrPV IRES +1-frame translation, we mutated U_{6186} and G_{6217} to other

bases. Mutating U_{6186} to C or G led to an approximate 18%-23% reduction in +1 frame

activity and mutating G_{6217} to C, A, or U resulted in roughly a 39%, 24%, and 36%

reduction in +1-frame activity, respectively. Although each mutation reduced +1-frame

translational activity to some degree, none of the mutations abolished it (Fig. 2B, lanes

4-9). These results suggest that unlike with IAPV, base pairing between nucleotides

6186 and 6217 is not absolutely required for CrPV IRES +1-frame translation.

To determine the potential initiation site of CrPV ORFx, we systematically

replaced codons downstream of the IRES with a stop codon and monitored 0- and +1-

frame translation *in vitro* using the bicistronic reporter construct (Fig. 2B, 2C). Overall,

stop codons placed in the +1-frame did not significantly affect 0-frame translation,

indicating that IRES activity was not compromised (Fig. 2B, 2C). Replacing individual

codons between the 1st and the 12th +1-frame codon with a stop codon inhibited to

varying extents (between 36%-71% reduction compared to wild-type) but did not

completely abolish +1 frame translation (Fig. 2B, lanes 11-16; 2C). Conversely, +1-

frame translation was completely inhibited when the 13th +1-frame codon and codons

thereafter were replaced with a stop codon (Fig. 2B, lanes 15-16; 2C). Replacing both

the 3rd and 5th +1-frame codons with stop codons reduced +1-frame translation by 33%

but did not eliminate it, suggesting ribosome read-through did not occur. To address the

possibility that IRES translation initiates in the 0-frame and then a fraction of translating

ribosomes shift into the +1-frame, we inserted a stop codon in the 0-frame downstream

of the IRES. As expected, a stop codon in the 0-frame 1st to 4th codons downstream of

the IRES abolished 0-frame translation (Fig. 2B, lanes 18-21). However, the 0-frame

stop codon insertions reduced by 52-77% but did not eliminate +1 frame translation

indicating that ribosomes likely do not shift from the 0 to +1 reading frame. Furthermore,

introducing stop codons in both the 0 and +1 frames did not abolish +1-frame translation,

though inhibited +1-frame translation by ~80% (Fig. 2C, 0 S4/+1 S5). We noted that the

adjacent 14th codon is an AUG methionine. Mutating the AUG to a CGG (M14R) or

GUG (M14V) decreased but did not abolish +1-frame translation (Fig. 2C). Mutating the

13th AAA codon to UUU, AAG or GCU did not abolish +1-frame translation (Fig. 2C),

suggesting that there is flexibility in the codon identity for CrPV ORFx +1-frame

translation. In summary, the mutational analysis indicated that CrPV ORFx translation

requires an intact IRES and a +1-frame 13th nonstop codon.

CrPV +1-frame translation requires 80S assembly and is edeine-insensitive

The current data led us to generate two hypotheses: i) a subset of 40S subunits

recruited to the CrPV IRES scan downstream to start translation at the 13th codon

(scanning hypothesis) or ii) 40S or 80S ribosomes recruited to the IGR IRES bypass the

spacer region to the 13th codon (bypass hypothesis). To address the scanning

hypothesis, we took two approaches. First, we utilized mutants of the CrPV IRES in the

L1.1 loop that is known to be deficient in recruitment of the 60S subunit (15, 23). If

scanning is occurring, 40S subunits recruited to the L1.1 IRES may scan downstream to

the downstream +1-frame initiation codon. Reporter constructs harbouring mutations in

the L1.1 loop were deficient in 0 and +1-frame translation (Fig. 3A), suggesting that 60S

recruitment by the IGR IRES specifically is necessary for translation in the +1-frame.

Secondly, we utilized the translational inhibitor edeine to assess if scanning was

occurring (Fig. 3B). Edeine prevents the 40S ribosomal subunit from recognizing an

AUG start codon (33, 34). Both 0- and +1-frame translation were resistant to edeine

relative to that of scanning-dependent translation (Fig. 3B), thus suggesting that 40S

scanning is not involved in +1 frame translation.

The integrity of the variable- loop region and pseudo-translocation of the IGR

IRES through the ribosome is required for ORFx expression

    If 80S ribosomes recruited to the IRES are indeed repositioning or "bypassing" to the downstream 13th +1-frame codon, we next looked to investigate the rules which govern this potential mechanism. Since the IGR IRES, which occupies the ribosomal A site upon ribosome binding, undergoes pseudotranslocation to the ribosomal P site to vacate the A site for delivery of the first aminoacyl-tRNA (17), we reasoned that the ribosome bound to the CrPV IGR IRES must have an empty ribosomal P or A site in order to reposition downstream and accommodate the +1 frame start codon. To address this, we introduced mutations in the variable loop region (VLR), which has been shown to disrupt the IGR IRES-mediated pseudotranslocation event (Fig. 1B) (35). Specifically, shortening the length of the VLR by 2 or 3 nucleotides ($\Delta2$ and $\Delta3$, respectively) inhibits the first pseudo-translocation event from the A site to the P site whereas altering the identity of nucleotides A_{6204}, and AA_{6208-9} to guanosines (G-rich) inhibited IRES translocation from the P site to the E site (35). Interestingly, all three VLR mutants decreased +1-frame activity (Fig. 4); both the G-rich and $\Delta3$ mutants demonstrated little to no activity, while the $\Delta2$ mutant still exhibited roughly 50% activity to that of WT. This result is consistent with previous data that the $\Delta2$ mutant IRESs are still able to accommodate a fraction of aminoacyl-tRNA in the A site (~25%), allowing translation to occur (35). Altogether, these results indicate that the pseudotranslocation event of the IGR IRES through the ribosome contributes to +1-frame translation downstream.

The spacer region downstream of the IGR IRES is necessary for +1 frame translation

We next investigated whether the spacer region located between PKI of the IGR

IRES and the downstream AAA codon contributes to IRES-mediated ribosome bypass.

Our results showed that inserting stop codons in the 'spacer region' (Fig. 2A) between

the IRES PKI domain and the +1-frame 13th codon of ORFx inhibited but did not

completely abolish +1-frame translation, suggesting that an element within this spacer

region may promote +1-frame translation. We first addressed whether the CrPV spacer

region is sufficient to direct +1-frame translation by generating a chimeric construct

whereby the PSIV IRES is fused with the CrPV spacer region (Fig. 5). The PSIV IRES

directs strong 0-frame but no or relatively weak +1-frame translation (Fig. 1A). The

PSIV-CrPV spacer chimeric reporter resulted in 0-frame translation (Fig. 5A), indicating

that the spacer region does not affect the activity of the PSIV IRES. In contrast to the

PSIV IRES construct, the chimeric PSIV-CrPV reporter resulted in +1-frame translation,

implying that the CrPV spacer region is sufficient to drive +1-frame translation in the

presence of a functional IGR IRES (Fig. 5A).

To delineate whether there is a specific element within the spacer region that is

required for CrPV +1 frame translation, we systematically deleted from the 3' end of the

spacer region (Fig. 5B). Interestingly, deleting 3 to 27 nucleotides did not affect +1

frame translation. However, deleting 30 nucleotides and leaving seven nucleotides

adjacent to the IRES abolished +1 translation (Fig. 5B). Δ3, Δ6 and Δ27 mutants appear

to have much higher Fluc/Rluc activity when compared to WT, an observation that

requires further examination. In summary, these results suggest that the specific

sequences and its context within the spacer region, particularly the sequences

immediately downstream of the PKI domain, are important for mediating IRES-

dependent +1 frame translation.

+2-frame translation mediated by the CrPV IRES

We hypothesized that if the ribosome is indeed bypassing to a specific 'landing

site' then it should be independent of frame, whereas if the ribosome merely begins

translation in the 0-frame before slipping into the +1-frame, then we expect to see no

ORFx translation. To address this, we inserted a series of nucleotides into our

bicistronic construct that shifts only the ORFx-Fluc into the +2-frame (Figure S3).

Specifically, we inserted either a single nucleotide or up to 7 proceeding the 13th AAA

codon. As a control, we inserted either 6 or 9 nucleotides in the same position that does

not introduce an additional frameshift. Insertion of a U creates a stop codon in the +1-

frame while an inserted C does not. To our surprise, we observed ORFx expression

with all insertions (Figure S3). These results suggest translation of ORFx is reading

frame-independent and indicates that the ribosome may be repositioned from the IRES

to the downstream 13th codon.

CrPV +1-frame ORFx is expressed yet not required for infection in *Drosophila* S2

cells

Our *in silico* and biochemical data indicate that ribosomes recruited to the CrPV

IRES may bypass downstream to the +1-frame translation from the 13th codon. The

CrPV ORFx is predicted to be 41 amino acids in length if ORFx is translated from the

+1-frame 13th codon (Fig. 6). To determine whether CrPV ORFx is synthesized during

infection, *Drosophila* S2 cells infected with CrPV (MOI 10) were harvested at 6 hours

post infection and lysed. Proteins were subsequently digested with trypsin and peptides

were analyzed by LC-MS/MS. We identified two peptides that correspond to CrPV

ORFx both of which were located downstream of the +1-frame 13th codon (Fig. 6).

Importantly, these peptides were not identified in mock-infected S2 cells. Thus, ORFx is

expressed in CrPV-infected S2 cells.

Given this, we sought to determine the influence of CrPV ORFx expression on

the outcome of viral infection. Using a recently developed CrPV infectious clone, termed

CrPV-2, we introduced mutations that would abolish ORFx expression (36). To this end,

we created two separate mutant clones: the +1-frame 12th codon (UUG$_{6251-3}$) was

altered to an amber stop codon (UAG; CrPV-S12) and the +1-frame 19th codon

(UUA$_{6272-4}$) changed to an ochre stop codon (UAA; CrPV-S19) (Fig. 7A). Both mutations

are synonymous in the 0-frame. Based on our translation data (Fig. 2), the +1-frame

S19 but not the S12 would inhibit ORFx expression.

First, we examined whether ORFx influences viral protein synthesis *in vitro*.

Incubation of *in vitro* transcribed CrPV-2 RNA in Sf21 translation extracts led to

synthesis and processing of all viral proteins as reported previously (Fig. S4A) (36).

While stop codons in the 0-frame of ORF1 or ORF2 inhibited synthesis of viral proteins,

both CrPV-S12 and CrPV-S19 RNAs resulted in viral protein synthesis that was

indistinguishable compared to wild-type CrPV -2 *in vitro*, demonstrating that CrPV ORFx

is not necessary for viral protein synthesis *in vitro* (Fig. S4B) (36).

We next assessed the viability of the CrPV-S12 and -S19 viruses in cell culture.

Harvested wild-type, mutant S12 or S19 CrPV-2 virus were used to infect naïve S2 cells

at a MOI 10 and 1 to follow the first round of infection and subsequent rounds of

infection, respectively. Infection with wild-type CrPV-2, CrPV-S12 or CrPV-S19 all

resulted in accumulation of viral proteins and RNA and shutdown of host translation in a

similar manner (Fig. S4C). Similarly, neither infection produced significantly different

titres between wild type CrPV-2 and either CrPV-S12 or CrPV-S19 at any time point,

apart from both mutant viruses resulting in higher titres than CrPV-2 after 24 hours post

infection (Fig. S4D). A similar result was observed with infections at a MOI 0.1 (data not

shown). Taken together, ORFx has no observable effect on the life cycle of CrPV in S2

cells.

ORFx contributes to CrPV infection in adult flies and associates with membranes

We addressed whether ORFx contributes to CrPV infection in adult fruit flies. To

test this, we injected adult flies intrathoracically with PBS, CrPV-2, CrPV-S12, or CrPV-

S19 and monitored fly mortality daily. Flies injected with CrPV-2 or CrPV-S12 exhibited

50% mortality by day 5 and 100% mortality by day 11 and 12 (Fig. 7B). By contrast, flies

injected with CrPV-S19 did not reach 50% mortality until day 9 and reached 100%

mortality at day 14 (Fig. 7B). These results demonstrate that ORFx contributes to CrPV

pathogenesis in adult flies.

To determine if the effect seen on CrPV pathogenesis is a result of defects in

viral replication, we measured viral titres and assessed viral protein levels in injected

adult flies at 5 days post infection. Interestingly, viral titres and viral protein levels

showed no significant differences between wild type CrPV-2, CrPV-S12, and CrPV-S19

(Fig. S5). Finally, using RT-PCR followed by sequencing, the S12 and S19 mutations

are stable during virus propagation in S2 cells (data not shown). In summary, our results

indicate that the defect in viral pathogenesis observed in the CrPV-S19 mutant virus is

not due to a defect in viral replication.

From our data, CrPV ORFx is predicted to be a 41 amino acid protein, mediated

by a ribosomes bypass mechanism from the 13th +1-frame AAA codon. Comparing the

sequence of ORFx from CrPV and other species shows no appreciable homology to

other proteins. Nevertheless, *in silico* topology predictions suggest that ORFx can adopt

an alpha helical transmembrane segment at its C-terminus (amino acids 22-39; Fig.

S6A). To examine ORFx function, we generated constructs containing either N- or C-

terminal HA-tagged ORFx. Transfection for 10 or 24 hours with the HA-tagged ORFx

constructs did not result in a dramatic decrease in S2 cell viability as measured by a

trypan blue exclusion assay (Fig. S6B). However, by 48 hours, transfection of ORFx-HA

and HA-ORFx led to a relatively minor reduction (13% and 15% decrease, respectively)

in cell viability, suggesting that ORFx expression is slightly toxic in S2 cells.

Immunoblotting for HA showed that HA-ORFx is expressed in S2 cells (Fig. S6C). To

determine the subcellular localization in S2 cells, we transfected the HA-tagged ORFx

constructs and monitored ORFx localization (pORFx-HA) by HA-antibody

immunofluorescence staining in comparison with cytoplasmic, ER, Golgi, and nuclear

marker protein antibodies. We also mutated two pairs of amino acids, LV and LI, to KK

and KR respectively, to disrupt the transmembrane domain (pORFx-TMmut-HA). Co-

staining showed that the wild-type HA-tagged ORFx overlaps mainly with ER protein

marker Calnexin, and partially overlaps with Golgi-associated protein, Golgin84 (Fig. 8,

S7, pORFx-HA). The wild-type HA-ORFx displayed little to no overlap with α-Tubulin

staining. By contrast, the mutant transmembrane HA-tagged ORFx staining showed no

overlap with Calnexin or Golgin84 (Fig. 8, S7, pORFx-TmMut-HA). Collectively, these

results suggest that ORFx associates with membranous organelles and potentially the

ER specifically. To examine this further, we used differential centrifugation to separate

subcellular components followed by immunoblotting of HA. As expected, tubulin is found

in the cytoplasmic fraction and the ER-associated KDEL protein and cytochrome C are

enriched in nuclei and ER fractions (Fig. S6C). Both HA-tagged ORFx are detected

within membranous fractions but not within the cytoplasmic fractions (Fig. S6C), further

supporting that ORFx resides within the membrane of cells that may be important for its

function.

DISCUSSION

Recoding mechanisms have illuminated diverse RNA structural elements that

interact with the ribosome to affect reading frame maintenance (2). In this study, we

have demonstrated a novel translation recoding mechanism by which an IRES may

promote ribosome repositioning at a downstream codon. In contrast to the IAPV IGR

IRES, which directs ribosome reading frame selection (6), the CrPV IGR IRES can

facilitate the expression of a downstream +1 overlapping frame, which we termed ORFx.

We provide extensive analysis using mutagenesis that translation of CrPV ORFx likely

occurs 37 nucleotides downstream at the 13th AAA (Lys) codon. Moreover, we show

that ORFx is expressed in CrPV-infected cells by mass spectrometry and that ORFx is

required for promoting CrPV pathogenesis in a *Drosophila* injection model. Our data

suggest a model whereby after 80S assembly on the CrPV IGR IRES, the majority of

the ribosomes translate in the 0-frame by delivery of the incoming Ala-tRNA^{Ala} to the 0-

frame GCU codon whereas a fraction of ribosomes bypasses or "slides" 37 nucleotides

downstream to direct translation at the +1-frame AAA (Lys) codon (Fig. S8). From our

results, the following rules appear to apply to +1-frame translation in CrPV: i) the PKI

must be intact, ii) both 40S and 60S ribosomal subunits are required to bind to the IRES,

iii) pseudotranslocation of the IRES through the ribosome is necessary, iv) the spacer

region between PKI and the 13th AAA codon is essential, and v) the nucleotide identity

of the spacer region is crucial for efficient +1 frame translation.

A unique feature of this repositioning mechanism is that an intact IRES is

essential for +1-frame translation (Fig. 2). The current model is that the PKI domain of

the IRES occupies the ribosomal A site upon ribosome binding to the IRES, followed by

a pseudotranslocation event in order to vacate the A site to allow delivery of the first

aminoacyl-tRNA in the 0-frame (17). Given this model, it is difficult to envision how the

IRES can direct the ribosome to bypass 37 nucleotides to initiate translation at a +1-

frame AAA Lys codon. A potential clue comes from our mutagenesis analysis that

suggests that translocation of the CrPV IRES through the ribosome is a prerequisite for

downstream +1-frame translation (Fig. 4). This is intuitive as the PKI domain must

vacate the A site in order to allow translation in both the 0 and +1 frames. At this point, it

is unclear whether the CrPV IGR IRES ribosome repositioning event requires delivery of

an aminoacyl-tRNA to the A site of the ribosome prior to the ribosome sliding

downstream to initiate translation at the 13th +1 frame codon or whether a vacant

ribosome bypasses from the IRES to the initiating +1 frame codon. Determination of the

aminoacyl-tRNA(s) that are delivered during the initial CrPV IGR IRES

pseudotranslocation event should provide insights whether the ribosomal A site is

vacant prior to ribosome sliding downstream to the 13th +1 frame codon.

A recent *in vitro* study using single molecule fluorescence spectroscopy

demonstrated that the CrPV IGR IRES can facilitate +1-frame translation approximately

5% of the time compared to 0-frame translation whereby reading frame selection is

dictated by the kinetics of tRNA binding in the first 0- or +1-frame codon (12). Using a

more physiologically-related system, our results suggest that the CrPV IRES directs +1-

frame translation using a different mechanism. Based on our systematic stop codon

mutational analyses, the most likely scenario is that ribosomes recruited to the IRES

must bypass downstream to translate in the +1-frame (Fig. 2). In support of this,

insertion of a stop codon at the 12th +1-frame codon, but not the 19th, attenuates CrPV-

mediated death in a Drosophila injection model, thus providing biological significance of

CrPV +1-frame translation (Fig. 7). We also note that the downstream authentic 'spacer'

sequence is necessary for +1-frame translation (Fig. 5), which is partially absent and/or

altered from previous studies, which may preclude detection of bypass (12).

The downstream spacer region appears to be a key feature that is necessary for

efficient CrPV IRES-mediated +1-frame synthesis (Fig. 5). Interestingly, the majority of

mutations within the spacer region causes a reduction ranging from 10-80% in the

amount of CrPV IRES-mediated +1-frame synthesis (Fig. 2), suggesting that nucleotide

or codon identity is crucial for triggering the bypass event. There are no obvious RNA

secondary structures within the spacer region, however, we cannot rule out any long

distance RNA:RNA interactions that may contribute to +1-frame translation.

Furthermore, sequences immediately downstream of the IRES PKI domain are

important for +1 frame translation (Fig. 5B). Investigation into how the spacer sequence

influences ribosomes bound to the IRES to start translation in the 0-frame or at the

downstream +1-frame initiating codon should shed light onto this atypical ribosome

bypass mechanism.

How does the ribosome reposition to the downstream +1-frame initiation codon

after IRES binding? It is possible that the repositioning of the ribosomes occurs via

mechanism similar to that observed with prokaryotic 70S 'scanning' (37) or 70S 'sliding'

that occurs in coupled translational reinitiation (38, 39) (Fig. S8). Indeed, a related

dicistrovirus PSIV IRES can direct translation using a scanning-like mechanism in

prokaryotes, which may suggest a similar property observed with our studies on the

CrPV IRES (40). Moreover, it has been reported and proposed that energy-independent

scanning or diffusion of the ribosome or ribosomal subunits (i.e. phaseless wandering)

can occur to locate an AUG codon (39, 41-43)(Fig. S8). Nevertheless, our study shows

that CrPV IRES +1-frame translation requires 80S ribosome binding to the IRES (Fig.

3A) and is edeine-insensitive (Fig. 3B), thus we favour a model that 80S ribosomes

reposition to the 13th +1-frame codon. This warrants comparison with the translational

bypassing observed in *gene 60* of T4 bacteriophage (44). In *gene 60*, translating

ribosomes stall in a non-canonical rotated state at a 'take-off' Gly codon with a peptidyl-

tRNA[Gly], which dissociates from the anticodon, and 'lands' at a matching Gly codon 50

nucleotides downstream and allowing translation to resume. This bypass mechanism is

dependent on a post-translocation step requiring a downstream 5' 'take-off' stem-loop

structure and a nascent translated peptide (45, 46). Although CrPV does not have an

obvious RNA structure within the spacer region, it is possible that the highly complex

structure of the IGR IRES itself may contribute to downstream +1 frame translation,

especially given its dynamic nature during movement through the ribosome (Fig. S8)(28,

47). Moreover, it is known that the ribosome bound to the IGR IRES is in a rotated state

and that the first pseudotranslocation step is rate limiting, which may contribute to CrPV

IRES reading frame selection (17, 29). Finally, it is possible is that CrPV IRES bypass

could be occurring through an RNA looping event (48); the downstream RNA is brought

into close proximity with the 80S ribosome allowing it to transition to 13[th] AAA codon

potentially by an unknown protein factor or a long-range RNA:RNA interaction. Taken

together, it is likely that it is a combination of tRNA kinetics and conformational changes

of the IRES and the downstream spacer region that lead to bypassing, of which the

contributions of each element require further investigation.

The biological relevance of CrPV +1-frame translation was initially evidenced by

the detection of ORFx peptides in CrPV-infected S2 cells (Fig. 6). To our surprise,

disruption of ORFx synthesis by stop codon insertion in the +1-frame did not perturb

viral infection in tissue culture cells but showed retarded mortality in adult flies even

though viral load remained similar between wild type and mutant viruses (Fig. 7). CrPV

infection is thought to cause death through paralysis, subsequently leading to

dehydration or starvation of the host (49, 50). CrPV can infect several tissues in the fly

including the trachea, midgut, and central nervous system although the latter has not

been demonstrated directly (48, 50, 51). How ORFx may contribute to CrPV

pathogenesis is an outstanding question. Our results indicate that ORFx is membrane

associated (Fig. 8, S7) and does not contribute directly to viral replication but rather to

the pathogenesis of CrPV infection in fruit flies (Fig. 7, S4). Furthermore, expression of

ORFx is slightly cytotoxic in Drosophila cells, a property that may also contribute to

pathogenesis of CrPV infection. Future studies into the localization of ORFx, potential

interacting partners, and tissue tropism of wild-type versus mutant virus infection in the

fly should provide insights into the role of ORFx.

Viruses continue to surprise us with their ability to manipulate the ribosome in

remarkable ways. Here, in addition to the previous findings on the honey bee

dicistrovirus IRES, we have revealed another recoding mechanism utilizing an IRES,

thus highlighting the strong selection to increase the coding capacity of the dicistrovirus

genome. Furthermore, an IRES that can direct ribosome repositioning to facilitate the

translation of a hidden +1 overlapping ORFx adds to the growing list of diverse

pathways of ribosome translational recoding. It will be of considerable interest to

investigate whether other IRESs direct reading frame selection by a similar CrPV IGR

IRES-mediated ribosome repositioning mechanism. Ribosome repositioning or bypass

is not specific to bacteria or mitochondria (52, 53) but may be a more general

phenomenon in eukaryotes that initially thought.

MATERIALS AND METHODS

Cell culture and virus. *Drosophila* Schneider line 2 (S2; Invitrogen) cells were maintained and passaged in Shield's and Sang medium (Sigma) supplemented with 10% fetal bovine serum.

Propagation of CrPV in *Drosophila* S2 cells has been previously described (54). CrPV-2 and mutant viruses were generated from *Drosophila* S2 cells using an adapted protocol (55). Briefly, 5.0×10^7 S2 cells were transfected with *in vitro* transcribed RNA derived from pCrPV-3 or mutant plasmids and incubated for 48h. Cells were dislodged into the media, treated with 0.5% Igepal CA-630 (Nonidet P-40) and 0.1% 2-mercaptoethanol, and incubated on ice for 10 min. Cell debris was cleared by centrifugation at 13,800 RCF for 15 min at 4°C. Viral particles were then concentrated by ultracentrifugation at 141,000 RCF for 2.5 h at 4°C. The pellet was resuspended in PBS and sterilized through a 0.2 µM filter. Viral titres and yield were determined as previously described (20). All viruses were sequence verified via RT-PCR with primers directed against the CrPV IGR IRES.

LC-MS/MS Analysis. Cell pellets harvested from CrPV-infected cells at 6 hpi were solubilized in 1% sodium deoxy cholate and 50 mM NH_4HCO_3. Protein concentrations were determined via BCA assay (Thermo). Proteins (100 µg) were reduced (2 µg DTT, 37°C, 30 min) and alkylated (5 µg iodoacetamide, RT, 20 min). Samples were digested with trypsin overnight at room temperature. Peptides were acidified with 1% TFA to pH <2.5 and the precipitated deoxycholate was remove via centrifugation. Peptides were desalted and concentrated on C18 STAGE-tips, eluted in 80% acetonitrile, 0.5% acetic acid, and dried in a vacuum concentrator (Eppendorf)(Rappsilber, Ishihama, & Mann,

2003). Samples were resuspended in 20% acetonitrile and 0.1% formic acid before

loading on an Agilent 6550 mass spectrometer.

Data were searched using MaxQuant (v1.5.3.30)(56). Parameters included:

carbamidomethylated cysteine (fixed), methionine oxidation (variable), glutamine and

asparagine deamidation (variable), and protein N-terminal acetylation (variable); trypsin

specific; maximum 2 missed cleavages; 10 ppm precursor mass tolerance; 0.05 Da

fragment mass tolerance; 1% FDR; +1 to +7 charge states; and common contaminants

were included. Both the *Drosophila* and CrPV protein databases used were the most

recent annotations downloaded from UniProt (www.uniprot.org).

Fly stocks and viral injections. Flies (*Isogenic w^{1118}*; Bloomington Drosophila Stock

Center) were maintained on standard cornmeal food at 25°C and 70% humidity with a

12 h light-dark cycle. Freshly eclosed virgin males and females were separated and

collected in groups of 10 each. Flies (10 males and 10 females) were injected with 200

nL of PBS, CrPV-2, CrPV-S12, or CrPV-S19 (5000 FFU) using a PV830 PicoPump

(World Precision Instruments) and transferred to standard food. Mortality was monitored

daily.

ACKNOWLEDGEMENTS

We thank John Atkins and the Jan lab for insightful discussions. Kathy Pan and

Seonghoon Lee contributed a subset of data in Fig. 1A and 2B, respectively. This study

was supported by a Canadian Institutes of Health Research (PJT-148761) to E.J. and to

D.A. (MOP-130517), an NSERC Discovery Grant to L.F. and an NSERC CGS-D

fellowship to C.K.

REFERENCES

1. Baranov PV, Atkins JF, & Yordanova MM (2015) Augmented genetic decoding: global, local and temporal alterations of decoding processes and codon meaning. *Nature reviews. Genetics* 16(9):517-529.
2. Dinman JD (2012) Mechanisms and implications of programmed translational frameshifting. *Wiley interdisciplinary reviews. RNA* 3(5):661-673.
3. Firth AE & Brierley I (2012) Non-canonical translation in RNA viruses. *J Gen Virol* 93(Pt 7):1385-1409.
4. Jan E, Mohr I, & Walsh D (2016) A Cap-to-Tail Guide to mRNA Translation Strategies in Virus-Infected Cells. *Annual review of virology* 3(1):283-307.
5. Au HH & Jan E (2014) Novel viral translation strategies. *Wiley interdisciplinary reviews. RNA* 5(6):779-801.
6. Ren Q, *et al.* (2012) Alternative reading frame selection mediated by a tRNA-like domain of an internal ribosome entry site. *Proc Natl Acad Sci U S A* 109(11):E630-639.
7. Jackson RJ, Hellen CU, & Pestova TV (2010) The mechanism of eukaryotic translation initiation and principles of its regulation. *Nature reviews. Molecular cell biology* 11(2):113-127.
8. Wilson JE, Pestova TV, Hellen CU, & Sarnow P (2000) Initiation of protein synthesis from the A site of the ribosome. *Cell* 102(4):511-520.
9. Jan E, Kinzy TG, & Sarnow P (2003) Divergent tRNA-like element supports initiation, elongation, and termination of protein biosynthesis. *Proc Natl Acad Sci U S A* 100(26):15410-15415.
10. Jan E & Sarnow P (2002) Factorless ribosome assembly on the internal ribosome entry site of cricket paralysis virus. *J Mol Biol* 324(5):889-902.
11. Pestova TV & Hellen CU (2003) Translation elongation after assembly of ribosomes on the Cricket paralysis virus internal ribosomal entry site without initiation factors or initiator tRNA. *Genes Dev* 17(2):181-186.
12. Petrov A, Grosely R, Chen J, O'Leary SE, & Puglisi JD (2016) Multiple Parallel Pathways of Translation Initiation on the CrPV IRES. *Mol Cell* 62(1):92-103.
13. Nishiyama T, *et al.* (2003) Structural elements in the internal ribosome entry site of Plautia stali intestine virus responsible for binding with ribosomes. *Nucleic Acids Res* 31(9):2434-2442.
14. Costantino DA, Pfingsten JS, Rambo RP, & Kieft JS (2008) tRNA-mRNA mimicry drives translation initiation from a viral IRES. *Nat Struct Mol Biol* 15(1):57-64.
15. Pfingsten JS, Costantino DA, & Kieft JS (2006) Structural basis for ribosome recruitment and manipulation by a viral IRES RNA. *Science* 314(5804):1450-1454.
16. Schuler M, *et al.* (2006) Structure of the ribosome-bound cricket paralysis virus IRES RNA. *Nat Struct Mol Biol* 13(12):1092-1096.
17. Fernandez IS, Bai XC, Murshudov G, Scheres SH, & Ramakrishnan V (2014) Initiation of Translation by Cricket Paralysis Virus IRES Requires Its Translocation in the Ribosome. *Cell*.
18. Au HH, *et al.* (2015) Global shape mimicry of tRNA within a viral internal ribosome entry site mediates translational reading frame selection. *Proc Natl Acad Sci U S A* 112(47):E6446-6455.
19. Muhs M, *et al.* (2015) Cryo-EM of ribosomal 80S complexes with termination factors reveals the translocated cricket paralysis virus IRES. *Mol Cell* 57(3):422-432.
20. Kerr CH, Ma ZW, Jang CJ, Thompson SR, & Jan E (2016) Molecular analysis of the factorless internal ribosome entry site in Cricket Paralysis virus infection. *Scientific reports* 6:37319.

21. Nakashima N & Uchiumi T (2009) Functional analysis of structural motifs in
dicistroviruses. *Virus Res* 139(2):137-147.
22. Jan E (2006) Divergent IRES elements in invertebrates. *Virus Res* 119 (1):16-28.

23. Jang CJ, Lo MC, & Jan E (2009) Conserved element of the dicistrovirus IGR IRES that
mimics an E-site tRNA/ribosome interaction mediates multiple functions. *J Mol Biol*
387(1):42-58.
24. Jang CJ & Jan E (2010) Modular domains of the Dicistroviridae intergenic internal
ribosome entry site. *RNA* 16(6):1182-1195.
25. Hatakeyama Y, Shibuya N, Nishiyama T, & Nakashima N (2004) Structural variant of the
intergenic internal ribosome entry site elements in dicistroviruses and computational
search for their counterparts. *RNA* 10(5):779-786.
26. Cevallos RC & Sarnow P (2005) Factor-independent assembly of elongation-competent
ribosomes by an internal ribosome entry site located in an RNA virus that infects
penaeid shrimp. *J Virol* 79(2):677-683.
27. Hertz MI & Thompson SR (2011) In vivo functional analysis of the Dicistroviridae
intergenic region internal ribosome entry sites. *Nucleic Acids Res* 39(16):7276-7288.
28. Abeyrathne PD, Koh CS, Grant T, Grigorieff N, & Korostelev AA (2016) Ensemble cryo-
EM uncovers inchworm-like translocation of a viral IRES through the ribosome. *eLife* 5.
29. Zhang H, Ng MY, Chen Y, & Cooperman BS (2016) Kinetics of initiating polypeptide
elongation in an IRES-dependent system. *eLife* 5.
30. Firth AE, Wang QS, Jan E, & Atkins JF (2009) Bioinformatic evidence for a stem-loop
structure 5'-adjacent to the IGR-IRES and for an overlapping gene in the bee paralysis
dicistroviruses. *Virol J* 6:193.
31. Ren Q, Au HH, Wang QS, Lee S, & Jan E (2014) Structural determinants of an internal
ribosome entry site that direct translational reading frame selection. *Nucleic Acids Res*
42(14):9366-9382.
32. Wang QS, Au HH, & Jan E (2013) Methods for studying IRES-mediated translation of
positive-strand RNA viruses. *Methods* 59(2):167-179.
33. Dinos G*, et al.* (2004) Dissecting the ribosomal inhibition mechanisms of edeine and
pactamycin: the universally conserved residues G693 and C795 regulate P-site RNA
binding. *Mol Cell* 13(1):113-124.
34. Kozak M & Shatkin AJ (1978) Migration of 40 S ribosomal subunits on messenger RNA
in the presence of edeine. *J Biol Chem* 253(18):6568-6577.
35. Ruehle MD*, et al.* (2015) A dynamic RNA loop in an IRES affects multiple steps of
elongation factor-mediated translation initiation. *eLife* 4.
36. Kerr CH*, et al.* (2015) The 5' untranslated region of a novel infectious molecular clone of
the dicistrovirus cricket paralysis virus modulates infection. *J Virol* 89(11):5919-5934.
37. Yamamoto H*, et al.* (2016) 70S-scanning initiation is a novel and frequent initiation mode
of ribosomal translation in bacteria. *Proc Natl Acad Sci U S A* 113(9):E1180-1189.
38. Nomura M, Gourse R, & Baughman G (1984) Regulation of the synthesis of ribosomes
and ribosomal components. *Annu Rev Biochem* 53:75-117.
39. Adhin MR & van Duin J (1990) Scanning model for translational reinitiation in eubacteria.
J Mol Biol 213(4):811-818.
40. Colussi TM*, et al.* (2015) Initiation of translation in bacteria by a structured eukaryotic
IRES RNA. *Nature* 519(7541):110-113.
41. Sarabhai A & Brenner S (1967) A mutant which reinitiates the polypeptide chain after
chain termination. *J Mol Biol* 27(1):145-162.
42. Skabkin MA, Skabkina OV, Hellen CU, & Pestova TV (2013) Reinitiation and other
unconventional posttermination events during eukaryotic translation. *Mol Cell* 51(2):249-
264.

43. Abaeva IS, Pestova TV, & Hellen CU (2016) Attachment of ribosomal complexes and
retrograde scanning during initiation on the Halastavi arva virus IRES. *Nucleic Acids Res*
44(5):2362-2377.
44. Wills NM*, et al.* (2008) Translational bypassing without peptidyl-tRNA anticodon
scanning of coding gap mRNA. *EMBO J* 27(19):2533-2544.
45. Chen J*, et al.* (2015) Coupling of mRNA Structure Rearrangement to Ribosome
Movement during Bypassing of Non-coding Regions. *Cell* 163(5):1267-1280.
46. Samatova E, Konevega AL, Wills NM, Atkins JF, & Rodnina MV (2014) High-efficiency
translational bypassing of non-coding nucleotides specified by mRNA structure and
nascent peptide. *Nature communications* 5:4459.
47. Pisareva VP, Pisarev AV, & Fernandez IS (2018) Dual tRNA mimicry in the Cricket
Paralysis Virus IRES uncovers an unexpected similarity with the Hepatitis C Virus IRES.
eLife 7.
48. Paek KY*, et al.* (2015) Translation initiation mediated by RNA looping. *Proc Natl Acad
Sci U S A* 112(4):1041-1046.
49. Reinganum C (1975) The isolation of cricket paralysis virus from the emperor gum moth,
Antheraea eucalypti Scott, and its infectivity towards a range of insect species.
Intervirology 5(1-2):97-102.
50. Chtarbanova S*, et al.* (2014) Drosophila C virus systemic infection leads to intestinal
obstruction. *J Virol* 88(24):14057-14069.
51. Bonning BC & Miller WA (2010) Dicistroviruses. *Annu Rev Entomol* 55:129-150.
52. Lang BF*, et al.* (2014) Massive programmed translational jumping in mitochondria. *Proc
Natl Acad Sci U S A* 111(16):5926-5931.
53. Herr AJ, Gesteland RF, & Atkins JF (2000) One protein from two open reading frames:
mechanism of a 50 nt translational bypass. *EMBO J* 19(11):2671-2680.
54. Garrey JL, Lee YY, Au HH, Bushell M, & Jan E (2010) Host and viral translational
mechanisms during cricket paralysis virus infection. *J Virol* 84(2):1124-1138.
55. Krishna NK, Marshall D, & Schneemann A (2003) Analysis of RNA packaging in wild-
type and mosaic protein capsids of flock house virus using recombinant baculovirus
vectors. *Virology* 305(1):10-24.
56. Cox J & Mann M (2008) MaxQuant enables high peptide identification rates,
individualized p.p.b.-range mass accuracies and proteome-wide protein quantification.
Nature biotechnology 26(12):1367-1372.

683
684

    FIGURE LEGENDS

Fig. 1. Biochemical analysis of potential +1-frame translation by Dicistrovirus IGR

IRESs. A) IGR IRES mediated +1-frame translation using an *in vitro* translation

assay. The translation of firefly luciferase (FLuc), which is fused in the 0-frame or +1-

frame, is driven by the individual IGR IRES within the bicistronic reporter construct.

Linearized reporter constructs were incubated in Sf21 *in vitro* translation insect cell

extract at 30 °C for 2 hours in the presence of [^{35}S] methionine/cysteine. Translation of

FLuc and RLuc was monitored by phosphorimager analysis after resolving on a 16%

SDS PAGE. On the bottom panel, the ratio of FLuc/RLuc are quantified and normalized

to the IAPV 0-frame translation. Shown are averages from at least three independent

biological experiments (± SD). **B) The secondary structure of CrPV IGR IRES.** The

ΔPKI, ΔPKII, ΔPKIII, L1.1A, and L1.1B mutants are shown. Comp denotes

compensatory mutations combining ΔPKI CC/GG and ΔPKI GG/CC. DM - double

mutation of ΔPKI and ΔPKII. TM - triple mutation of ΔPKI, ΔPKII, and ΔPKIII. The

potential U6186/G6617 base pair to direct +1 frame translation from a CUA leucine

codon.

Fig. 2. Translation of +1 frame CrPV ORFx is IGR IRES-dependent and initiates

downstream of the PKI region. A) Schematic of mutations introduced downstream

of the IGR IRES. The PKI region and downstream 'spacer' sequence of CrPV IGR

IRES are shown. A series of mutations and stop codons were introduced either on the

0-frame or +1-frame. **B) Analysis of the IGR IRES translation *in vitro*.** Linearized

reporter constructs are incubated in Sf21 translation extract at 30°C for 2 hours in the

present of [^{35}S]-methionine/cysteine. Translation of FLuc and RLuc was monitored by

autoradiography after resolving on a 16% SDS PAGE. A representative gel from at least

three independent biological experiments is shown. * = p-value < 0.05. **C) Quantitation**

of IRES-mediated translation. Translation of FLuc and RLuc monitoring FLuc/RLuc

enzymatic activity were quantified and normalized to wild-type (WT) IAPV IRES. RLuc

monitors scanning-dependent translation acting as an internal control. RLA- relative

light units. Shown are averages from at least three independent biological experiments

(± SD). * = p-value < 0.05.

Fig. 3. Translation of CrPV ORFx is dependent on 80S binding and scanning-

independent. A) Effect of L1.1 mutation on +1 frame translation. Comparison of

luciferase activities between WT IGR IRES and L1.1 mutants (L1.1A and L1.1B) that do

not bind the 60S subunit. The value of RLuc and FLuc are normalized to WT. **B) +1-**

frame translation is relatively insensitive to edeine. CrPV IRES-mediated 0- and +1-

frame translation is monitored in the absence or presence of 2 µM edeine. The ΔPKI

mutants in both 0- and +1-frame are used as controls for IGR IRES dependent

translation. The luciferase activities of RLuc and FLuc on the left panel are normalized

to the WT of the 0-frame, whereas the luciferase activities of RLuc and FLuc on the right

panel are normalized to the WT of the +1-frame. Shown are averages from at least

three independent biological experiments (± SD).

Fig. 4. Pseudotranslocation is necessary for +1-frame translation. Constructs

containing mutations in the variable loop region (VLR) of the IGR IRES known to disrupt

pseudotranslocation were monitored for translation activity. Luciferase activities were

compared between WT IGR IRES and VLR mutants after *in vitro* translation assays. "x"

denotes deleted nucleotides within VLR. Shown are averages from at least three

independent biological experiments (± SD).

Fig. 5. The downstream sequence of CrPV IGR IRES promotes +1-frame

translation. A) Chimeric IRES displays +1-frame translation activity. The PSIV IGR

IRES, which shows no +1-frame translation activity (Fig. 1 and Fig. 3A), is combined

with the downstream sequence of CrPV IGR IRES PKI region. The luciferase activities

are normalized to the wild-type CrPV constructs in both +1- and 0-frame. The PSIV-

CrPV chimera in the 0-frame is used as control to demonstrate that PSIV IGR IRES

functions under this context. **B) Truncation of the spacer region impairs +1-frame**

translation from the CrPV IGR IRES. Schematic displaying truncations within the

spacer region between the PKI domain and the +1 frame start site is shown above. *In*

vitro translation assay of the CrPV IRES bicistronic construct with the respective

mutations were monitored by luciferase activities. Both RLuc and FLuc activities are

normalized to that of WT constructs. Results are shown as a normalized ratio of

FLuc/RLuc activity (left) and as separated RLuc and Fluc activities (right) Shown are

averages from at least three independent biological experiments (± SD). RLA = relative

luciferase activity

Fig. 6. ORFx is expressed in CrPV infected S2 cells. The predicted +1 frame amino

acid sequence is shown above. Residues that are italicized represent the amino acid

sequence if initiation occurred adjacent to the IGR IRES, whereas the bolded K lysine

residue denotes the position of the 13th amino acid. *Drosophila* S2 cells were infected

with CrPV (MOI 10). Proteins were extracted, digested with trypsin and subjected to LC-

MS/MS analysis. Peptide fragment spectra were searched against a *Drosophila* uniprot

database plus CrPV proteins. Two peptides (highlighted) were detected from the trypsin

digestion of S2 cell lysate at 6 hours post infection. Individual fragment ions are

annotated in the spectra and in the sequence representation.

Fig. 7. Pathogenesis of a CrPV mutant lacking ORFx is attenuated in adult

***Drosophila melanogaster* flies. A) Schematic of mutations in the CrPV infectious**

clone, CrPV-2. The downstream nucleotide sequence of CrPV-2 IGR IRES and its

potential amino acid sequence of +1-frame ORFx are shown. The ORF2-Stop mutant

depletes the synthesis of 0-frame viral structural proteins. Mutants S12 and S19 place a

stop codon on the +1-frame with a synonymous mutation in the 0-frame. Note that

CrPV-2 (Accession: KP974706) sequence has a $C_{6279}A$ mutation (denoted with a black

arrowhead) compared to the original CrPV (Accession: AF218039) sequence. Residues

that are italicized, underlined and in bold are the peptides identified by mass

spectrometry in Fig. 6. **B)** Adult flies (Iso w^{1118}; 10 male, 10 female) were injected

intrathoracically with 5000 FFU of CrPV-2, CrPV-S12, CrPV-S19, or PBS. Subsequently,

flies were flipped onto standard media and survival was monitored daily. Shown is a

graph representing the average from three separate biological experiments (±SEM).

Fig. 8. Subcellular localization of CrPV ORFx. S2 cells were transfected with a

construct expressing C-terminally HA-tagged ORFx or a mutant version of ORFx in the

transmembrane region and incubated for 48 h. Following incubation, cells were fixed,

permeabilized and co-stained with HA antibody and antibodies against the ER (Calnexin

99A) and cytoplasm (alpha-Tubulin). Shown are representative Z-stack micrographs

from three independent experiments. Scale bars represent 15 μm.

A Kerr_Figure 1

B Cricket paralysis virus IGR IRES

Kerr_Figure 2

A Kerr_Figure 3

B

Kerr_Figure 4

Kerr_Figure 5

A

CrPV IRES | PSIV IRES | Chimera IRES

0 frame +1 frame | 0 frame No +1 frame | 0 frame +1 frame?

(+1) frame: CrPV, ΔPKI, PSIV, PSIV/CrPV — RLuc / FLuc

(0) frame: CrPV, ΔPKI, PSIV, PSIV/CrPV — RLuc / FLuc

B

6185

+1-frame

a u g g a
u a c c u
6212

0-frame

g cua cau uuc aag aua aac aag aaa auu cac aca uug aaaau g... ORF2

Δ30 Δ27 Δ21 Δ18 Δ15 Δ6 Δ3

Normalized Fluc/Rluc Activity: WT, ΔPKI, Δ3, Δ6, Δ15, Δ18, Δ21, Δ27, Δ30

Normalized RLA: WT, ΔPKI, Δ3, Δ6, Δ15, Δ18, Δ21, Δ27, Δ30 — RLuc / FLuc

Kerr_Figure 6

Predicted Sequence: N - **K**MKIKDLCQNRKK **LYILLAK ELPLVLLR** SLISLIFQLIIW - **C**

A

```
                              0 frame
                              ORF2-stop              S12 (+1 frame)
                                 uaa                    uag        14
                                                                   •
(+1) frame     L H F K I N K K I H T L K M K I K D
 (0) frame     A T F Q D K Q E N S H I E N E D K R
           UACCU gcuacauuucaagauaaacaagaaaauucacacauugaaaaugaagauaaaaga
           PKI   6218           6232              6251
                   •              •                •
```

S19 (+1 frame)
```
              uaa                                       34
                                                        •
(+1) frame    L C Q N R K K  *L Y I L L A K*  *E L P L V L*
 (0) frame    L M S E Q K E I V H F V S E G V T P S T
            cuuaugucagaacagaaagaaauuguacauuuuguuagcgaaggaguuaccccuaguacu
            6272           ↑                           6317
              •                                          •
```

```
                              51
                              •
(+1) frame    *L R* S L I S L I F Q L I I W T
 (0) frame    T A L P D I V N L S T N Y L D M
            acugcgcucccugauaucguuaaucuuucaacuaauuaucuggacauga
            6332                      6368        6379
              •                        •           •
```

B

% Survival vs Day

● PBS   ○ CrPV-2   △ CrPV-S12   □ CrPV-S19

AlexFluor-468 α-HA Hoechst Merge

Calnexin99a

Tubulin

pORFx-HA

Calnexin99a

Tubulin

pORFx-TmMut-HA