Determining protein structures using genetics

Jörn M. Schmiedel[1,2], Ben Lehner[1-3*]

[1] Systems Biology Program, Centre for Genomic Regulation (CRG), The Barcelona Institute of

Science and Technology, Dr. Aiguader 88, Barcelona 08003, Spain

[2] Universitat Pompeu Fabra (UPF), Barcelona 08003, Spain

[3] Institució Catalana de Recerca i Estudis Avançats (ICREA), Pg. Lluís Companys 23, 08010

Barcelona, Spain

*Corresponding Author and Lead Contact: ben.lehner@crg.eu

Summary

Determining the three dimensional structures of macromolecules is a major goal of biological

research because of the close relationship between structure and function. Structure

determination usually relies on physical techniques including x-ray crystallography, NMR

spectroscopy and cryo-electron microscopy. Here we present a method that allows the high-

resolution three-dimensional structure of a biological macromolecule to be determined only from

measurements of the activity of mutant variants of the molecule. This genetic approach to

structure determination relies on the quantification of genetic interactions (epistasis) between

mutations and should be widely applicable, including to macromolecules whose structures are

difficult to determine by physical techniques.

Introduction

Mutations within a protein or RNA can have non-independent effects on folding, stability and activity (Fowler and Fields, 2014; Lehner, 2011; Starr and Thornton, 2016; Tokuriki and Tawfik, 2009). Consistently, quantifying the evolutionary couplings between amino acid and nucleotide occurrences in protein and RNA families yields structural and functional information tha has been used to predict macromolecular structures, physical interactions and mutational effect sizes (Altschuh et al., 1987; Burger and van Nimwegen, 2010; Dunn et al., 2008; Ekeberg et al., 2013; Gloor et al., 2005; Göbel et al., 1994; Halabi et al., 2009; Hopf et al., 2012; Hopf et al., 2017; Jones et al., 2012; Kamisetty et al., 2013; Lockless and Ranganathan, 1999; Marks et al., 2011; McLaughlin Jr et al., 2012; Michel et al., 2014; Morcos et al., 2011; Ovchinnikov et al., 2015; Ovchinnikov et al., 2017; Seemayer et al., 2014; Tóth-Petróczy et al., 2016; Weigt et al., 2009; Weinreb et al., 2016). Current state of the art structure prediction algorithms routinely incorporate evolutionary sequence information to predict tertiary protein contacts (Schaarschmidt et al., 2018). However, the success of this strategy depends on the depth and diversity of sequence alignments, with the approach failing for many proteins because of insufficient number of homologs (Marks et al., 2012; Ovchinnikov et al., 2017; Tóth-Petróczy et al., 2016).

An alternative approach to structure prediction would be to perform experiments that directly measure the dependencies between mutations in a molecule (Fowler and Fields, 2014) (Figure 1A). Such an approach, if successful, would provide a general strategy to determine macromolecular structures, including for molecules without sufficient numbers of homologs in sequence databases and for molecules whose structures are difficult to determine by physical techniques. All that would be required would be a selection assay that allows the molecular activity of a sufficiently large number of individual mutations and combinations of mutations to be quantified. However, whether such an approach is actually feasible is unknown.

Here we show that deep mutagenesis of a molecule can indeed provide sufficient information to determine its macromolecular structure. Our method quantifies how often residue pairs interact epistatically – that is non-independently – in experimental data and how such epistatic interaction patterns correlate. These data accurately identify individual tertiary structure contacts as well as secondary structure elements and their interactions in a protein domain. Moreover, we show that these data can be used to predict the structure of the protein domain at 1.9Å root mean squared deviation (RMSD) compared to a reference crystal structure.

Results

Epistatic variants within a protein domain are enriched for structural proximity

We used deep mutational scanning data for the immunoglobulin-binding protein G domain B1 (Olson et al., 2014) to investigate how genetic – or epistatic – interactions between mutations in a protein relate to structure. This dataset is the most complete double mutant deep mutagenesis of a protein domain reported to date and was generated by replacing each of 55 residues of the wild-type domain with all 19 alternative amino acids both individually and as double mutant pairwise combinations (Figure 1B, step 1). The resulting library consists of more than half a million variants (55*19 = 1,045 single mutants plus 55*54/2*19*19 = 536,085 double mutants). mRNA display was then used to combine an in vitro immunoglobulin G binding assay with a sequencing readout of protein fitness (here defined as propensity to bind immunoglobulin, Figure 1B, step 2). The protein fitness of each mutant was calculated as the change in read frequencies in the library due to selection relative to the wild-type variant (Figure 1B, steps 3 & 4). The assay provides a measurement range of more than two orders of magnitude, before background effects dominate the measured fitness values (Figure S1A), and a median error relative to the measurement range of 3.2%, as judged from count-based error estimates (see Methods).

We first computed which double mutant variants show epistatic fitness effects, i.e. non-independent fitness effects of the constituting single mutant variants (Figure 1). We applied a non-parametric null model - the running median of double mutant fitness values given the constituting single mutant fitness values - for the independence of mutations (Figure 2A). We then similarly calculated 5th and 95th percentile fitness surfaces; and classified double mutants as positive epistatic if they were fitter than the 95th percentile fitness surface and as negative epistatic if they were less fit than the 5th percentile fitness surface (Figure 2A). Basing the assessment of epistasis on this non-parametric null model accounts for nonlinearities introduced by the finite measurement range of the fitness assay, variable magnitudes of measurement error across the fitness landscape (Figure S1B), as well as non-specific epistatic behavior resulting from, for example, thermodynamic stability threshold effects (Diss and Lehner, 2018;

Tokuriki and Tawfik, 2009). We restricted the evaluation of positive or negative epistasis to
specific subsets of the data, where measurement errors do not impede epistasis classification
(Figure S1D, see Methods). Most importantly, we classified negative epistasis only for double
mutants whose expected fitness is significantly above the lower measurement limit of the fitness
assay. As a result of these restrictions as well as differences in initial coverage, the number of
double mutant variants that can be used to assess positive and negative epistasis varies
substantially across position pairs (with an average of 252 and 187 variants per position pair in
the positive and negative epistasis subsets, respectively; Figure S2). Consistent with our initial
hypothesis and previous observations (Melamed et al., 2013; Olson et al., 2014; Sahoo et al.,
2015), both positive and negative epistatic double mutants are enriched for proximal variants;
and this is not due to the specific subsets of the data being analyzed (Figure 2B).

Figure 1

A working hypothesis

B deep mutational scanning experiment

1) mutate protein of interest

2) competitive selection assay

3) genotype frequencies from sequencing

computational analysis

4) calculate fitness F_i and error $\sigma_{F,i}$
   from sequencing read counts

<div>
Dataset: Olson <i>et al.</i> 2014
protein G B1 domain, 56 aa
NNS mutagenesis
selection assay: <i>in vitro</i> IgG binding
100% possible single mutants
99.97% possible double mutants
(after QC: 78% / 50% for pos. / neg.
epistasis quantification)
</div>

5) 1000x re-sampling procedure
5a) sample variant fitness from $f_i \sim N(F_i, \sigma_i)$ & classify epistasis (see panel C)

6) compute **interaction scores** by
   merging positive & negative epistasis information

Figure 1: Extracting epistatic mutational effects from deep mutational

scanning of a protein domain

A. We hypothesize that epistatic interactions mainly result from structural interactions.
Epistasis is the non-independence of single mutant fitness effects, i.e. fitness of the
double mutant deviates from (multiplicative) expectation.

B. Workflow to predict interacting position pairs from deep mutational scanning datasets
(see Methods and Results).

QC: quality control filtering of double mutant variants.

Figure 2

Figure 2: Epistatic variants are enriched for structural proximity

A. Non-parametric null model of epistasis, based on running quantile surfaces, accounts for non-linear dependencies due to lower fitness limit of assay and unspecific epistatic effects. Variants above the 95th or below the 5th percentile of double mutant fitness given their single mutant fitness values were classified as positive (red, ε^+) or negative (yellow, ε^-) epistatic, respectively. Shown is a random sample of 10^4 variants.

B. Cumulative probability of minimal distances between side chain heavy atoms of double mutant variant pairs. Positive and negative epistasis subsets refer to the sets of variants applicable for epistasis analysis, see Figure S1D. All variants, n = 497011; positive

116 epistasis subset, n = 378884; positive epistatic variants ε^+, n = 16907; negative

117 epistasis subset, n = 262768; negative epistatic variants ε^-, n = 10708.

118 C. Upper panel: Examples of double mutant variant distribution of position pairs in rescaled

119 fitness space (fitness of variants below median rescaled by 5th percentile ("5%"), fitness

120 of variants above median fitness rescaled by 95th percentile ("95%")). Lower panel:

121 Position pair-wise fractions of positive and negative epistatic variants. Shown are mean

122 epistatic fractions across 1.000 resampling runs. Black circles mark position pairs with

123 highly significant fractions (mean fraction (either positive or negative epistatic)

124 normalized by standard deviation of fraction over resampling runs greater than 4). Red

125 marks position pairs with significantly larger positive than negative epistatic fractions.

126 Yellow marks opposite position pairs. Blue marks positions pairs with non-significant

127 differences. "x1-5" mark example pairs shown in upper panel.

Position pairs enriched for epistatic interactions predict tertiary structure contacts

131 For each position pair, we next calculated the fraction of positive or negative epistatic double

132 mutant variants (Figure 1B, step 5). Even moderate enrichments for positive and negative

133 epistatic variants appear to be mutually exclusive (Figure 2C). We find that the reason for this

134 are two clusters of positions that exhibit mostly positive or negative interactions among

135 themselves as well as towards subsets of non-cluster positions (Figure S3; as also noted before

136 by Olson et al. (2014)). Both clusters of positive and negative interactions form blocks on the

137 contact map in regions that are enriched for teritary contacts (Figure 3A). We calculated the

138 number of edges of distance smaller than 8 Ångström (Å) that connect position pairs. Consistent

139 with interaction clusters forming a dense network of proximal positions, we find that 45% of the

140 top 100 positions pairs with high fractions of positive or negative epistatic interactions are direct

141 contacts (one edge, 2.4-fold over expectation) and another 41% or 45% share a common

142 neighbour (two edges), while interactions across more edges are depleted (Figure 3C).

143 We attempted to derive a single *epistasis score* per position pair from positive and negative

144 epistasis interations. We therefore first estimated the uncertainty in the fraction of positive or

145 negative epistatic variants per position pair due to measurement errors and the finite number of

146 observed double mutant variants (Figure 1B, step 5). We implemented a resampling approach

where, in each of 1000 runs, each variant's fitness is drawn from a normal distribution with the
measured fitness as mean and the uncertainty due to sequencing coverage as standard
deviation (step 5a). In each resampling run we re-classified positive and negative epistasis of
variants given the drawn fitness values. Finally, the fraction of positive (or negative) epistatic
variants for each position pair was drawn from the posterior probability distribution of epistatic
fractions (step 5b). The posterior probability distribution is the product of the prior probability
distribution - the distribution of epistatic fractions over all position pairs - and the likelihood
function – how likely a given fraction is to generate the observed fraction of epistatic variants.
The uncertainty of the fraction of epistatic variants per position pair is then the standard
deviation of drawn fractions over all resampling runs.

We then combined positive and negative epistasis information by computing the weighted
average of mean epistatic fractions over all resampling runs given their uncertainty. The final
epistasis score is a z-score, which is obtained by further normalizing this weighted average by
its uncertainty, to give priority to high confidence enrichments (step 6).

The position pairs with highest *epistasis scores* are well distributed across the contact map
(Figure 3B), and the number of direct contacts (one edge < 8Å) among the top 100 *epistasis*
score pairs increases to 57% (Figure 3C), thus showing that the *epistasis score* succesfully
incorporates information from both positive and negative epistasis. Additionally, *epistasis scores*
generally correlate with proximity in the tertiary structure (minimal distance between side-chain
heavy atoms, rank correlation coefficient = -0.39, $p < 10^{-6}$, n = 1485; Figure 3D).

We examined the precision to predict direct contacts of amino acids spaced more than five
positions apart (commonly used as restraints for structural simulations, Fig. 3E). Precision is
82% for the top L/5 (n=11, L as length of the protein, 56aa) *epistasis score* pairs, 71% for the
top L/2 pairs and 61% for the top L pairs. Of note, the precision to predict direct contacts from
epistasis scores is 10-20% higher than that of the non-normalized epistasis score, thus showing
the need to account for experimental uncertainity.

Together, our analyses suggest that physical interactions are a main determinant of epistasis in
this protein domain. Nonetheless, the large fractions of epistatically interacting position pairs
separated by a common neighbor (two edges), as well as their connectivity observed on the
contact map, highlight the fact that strong epistatic interactions do also arise from indirect
interactions. Nonetheless, epistatic interaction data suffice to predict tertiary contacts with high
precision.

Figure 3

Figure 3: Position pairs enriched for epistatic interactions predict tertiary structure contacts

    A. Interaction score map for top 100 position pairs with highest positive (lower triangle, red) and negative (upper triangle, yellow) epistatic fractions. Dot size indicates mean epistatic fractions across 1000 re-sampling runs relative to average over top 100 pairs, dot fill indicates distance below 8Å. Underlying in grey is the contact map of the crystal

structure (PDB entry 1pga, distance < 8Å) and shown on top and to the right are its

secondary structure elements (wave – alpha helix, arrow – beta strand).

B. Interaction score map for top 100 pairs with highest merged epistatic enrichments before

normalization by uncertainty (light blue, lower triangle) and highest *epistasis scores*

(black, upper triangle). Dot size indicates interaction score relative to mean of top 100

pairs, everything else as in panel A.

C. Minimal number of edges (contact with distance < 8Å in crystal structure 1pga)

connecting a position pair.

D. Correlation of pair-wise *epistasis score* and minimal heavy atom distance. Fill of contour

map indicates density (red – dense, white – sparse) of position pairs. Spearman

correlation coefficient is indicated.

E. Precision of interaction scores to predict direct contacts (distance < 8Å in crystal

structure 1pga) as a function of the number of top scoring position pairs. Only position

pairs with linear chain distance greater than 5 amino acids are considered (n = 1225

pairs, n = 131 direct contacts). Horizontal dashed line indicates random expectation.

Protein length is 56 amino acids.

Tertiary structure neighborhood leads to correlated epistatic patterns

We reasoned that, if epistasis arises mainly from physical interactions within the protein domain,

proximal positions should also exhibit similar epistasis interactions with all other positions in the

protein (Figure 4A). To test this, we estimated the correlation matrix of positive (or negative)

epistatic fractions across all position pairs in each run of our resampling procedure (Figures 1B

(step 5c) and 4A, see Methods).

Indeed, the top 100 position pairs with the highest epistasis pattern correlation are enriched for

being direct neighbors (51%) and for sharing a common neighbor (Figure 4B) and, conversely,

more proximal position pairs have more highly correlated epistasis patterns (rank correlation

coefficient = -0.47, $p < 10^{-6}$, n = 1485, Figure 4D). Nonetheless, correlation patterns do show a

strong position-dependency, with positions close in the linear sequence forming clusters of

lower or higher correlations (Figures 1B (step 5d lower triangles in the heat maps) and 4C

(lower triangle)). Thus, similar to individual epistatic interactions, correlations of epistatic

interaction patterns are highly enriched for proximal pairs but also contain information on
indirect structural interactions.

We tested whether partial correlations of epistasis patterns, that is dependencies of epistasis
patterns between position pairs after accounting for the epistasis patterns of all other positions,
would minimize indirect interactions and reveal more direct interactions. We derived partial
correlations by inversion of the correlation matrix, estimated the uncertainty of individual values
from resampling, merged values from positive and negative epistatic patterns, and ranked these
by their z-scores, which we refer to as *association scores* (Figure 1B, steps 5c and 6).

In contrast to the correlation of epistatic patterns, *association scores* are slightly less correlated
with proximity (rank correlation coefficient R = -0.36, p < 10^{-6}, n = 1485), with most position pairs
having an *association score* around 0 and only few pairs deviating to higher values (Figure 4D).
These pairs with highest *association scores*, however, are strongly enriched for being direct
neighbors in the structure (71%, Figure 4B). Furthermore, top pairs are well distributed across
the contact map and are enriched for pairs close in the linear chain (Figure 4C). These results
are consistent with a more binary classification of being distant (*association score* around zero)
or being in direct contact (positive *association score*).

We further derived a *combined score* by summing the standardized *epistasis* and *association*
scores, to explore whether combining information from individual epistatic interactions and
epistasis interaction patterns can improve proximity estimates.

We evaluated the precision of the three interaction scores in predicting direct contacts in the
protein domain. For all pairs at least 5 amino acids apart, the *association score* predicts direct
contacts with similar precision to the *epistasis score* (Figure 4F). The *combined score* is, apart
from the very top contacts, consistently more precise than the individual scores (L/5 – 72%, L/2
– 79%, L – 68%, 2*L – 53% precision), showing that information from epistasis interactions and
their interaction patterns across the protein yield complementary results. Finally, we also find
that the interaction scores precisely predict direct contacts over long ranges in the linear chain.
For pairs that are at least 24 amino acids apart, precision is as high as 91% for the top L/5
contacts, 64% for the top L/2 contacts and 48% for the top L contacts (Figure 4G).

Figure 4

Figure 4: Tertiary structure neighborhood leads to correlated epistatic patterns

    A. Example of epistatic interaction patterns of the two proximal positions 7 and 33. Matrix with fraction of positive epistatic variants is shown on the right. Epistatic interaction patterns across the protein correspond to columns of the matrix. Epistatic interaction patterns for positive (upper) and negative (lower) epistatic fractions of positions 7 and 33 are shown on the right. Pearson correlation coefficients for epistatic interaction patterns are given on the right, note that these do not allow a direct comparison to epistasis correlation values given in panel D. Grey vertical bars mark the positions under consideration.

B. Minimal number of edges (contact with distance < 8Å in crystal structure 1pga) connecting a position pair.

C. Interaction score map for top 100 position pairs with highest correlation (lower triangle, orange) and association scores (upper triangle, green). Dot size indicates score across 1000 re-sampling runs relative to average over top 100 pairs, dot fill indicates distance below 8Å. Underlying in grey is the contact map of the crystal structure (PDB entry 1pga, distance < 8Å) and shown on top and to the right are its secondary structure elements (wave – alpha helix, arrow – beta strand).

D. Correlation of pair-wise epistasis pattern correlation to minimal heavy atom distance. Shown are weighted averages of shrunk correlation coefficients for positive and negative epistatic patterns (means across re-sampling runs), with weights as the inverse variances across resampling runs (equivalent to epistatic fraction aggregation). Fill of contour map indicates density (red – dense, white – sparse) of position pairs. Spearman correlation coefficient is indicated.

E. Correlation of pair-wise *association scores* to minimal heavy atom distance. As in panel D.

F. Precision of interaction scores to predict direct contacts (distance < 8Å in crystal structure 1pga) as a function of top scoring position pairs. Only position pairs with linear chain distance greater than 5 amino acids are considered (n = 1225 pairs, n = 131 direct contacts). Horizontal dashed line indicates random expectation.

G. Precision of interaction scores to predict direct contacts (distance < 8Å in crystal structure 1pga) greater than 24 amino acids apart (n = 465 pairs, n = 53 direct contacts). Horizontal dashed line indicates random expectation.

Periodic epistatic patterns reveal secondary structure arrangements

We investigated whether the periodic geometrical arrangement of amino acid residues in secondary structures results in periodic epistasis patterns. This might be expected both locally as well as across the full length of the protein if structural interactions are a primary determinant of epistasis (Andreani and Söding, 2015; Tóth-Petróczy et al., 2016).

Within an alpha helix with 3.6 residues per helical turn, a helical position would be predicted to

interact epistatically with the third or fourth neighbor along the linear amino acid chain. Moreover

these positions should also have similar epistasis patterns towards all other positions in the

protein due to similar side chains orientations (Figure 5A). Similarly, within a beta strand,

positions should interact epistatically with the next-but-one neighbor with whom they should also

share similar epistasis patterns towards all other positions in the protein.

We used a two dimensional kernel smoothing approach to estimate the positions of alpha

helices and beta strands from the deep mutational scanning data (Figure 5B). Here, the

propensity of a position to belong to an alpha helix or a beta strand is calculated by taking into

account how the position interacts with neighboring positions as well as how neighboring

positions interact with each other. For alpha helices, the kernel has a sinusoidal profile with a

period of 3.6 perpendicular to the diagonal, to weight interactions with neighboring positions,

and a Gaussian profile along the diagonal to integrate over multiple adjacent positions. For beta

strands, the kernel has an alternating profile perpendicular to the diagonal, down-weighting

interactions with adjacent positions and up-weighting interactions with the next-but-one position,

and a Gaussian profile along the diagonal to integrate over multiple adjacent positions.

We calculated a p-value for each position to belong to an alpha helix or a beta strand by

comparing the value obtained from kernel smoothing the interaction score data to values

obtained when applying the kernels to 10.000 randomized data sets. To determine the range of

secondary structure elements we combined the p-values of individual positions by Fisher's

method, extending stretches of positions with highly significant scores for alpha helices or beta

strands until the combined p-values drop off when adding additional adjacent positions (see

Methods).

Predicting secondary structures using *combined scores* yielded secondary structure elements

that coincide well with secondary structure elements in the crystal structure (Figure 5C). Using

epistasis and *association scores* to predict secondary structures elements gave similar results,

although the epistasis score analysis failed to detect both beta strands at the end of the domain

due to low statistical power (Figure S4A).

We further used two-dimensional kernel smoothing to detect parallel and anti-parallel beta sheet

interactions, by applying beta strand kernels to off-diagonal entries on the interaction score

matrices (Figure 5D). Using *combined scores* we detected several putative stretches of parallel

and anti-parallel beta sheet interactions throughout the protein domain (Figure 5E). We defined

a set of rules that iteratively fix the most significant stretches of beta sheet interaction when they

are consistent with previously determined secondary structure elements and already fixed beta
sheet interactions (see Methods). This yielded an accurate prediction of beta sheet interactions,
in terms of the number and orientation of sheet interactions as well as the exact pairwise
interactions of all but two position pairs (Figure 5E). Furthermore, updating beta strands
according to inferred beta sheet pairings improved beta strand prediction itself, notably
introducing a correct split of beta strand 1 and 2 and adjusting the length of beta strands 3 and
4. Together, predictions of secondary structures and beta sheet interactions from *combined*
scores have a precision greater than 95% and a recall of greater than 85% (Figure 5F). Similar,
but slightly worse, results are obtained when applying this workflow to *epistasis* and *association*
score data instead (Figures S4B,C).

Figure 5: Periodic epistatic patterns reveal secondary structure arrangements

a) Local interactions reveal signatures of secondary structure elements. Middle line is diagonal of interaction score map (rotated by 45 degree) and shows secondary structure elements of crystal structure (PDB entry 1pga). Data above diagonal shows *combined score* data close to the diagonal, i.e. local interactions. Below the diagonal the same data smoothed with a Gaussian kernel along the direction of the diagonal (i.e. horizontally, length of Gaussian kernel as for kernels in panel b) is shown.

b) Two-dimensional kernels for alpha helix and beta strand detection. Kernel has a Gaussian profile in the direction of the diagonal, and a sinusoidal or alternating profile in the off-diagonal direction for alpha helices and beta strands, respectively.

c) Workflow for secondary structure element prediction and results from *combined score* data compared to known secondary structure from crystal structure 1pga (see Figure S4 for other interaction score data). In workflow, "sumlog(p)" referes to p-value combination by Fisher's method (see Methods for more detail on workflow). In plot, lines give position-wise p-value of secondary structure propensity; dashed lines indicate data was not used (p-value for other secondary structure type lower); boxes indicate final predictions, lower limit of box indicates p-value of final predictions (limited to minimum of 10^{-4}). Note that sumlog(p) of seeds length 3 is not shown.

d) Left: *Combined score* matrix. Right: Position-wise p-values of beta sheet propensity after kernel smoothing off-diagonal elements with beta sheet kernels (shown in the middle) and comparing to 10^4 random datasets (lower triangle - parallel beta sheet propensity, upper triangle – anti-parallel beta sheet propensity). Secondary structure element predictions from panel c) indicated on both matrices (green – alpha helix, red – beta sheet).

e) Workflow to detect beta sheet interactions. Plot shows matrix of combined p-value for extended (beta sheet) seeds, the final predictions and the reference elements in the crystal structure (lower triangle – parallel beta sheets, upper triangle – anti-parallel beta sheet, diagonal –secondary structure elements).

f) Precision and recall for position-wise secondary structure element and beta sheet interaction predictions as shown in panel e).

Folding a protein domain from epistatic mutational patterns

Together, these findings show that genetic interaction data contain information on a protein's secondary and tertiary structure. We therefore tested whether the information extracted from the genetic data would suffice to predict *ab initio* the structure of protein G domain B1. We performed simulated annealing with structural restraints derived from the deep mutational scanning data. Structural restraints were defined for tertiary contacts, secondary structure elements, and beta sheet interactions. In particular, 56 distance restraints (distance < 8Å

between Cβ atoms) for tertiary contacts were defined from the top scoring position pairs with a
linear chain distance greater than five amino acids. We found that using fewer or more contacts
generally decreases the accuracy of the predicted models (Figures S5A,B). Furthermore, we
defined dihedral angle restraints for predicted secondary structure elements. Finally, we defined
restrictive distance restraints (distance smaller than 2.1Å for N-H : C=O atom pairs) for beta
sheet positions that form hydrogen bonds with each other.

We performed structural simulations by simulated annealing using the Xplor-NIH modeling suite
(Schwieters et al., 2003). We set up a three-stage approach, with 500 structural models
generated per stage. Stages 1 and 2 served to identify restraints inconsistent with the most
plausible structural models. These restraints were subsequently removed, thus improving the
precision of the utilized restraints, especially those for tertiary contacts, for which precision
improved from ~65% up to 85% for the top 56 contacts (Figures 6A,B). Stage 3 then served to
refine an average structure for the top 10% of models from stage 2 to obtain a final set of
structural models.

We evaluated the top 5% structural models generated in the third simulation stage against the
known crystal structure of protein G domain B1 (PDB entry 1pga) (Figures 6C and S5A,B).
Models predicted from *combined score* data performed best, with an average Cα-root mean
squared deviation of the top models ($\langle C\alpha - RMSD \rangle$) of 1.9Å and an average template modeling
score of 0.75 (Figure 6D). This accuracy is similar to the resolution of the crystal structure
(2.07Å) and very close to the optimum achievable with our simulation protocol (using true
contacts, secondary structure elements and beta sheet interactions, $\langle C\alpha - RMSD \rangle = 1.4\text{Å}$ and
TM score = 0.79). Consistent with lower precision of contact and secondary structure
predictions, models generated with restrains from *epistasis* or *association scores* have on
average a somewhat lower accuracy ($\langle C\alpha - RMSD \rangle = 2.9\text{Å}$ and $\langle C\alpha - RMSD \rangle = 2.6\text{Å}$,
respectively), with association score models performing slightly but consistently better (Figures
6C and S5A,B). Defining restraints for beta sheet interactions results in an improvement of
about 50% average accuracy for the top structural models (Figure 6C, $\langle C\alpha - RMSD \rangle = 5.7\text{Å}$ for
epistasis score, $\langle C\alpha - RMSD \rangle = 5.4\text{Å}$ for *association score*, and $\langle C\alpha - RMSD \rangle = 3.8\text{Å}$ for
combined score without including restraints for beta sheet interactions).

Together, this shows that genetic interaction data alone is sufficient to accurately predict the
structure of a protein domain.

Figure 6: *Ab initio* folding of protein G domain B1

A. Map of structural restraints derived from *combined scores* (lower triangle - initial scores, upper triangle – scores after reweighting in structural simulations, scores below 0.1*mean(scores) were discarded). Top 56 tertiary contacts (greater than 5 amino acid linear chain distance) are shown as blue circles, alpha helices as squares (score is average weight of dihedral angle restraints), beta strands as diamonds (score is average weight of dihedral angle restraints) and beta sheet interactions as triangles. Underlying in grey is the contact map of the crystal structure (PDB entry 1pga, distance < 8Å) and shown on top and to the right are its secondary structure elements (wave – alpha helix, arrow – beta strand)

B. Improvements in precision of contact predictions due to consistency checking in structural simulations. Solid lines give precision as initially determined from interaction

scores (same as shown in Figure 4F). Dashed lines give precision after simulation stage

2 rescoring of contacts based on consistency with generated structural models for

simulations that were started with top 2*L contacts.

C. Accuracy ($C\alpha$ root mean squared deviation) of top 5% structural models generated from

different sets of restraints compared to crystal structure 1pga. Negative control – only

restraints for secondary structure predicted by PSIPRED (Jones, 1999). Positive control

– 56 true contacts (random subset), secondary structure elements and beta sheet

interactions from crystal structure. All restraints – top 56 contacts, secondary structure

element and beta sheet interaction predictions from the three interaction scores, as

indicated by color. Only contact restraints – top 56 contacts from interaction scores and

secondary structure elements as predicted by PSIPRED.

D. Best structural model generated with all restraints derived from *combined score* (left)

and crystal structure (PDB entry 1pga) (right). Visualization generated in PyMOL (The

PyMOL Molecular Graphics System, 2015).

Figure 7

Figure 7: Requirements for structure determination from deep mutational scanning data

Down-sampling analysis to test data quality requirements for accurate prediction of

protein structures from deep mutational scanning datasets (see also Supplementary

Text). Results from structural simulations when using restraints for tertiary contacts,

secondary structure elements and interactions (left panel) or only tertiary contacts

supplemented with secondary structure elements predicted by PSIPRED (right panel)

which were derived from *combined scores*. Additionally we also varied the calculation of

combined scores, either calculated from full epistasis information (blue) or only positive

(red) or only negative (yellow) epistatic information. Full dataset refers to dataset used

throughout the study. Noisy dataset is down-sampled to 25% of sequencing read

coverage. "Doped" datasets contains only amino acid changes achieved by single

nucleotide mutations (31% of possible single mutants and 10% of possible double

mutants).

Boxplots show: center line, median; box limits, upper and lower quartiles; whiskers, 1.5x

interquartile range; points, outliers.

Discussion

We have shown here that the structure of a protein domain can be accurately predicted simply by quantifying the activity of a large number of single and double mutant variants.

For macromolecules with very large numbers of homologs available in sequence databases, correlated changes in sequence can provide sufficient information for structure determination (Hopf et al., 2012; Kosciolek and Jones, 2014; Marks et al., 2011; Michel et al., 2014; Ovchinnikov et al., 2015; Ovchinnikov et al., 2017; Tóth-Petróczy et al., 2016; Weinreb et al., 2016). However, for many proteins and RNAs this is not the case, and for evolutionary novel folds and families this may never be so (Marks et al., 2012; Ovchinnikov et al., 2017; Tóth-Petróczy et al., 2016). In contrast, our approach should work for any molecule that can be subjected to deep mutagenesis and selection *in vitro* or *in vivo* (Araya et al., 2012; Diss and Lehner, 2018; Fowler et al., 2010; Li et al., 2016; Matreyek et al., 2017; Poelwijk et al., 2017; Rocklin et al., 2017; Starita et al., 2013; Starita et al., 2015; Starr et al., 2017; Weile et al., 2017). However, the same fundamental principle underlies both approaches – that mutations affect each other because of specific structural interactions (Poelwijk and Ranganathan, 2017). The successful application of evolutionary coupling analysis to many different molecules and problems (Hopf et al., 2012; Kosciolek and Jones, 2014; Marks et al., 2011; Michel et al., 2014; Ovchinnikov et al., 2015; Ovchinnikov et al., 2017; Tóth-Petróczy et al., 2016; Weinreb et al., 2016) therefore provides very strong evidence that the approach that we have outlined here will work for many different macromolecules. We thus envisage broad applicability of our framework, including to macromolecules whose structures are difficult to determine by physical methods, such as membrane proteins and intrinsically disordered proteins that only fold upon binding or in particular conditions.

 As a proof-of-principle we used only information extracted from a deep mutational scanning experiment to predict the structure of protein G domain B1. However, taking advantage of recent advances in contact predictions, such as machine learning of sequence-derived protein features, (Schaarschmidt et al., 2018) and homology-driven structure modeling (Rohl et al., 2004) is likely to further improve structure prediction from genetic data, especially for more complex folds than the one tested here.

An analysis of incomplete and down-sampled variants of the deep mutational scanning dataset used here suggests that a high signal-to-noise ratio (measurement range relative to

experimental error of fitness estimates), which allows both positive and negative epistasis to be

quantified, is an important factor for generating datasets with a quality sufficient for protein

structure prediction (Figure 7 and Supplementary Text). At sufficient signal-to-noise ratios,

however, even deep mutational scanning datasets containing an incomplete (random) subset of

double mutants or sequenced at lower coverage might be amenable to protein structure

prediction (Figure 7), thus suggesting that our framework should be easily applicable to longer

molecules.

Finally, we note that our results strongly support the hypothesis (Diss and Lehner, 2018; Fowler

and Fields, 2014; Melamed et al., 2013; Olson et al., 2014; Sahoo et al., 2015; Starr and

Thornton, 2016) that structural interactions are the main cause of specific genetic interactions

(epistasis) within macromolecules.

Acknowledgements

We thank members of the Lehner lab, Torsten Gross and Gregor Mönke for discussions and feedback. This work was supported by a European Research Council (ERC) Consolidator grant (616434), the Spanish Ministry of Economy and Competitiveness (BFU2011-26206 and SEV-2012-0208), the AXA Research Fund, the Bettencourt Schueller Foundation, Agencia de Gestio d'Ajuts Universitaris i de Recerca (AGAUR, SGR-831), the EMBL-CRG Systems Biology Program, and the CERCA Program/Generalitat de Catalunya. J.M.S. was supported by an EMBO Long-Term Fellowship (ALTF 857-2016).

Author Contributions

Conceptualization, B.L. and J.M.S.; Methodology, J.M.S.; Investigation, J.M.S.; Writing - Original Draft, J.M.S.; Writing - Review & Editing, J.M.S. and B.L.; Supervision, B.L.

Declaration of Interests

The authors declare no competing interests.

References

Adhikari, B., Bhattacharya, D., Cao, R., and Cheng, J. (2015). CONFOLD: Residue-residue contact-guided ab initio protein folding. Proteins *83*, 1436-1449.

Altschuh, D., Lesk, A.M., Bloomer, A.C., and Klug, A. (1987). Correlation of co-ordinated amino acid substitutions with function in viruses related to tobacco mosaic virus. Journal of molecular biology *193*, 693-707.

Andreani, J., and Söding, J. (2015). bbcontacts: prediction of β-strand pairing from direct coupling patterns. Bioinformatics *31*, 1729-1737.

Araya, C.L., Fowler, D.M., Chen, W., Muniez, I., Kelly, J.W., and Fields, S. (2012). A fundamental protein property, thermodynamic stability, revealed solely from large-scale measurements of protein function. Proceedings of the National Academy of Sciences *109*, 16858-16863.

Burger, L., and van Nimwegen, E. (2010). Disentangling Direct from Indirect Co-Evolution of
Residues in Protein Alignments. PLoS Computational Biology *6*, e1000633.

Diss, G., and Lehner, B. (2018). The genetic landscape of a physical interaction. Elife *7*,
e32472.

Dunn, S.D., Wahl, L.M., and Gloor, G.B. (2008). Mutual information without the influence of
phylogeny or entropy dramatically improves residue contact prediction. Bioinformatics *24*, 333-
340.

Ekeberg, M., Lövkvist, C., Lan, Y., Weigt, M., and Aurell, E. (2013). Improved contact prediction
in proteins: Using pseudolikelihoods to infer Potts models. Physical Review E *87*, 012707.

Fowler, D.M., Araya, C.L., Fleishman, S.J., Kellogg, E.H., Stephany, J.J., Baker, D., and Fields,
S. (2010). High-resolution mapping of protein sequence-function relationships. Nature Methods
7, 741-746.

Fowler, D.M., and Fields, S. (2014). Deep mutational scanning: a new style of protein science.
Nature Methods *11*, 801-807.

Gallagher, T., Alexander, P., Bryan, P., and Gilliland, G.L. (1994). Two crystal structures of the
B1 immunoglobulin-binding domain of streptococcal protein G and comparison with NMR.
Biochemistry *33*, 4721-4729.

Gloor, G.B., Martin, L.C., Wahl, L.M., and Dunn, S.D. (2005). Mutual information in protein
multiple sequence alignments reveals two classes of coevolving positions. Biochemistry *44*,
7156-7165.

Göbel, U., Sander, C., Schneider, R., and Valencia, A. (1994). Correlated mutations and residue
contacts in proteins. Proteins *18*, 309-317.

Halabi, N., Rivoire, O., Leibler, S., and Ranganathan, R. (2009). Protein sectors: evolutionary
units of three-dimensional structure. Cell *138*, 774-786.

Hopf, T.A., Colwell, L.J., Sheridan, R., Rost, B., Sander, C., and Marks, D.S. (2012). Three-
Dimensional Structures of Membrane Proteins from Genomic Sequencing. Cell *149*, 1607-1621.

Hopf, T.A., Ingraham, J.B., Poelwijk, F.J., Schärfe, C.P.I., Springer, M., Sander, C., and Marks,
D.S. (2017). Mutation effects predicted from sequence co-variation. Nature Biotechnology.

Jones, D.T. (1999). Protein secondary structure prediction based on position-specific scoring
matrices. Journal of molecular biology *292*, 195-202.

Jones, D.T., Buchan, D.W.A., Cozzetto, D., and Pontil, M. (2012). PSICOV: precise structural
contact prediction using sparse inverse covariance estimation on large multiple sequence
alignments. Bioinformatics *28*, 184-190.

Kamisetty, H., Ovchinnikov, S., and Baker, D. (2013). Assessing the utility of coevolution-based
residue-residue contact predictions in a sequence- and structure-rich era. Proceedings of the
National Academy of Sciences *110*, 15674-15679.

Kosciolek, T., and Jones, D.T. (2014). De novo structure prediction of globular proteins aided by
sequence variation-derived contacts. PLoS ONE *9*, e92197.

Lehner, B. (2011). Molecular mechanisms of epistasis within and between genes. Trends Genet
27, 323-331.

Li, C., Qian, W., Maclean, C.J., and Zhang, J. (2016). The fitness landscape of a tRNA gene.
Science *352*, 837-840.

Lockless, S.W., and Ranganathan, R. (1999). Evolutionarily conserved pathways of energetic
connectivity in protein families. Science *286*, 295-299.

Marks, D.S., Colwell, L.J., Sheridan, R., Hopf, T.A., Pagnani, A., Zecchina, R., and Sander, C.
(2011). Protein 3D structure computed from evolutionary sequence variation. PLoS ONE *6*,
e28766.

Marks, D.S., Hopf, T.A., and Sander, C. (2012). Protein structure prediction from sequence
variation. Nature Biotechnology *30*, 1072-1080.

Matreyek, K.A., Stephany, J.J., and Fowler, D.M. (2017). A platform for functional assessment
of large variant libraries in mammalian cells. Nucleic Acids Research *45*, e102.

McLaughlin Jr, R.N., Poelwijk, F.J., Raman, A., Gosal, W.S., and Ranganathan, R. (2012). The
spatial architecture of protein function and adaptation. Nature *491*, 138-142.

Melamed, D., Young, D.L., Gamble, C.E., Miller, C.R., and Fields, S. (2013). Deep mutational
scanning of an RRM domain of the Saccharomyces cerevisiae poly(A)-binding protein. RNA *19*,
1537-1551.

Michel, M., Hayat, S., Skwark, M.J., Sander, C., Marks, D.S., and Elofsson, A. (2014).
PconsFold: improved contact predictions improve protein models. Bioinformatics *30*, i482-488.

Morcos, F., Pagnani, A., Lunt, B., Bertolino, A., Marks, D.S., Sander, C., Zecchina, R., Onuchic,
J.N., Hwa, T., and Weigt, M. (2011). Direct-coupling analysis of residue coevolution captures
native contacts across many protein families. Proceedings of the National Academy of Sciences
108, E1293-1301.

Olson, C.A., Wu, N.C., and Sun, R. (2014). A comprehensive biophysical description of pairwise
epistasis throughout an entire protein domain. Current biology : CB *24*, 2643-2651.

Ovchinnikov, S., Kinch, L., Park, H., Liao, Y., Pei, J., Kim, D.E., Kamisetty, H., Grishin, N.V.,
and Baker, D. (2015). Large-scale determination of previously unsolved protein structures using
evolutionary information. eLife *4*, e09248.

Ovchinnikov, S., Park, H., Varghese, N., Huang, P.-S., Pavlopoulos, G.A., Kim, D.E., Kamisetty,
H., Kyrpides, N.C., and Baker, D. (2017). Protein structure determination using metagenome
sequence data. Science *355*, 294-298.

Poelwijk, F.J., and Ranganathan, R. (2017). The relation between alignment covariance and
background-averaged epistasis arXiv, 1-7.

Poelwijk, F.J., Socolich, M., and Ranganathan, R. (2017). Learning the pattern of epistasis
linking genotype and phenotype in a protein. bioRxiv, 213835.

Rocklin, G.J., Chidyausiku, T.M., Goreshnik, I., Ford, A., Houliston, S., Lemak, A., Carter, L.,
Ravichandran, R., Mulligan, V.K., Chevalier, A.*, et al.* (2017). Global analysis of protein folding
using massively parallel design, synthesis, and testing. Science *357*, 168-175.

Rohl, C.A., Strauss, C.E., Misura, K.M., and Baker, D. (2004). Protein structure prediction using
Rosetta. Methods Enzymol *383*, 66-93.

Sahoo, A., Khare, S., Devanarayanan, S., Jain, P.C., and Varadarajan, R. (2015). Residue
proximity information and protein model discrimination using saturation-suppressor
mutagenesis. eLife *4*, 371.

Schaarschmidt, J., Monastyrskyy, B., Kryshtafovych, A., and Bonvin, A. (2018). Assessment of
contact predictions in CASP12: Co-evolution and deep learning coming of age. Proteins *86*
Suppl 1, 51-66.

Schäfer, J., and Strimmer, K. (2005). A shrinkage approach to large-scale covariance matrix
estimation and implications for functional genomics. Statistical applications in genetics and
molecular biology *4*, Article32.

Schwieters, C.D., Kuszewski, J.J., Tjandra, N., and Clore, G.M. (2003). The Xplor-NIH NMR
molecular structure determination package. Journal of Magnetic Resonance *160*, 65-73.

Seemayer, S., Gruber, M., and Söding, J. (2014). CCMpred--fast and precise prediction of
protein residue-residue contacts from correlated mutations. Bioinformatics *30*, 3128-3130.

Starita, L.M., Pruneda, J.N., Lo, R.S., Fowler, D.M., Kim, H.J., Hiatt, J.B., Shendure, J., Brzovic,
P.S., Fields, S., and Klevit, R.E. (2013). Activity-enhancing mutations in an E3 ubiquitin ligase
identified by high-throughput mutagenesis. Proceedings of the National Academy of Sciences
110, E1263-1272.

Starita, L.M., Young, D.L., Islam, M., Kitzman, J.O., Gullingsrud, J., Hause, R.J., Fowler, D.M.,
Parvin, J.D., Shendure, J., and Fields, S. (2015). Massively Parallel Functional Analysis of
BRCA1 RING Domain Variants. Genetics *200*, 413-422.

Starr, T.N., Picton, L.K., and Thornton, J.W. (2017). Alternative evolutionary histories in the
sequence space of an ancient protein. Nature *5*, e16965.

Starr, T.N., and Thornton, J.W. (2016). Epistasis in protein evolution. Protein Science *25*, 1204-
1218.

The PyMOL Molecular Graphics System, V.S., LLC. (2015). The PyMOL Molecular Graphics
System, Version 1.8.

Tokuriki, N., and Tawfik, D.S. (2009). Stability effects of mutations and protein evolvability.
Current Opinion in Structural Biology *19*, 596-604.

Tóth-Petróczy, A., Palmedo, P., Ingraham, J., Hopf, T.A., Berger, B., Sander, C., and Marks,
D.S. (2016). Structured States of Disordered Proteins from Genomic Sequences. Cell *167*, 158-
170.e112.

Weigt, M., White, R.A., Szurmant, H., Hoch, J.A., and Hwa, T. (2009). Identification of direct
residue contacts in protein-protein interaction by message passing. Proceedings of the National
Academy of Sciences *106*, 67-72.

Weile, J., Sun, S., Cote, A.G., Knapp, J., Verby, M., Mellor, J.C., Wu, Y., Pons, C., Wong, C.,
van Lieshout, N.*, et al.* (2017). A framework for exhaustively mapping functional missense
variants. Molecular Systems Biology *13*, 957.

Weinreb, C., Riesselman, A.J., Ingraham, J.B., Gross, T., Sander, C., and Marks, D.S. (2016).
3D RNA and Functional Interactions from Evolutionary Couplings. Cell *165*, 963-975.

Zhang, Y., and Skolnick, J. (2004). Scoring function for automated assessment of protein
structure template quality. Proteins *57*, 702-710.

Methods:

Datasets and preprocessing

Protein G domain B1 deep mutational scanning data was obtained from the supplementary information of Olson et al. (2014). The data consists of summed read counts of three replicate experiments assaying the binding affinity of protein G domain B1 variants to immunoglobulin G (IgG).

Read frequencies of each single or double mutant variant in input library and output library (after binding affinity assay) were calculated as variant read counts relative to wild-type variant read counts. A variant's fitness was calculated as the natural logarithm of the ratio of output to input read frequency, i.e. $f_i = log(\frac{n_i^{out}/n_{wt}^{out}}{n_i^{in}/n_{wt}^{in}})$, with n as read counts, superscripts denoting input or output sequencing library and subscripts denoting variant i or wild-type variant.

Additionally, a correction factor for wild-type fitness was derived, based on the observation that the upper modes of single and double mutant variant fitness distributions were not centered on the wild-type fitness value and that the epistasis distribution was highly non-symmetric. The kernel density estimate of the single mutant fitness distribution was calculated (using the R function *density* with parameter *bw* set to 0.15) and the correction factor was derived as the position of the upper fitness mode as $cf = 0.198$. Each fitness value was corrected by this factor, which centered the upper modes of single and double mutant fitness distributions on the wild-type fitness value and resulted in a symmetric epistasis distribution. We note, however, that this correction has little effect on the downstream analysis, because of the use of quantile-based estimates to call positive or negative epistasis. This would be different if the fitness deviation from expectation would have been used to calculate epistasis.

The standard error of fitness estimates was calculated from read counts under Poissonian assumptions, i.e. $\sigma_i = \sqrt{\frac{1}{n_i^{in}} + \frac{1}{n_i^{out}} + \frac{1}{n_{wt}^{in}} + \frac{1}{n_{wt}^{out}}}$. We note that this is a lower bound estimate of the actual error, due to the lack of replicate information.

Each measurement assay has a lower measurement limit due to unspecific background effects (Figure S1A). In the case of the IgG-binding assay for protein G domain B1, this is presumably mainly due to unspecific carry-over on beads (Olson et al., 2014). The fitness values derived from the measurement are therefore a convolution of the actual binding affinities to IgG and

nonspecific carryover, i.e. $exp(f_i^{measured}) = exp(f_i^{binding}) + exp(f^{carryover})$. Fitness values of
variants close to the lower measurement limit of the assay are therefore dominated by
unspecific carryover effects. The lower measurement limit of the assay was estimated by two
approaches that yielded very similar estimates. One, from a kernel density estimate of the single
mutant fitness distribution (R function *density* with parameter *bw* set to 0.15), where the position
of the lower mode of the data corresponded to $f^{carryover} = -5.85$ (~0.3% on linear scale). Two,
from examining the fitness distribution of double mutants with expected fitness lower than -8, i.e.
double mutants resulting from two lethal or nearly lethal single mutant variants, whose fitness
values are thus expected to be dominated by background effects. The median of this
background fitness distribution yielded an estimate of $f^{carryover} = -5.89$. The mean of the two
estimates was used in downstream analyses.

7% of double mutant variants were discarded due to too low sequencing coverage in input or
output libraries (Figure S1C). That is, variants that had less than 200 input reads and no output
reads were discarded, because it is not possible to determine their fitness. Above 200 input
reads, variants without output reads are certain to be dominated by nonspecific carryover
effects. These variants were retained and their fitness was calculated by setting their output
read count to 0.5.

Epistasis classification

Epistasis was calculated from a non-parametric null model (Figure 2A) in order to account for
nonlinearities close to the lower limit of the fitness assay measurement range, non-specific
epistatic behavior resulting from e.g. thermodynamic stability thresholds as well as differential
uncertainty of fitness measurements across the fitness landscape (Figure S1B).

First, double mutant fitness values were corrected by subtracting the average local fitness
computed using a two-dimensional generalized additive model (using R function *gam* with
tensor product smooths, with the two constituting single mutant fitness values as independent
variables and the resulting double mutant fitness value as dependent variable). This was
necessary to avoid boundary effects of quantile-based fits in boundary regions with non-zero
slopes. 5th and 95th percentile surfaces were then fit to these corrected double mutant fitness
values, by computing for each double mutant variant the 5th and 95th percentile of the fitness
distribution made up of the 1% closest neighbors in single mutant fitness space. Double mutant

variants with fitness values below the 5th or above the 95th percentile were categorized as
negative or positive epistatic, respectively (Figure 2A).

The evaluation of positive or negative epistasis was, however, restricted to specific subsets of
the data where measurement errors do not impede epistasis classification (Figure S1D). The
data subset in which variants were potentially classified as positive epistatic is limited to data
regions where

● the 95th percentile fitness surface is below wild-type fitness

● at least one single mutant fitness value is significantly smaller than wild-type fitness at
five standard errors

● the expected fitness (sum of both single mutant fitness values) is not significantly higher
than wild-type at five standard errors

● the expected fitness is significantly higher than twice the lower fitness limit of the assay

The rationale for criteria 1-3 is to avoid double mutants from two neutral single mutants,
because these are dominated by measurement noise of wild-type like variants. Minimal
restrictions were instead applied to the lower limits of the measurement range (criterion 4),
because otherwise no/little epistasis quantification would have been available for several
positions with very strong detrimental effects as well as because strong positive epistatic effects
are observed in these regions, despite the dominance of background measurement effects.

The data subset in which variants were potentially classified as negative epistatic is limited to
data regions where

● the 5th percentile fitness surface is above the 95th percentile of the background effect
distribution; this value is derived from the 95th percentile of double mutant fitness values
with expected fitness below -8 (analogous to lower fitness limit estimation, see above).

● both single mutant fitness values are significantly higher than the lower limit of the
fitness assay measurement range at five standard errors

● the expected fitness (sum of both single mutant fitness values) is not significantly higher
than wild-type at five standard errors

The rationale for criteria 1 and 2 is to avoid background measurement effects that make
negative epistasis quantification unreliable.

As a result of these restrictions as well as differences in initial coverage, the number of double
mutant variants that can be used to assess positive and negative epistasis varies substantially
across position pairs (with an average of 252 and 187 variants per position pair in the positive
and negative epistasis subsets, respectively; Figure S2).

Epistatic interactions

We derived several interaction scores to estimate which position pairs are in close contact in the
tertiary structure (see Figure 1B). These scores are based on summarizing epistasis information
on the position pair-level and accounting for the uncertainty inherent in the summarized
estimates due to differential error of fitness estimates across the measurement range (Figure
S1B) as well as varying numbers of double mutants amenable to epistasis classification
(Figures S1D and S2).

To summarize epistasis information on the position pair-level, we calculated the fraction of
positive or negative epistatic variants per position pair. The fraction of positive epistatic variants
per position pair is the number of positive epistatic variants divided by the number of all variants
that lie in the double mutant space amenable to positive epistasis classification (Figure 1B, step
5b, equivalent calculation for negative epistasis fraction). Because enrichments with positive
and negative epistatic variants per position appear mutually exclusive (Figure S3B), we treated
both separately and only aggregated them to derive the final interaction scores.

To estimate the uncertainty in epistatic fractions per position pair for downstream analyses we
implemented a re-sampling approach (Figure 1B, step 5, described here for positive epistatic
variants, but equivalent for negative epistatic variants). In each of 1000 re-sampling runs:

- 758 ● each variant's fitness was drawn from a normal distribution with the measured fitness as
- 759 mean and the uncertainty due to sequencing coverage as standard deviation $f_{ij}^{sampled} =$
- 760 $\mathcal{N}(f_{ij}, \sigma_{ij} + \sigma_i + \sigma_j)$ (step 5a)
- 761 ● positive epistasis of variants was re-classified given the drawn fitness values (also step
- 762 5a, see also Figure 2A)
- 763 ● each position pair's fraction of positive epistatic variants was drawn from the posterior
- 764 probability distribution of how likely an underlying true fraction of epistatic variants E_{xy}^+ is
- 765 to generate the observed fraction of epistatic variants given the finite number of overall

variants, i.e. $e_{xy}^+ \sim p(E_{xy}^+ \mid \# \varepsilon_{xy}^+, \# \text{variants}_{xy})$ (step 5b). The posterior probability

distribution is the product of a prior probability distribution – the kernel density estimate

of the expected epistatic fractions across all position pairs (calculated using R function

density with parameter *bw* set to 0.05) – and the likelihood function for the underlying

true fraction of epistatic variants given the observed fraction of epistatic variants and the

overall number of variants under binomial sampling assumptions

To derive an interaction score from the epistatic fractions per position pair, mean positive and

negative epistatic fractions across resampling runs were combined by weighted averaging, with

weights as the inverse variances of epistatic fractions across resampling runs, i.e. $\langle e_{xy} \rangle =$

$\dfrac{\langle e_{xy}^+ \rangle * \sigma_{e_{xy}^+}^{-2} + \langle e_{xy}^- \rangle * \sigma_{e_{xy}^-}^{-2}}{\sigma_{e_{xy}^+}^{-2} + \sigma_{e_{xy}^-}^{-2}}.$

To arrive at the final *epistasis score,* the mean epistatic fractions were further normalized by

their combined uncertainty, $E_{xy} = \langle e_{xy} \rangle \big/ \sigma_{xy}$, with $\sigma_{xy} = \left(\sigma_{e_{xy}^+}^{-2} + \sigma_{e_{xy}^-}^{-2} \right)^{-1/2}$ (step 6).

Epistasis patterns

In addition to the *epistasis score* we derived an interaction score, termed *association score,*

from the partial correlation of epistasis patterns between position pairs. The rationale behind this

score is that proximal positions in the protein should have similar distances and geometrical

arrangements towards all other positions in the protein and should therefore also have similar

patterns of epistatic interactions with all other positions.

In each re-sampling run we constructed a symmetric matrix of the drawn positive epistatic

fractions e_{xy}^+ with number of rows and columns as the number of mutated positions (Figure 1B,

step 5c, equivalent for negative epistatic fractions). Missing values (positions pairs without

observed variants) were supplemented by drawing a random value from the overall distribution

of epistatic fractions. A pseudo count equal to the first percentile of the epistatic fraction

distribution was added to each epistatic fraction. Diagonal elements (epistatic fractions of a

position with itself) were set to 1. The matrix values were transformed by the natural logarithm

(to make distributions more symmetric, thus correlations are not dominated by few position pairs

with large epistatic fractions) and for each pair of columns the Pearson correlation coefficient

was calculated to arrive at the correlation matrix (step 5d).

A shrinkage approach was used to improve the estimate of the correlation matrix (Schäfer and
Strimmer, 2005). In brief, the empirical correlation matrix is shrunk towards the identity matrix in
order to minimize the mean-squared error between estimated and true correlation matrix.
Additionally, this yields a positive definite and well-conditioned correlation matrix, suitable for
inversion. All computations on correlation matrices, shrinkage and matrix inversion were
performed with the R package *corpcor*, functions *cor.shrink* and *pcor.shrink* (Schäfer and
Strimmer, 2005).

Partial correlations of epistatic patterns between each position pair were calculated by inverting
the shrunk correlation matrix and normalizing each off-diagonal entry of the inverted matrix by

the geometric mean of the two respective diagonal entries, i.e. $a_{xy}^+ = \dfrac{r_{xy}^{-1}}{\sqrt{r_{xx}^{-1} * r_{yy}^{-1}}}$, with r_{xy}^{-1} as the

(x,y)-entry of the inverted correlation matrix (Figure 1B, step 5d). Equivalent to the *epistasis*
score, positive and negative partial correlation estimates were merged by calculating weighted
averages of their mean estimates across re-sampling runs, with weights as the inverse

variances across resampling runs, i.e. $\langle a_{xy} \rangle = \dfrac{\langle a_{xy}^+ \rangle * \sigma_{a_{xy}^+}^{-2} + \langle a_{xy}^- \rangle * \sigma_{a_{xy}^-}^{-2}}{\sigma_{a_{xy}^+}^{-2} + \sigma_{a_{xy}^-}^{-2}}$, and the final *association*

score normalized by the combined uncertainty, $A_{xy} = \langle a_{xy} \rangle \big/ \sigma_{xy}$, with

$\sigma_{xy} = \left(\sigma_{a_{xy}^+}^{-2} + \sigma_{a_{xy}^-}^{-2} \right)^{-1/2}$ (step 6).

We further derived a *combined score* by summing the standardized *epistasis* and *association*
scores, i.e. $C_{xy} = \dfrac{E_{xy} - \langle E \rangle}{\sigma_E} + \dfrac{A_{xy} - \langle A \rangle}{\sigma_A}$, to explore whether combining information from individual
epistatic interactions and epistasis interaction patterns can further improve proximity estimates.
We note that this is a naïve approach to combining the information from these two sources, and
surely more sophisticated approaches that further improve proximity estimates can be
developed. Additionally, we note that it should be possible to also solve the inverse problem for
epistatic fraction matrix, thereby potentially improving proximity estimates from epistatic
interactions.

820 Secondary structure prediction

We used a two dimensional kernel smoothing approach to predict secondary structure elements
from deep mutational scanning data (Figure 5). For a given position along the linear chain (on

the diagonal of the interaction score matrix), interaction scores (off the diagonal) are integrated

with distance-specific weighting according to the kernel, which reflects the known geometry of

secondary structures.

The alpha kernel takes on a sinusoidal profile perpendicular to the diagonal to weight

interactions according to whether the position pair considered should have congruent side-chain

orientations (see diagonal and perpendicular profiles in Figure 5B). The kernel was defined as

$k_\alpha(d,p) = \left(\cos\left(p * \frac{2\pi}{3.6} \right) + 1/3 \right) * e^{-\frac{d^2}{c^2}}$, with $d = |2x - i - j|$ as the diagonal distance of the

interaction ij (off the diagonal) to the reference position x (on the diagonal) and $p = |i - j|$ as the

perpendicular distance of the interaction off the diagonal. The kernel weight for positions with p

> 5 was set to 0, thus only including interactions across little more than the first helical turn.

Finally, $c = 4$ is the integration scale for the Gaussian kernel along the diagonal. While in

principle smaller integration scales do yield nosier results and longer integration scales can lead

to non-detection of shorter secondary structure stretches, we found that in practice our whole

approach (including the actual detection algorithm described below) is robust to alterations of

the integration length.

The kernel smoothed alpha value for a given position x on the diagonal is then calculated as the

sum over all interaction scores times their kernel weights $K_{\alpha,x} = \sum_i \sum_j k_\alpha(d,p) * S_{ij}$, where S_{ij} is

one of the interaction scores (*epistasis, association* or *combined score*) at position pair *ij*.

The beta kernel takes an alternating profile perpendicular to the diagonal to weight interactions

according to alternating side-chain orientations in a beta strand and was defined as $k_\beta(d,p) =$

$\left((p + 1) \bmod 2 - 1/3 \right) * e^{-\frac{d^2}{c^2}}$, with $c = 4$. Only interactions with perpendicular distances equal

or smaller than two (i.e. $k_\beta(d, p > 3) = 0$) were included.

To calculate whether kernel-weighted interaction scores of a specific position are larger than

expected, they were compared to those obtained from 10^4 randomized control datasets.

Randomization was performed by shuffling all interaction scores, while preserving matrix

symmetry, and kernel-weighted interaction scores from randomized control datasets were

calculated for each position independently to control for possible boundary effects in positions

close to the borders of the protein chain. A p-value for each position was calculated as the

fraction of random controls with kernel smoothed values above that of the real data.

Secondary structure elements were identified by searching for continuous stretches of positions
with high propensities to belong to either alpha helices or beta strands. The following workflow
was implemented:

1. calculate a combined p-value for seeds of length 3 by combining position-wise p-values
using Fisher's method for both alpha and beta kernel smoothed interaction scores
2. separate positions according to whether combined p-values of seeds from alpha or beta
kernels are more significant, i.e.
2.1. for alpha helical propensity calculations only consider stretches of at least 5 positions
where the combined p-value of seeds for alpha kernel smoothing is smaller than that
from beta kernel smoothing (thus setting the lower size limit of alpha helical elements to
five)
2.2. for beta strand propensity calculations only consider stretches of at least 3 positions
where the combined p-value of seeds for beta kernel smoothing is smaller than that
from alpha kernel smoothing (thus setting the lower size limit of beta strands to three)
For alpha helices and beta strands separately and while combined p-values of seeds < 0.05

3. select most significant seed
4. test whether extension to any side yields lower combined p-value
4.1. if yes: extend seed in this direction and repeat step 4
4.2. else: repeat step 4 once to see whether further extension in same direction yields lower
combined p-value
4.2.1. if yes: extend and repeat step 4
4.2.2. else: proceed to step 5
5. fix as secondary structure element and delete all 'used' p-values (and combined seed p-
values), such that other elements cannot incorporate them
6. check whether other already fixed elements are adjacent or at most one position away
6.1. if yes: merge both elements
7. repeat steps 3-6 until no more seeds with combined p-value < 0.05 are left
This yields a list of predicted locations of secondary structure elements. We note that the
secondary structure elements predicted from deep mutational scanning data could be compared
to and combined with predictions derived from other tools, such as PSIPRED (Jones, 1999), to
further improve reliability.

To detect beta sheet interactions a modified beta strand kernel was used. In contrast to beta
strand detection, the beta sheet interaction kernel is centered on each off-diagonal position. For

beta sheet kernels diagonal and perpendicular distances are therefore modified as $d = $

$|x + y - i - j|$ and $p = |x - i - (y - j)|$. The kernels to detect parallel and anti-parallel beta

sheets differ in which is their 'diagonal' direction, i.e. the direction at which consecutive position

pairs interact in the beta sheet (Figure 5D). Therefore parameters *d* and *p* were swapped for the

anti-parallel beta sheet kernel. Also, because these positions can be deemed the most crucial

for deciding whether a position participates in a beta sheet interaction or not, we up-weighted

these positions (those with $p = 0$) in the kernel by a factor of two, i.e. $K_\beta(d, 0) = 4/3 * e^{-\frac{d^2}{c^2}}$.

Beta sheet interactions were identified by searching for the most significant stretches of parallel

and anti-parallel interactions (similar to workflow for alpha helices and beta strands), then

identifying the set of most significant interactions that is consistent with previously predicted

secondary structure elements.

In particular, step 1 & 3-7 from the above-described workflow were performed for the parallel

beta sheet kernel on each sub-diagonal (parallel to the main diagonal) of the interaction score

matrix separately; and for the anti-parallel beta sheet kernel on each perpendicular diagonal of

the interaction score matrix separately.

The steps were modified as follows:

● the minimal combined p-value to consider a seed was lowered to 0.001

● for anti-parallel beta sheet interactions, only positions with a distance greater than two

from the main diagonal were used to calculate seed p-values (assuming anti-parallel

beta sheet interactions need a turn of at least length two to be connected)

● for parallel beta sheet interactions, only sub-diagonals with a distance greater than six

from the main diagonal were considered (assuming parallel beta sheet interactions of

two adjacent beta strands need a linker region)

We extended the workflow with the following steps to predict beta sheet interactions within the

protein domain:

8. compute association of seeds with known beta strands (e.g. seed positions overlap strand 1

on one side and coincide with strand 3 on the other side)

9. while there are seeds with p < 0.001: pick most significant seed from either the parallel or

anti-parallel sheet subset

10. check consistency with secondary structure elements

10.1.   discard the seed and jump back to step 9 if

916   10.1.1. it is overlapping or too close to an alpha helix or the linker region between two

917   beta strands that interact (minimal distance smaller one)

918   10.1.2. at least one of the two strands it is associated with already has two other beta

919   sheet interactions or the total number of beta sheet interactions exceeds 2*(#beta

920   strands – 1)

921   10.2.   modify secondary structure elements and start anew from step 3 if

922   10.2.1. one sides of the seed is not associated to a known beta strand: create this beta

923   strand

924   10.2.2. if both sides of the seed are associated with the same known strand: split the

925   strand and create a linker region in-between the strands

926   10.3.   else fix the beta sheet interaction and delete all other interactions that are

927   associated with the same strands and haven't been fixed yet, jump back to step 9

928 11. if no more seeds with p<0.001, finish

929 12. update beta strands: keep only those positions that are part of a beta sheet interaction

Protein distance metrics

The minimal side chain heavy atom distance, i.e. the minimal distance between any two side

chain heavy atoms of a position pair (in case of glycine, $C\alpha$), was used as the general distance

measure. The reference structure used was PDB entry 1pga, which was determined by X-ray

diffraction (Gallagher et al., 1994). A direct contact or edge was defined as minimal side chain

heavy atom distance < 8Å. The Floyd-Warshall algorithm (implemented as custom script in R)

was used to calculate the minimal number of edges that connect any two positions in the

protein.

Protein folding

To *ab initio* predict protein structures, we performed simulated annealing using the Xplor-NIH

modeling suite (Schwieters et al., 2003) with structural restraints derived from the deep

mutational scanning data. Simulations were performed in three stages, in each of which 500

structural models were generated. Stages 1 and 2 served to identify inconsistencies among

defined structural restraints. Additionally, in stage 2 an average structure of the best 10% of

models was calculated. Stage 3 served to refine this average structure to obtain a final set of
best models.

Restraints from top predicted contacts (position pairs with highest interaction scores) were
implemented by setting Cβ-Cβ atom distances (Cα in case of Glycine) between positions to
range between 0 and 8Å and weighting the restraints according to their relative interaction score
(interaction score divided by mean interaction score of all predicted contacts used).

Restrains from secondary structures elements were implemented as dihedral angle restraints.
Dihedral angles of both beta strands and alpha helices were set to range between values
commonly observed in crystal structures (Adhikari et al., 2015), for alpha helices $\Phi_\alpha =$
$-63.5° \pm 4.5°$ and $\Psi_\alpha = -41.5° \pm 5°$ and for beta strands $\Phi_\beta = -118° \pm 10.7°$ and $\Psi_\beta = 134° \pm$
$8.6°$.

Restraints for beta sheet interactions were implemented by setting H-N:O=C hydrogen bond
distances between interacting positions to range between 1.8 and 2.1Å (Adhikari et al., 2015),
with weight one. Predictions of beta sheet interactions derived from deep mutational scanning
data yield a string of interacting positions, but hydrogen bonding in beta sheets occurs in
specific non-continuous patterns between position pairs (between alternating positions off the
interaction diagonal in parallel beta sheets and between every second set of position pairs in
anti-parallel beta sheets). Specifically, for each set of interacting positions there are two
alternative patterns of hydrogen bonding possible. These alternative possibilities of pairing were
implemented as mutually exclusive selection pairs with the "assign … or" syntax in Xplor-NIH.

Distance restraints were implemented in Xplor-NIH as NOE (nuclear Overhausser effect)
potential, with potential type set to "soft" for stages 1 and 2 and "hard" for the final simulation
stage. Dihedral angle restraints were implemented via the CDIH potential.

After simulation stages 1 and 2 restraints were checked for their consistency with predicted
structural models. Frist, structural models were clustered according to their violations of
distance and dihedral angle restraints (k-means clustering, k = 4). Clusters were ranked by the
mean total energy (from all energy potentials used) of their 50 models with lowest total energy
(or all of their structures if clusters are smaller 50 models). From the 50 models with the lowest
total energy from the top-ranked cluster (or however many top-ranked clusters were necessary
to arrive at 50 models) the fraction of models that violate specific restraints was recorded. For
the subsequent simulation stage, distance restraints were down-weighted according to the
fraction of models that violated them, $w_{x,i} = w_{x,i-1} * (1 - f_x)^2$, and distance restraints with a

weight below 0.1 were discarded. There is no option to weight dihedral angle restraints, thus
instead dihedral angle restraints with a 'weight' below 1/3 were discarded for the subsequent
simulation stages.

The top 5% structural models from simulation stage 3 were evaluated against the known crystal
structure of protein G domain B1. The TM-score program (update 2016/03/23) was used to
calculate the Cα root mean squared deviation and the template modeling score (Zhang and
Skolnick, 2004).

Several types of control simulations were performed to judge the predictive power of restraints
derived from deep mutational scanning data. Simulations without restraints from predicted
contacts and beta sheet interactions, but with restraints from secondary structure elements
predicted by PSIPRED (version 2.6, (Jones, 1999)), which serve as a negative control.
Simulations with restraints derived from the known crystal structure of protein G domain B1
(PDB entry 1pga). Here, 56 true contacts of position pairs with linear chain distance greater than
5 amino acids were randomly sampled and beta sheet interactions were determined by PyMOL
(The PyMOL Molecular Graphics System, 2015). These simulations serve as a positive control
and give the maximally achievable accuracy of our Xplor-NIH workflow. Finally, a set of
simulations that only used contact restraints derived from top predicted interaction sores and
secondary structure restraints derived from PSIPRED predictions but no restraints for beta
sheet interactions. These simulations serve to judge the power of predicting secondary structure
elements and beta sheet interactions from deep mutational scanning data.

Code availability

1000 Data was analysed with custom scripts written and executed in R programming language,
1001 version 3.4.3. Structural simulations were performed with Xplor-NIH modeling suite version
1002 2.46. Custom R code is available upon request from the corresponding author.

1003

Data availability

1005 No primary data was generated in this study. Processed data is available upon request from the
1006 corresponding author.

Supplementary Text

Down sampling analysis

We evaluated how robust our prediction strategy is to lower data quality (Figure 7), thus assessing what might be the minimal requirements for deep mutational scanning datasets to be useful for structure prediction.

The protein G domain B1 dataset used here has a high signal-to-noise ratio of about 35x (measurement range relative to the median experimental error of fitness estimates). Assays based on cellular growth or phage display typically have smaller measurement ranges (Araya et al., 2012; Fowler et al., 2010; Li et al., 2016; Matreyek et al., 2017; Starita et al., 2013; Starita et al., 2015; Weile et al., 2017), thus in tendency smaller signal-to-noise ratios, which might render the majority of variants unsuitable for negative epistasis classification. We thus tested how our prediction strategy performs when only positive epistasis information is available. Using *combined scores* we find that accurate prediction of protein structure is still feasible with positive epistatic information only $\langle C\alpha - RMSD \rangle = 3.4\mathring{A}$). This is mainly due to the prediction of beta sheet interactions, as the accuracy drop is larger if only tertiary contacts and secondary structure element prediction are used to model the domain ($\langle C\alpha - RMSD \rangle = 6.4\mathring{A}$). Interestingly, using only negative epistasis information results in a complete loss of predictive power for the structure of the protein domain.

Furthermore, we tested the effect of a lower sequencing coverage. The dataset used here was produced from 600 million reads, roughly the equivalent of four flow cell lanes on an Illumina HiSeq sequencer (Olson et al., 2014). Because the number of double mutant variants scales quadratically with the protein length assayed, it is important to understand the effect of sequencing depth on prediction accuracy in order to assess whether scaling the approach to longer proteins is feasible. We artificially down-sampled the sequencing read coverage of the datasets by a factor four, thus doubling the error of fitness estimates ("noisy dataset"). The effect on accuracy was minor if using both positive and negative epistatic values as well as predicting beta sheet interactions $\langle C\alpha - RMSD \rangle = 2.3\mathring{A}$), while the negative impact on accuracy increased if using only positive epistasis information or omitting restraints for beta sheet interactions ($\langle C\alpha - RMSD \rangle = 4.1\mathring{A}$ and $\langle C\alpha - RMSD \rangle = 5.6\mathring{A}$, respectively). We note that the reason for minor effects of down-sampling sequencing read coverage might come from the fact

that for most variants the error from the fitness measurements itself might be the dominant
source of uncertainty. Unfortunately, for the present dataset we cannot assess this due to the
lack of replicate information. This does not affect the conclusions about the effects of lower
sequencing coverage, but highlights the importance of performing the fitness experiments in
replicate.

Finally, we tested the effect of a lower coverage of double mutants in the dataset. Fewer double
mutants will increase the uncertainty in estimating epistatic fractions per position pair, but
limiting the number of double mutants constructed in an experiment would also allow to assay
longer proteins. The present dataset was constructed by NNS-mutagenesis, i.e. replacing each
wild-type amino acid with all other 19 amino acids. Synthesis strategies relying on doping or
error-prone template replication can effectively only give rise to a fraction of all possible amino
acid changes (Araya et al., 2012; Fowler et al., 2010; Starita et al., 2013; Starita et al., 2015;
Weile et al., 2017). We thus considered only those single amino acid mutations in the dataset
that can be reached by one mutation in the nucleotide sequence, thereby lowering the coverage
of single mutants to 31% and the coverage of double mutants to 10% ("doped dataset"). Again,
we find only a minor effect on accuracy if using the full epistatic information and predicted beta
sheet interactions ($\langle C\alpha - RMSD \rangle = 2.1 \text{Å}$), but greater negative effects if using only positive
epistasis information or omitting restraints for beta sheet interactions ($\langle C\alpha - RMSD \rangle = 5.6 \text{Å}$ and
$\langle C\alpha - RMSD \rangle = 5.1 \text{Å}$, respectively).

Supplemental Figures

Supplemental Figure 1

1057

Figure S1: Deep mutational sequencing data processing

A. Distribution of fitness values for single and double mutant variants. Lower peak in distributions indicates lower limit of fitness assay measurement range (see Methods).

B. Two-dimensional variant density showing relationship between variant fitness and their error estimates. Red line shows *loess* fit of data. Dashed horizontal line indicates median error estimate. Fitness values and their error estimates are linked due to common sequencing read count dependencies.

C. Two-dimensional variant density showing dependency of fitness values on sequencing read counts in input library. For variants with too low coverage in the input library low

fitness values cannot be accurately estimated. Red line shows sequence read cutoff

used for variant inclusion (93% of variants included for downstream analysis). Horizontal

dashed line indicates lower limit of fitness assay measurement range.

D. Reliable quantification of positive and negative epistasis is limited to subsets of the data.

Upper plot shows two-dimensional double mutant variant density in single mutant fitness

space. Red outline shows single mutant fitness space that enables positive epistasis

quantification. Yellow outline shows single mutant fitness space that enables negative

epistasis quantification. Lower plot shows an example slice through fitness landscape at

single mutant fitness = -1.5. Red, grey and yellow curves show slices through quantile

fitness surfaces of 95th percentile, median and 5th percentile, respectively (see Figure

2A).   Diagonal   dashed   line   shows

$double\ mutant\ fitness\ =\ single\ mutant\ 2\ fitness\ –\ 1.5$ (expected fitness = observed

fitness). Horizontal dashed lines give lower limit of fitness assay measurement range

and the 95th percentile of fitness values of variants dominated by background fitness

effects. Red and yellow boxes indicate the range that includes 99% of variants within the

slice that are suitable for positive or negative epistasis quantification, respectively.

1083

Supplemental Figure 2

1084

Figure S2: Variant number distributions across position pairs

A. Upper panel: Position-wise distributions of single mutant fitness values. Lower panel: Distributions of the number of double mutant variants for each position pair a position is involved in. Shown are distributions for all variants (blue), i.e. coverage after quality control (93% of double mutants), variants suitable to positive epistasis quantification (red) and variants suitable to negative epistasis quantification (yellow). Note how number of double mutant variants suitable to epistasis quantification relates to single mutant fitness distributions shown in upper panel.

B. Distribution of number of double mutant variants across all position pair. Colors as in panel A.

1095

Supplemental Figure 3

averaged epistatic
fraction
· 0.2
● 0.4
● 0.6
● 0.8

directionality:
● similar
● positive > negative
● positive < negative

1096

Figure S3: Positive and negative epistasis enrichments across position
pairs

Hierarchical cluster analysis of epistatic fraction patterns. Positions are clustered
according to the Euclidean distance of their mean epistatic fractions (weighted average
of positive and negative epistatic fractions, weights are inverse variances of fractions in
resampling runs) to all other positions. Note that directionality of interactions (positive or
negative fractions more significant) was not used for clustering but only marked post-
analysis. Clustering shows two highly interconnected subsets of positions that interact
mostly positively or negative with each other and other non-subset positions.

1106

Figure S4: Secondary structure and beta sheet prediction from *epistasis* and *association scores*

A. Secondary structure element detection from *epistasis* (upper panel) and *association scores* (lower panel). Lines give position-wise p-value of secondary structure propensity; dashed lines indicate data was not used (p-value for other structure element lower); boxes indicate final predictions, lower limit of box indicates p-value of final predictions (limited to minimum of 10^{-4}). Secondary structure elements in crystal structure 1pga is shown for comparison on top.

B. Beta sheet interaction predictions from *epistasis* (upper panel) and *association scores* (lower panel). Plots show matrix of combined p-values for extended (beta sheet) seeds,

1118   the final predictions and the reference elements in the crystal structure (lower triangle –
1119   parallel beta sheets, upper triangle – anti-parallel beta sheets, diagonal – secondary
1120   structure elements).

1121   C. Precision and recall for position-wise secondary structure element and beta sheet
1122   interaction predictions from panel B.

1123

Supplemental Figure 5

1124

Figure S5: *Ab initio* folding of protein G domain B1

A. Template modeling score of top 5% structural models compared to crystal structure
1pga and the dependency on number of long-range contacts used. Negative control –
only restraints for secondary structure predicted by PSIPRED. Positive control – 0.5-2*L
ture contacts (random subset), secondary structure elements and beta sheet interactions
from crystal structure. All other: all restraints – top 0.5-2*L contacts, secondary structure
element and beta sheet interaction predictions from the three interaction scores, as
indicated by color.

B. Dependency of accuracy ($C\alpha$ root mean squared deviation) of top 5% of structural
models on number of contacts used. Colors as in panel A.

1135