Dynamic and causal contribution of parietal cortex to perceptual decisions during category learning

Lin Zhong[1,2], Yuan Zhang[1,2], Chunyu A. Duan[1], Jingwei Pan[1], Ning-long Xu[1,2]*

[1] Institute of Neuroscience, State Key Laboratory of Neuroscience, CAS Center for Excellence in Brain Science and Intelligence Technology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, China.

[2] University of Chinese Academy of Sciences, Shanghai 200031, China.

*Correspondence to: Ning-long Xu, xunl@ion.ac.cn

Making perceptual decisions to classify unknown sensory stimuli into behaviorally relevant categories, known as category learning, is a fundamental cognitive function[1–4]. The posterior parietal cortex (PPC), although has been intensively studied for its role in perceptual and categorical decision-making[5–10], as well as in many other cognitive functions[11–17], its functional significance remains controversial due to lack of causal effects on choice behaviors[10,18–21]. Here we show that PPC neurons exhibit characteristic dynamics of category learning, and are causally involved specifically in the decision process of categorizing new sensory stimuli, which is attributable to the PPC to sensory cortex feedback circuits. We trained mice to perform an auditory category learning task, where mice made categorical decisions classifying new tone stimuli into learned categories. Using *in vivo* two-photon imaging tracking the activity of the same population of PPC neurons over learning, we found that PPC neurons show dynamic representations adaptively classifying new stimuli while maintaining stable

representation for learned categories. Inactivation of PPC using pharmacology or optogenetics led to significant impairment in categorical decisions for new stimuli but not for well-learned stimuli. In addition, PPC inactivation led to a stronger bias of choices by previous trials, revealing an additional causal contribution of PPC counterbalancing recent history bias. Circuit-level perturbations using chemogenetics to silence the PPC to dorsal auditory cortex projections also impaired categorical decisions during learning. Our results reveal a dynamic and causal role of PPC circuits in decision-making when categorizing new sensory stimuli, which not only provide a potential reconciliation to the widespread controversies, but also provide a circuit-level mechanism for the key cognitive function of perceptual classification of new sensory information.

We trained head-fixed mice to perform a two-alternative-forced-choice (2AFC) auditory discrimination task using two exemplar tones (training stimuli, 2 octaves apart, 8 kHz and 32 kHz; **Fig. 1a**). Mice report their perceptual decisions on tone stimuli as "low" or "high" frequency by licking the left or right lick-port (**Fig. 1a**; see **Extended Data Fig. 1** and **movie 1**). We then introduced new "test" tones in randomly interleaved trials (30%) to examine whether mice can make categorical judgements on new sensory stimuli based on prior trainings (**Fig. 1a**). Remarkably, mice exhibit sharp categorical decisions for the test stimuli (**Fig. 1b, c**), indicated by the peak discriminability (the slope value at the categorization boundary) significantly greater than that for an otherwise linear relation between choice and tone frequency (**Fig. 1c, d**). Mice's subjective categorization boundaries (points of subjective equality, PSE) are distributed in close proximity of experimentally defined category boundary

(16 kHz; **Fig. 1e**; also see **Extended Data Fig. 1e**). These data indicate that mice are able to make categorical decisions to classify new tone stimuli as "high" or "low" based on previous trainings, a key process in category learning[3].

Although the dynamics of PPC activity has been extensively studied[9,11,16,22,23], the precise relationship between PPC activity and specific learning process is still unclear. Here we investigate the relationship between PPC activity dynamics and perceptual category learning, using chronic *in vivo* two-photon calcium imaging from PPC neurons across different sessions/days. Genetically encoded calcium indicator GCaMP6s[24] was expressed in PPC of left hemisphere using AAV vectors to indicate the intracellular calcium dynamics, followed by implantation of a chronic imaging window[25]. The same population of layer 2/3 neurons were imaged during task performance repeatedly across days starting from the session when the test stimuli were first introduced (**Fig. 2a;** see **Methods** and **Extended Data Fig. 2**). Consistent with previous studies[11,17,22,26], we found that PPC neurons show complex response selectivity to multiple behavioral variables (**Fig. 2b**; **Extended Data Fig. 3** and **Table 1**).

Here we ask whether the PPC neuronal dynamics exhibits characteristics of category learning. With the progress of learning, new sensory stimuli are classified into existing categories, such that the perceptual distinction between new sensory stimuli and exemplars of the same category tend to diminish. We thus focused on whether PPC neuronal activity could distinguish the test stimuli (new sensory information) and the training stimuli (prior knowledge) of the same category, and how such distinction changes with learning. The high performance of perceptual categorization in mice for the test stimuli on day 1 (the first session introducing test stimuli; **Fig. 1**) suggests that significant category learning likely

occurred rapidly. Indeed, we found rapid changes in PPC neuronal responses to test stimuli across the first two sessions (days) following introducing test stimuli. On day 1, a significant fraction of neurons responded more strongly to test stimuli than to training stimuli, however, such difference was markedly reduced on day 2 (**Fig. 2c, d**; also see **Extended Data Fig. 4**).

To quantify the distinction in neuronal responses between test stimuli and training stimuli of the same category, we use receiver operator characteristic (ROC) analysis to define a selectivity index (SI), with negative SI indicating stronger preference to training trials, and positive SI, stronger preference to testing trials (see Methods)[27]. For learning, the activity in the entire trial is likely to be important, we therefore computed the SI of each neuron in an imaging field for every imaging frame across the time course of the trial (**Fig. 2e**). We found that on day 1, when the test stimuli were first introduced, more neurons significantly distinguish test stimuli and training stimuli, with stronger preference to test stimuli. However, on day 2 and day 3, the preference to test stimuli were largely reduced. This trend is also evident over the entire neuronal population (757 neurons from 6 mice), where the population distribution of SIs also shifted from significantly positive values on day 1 ($P<0.001$, signed Wilcoxon rank sum test) toward zero on day 2 (**Fig. 2f**). For neurons showing significant preference to test stimuli on day 1, the neuronal discrimination (measured by the area under ROC curve) between test stimuli and training stimuli of the same category also significantly decreased in day 2 and 3 (**Fig. 2g**), indicating that the distinction between the test stimuli and training stimuli of the same category was largely reduced. These results indicate that in parallel to perceptual categorization, PPC neuronal representations exhibit the characteristic dynamics of category learning, incorporating new sensory stimuli into existing categories.

During learning, the circuits not only need to dynamically integrate new information but also need to maintain a stable representation of prior knowledge. We thus ask whether the representation of established "high" and "low" categories in PPC population is relatively stable. First, we found that the number of neurons selective to training stimuli was almost unchanged (**Extended Data Fig. 5e;** 21% in day 1 vs. 20% in day 2). Second, using ROC analysis, we computed the selectivity index of each neuron for "high" vs. "low" frequency categories for all the neurons (757 neurons) over the time course of the behavioral trial across consecutive days (**Fig. 2h**; see **Methods**). We sort all the neurons according to their SI values on day 1. We found that across the entire population and trial time course, neuronal selectivity for "high" and "low" categories is highly preserved across days, as indicated by the resemblance of the profiles of the SIs between the two consecutive days (**Fig. 2h**; Pearson's $r = 0.83$). The mean SI values within the time epoch of stimulus and choice (1 s time window after stimulus onset) are also strongly correlated between two consecutive days (**Fig. 2i**; $r = 0.91$). Thus, the neuronal selectivity for established stimulus categories is stable across days. Third, we asked whether the population coding for the established categories was also stable. We used a linear classifier[28] to examine the population decoding accuracy of simultaneously imaged neurons in each imaging field for categorical discrimination over the trial time. We used the population activity in 80% of training trials on day 1 as training set to predict either the rest of training trials in day 1, or the training trials in day 2 and day 3 respectively (**Fig. 2j)**. We found that the decoding accuracies around the decision time (first lick) were not significantly different for day 1, day 2, and day 3 (**Fig. 2j**), indicating that the population coding for the stimulus categories was stable across days over category learning. Taken together, these results show that PPC neurons exhibit characteristic dynamics of

category learning, adaptively classifying new sensory stimuli into existing categories, while maintaining a stable representation for prior learned categories (**Fig. 2k**).

Next, we sought to determine whether PPC activity is important in the decision process during category learning. Our imaging and behavioral data suggest that the learning occurred primarily when the test stimuli were initially introduced in day 1, therefore, it is likely that PPC activity may play a critical role during this period. To test this, we reversibly silenced PPC by bilateral microinjection of muscimol, a selective $GABA_A$ receptor agonist, when the new test stimuli were first introduced (**Fig. 3a, c**). In contrast to previous reports in rodents[10,17,19,29] and primates[18,20], we found that mice's decisions on newly introduced test stimuli were markedly impaired, comparing to control mice microinjected with saline also on day 1. The impairment is measured as a significant reduction in the peak discriminability (slope at the category boundary; saline, 1.46 ± 0.16; muscimol, 0.97 ± 0.09; $P < 0.05$, Wilcoxon rank sum test; **Fig. 3d, f**).

Since muscimol inactivation does not have the temporal resolution to distinguish whether it was during training trials or during testing trials that silencing PPC led to impaired categorization, we employed optogenetics to inactivate PPC in a more temporally controlled fashion[21]. An adeno-associated virus (AAV) encoding Cre-inducible channelrhodopsin 2 (ChR2)[30] was injected bilaterally in PPC of VGAT-Cre mice[31] to express ChR2 in GABAergic inhibitory neurons. Glass optical windows were implanted above PPC of both hemispheres following virus injection. Collimated blue laser beams were used to illuminate the glass window to activate ChR2 in GABAergic neurons to inhibit the illuminated cortical areas (**Fig. 3b, c;** see **Extended Data Movie 1**)[21]. During task performance, photoinhibition was applied in all testing trials, but only in 10% of training trials for comparison, leaving

majority (90%) of training trials unaffected. We found that this manipulation also significantly reduced mice's categorization performance for the new tone stimuli (control, 2.43 ± 0.46; photoinhibition, 1.17 ± 0.37; $P < 0.05$, Wilcoxon rank sum test; **Fig. 3e, f**), indicating that PPC activity during testing trials rather than during training trials is required for categorical decisions on the new test stimuli. Moreover, we found that when photoinhibition of PPC was applied in 50% of the testing trials, mice's categorization performance was not significantly different in photostimulated trials comparing to that in control trials (**Extended Data Fig. 7**), suggesting that PPC activity in partial of the testing trials on day 1 was sufficient for category learning of the newly introduced tone stimuli.

These results, together with our imaging data (**Fig. 2**), point to the possibility that PPC activity is specifically required only during category learning for new stimuli, but not after learning. Consistent with this idea, for the training stimuli, neither muscimol silencing (**Fig. 3h**), nor photoinhibition (**Fig. 3i)** influenced mice's performance (**also see Extended Data Fig. 8**). To further test this possibility, we performed bilateral PPC perturbation in a separate cohort of mice after the test stimuli had been well-experienced following continued training for two additional sessions (>1000 trials; **Fig. 3j**). We found that when the test stimuli were well-experienced, inactivation of PPC did not significantly influence mice's categorization performance (**Fig. 3k-m**), agreeing with previous studies in rodents and non-human primates assessing the effect of PPC inactivation on choice behavior, where the stimuli were likely also well-learned[10,18,19,21]. Taken together, these data indicate that PPC is causally involved in categorical decisions specifically on new sensory stimuli during category

learning, but not after learning when the stimuli have been already incorporated in to existing categories.

Decision-making in humans and animals is often biased by recent behavioral history, which usually produce a negative impact on performance in complex and unpredictable settings[16,17,32–36]. It has been reported that PPC activity reflected both accumulated behavioral history[16,17,22,37] and systematic bias from long-term training history[12]. Meanwhile, our imaging data, as well as previous studies[5,8,20,37], show that even after learning, PPC neurons continue to encode stimulus category and choice information. To understand how the accumulated prior knowledge encoded in PPC may influence the bias from recent behavioral history, we examined the effect of PPC inactivation on trial-by-trial history bias during the auditory categorization task. We quantified the influence of the preceding training trials on choices for testing stimuli by comparing mice's performance on test stimuli when the stimulus in the immediate preceding training trial was of the same vs. different category as the current test stimuli. Under control condition, preceding training trials did not significantly bias mice's performance on the testing trials (**Fig. 4b,** 77.3 ± 4.6% for same category vs. 74.3 ± 3.2% for different category, P = 0.32), consistent with a previous study, in cases where decisions did not depend on short-term memory[17]. However, when PPC was inactivated by either muscimol or photoinhibition, preceding training trials significantly biased mice's choices on test stimuli *towards* the preceding categories (**Fig. 4a**, muscimol, same 71.8 ± 3.4% vs. different 63.5 ± 3.7%, P = 0.017; photoinhibition, same 70.3 ± 2.4% vs. different 55.9±3.2%, P = 0.003.) Thus, whereas mice's decisions were minimally influenced by preceding events when PPC was intact, inactivation of PPC led to a significant bias by short-term behavioral history (**Fig. 4c, d**). Moreover, when the newly introduced test stimuli

became well-experienced after continued training, inactivation of PPC no longer resulted in significant bias by preceding training trials (**Extended Data Fig. 12e**, **f**). These results indicate that PPC activity counterbalances the short-term history bias when categorizing novel sensory stimuli.

To simultaneously examine PPC's contributions to category learning (**Fig. 3**) and to short-term history bias (**Fig. 4a, c**), we modeled animals' performance in control and silencing conditions using a series of logistic regression models (Generalized Linear Mixed-Model, GLMM; **Fig. 4e**; also see Methods)[34]. Our models confirmed the lack of history bias effect in control data, in that including of previous choice as a regressor term did not significantly improved the model prediction of the behavioral data (**Fig. 4f**). The models also confirmed the effect of PPC inactivation on changing the psychometric slopes, in that including of PPC silencing as a regressor significantly improved the model prediction (**Fig. 4g**; also see **Extended Data Fig. 13a-d**). Interestingly, when including an interaction term between previous choice and the silencing condition as a regressor, the model predicted animals' behavior significantly better comparing to the model that only quantified the effect of silencing on the psychometric slope changes (**Fig. 4h**; see **Extended Data Fig. 14a-c**), indicating that the effect of PPC silencing on short-term history bias is in addition to its effect on categorization slopes. Thus, PPC causally contributes to the decision process in category learning by both influencing categorization slopes and counterbalancing short-term history bias when categorizing novel stimuli.

PPC neurons consist of diverse populations interconnected with many brain regions[38], and are involved in many behavioral functions[5–7,11,13,39]. It is likely that for a given task, specific circuits interconnecting PPC and task-specific brain regions are critically involved.

We thus further examined the circuit-level specificity of the causal role of PPC neurons in auditory categorical decision-making by manipulating PPC to auditory cortex projections using chemogenetics. We first determined the projection patterns between PPC and the auditory cortex by expressing green fluorescent proteins (GFP) in PPC, and red fluorescent proteins (tdTomato) in the primary auditory cortex (AUDp), using AAV vectors. We found that PPC receives projections from AUDp while sending projections to the dorsal portion of auditory cortex (AUDd; **Fig. 5a**), implying an interconnected circuit between PPC and the auditory cortex (**Fig. 5b**). We next expressed hM4Di, a designer receptor exclusively activated by designer drug (DREADD), in PPC of both hemispheres using an AAV vector (AAV-Syn-hM4Di). On day 1 (test tones first introduced), we injected Clozapine-N-oxide (CNO) to bilateral AUDd to silence the projections from PPC prior the behavioral session (**Fig 5c**). As control, in a separate group of mice we injected AAV-Syn-mCherry in PPC, and then injected CNO to AUDd during task performance on day 1. We found that comparing to the control group, silencing PPC to AUDd projections impaired mice's categorization performance for the new tone stimuli, as indicated by a significant reduction in the slopes of psychometric functions (**Fig. 5d**). Quantification using GLMM test also shows a significantly better prediction of the model with the term of projection silencing than without the term (**Fig. 5e**). These results indicate that the specific projections from PPC to secondary auditory cortex play a critical role in the auditory categorical decisions on new sensory stimuli.

The neural basis for how organisms integrate novel sensory information with prior knowledge to assign behavioral meanings to physical world remains poorly understood. Here we provide functional and causal evidence to support a critical role of PPC circuits in this process. We show that PPC neurons exhibit characteristic dynamics related to category

learning, adaptively incorporating new sensory information while maintaining stable representations for learned category to choice mappings, a balance in flexibility and stability. Importantly, we provide, to our knowledge, the first evidence supporting that PPC causally contributes to perceptual decision-making in categorizing new sensory information, a key process in category leaning. Furthermore, at the circuit level, we identified a necessary role of PPC to auditory cortex feedback projections, providing a critical foundation for further dissection of circuit-level mechanisms of category learning. After the stimuli were well-learned, PPC continued to represent stimulus categories (**Fig. 2**) and behavioral choices[5,8,9,18,20]. This is likely to be important for guiding future decisions on new sensory stimuli, and reducing biasing effects from random factors (**Fig. 3**). Some of recent studies showed that silencing PPC impaired visually guided[10,11,26,29] but not auditory guided[10,19,29] choice behaviors in rodents, while in primates, silencing parietal cortex did not influence either visual motion[18] or self-motion[20] guided choice behavior, suggesting that that PPC may also play certain visual-specific functions in rodents. Our results, however, support a more general functional role of PPC in perceptual decision-making when facing novel conditions. More recent studies in rodents also using well-trained stimuli showed increased performance in tasks requiring short-term memory or working memory following inactivation of PPC[16,17], likely attributable to an interference with short-term memory by the effect of PPC on recent choice history, but not an direct effect on perceptual decision-making *per se*. This is consistent with our observation that PPC activity counterbalances recent history bias. Our results thus provide a framework for mechanistic understanding of neuronal circuit basis of decision-making integrating new sensory information with prior knowledge in a dynamic environment.

References:

1.   Ashby, F. G. & Maddox, W. T. Human Category Learning. *Annu. Rev. Psychol.* **56,** 149–178 (2005).

2.   Seger, C. A. & Miller, E. K. Category Learning in the Brain. *Annu. Rev. Neurosci.* **33,** 203–219 (2010).

3.   Ohl, F. W., Scheich, H. & Freeman, W. J. Change in pattern of ongoing cortical activity with auditory category learning. *Nature* **412,** 733–736 (2001).

4.   *Categorical Perception: The Groundwork of Cognition*. (Cambridge University Press, 1990).

5.   Shadlen, M. N. & Newsome, W. T. Neural Basis of a Perceptual Decision in the Parietal Cortex (Area LIP) of the Rhesus Monkey. *J. Neurophysiol.* **86,** 1916–1936 (2001).

6.   Gold, J. I. & Shadlen, M. N. The Neural Basis of Decision Making. *Annu. Rev. Neurosci.* **30,** 535–574 (2007).

7.   Freedman, D. J. & Assad, J. A. A proposed common neural mechanism for categorization and perceptual decisions. *Nat. Neurosci.* **14,** 143–146 (2011).

8.   Freedman, D. J. & Assad, J. A. Experience-dependent representation of visual categories in parietal cortex. *Nature* **443,** 85–88 (2006).

9.   Hanks, T. D. *et al.* Distinct relationships of parietal and prefrontal cortices to evidence accumulation. *Nature* **520,** 220–223 (2015).

10.   Raposo, D., Kaufman, M. T. & Churchland, A. K. A category-free neural population supports evolving demands during decision-making. *Nat. Neurosci.* **17,** 1784–1792 (2014).

11.	Harvey, C. D., Coen, P. & Tank, D. W. Choice-specific sequences in parietal cortex during a virtual-navigation decision task. *Nature* **484,** 62–68 (2012).

12.	Yang, T. & Shadlen, M. N. Probabilistic reasoning by neurons. *Nature* **447,** 1075–1080 (2007).

13.	Andersen, R. A. & Buneo, C. A. Intentional Maps in Posterior Parietal Cortex. *Annu. Rev. Neurosci.* **25,** 189–220 (2002).

14.	Andersen, R. A. & Cui, H. Intention, Action Planning, and Decision Making in Parietal-Frontal Circuits. *Neuron* **63,** 568–583 (2009).

15.	Cui, H. & Andersen, R. A. Posterior Parietal Cortex Encodes Autonomously Selected Motor Plans. *Neuron* **56,** 552–559 (2007).

16.	Akrami, A., Kopec, C. D., Diamond, M. E. & Brody, C. D. Posterior parietal cortex represents sensory history and mediates its effects on behaviour. *Nature* **554,** 368–372 (2018).

17.	Hwang, E. J., Dahlen, J. E., Mukundan, M. & Komiyama, T. History-based action selection bias in posterior parietal cortex. *Nat. Commun.* **8,** 1242 (2017).

18.	Katz, L. N., Yates, J. L., Pillow, J. W. & Huk, A. C. Dissociated functional significance of decision-related activity in the primate dorsal stream. *Nature* **535,** 285–288 (2016).

19.	Erlich, J. C., Brunton, B. W., Duan, C. A., Hanks, T. D. & Brody, C. D. Distinct effects of prefrontal and parietal cortex inactivations on an accumulation of evidence task in the rat. *eLife* **4,** e05457 (2015).

20.	Chen, A., Gu, Y., Liu, S., DeAngelis, G. C. & Angelaki, D. E. Evidence for a Causal Contribution of Macaque Vestibular, But Not Intraparietal, Cortex to Heading Perception. *J. Neurosci.* **36,** 3789–3798 (2016).

21.   Guo, Z. V. *et al.* Flow of Cortical Activity Underlying a Tactile Decision in Mice. *Neuron* **81,** 179–194 (2014).

22.   Morcos, A. S. & Harvey, C. D. History-dependent variability in population dynamics during evidence accumulation in cortex. *Nat. Neurosci.* **advance online publication,** (2016).

23.   Driscoll, L. N., Pettit, N. L., Minderer, M., Chettih, S. N. & Harvey, C. D. Dynamic Reorganization of Neuronal Activity Patterns in Parietal Cortex. *Cell* **170,** 986–999.e16 (2017).

24.   Chen, T.-W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499,** 295–300 (2013).

25.   Xu, N. *et al.* Nonlinear dendritic integration of sensory and motor input during an active sensing task. *Nature* **492,** 247–251 (2012).

26.   Goard, M. J., Pho, G. N., Woodson, J. & Sur, M. Distinct roles of visual, parietal, and frontal motor cortices in memory-guided sensorimotor decisions. *eLife* **5,** e13764 (2016).

27.   Feierstein, C. E., Quirk, M. C., Uchida, N., Sosulski, D. L. & Mainen, Z. F. Representation of Spatial Goals in Rat Orbitofrontal Cortex. *Neuron* **51,** 495–507 (2006).

28.   Hung, C. P., Kreiman, G., Poggio, T. & DiCarlo, J. J. Fast Readout of Object Identity from Macaque Inferior Temporal Cortex. *Science* **310,** 863–866 (2005).

29.   Licata, A. M. *et al.* Posterior Parietal Cortex Guides Visual Decisions in Rats. *J. Neurosci.* **37,** 4954–4966 (2017).

30.   Zhang, F., Wang, L.-P., Boyden, E. S. & Deisseroth, K. Channelrhodopsin-2 and optical control of excitable cells. *Nat. Methods* **3,** 785–792 (2006).

31.   Chao, H.-T. *et al.* Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature* **468,** 263–269 (2010).

32.   Bertelson, P. Serial Choice Reaction-time as a Function of Response versus Signal-and-Response Repetition. *Nature* **206,** 217–218 (1965).

33.   Gold, J. I., Law, C.-T., Connolly, P. & Bennur, S. The Relative Influences of Priors and Sensory Evidence on an Oculomotor Decision Variable During Perceptual Learning. *J. Neurophysiol.* **100,** 2653–2668 (2008).

34.   Busse, L. *et al.* The Detection of Visual Contrast in the Behaving Mouse. *J. Neurosci.* **31,** 11351–11361 (2011).

35.   Abrahamyan, A., Silva, L. L., Dakin, S. C., Carandini, M. & Gardner, J. L. Adaptable history biases in human perceptual decisions. *Proc. Natl. Acad. Sci.* **113,** E3548–E3557 (2016).

36.   Akaishi, R., Umeda, K., Nagase, A. & Sakai, K. Autonomous Mechanism of Internal Choice Estimate Underlies Decision Inertia. *Neuron* **81,** 195–206 (2014).

37.   Runyan, C. A., Piasini, E., Panzeri, S. & Harvey, C. D. Distinct timescales of population coding across cortex. *Nature* **advance online publication,** (2017).

38.   Zingg, B. *et al.* Neural Networks of the Mouse Neocortex. *Cell* **156,** 1096–1111 (2014).

39.   Bisley, J. W. & Goldberg, M. E. Attention, Intention, and Priority in the Parietal Lobe. *Annu. Rev. Neurosci.* **33,** 1–21 (2010).

Supplementary Information is available in the online version of the paper.

Acknowledgments: We thank R. Egnor for advice on auditory behavioral apparatus; A. Kerlin for communication on behavioral control software; N. Andersen for pilot optogenetic experiments; T.T. Zhou for technical support; L. Cui, S. Tang and J. Xiao for helping on

behavior training; Y. Xin for discussion on data analysis; and M. M. Poo for comments on the manuscript.

Funding: This work was supported by National Natural Science Foundation of China (grant No. 31571081); the Strategic Priority Research Program of the Chinese Academy of Sciences (grant No. XDB02040008); Key Research Program of Frontier Sciences, CAS, Grant No. QYZDB-SSW-SMC045; the Youth Thousand Talents Plan (to N.L.X.). C.A.D. is supported by the Simons Collaboration on the Global Brain Postdoctoral Fellowship and the CPSF-CAS Joint Foundation for Excellent Postdoctoral Fellows.

Author contributions: L.Z. and N.L.X. conceived the project and designed the experiments. L.Z. performed all the experiments and data analysis. Y.Z. collected part of the behavioral data. L.Z. and C.A.D. conceived and performed the GLMM analysis. L.Z., J.P. and N.L.X. developed the hardware and software of the behavior and imaging system. L.Z. and N.L.X. wrote the manuscript with contributions from C.A.D.

Competing interests: The authors declare no competing financial interests.

Materials & Correspondence: Correspondence and requests for materials should be addressed to N.L.X. (xunl@ion.ac.cn).

Data and materials availability: All data to understand and assess the conclusions of this study are available in the main text or supplementary materials. All the original behavioral, optogenetic, imaging, histochemical data and analysis code are archived in the Institute of Neuroscience, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences.

Figure 1 | Perceptual categorization of auditory stimulus in head-fixed mice. a,

Schematic showing the configuration of behavioral paradigm and trial structure. **b**,

Psychometric function from an example behavioral session where the test stimuli were first

introduced. **c**, Slopes of the psychometric function and hypothetical linear relation between

stimulus and behavioral choice shown in **b**. The peak of the slope of the psychometric

function (black) around the point of subjective equality (PSE) indicates perceptual

categorization. The slope of the linear function (grey) shows a constant value. **d**, Scatter plot

of group data comparing peak psychometric slopes (1.66 ± 0.16) and linear relation slopes

(0.45 ± 0.01) within each animal (*P* < 0.0001, n = 25 mice). **e**, Histogram showing

distribution of the PSE of 25 animals from the first session introducing test stimuli (PSE =

16.5 ± 0.3 kHz). Red error bars indicate median and interquartile range.

Figure 2 | PPC representations show characteristic dynamics of category learning. a,

Schematic and images showing in vivo two-photon imaging from PPC populations using

GCaMP6s. The same imaging fields were repeatedly imaged across days. **b,** Example Ca²⁺

signal traces of PPC neurons showing high signal to noise ratio with diverse selectivity and

rich dynamics. **c,** Ca²⁺ signal of an example PPC neuron imaged during behavioral sessions

on consecutive days. Upper, color indicates % $\Delta F/F_0$. Blocks of trials presenting training and test stimuli of low and high frequency categories were segregated. Trials within each block were aligned by stimulus onset time and sorted by time of first answer lick. Lower, mean Ca^{2+} signal traces averaged from blocks of trials in the upper panel. Color corresponding to tone frequencies. **d,** Mean response amplitudes in 1 s window after stimulus onset averaged from different blocks shown in **c**. Error bars, s.e.m. **e**, Selectivity index (SI) between training and test stimuli across time based on ROC analysis using the neuronal responses as shown in **c**, for all neurons in an example imaging field. Positive values of SI represent preference to test stimuli and negative value for training stimuli. **f**, Histogram showing distributions of SI values (mean value in 1 s window after stimulus onset) for training vs. test stimuli from the total of 757 neurons across two consecutive days. Significant preference to test stimuli on day 1 ($P < 10^{-23}$, Wilcoxon signed rank test), but to training stimuli on day 2 ($P < 10^{-6}$, Wilcoxon signed rank test). **g,** Averaged area under ROC curve, discriminating training and test stimuli, across consecutive days for neurons with significant preference to test stimuli on day 1. Error bars, s.e.m. **h**, Color raster plot showing index of selectivity between high and low frequency categories across trial time, for all neurons as in **f** across consecutive days. Neurons are sorted according to SI values on day 1. **i**, Scatter plot of the mean SI values in the 1 s trial time window after stimulus onset as in **h**. *r*, correlation coefficient. *Slope*, slope of the linear regression line (red). **j,** Decoding accuracy across trial time for high and low training tones by population activity of simultaneously imaged neurons in each imaging field using a linear classifier. The classifier was trained using neuronal activity in 80% of training trials on day 1 to decode stimulus categories on consecutive 3 days. Insert, mean decoding accuracies in a 1 sec time window following stimulus onset. **k**, Schematic showing a conceptual model for differential dynamic representations in PPC reflecting category learning.

Figure 3 | PPC activity is necessary for categorical decision-making on new sensory stimuli. a, Schematic and histology image showing the locations of muscimol or saline injection in PPC. Red color in the histology image is the fluorescence of co-injected red beads to mark the injection sites. **b**, Schematic and histology image showing the locations of photo stimulation and viral expression of ChR2-mCherry in GABAergic neurons in bilateral PPC of Vgat-Cre mice. Red fluorescence represents mCherry expression. **c**, Schematic showing experimental schedules for PPC inactivation and control experiments on the first day introducing test stimuli. Symbols represent the 4 types of experiments as indicated in **a** and **b**. Frequencies of tone stimulation at different stages were indicated. **d**, Psychometric functions (upper) and corresponding slopes (lower) from an example muscimol injection session and control session as indicated in **a** and **c**. Error bars, 95% confidence interval. See Extended Data Fig. 9 for individual cases. **e**, Similar as in **d** for an photoinhibition and control sessions as indicated in **b** and **c**. See Extended Data Fig. 10 for individual cases. **f**, Summary of grouped data of peak psychometric slopes from experiments indicated in **c**. Muscimol, $n = 13$, vs. saline, $n = 14$, $P < 0.05$; photoinhibition, $n = 7$, vs. mask only, $n = 6$, $P < 0.05$. Error bars, s.e.m. **g**, Summary of PSE for experiments indicated in **c** (muscimol vs. saline, $P = 0.68$; photoinhibition with mask vs. mask only, $P = 0.73$; Error bars, s.e.m.). **h**, Summary of mice's performance (percent correct) for training stimuli, compared between muscimol injection and control experiments. **i**, Similar as in **h** for photoinhibition vs. control experiments. Error bars, s.e.m. Paired *t*-test. **j**, Experimental schedules, similar as in **c**, for PPC inactivation on the third day after introducing test stimuli. **k**, Similar plot as in **e** for photoinhibition experiments indicated in **j**. See Extended Data Fig. 11 for individual cases. **l** and **m**, Summary of grouped data of peak psychometric slopes for experiments indicated in **k**.

Figure 4 | PPC activity counterbalances short-term history bias on perceptual decisions.

a, Comparison of behavioral performance on testing trials when the stimuli in preceding training trial was of different categories (diff. prev.) or the same category (same prev.) as the current trial. Mice's choices were significantly biased by the previous trial when PPC was silenced by muscimol ($P = 0.017$, $n = 13$) or by photoinhibition ($P = 0.003$, $n = 7$). **b**, Same plot as in **a**. Mice's choices were not significantly biased by previous trials under control

conditions (saline, $P = 0.14$, $n = 14$; light mask, $P = 0.32$, $n = 6$). **c**, Summary of changes in performance attributable to previous trials as in **a** and **b**. Data from a separate group of control mice without any manipulation were also included. *, $P < 0.05$; **, $P < 0.01$. Two sample t-test. Error bars, s.e.m. **d**, Schematic showing the bias effect on categorical decisions by the previous trials when PPC was silenced. **e**, Illustration of linear regression models to simultaneously quantify the effects of PPC silencing on categorization slopes and short-term history bias. **f-h**, Likelihood ratio test comparing different regression models using data from the control group for models with or without previous trial choices (**f**), using all data from both control and PPC silencing groups for models with or without the interaction term between sensory stimuli and PPC silencing (**g**), and using data from both groups for models with or without the interaction term between PPC silencing and previous choices (**h**).

Figure 5 | PPC to auditory cortex projections are necessary for categorical decision-making on new sensory stimuli.

a, Histology images showing the projection patterns between PPC and the auditory cortex. Left, red fluorescence indicates the site of injection of AAV-syn-mCherry in the primary auditory cortex. Green fluorescence in the dorsal auditory cortex indicates axons projected from PPC. Insert, showing site of injection of AAV-syn-GFP in PPC. Right, red fluorescence indicates axons projected from the primary auditory cortex in PPC. **b**, Schematic showing the projection patterns between PPC and the auditory cortex. **c**, Schematic showing chemogenetic silencing of the projections from PPC to the auditory cortex. AAV-syn-hM4D-mCherry (experimental group) or AAV-syn-mCherry (control group) were injected in bilateral PPC. CNO was microinjected in bilateral auditory cortex prior the first session introducing test tone stimuli. **d**, Summary of the effect of silencing PPC to auditory cortex projections on psychometric slopes. $P < 0.05$, Wilcoxon rank sum test, n (control) = 7, n

(silencing) = 8. Error bars, s.e.m. **e**, Likelihood ratio test comparing regression models with or without projection silencing as a regressor.

Extended Data Figure 1 | Behavioral control system and auditory psychophysical task in mice.

a, Schematic diagram of custom designed behavior control hardware system for two choice auditory decision task in head-fixed mice, named as PX-Behavior System. **b**, Schematic diagram of the auditory two-alternative forced choice task in head-fixed mice. **c**, Example of auditory 2AFC task performance over the time course of training. **d**, Summary of mice performance (percent correct, mean ± s.e.m.) upon training stimuli at the beginning of training and at well-trained stage. **e**, Psychometric curves from a separate group of mice with randomly rewarded testing trials, i.e., the reward in the trials presenting intermediate tone frequencies were rewarded randomly, not according to the experimentally defined categorization boundary. Data from 5-7 continuous sessions following test stimuli were first introduced were grouped for each animal. Circles with bars, mean ± s.e.m.

Extended Data Figure 2 | Overview of population response profiles in PPC across consecutive days recorded using two-photon calcium imaging.

 a, Expression of GCaMP6s in PPC using AAV-SYN-GCaMP6s. **b**, An example imaging field across three days. **c**, Normalized mean Ca^{2+} signals from all neurons (736 neurons from 5 mice) in different trial types (training and testing, with "low" or "high" stimulus categories). Neurons were sorted by the response peak time in each trial type on day 1 (test stimuli were first introduced). Upper, data from day 1. Lower, data from day 2. Dash lines indicate stimulus onset time.

Extended Data Figure 3 | Example neurons showing diverse response patterns during auditory categorization task. Testing trials with the same categorical frequencies were grouped together. Dash lines indicate stimulus onset. Ca^{2+} signals ($\Delta F/F$) in each trial were aligned to stimulus onset, and sorted according to first lick time after stimulus onset. Traces are mean across trials.

Extended Data Figure 4 | Extra example neurons showing preferred selectivity to testing stimuli on day 1. a-d, Extra example neurons showing preferred selectivity to test stimuli reduced on day 1 but reduced in the following session.

Extended Data Figure 5 | Changes in selectivity for training vs. test stimuli in longitudinally imaged neuron populations.

a, Selectivity index for training vs. test stimuli across trial time in simultaneously imaged PPC neurons from an example imaging field across two consecutive days. Upper, data from trials presenting tones of "low" frequency category. Lower, data from trials presenting tones of "high" frequency category. Dash line, stimulus onset. **b**, Similar as in **a** from another imaging field across three consecutive days, with test stimuli first introduced on day 1. **c**, Distribution of SI for training vs. testing across four days. Data in day 1 to day 3 were from the same population of neurons, n (mice) = 4, n (neurons) = 582. Data on day 4 were from neurons in different imaging fields in the same 4 mice, n (neurons) = 1055. Only on day 1 was the population selectivity significant biased to testing trials. **c**, Percentage of neurons showing SI > 0 across 4 days following introduction of test stimuli. **d**, Selectivity index for test and training stimuli across 3 days for neurons preferring test stimui on day 1. **e**, Fraction of neurons responded more strongly (*t*-test) training stimuli vs test stimuli on day 1 and day 2.

Extended Data Figure 6 | Selectivity for "low" vs. "high" stimulus categories in PPC neuron populations across consecutive days.

a, Selectivity index of "low" vs. "high" categories in training trials over trial time in all neurons in two consecutive days. Neurons were sorted by SI values in sequential time windows on day 1. Note that the pattern of SI is preserved in day 2, even using the same neuronal sorting as in day 1. **b**, Similar in **a** for SI computed using testing trials. **c**, Correlation of SI value in the time window of 1 s after stimulus onset computed using training trials ($P < 0.0001$). **d**, Similar in (C) using data from testing trials ($P < 0.0001$).

Extended Data Figure 7 | PPC activity in partial of testing trials is sufficient for category learning

a, Psychometric curves for individual animals of the control group, expressed mCherry in PPC. Photostimulation was delivered in 50% of the testing trials, and in 10% of the training trials. Blue curve represents psychometric curves for photo stimulated trials. **b**, Same plot as in **a**, for individual animals of the photoinhibition group, with PPC silenced in 50% of the testing trials, and in 10% of the training trials. **c**, Summary of the mean impairment ratio, *(control-silence) / control*, for data shown in **a** and **b**. Control, *P* = 0.27. Partial silencing, *P* = 0.97, one sample *t*-test.

Extended Data Figure 8 | Effects of PPC inactivation on behavioral performance for well-trained tones.

a, Behavioral performance for training tones during PPC inactivation by muscimol on the first day introducing test stimuli ($n = 13$), compared with control group (injected with saline, left, $n = 14$, two-sample t-test, $P < 0.05$) and compared within group (same animals on the 2nd day when injected with saline, right, $n = 13$, paired t-test, $P > 0.05$). Bars, mean ± s.e.m. Light lines, individual mice. **b**, As in **a** for photoinhibition experiments. Left, comparison between photoinhibition group (blue light + mask, $n = 7$) with control group (mask only, $n = 6$), two-sample t-test, $P > 0.05$. Right, comparison for the same animals on day 1 with photoinhibition with day 2 with light mask. $P > 0.05$, paired t-test, $n = 7$. **c**, Behavioral performance for training tones of with muscimol silencing on the 3rd day/session following introduction of test stimuli. $P > 0.05$ between consecutive days, paired t-test, $n = 7$. **d**, Left, similar as in **c** for photoinhibition, $P > 0.05$ between consecutive days, paired t-test, $n = 5$. Right, comparison of behavioral performance for training stimuli between photoinhibition trials (10%) and control trials (90%) on day 3. $P > 0.05$, paired t-test, $n = 5$.

Extended Data Figure 9 | Individual psychometric curves from all animals used for muscimol silencing and saline injection.

a, Control group ($n = 14$). Mice received saline injections in PPC on day 1 (first introducing test stimuli). **b**, PPC silencing group ($n = 13$). Mice received muscimol injections in PPC on day 1, and saline injection on day 2. Error bars, 95% confidence interval.

Extended Data Figure 10 | Individual psychometric curves from all animals used for photoinhibition and control.

a, Control group ($n = 6$). VGAT-Cre mice received light mask only on day 1 introducing test stimuli. **b**, Photoinhibition group ($n = 7$). VGAT-Cre mice received blue laser stimulation plus light mask on day 1 and mask only on day 2. Error bars, 95% confidence interval.

Extended Data Figure 11 | Individual psychometric curves from all animals used for PPC silencing and control on day 3.

a, PPC inactivation with muscimol, $n = 7$. **b**, PPC inactivation with optogenetics, $n = 5$. Error bars, 95% confidence interval.

Extended Data Figure 12 | Previous trial history bias during PPC silencing and control conditions in the same animal across different days.

a and **b**, Difference in performance for test trials when the previous training trials were of the same or different categories, presented as scatter plot **a** and the values of difference **b**. Blue symbols, day 1 with photoinhibition ($P < 0.05$), and day 2 with mask only ($P > 0.05$). $n = 7$, paired t-test. **c** and **d**, Similar as in **a** and **b** for muscimol silencing group (day 1, muscimol, $P < 0.05$; day 2, saline, $P > 0.05$; $n = 13$, paired t-test). **e**, Effect of PPC silencing with muscimol on previous trial history bias on day 3 following introduction of testing trials. Bias is expressed as the difference values as in **b** across 3 days. **f**, As in **e** for the photoinhibition animal group. n.s., not significant.

model 1: no sensory interaction with silence term
model 2: with sensory interaction with silence term

Extended Data Figure 13 | Comparison of GLM models with or without the term of interaction between sensory stimuli and PPC silencing.

a, Visualization of slope change in GLM fitting using model with the interaction of stimulus and PPC silencing term. **b**, Model comparison using likelihood ratio test for models with (model 2) and without (model 1) the interaction of stimulus and silencing term to account for slope change effect during PPC silence on day 1 (first introduction of testing trials). Data from muscimol and saline injection groups ($n = 27$, mean –logLR = -8.92 ± 0.21, $P < 10^{-129}$). **c**, As in **a** for photoinhibition experiments. **d**, As in **b** for photoinhibition and control groups ($n = 13$, mean –logLR = -7.61 ± 0.17, $P < 10^{-133}$). **e**, Comparison of model 1 and model 2 to account for slope changes for PPC silencing by muscimol on day 3 (testing trials were introduced on day 1). Data from the same animal group on day 1 (control) and day 3 (muscimol injection). $n = 7$, mean –logLR = 0.75 ± 0.2, $P < 0.001$. **f**, As in **e** for photoinhibition on day 3. $n = 5$, mean –logLR = 0.17 ± 0.26, $P > 0.05$. Significance test, paired t-test.

model 1: no previous choice history
model 2: with previous choice history

Extended Data Figure 14 | Comparison of GLM models with or without the term of interaction between PPC silencing and previous trial bias.

a, Visualization of previous trial choice bias in GLM fitting using model 1 (without the interaction of previous trial choice and PPC silence term). Red, data from trials with previous choice being right licking. Blue, data from trials with previous choice being left licking. Left, data from control condition. Right, data from PPC silencing (including both muscimol and photoinhibition). **b**, As in **a**, using model 2 (with the interaction of previous trial choice and PPC silencing term). Note a clear shifting in the data points during PPC silencing that is better captured by model 2. **c**, Model comparison using likelihood ratio test for models 1 and 2 in accounting for previous trial choice bias effect when PPC was silenced on day 1 introducing testing trials. Data from all groups ($n = 40$, mean $-\log LR = -3.71 \pm 0.15$, $P < 10^{-71}$, paired t-test). **d** and **e**, Model comparison using likelihood ratio test for models 1 and 2 fitted to the data of PPC silencing on day 3, using muscimol **d**, mean $-\log LR = -0.3 \pm 0.2$, $P = 0.12$, $n = 6$, and photoinhibition **e**, mean $-\log LR = 0.35 \pm 0.03$, $P < 10^{-32}$, $n = 5$. Paired t-test.

Extended Data Figure 15 | Effect of PPC silencing on reaction time and licking actions.
a, Comparison of reaction time (first lick after stimulus onset) between muscimol injection (day 1) and saline injection (day 2) conditions of the same animals. Data were segregated based on trial types. $P > 0.05$ for all comparisons, $n = 13$. **b**, Similar in **a** for data in photoinhibition of PPC. $P > 0.05$, $n = 7$. **c**, Similar plot as in **a** for mean lick number per trial in 1.5 s window after stimulus onset, compared between muscimol silencing and control conditions. $P > 0.05$ for all comparisons, $n = 13$. **d**, Similar in **c** compared between photoinhibition and control conditions. $P > 0.05$ for all comparisons, $n = 7$. Paired t-test. **e**, Raster plot of lick times in an example session with PPC silenced.

Table S1. Overlap of category selectivity

	n / Total	Percentage
No selectivity	29/758	4%
Sensory	21/758	3%
Choice	174/758	23%
Late response	161/758	21%
Sensory + choice	156/758	21%
Sensory + late response	93/758	12%
Choice + late response	268/758	35%
Sensory + choice+ late response	72/758	9%

Extended Data Table 1

Summary of fraction of neurons showing response selectivity to different behavioral variables.

Neurons are classified based on alignment of response time regarding to stimulus onset time, answer lick time, occurrence in correct trials and error trials respectively, or occurrence in subset of trials with certain tone frequencies. Neurons with multiple response peaks are classified as mixed selectivity, e.g., "Sensory + choice". Some neurons show large responses late in trials but still correlate with stimulus or choice are classified as "Sensory + late response" or "Choice + late response".

Movie S1.

Example of behavioral performance from a session with optogenetic silencing of PPC.

The mouse was trained to report categorical decisions on low frequency tones or high frequency tones by licking the left or right lickport. During a session of photoinhibition (optogenetic silencing of PPC), photoinhibition trials (blue light stimulation of bilateral PPC plus LED mask light, "optostim + LED", trial 3 and 5 in the movie) were randomly interleaved with control trials ("LED only", trial 1, 2, and 4 in the movie).