1

Human cytomegalovirus tropism modulator UL148 interacts with SEL1L, a cellular factor

that governs ER-associated degradation of the viral envelope glycoprotein, gO.

Christopher C. Nguyen,[a] Mohammed N. Siddiquey,[a] Hongbo Zhang,[a] and Jeremy P.

Kamil[a, #]

Department of Microbiology and Immunology, LSU Health Sciences Center, Shreveport,

Louisiana, USA[a]

Running Head: ER-associated degradation of HCMV glycoprotein O

#Address correspondence to Jeremy P. Kamil, jkamil@lsuhsc.edu

1501 Kings Highway, Shreveport, Louisiana 71103

ABSTRACT.

UL148 is a viral endoplasmic reticulum (ER)-resident glycoprotein that contributes to

human cytomegalovirus (HCMV) cell tropism. The influence of UL148 on tropism

correlates with its potential to promote the expression of glycoprotein O (gO), a viral

envelope glycoprotein that participates in a heterotrimeric complex with glycoproteins H

and L that is required for infectivity. In an effort to gain insight into mechanism, we used

mass spectrometry to identify proteins that co-immunoprecipitate from infected cells

with UL148. This approach led us to identify an interaction between UL148 and SEL1L,

a factor that plays key roles in ER-associated degradation (ERAD). In pulse-chase

experiments, gO was less stable in cells infected with a *UL148*-null mutant HCMV than

during wild-type infection, suggesting a potential functional relevance for the interaction

with SEL1L. To investigate whether UL148 regulates gO abundance by influencing

ERAD, siRNA silencing of either SEL1L or its partner, Hrd1, was carried out in the

context of infection. Knockdown of these ERAD factors strongly enhanced levels of gO,

but not other viral glycoproteins, and the effect was amplified in the presence of UL148.

Furthermore, pharmacological inhibition of ERAD showed similar results. Silencing of

SEL1L during infection also stabilized an interaction of gO with the ER lectin OS-9,

which likewise suggests that gO is an ERAD substrate. Taken together, our results

identify an intriguing interaction of UL148 with the ERAD machinery, and demonstrate

that gO behaves as a constitutive ERAD substrate during infection. These findings have

implications for understanding the regulation of HCMV cell tropism.

IMPORTANCE.

Viral glycoproteins in large part determine the cell types that an enveloped virus can

infect, and hence play crucial roles in transmission and pathogenesis. The glycoprotein

H/L heterodimer (gHgL) is part of the conserved membrane fusion machinery that all

herpesviruses use to enter cells. In human cytomegalovirus, gHgL participates in

alternative complexes in virions, one of which is a trimer of gHgL with glycoprotein O

(gO). Here, we show that gO is constitutively degraded during infection by the

endoplasmic reticulum-associated degradation (ERAD) pathway, and that UL148, a viral

factor that regulates HCMV cell tropism, interacts with the ERAD machinery and slows

gO decay. Since gO is required for cell-free virus to enter new host cells, but

dispensable for cell-associated spread that resists antibody neutralization, our findings

imply that the post-translational instability of a viral glycoprotein provides a basis for

viral mechanisms to modulate tropism and spread.

INTRODUCTION.

Human cytomegalovirus (HCMV) is a β-herpesvirus that establishes life-long infection in the human host and causes significant morbidity and mortality in immunocompromised patients. The virus exhibits a remarkably broad cell tropism, infecting a wide array of cell types that include epithelial cells, endothelial cells, fibroblasts, smooth muscle cells, hepatocytes, leukocytes, and hematopoietic cells [reviewed in (1, 2)]. Primary infection is thought to commence with viral replication in mucosal epithelia. The virus then transfers to circulating leukocytes to establish latent infection in hematopoietic progenitor cells (3-5). For horizontal transmission, HCMV productively infects the epithelium of several secretory tissues, allowing for the release of virus into bodily fluids including saliva, breastmilk, and urine (2, 6).

HCMV expresses two alternative envelope glycoprotein H / glycoprotein L (gHgL) complexes that impact its cell tropism: gHgLgO ("trimer") and gHgL/UL128-UL130-UL131 ("pentamer") [reviewed in (7)]. Although the pentamer is dispensable for infection of fibroblasts, it is required to infect epithelial cells, endothelial cells, and leukocytes (8-10). Recently, Murrell *et al.* showed that the pentamer promotes cell-to-cell spread that is resistant to antibody neutralization (11). On the other hand, the trimer appears to be essential for the infectivity of cell-free HCMV virions, since *gO*-null mutant viruses are defective for entry into all cell types (12, 13). While the platelet-derived growth factor receptor alpha has been identified as a cellular receptor for the trimer that is required for infection of fibroblasts (14-16), pentamer receptor(s) remain to be demonstrated. Evidence from the murine cytomegalovirus (MCMV) *in vivo* infection model suggests that the MCMV ortholog of the HCMV trimer (gHgL with m74) promotes initial infection

of tissues via a cell-free route but is dispensable for subsequent intratissue spread (17).

While these and other studies have significantly advanced the understanding of HCMV

envelope glycoprotein complexes involved in cell-type specific entry, the mechanisms

driving production of virions with distinct glycoprotein profiles remain unclear, despite

previous reports that imply that such mechanisms exist (18, 19).

In 2015, we reported the identification of a viral regulator of HCMV envelope

gH/gL complexes encoded by the *UL148* gene (20). Disruption of *UL148* in strain

TB40/E led to markedly enhanced tropism for epithelial cells accompanied by reduced

levels of trimer on virions, and an overall reduction in virion levels of gHgL. These

findings, together with results from two studies by Zhou et al. (21, 22), imply that HCMV

cell tropism might be influenced by the ratio of pentamer-to-trimer in the viral envelope,

and/or by the overall amount of gH complexes in virions. Although our previous work

suggested that UL148 increases the amount of trimer that matures beyond the

endoplasmic reticulum (ER) to the post-Golgi compartments where virions acquire their

infectious envelopes, the molecular details have remained unknown. UL148 localizes

exclusively to the ER, and *UL74* (gO) transcript levels were unaffected by disruption of

UL148 (20). Hence, it seems likely that UL148 acts within the ER to either stabilize gO

or promote assembly of the trimer. To gain more detailed information concerning

potential mechanisms, we embarked to identify proteins that interact with UL148 during

infection. Here, we show that UL148 interacts with SEL1L, a core element of the

cellular ER-associated degradation (ERAD) pathway. Furthermore, results of

experiments carried out to address the functional relevance of the interaction reveal that

gO behaves as a constitutive ERAD substrate during HCMV infection.

RESULTS

HCMV tropism factor UL148 physically associates with the ERAD adapter SEL1L

during infection. To investigate the mechanism by which UL148 promotes gO

expression, we sought to identify its interaction partners. To this end, we infected

primary human foreskin fibroblasts (HFF) with an recombinant HCMV strain TB40/E that

expresses an HA-epitope tagged UL148 (TB_148^{HA}). A parallel infection was

conducted using a control virus, TB_16^{HA}, which expresses HA-tagged UL16. Like

UL148, UL16 is an ER resident glycoprotein with type I transmembrane topology (36).

Infected cells were harvested for anti-HA immunoprecipitation (IP) at 72 h post infection

(hpi). After resolving eluates by SDS-PAGE, we excised silver-stained bands for protein

identification by mass spectrometry (MS) analysis (Fig. 1A). The full list of cellular

proteins identified from IPs of TB_148^{HA} infections in two independent experiments, but

not from IPs of TB_16^{HA}, is shown in Table S1. We were intrigued to observe peptides

that mapped to factors in the ERAD pathway, including SEL1L, XTP3-B, and OS-9.

Although co-IP experiments failed to validate several of the putative hits (not

shown), we found evidence that suggested a physical association between UL148 and

SEL1L (Fig. 1). Results of co-IP experiments from HCMV infected fibroblasts (Fig. 1B),

and from plasmid transfected 293T cells (Fig. 1C) supported the possibility that UL148

associates with endogenous SEL1L. HCMV US11, a viral immune-evasin that was

included as a positive control known to interact with SEL1L (37), was readily detected in

anti-SEL1L IPs, as was UL148. However, detection of SEL1L was less pronounced

following IP of HA-tagged US11 than following IP of HA-tagged UL148, which may

indicate a more robust interaction with UL148. UL16 and SEL1L were found to co-IP

each other poorly or undetectably, consistent with the negative result from infected cells

(Fig. 1B-1C). From these results, we concluded that UL148 may physically associate

with SEL1L, or with a protein complex that contains it.

Pulse-chase analysis of gO during WT- and *UL148*-null HCMV infection.

SEL1L plays key roles in the degradation of misfolded ER proteins [reviewed in

(38, 39)]. In the canonical ERAD pathway, SEL1L acts as an adapter between the ER

lectins OS-9 and XTP3-B and the Hrd1 complex (40), of which SEL1L is a stable

component (41, 42). The ER lectins are posited to deliver terminally misfolded

glycoproteins to the SEL1L-Hrd1 complex, which mediates ubiquitination and

translocation of ERAD substrates to the cytosol for proteasomal degradation (40). Given

that UL148 is required for high level expression of gO during infection (20) and appears

to physically associate with SEL1L (Fig. 1), we sought to determine (i) whether the

ERAD pathway targets gO for degradation and (ii) whether UL148 limits degradation of

gO.

We therefore carried out pulse-chase experiments to compare gO stability

between WT and *UL148*-null HCMV infected cells, using recombinant viruses that

express an S-peptide tag at the C-terminus of gO in a manner that avoids disruption of

the overlapping *UL73* gene encoding gN (Fig. S1A). Importantly, the *UL148*-null mutant

virus used herein, which carries nonsense codons in *UL148*, phenocopied the

enhanced epithelial cell tropism of the previously characterized *UL148* deletion mutant

(20) (Fig. S2). Further, incorporation of the S-tag, which allows for S-affinity purification

(S-AP) of gO during pulse-chase experiments, had no effect on viral replication kinetics

even during low MOI infection (Fig. S1B). To visualize the global stability of gO, and to

discriminate between immature and mature gO glycoforms, we subjected S-AP eluates

to PNGase F and endoglycosidase H (endoH) digestion, respectively. After resolving S-

AP eluates by SDS-PAGE following pulse-chase and subjecting dried gels to

autoradiography, we quantified the ~55 kD bands that corresponded to

endoglycosidase-digested gO (Fig. 2).

Notably, we found that gO decayed approximately 2-fold more rapidly during

infection with the *UL148*-null virus, TB_148STOP_gO-S (Fig. 2, "norm" signal in both

endoH and PNGase F treatments). We also observed that infections with the wild-type

(WT) comparator virus, TB_gO-S, exhibited higher absolute levels of labeled gO than

TB_148STOP_gO-S (Fig. 2, "abs" signal), even at the earliest chase time point (0 h).

Weaker baseline signals for ERAD substrates have also been observed by others

during conditions of more rapid decay, e.g. (43). We thus interpreted these results to

indicate that gO undergoes accelerated ERAD in the absence of UL148.

Knockdown of SEL1L leads to an increase in steady-state gO levels.

We next sought to investigate whether gO is subjected to SEL1L-dependent

ERAD during HCMV infection. To deplete cells of SEL1L, we transfected cells with

siRNA pools targeting *SEL1L,* and then infected with either wild-type (TB_WT) or

UL148-null mutant (TB_148STOP) HCMV strain TB40/E viruses. Depletion of SEL1L led

to a robust increase in steady-state gO levels, while levels of other viral proteins such

as pp150 and UL44 were found to be decreased or unaffected (Fig. 3). SEL1L

knockdown likewise failed to stabilize any of the other three viral glycoproteins whose

expression we monitored, gB, gH, and gL, and instead appeared to decrease their

expression (Fig. 3). The observed decrease in levels of other viral gene products may

due to the unfolded protein response (UPR), which can be activated during SEL1L

depletion (44). Intriguingly, stabilization of gO in response to SEL1L siRNA treatment

was more pronounced during WT virus infection than in the *UL148*-null setting.

Endoglycosidase H (EndoH) treatment, which removes N-linked glycans to cause

ER glycoforms of gO to migrate as single species of M_r ~50 kD, confirmed that the

major immunoreactive bands we detected during SEL1L depletion were indeed gO, and

also indicated that SEL1L knockdown stabilized endoH-sensitive ER forms of gO, as

would be consistent with the interpretation that gO is an ERAD substrate (Fig. S3A).

Similar results were also seen when S-tag antibodies were used to detect SEL1L

knockdown stabilization of gO in TB_gO-S infected cells (Fig. S3B). Arguing against

the possibility that the effects of SEL1L depletion on gO levels might be a strain-specific

phenomena, we found similar results for the HCMV laboratory strain AD169 [which lost

UL148 and several other viral genes during tissue culture adaptation (45)], and as was

the case in WT vs. *UL148*-null TB40/E, we likewise observed amplified gO stabilization

for an AD169 derivative restored for *UL148* (Fig. S4). Finally, siRNA knockdown of the

SEL1L partner protein, the E3 ubiquitin ligase Hrd1, led to increased gO levels, as

would be expected based on the effects of SEL1L knockdown (Fig. S5). We interpreted

these results to strongly suggest that gO is a constitutive ERAD substrate during HCMV

infection.

Kifunensine treatment increases steady-state gO levels. We next tested whether

pharmacological inhibition of ERAD would similarly stabilize gO. Kifunensine (KIF) is a

potent inhibitor of ER mannosidase I (46), which demannosylates the N-linked glycans

on misfolded glycoproteins to target them for recognition by ER lectins such as OS-9

and XTPB-3, which in turn deliver the misfolded glycoproteins to the SEL1L-Hrd1

complex for ERAD [(40), reviewed in (39)]. After treating HCMV-infected fibroblasts

with KIF at 72 hpi and harvesting lysates at 96 hpi, we observed a robust, dose-

dependent increase in the abundance of an intermediate-size gO glycoform that

migrated at ~90 kD SDS-PAGE (Fig. 4, Fig. S6). Furthermore, as in the SEL1L and

Hrd1 knockdown experiments, the presence of UL148 appeared to further enhance gO

levels in both control and KIF-treated cells. In contrast to SEL1L depletion, however, we

observed no significant decrease in levels of gH or gB after KIF treatment, suggesting

that KIF may be less toxic to infected cells than siRNA depletion of SEL1L or Hrd1.

These results further suggested to us that gO is a constitutive ERAD substrate during

infection.

Pulse-chase analysis of gO during SEL1L and Hrd1 depletion.

To evaluate how blockade of ERAD would affect gO decay in the presence versus the

absence of UL148, we carried out pulse-chase experiments in fibroblasts that were

siRNA depleted for SEL1L or Hrd1 and subsequently infected with WT or *UL148*-null

viruses expressing S-tagged gO. To fully resolve gO from gH, we subjected S-AP

eluates to PNGase F digestion. As expected, gO decayed more slowly during

knockdown of either SEL1L or Hrd1 compared to non-targeting control (NTC)

conditions, regardless of whether UL148 was present (Fig. 5A-B). Absolute levels of gO,

however, were most markedly amplified—by up to 7-fold at the 4 h chase time point,

when knockdown of either Hrd1, or to a lesser extent, SEL1L, occurred in presence of

UL148 (Fig. 5B). The remarkable degree of gO stabilization during Hrd1 siRNA

treatment made the roughly 2- to 3-fold effect of UL148 on gO stability, as seen in NTC

treatments at 4 h post chase, appear modest by comparison (Fig. 5A-B). The

observation that UL148 appeared to amplify stabilization of gO during ERAD blockade

may suggest that UL148 acts upstream of Hrd1/SEL1L. Alternatively, this effect may

simply reflect that substantial amounts of gO are still getting degraded by residual levels

of SEL1L/Hrd1, despite efficient siRNA knockdown (Figs. 3, S4, S5), and that these

residual levels of ERAD are readily squelched by UL148, which would be present at

higher ratio to Hrd1 during the siRNA treatment. The latter possibility would be

consistent with a functional relevance for the SEL1L interaction. Overall, these data

corroborate the hypothesis that gO is targeted by ERAD during infection, and depending

on the interpretation, might be taken to suggest that UL148 functions to inhibit ERAD of

gO.

Knockdown of SEL1L stabilizes an association between gO and the ER lectin OS-

9.

As a final complementary approach to verify whether gO behaves as an ERAD

substrate during infection, we asked whether gO associates with an ER lectin involved

in upstream events during ERAD. Depletion of SEL1L is known to stabilize what is

otherwise a transient association of the ER lectin OS-9 with ERAD substrates (40). We

thus tested whether gO associates with OS-9 during knockdown of SEL1L in the context

of the HCMV infected cell. We failed to observe co-purification of OS-9 with S-tagged

gO from infected cells that were treated with non-targeting control siRNA (NTC) (Fig. 5).

In contrast, we repeatedly detected co-purification of OS-9 with gO-S from infected cells

that were knocked down for SEL1L (e.g., Fig. 5). Furthermore, gB failed to co-IP OS-9

during either NTC or SEL1L knockdown, further arguing that the association that we

detected between OS-9 and gO is specific. That we were able to detect co-purification

of endogenous OS-9 with a putative ERAD substrate is notable. To our knowledge,

recovery of detectable levels of endogenous OS-9 in this type of experiment has been

observed only in one other study (47); most studies using this approach have relied

upon OS-9 over-expression to enhance the sensitivity of the assay (40, 44, 48, 49).

Overall, we interpreted this result to suggest that a physical association occurs between

gO and OS-9 during HCMV infection, which further argues that gO is an ERAD

substrate.

DISCUSSION

Despite that this study was initiated to address the mechanisms underlying the

influence of UL148 on trimer (gHgLgO) expression, one of the most striking findings is

the unexpectedly large role for ERAD in regulating gO expression. Newly synthesized

gO appears to be very unstable within the ER, much more so than other viral

glycoproteins, as we did not find that pharmacologic or siRNA mediated inhibition of

ERAD to stabilize any other viral glycoproteins. In fact, siRNA knockdown of either

SEL1L or Hrd1 each led to decreased expression of gH, gL, and gB (Fig. 3) but

amplified gO levels by up to 7-fold (Fig. 5B). Although the literature also suggests the

existence of post-ER processes that regulate virion incorporation of gHgL complexes

(50), because gO participates in alternative gHgL complexes that contribute to HCMV

cell tropism, our findings underscore the large potential for ERAD to sculpt the

glycoprotein composition of the virion envelope, and hence, to influence the infectivity of

HCMV for different cell types.

Why would gO be an ERAD substrate? gO is a heavily glycosylated protein, with at

least half of its ~100 kD molecular mass derived from N-linked glycans. Although the

exact number varies between HCMV strains, its primary amino acid sequence contains

approximately 20 N-x-(S/T) sequons that might serve as sites for N-linked glycosylation.

Such a large degree of glycosylation may serve a number of purposes. For instance,

gO also been reported to attenuate antibody-mediated neutralization of gH and gB (51).

Hence, analogous to the "glycan shield" provided by gp120 in HIV (52), extensive

glycosylation may contribute to the capacity of gO to protect the viral fusion machinery

from neutralizing antibodies. However, given the roles that N-linked glycans play in

glycoprotein quality control (39), the heavy degree of glycosylation may also predispose

gO to be especially susceptible to ERAD targeting by ER mannosidases.

Additionally, inefficient folding may be an inherent property of the gO polypeptide.

Previous pulse-chase studies have shown that gH forms a complex with gL within 30

min of its de novo synthesis, but that assembly of gHgLgO does not occur until

approximately 2 h (53). This observation may imply that folding of gO occurs slowly and

may be rate limiting for trimer assembly. Moreover, because gO must assemble into

trimer in order to traffic further through the secretory pathway, the relatively slow

kinetics of trimer assembly might compel a certain degree of constitutive degradation.

Indeed, orphan subunits of multisubunit complexes, such as the T-cell receptor, are

retained in the ER and destroyed via ERAD (54, 55). Competition with the pentamer-

specific subunits UL128, UL130 and UL131 for assembly onto gHgL (9, 56-58) may

further increase levels of unassembled gO that would targeted for disposal. Accordingly,

upon repair of pentamer expression to HCMV strain AD169, Wang and Shenk showed

decreased amounts of gO co-immunoprecipitated with gH [(9), Fig. 1C], and Zhou *et al.*

found transcriptional suppression of the *UL128* locus to increase levels of gHgLgO in

virions (21). It thus seems likely that both of these mechanisms, intrinsic instability and

inefficient trimer assembly, may contribute to limiting trimer expression during HCMV

infection.

Notably, we find that the homologs of gO from murine cytomegalovirus (MCMV)

and rhesus cytomegalovirus also appear to be stabilized in infected cells during

treatment with ERAD inhibitors or SEL1L knockdown (Zhang, H and Kamil, JP;

unpublished results). Therefore, constitutive ERAD of gO homologs may prove to be a

conserved feature among β-herpesviruses. Since the expression of alternative gHgL

complexes is also shared across the β-herpesvirus subfamily, the intrinsic instability of

gO and its homologs could provide a basis for mechanisms to modulate the composition

of alternative gHgL complexes in a cell-type specific manner.

How does UL148 influence ERAD of gHgLgO? Our results suggest that gO is

targeted by ERAD during infection, and that UL148 limits its degradation, with

implications for HCMV cell tropism that are summarized in our model (Fig. 7). UL148

could accomplish its effects on gO by at least two distinct but related mechanisms: (i) by

promoting the folding and/or assembly of gO into mature trimer complexes, thereby

preventing gO from being shunted to ERAD, or (ii) by dampening the ERAD pathway

generally, perhaps through an interaction with SEL1L. The first mechanism would be

consistent with our previous data suggesting that UL148 interacts with immature gHgL

complexes during their transit through the ER (20). Although we cannot yet exclude the

second possibility, determining whether UL148 stabilizes classical ERAD substrates is a

foremost experimental priority. Moreover, the two mechanisms are not mutually

exclusive. For instance, UL148 may bind to the gHgL dimer and inhibit the ERAD

machinery to allow time for gO to assemble onto the complex. Interestingly, we

consistently observed a UL148-dependent increase in the absolute levels of gO,

whether or not ERAD was disrupted, which might be taken to favor the possibility that

UL148 enhances gO levels via mechanisms upstream of ERAD. On the other hand, the

observation of enhanced gO stabilization when UL148 is present during ERAD blockade

may simply reflect an influence of UL148 against the residual levels of ERAD that

remain after siRNA- or pharmacologic disruption.

Implications of gO instability and trimer regulation by UL148. Tissue culture models

have consistently demonstrated the importance of gHgLgO for infection of several cell

types (12, 13, 22). However, *in vivo* studies using MCMV suggest that while the

analogous complex of m74 with gHgL is required for initial infection of tissues by cell-

free virions, it is dispensable for subsequent intra-tissue spread (17). By analogy, it

seems plausible that *in vivo* most HCMV-infected cells constitutively degrade gO to

promote "immunologically covert" cell-associated modes of virus spread that are driven

by the pentameric gHgL complex (11), thereby mimicking the *UL148*-null phenotype

shown in our model (Fig. 7). In cell types relevant for horizontal spread, however,

UL148 expression or activity may be enhanced to promote higher levels of gO, as would

be consistent with previous reports suggesting that the type of cell producing virions

influences the processing of virion components and virion assembly to yield distinct

virus populations (18, 19). In support of this notion, the repaired HCMV strain Merlin

(59)—which is thought to recapitulate features of HCMV that are rapidly lost during

adaptation to growth in cultured cells, and which expresses trimer very poorly, is found

to express UL148 at much lower levels than a HCMV clinical strain known to express

high levels of trimer (60). Given that HCMV is highly cell-associated *in vivo* (2), it is

intriguing to consider that UL148 abundance might be regulated so that trimer-rich

virions are produced only in settings such as glandular or ductal epithelial cells where

production of cell-free virions would be necessary for horizontal spread.

MATERIALS AND METHODS.

Cells. Human foreskin fibroblasts (HFF) were immortalized by transducing primary HFFs (ATCC #SCRC-1041) with lentivirus encoding human telomerase (hTERT) to yield HFFT cells (see below). 293T cells were purchased from Genhunter Corp. (Nashville, TN). ARPE-19 retinal pigment epithelial cells were purchased from ATCC (CRL-2302). All cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM, Corning) supplemented with 25 µg/mL gentamicin (Invitrogen), 10 µg/mL ciprofloxacin (Genhunter), and either 5% fetal bovine serum (FBS, Sigma #F2442) or 5% newborn calf serum (NCS, Sigma #N4637).

Viruses. Virus was reconstituted by electroporation of HCMV bacterial artificial chromosomes (BACs) into HFFTs, as described previously (20, 23). HCMV strain AD169rv repaired for *UL131* (ADr131) (20) and recombinants derived from it were amplified at low MOI on ARPE-19 cells until 100% cytopathic effect (CPE) was observed. Virus-containing culture supernatants were then subjected to centrifugation (1000g) for 10 minutes to pellet any cellular debris. The supernatant was then ultracentrifuged (85,000g, 1 h, 4˚C) through a 20% sorbitol cushion, and the resulting virus pellet was resuspended in DMEM containing 20% NCS. For TB40/E and related recombinants, virus was grown on HFFTs until 100% CPE was observed. Cell-associated virus was then released by Dounce-homogenization of pelleted infected cells, clarified of cell debris by centrifugation (1000g, 10 min), and combined with the cell-free medium before ultracentrifugation and resuspension as above.

Virus titration. Infectivity of virus stocks and samples were determined by the tissue

culture infectious dose 50% ($TCID_{50}$) assay. Briefly, serial dilutions of virus were used to

infect multiple wells of a 96-well plate. After 9 days, wells were scored as positive or

negative for CPE, and $TCID_{50}$ values were calculated according to the Spearman-

Kärber method (24, 25).

Virus growth kinetics. For viral yield on fibroblasts (HFFT) vs ARPE-19 epithelial cells

(Fig. S1), cells were seeded in a 24-well plate at 6.5×10^4 cells per well. For multi-cycle

growth kinetics of TB_gO-S (Fig. S3), HFFT were seeded at 1.5×10^5 cells per well.

Wells were then infected in duplicate at the indicated MOIs in in 0.3 mL per well, and

back-titration of the inocula was initiated in parallel. Inocula were removed after 16 hr,

and the cells were washed three times with 1 mL of Dulbecco's phosphate-buffered

saline (DPBS, Lonza Biowhittaker, Inc.). Cell-free supernatants were collected at

indicated times post-infection and stored at -80°C until analysis. Infectivity titers

($TCID_{50}$) were determined in parallel.

Antibodies. Mouse monoclonal antibodies (mAbs) specific for gH (AP86) (26), gB (27-

156) (27), and pp150 (28) were generous gifts from William J. Britt (University of

Alabama, Birmingham). Additional gB and UL44 mouse mAbs were purchased from

Virusys (#P1201 and #P1202-2). Rabbit antibodies were used to detect the following

proteins: UL148 (20), SEL1L (Sigma #S3699), OS-9 (Cell Signaling Tech #12497), HA

epitope tag: YPYDVPDYA (Bethyl #A190-108A), S-peptide: KETAAAKFERQHMDS

(Bethyl #A190-135A). Additional rabbit sera immunoreactive to gL, and to gO variants

from HCMV strains TB40/E and AD169 were generous gifts from Brent J. Ryckman

(University of Montana, Missoula, MT) (21).

Construction of recombinant viruses. Synthetic dsDNAs (gBlocks) and

oligonucleotides were purchased from Integrated DNA Technologies (Coralville, IA); full

sequence details are provided in Table S2. New recombinant HCMVs were constructed

in the context of the infectious bacterial artificial chromosome (BAC) clones of HCMV

strains TB40/E, TB40-BAC4 (29), and AD169, AD169rv (30), as previously described

(20, 23, 31), with full details provided in Supplemental Text S1. Briefly, two-step Red

"*en passant*" recombination (32, 33) was used in conjunction with GS1783 *E. coli* (a gift

of Greg Smith, Northwestern University) harboring BAC-cloned HCMV genomes.

Recombinant BACs were confirmed by Sanger-sequencing (Genewiz, Inc.) of modified

loci (not shown), and by BamHI and EcoRI restriction analysis (not shown) to exclude

spurious rearrangements.

TB40/E expressing UL148^{HA} or UL16^{HA}. TB_148^{HA} was described previously (20). Text

S1 describes construction of recombinant TB_UL16^{HA}.

UL148-null recombinant TB_148$_{STOP}$. TB_148$_{STOP}$ was constructed by replacement of

the UL148 coding sequence (CDS) with a synthetic version in which all in-frame

methionine codons were replaced with nonsense codons.

TB40/E and TB_148$_{STOP}$ expressing S-tagged gO. To add a bovine pancreatic

ribonuclease A S-peptide tag (S-tag) (34) (KETAAAKFERQHMDS) to the *UL74* (gO)

ORF without disrupting the overlapping gene, *UL73 (gN)*, a codon-optimized sequence

encoding the following elements was inserted between nucleotides 139966 and 139967

(numbering per Genbank Accession #EF2999921.1): (i) the 30 C-terminal amino acids

of gO, (ii) triple (Gly-Gly-Gly-Ser) linker, and (iii) the S-tag, followed by a tandem pair of

stop codons.

ADr131 expressing UL148[HA]. Repair of *UL131* in strain AD169rv (ADr131) was

described previously (20). ADr131_148 was constructed by replacing the vestigial

UL148 remnant with full-length *UL148* from HCMV strain TB40/E*.*

Plasmids.

The construction of new plasmids for this study is described in Text S1.

hTERT lentivirus production and transduction. hTERT lentivirus was isolated and

used to transduce HFF according to a modified version of the Addgene pLKO.1

protocol, as described previously (23). See Text S1 for details.

siRNA treatments.   siGenome SMARTpool siRNAs targeting SEL1L, Hrd1, and non-

targeting control (NTC) siRNA were purchased from Dharmacon. Details are provided in

Table S3. siRNAs were reverse-transfected into cells using Lipofectamine RNAiMAX

reagent (Thermo Fisher) as per the manufacturer's instructions. Briefly, two mixes were

prepared separately: Mix #1 was prepared by adding 30 pmol of siRNA to 150 µL non-

supplemented OptiMEM medium (Thermo Fisher) and gently mixing. Mix #2 was

prepared by adding 9 µL of RNAiMAX reagent to 150 µL non-supplemented OptiMEM.

Mix #1 and #2 were then combined and immediately transferred to an empty well of a 6-

well plate and incubated at room temperature for 5 min. Approximately 1.0 million cells

were then added to the well in 2.2 mL of DMEM containing 5% NCS (10 nM siRNA

final).

Endoglycosidase H analysis. EndoH digestion of 20 µg total protein was performed

according to the manufacturer's instructions (New England Biolabs).

Pulse-chase. 4×10^6 HFFTs were infected at an MOI of 1 $TCID_{50}$ per cell with TB_gO-

S and TB_148$_{STOP}$_gO-S. In experiments involving knockdown of SEL1L or Hrd1, the

cells were first reverse-transfected with siRNAs targeting SEL1L, Hrd1, or NTC, as

described above; at 8 hours post-transfection (hpt), the cells were washed with DPBS

and infected as above. At 96 hpi, the cells were washed twice with 10 mL DPBS and

incubated for 30 mins in 5 mL of Met/Cys starving medium [DMEM lacking Met, Cys,

and Glu (Gibco #21013024) plus 2 mM glutamate and 5% dialyzed FBS (Sigma

#F0392)]. Cells were then pulsed for 20 mins with 2 mL of starving medium plus 200

μCi/mL ^{35}S-Met/Cys (PerkinElmer #NEG772). To quench radiolabeling, 6 mL of chase

medium [DMEM (Corning #10013CV) plus 2 mM Met, 2 mM Cys, and 5% NCS] was

added directly to each dish. At the indicated time-points, the chase medium was

replaced with ice-cold PBS. Cells were scraped, pelleted (4000g for 1 min), and lysed in

RIPA buffer containing 25 mM HEPES (pH 7.5), 400 mM NaCl, 0.1% SDS, 0.5%

sodium deoxycholate, 1% NP-40, 1% bovine serum albumin (BSA), and 1X protease

inhibitor cocktail (PIC, CST #5871). Lysis continued for 1 h (4°C with rotation), and

lysates were clarified of insoluble material by centrifugation (20,000g, 30 mins) and

discarding of the pellet.

To quantify radiolabel incorporation into protein, 10 μL of lysate was added to 1 mL of

10% trichloroacetic acid (TCA) in triplicate, vortexed, and incubated on ice for 30 mins.

Lysate/TCA solution was pipetted onto a glass disk (Whatman #1821025), and the disk

was washed with 10 mL ice-cold 10% TCA and 10 mL ice-cold 100% ethanol. Disks

were air-dried for 30 mins and shaken overnight in 5 mL scintillation fluid (Fisher

#SX12-4). Counts per minute (CPMs) were measured on a scintillation counter

(Beckman LS 6500) and averaged across triplicates.

S-protein affinity purification of radiolabeled gO. Equal CPMs of lysate were combined

with 30 µL of S-protein agarose bead slurry (EMD Millipore) and RIPA buffer to a final

volume of 400 µL. S-AP mixes were rotated overnight at 4˚C. Beads were pelleted

(1500g, 5 mins) and washed twice with 400 µL RIPA buffer containing 1% BSA, then

washed twice with 400 µL RIPA buffer without BSA.

Endoglycosidase-digestion of S-AP eluates and autoradiography. To more effectively

resolve co-purified gO and gH during SDS-PAGE, S-AP eluates were first digested with

either peptide:N-glycosidase F (PNGase F, NEB) or endoglycosidase H (Endo H, NEB)

as follows. After the final wash of S-AP above, the beads were boiled (85°C, 5 min) in

1X glycoprotein denaturing buffer, chilled on ice, then incubated with PNGase F or Endo

H digestion mix (final volume 21 µL, 37˚C, 1 h) according to the manufacturer's

instructions. 10 µL of 4X Laemmli buffer containing 20% β-mercaptoethanol (BME) was

added to each sample, followed by boiling (85˚C, 5 mins). Samples were resolved by

sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), as described

previously (23). Gels were dried using a Bio-Rad Model 583 Gel Dryer, and radioactive

signal was captured on a storage phosphor screen for a minimum of 72 h prior to data

acquisition on Bio-Rad Molecular Imager FX. Band intensities were quantified using

Quantity One 1-D Analysis Software (Bio-Rad). In brief, volume rectangles were drawn

closely around the 55 kD gO band to generate a report of adjusted volume (CNT*mm^2)

values after local background subtraction. These values were normalized and reported

as "gO signal".

Mass spectrometry. 2×10^7 fibroblasts (HFF) were infected at an MOI of 1 TCID$_{50}$ per

cell with TB_148^{HA}. Cells were lysed at 120 hpi in modified RIPA lysis buffer [50 mM

HEPES (pH 7.5), 1% TritonX-100, 400 mM NaCl, 0.5% sodium deoxycholate, 10%

glycerol, and 1X PIC]. Anti-HA magnetic beads (Pierce #8837) were incubated (rotation

overnight, 4°C) with lysates plus 100 µg/mL BSA, washed three times with lysis buffer,

and eluted by boiling in Laemmli buffer (65°C, 5 min). Eluate was resolved by SDS-

PAGE, and the gel was silver-stained according to the manufacturer's protocol

(ThermoFisher #24600). Silver-stained bands were excised from SDS-PAGE gels and

sent to the Taplin Biological Mass Spectrometry Facility at Harvard Medical School

(Boston, MA). Processing of gel slices for trypsin digestion, and resolution of peptides

by nano-scale capillary reverse-phase HPLC are described in SI Methods. Eluted

peptides were subjected to electrospray ionization and entered an LTQ Orbitrap Velos

Pro ion-trap mass spectrometer (Thermo Fisher). Peptides were fragmented, and

specific fragment ions were detected to generate a tandem pass spectrum for each

peptide. Each spectrum was matched to a fragmentation pattern database by the

program Sequest (Thermo Fisher) to determine the sequence of each peptide and

hence protein identity (35). All databases included forward and reverse versions of

peptide sequences, and data was filtered on a 1-2% peptide false discovery rate.

Immunoprecipitation from transfected cells.

5×10^5 293T cells were transfected with 2.5 µg of pEF1α V5 C vector (Invitrogen)

encoding codon-optimized *US11^{HA}*, *UL16^{HA}*, or *UL148^{HA}* using Mirus TransIT-2020

(#MIR 5404) according to the manufacturer's instructions. At 48 hpt, cells were lysed in

300 µL of lysis buffer [50 mM HEPES (pH 7.5), 400 mM NaCl, 0.5% sodium

deoxycholate, and 1X PIC]. 200 µL of lysate was rotated overnight (4°C) with 25 µL of

anti-HA magnetic bead slurry (Pierce #8837). Beads were washed three times with lysis

buffer and eluted by heating (50˚C, 10 min) in 2X Laemmli buffer containing 5% (v/v)

BME. Eluates and whole cell lysates were analyzed by western blot.

OS-9 co-purification assay. 1×10^6 HFFT were reverse-transfected with siRNAs

targeting SEL1L or NTC, as above. At 48 hpt, cells were infected with TB_gO-S virus at

an MOI of 1 $TCID_{50}$ per cell. At 96 hpi, cells were lysed in 50 mM HEPES (pH 7.5), 1%

TritonX-100, 400 mM NaCl. Equal µg of protein (determined by Pierce BCA assay) were

subjected to S-AP [30 µL of S-protein agarose slurry] or glycoprotein B (gB)-IP [30 µL

Protein G magnetic bead slurry (EMD Millipore) plus anti-gB mAb 27-156 (27)]. After

overnight rotation (4˚C), beads were washed three times with lysis buffer and eluted in

1X Laemmli buffer plus 5% BME. Eluates and whole cell lysates were analyzed by

western blot.

ACKNOWLEDGMENTS

This project was supported by NIH Grants R01-AI116851 and P30GM110703. Its

contents are solely the responsibility of the authors and do not necessarily represent the

official views of the funding agencies. C.C.N. was supported by a fellowship from the

Center of Cardiovascular Diseases and Sciences at LSU Health Sciences Center –

Shreveport. We thank William Britt, Brent Ryckman, Greg Smith, Christian Sinzger, and

Ulrich Koszinowski for generously sharing reagents.

Author contributions are as follows: C.C.N. performed the majority of experiments.

C.C.N. and J.P.K. designed experiments, interpreted results, and wrote the manuscript.

J.P.K. designed HCMV recombinants. M.N.S. constructed recombinant TB_148$_{STOP}$,

performed UL148/SEL1L and gO/OS-9 co-purification studies, and prepared samples

for MS analysis. H.Z. constructed recombinant viruses TB_gO-S and TB_148$_{STOP}$_gO-S

and assisted with replication of virus yield experiments.

FIG. 1. Viral tropism factor UL148 associates with ERAD adapter SEL1L during infection. (A) Silver stain. Fibroblasts (HFF) infected with TB_148HA were lysed at 72 hpi and subjected to HA-IP. Visible silver-stained bands were excised and analyzed by mass spectrometry. The putative association between UL148 and SEL1L was confirmed by HA-IP and western blot of UL148HA complexes from (B) infected cells or (C) 293T cells transfected with plasmids expressing the indicated HCMV ER-resident proteins.

FIG. 2. Pulse-chase analysis of gO during *UL148*-null HCMV infection. Fibroblasts (HFFTs) were infected at an MOI of 1 $TCID_{50}$ per cell with TB40/E-derived recombinant TB_gO-S expressing S-tagged gO (WT) or its *UL148*-null derivative, TB_148$_{STOP}$_gO-S (148$_{STOP}$). At 96 hpi, the cells were pulsed for 20 mins with 200 µCi/mL ^{35}S-Met/Cys and chased for the indicated times before lysis. Equal TCA-precipitable CPMs of each sample were subjected to S-AP, followed by endoglycosidase digestion and SDS-PAGE. The dried gel was exposed to a phosphor screen to produce an autoradiograph. The densities of the indicated gO bands (*) were calculated and expressed either as absolute signal ("abs") in relation to the WT/0 h band or as signal normalized ("norm") across WT or *UL148*-null conditions relative to the respective 0 h band.

FIG. 3. Depletion of SEL1L in HCMV-infected cells increases steady-state gO levels. (A) Fibroblasts (HFFTs) were reverse-transfected with siRNA targeting SEL1L (SEL) or non-targeting control (NTC). At 48 h post transfection, the cells were infected

with HCMV strain TB40/E (WT) or its *UL148*-null derivative, TB_148$_{STOP}$ (148$_{STOP}$) at an

MOI of 1 TCID$_{50}$ per cell. HCMV glycoprotein levels at 96 hpi were analyzed by western

blot.

FIG. 4. Treatment of HCMV-infected cells with an ERAD inhibitor increases

steady-state gO levels. Fibroblasts (HFFTs) were infected at an MOI of 1 TCID$_{50}$ per

cell with TB_gO-S (WT), TB_148$_{STOP}$_gO-S (148$_{STOP}$), or mock infection (M). Cells

were treated with kifunensine (KIF) at 2.5 µM or carrier-alone (water) at 72 hpi. At 96

hpi, HCMV glycoprotein levels were analyzed by western blot.

FIG. 5. Depletion of SEL1L or Hrd1 in HCMV-infected cells stabilizes gO during

WT and *UL148*-null infection. Fibroblasts (HFFTs) were reverse-transfected with

siRNAs targeting (A) SEL1L or (B) Hrd1. At 6 hpt, the cells were infected at an MOI of 1

TCID$_{50}$ per cell with TB_gO-S (WT) or TB_148$_{STOP}$_gO-S (148$_{STOP}$). At 96 hpi, the cells

were pulsed for 20 mins with 200 µCi/mL ^{35}S-Met/Cys and chased for the indicated

times before lysis. Equal TCA-precipitable CPMs of each sample were subjected to S-

AP, followed by PNGase F digestion to allow better resolution of gH and gO by SDS-

PAGE. The dried gel was exposed to a phosphor screen to produce an autoradiograph.

The densities of gO bands were calculated and expressed either as absolute signal

("abs") in relation to the WT/NTC/0 h band or as normalized signal ("norm") for each

virus condition relative to the respective NTC/0 h band.

FIG. 6. Depletion of SEL1L in HCMV-infected cells stabilizes a physical

association between gO and ER lectin OS-9. Fibroblasts (HFFTs) were reverse-

transfected with siRNAs targeting SEL1L (SEL) or NTC. 48 h later, the cells were

infected with TB_gO-S at an MOI of 1 $TCID_{50}$ per cell. At 96 hpi, cell lysates were

subjected to S-AP or gB-IP and analyzed by western blot.

FIG. 7. Model. HCMV gO, either through unfavorable folding or inefficient assembly

onto gH/gL, is targeted for destruction by ERAD, which is mediated by SEL1L/Hrd1.

UL148, perhaps through a functional interaction with SEL1L, slows the degradation of

gO.

SUPPLEMENTAL MATERIAL

TEXT S1: Supplemental Methods and Supplementary Figure Legends.

FIG. S1: Construction and characterization of TB_gO-S.

FIG. S2: Growth kinetics of TB_148^{HA} and TB_148$_{STOP}$ on HFFT and ARPE-19 cells.

FIG. S3: EndoH analysis and detection of gO levels during SEL1L knockdown.

FIG. S4: Depletion of SEL1L in HCMV strain AD169-infected cells recapitulates effects
on HCMV glycoproteins.

FIG. S5: Hrd1 depletion in HCMV-infected cells increases gO levels.

FIG. S6: Dose-responsive effect of kifunensine on HCMV gO levels.

TABLE S1: Mapping of peptide hits from UL148HA-IP eluates to human gene products.

TABLE S2: Synthetic DNAs used in the construction and sequence-confirmation of BAC
recombinants and plasmids

TABLE S3: Dharmacon SMARTpool siRNA sequences used in this study.

REFERENCES

1.	Sinzger C, Digel M, Jahn G. 2008. Cytomegalovirus cell tropism. Curr Top Microbiol Immunol 325:63-83.
2.	Mocarski ES, Shenk T, Griffiths PD, Pass RF. 2013. Cytomegaloviruses, p 1960-2014. *In* Knipe DM, Howley PM (ed), Fields' Virology, 6th ed, vol 2. Wolters Kluwer Health/Lippincott Williams & Wilkins, Philadelphia.
3.	Goodrum FD, Jordan CT, High K, Shenk T. 2002. Human cytomegalovirus gene expression during infection of primary hematopoietic progenitor cells: a model for latency. Proc Natl Acad Sci U S A 99:16255-60.
4.	Maciejewski JP, Bruening EE, Donahue RE, Mocarski ES, Young NS, St Jeor SC. 1992. Infection of hematopoietic progenitor cells by human cytomegalovirus. Blood 80:170-8.
5.	Smith MS, Bentz GL, Alexander JS, Yurochko AD. 2004. Human cytomegalovirus induces monocyte differentiation and migration as a strategy for dissemination and persistence. J Virol 78:4444-53.
6.	Cannon MJ, Hyde TB, Schmid DS. 2011. Review of cytomegalovirus shedding in bodily fluids and relevance to congenital cytomegalovirus infection. Rev Med Virol 21:240-55.
7.	Li G, Kamil JP. 2015. Viral Regulation of Cell Tropism in Human Cytomegalovirus. J Virol 90:626-9.
8.	Wang D, Shenk T. 2005. Human cytomegalovirus UL131 open reading frame is required for epithelial cell tropism. J Virol 79:10330-8.
9.	Wang D, Shenk T. 2005. Human cytomegalovirus virion protein complex required for epithelial and endothelial cell tropism. Proc Natl Acad Sci U S A 102:18153-8.
10.	Hahn G, Revello MG, Patrone M, Percivalle E, Campanini G, Sarasini A, Wagner M, Gallina A, Milanesi G, Koszinowski U, Baldanti F, Gerna G. 2004. Human cytomegalovirus UL131-128 genes are indispensable for virus growth in endothelial cells and virus transfer to leukocytes. J Virol 78:10023-33.
11.	Murrell I, Bedford C, Ladell K, Miners KL, Price DA, Tomasec P, Wilkinson GWG, Stanton RJ. 2017. The pentameric complex drives immunologically covert cell-cell transmission of wild-type human cytomegalovirus. Proc Natl Acad Sci U S A 114:6104-6109.
12.	Jiang XJ, Adler B, Sampaio KL, Digel M, Jahn G, Ettischer N, Stierhof YD, Scrivano L, Koszinowski U, Mach M, Sinzger C. 2008. UL74 of human cytomegalovirus contributes to virus release by promoting secondary envelopment of virions. J Virol 82:2802-12.
13.	Wille PT, Knoche AJ, Nelson JA, Jarvis MA, Johnson DC. 2010. A human cytomegalovirus gO-null mutant fails to incorporate gH/gL into the virion envelope and is unable to enter fibroblasts and epithelial and endothelial cells. J Virol 84:2585-96.
14.	Kabanova A, Marcandalli J, Zhou T, Bianchi S, Baxa U, Tsybovsky Y, Lilleri D, Silacci-Fregni C, Foglierini M, Fernandez-Rodriguez BM, Druz A, Zhang B, Geiger R, Pagani M, Sallusto F, Kwong PD, Corti D, Lanzavecchia A, Perez L. 2016. Platelet-derived

growth factor-alpha receptor is the cellular receptor for human cytomegalovirus
gHgLgO trimer. Nat Microbiol 1:16082.
15.　　Wu Y, Prager A, Boos S, Resch M, Brizic I, Mach M, Wildner S, Scrivano L, Adler B.
2017. Human cytomegalovirus glycoprotein complex gH/gL/gO uses PDGFR-alpha
as a key for entry. PLoS Pathog 13:e1006281.
16.　　Stegmann C, Hochdorfer D, Lieber D, Subramanian N, Stohr D, Laib Sampaio K,
Sinzger C. 2017. A derivative of platelet-derived growth factor receptor alpha binds
to the trimer of human cytomegalovirus and inhibits entry into fibroblasts and
endothelial cells. PLoS Pathog 13:e1006273.
17.　　Lemmermann NA, Krmpotic A, Podlech J, Brizic I, Prager A, Adler H, Karbach A, Wu
Y, Jonjic S, Reddehase MJ, Adler B. 2015. Non-redundant and redundant roles of
cytomegalovirus gH/gL complexes in host organ entry and intra-tissue spread. PLoS
Pathog 11:e1004640.
18.　　Patrone M, Secchi M, Fiorina L, Ierardi M, Milanesi G, Gallina A. 2005. Human
cytomegalovirus UL130 protein promotes endothelial cell infection through a
producer cell modification of the virion. J Virol 79:8361-73.
19.　　Scrivano L, Sinzger C, Nitschko H, Koszinowski UH, Adler B. 2011. HCMV spread and
cell tropism are determined by distinct virus populations. PLoS Pathog 7:e1001256.
20.　　Li G, Nguyen CC, Ryckman BJ, Britt WJ, Kamil JP. 2015. A viral regulator of
glycoprotein complexes contributes to human cytomegalovirus cell tropism. Proc
Natl Acad Sci U S A 112:4471-6.
21.　　Zhou M, Yu Q, Wechsler A, Ryckman BJ. 2013. Comparative analysis of gO isoforms
reveals that strains of human cytomegalovirus differ in the ratio of gH/gL/gO and
gH/gL/UL128-131 in the virion envelope. J Virol 87:9680-90.
22.　　Zhou M, Lanchy JM, Ryckman BJ. 2015. Human Cytomegalovirus gH/gL/gO
Promotes the Fusion Step of Entry into All Cell Types, whereas gH/gL/UL128-131
Broadens Virus Tropism through a Distinct Mechanism. J Virol 89:8999-9009.
23.　　Wang D, Li G, Schauflinger M, Nguyen CC, Hall ED, Yurochko AD, von Einem J, Kamil
JP. 2013. The ULb' region of the human cytomegalovirus genome confers an
increased requirement for the viral protein kinase UL97. J Virol 87:6359-76.
24.　　Kärber G. 1931. Beitrag zur kollektiven Behandlung pharmakologischer
Reihenversuche. Archiv f experiment Pathol u Pharmakol 162: 480-483.
25.　　Spearman C. 1908. The Method of "Right and Wrong Cases" (Constant
Stimuli) without Gauss's Formula. . Br J Psychol 2:227-242
26.　　Urban M, Britt W, Mach M. 1992. The dominant linear neutralizing antibody-binding
site of glycoprotein gp86 of human cytomegalovirus is strain specific. J Virol
66:1303-11.
27.　　Spaete RR, Saxena A, Scott PI, Song GJ, Probert WS, Britt WJ, Gibson W, Rasmussen L,
Pachl C. 1990. Sequence requirements for proteolytic processing of glycoprotein B
of human cytomegalovirus strain Towne. J Virol 64:2922-31.
28.　　Sanchez V, Greis KD, Sztul E, Britt WJ. 2000. Accumulation of virion tegument and
envelope proteins in a stable cytoplasmic compartment during human
cytomegalovirus replication: characterization of a potential site of virus assembly. J
Virol 74:975-86.
29.　　Sinzger C, Hahn G, Digel M, Katona R, Sampaio KL, Messerle M, Hengel H,
Koszinowski U, Brune W, Adler B. 2008. Cloning and sequencing of a highly

90   productive, endotheliotropic virus strain derived from human cytomegalovirus
91   TB40/E. J Gen Virol 89:359-68.

92   30.   Hobom U, Brune W, Messerle M, Hahn G, Koszinowski UH. 2000. Fast screening
93   procedures for random transposon libraries of cloned herpesvirus genomes:
94   mutational analysis of human cytomegalovirus envelope glycoprotein genes. J Virol
95   74:7720-9.

96   31.   Li G, Rak M, Nguyen CC, Umashankar M, Goodrum FD, Kamil JP. 2014. An epistatic
97   relationship between the viral protein kinase UL97 and the UL133-UL138 latency
98   locus during the human cytomegalovirus lytic cycle. J Virol 88:6047-60.

32.   Tischer BK, Smith GA, Osterrieder N. 2010. En passant mutagenesis: a two step
markerless red recombination system. Methods Mol Biol 634:421-30.

33.   Tischer BK, von Einem J, Kaufer B, Osterrieder N. 2006. Two-step red-mediated
recombination for versatile high-efficiency markerless DNA manipulation in
Escherichia coli. Biotechniques 40:191-7.

34.   Richards FM, Vithayathil PJ. 1959. The preparation of subtilisn-modified
ribonuclease and the separation of the peptide and protein components. J Biol Chem
234:1459-65.

35.   Eng JK, McCormack AL, Yates JR. 1994. An approach to correlate tandem mass
spectral data of peptides with amino acid sequences in a protein database. J Am Soc
Mass Spectrom 5:976-89.

36.   Dunn C, Chalupny NJ, Sutherland CL, Dosch S, Sivakumar PV, Johnson DC, Cosman D.
2003. Human cytomegalovirus glycoprotein UL16 causes intracellular sequestration
of NKG2D ligands, protecting against natural killer cell cytotoxicity. J Exp Med
197:1427-39.

37.   van de Weijer ML, Bassik MC, Luteijn RD, Voorburg CM, Lohuis MA, Kremmer E,
Hoeben RC, LeProust EM, Chen S, Hoelen H, Ressing ME, Patena W, Weissman JS,
McManus MT, Wiertz EJ, Lebbink RJ. 2014. A high-coverage shRNA screen identifies
TMEM129 as an E3 ligase involved in ER-associated protein degradation. Nat
Commun 5:3832.

38.   Brodsky JL. 2012. Cleaning up: ER-associated degradation to the rescue. Cell
151:1163-7.

39.   Olzmann JA, Kopito RR, Christianson JC. 2013. The mammalian endoplasmic
reticulum-associated degradation system. Cold Spring Harb Perspect Biol 5.

40.   Christianson JC, Shaler TA, Tyler RE, Kopito RR. 2008. OS-9 and GRP94 deliver
mutant alpha1-antitrypsin to the Hrd1-SEL1L ubiquitin ligase complex for ERAD.
Nat Cell Biol 10:272-82.

41.   Lilley BN, Ploegh HL. 2005. Multiprotein complexes that link dislocation,
ubiquitination, and extraction of misfolded proteins from the endoplasmic reticulum
membrane. Proc Natl Acad Sci U S A 102:14296-301.

42.   Mueller B, Lilley BN, Ploegh HL. 2006. SEL1L, the homologue of yeast Hrd3p, is
involved in protein dislocation from the mammalian ER. J Cell Biol 175:261-70.

43.   Lilley BN, Ploegh HL. 2004. A membrane protein required for dislocation of
misfolded proteins from the ER. Nature 429:834-40.

44.   Alcock F, Swanton E. 2009. Mammalian OS-9 is upregulated in response to
endoplasmic reticulum stress and facilitates ubiquitination of misfolded
glycoproteins. J Mol Biol 385:1032-42.

45.   Cha TA, Tom E, Kemble GW, Duke GM, Mocarski ES, Spaete RR. 1996. Human
137   cytomegalovirus clinical isolates carry at least 19 genes not found in laboratory
138   strains. J Virol 70:78-83.
139   46.   Elbein AD, Tropea JE, Mitchell M, Kaushal GP. 1990. Kifunensine, a potent inhibitor
140   of the glycoprotein processing mannosidase I. J Biol Chem 265:15599-605.
141   47.   Bernasconi R, Galli C, Calanca V, Nakajima T, Molinari M. 2010. Stringent
142   requirement for HRD1, SEL1L, and OS-9/XTP3-B for disposal of ERAD-LS substrates.
143   J Cell Biol 188:223-35.
144   48.   Hosokawa N, Kamiya Y, Kamiya D, Kato K, Nagata K. 2009. Human OS-9, a lectin
145   required for glycoprotein endoplasmic reticulum-associated degradation,
146   recognizes mannose-trimmed N-glycans. J Biol Chem 284:17061-8.
147   49.   Ron E, Shenkman M, Groisman B, Izenshtein Y, Leitman J, Lederkremer GZ. 2011.
148   Bypass of glycan-dependent glycoprotein delivery to ERAD by up-regulated EDEM1.
149   Mol Biol Cell 22:3945-54.
150   50.   Luganini A, Cavaletto N, Raimondo S, Geuna S, Gribaudo G. 2017. Loss of the Human
151   Cytomegalovirus US16 Protein Abrogates Virus Entry into Endothelial and Epithelial
152   Cells by Reducing the Virion Content of the Pentamer. J Virol 91.
153   51.   Jiang XJ, Sampaio KL, Ettischer N, Stierhof YD, Jahn G, Kropff B, Mach M, Sinzger C.
154   2011. UL74 of human cytomegalovirus reduces the inhibitory effect of gH-specific
155   and gB-specific antibodies. Arch Virol 156:2145-55.
156   52.   Wei X, Decker JM, Wang S, Hui H, Kappes JC, Wu X, Salazar-Gonzalez JF, Salazar MG,
157   Kilby JM, Saag MS, Komarova NL, Nowak MA, Hahn BH, Kwong PD, Shaw GM. 2003.
158   Antibody neutralization and escape by HIV-1. Nature 422:307-12.
159   53.   Huber MT, Compton T. 1999. Intracellular formation and processing of the
160   heterotrimeric gH-gL-gO (gCIII) glycoprotein envelope complex of human
161   cytomegalovirus. J Virol 73:3886-92.
162   54.   Yang M, Omura S, Bonifacino JS, Weissman AM. 1998. Novel aspects of degradation
163   of T cell receptor subunits from the endoplasmic reticulum (ER) in T cells:
164   importance of oligosaccharide processing, ubiquitination, and proteasome-
165   dependent removal from ER membranes. J Exp Med 187:835-46.
166   55.   Bonifacino JS, Suzuki CK, Lippincott-Schwartz J, Weissman AM, Klausner RD. 1989.
167   Pre-Golgi degradation of newly synthesized T-cell antigen receptor chains: intrinsic
168   sensitivity and the role of subunit assembly. J Cell Biol 109:73-83.
169   56.   Chandramouli S, Malito E, Nguyen T, Luisi K, Donnarumma D, Xing Y, Norais N, Yu D,
170   Carfi A. 2017. Structural basis for potent antibody-mediated neutralization of
171   human cytomegalovirus. Sci Immunol 2.
172   57.   Ciferri C, Chandramouli S, Donnarumma D, Nikitin PA, Cianfrocco MA, Gerrein R,
173   Feire AL, Barnett SW, Lilja AE, Rappuoli R, Norais N, Settembre EC, Carfi A. 2015.
174   Structural and biochemical studies of HCMV gH/gL/gO and Pentamer reveal
175   mutually exclusive cell entry complexes. Proc Natl Acad Sci U S A 112:1767-72.
176   58.   Ryckman BJ, Rainish BL, Chase MC, Borton JA, Nelson JA, Jarvis MA, Johnson DC.
177   2008. Characterization of the human cytomegalovirus gH/gL/UL128-131 complex
178   that mediates entry into epithelial and endothelial cells. Journal of Virology 82:60-
179   70.
180   59.   Stanton RJ, Baluchova K, Dargan DJ, Cunningham C, Sheehy O, Seirafian S, McSharry
181   BP, Neale ML, Davies JA, Tomasec P, Davison AJ, Wilkinson GW. 2010.

182   Reconstruction of the complete human cytomegalovirus genome in a BAC reveals
183   RL13 to be a potent inhibitor of replication. J Clin Invest 120:3191-208.
184 60.   Zhang L, Zhou M, Stanton R, Kamil JP, Ryckman BJ. 2018. Expression levels of
185   glycoprotein O (gO) vary between strains of human cytomegalovirus, influencing the
186   assembly of gH/gL complexes and virion infectivity. bioRxiv doi:10.1101/299222.
187

wild-type

UL148-null

ER lumen

gO

SEL1L UL148

Hrd1

gH/gL

SEL1L

Hrd1

ERAD

trimer/pentamer ratio ↑

epithelial-tropism ↓

cell-free spread

trimer/pentamer ratio ↓

epithelial-tropism ↑

intratissue spread