In silico analysis of piRNAs in retina reveals potential targets in intracellular transport and retinal degeneration

Suganya Sivagurunathan[a, b†], Nagesh Srikakulam[c†], Jayamuruga Pandian Arunachalam[d], Gopal Pandi[c], Subbulakshmi Chidambaram[a, e*]

[a]RS Mehta Jain Department of Biochemistry and Cell Biology, Vision Research Foundation, Chennai, India.

[b]School of Chemical and Biotechnology, SASTRA University, Thanjavur, India.

[c]Department of Plant Biotechnology, School of Biotechnology, Madurai Kamaraj University, Madurai 625021, Tamil Nadu, India.

[d]Central Inter-Disciplinary Research Facility (CIDRF), Sri Balaji Vidyapeeth University, Mahatma Gandhi Medical College and Research Institute Campus, Pondicherry, India.

[e]Department of Biochemistry and Molecular Biology, Pondicherry University, Puducherry, India.

[†]Both authors contributed equally to this work

*Email ID – csubbulakshmi@gmail.com

Abstract

Long considered to be active only in germline, PIWI/piRNA pathway is now known to play significant regulatory function in somatic cells, especially neurons. Nonetheless, so far there is no evidence for the presence of piRNAs in the neurosensory retina. In this study, we have uncovered 102 piRNAs in human retina and retinal pigment epithelium (RPE) by analysing RNA-seq data. The identified piRNAs were enriched with three motifs predicted to be involved in rRNA processing and sensory perception. Computational analysis revealed complete and partial seed sequence similarity between piR-62011 and sensory organ specific miR-183/96/182 cluster. Further, expression of piRNAs in donor eyes were assessed by qRT-PCR. Intriguingly, loss of piRNAs in HIWI2 knockdown ARPE19 cells downregulated specific targets implicated in intracellular transport (SNAREs and *Rabs*), circadian clock (*TIMELESS*) and retinal degeneration (*LRPAP1* and *RPGRIP1*). Moreover, under oxidative stress ARPE19 cells expressed elevated levels of piRNAs indicating their potential role in retinal pathology. Thus, our data provides a clear evidence for the expression of piRNAs in human retina and RPE. Collectively, our work demonstrates the pleiotropic effects of piRNAs illustrating their dynamic regulation of distinct molecular events in the maintenance of retinal homeostasis.

Introduction

Retina, a neural layer of the eye, is developed from the inner layer of the optic cup regulated by the eye field specific transcription factors (EFTF) whereas the Retinal Pigment Epithelium (RPE) develops from outer layer of the optic cup. Photoreceptors of the retina converts the photons of light into electrical impulses where the crosstalk with RPE is necessary. Recent studies have indicated the involvement of non-coding RNAs in eye development plausibly by targeting the EFTF[1]. Besides their role in development, the importance of non-coding RNAs in visual function was confirmed when the depletion of miRNAs resulted in loss of outer segments of cone photoreceptors causing reduced light response[2]. Further, non-coding RNAs are also linked with retinal pathologies and are suggested to be novel targets for therapy[3].

piRNAs are a group of small noncoding RNAs which bind with PIWI subclade of Argonaute family proteins[4–6]. They are 24 – 32 nucleotides (nt) long, methylated at their 3' ends[7,8] and are produced by a pathway independent of Dicer, making them distinct from other small noncoding RNAs. These group of RNAs are essential for maintaining the genomic integrity in germline cells by repressing the transposable elements[9]. Initially, it was thought to be restricted to germ cells, however, recent reports indicate their presence in various somatic tissues[10,11] and cancer cells[12–14]. In addition, altered expression of piRNAs is observed during various pathological conditions like cardiac hypertrophy[15] and liver regeneration[10] in rats. Thus, growing evidence on the role of piRNAs in various non-germline tissues underscores the need to explore their importance in specific somatic cells as well.

Recent years has seen a surge of research findings on the existence of piRNAs in neurons. Initially, Lee *et al* has suggested roles for piRNAs in modulating the dendritic spine development in mouse hippocampal neurons[16]. In addition, altered expression of piRNAs were reported in rat brain after transient focal Ischemia[17]. Interestingly, a mechanistic role for

piRNAs in neurons has been demonstrated by Rajasethupathy *et al* where PIWI/piRNA complexes in *Aplysia* enhanced memory-related synaptic plasticity through epigenetic regulation of CREB2[18]. Recently, piRNAs have shown to attenuate the axonal regeneration of adult sensory neurons in rats[19]. Moreover, piRNAs are deregulated in Alzheimer's disease and are known to affect the disease associated pathways[20]. Nevertheless, there is no study on the presence and role of PIWI-interacting RNAs (piRNAs) in retina.

In our recent reports, we have shown that PIWI-like proteins are expressed in human retina and HIWI2 (PIWIL4) regulates tight junctions in maintaining the integrity of RPE[21]. HIWI2 also alters the expression of OTX2, an eye field specification marker in human retinoblastoma cells[22]. The presence of PIWI-like proteins in retina strongly implies the existence of piRNAs. Therefore, we investigated the expression and role of piRNAs in retina and RPE. The present study identified 102 piRNAs by computational analysis of RNA-seq data obtained from human retina and RPE tissues. Additionally, novel targets of piRNAs in multiple molecular processes have been validated. Our data presents new insights into the functions of piRNAs in retina and RPE.

Results

Expression of piRNAs in human retina and RPE

We examined whether piRNAs are expressed in retina and RPE by analysing the small RNA sequencing data retrieved from ArrayExpress (Accession ID: E-MTAB-3792)[23]. It consisted of 16 samples of human retina and 2 samples of RPE. The small RNA sequences were checked for their quality and the reads were mapped with the annotated small RNAs (tRNA, snRNA, snoRNA, miRNA, rRNA, mtRNA). The mapped reads were discarded to filter out the known small RNAs and the unmapped reads were aligned to the human genome. The resulting reads were considered as potential piRNA candidates and were mapped with the sequences of

piRNAs downloaded from piRNAQuest[24] (Fig.1). Interestingly, we observed 102 piRNAs with transcript per million (TPM) values > 1. piRNAs that were expressed in retina and RPE are listed in Table S1. A heatmap on the expression of the identified piRNAs indicated that piR-31068, piR-35982, piR-35284 were in abundance in both the tissues analysed. Interestingly, piR-62011, piR-36743 and piR- 36742 were highly expressed only in retina. Similarly, the expression of piR-35411, piR-35412, piR-35413, piR-31104 and piR-31103 were higher only in RPE tissue (Fig. 2a & b). Together, the results showed that piRNAs were expressed in ocular tissues and indeed differential expression of piRNAs were observed between retina and RPE.

Tissue enriched piRNAs

Having observed the piRNAs in ocular tissues, we next investigated whether the identified piRNAs are tissue specific. To this end, our data was compared with the piRNAs so far reported in human brain, cardiac progenitors and cardiac fibroblasts, ovary and testis[20,25–27]. We observed 4 piRNAs (piR-30855, piR-34535, piR-44719 and piR-61647) to be expressed only in retina and RPE. All the analysed tissues shared 5 piRNAs (piR-33382, piR-33468, piR-35284, piR-35413 and piR-36225) in common. Moreover, 3 piRNAs (piR-32376, piR-43771 and piR-62011) were found to be commonly present in retina, RPE and testis. However, piR-62011 is abundantly expressed in retina (TPM value - 112874) when RPE is considered (TPM value - 1003) (Fig 2b). Notably, around 75% of the piRNAs in the retina/RPE tissues matched with the piRNAs in brain, followed by 72% with ovary. In contrast, piRNAs of ocular tissues showed only 35% and 34% match with testis and cardiomyocytes (cardiac progenitors and cardiac fibroblasts) respectively (Fig. 2c). Thus, a significant overlap was found amongst piRNAs expressed in human brain and retina/RPE; and piR-62011 appeared to be a retina enriched piRNA.

Genomic origin of piRNAs

piRNAs are shown to originate originally from the transposon sequences. The origin of the identified piRNAs indicated that majority of them mapped to tRNA family repeats, especially to tRNAs Val and Gly followed by Ala. This is in accordance with the earlier report on somatic piRNAs derived from the tRNA fragments[28]. Moreover, protein coding genes were also known to be a source of piRNAs[29] and likewise our data specified 28 protein coding genes as origin for a set of piRNAs. Besides, 20 of the piRNAs originated from SNORD gene family which encodes small nucleolar RNAs (snoRNA). Recently, small nucleolar RNAs were reported to be precursor of piRNA[30]. Strikingly, the retina enriched piR-62011 had its origin in the 3' UTR of miR-182 (Fig. 3a & b, Table S2).

Somatic piRNAs are reported to map only lesser percentage to piRNA clusters unlike the piRNAs expressed in germline cells[31]. In order to understand the genomic localization of piRNAs in retina/RPE, we considered minimum 3 piRNAs originating within 20 kb distance to form a cluster. Our analysis indicated that chromosome (chr) 6 harboured four piRNA clusters followed by chr 1 with three piRNA clusters. Chr 2, 5 and 16 had two piRNA clusters in each while chr 3,11,14,17 and 18 expressed one piRNA cluster. It is important to note that genomic locus of piR-62011 was present in chr 7. Interestingly, 2 piRNA clusters had its origin in mitochondrial genome, which is in line with the identification of piR-36707 and piR-36741 in mitochondria of the mammalian cancer cells[32] (Table 1).

Long non-coding RNA (lncRNA) _piRNA_microRNA (miRNA) alignment

Since the crosstalk between the non-coding RNAs is evolving[33], we investigated the lncRNAs and miRNAs that might be related to the identified piRNAs. Alignment of the piRNA sequences with miRNAs showed that piR-35059, piR-36741 and piR-62011 matched with the seed sequences of miR-4420, miR-4284 and miR-182 respectively. It is known that miR-

183/96/182 forms a cluster in chromosome 7 and piR-62011 displayed seed sequence match with miR-96 and miR-183 as well but with 1 and 2 mismatches respectively. It was interesting to note that the sequence of piR-62011, an abundant piRNA in retina was exactly similar to miR-182, which was reported to be the highly expressed miRNA in human retina[23]. Additionally, some piRNAs showed an exact sequence match with 20 lncRNAs and few with either 1 or 2 mismatches. Intriguingly, many of the lncRNAs that mapped with the piRNAs were uncharacterised (Table S3). Apparently, the sequence similarities between miR/piR/lncRNA hint a potential crosstalk among the non-coding RNAs in the retina.

Enriched motifs

Though piRNAs are not conserved across species, these small RNAs have been observed to possess short motifs. We identified 3 conserved motifs in piRNAs using motif predictor MEME and the first motif (21 nt long) with an E-value of $5.4 \, e^{-127}$ was conserved in 38 piRNA sequences. Further, the function and localization of the nucleotide motif was predicted using GOMo which showed that the nucleotide sequences in the first motif was found to be involved in rRNA processing and localized to nucleolus and cytoplasm. The second motif (21 nt long) with an E-value of $1.3 \, e^{-023}$ was conserved in 14 piRNAs and was predicted to be localized in mitochondria. The third motif (28 nt long) with an E-value of $2.2 \, e^{-016}$, speculated to reside in extracellular region, was conserved in 21 piRNA sequences. This particular motif is predicted to be involved in G-protein coupled protein receptor signalling pathway and sensory perception (Fig. 4a).

Transcription factors regulating piRNAs

Given the presence of piRNAs in human retina and RPE, we further attempted to describe the possible regulatory factors of piRNA. Sequences upstream of the piRNA loci were searched for the presence of transcription factor binding sites (Fig. 4b). It is interesting to note that

transcription factor c-Rel showed higher number of binding sites in the promoters of piRNAs followed by Hepatic Leukemia factor (HLF). c-Rel is a member of NFκB transcription factor family and we also observed binding sites for NFκB in the promoters of piRNA. Though c-Rel defects are mainly associated with inflammatory disorders they are also associated with memory and synaptic plasticity[34]. Meanwhile, piRNAs were also reported to be crucial for synaptic plasticity and memory storage by regulating CREB2[18]. Our data indicated that there were binding sites for CREB in the promoter regions of the 102 piRNAs. c-Rel, HLF, Evi-1, CCAAT-displacement protein (CDP) and Nkx2-5 were the transcription factors which contained the most number of binding sites in the analysed regions.

Evaluation of the identified piRNAs in human retina and RPE samples

Out of the 102 identified piRNAs, we evaluated the expression of piR-31068 (abundant in both retina and RPE), piR-35411 (abundant in RPE), piR-36742, piR-62011(abundant in retina) and piR-60576 (moderate expression in both the tissues). qPCR was done in human retina and RPE tissues dissected from donor eyes which confirmed the expression of these piRNAs (Fig. 5a). cDNA conversion without the addition of reverse transcriptase (or) the cDNA that were converted using random primers did not result in amplification indicating the specificity of the reaction. Among the piRNAs that were evaluated, piR-31068 is abundantly expressed (with low ΔCq), which is in accordance with the TPM values obtained (Table S1).

To validate whether the amplicons are indeed piRNAs, we used Reverse Transcription at Low dNTP concentration followed by PCR (RTL-P) analysis. This method investigated 2' O - methylation at the 3` ends of piRNAs. Reverse transcriptase stalls at methylated sites at low dNTP concentration. Exploiting this feature of the enzyme, piRNAs and miR-26a (control) were amplified at both high and low concentration of dNTP. During amplification, two different primers were used – unanchored[35] (3` methylation site of the piRNA will not be

covered) and anchored (3` methylation site of the piRNA will be covered by the primer) primers. Amplification of piRNAs in the presence of unanchored primer at low dNTP concentration was low whereas it was more at high dNTP concentration. miR-26a which was not methylated at its 3` ends did not show such changes (Fig.5b). Hence, the methylation at 3` ends of the amplicons further confirmed that the obtained bands were indeed piRNAs. Altogether, our data confirmed the expression of piRNAs in human retina and RPE tissues.

Putative targets of piRNAs

We used miRanda algorithm to identify the putative targets of piRNAs, to gain insight into the possible regulatory function. We obtained a list of potential targets with a threshold score of 200 and a minimum energy value of -20 kCal/Mol (Table S4). When the targets were functionally annotated, we observed that 24% of them were uncharacterised proteins. Majority of the putative targets were channels, transporters and proteins necessary for signal transduction followed by 13% and 14% of proteins essential for metabolism and nucleotide binding respectively. Around 9% of the proteins are involved in intracellular trafficking and 7% belongs to proteins important for adhesion, cytoskeletal arrangements and cell-cell contact including tight junction proteins. The other predicted targets are known to be crucial for immune and stress response, sensory perception, neuronal activity and DNA repair (Fig. 6a).

Evaluation of the predicted targets of piRNA

Among the predicted targets of piRNAs, the possible influence of piRNAs in cell signalling and cell-cell contact proteins can be conjectured from our previous study[21]. We demonstrated the function of HIWI2 in altering the tight junction proteins through Akt/GSK3β signalling pathway[21]. Since PIWI proteins bind with piRNAs, effect of HIWI2 on tight junctions might be plausibly through piRNAs. Next, we analysed whether absence of piRNAs influenced the expression of putative targets. HIWI2, a ubiquitously present member of 4 PIWI-like proteins

in human, was efficiently silenced in ARPE19 (RPE) cell line as indicated by the Western blot (Fig. 6b). Accordingly, the piRNAs were downregulated in Si-HIWI2 cells (Fig. 6c) and the interesting novel targets from intracellular trafficking machinery were analysed. We screened for the expression of *SNARE*s and *Rab* transcripts upon HIWI2 silencing in ARPE19 by semi RT-PCR (Fig. 6d). Further qRT-PCR indicated that *STX1A, Rab5, Rab8, VAMP7* and *VAMP8* were significantly downregulated in Si-HIWI2 cells (Fig. 6e). In addition, the expression of other predicted targets such as *TIMELESS, RPGRIP1* and *LRPAP1* were also evaluated. Strikingly, *TIMELESS* transcripts were 4.34-fold downregulated in the absence of HIWI2 (Fig. 6f). Both, *RPGRIP1* and *LRPAP1* transcripts were 3.57-fold and 1.51-fold downregulated in Si-HIWI2 cells respectively (Fig. 6f). Notably, it was reported that *RPGRIP1* mutant of Zebrafish did not develop rod outer segments in the retina and were suggested to regulate ciliary protein trafficking[36]. *LRPAP1* is known to be associated with myopia[37] and the variations in LRPAP1 were observed to be associated with late onset of Alzheimer's disease[38]. Thus, loss of piRNAs downregulated the expression of *SNARE*s, *Rab*s and *RPGRIP1* which are important for ciliary trafficking. In addition, *LRPAP1* and the expression of key circadian clock protein, *TIMELESS* were also reduced in the absence of piRNAs.

Elevated levels of piRNA under oxidative stress

Oxidative stress is a major cause in the pathogenesis of ocular diseases such as keratoconus, cataractogenesis and age-related macular degeneration[39]. To gain insight into the importance of piRNAs in pathological conditions, ARPE19 cells were subjected to oxidative stress. Upon treatment of the cells with 200 μM of H_2O_2, the piRNAs were significantly upregulated. We observed 2.3-fold upregulation of piR-36742 followed by 1.9-fold increase in piR-35411, piR-31068, piR-62011 and 1.7-fold upregulation in piR-60576 (Fig. 7). Clearly, the expression of piRNAs were altered under oxidative stress.

Discussion

Recently, we reported the presence of PIWI-like proteins in ocular tissues and the importance of HIWI2 in the maintenance of epithelial integrity of RPE. Since PIWI proteins are associated with piRNAs, it was compelling to examine whether piRNAs play a role in retinal function. The present study confirmed the expression of piRNAs in retina and RPE. Moreover, loss of piRNAs altered the expression of specific transcripts involved in intracellular trafficking, sensory perception and circadian rhythm. We have also observed elevated levels of piRNAs under oxidative stress denoting their association with pathological conditions.

Our analysis showed that majority of the piRNAs in retina and RPE originate from tRNA fragments of Val and Gly, which is in line with the findings in human brain[20] and ovarian cancer tissues[26]. Although, recent studies demonstrated the existence of small nucleolar-RNA (sno-RNA) derived piRNAs and a novel pathway for gene regulation[30,40], still, far less is known about this group of piRNAs. The SNORD family derived piRNAs in retina/RPE may play important functions which warrants further investigation.

Interestingly, seed sequence similarity between miR-17-5p and piRNAs has been shown to reciprocally regulate each other during mouse embryonic development[41]. It was surprising to find that piR-62011, an abundant piRNA in retina displayed seed sequence match and aligned completely with miR-182. In addition, piR-62011 exhibited seed sequence match with miR-96 and also with miR-183 but with one and two mismatches respectively. It is plausible that PIWI is essential for the functioning of these miRNAs as MIWI is shown to be required for the expression of a subset of miRNAs in mouse[42]. Furthermore, it will be interesting to investigate the functions of two classes of small non-coding RNAs with similar sequences. According to miRbase[43], miR-183/96/182 are in the same cluster within 5 kb on chromosome 7. Strikingly, piR-62011 is also present on chromosome 7, although originates roughly 360kb away from this

miRNA cluster. Importantly, studies in mouse retina demonstrated miR-183/96/182 as 'sensory organ specific' miRNA cluster abundantly expressed in retina[44]. Moreover, this cluster is modulated by light[45]. In support to these findings, absence of miR-182 and miR-183 lead to the loss of cone outer segments and were described to be necessary for the formation of inner segments, connecting cilia and short outer segments of photoreceptor in adult mice[2]. Moreover, miR-183/96/182 has been downregulated in a mouse model for inherited blindness[46]. Further, miR-182 is associated with an ocular pathology called primary open-angle glaucoma[47]. Hence, we speculate a reciprocal regulation between piR-62011 and the sensory organ specific miR-183/96/182 through seed sequence match and it would be relevant to investigate the role of piR-62011 in glaucoma and retinal degeneration.

It is interesting to note the three conserved motifs in retina and RPE. Earlier, Grivna *et al* have observed an 8-nt motif shared by 17 sequences and a 21-nt motif shared by 5 sequences, among the 40 piRNAs they have cloned from germline of male mouse[6]. Watanabe *et al* also observed 5 motifs in the germline small RNAs (gsRNA) of mouse[5]. Our bioinformatic analysis predicted, nucleus and mitochondria as the localization signal for the first motif. Accordingly, the piRNAs are reported to be observed in mitochondria of cancer cells[32] and nucleus of *Aplysia* neurons[18]. Out of the 102 piRNAs, 7 of them mapped to the ribosomal proteins (Table S1) implicating their probable role in rRNA processing, which is in line with the functional prediction of piRNAs. The third motif identified from our study is predicted to be involved in sensory perception which could be one of the novel functions of piRNAs in retina.

As the expression of piRNAs were reported to be tissue-specific[31], our data also indicates that there are only 5 piRNAs in common among so far reported human tissues – brain[20], cardiac progenitors and cardiac fibroblasts[25], ovary[26] and testis[27]. It is interesting to note that 75% of piRNAs in retina/RPE showed similarities with the piRNAs reported in human brain (number of reported piRNAs - 668) although testis had a larger number of annotated piRNAs (11680)

suggesting their expression to be enriched in neurons. Our findings aligned with the study of Martinez *et al* which described the tissue specific expression of piRNAs with the potential of special functions in various somatic tissues[31]. Tissue enrichment analysis showed that piR-62011 is enriched in retina indicating its potential as a biomarker or a therapeutic target in retinal diseases. Although, we reported only the annotated piRNAs; identification of *de novo* piRNAs in retina and RPE should shed light on more ocular specific piRNAs. In this context, investigation of piRNAs in other non-gonadal tissues is necessary, in order to categorise the tissue specific piRNAs.

Computational annotation of piRNA target sites showed potential pleiotropic roles of piRNA in retina and RPE. It is intriguing that the majority of the identified putative targets of piRNAs are yet to be characterized for their function. The largest class of the targets is involved in signal transduction. For instance, piR-Hep1, which is conserved with piR-43771, piR-43772 and piR-60565 reduces the phosphorylation levels of Akt in hepatocellular carcinoma[48] and Rajan *et al* speculated the role of piRNAs in Akt pathway[49]. Evidently, our initial work showed that HIWI2 silencing increased the phosphorylation of Akt and GSK3β. We have screened 43 kinases in Si-HIWI2 cells, involved in 4 different pathways – JAK/STAT, MAPK, Akt and AMPK. Apart from modulation of Akt pathway, we observed significant variation in the phosphorylation levels of STAT3, p53, AMPKα and HSP90[21]. In the similar line, Wang *et al* have also shown the influence of HIWI2 in the activation of MAPK/ERK, TGFβ and FGF signalling pathways[50]. It is plausible that these changes could be mediated through piRNAs, suggesting its involvement in cell signalling and cell-cell contact.

Other key group of identified piRNA targets were related to specific functions of retina and we have validated the expression of selective novel targets in the absence of piRNAs. Downregulation of *STX1A, Rab5, Rab8, VAMP7* and *VAMP8* indicates the likely association of piRNAs in intracellular trafficking. Rab8 is suggested to participate in ciliary trafficking of

rhodopsin and in the morphogenesis of rod outer segment disk[51]. Moreover, Rab8 and Rab11 coordinates the primary ciliogenesis mainly by regulating the vesicular trafficking[52]. Primary cilia is crucial for sensory perception where defects in ciliogenesis have been observed in Bardet-Biedl syndrome (BBS) and retinal degeneration is one of the primary characteristics of BBS[53]. Rabin8 directly interacts with the BBSome[54] which is stimulated by Rab11[52] underscoring the importance of trafficking proteins in sensory perception. In line with our study, piR-34393 is known to target *Rab11A* in human brain[20]. Remarkably, mutations in *RPGRIP1*, an interacting protein of RPGR (retinitis pigmentosa GTPase regulator), is associated with diseases like leber congenital amaurosis, retinitis pigmentosa and cone-rod dystrophy. It is also known to be essential for the development of rod outer segments by regulating the ciliary protein trafficking and Rab8 was observed to be mislocalized in *rpgrip1* mutants[36]. Our data shows significant downregulation of both *Rab8* and *RPGRIP1* transcripts in the absence of piRNAs. Interestingly, expression of *RPGRIP1* was readily detected only in retina and testis when a panel of RNA from human tissues were screened[55]. piRNAs, which were thought to be specific to testis, might regulate *RPGRIP1* and thus may effect neurodegeneration which needs to be confirmed by further functional experiments.

LRPAP1 (LDL receptor related protein associated protein 1), another predicted piRNA target, is also downregulated in Si-HIWI2 cells. *LRPAP1* is known to be associated with high myopia by influencing the activity of TGFβ and thereby remodelling the sclera[56]. Alterations in *LRPAP1* is also related with late-onset of Alzheimer's disease (AD)[38]. Esposito *et al* have shown the expression of piRNAs in Alzheimer's disease[57] and a further study demonstrated dysregulation of piRNAs in AD[20]. Together with *RPGRIP1*, downregulation of *LRPAP1* indicates a possible role of piRNAs in retinal degeneration.

Our data also showed that TIMELESS, a protein needed for the regulation of circadian rhythm in mammals, is observed to be affected in the absence of piRNAs. Activation of circadian

piRNA expression is suggested to be a strategy adopted by *Drosophila melanogaster* to maintain the genomic integrity during aging[58]. Furthermore, the role of PIWIL2 in suppressing the degradation of circadian proteins CLOCK and BMAL1 through phosphorylation of Akt/GSK3β and by binding to the E-box sequences of BMAL1/CLOCK complex, confirms the importance of PIWI-like proteins in regulating circadian rhythms[59]. It is worth noting that miR-182 is known to target the 3`UTR of *CLOCK* mRNA[60,61] and is inhibited by GSK3β[62]. Similarly, HIWI2 also affects the Akt/GSK3β pathway[21] and as mentioned earlier, piR-62011 aligns exactly with the seed sequence of miR-182 indicating a potential functional interaction between piR-62011/miR-182 in influencing the activity of CLOCK-controlled genes. In addition, improper processing of pre-miR-182 is documented to be the major cause of depression in insomnia patients[61]. Hence, it would be interesting to explore the regulatory role played by piRNAs and miRNAs in the circadian rhythm related disorders.

It is intriguing that the examined targets were all downregulated at transcript level although they were selected based on complementarity using miRanda. It is known that piRNAs regulate the expression of genes by imperfect base-pairing rule[63], a mechanism similar to miRNAs. The possible mechanism behind the repression of the transcripts could be epigenetic activation[64] or silencing[18] through repressive epigenetic marks by piRNAs at the transcriptional level. Nevertheless we cannot rule out the mechanism of posttranscriptional regulation by piRNAs as shown in transposon-driven mRNAs[65].

Our data showed that expression of piRNAs are elevated under oxidative stress. Martinez *et al* recently reported 40 piRNAs that were regulated by hypoxia in tumors[66]. It is possible that these piRNAs might have been influenced by the oxidative stress/reactive oxygen species generated by the hypoxic conditions. In eye, oxidative stress plays a major role in the pathogenesis of many diseases such as keratoconus, Fuchs' endothelial corneal dystrophy, Leber's hereditary optic neuropathy and cataractogenesis[39]. ROS and oxidative stress is also

known to stimulate inflammation and pathological angiogenesis in diabetic retinopathy and it plays a pivotal role in the pathogenesis of age-related macular degeneration[39]. Thus, overall several evidences are indicating a potential role of piRNAs in retinal pathologies.

Conclusion

Our study showed the first line of evidence for the presence of piRNAs in human retina and RPE. We have further identified and validated novel target proteins of piRNAs which are essential for fundamental functions of retina and RPE. The intriguing finding of our analysis is the seed sequence similarity between piR62011 and sensory organ specific miR183/96/182 cluster. The crosstalk may modulate the unique cellular demands of retina, such as, circadian expression of phototransduction components and polarized trafficking in photoreceptors. In addition, altered expression of piRNAs under oxidative stress implies piRNAs in pathological conditions. In summary, our finding highlights the significance of PIWI/piRNAs in visual function and retinal degeneration.

Materials and methods

Data source

Small RNA sequencing datasets of human retina and retinal pigment epithelium was downloaded from (Accession ID: E-MTAB-3792)[23]. The dataset consisted of 16 human retinal and 2 retinal pigment epithelial tissue samples. The datasets ERR973601-ERR973608 and ERR973611-ERR673618 were retinal samples. ERR973609 and ERR973610 were RPE datasets.

Data processing

Adapter sequences present in the datasets were removed using Cutadapt[67]with PHRED score 30 and the sequences less than 20 nt were filtered out. The reads which mapped to other non-

coding RNAs (tRNA, snRNA, snoRNA, miRNA, rRNA, mtRNA) were discarded and the unmapped reads were taken for further analysis. miRNA sequences were downloaded from miRBase[43,68] and all other RNA sequences were retrieved from Pfam[69]. The reads were mapped using BOWTIE[70].

Identification and differential expression of known piRNAs

After eliminating the known annotated small RNAs, unique reads and read count from the data were retrieved using unique_reads_for_mapping.pl from miRGrep tool[71]. Unique reads were aligned to the known piRNA database downloaded from piRNAquest[24] using soap.short tool[71] with default parameters (allowed 2bp mismatches). Output file was used to calculate the final read count using 'awk' command and retain the piRNAs which are having >=10 read count in any one library. Details of the origin of piRNAs were retrieved from piRNAQuest[24] and the piRNAs were categorised as clusters when minimum of 3 piRNAs originated within 20 Kbp.

The quantification of each piRNA in each library was normalized to TPM (Tags per million reads) values and further converted to Log2 values. Before Log2 conversion, piRNAs which were having 0 TPM in any one of the library were removed. Log2 values were used in Morpheus to generate heatmap.

Tissue enrichment analysis

piRNAs annotated in brain[20], cardiac progenitors and fibroblasts[25], ovary[26] and testis[27] were downloaded. In case of human brain, piRNAs reported in both healthy brain and brain with Alzheimer's disease were combined and the duplicates were removed which amounted to 668 in total. For piRNAs in cardiac progenitors and cardiac fibroblasts, the piRNAs listed in cardiospheres, cardiosphere-derived cells and cardiac fibroblasts were combined and the redundant entries were eliminated resulting to 2366 piRNAs. Out of 20121 piRNAs in human

testis, same piRNAs of varying lengths were considered as one and a total of 11680 piRNAs were compared with retina/RPE. InteractiVenn[72] was used for tissue enrichment analysis.

Motif prediction and lncRNA/miRNA/piRNA alignment

Motifs in the piRNA sequences were predicted using MEME software[73] with site distribution parameter to be any number of repetitions and the putative functions of the motifs were analysed by GOMo[73] with the default factors. Soap.short tool[71] was used to compare the identified piRNA sequences with the lncRNA sequences downloaded from the database LNCipedia[74,75]. The alignment was performed with the default parameters allowing maximum number of 2 bp mismatches. Human miRNA sequences from miRbase were aligned with the identified piRNAs to analyse seed sequence similarity and the alignments with the maximum of 2 mismatches are listed in Supplementary table S3.

Identification of transcription factor binding sites

We analysed the Transcription start site (TSS), promoter and TFBS for the piRNA genes. Human promoter prediction tool, FPROM[76] was used to predict the TSS and promoter site within the 10 kb upstream of the piRNA locus, extracted from the UCSC table browser. TFBS scanned for the 1-kb upstream of the each piRNA using the P-Match 1.0[77] with the vertebrate matrix groups and cut-off has chosen to minimize the sum of both false positives and false negatives error rates.

Prediction of piRNA targets

Targets of the piRNA were predicted using miRanda v3.3a algorithm[78–80] with the following parameters: Gap Open Penalty (-9.0); Gap Extend Penalty (-4.0); Score Threshold (200.0); Energy Threshold (1.0 kcal/mol); Scaling Parameter (4.0). The hits which had score > 200 and energy value < -20 kcal/Mol were considered to be the putative targets.

Ethics statement

Donor eyes were obtained from CU Shah eye bank, Sankara Nethralaya with prior approval from the Institutional Review Board and Ethics Sub-Committee of Vision Research Foundation, Chennai, India. The experiments involving the usage of the ocular tissues were carried out according to the Tenets of the Declaration of Helsinki.

Cell culture

ARPE19 cells (ATCC-CRL-2302) (ATCC, USA) were cultured using DMEM-F12 (Sigma Aldrich, USA) with 10% (v/v) heat-inactivated Fetal Bovine Serum (ThermoFisher Scientific, USA) and was maintained at 37ºC in 5% CO_2 incubator. The cells were passaged the day before transfection and 2×10^5 cells were seeded in a 6-well plate. DsiRNA specific to 3' UTR of HIWI2 (Integrated DNA Technologies, USA) were transfected using Lipofectamine RNAiMAX (ThermoFisher Scientific, USA) according to manufacturer's instruction. Scrambled sequence was used as experimental control and RNA was isolated from Si-Control and Si-HIWI2 cells, 48 h post transfection. To induce oxidative stress, the cells were treated with 200 μM H_2O_2 for 24 h and RNA was isolated from the same to check for the expression of piRNAs.

Real time PCR

Primers for piRNA amplification were designed using miR2 primer software[81]. cDNA conversion was done with 1 μg of RNA based on the protocol by Busk and Cirera, 2014[82]. The cocktail for cDNA conversion contained Poly A polymerase (New England Biolabs, USA) for adding poly A tail; M-MuLV reverse transcriptase (New England Biolabs, USA) and a RT-primer of sequence - 5′-CAGGTCCAGTTTTTTTTTTTTTTTGC-3', to convert the poly A tailed RNA to cDNA. 25 ng of cDNA was used for amplification of piRNAs listed in Table 2.

Transcripts other than small RNAs were amplified from cDNA converted using iScript cDNA synthesis kit (Biorad, USA).

Reverse Transcription at Low dNTP concentration followed by PCR (RTL-P) analysis

RTL-P analysis was carried out as described by Dong *et al*, 2012[35]. RNA was converted to cDNA with low concentration of dNTP (0.4 µM) in the presence of either unanchored primer (UAP) or anchored primer (AP). The same was done at high concentration of dNTP (40 µM) as well. Four different cDNA ((1) Low dNTP with UAP; (2) Low dNTP with AP; (3) High dNTP with UAP; (4) High dNTP with AP) were used to amplify piRNA and miRNA and the amplicons were run in 2% agarose gel. Primers used in the study are listed in Table 2.

Data availability

The datasets analysed were downloaded from the ArrayExpress (Accession No: E-MTAB-3792) (https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-3792/).

References

1. Graw, J. Eye development. *Curr Top Dev Biol* **90,** 343–386 (2010).

2. Busskamp, V. *et al.* miRNAs 182 and 183 Are Necessary to Maintain Adult Cone Photoreceptor Outer Segments and Visual Function. *Neuron* **83,** 586–600 (2014).

3. Zelinger, L. & Swaroop, A. RNA Biology in Retinal Development and Disease. *Trends Genet* (2018). doi:10.1016/j.tig.2018.01.002

4. Aravin, A. *et al.* A novel class of small RNAs bind to MILI protein in mouse testes. *Nature* **442,** 203–7 (2006).

5. Watanabe, T. *et al.* Identification and characterization of two novel classes of small RNAs in the mouse germline: retrotransposon-derived siRNAs in oocytes and germline small RNAs in testes. *Genes Dev* **20,** 1732–1743 (2006).

6. Grivna, S. T., Beyret, E., Wang, Z. & Lin, H. A novel class of small RNAs in mouse spermatogenic cells. *Genes Dev* **20,** 1709–14 (2006).

7. Ohara, T. *et al.* The 3′ termini of mouse Piwi-interacting RNAs are 2′-O-methylated. *Nat Struct Mol Biol* **14,** 349–350 (2007).

8. Kirino, Y. & Mourelatos, Z. Mouse Piwi-interacting RNAs are 2′-O-methylated at their 3′ termini. *Nat Struct Mol Biol* **14,** 347–348 (2007).

9. Brennecke, J. *et al.* Discrete small RNA-generating loci as master regulators of

transposon activity in Drosophila. *Cell* **128,** 1089–103 (2007).

10. Rizzo, F. *et al.* Timed regulation of P-element-induced wimpy testis-interacting RNA expression during rat liver regeneration. *Hepatology* **60,** 798–806 (2014).

11. Henaoui, I. S. *et al.* PIWI-interacting RNAs as novel regulators of pancreatic beta cell function. *Diabetologia* **60,** 1977–1986 (2017).

12. Yin, J. *et al.* piR-823 contributes to colorectal tumorigenesis by enhancing the transcriptional activity of HSF1. *Cancer Sci* **108,** 1746–1756 (2017).

13. Li, D. *et al.* piR-651 promotes tumor formation in non-small cell lung carcinoma through the upregulation of cyclin D1 and CDK4. *Int J Mol Med* **38,** 927–36 (2016).

14. Peng, L. *et al.* piR-55490 inhibits the growth of lung carcinoma by suppressing mTOR signaling. *Tumour Biol* **37,** 2749–56 (2016).

15. Rajan, K. S. *et al.* Abundant and Altered Expression of PIWI-Interacting RNAs during Cardiac Hypertrophy. *Hear Lung Circ* **25,** 1013–1020 (2016).

16. Lee, E. J. *et al.* Identification of piRNAs in the central nervous system. *RNA* **17,** 1090–1099 (2011).

17. Dharap, A., Nakka, V. P. & Vemuganti, R. Altered Expression of PIWI RNA in the Rat Brain After Transient Focal Ischemia. *Stroke* **42,** 1105–1109 (2011).

18. Rajasethupathy, P. *et al.* A Role for Neuronal piRNAs in the Epigenetic Control of Memory-Related Synaptic Plasticity. *Cell* **149,** 693–707 (2012).

19. Phay, M., Kim, H. H. & Yoo, S. Analysis of piRNA-Like Small Non-coding RNAs Present in Axons of Adult Sensory Neurons. *Mol Neurobiol* **55,** 483–494 (2018).

20. Roy, J., Sarkar, A., Parida, S., Ghosh, Z. & Mallick, B. Small RNA sequencing revealed dysregulated piRNAs in Alzheimer's disease and their probable role in pathogenesis. *Mol Biosyst* **13,** 565–576 (2017).

21. Sivagurunathan, S., Palanisamy, K., Arunachalam, J. P. & Chidambaram, S. Possible role of HIWI2 in modulating tight junction proteins in retinal pigment epithelial cells through Akt signaling pathway. *Mol Cell Biochem* **427,** 145–156 (2017).

22. Sivagurunathan, S., Arunachalam, J. P. & Chidambaram, S. PIWI-like protein, HIWI2 is aberrantly expressed in retinoblastoma cells and affects cell-cycle potentially through OTX2. *Cell Mol Biol Lett* **22,** 17 (2017).

23. Karali, M. *et al.* High-resolution analysis of the human retina miRNome reveals isomiR variations and novel microRNAs. *Nucleic Acids Res* **44,** 1525–40 (2016).

24. Sarkar, A., Maji, R. K., Saha, S. & Ghosh, Z. piRNAQuest: searching the piRNAome for silencers. *BMC Genomics* **15,** 555 (2014).

25. Vella, S. *et al.* PIWI-interacting RNA (piRNA) signatures in human cardiac progenitor cells. *Int J Biochem Cell Biol* **76,** 1–11 (2016).

26. Singh, G., Roy, J., Rout, P. & Mallick, B. Genome-wide profiling of the PIWI-interacting RNA-mRNA regulatory networks in epithelial ovarian cancers. *PLoS One* **13,** e0190485 (2018).

27. Yang, Q. *et al.* MicroRNA and piRNA Profiles in Normal Human Testis Detected by

Next Generation Sequencing. *PLoS One* **8,** e66809 (2013).

28. Keam, S. P. *et al.* The human Piwi protein Hiwi2 associates with tRNA-derived piRNAs in somatic cells. *Nucleic Acids Res* **42,** 8984–95 (2014).

29. Robine, N. *et al.* A Broadly Conserved Pathway Generates 3′UTR-Directed Primary piRNAs. *Curr Biol* **19,** 2066–2076 (2009).

30. Zhong, F. *et al.* A SnoRNA-derived piRNA interacts with human interleukin-4 pre-mRNA and induces its decay in nuclear exosomes. *Nucleic Acids Res* **43,** gkv954 (2015).

31. Martinez, V. D. *et al.* Unique somatic and malignant expression patterns implicate PIWI-interacting RNAs in cancer-type specific biology. *Sci Rep* **5,** 10423 (2015).

32. Kwon, C. *et al.* Detection of PIWI and piRNAs in the mitochondria of mammalian cancer cells. *Biochem Biophys Res Commun* **446,** 218–223 (2014).

33. Yamamura, S., Imai-Sumida, M., Tanaka, Y. & Dahiya, R. Interaction and cross-talk between non-coding RNAs. *Cell Mol Life Sci* **75,** 467–484 (2018).

34. Gilmore, T. D. & Gerondakis, S. The c-Rel Transcription Factor in Development and Disease. *Genes Cancer* **2,** 695–711 (2011).

35. Dong, Z.-W. *et al.* RTL-P: a sensitive approach for detecting sites of 2'-O-methylation in RNA molecules. *Nucleic Acids Res* **40,** e157 (2012).

36. Raghupathy, R. K. *et al.* Rpgrip1 is required for rod outer segment development and ciliary protein trafficking in zebrafish. *Sci Rep* **7,** 16881 (2017).

37. Khan, A. O., Aldahmesh, M. A. & Alkuraya, F. S. Clinical Characterization of LRPAP1-Related Pediatric High Myopia. *Ophthalmology* **123,** 434–435 (2016).

38. Sánchez, L. *et al.* Variation in the LRP-associated protein gene (LRPAP1) is associated with late-onset Alzheimer disease. *Am J Med Genet* **105,** 76–8 (2001).

39. Nita, M. & Grzybowski, A. The Role of the Reactive Oxygen Species and Oxidative Stress in the Pathomechanism of the Age-Related Ocular Diseases and Other Pathologies of the Anterior and Posterior Eye Segments in Adults. *Oxid Med Cell Longev* **2016,** 3164734 (2016).

40. He, X. *et al.* An Lnc RNA (GAS5)/SnoRNA-derived piRNA induces activation of TRAIL gene by site-specifically recruiting MLL/COMPASS-like complexes. *Nucleic Acids Res* **43,** 3712–25 (2015).

41. Du, W. W. *et al.* Reciprocal regulation of miRNAs and piRNAs in embryonic development. *Cell Death Differ* **23,** 1458–1470 (2016).

42. Grivna, S. T., Pyhtila, B. & Lin, H. MIWI associates with translational machinery and PIWI-interacting RNAs (piRNAs) in regulating spermatogenesis. *Proc Natl Acad Sci* **103,** 13415–13420 (2006).

43. Griffiths-Jones, S., Grocock, R. J., van Dongen, S., Bateman, A. & Enright, A. J. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res* **34,** D140-4 (2006).

44. Xu, S., Witmer, P. D., Lumayag, S., Kovacs, B. & Valle, D. MicroRNA (miRNA)

Transcriptome of Mouse Retina and Identification of a Sensory Organ-specific miRNA Cluster. *J Biol Chem* **282,** 25053–25066 (2007).

45. Krol, J. *et al.* Characterizing Light-Regulated Retinal MicroRNAs Reveals Rapid Turnover as a Common Property of Neuronal MicroRNAs. *Cell* **141,** 618–631 (2010).

46. Palfi, A. *et al.* microRNA regulatory circuits in a mouse model of inherited retinal degeneration. *Sci Rep* **6,** 31431 (2016).

47. Liu, Y. *et al.* A Common Variant in *MIR182* Is Associated With Primary Open-Angle Glaucoma in the NEIGHBORHOOD Consortium. *Investig Opthalmology Vis Sci* **57,** 4528 (2016).

48. Law, P. T.-Y. *et al.* Deep sequencing of small RNA transcriptome reveals novel non-coding RNAs in hepatocellular carcinoma. *J Hepatol* **58,** 1165–73 (2013).

49. Rajan, K. S., Velmurugan, G., Pandi, G. & Ramasamy, S. miRNA and piRNA mediated Akt pathway in heart: antisense expands to survive. *Int J Biochem Cell Biol* **55,** 153–6 (2014).

50. Wang, Z., Liu, N., Shi, S., Liu, S. & Lin, H. The Role of PIWIL4, an Argonaute Family Protein, in Breast Cancer. *J Biol Chem* **291,** 10646–58 (2016).

51. Deretic, D. *et al.* rab8 in retinal photoreceptors may participate in rhodopsin transport and in rod outer segment disk morphogenesis. *J Cell Sci* **108 (Pt 1),** 215–24 (1995).

52. Knödler, A. *et al.* Coordination of Rab8 and Rab11 in primary ciliogenesis. *Proc Natl Acad Sci U S A* **107,** 6346–51 (2010).

53. Marshall, W. F. The cell biological basis of ciliary disease. *J Cell Biol* **180,** 17–21 (2008).

54. Nachury, M. V *et al.* A core complex of BBS proteins cooperates with the GTPase Rab8 to promote ciliary membrane biogenesis. *Cell* **129,** 1201–13 (2007).

55. Boylan, J. P. & Wright, A. F. Identification of a novel protein interacting with RPGR. *Hum Mol Genet* **9,** 2085–93 (2000).

56. Aldahmesh, M. A. *et al.* Mutations in LRPAP1 are associated with severe myopia in humans. *Am J Hum Genet* **93,** 313–20 (2013).

57. Esposito, T., Magliocca, S., Formicola, D. & Gianfrancesco, F. piR_015520 belongs to Piwi-associated RNAs regulates expression of the human melatonin receptor 1A gene. *PLoS One* **6,** e22727 (2011).

58. Kuintzle, R. C. *et al.* Circadian deep sequencing reveals stress-response genes that adopt robust rhythmic expression during aging. *Nat Commun* **8,** 14529 (2017).

59. Lu, Y. *et al.* Cancer/testis antigen PIWIL2 suppresses circadian rhythms by regulating the stability and activity of BMAL1 and CLOCK. *Oncotarget* **8,** 54913–54924 (2017).

60. Ding, X. *et al.* The role of miR-182 in regulating pineal CLOCK expression after hypoxia-ischemia brain injury in neonatal rats. *Neurosci Lett* **591,** 75–80 (2015).

61. Saus, E. *et al.* Genetic variants and abnormal processing of pre-miR-182, a circadian clock modulator, in major depression patients with late insomnia. *Hum Mol Genet* **19,** 4017–4025 (2010).

62. Tang, X. *et al.* Glycogen synthase kinase 3 beta inhibits microRNA-183-96-182 cluster via the β-Catenin/TCF/LEF-1 pathway in gastric cancer cells. *Nucleic Acids Res* **42,** 2988–98 (2014).

63. Gou, L.-T. *et al.* Pachytene piRNAs instruct massive mRNA elimination during late spermiogenesis. *Cell Res* **24,** 680–700 (2014).

64. Yin, H. & Lin, H. An epigenetic activation role of Piwi and a Piwi-associated piRNA in Drosophila melanogaster. *Nature* **450,** 304–308 (2007).

65. Watanabe, T. & Lin, H. Posttranscriptional Regulation of Gene Expression by Piwi Proteins and piRNAs. *Mol Cell* **56,** 18–27 (2014).

66. Martinez, V. D. *et al.* Non-coding RNAs predict recurrence-free survival of patients with hypoxic tumours. *Sci Rep* **8,** 152 (2018).

67. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17,** (2011).

68. Griffiths-Jones, S. The microRNA Registry. *Nucleic Acids Res* **32,** 109D–111 (2004).

69. Finn, R. D. *et al.* The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res* **44,** D279-85 (2016).

70. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* **10,** R25 (2009).

71. Li, N. *et al.* Global profiling of miRNAs and the hairpin precursors: insights into miRNA processing and novel miRNA discovery. *Nucleic Acids Res* **41,** 3619–34 (2013).

72. Heberle, H., Meirelles, G. V., da Silva, F. R., Telles, G. P. & Minghim, R. InteractiVenn: a web-based tool for the analysis of sets through Venn diagrams. *BMC Bioinformatics* **16,** 169 (2015).

73. Bailey, T. L. *et al.* MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* **37,** W202-8 (2009).

74. Volders, P.-J. *et al.* LNCipedia: a database for annotated human lncRNA transcript sequences and structures. *Nucleic Acids Res* **41,** D246-51 (2013).

75. Volders, P.-J. *et al.* An update on LNCipedia: a database for annotated human lncRNA sequences. *Nucleic Acids Res* **43,** D174–D180 (2015).

76. Solovyev, V., Kosarev, P., Seledsov, I. & Vorobyev, D. Automatic annotation of eukaryotic genes, pseudogenes and promoters. *Genome Biol* **7 Suppl 1,** S10.1-12 (2006).

77. Chekmenev, D. S., Haid, C. & Kel, A. E. P-Match: transcription factor binding site search by combining patterns and weight matrices. *Nucleic Acids Res* **33,** W432-7 (2005).

78. Enright, A. J. *et al.* MicroRNA targets in Drosophila. *Genome Biol* **5,** R1 (2003).

79. Betel, D., Wilson, M., Gabow, A., Marks, D. S. & Sander, C. The microRNA.org resource: targets and expression. *Nucleic Acids Res* **36,** D149-53 (2008).

80. John, B. *et al.* Human MicroRNA Targets. *PLoS Biol* **2,** e363 (2004).

81. Busk, P. K. A tool for design of primers for microRNA-specific quantitative RT-qPCR. *BMC Bioinformatics* **15,** 29 (2014).

82. Cirera, S. & Busk, P. K. in *Methods in molecular biology (Clifton, N.J.)* **1182,** 73–81 (2014).

Acknowledgements

We acknowledge the financial assistance provided by Department of Science and Technology under the project ''SR/FT/LS-145/2010'' and Department of Biotechnology under the project DBT-Neurobiology Taskforce - "BT/PR5055/MED/30/761/2012". The funding body is not involved in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript. We thank Indian Council of Medical Research project ''BMS/ FW/BIOCHEM/2015-23270/OCT-2015/24/TN/PVT'' and Max Planck-India mobility grant 'IGSTC/MPG/FS (SC)2012'. CS thanks the support from University Grants Commission, New Delhi for the award of Assistant Professorship under its Faculty Recharge Program (UGC-FRP) and for the start-up grant.

Author contributions

SS designed, performed the *in vitro* experiments; part of *in silico* experiments and wrote the manuscript. SN performed the *in silico* experiments. PG and AJ interpreted the data and reviewed the manuscript. CS conceived the idea, received the funding, planned the experiments and reviewed the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Figure legends

Fig 1. Workflow of piRNA identification. The flowchart describes the step by step protocol of piRNA analysis using the small RNA sequencing dataset of human retina and retinal pigment epithelium.

Fig 2. In silico analysis for identification of piRNAs in human retina and retinal pigment epithelium (RPE). a) Heatmap shows the expression of the identified 102 piRNAs in human retina (16 datasets) and RPE (2 dataset). b) Heatmap depicts the differential expression of piRNAs in retina and RPE. Transcript per million values (TPM) are used to generate heatmap using Morpheus. c) Venn diagram depicts the unique and commonly expressed piRNAs in human retina and RPE, brain, cardiac progenitors and cardiac fibroblasts, ovary and testis.

Fig 3. Genomic origin of piRNAs. a) Bar graph depicts the data obtained from piRNAQuest with the number of piRNAs that originate from the mentioned genomic regions. b) A representative list of identified piRNAs with piRNABank ID, sequence and genomic origins are listed in the table. This list also includes the 5 piRNAs validated in the study.

Fig 4. Enriched motifs and regulators of piRNAs. a) Logos with the corresponding E-values represent the motifs identified by MEME. Function and cellular localization of the motifs predicted by GOMo are also shown. b) Bar graph represents the number of transcription factor binding sites observed in the upstream sequences of 102 piRNAs.

Fig 5. Validation of piRNA expression in human retina and RPE tissues. a) Bar graph represents the average ΔCq values of piRNAs (*piR-31068, piR-35411, piR-36742, piR-60576 and piR-62011*) obtained using qRT-PCR from retina and RPE tissues. Since the bar graph shows the average ΔCq values, the smaller bar represents higher expression and vice versa. The Cq values were normalized with U6. b) RTL-P analysis confirms the 3`-terminal 2`-O-methylation of piRNAs. Amplification of piRNAs with unanchored primer at low dNTP

concentration was reduced whereas the miRNAs did not show any significant difference between unanchored and anchored primers. (RTL-P – Reverse Transcription at Low dNTP concentration followed by PCR; UAP – Unanchored primer; AP – Anchored primer)

Fig 6. Potential targets of piRNAs. a) Piechart shows the percentage of putative targets of piRNA in each category predicted by miRanda algorithm. b) Downregulation of HIWI2 is shown by Western blot in Si-Control and Si-HIWI2 ARPE19 cells. βACTIN is used as loading control. c) Expression of piRNAs in Si-HIWI2 ARPE19 cells is shown as fold change obtained by qRT-PCR. U6 levels are used for normalization. d) Agarose gel image showing the expression of *SNARE*s and *Rab*s in Si-Control and Si-HIWI2 cells. *GAPDH* is used as the loading control. e) Bar graph represents the fold change of *STX1A, STX6, STX16, Rab5, Rab8, VAMP7* and *VAMP8* in Si-HIWI2 ARPE19 cells obtained by qRT-PCR. *18S rRNA* is used for normalization. f) Bar graph shows the fold change of *LRPAP1, RPGRIP1* and *TIMELESS* in Si-HIWI2 ARPE19 cells. The expression is normalized with *18S rRNA*. (*p<0.05; **p<0.01; ***p<0.001; Student's *t* test was used to analyse the statistical significance between Si-Control and Si-HIWI2)

Fig 7. piRNAs are elevated under oxidative stress. Fold change of piRNAs obtained by qRT-PCR from the cells subjected to oxidative stress using 200 μM H_2O_2 are depicted as bar graph. U6 is used for normalization. (*p<0.05; **p<0.01; Student's *t* test was used to analyse the statistical significance between control and H_2O_2 treated cells)

Fig 8. Hypothetical model for pleiotropic functions of piRNAs. The model depicts the multiple functions of piRNAs in RPE in (1) altering the expression of tight junctions; (2 & 5) the regulation of sensory perception by influencing *RPGRIP1* and *LRPAP1;* (3) intracellular trafficking by affecting the expression of *STX1A, Rab5, Rab8, VAMP7 and VAMP8*; (4) Regulation of circadian clock by acting on *TIMELESS*. Regulation of these processes by

piRNAs may include miRNAs as well. Gaining deeper understanding of the exact molecular

mechanism will expand our knowledge on pleiotropic effects of piRNAs in somatic cells. (SE:

Sorting Endosome; FR: Fast Recycling; SR: Slow Recycling; LE: Late Endosome)

Table 1. Chromosomal distribution of piRNAs.

Chromosome	Number of piRNA clusters	Number of piRNAs in each cluster
Chromosome 1	3	Cluster 1 – 6 piRNAs Cluster 2 – 4 piRNAs Cluster 3 – 5 piRNAs
Chromosome 2	2	Cluster 1 – 5 piRNAs Cluster 2 – 4 piRNAs
Chromosome 3	1	Cluster 1 – 3 piRNAs
Chromosome 5	2	Cluster 1 – 5 piRNAs Cluster 2 – 9 piRNAs
Chromosome 6	4	Cluster 1 – 7 piRNAs Cluster 2 – 7 piRNAs Cluster 3 – 5 piRNAs Cluster 4 – 4 piRNAs
Chromosome 11	1	Cluster 1 – 4 piRNAs
Chromosome 14	1	Cluster 1 – 3 piRNAs
Chromosome 16	2	Cluster 1 – 6 piRNAs Cluster 2 – 4 piRNAs
Chromosome 17	1	Cluster 1 – 5 piRNAs
Chromosome 18	1	Cluster 1 – 3 piRNAs
Mitochondrial genome	2	Cluster 1 – 6 piRNAs Cluster 2- 4 piRNAs
Chromosome 4, 7, 8, 9, 10, 12, 13, 15, 19,20, 21, 22, X and Y chromosome	No clusters.	0-4 piRNAs

Table 2. Primer sequences used for RT and qRT-PCR

Name	Forward primer (5'- 3')	Reverse primer (5'- 3')
piR-31068	cattggtggttcagtggtag	ccagtttttttttttttttgcgaga
piR-35411	gctcagtcggtagagca	ggtccagttttttttttttttctca
piR-36742	tttccgtagtgtagtggtca	tccagtttttttttttttttgcga
piR-60576	ccgtagtgtagtggttatcac	cagtttttttttttttttggcgaa
piR-62011	gtttggcaatggtagaactca	gtccagttttttttttttttacctca
miR-26a	gcagttcaagtaatccaggatag	ggtccagtttttttttttttttagc
5.8S rRNA	tcgtgcgtcgatgaagaa	gaagtgtcgatgatcaatgtgtc
U6	gcttcggcagcacatatactaaaat	cgcttcacgaatttgcgtgtcat
LRPAP1	ggcgaagacctctgggaaat	gtctccagcaggacgttgta
RPGRIP1	cacagcccctccatcgttta	tgatgtgggcactgttaggc
TIMELESS	atgagacacgagatgtgcgg	gtgaggatgggcagaaggtc
STX1A	gaccgcttcatggatgagtt	cttggaacgaactttgtttgc
STX2	ggaagaatgatgatggagac	agaaagaatgatgctgtggt
STX3	tgctatcgacaacacggctt	ctggaatcggtgcagagaga
STX4	tgagctgcacgacatattca	tgtgcgacattatccaacca
STX6	ggtacagaaagcagtcaacac	tcatcaaggtcctctagatcc
STX7	aacacctcaagattcacctg	gttgtgaactctgccactaag
STX8	aagcttaccgtgacaatcag	aaggatgccagaagtagtctc
STX12	atgagccagctaggaactaag	gttccttctgaagtctctgct
STX16	cagcttagatccagaagcag	ctgagtgatctcttgggtagtt
STX17	gtccgaaaggatgacctagta	agtcatggatctggtcaaag
SNAP23	ggacatgagagagacagagaa	gcctggctgtttagatactac
SNAP29	agaacaggaagcaaagtacc	ctgtcgatcttctggtgatag
Rab1	gttacttctgattggcgact	catgggctcctctgtaataa
Rab3	cggatcagaacttcgactac	agcttgatcctcttgtcgt
Rab5	atacagctggtcaagaacga	ttaggacttgcttgcctct
Rab6	tagcctcattcccagttaca	cacttcctctttctgttctgac
Rab8	atcacaacggcctactacag	ccttggaaacttgtctcttg
Rab11	gagctgtaggtgccttattg	gctcttgcttcatctgtagg
VAMP3	aatcgaagacttcagcagac	cagagagcttctggtctcttt
VAMP7	cgagttctcaagtgtcttagc	agcaagatttctgctggtag
VAMP8	gaggtggagggagttaagaat	acaatcacgcagataaggac
18S rRNA	aacccgttgaaccccatt	ccatccaatcggtagtagcg
UAP	caggtccagtttttttttttttttgc	
AP	caggtccagttttttttttttttt	

Fig 1

NGS datasets of human retina and retinal pigment epithelium were retrieved from ArrayExpress (E-MTAB-3792)

Cutadapt was used to trim the adapters and to extract the reads with PHRED score above 30.

The extracted reads were mapped with human non-coding RNA sequences using BOWTIE.

Mapped reads were eliminated to filter out the known miRNAs and other small ncRNAs.

Unmapped reads were mapped with the human genome.

Mapped reads were compared with the known piRNA sequences and were further taken for analysis.

Fig 2

Fig 3

a

b

piRNA bank ID	Sequence	Gene	Intron	Intergenic	5'UTR	3'UTR	Repeat
piR-31068	agcattggtggttcagtggtagaattctcgc	Vac14	lies within the intron	-	-	-	L1PA5/LINE/L1 and tRNAs Gly
piR-35411	gcccggctagctcagtcggtagagcatgag	NBPF10	lies within the intron	intergenic	-	-	tRNA Lys
piR-36742	gtttccgtagtgtagtggtcatcacgttcgc	-	-	intergenic	-	-	tRNA Val
piR-60576	ttccgtagtgtagtggttatcacgttcgcc	-	-	intergenic	-	-	tRNA Val
piR-62011	tttggcaatggtagaactcacactggtgaggt	MIR182	-	-	-	3'UTR of miR 182	-
piR-31104	agcccggctagctcagtcggtagagcatgaga	-	-	-	-	-	tRNA Lys
piR-31925	attggtggttcagtggtagaattctcgcctg	VAC14	lies within the intron	intergenic	-	-	tRNA Gly
piR-35982	ggcattggtggttcagtggtagaattctcgc	VAC14, COLQ	lies within the intron	intergenic	-	-	tRNA Gly
piR-43993	tccgtagtgtagtggttatcacgttcgcct		-	intergenic	-	-	tRNA Val
piR-60565	ttccctggtggtctagtggttaggattcggc	NBPF10, PLCH1	lies within the intron	Intergenic	-	-	tRNA Glu

Fig 4

a

S.No	Enriched motifs	E-value	piRNAs with the motif	Predicted function and cellular localization
1.

5.4 e^{-127}	38	Localizes in nucleolus and cytoplasm. Function: rRNA processing
2.		1.3 e^{-023}	14	Localizes in mitochondria
3.		2.2 e^{-016}	20	Localizes in extracellular region. Function: 1. G-protein coupled protein receptor protein signalling pathway 2. Sensory perception of smell.

b

Number of transcription factor binding sites:

c-Rel 101, HLF 91, Evi-1 86, CDP 83, Nkx2-5 83, CREB 81, AREB6 79, ARP-1 79, YY1 76, HFH-1 73, XFD-1 69, ZID 68, BSAP 66, Myogenin/NF-1 66, XFD-2 66, E47 65, Elk-1 65, NFκB 64, XFD-3 63, COUP-TF,HNF-4 62, NRF-2 60, AP-4 57, ATF 57, CDP-CR3 57, Oct-1 56, RORα1 56, STATx 54, RREB-1 53, XBP-1 53, E2F 51, NFκB(p65) 45, NFκB(p50) 42, Hox-1.3 39, E2 31, IRF-1 29, NF-E2 25, Olf-1 23

Fig 5

Fig 6

Fig 7

Fig. 8