# LCA robustly reveals subtle diversity in large-scale single-cell RNA-seq data

Changde Cheng[1], John Easton[1], Celeste Rosencrance[1], Yan Li[2], Bensheng Ju[1], Justin Williams[1], Heather L Mulder[1], Wenan Chen[1] & Xiang Chen*[1]

Author affiliation:

[1]Department of Computational Biology, St. Jude Children's Research Hospital, Memphis, Tennessee 38105, USA

[2]Division of Biostatistics, University of Minnesota, Minneapolis, Minnesota 55455, USA

*Correspondence should be addressed to Xiang Chen (xiang.chen@stjude.org).

## Abstract

Single-cell RNA sequencing has emerged as a powerful tool for characterizing the cell-to-cell variation and dynamics. We present Latent Cellular Analysis (LCA), a machine learning–based analytical pipeline that features a dual-space model search with inference of latent cellular states, control of technical variations, cosine similarity measurement, and spectral clustering. LCA has proved to be robust, accurate, scalable, and powerful in revealing subtle diversity in cell populations.

Single-cell RNA sequencing (scRNA-seq) enables the characterization of cell-to-cell

variation in transcript abundance, leading to a deeper understanding of the diversity of

cell types and the dynamics of cell states in populations of thousands to tens of

thousands of single cells[1-3]. Although scRNA-seq offers enormous opportunities and

has inspired a tremendous explosion of data-analysis methods for identifying

heterogeneous subpopulations, significant challenges can arise because of the

inherently high noise associated with data sparsity and the ever-increasing number of

cells sequenced. The current state-of-the-art algorithms have significant limitations. The

biological meaning of the cell-to-cell similarity learned by most machine learning–based

tools (such as Seurat[4], Monocle2[5], SIMLR[6], and SC3[7]) is difficult to interpret. Although

certain technical biases (e.g., cell-specific library complexity) have been recognized as

major confounding factors in scRNA-seq analyses[8], other technical variations (e.g.,

batch effects and systematic technical variations that are irrelevant to the biological

hypothesis being evaluated) have received minimal attention, even though they present

major challenges to the analyses[9]. Most methods employ a gene-selection step before

clustering analysis, based on the assumption that a small subset of highly variable

genes is useful for revealing cellular diversity. Although this assumption is valid in most

scenarios and reduces the data dimensionality, it potentially excludes genes that are

informative for separating subpopulations with subtle diversity. Also, in datasets with

strong batch effects, this selection step may result in a small set of retained genes being

dominated by batch effects. Several methods require the user to provide an estimation

of the number of clusters in the data, and this may not be readily available.

Furthermore, most methods have a high computational cost that will be prohibitive for

datasets representing large numbers of cells.

We describe Latent Cellular Analysis (LCA), an accurate, robust, and scalable

computational pipeline that will facilitate a deep understanding of the transcriptomic

states and dynamics of single cells in large-scale scRNA-seq datasets. LCA makes a

robust inference of the number of populations directly from the data (a user can specify

this with a priori information), rigorously models the contributions from potentially

confounding factors, generates a biologically interpretable characterization of the

cellular states, and better reveals the underlying population structures. Furthermore,

LCA addresses the scalability problem by learning a model from a subset of the sample,

after which a theoretical scheme is used to assign the remaining cells to identified

populations.

LCA employs a heuristic dual-space search approach (**Fig. 1**) to provide an optimal

model for subpopulation structure identification (including the removal of confounding

factors, the inference of the number of clusters and informative states, and a mapping

function from the expression vector to cluster membership). In the primary latent cellular

(LC) space, LCA bypasses the gene selection and performs LC state inference from the

global gene expression matrix, removes cellular states representing known confounding

factors (e.g., cell-specific library complexity and batch information), and measures cell-

to-cell similarity by using the cosine of the angle between the low-dimensional cellular-

state vectors. Spectral clustering is employed to derive a set of candidate clustering

models with a range of cluster numbers. Meanwhile, in the dual principle component

(PC) space, cells are projected into a low-dimensional space based on principal

component analysis (PCA) of the expression vector. The list of significant components

is determined by the Tracy-Widom test[10], and cell-to-cell similarity is measured by

correlation similarity in the PC space. LCA then ranks candidate clustering models

derived in the LC space by the silhouette index[11] measured in the PC space. Finally,

LCA retains informative cellular states from the selected clustering model(s) and uses

these states to update the final clustering solution(s). Although the "optimal model" is

not necessarily the solution with the best numerical score, it should be among the top-

ranked models. Therefore, LCA provides users with multiple top-ranked solutions for

biology-based evaluations.

We benchmarked LCA with four commonly used state-of-the-art scRNA-seq clustering

algorithms (SC3[7], Seurat[4], Monocle2[5], and SIMLR[6]) in 3000 simulated datasets. LCA

outperformed competing methods in subpopulation number inference, cluster

membership inference, and running time (**Supplementary Result 1**). By using publicly

available large-scale scRNA-seq datasets, we demonstrated that LCA identified

orthogonal LC states of biological importance and produced parsimonious and accurate

models that were highly consistent with biological knowledge (**Supplementary Result**

**2**).

LCA groups cells based on orthogonal LC states aligned with major differences among

cells (**Supplementary Result 2**), which enables control of technical variations (e.g.,

batch effects) without an explicit need of gene filtering. We evaluated the efficiency of

batch effect removal in a cancer cell dataset and further experimentally validated the

subpopulations revealed by LCA. Rhabdomyosarcoma is the most common soft-tissue

tumor in children and has two major histologic subtypes with different genomic

landscapes: *PAX3/PAX7-FOXO1* fusion–positive alveolar rhabdomyosarcoma (FP-

ARMS) and fusion-negative embryonal rhabdomyosarcoma (ERMS)[12]. LCA identified

two subpopulations in a scRNA-seq dataset for Rh41 cells (a commonly used human

*PAX3-FOXO1* FP-ARMS cell line). The *CD44* gene, which encodes a commonly used

cell surface marker with great prognostic and therapeutic potential[13, 14], appeared at the

top of the differentially expressed genes (DEGs) of the two subpopulations

(**Supplementary Fig. 1A**). Flow cytometry confirmed a bimodal expression pattern of

CD44 in Rh41 cells (**Supplementary Fig. 1B**). We first used bulk RNA-seq to profile

unsorted populations and CD44$^{high}$ and CD44$^{low}$ subpopulations sorted by fluorescence-

activated cell sorting. In addition to the differences among the sorted CD44$^{high}$, CD44$^{low}$,

and unsorted populations, the analysis revealed strong batch effects. Specifically,

samples in Batch 1 and those in Batches 2 and 3 were separated on the first PC,

whereas the biologically different populations (the unsorted population and sorted

subpopulations) were separated on the second PC (**Supplementary Fig. 1C**). We

collected scRNA-seq data for the unsorted populations in Batch 1 and the sorted

CD44$^{high}$ and CD44$^{low}$ subpopulations in Batch 2 (10x Genomics, Pleasanton, CA). The

libraries were sequenced at different depths for the three samples. The median

numbers of UMIs captured per cell were 8278, 6678, and 12,850 for the CD44$^{high}$

subpopulation, CD44$^{low}$ subpopulation, and unsorted population, respectively (Kruskal-

Wallis rank sum test, $P < 2.2 \times 10^{-308}$). With both batch effects and biological

difference among cell populations, LCA inferred five clusters, with cells in Batch 1 being

represented by Clusters 1 and 2 and cells in Batch 2 being grouped in Clusters 3 to 5

(**Supplementary Fig. 1D**). Similarly, SC3 inferred a 12-cluster structure. Consistent

with the strong batch effects observed in bulk RNA-seq analysis, when required to infer

a two-cluster model, both LCA and SC3 achieved a perfect separation based on the

batch information (NMI $=1$ and 0.99, respectively). Next, we evaluated the

association between the batch covariate and individual LC states. When controlled at

FDR $\leq 0.05$, the first LC state was the only one significantly associated with the batch

information ($P < 2.2 \times 10^{-308}$) (**Supplementary Fig. 1E**). After excluding this state,

LCA retrieved an optimal structure with two clusters. Of 7005 cells profiled in the sorted

CD44$^{low}$ subpopulation, 6733 (96.1%) were clustered into Cluster 2, suggesting that the

subpopulation was relatively pure. However, 1796 (26.6%) of 6757 of cells profiled in

the sorted CD44$^{high}$ subpopulation were grouped in Cluster 2 with the CD44$^{low}$ cells

(**Fig. 2A**). We evaluated the expression signatures for DEGs identified in bulk RNA-seq

($\log_2$FC $\geq 1$, FDR $\leq 0.05$, FPKM $\geq 1$ in at least one subpopulation). Of 1709 DEGs

(**Supplementary Table 1**), 356 were captured in at least 10% of the cells in one

sequenced population or inferred cluster. As expected, Clusters 1 and 2 had higher

average expression levels of genes overexpressed in the bulk-sorted CD44$^{high}$ and

CD44$^{low}$ subpopulations, respectively (**Fig. 2B**). Importantly, Cluster 2 cells from the

sorted CD44$^{high}$ subpopulation had significantly lower expression of those DEGs

overexpressed in the bulk CD44$^{high}$ subpopulation than did Cluster 1 cells from the

sorted CD44$^{high}$ subpopulation ($P < 2.2 \times 10^{-308}$) or the sorted CD44$^{low}$

subpopulation ($P = 7.5 \times 10^{-71}$). Analysis of DEGs overexpressed in the bulk

CD44[low] subpopulation showed the same pattern (**Fig. 2C**). These results suggested

that Cluster 2 cells in the sorted CD44[high] subpopulation more closely resembled

CD44[low] cells. Moreover, the differential expression pattern was essentially captured by

the first remaining LC state (Spearman correlation coefficient $= -0.90$,

$P < 2.2 \times 10^{-308}$), confirming the biological importance of the inferred LC states.

Of the three established molecular markers (*TFAP2B*, *MYOG,* and *NOS1*) for FP-

ARMS[15], *TFAP2B* and *MYOG* were overexpressed in the sorted CD44[low] subpopulation

(**Supplementary Table 1**). Moreover, known *PAX3-FOXO1* target genes were

significantly enriched in DEGs overexpressed in the CD44[low] subpopulation

(**Supplementary Table 2**). These results suggest that the CD44[high] subpopulation

represents a less-differentiated, stem-like cell subpopulation.  Although the exact

mechanism by which the distinct subpopulations develop warrants further investigation,

we conclude that LCA can control technical variations and reveal reliable transcriptome-

based heterogeneity.

**Acknowledgements**

This study was supported in part by the National Cancer Institute of the National

Institutes of Health under Award Number P30CA021765 and by ALSAC. We thank

Keith A. Laycock for editing the manuscript.

**References**

1.    Tang, F. et al. mRNA-Seq whole-transcriptome analysis of a single cell. *Nat*
*Methods* **6**, 377-382 (2009).
2.    Shapiro, E., Biezuner, T. & Linnarsson, S. Single-cell sequencing-based
technologies will revolutionize whole-organism science. *Nat Rev Genet* **14**, 618-
630 (2013).
3.    Stegle, O., Teichmann, S.A. & Marioni, J.C. Computational and analytical
challenges in single-cell transcriptomics. *Nat. Rev. Genet.* **16**, 133-145 (2015).
4.    Macosko, E.Z. et al. Highly parallel genome-wide expression profiling of
individual cells using nanoliter droplets. *Cell* **161**, 1202-1214 (2015).
5.    Trapnell, C. et al. The dynamics and regulators of cell fate decisions are revealed
by pseudotemporal ordering of single cells. *Nat. Biotechnol* **32**, 381–386 (2014).
6.    Wang, B., Zhu, J., Pierson, E., Ramazzotti, D. & Batzoglou, S. Visualization and
analysis of single-cell RNA-seq data by kernel-based similarity learning. *Nat.*
*Methods* **14**, 414-416 (2017).
7.    Kiselev, V.Y. et al. SC3: consensus clustering of single-cell RNA-seq data. *Nat*
*Methods* **14**, 483-486 (2017).
8.    Wagner, A., Regev, A. & Yosef, N. Revealing the vectors of cellular identity with
single-cell genomics. *Nat Biotechnol* **34**, 1145-1160 (2016).
9.    Hicks, S.C., Townes, F.W., Teng, M. & Irizarry, R.A. Missing data and technical
variability in single-cell RNA-sequencing experiments. *Biostatistics* (2017).
10.   Patterson, N., Price, A.L. & Reich, D. Population structure and eigenanalysis.
*PLoS Genet* **2**, e190 (2006).
11.   Rousseeuw, P.J. Silhouettes - a graphical aid to the interpretation and validation
of cluster-analysis. *Journal of Computational and Applied Mathematics* **20**, 53-65
(1987).
12.   Chen, X. et al. Targeting oxidative stress in embryonal rhabdomyosarcoma.
*Cancer Cell* **24**, 710-724 (2013).
13.   Li, F., Tiede, B., Massague, J. & Kang, Y. Beyond tumorigenesis: cancer stem
cells in metastasis. *Cell Res* **17**, 3-14 (2007).
14.   Yan, Y., Zuo, X. & Wei, D. Concise Review: Emerging Role of CD44 in Cancer
Stem Cells: A Promising Biomarker and Therapeutic Target. *Stem Cells Transl*
*Med* **4**, 1033-1043 (2015).
15.   Rudzinski, E.R. et al. Myogenin, AP2beta, NOS-1, and HMGA2 are surrogate
markers of fusion status in rhabdomyosarcoma: a report from the soft tissue
sarcoma committee of the children's oncology group. *Am J Surg Pathol* **38**, 654-
659 (2014).
195

**ONLINE METHODS**

**Latent cellular states**

The input to LCA is a gene expression matrix in which each column is a cell, and each row is a gene/transcript. In UMI-based platforms, the expression level of a gene in a cell is divided by the total expression in that cell to generate a relative expression matrix (T). In read-count based platforms, T can be derived from size factor normalized expression measures. The relative expression matrix is then log-transformed after adding a zero-correction term:

$$X = \log(T + \epsilon),$$

where $\epsilon$ is an arbitrarily small number.

We obtain the LC states from a singular value decomposition (SVD) of $\mathbf{X}$.

$$\mathbf{X} = \mathbf{G} \, \mathbf{\Lambda} \, \mathbf{S}^T,$$

where $\mathbf{S}$ is a cell–by–LC states matrix. We note that under certain conditions, $\mathbf{S}$ is the same as the loading matrix from the principle component analysis (PCA) result of $\mathbf{X}$[16, 17].

**Determination of significant LC states**

We apply the Tracy-Widom test to associated eigenvalues to determine which LC states are significant[10, 18, 19]. The LC state associated with the eigenvalue $\lambda$ is significant if it is significantly different ($P < 0.05$) from the Tracy-Widom distribution, with

$$\mu(n,m) = \frac{(\sqrt{m-1}+\sqrt{n})^2}{m}$$

$$\sigma(n,m) = \frac{(\sqrt{m-1}+\sqrt{n})}{m}\left(\frac{1}{\sqrt{m-1}+\sqrt{n}}\right)^{\frac{1}{3}},$$

where $n$ is the total number of genes and $m$ is the total number of LC states. We then

discard all the LC states that are not significant or strongly associated with known

technical variations (e.g., cell-specific library complexity and batch information), leading

to a lower-dimensional cell–by–LC states matrix $\mathbf{S}$.

**Distance calculation**

Distances between cells in $\mathbf{S}$ (the cell–by–LC states matrix) are calculated using cosine

distance:

$$K_{a,b} = 1 - \frac{\sum_{i=1}^{p} S_{a,i}\, S_{b,i}}{\sqrt{\sum_{i=1}^{p} S_{a,i}^2}\,\sqrt{\sum_{i=1}^{p} S_{b,i}^2}}$$

where $S_{a,i}$ and $S_{b,i}$ represent LC state $i$ for cells $a$ and $b$, and $p$ is the total number of

retained LC states.

**Spectral clustering**

We perform spectral clustering [20] on the resulted distance matrix to derive a set of

candidate clustering models with a range of cluster numbers (i.e., 2 – 20 by default).

**Distance measure in the PC space**

The cell-gene relative expression matrix was decomposed by PCA and the significant

components were determined using the Tracy-Widom test. Distance between cells was

measured by the correlation distance of significant components.

**The optimal number of clusters and informative cellular states**

We rank the candidate clustering solutions (with different number of clusters) by the

silhouette score[11] measured in the PC space. With 2 or more clusters, the silhouette

measures the similarity of an individual to its cluster, as compared to other clusters. For

each cell, let $d_b$ be the lowest dissimilarity to any other cluster and let $d_w$ be the

average dissimilarity to other cells in its cluster. We calculate the silhouette as

$$Silhouette = \frac{d_b - d_w}{\max(d_b, d_w)}.$$

We assign a silhouette score of zero for the default solution of one cluster. An end-user

may evaluate the top candidate solutions to determine the optimal number of solutions

or specify it with a priori biological knowledge.

With the selected number of clusters, we retain LC states that show significant

difference among candidate clusters, update the distance matrix and derive the final

clustering solution.

**Fast processing of large numbers of cells**

We classify unknown cells $\mathbf{X}'$ efficiently by projecting them into space spanned by the

informative LC states learned from a representative sample:

$$S' = \Lambda_l^{-1} G_l^T X',$$

250    where $\Lambda_l$ is calculated from the original $\Lambda$ by removing those rows and columns that are

251    not associated with the informative LC states, and $G_l$ is calculated from $G$ by removing

252    those columns that are not associated with the informative LC states. We can then

253    calculate the cosine similarity between $S'$ and an "average" cell from each cluster and

254    find the cluster with the maximum similarity for each unknown cell.

**Simulation of different cell types**

We simulated different cell types from a relatively homogeneous population of cells. We

started from the scRNA-seq dataset for sorted CD44$^{\text{high}}$ Rh41 cells. The data

represented 6757 cells with 20,709 genes detected, with 86% of the entries in the data

matrix being zeroes. We considered two factors that have substantial effects on the

performance of clustering analysis: the sample size and the degree of difference

between cell types.

We tested five different sample sizes (the total number of cells analyzed): 250, 500,

1000, 2000, and 4000 cells. We sampled the selected number of cells from the original

set of cells then randomly partitioned them into three sets of variable size, ranging

between 10% and 50% of the total number of cells.

To test the effect of the degree of difference between cell types, we varied the number

of DEGs and the simulated difference in gene expression. We chose three sets of

genes such that the average expression levels for each set were $g_1$, $g_2$, and $g_3$, with

$\frac{g_{i+1}}{g_i} = 2, i = 1,2$. We then rotated the labels of the three sets of genes to make three

different sets of cells with perturbed expression: $g_1, g_2, g_3$; $g_2, g_3, g_1$; and $g_3, g_1, g_2$. We

tested two different numbers of DEGs: 300 and 1200. The simulation setting is

summarized in **Supplemental Table 3**. For each set of parameters, we generated 100

random samples for testing.

**Normalized mutual information and variation of information**

We evaluate the performance of clustering against true labels of cells by using

normalized mutual information (NMI)[21] and variation of information (VI)[22]. We use the R

package *igraph* to calculate both metrics[23].

**Benchmarking**

We used SC3 (Version 1.4.2), Seurat (Version 2.0.0), Monocle2 (Version 2.4.0) and

SIMLR (Version 1.2.1) for comparisons with LCA. We used the default settings for all

the packages tested. For SIMLR, when the size of the sample was 1000 or larger, we

used SIMLR_Large_Scale() instead of SIMLR().

**Rh41 single-cell dataset**

The human alveolar rhabdomyosarcoma cell line Rh41 was grown in culture in a 5%

$CO_2$ incubator in 75-cm$^2$ vented flasks containing DMEM medium supplemented with

10% FBS and 2× glutamine until the cells reached 75% confluence at approximately

$3.6 \times 10^6$ cells. The cells were detached from the flask with 7 mL of 1× citrate saline to

which 7 mL of DPBS was added. The cell suspension was then centrifuged at $300 \times g$

for 7 min, and the cell pellet was resuspended in 300 µL of blocking buffer (rat IgG/PBS)

and incubated on ice for 30 min. An aliquot of 50 µL of the cells in blocking buffer was

transferred to a separate tube for the isotype control. The cells were washed with 1 mL

of staining buffer (5% BSA/PBS) and centrifuged at $300 \times g$ for 5 min. The pellet,

which contained approximately $3 \times 10^6$ cells, was then incubated with rat IgG2B anti-

CD44–Alexa Fluor 488 antibody (R&D Systems) in staining buffer (15 µL antibody + 135

µL of staining buffer) on ice for 30 min. For the isotype control, approximately 600,000

cells were incubated with 5 µL of rat IgG2B–Alexa Fluor 488 (R&D Systems,

Minneapolis, MN) + 45 µL of staining buffer on ice for 30 min. After the incubation, both

sets of cells were collected by centrifugation, washed with 1 mL of staining buffer as

described above, and resuspended in staining buffer. Flow cytometric analysis was then

used to identify the fractions corresponding to the CD44 [high] and CD44 [low] populations.

For the single-cell experiment, Rh41 cells were grown in culture, harvested, and

washed in DPBS as described above. They were then resuspended in PBS/0.2% BSA

at a concentration of $1 \times 10^6$ cells/mL. The 10x Genomics Single Cell platform

performs 3′ gene expression profiling by poly-A selection of mRNA within a single cell,

which uses a cell barcode and UMIs (unique molecular identifiers) for each transcript.

Single-cell suspensions were loaded onto the Chromium Controller according to their

respective cell counts to generate approximately 6000 partitioned single-cell GEMs (Gel

Bead-in-EMulsions). The library was prepared using the Chromium Single Cell 3′ v2

Library and Gel Bead Kit (10x Genomics) in accordance with the manufacturer's

protocol. The cDNA content of each sample after cDNA amplification for 12 cycles was

quantified, and the quality was checked by High-Sensitivity DNA chip analysis on an

Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA) at a dilution of 1:6.

This quantification was used to determine the final library amplification cycles in the

protocol, which were calculated out to 12 cycles. After library quantification and a quality

check by DNA 1000 chip (Agilent Technologies), samples were diluted to 3.5 nM for

loading onto the HiSeq 4000 sequencer (Illumina) with a 2 × 75-bp paired-end kit,

using the following read length: 26 bp Read1 (10x cell barcode and UMI), 8 bp i7 Index

(sample index), and 98 bp Read2 (insert). In total, 518 million, 237 million, and 154

million reads were obtained for unsorted, CD44$^{low}$, and CD44$^{high}$ populations,

respectively. The Cell Ranger 2.0.1 Single-Cell Software Suite (10x Genomics) was

implemented to process the raw sequencing data from the Illumina HiSeq run. This

pipeline performed de-multiplexing, alignment (GRCh38/STAR), and barcode

processing to generate gene-cell matrices used for downstream analysis.

After matrix generation, the ribosomal and mitochondria-related genes were filtered.

**Rh41 bulk RNA-seq dataset**

RNA was isolated from the sorted subpopulations by using Trizol (Thermo Fisher

Scientific) in accordance with the manufacture's recommendations. RNA libraries were

prepared with the KAPA RNA HyperPrep Kit with RiboErase (Roche), using the

recommended conditions. Briefly, 200 ng of total RNA was used as input for

fragmentation, reverse transcription, and second-strand synthesis. After clean-up, end

repair, and A-tailing, NEXTflex adapters (Bioo Scientific, Austin, TX) were ligated to the

fragments, and this was followed by 12 cycles of PCR amplification on a C1000

Thermal Cycler (Bio-Rad). Paired-end sequencing was performed (151 bases per read)

on a HiSeq 4000 system (Illumina). Three replicates were generated. HTSeq[24] was

used to produce the count data, and edgeR[25] was used for the DE analysis with TMM

normalization. Each replicate was coded as a pair of CD44$^{high}$ and CD44$^{low}$ in the

analysis.

**Statistical analysis**

If not specifically stated, the statistical significance was evaluated by the Wilcoxon rank

sum test.

**Availability of data and material**

The Rh41 scRNA-seq dataset generated during the current study is available from the

corresponding author upon reasonable request. The bulk RNA-seq data for sorted

CD44$^{high}$ and CD44$^{low}$ subpopulations have been deposited in the European

Bioinformatics Institute (EMBL-EBI) under accession number EGAS00001002868. The

LCA framework is described in detail in supplementary methods. An R package for

implementing LCA has been developed and will be made generally available in

conjunction with publication of this manuscript.

16.    Pearson, K. On lines and planes of closest fit to systems of points in space. *The
London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*
**2**, 559-572 (1901).

17.  Hotelling, H. Analysis of a complex of statistical variables into principal
components. *Journal of Educational Psychology* **24**, 417-441 (1933).
18.  Tracy, C.A. & Widom, H. Level-spacing distributions and the Airy kernel. *Physics*
*Letters B* **305**, 115-118 (1993).
19.  Johnstone, I.M. On the distribution of the largest eigenvalue in principal
components analysis. *Ann Stat* **29**, 295-327 (2001).
20.  Ng, A.Y., Jordan, M.I. & Weiss, Y. in Advances in neural information processing
systems 849-856 (2002).
21.  Danon, L., Diaz-Guilera, A., Duch, J. & Arenas, A. Comparing community
structure identification. *J Stat Mech-Theory E* (2005).
22.  Meilă, M. Comparing clusterings—an information based distance. *Journal of*
*Multivariate Analysis* **98**, 873-895 (2007).
23.  Csardi, G. & Nepusz, T. The igraph software package for complex network
research. *InterJournal* **Complex Systems**, 1695 (2006).
24.  Anders, S., Pyl, P.T. & Huber, W. HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).
25.  Robinson, M.D., McCarthy, D.J. & Smyth, G.K. edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics*
**26**, 139-140 (2010).
382
383

**Figure Legends**

**Figure 1**. Overview of the workflow of LCA. LCA infers LC states from full expression

matrices. Explicit gene filtering is not necessary. LCA converts the raw transcript count

data to gene fractions and then performs log transformations. The algorithm features a

dual-space model search, generating candidate clustering models based on the cosine

similarity matrix in the LC space. Candidate models are then ranked based on the

silhouettes measured in the PC space.

**Figure 2**. LCA analysis of Rh41 cells, correcting for batch effects. (A) t-SNE plots of

clustering results, colored according to source ID or cluster ID. (B) Cellular expression

patterns of DEGs identified in bulk RNA-seq. (C) Violin plot of expression patterns of

DEGs from bulk RNA-seq in cells classified by source and cluster ID.

**Figure 1.** Overview of the workflow of LCA.

**Figure 2.** LCA analysis of Rh41 cells, correcting for batch effects.

**Supplementary Information**

**Supplementary Result 1**

**Benchmarking of LCA on simulated datasets**

By using extensive simulations, we evaluated the performance of LCA with four

commonly used state-of-the-art scRNA-seq clustering algorithms (SC3[7], Seurat[4],

Monocle2[5], and SIMLR[6]). Instead of generating artificial datasets from a theoretical

distribution, we simulated three equally spaced cell subpopulations of various sizes by

shuffling a predetermined number of genes across cells from high-quality, UMI-based

scRNA-seq data. We simulated seven scenarios with increasing signal strength, varying

the number of simulated differentially expressed genes (DEGs), the fold change among

groups, and the average expression levels of the simulated DEGs. We also investigated

the effect of the sample size (from 250 to 4000 cells) on the performance of the

algorithms.

A typical clustering analysis partitions data points into meaningful subgroups without a

priori knowledge of the number of subsets or their compositions. An essential

component of clustering analysis is inferring the number of clusters from the data[26],

which presents an analytical challenge distinct from that posed by clustering with a

given number of clusters. Therefore, we first analyzed the accuracy and robustness of

the subpopulation number inference. Monocle2 and SIMLR were excluded from this

analysis because both algorithms require end-user input of the number of clusters.

**Supplementary Figure 2A** shows that although LCA started with a slightly off-target

estimation of the cluster number in datasets with very weak signal strength, it stably

converged to the correct number with increasing signal strength from the number of

cells, the number of DEGs, the fold difference, or the expression level. In contrast, using

SC3 or Seurat resulted in an upward bias in the estimation of the number of clusters,

and this bias increased with both the sample size and the signal strength. For example,

in simulated model #6, with the strongest signal strength, the actual cluster number

remained the same (three) for all sample sizes, but SC3 estimated medians of four,

eight, and sixteen clusters in datasets with 250, 1000, and 4000 cells, respectively.

Seurat estimated medians of three, four, and nine subpopulations in the same

respective datasets. Combining the results of all simulations, we find that LCA arrived at

the correct number of clusters in 70.2% of simulations, which is a significantly higher

percentage than was achieved with SC3 (5.6%, $P < 2.2 \times 10^{-16}$ [proportion test])

and Seurat (22.7%, $P < 2.2 \times 10^{-16}$ [proportion test]).

We used variation of information (VI[22], 0 for perfect matching) (**Supplementary Fig. 2B**)

and normalized mutual information (NMI[21], 1 for perfect matching) (**Supplementary Fig.**

**2C**) to measure the clustering accuracy. For methods with a feature for inferring the

number of clusters from the data (LCA, SC3, and Seurat), we compared the ground-

truth labels with inferred cluster membership from the optimal clustering models. For

methods without this inference feature (Monocle2 and SIMLR), the optimal models were

derived with the correct number of clusters (three). The comparison showed that LCA

outperformed other methods in most simulations (**Supplementary Fig. 2**). When all

3000 simulation datasets were combined, LCA achieved the highest overall accuracy

(average NMI = 0.756; VI = 0.555). Specifically, when LCA correctly inferred the

number of clusters (in 70.2% of all simulations), it produced near-perfect accuracy

(average NMI□=□0.941, VI□=□0.122). The accuracies of the competing algorithms

were consistently lower. Notably, SC3 achieved a relatively good performance

(NMI□=□0.807, VI□=□0.389) when it happened to find the correct number of clusters

(mostly in small datasets). However, its performance deteriorated for other datasets,

largely due to an overestimation of cluster numbers. The overall accuracy of SC3 was

significantly lower than that of LCA (NMI□=□0.583, $P□<□2.2□×□10^{-16}$; VI□=□1.203,

$P□<□2.2□×□10^{-16}$ [Wilcoxon signed-rank test]). More strikingly, even if provided with

the correct cluster number, Monocle2 attained high accuracy only for the dataset with

the strongest signal strength (simulated model #6), and its performance was suboptimal

compared to that of LCA (NMI□=□0.190, $P□<□2.2□×□10^{-16}$; VI□=□1.719,

$P□<□2.2□×□10^{-16}$ [Wilcoxon signed-rank test]). Similarly, the overall accuracies of

Seurat (NMI□=□0.415, $P□<□2.2□×□10^{-16}$; VI□=□1.545, $P□<□2.2□×□10^{-16}$ [Wilcoxon

signed-rank test]) and SIMLR (NMI□=□0.231, $P□<□2.2□×□10^{-16}$; VI□=□1.353,

$P□<□2.2□×□10^{-16}$ [Wilcoxon signed-rank test]) were significantly lower than that of

LCA.

In addition to the superior accuracy of LCA with respect to both cluster number and

membership assignment, its implementation renders exceptional scalability, which is an

attractive feature for scRNA-seq analysis of an ever-increasing number of cells. LCA

can derive a cell subpopulation structure by using a relatively small representative set

(training cells) sampled from the full data. LCA provides mathematical formulae with

which to project the remaining cells (testing cells) directly to the inferred low-

dimensional LC space, after which individual cells are assigned to the subpopulation

with the best similarity. Consequently, LCA runs at a low level of computational

complexity. The median overall running times for a 4000-cell dataset with 10% or 25%

training cells were 40.6 s and 202.9 s, respectively. We compared the accuracy of

clustering between the training cells (5%–25% of all cells) and the remaining testing

cells in two simulated models (**Supplementary Fig. 2C, D**, models #2 and #6). As

expected, the accuracy of the clustering model improved with the increase of training

cells in the simulated dataset #2 (weak signal strength) and remained at a near-perfect

level in simulated dataset #6 (strong signal strength). Strikingly, LCA achieved

comparable accuracies in the testing cells in both models and outperformed SC3 in

terms of accuracy and running time.

## Supplementary Result 2

### Evaluation of LCA in publicly available datasets

We evaluated the performance of LCA with large-scale datasets published by Zheng et al.[27] and Tirosh et al.[28]. Using the GemCode platform (10x Genomics, Pleasanton, CA), Zheng et al. generated reference scRNA-seq transcriptomes for subpopulations of peripheral blood mononuclear cells (PBMCs) purified via well-established cell surface markers[27]. We generated a large-scale purified T-cell dataset (representing 55,000 cells) by combining the CD4+ T-helper (CD4+ helper), CD4+/CD25+ regulatory T ($T_{reg}$), CD4+/CD45RO+ memory T ($T_{mem}$), CD4+/CD45RA+/CD25– naïve T (CD4+ ab T), CD8+ cytotoxic T (cytotoxic T), and CD8+/CD45RA+ naïve cytotoxic T (CD8+ ab T) cell subpopulations. We inferred the cellular states and subpopulation structure by using 10% and 25% of the full dataset, and we assigned the remaining cells to one of the inferred subpopulations. We evaluated the top three models for both runs, which were the 3-population, 4-population, and 5-population models. Despite a 9.6-fold difference in running time, both the 3-population and 5-population models from the two runs achieved high consistency (NMI $=$ 0.87 and 0.84, respectively). Although the 4-population models differed between runs, they represented two different subpopulation-merging orders from the 5-population model to the 3-population model. We selected the 5-population model learned from 25% cells (with 19 LC states) for further biological inference. Whereas the purified $T_{mem}$, CD4+ ab T, and CD8+ ab T cells contained cells mostly from a single subpopulation (Clusters 1, 3, and 4, respectively), different levels of heterogeneity were detected in the remaining purified populations, especially the CD4+ helper and cytotoxic T cells. Given the single surface-marker settings in the purification

of CD4+ helper and cytotoxic T cells, it is not surprising that we found substantial

heterogeneity in the two populations. Nevertheless, LCA inferred a parsimonious

subpopulation structure in this large dataset (compared to the 1108 subpopulations

predicted by SC3). We selected representative genes encoding surface markers,

transcription factors, and secreted effector molecules for 19 usual T-cell subsets[29] and

derived a PCA projection of the six purified populations and five inferred clusters based

on the average population/cluster expression level of individual marker genes. The first

PC largely described the differentiation status, and the second PC represented the

difference between the CD4+ subsets and CD8+ subsets. As expected, Clusters 3 and

4 were found next to the CD8+ ab T and CD4+ ab T cells. Cluster 1 was found near the

$T_{mem}$ population. Cluster 2 consisted mostly of $T_{reg}$ cells and a smaller fraction of CD4+

helper cells that was located adjacent to the $T_{reg}$ population. Both the CD4+ helper and

cytotoxic T-cell populations were split approximately equally between naïve and

differentiated cells and were spotted between their corresponding naïve clusters

(Cluster 3 for CD8+ cells and Cluster 4 for CD4+ cells) and differentiated clusters

(Clusters 1 and 2 for CD4+ cells and Cluster 5 for CD8+ cells). Analysis of selected

genes validated the separation of naïve and differentiated cells in the CD4+ helper and

cytotoxic T cells (**Supplementary Fig. 3**). An evaluation of the first LC state inferred

from the full expression data revealed a striking approximation to the expected

distribution of naïve and differentiated cells in both purified populations and inferred

clusters. Similarly, the second LC state recaptured the CD4+ and CD8+ difference in

the dataset. These results showed that LCA identified LC states of biological importance

and consequently produced a parsimonious and accurate model that was highly

consistent with biological knowledge.

We evaluated the performance of LCA in a second dataset published by Tirosh et al.[28],

who employed a stepwise approach to analyze separately the malignant and stromal

cells of 19 melanoma tumors captured on the C1 Fluidigm platform. We applied those

authors' cell-selection criteria[28] to generate a dataset with 1169 malignant cells from

eight tumors and 2588 nonmalignant (stromal) cells. LCA inferred 18 clusters with 53

LC states (**Supplementary Fig. 4**). Malignant cells dominated eight clusters, which

were further separated by the patient origin of the tumors (**Supplementary Fig. 4A**).

Among the stromal cells, distinct clusters were identified for B cells, macrophages,

cancer-associated fibroblasts, and endothelial cells (**Supplementary Fig. 4B**).

Moreover, LCA divided tumor-infiltrating T cells and natural killer cells into six clusters,

which was concordant with the supervised analysis of T cells based on surface markers

by Tirosh et al.[28]. Using the pre-defined marker genes for T-cell subsets and *MKI67* for

cell-cycle activities, we revealed characteristics of these T-cell populations

(**Supplementary Fig. 4C**). Cluster 1 was enriched for naïve CD4+ cells, whereas

Cluster 2 harbored mostly $T_{reg}$ and $T_{FH}$ cells. Although Clusters 3 and 5 both had

enriched signatures for cytotoxic T and exhaustive T cells, Cluster 3 had greater signal

strength in both signatures, consistent with the reported high correlation between

exhaustion markers and cytotoxic markers[28]. Cells in active cell cycles were grouped in

Cluster 4, which showed unique enrichment of *MKI67* expression. Lastly, natural killer

cells and T cells with weak cytotoxic activity and no exhaustion signatures were found in

Cluster 6. As a comparison, we applied SC3 to the same data, which resulted in an

estimation of 43 clusters. Although SC3 clustering of malignant cells was consistent with

the separation by patient, accuracy in the stromal component was lower than with LCA

(**Supplementary Table 4**). These results demonstrate the power of LCA to reveal

subtle diversities in different subpopulations of tumor-infiltrating T cells in the presence

of strong transcriptomic variations among malignant cells (from different patients) and

various stromal cells from different lineages (cancer-associated fibroblasts,

macrophages, B cells, and endothelial cells).

26. Tibshirani, R., Walther, G. & Hastie, T. Estimating the number of clusters in a
data set via the gap statistic. *Journal of the Royal Statistical Society Series B-*
*Statistical Methodology* **63**, 411-423 (2001).
27. Zheng, G.X. et al. Massively parallel digital transcriptional profiling of single cells.
*Nat Commun* **8**, 14049 (2017).
28. Tirosh, I. et al. Dissecting the multicellular ecosystem of metastatic melanoma by
single-cell RNA-seq. *Science* **352**, 189-196 (2016).
29. Dong, C. & Martinez, G.J. T cells: the usual subsets (poster).  (2010).
30. Kuleshov, M.V. et al. Enrichr: a comprehensive gene set enrichment analysis
web server 2016 update. *Nucleic Acids Res* **44**, W90-97 (2016).
568

**Supplementary Figure 1.** Single-cell analysis of Rh41 cells with strong batch effects.

(A) Distance matrix visualized as a heat map. (B) Flow cytometry confirmed a bimodal

expression pattern of CD44 in Rh41 cells. (C) The first two PCs for the bulk RNA-seq

data showed strong batch effects. (D) t-SNE plot of cells, including sorted and unsorted

samples, colored according to source/cluster ID. (E) The first LC state was significantly

associated with batch information.

**Supplementary Figure 1.**

**Supplementary Figure 2**. Benchmarking of LCA against four other methods with

simulated datasets. (A) The number of clusters inferred by LCA, SC3, and Seurat was

compared for seven simulated datasets: the signal strength increases from simulated

datasets 1 to 6. Each simulated dataset contained 100 randomly generated samples.

(B) Comparison of the clustering performance of LCA, SC3, Seurat, Monocle2, and

SIMLR, as measured by variation of information, across the seven simulated datasets.

For LCA, SC3, and Seurat, the number of clusters was inferred during the analysis. For

Monocle2 and SIMLR, we used the correct number of clusters. (C) Comparison of the

clustering performance as measured by normalized mutual information. (D) Clustering

performance scalability of LCA and SC3. Each panel represents the results obtained

with a different percentage of cells used for training (from 5% to 25%). The total number

of cells in the sample was 4000. The signal in this comparison was relatively weak. (E)

Comparison of the clustering performance scalability of LCA and SC3 when the signal

was strong. (F) Comparison of the running times of LCA and SC3.

**Supplementary Figure 2.** Benchmarking of LCA against four other methods with

simulated datasets.

**Supplementary Figure 3.** Investigation of PBMC heterogeneity by using LCA. We

applied LCA to data from Zheng et al.[25] for 55,000 cells representing a combination of

six sorted T-cell populations. LCA identified five cell populations with biologically

meaningful LC states. (A) Cell distance matrix, sorted by source ID. (B). Cell distance

matrix, sorted by LCA clustering results. (C) The first LC state captured the difference

between the naïve and differentiated T cells, whereas the second LC state captured the

difference between the CD4+ and CD8+ cells. (D) LCA further revealed heterogeneity in

the CD4+ helper T cells and CD8+ cytotoxic T cells.

**Supplementary Figure 3.** Investigation of PBMC heterogeneity by using LCA.

**Supplementary Figure 4.** Reanalysis of melanoma cellular data[28] with LCA. We

applied LCA to 1169 malignant cells from eight tumors and to 2588 stromal cells. LCA

identified 18 clusters with a clear biological interpretation. LCA further revealed subtle

diversity among the infiltrating T cells. (A) t-SNE plot of the clustering results for

malignant cells, contrasting the LCA results and original results. (B) t-SNE plot of the

clustering results for stromal cells, contrasting the LCA results and original results. (C)

Enrichment of gene markers in T-cell clusters.

**Supplementary Figure 4.** Reanalysis of melanoma cellular data with LCA.

622    **Supplementary Table 1**. Expression profile of differential expressed genes (in a

623    standalone excel file).

625    **Supplementary Table 2**. ChEA analysis by Enrichr[30] of 361 overexpressed genes in

626    the CD44[low] subpopulation of Rh41 (in a standalone excel file).

628    **Supplementary Table 3**. Summary of simulated datasets.

| Signal Scale | Number of Genes Perturbed | Mean Gene Fraction in Group 1 | Mean Gene Fraction in Group 2 | Mean Gene Fraction in Group 3 |
|---|---|---|---|---|
| Sim. #1 | 300 | 1e-5 | 2e-5 | 4e-5 |
| Sim. #2 | 300 | 2e-5 | 4e-5 | 8e-5 |
| Sim. #3 | 300 | 4e-5 | 8e-5 | 1.6e-4 |
| Sim. #4 | 1200 | 1e-5 | 2e-5 | 4e-5 |
| Sim. #5 | 1200 | 2e-5 | 4e-5 | 8e-5 |
| Sim. #6 | 1200 | 4e-5 | 8e-5 | 1.6e-4 |

631 **Supplementary Table 4**. The results of applying LCA and SC3 to the dataset of Tirosh

632 et al.[26], using normalized mutual information.

| | LCA | SC3 |
|---|---|---|
| All cells | 0.77 | 0.69 |
| Malignant cells | 0.99 | 0.99 |
| All stromal cells | 0.53 | 0.42 |
| All cells excluding T cells | 0.97 | 0.92 |
| Stromal excluding T cells | 0.90 | 0.71 |