Whole Exome Sequencing in 20,197 Persons for Rare Variants in Alzheimer Disease

Neha S. Raghavan, PhD [1,2], Adam M. Brickman, PhD[1,2,3], Howard Andrews, PhD[1,2,4], Jennifer J. Manly, PhD[1,2,3], Nicole Schupf, PhD[1,2,3,], Rafael Lantigua, MD[1,6], The Alzheimer's Disease Sequencing Project*, Charles J. Wolock, BA[8], Sitharthan Kamalakaran, PhD[8], Slave Petrovski, PhD[8,9], Giuseppe Tosto, MD, PhD [1,2], Badri N. Vardarajan, PhD[1,2,3], David B. Goldstein, PhD[3,6,8] and Richard Mayeux, MD[1,2,3,4,7]

KEYWORDS: Alzheimer's disease, Mutations, Whole Exome Sequencing

RUNNING TITLE: Ultra-Rare Variants and Alzheimer's disease

WORD COUNT: Abstract 250; Introduction 344; Discussion 1,125; Body 2,440

FIGURES AND TABLES: Figures 1; Tables 4; Supplemental Tables 2

CHARACTERS: Title 122; Running Head 43

[1]The Taub Institute for Research on Alzheimer's Disease and the Aging Brain, [2]The Gertrude H. Sergievsky Center, The Departments of [3]Neurology, [4]Psychiatry, [5]Systems Biology and [6]Medicine, at the College of Physicians and Surgeons, Columbia University, The New York Presbyterian Hospital, [7]The Department of Epidemiology, Mailman School of Public Health, [8]Institute of Genomic Medicine, Columbia University, The New York Presbyterian Hospital, New York, NY, USA, [9]AstraZeneca Centre for Genomics Research, Precision Medicine and Genomics, IMED Biotech Unit, AstraZeneca, Cambridge CB2 0AA, UK

*Members listed by Institution in the supplement

Correspondence:
Richard Mayeux, MD, MSc.
Department of Neurology
710 West 168th Street
Columbia University
New York, NY 10032
Phone: 212-305-2391
Email: rpm2@cumc.columbia.edu

Abstract

Objective

The genetic bases of Alzheimer's disease remain uncertain. An international effort to fully

articulate genetic risks and protective factors is underway with the hope of identifying potential

therapeutic targets and preventive strategies. The goal here was to identify and characterize

the frequency and impact of rare and ultra-rare variants in Alzheimer's disease using whole

exome sequencing in 20,197 individuals.

Methods

We used a gene-based collapsing analysis of loss-of-function ultra-rare variants in a case-

control study design with data from the Washington Heights-Inwood Columbia Aging Project,

the Alzheimer's Disease Sequencing Project and unrelated individuals from the Institute of

Genomic Medicine at Columbia University.

Results

We identified 19 cases carrying extremely rare *SORL1* loss-of-function variants among a

collection of 6,965 cases and a single loss-of-function variant among 13,252 controls (p = 2.17 x

10^{-8}; OR 36.2 [95%CI 5.8 - 1493.0]). Age-at-onset was seven years earlier for patients with

SORL1 qualifying variant compared with non-carriers. No other gene attained a study-wide

level of statistical significance, but multiple top-ranked genes, including *GRID2IP*, *WDR76* and

GRN, were among candidates for follow-up studies.

Interpretation

This study implicates ultra-rare, loss-of-function variants in *SORL1* as a significant genetic risk

factor for Alzheimer's disease and provides a comprehensive dataset comparing the burden of

rare variation in nearly all human genes in Alzheimer's disease cases and controls. This is the

first investigation to establish a genome-wide statistically significant association between

multiple extremely rare loss-of-function variants in *SORL1* and Alzheimer's disease in a large

whole-exome study of unrelated cases and controls.

Introduction

Alzheimer's disease (AD) is a highly prevalent disorder that dramatically increases in frequency with age, and has no effective treatment or means of prevention. While three causal genes, Amyloid Precursor Protein (*APP*), Presenilin 1 and 2 (*PSEN1* and *PSEN2*), have been established for early-onset AD (age of onset <65 years of age), the rest of the heritability is still unknown. Further, beyond Apolipoprotein E (*APOE*), which confers the greatest risk for late-onset AD (age of onset ≥65 years of age), there remains a large gap in the understanding of its causes. Identifying genetic variants that increase risk or protect against AD is considered an international imperative because of the potential therapeutic targets that may be revealed. Recent technological advances in genome-wide association studies and high throughput next-generation sequencing may help to implicate variants in genes in specific molecular pathways relevant to AD.

In this study, we used whole-exome sequencing to investigate all protein-coding genes in the genome focusing on ultra-rare (allele frequency less than 0.01%) and putatively deleterious variants. Rare variants are hypothesized to contribute to disease [1,2], and studies of complex traits in population genetic models indicate an inverse relationship between the odds ratio and effect size conferred by rare variants and low allele frequencies [3]. Thus, we searched for large effects conferred by putatively causal ultra-rare variants. Traditional single variant statistics can be underpowered because patients with similar clinical presentations possess distinct rare variants that inflict similar effects on the gene [4]. Gene-based collapsing analyses increase signal detection by aggregating individual qualifying variants within an *a priori* region (e.g., a gene), facilitating detection of genes associated with disease through a specific class of genetic variation (e.g., loss-of-function variants).

In order to maximize the ability to detect ultra-rare variants associated with AD, exome-

sequencing data of 20,197 cases and controls from the Washington Heights-Inwood Community

Aging Project (WHICAP), the Alzheimer's Disease Sequencing Project (ADSP) and unrelated

controls from the Institute of Genomic Medicine were systematically combined and analyzed

using a collapsing method with proven prior success in identifying disease associated genes [5, 6].

Methods

The three groups used in this study and their sequencing information are described below.

Washington Heights-Inwood Community Aging Project. The WHICAP study consisted of a

multi-ethnic cohort of 4,100 individuals followed over several years The cohort participants were

non-demented initially, 65 years of age or older, and comprised of non-Hispanic whites, African

Americans, and Caribbean Hispanics from the Dominican Republic. During each assessment,

participants received a neuropsychological test battery, medical interview, and were re-

consented for sharing of genetic information and autopsy. A consensus diagnosis was derived

for each participant by experienced clinicians based on NINCDS-ADRDA criteria for possible,

probable, or definite AD, or moderate or high likelihood of neuropathological criteria of AD [7,

8]. Every individual with whole-exome sequencing has at least a baseline and one follow-up

assessment and examination, and for those who have died, the presence or absence of

dementia was determined using a brief, validated telephone interview with participant

informants: the Dementia Questionnaire (DQ) [9] and the Telephone Interview of Cognitive Status

(TICS) [10]. 3,702 exome-sequenced WHICAP individuals were designated with case or control

status and included in this analysis. From the sequenced cohort, 27% died and less than 1%

were lost at follow-up.

Alzheimer's Disease Sequencing Project. The ADSP, developed by the National Institute on

Aging (NIA) and National Human Genome Research Institute (NHGRI) includes a large case-

control cohort of approximately 10,000 individuals [7]. The recruitment of these individuals was in

collaboration with the Alzheimer's Disease Genetics Consortium and the Cohorts for Heart and

Aging Research in Genomic Epidemiology Consortium. The details and rationale for the case-

control selection process have been previously described [7]. All cases and controls were at least

60 years old and were chosen based on sex, age and *APOE* status: 1) controls were evaluated

for their underlying risk for AD and for their likelihood of conversion to AD by age 85, based on

age at last examination, sex, and *APOE* genotype, and those with the least risk for conversion

to AD were selected, and 2) cases were evaluated for their underlying risk for AD based on age

at onset, sex, and *APOE* genotype and those with a diagnosis least explained by these factors

were selected [7]. Cases were determined either because they met NINCDS-ADRDA clinical

criteria for AD, or postmortem findings met moderate or high likelihood of neuropathological

criteria of AD [7, 8]. Autopsy data was available for 28.7% of the cases and controls used in the

analysis. Further, some cases were originally diagnosed clinically, subsequently died and had

neuropathological findings available after postmortem examination. Cases had documented age

at onset or age at death (for pathologically determined cases). Controls were free of dementia

by direct, documented cognitive assessment or neuropathological results. The ADSP group

consisted of European-Americans and Caribbean Hispanics. All data were available for

download for approved investigators at The National Institute on Aging Genetics of Alzheimer's

Disease Data Storage Site website (https://www.niagads.org/adsp/content/home). As part of the

ADSP, 116 non-Hispanic white WHICAP controls and 34 cases previously sequenced were

included here.

Additional Controls. The Institute for Genomic Medicine (IGM) (Columbia University Medical

Center, New York, NY) hosts an internal database of sequencing data collected from previously

exome-sequenced material. In this study, exome-sequencing data from 6,395 IGM controls

were utilized. All data used were previously consented for future control use from multiple

studies of various phenotypes. The cohort was made up of 55.7% healthy controls and 46.3%

with diseases not co-morbid with AD (disease classifications shown in Supplemental Table 1).

Although the cohort of controls were not enriched for any neurological disorder or diseases with

a known co-morbidity with AD, presence or future possibility of AD could not be excluded based

on the available clinical data. individuals with Age and *APOE* status were not available for these

participants. The cohort comprised of 70% non-Hispanic white individuals along with those of

African American, Hispanic, Middle Eastern, Asian and unknown descent.

Sequencing, Quality Control and Variant Calling

Whole-exome sequencing of the WHICAP cohort was performed at Columbia University. The

additional controls were sequenced at Duke University and Columbia University. Whole-exome

sequencing of the ADSP cohort was performed at The Human Genome Sequencing Center,

Baylor College of Medicine, Houston, Texas; The Broad Institute Sequencing Platform, The Eli

& Edythe L. Broad Institute of the Massachusetts Institute of Technology and Harvard

University, Cambridge Massachusetts and Washington University Genome Sequencing Center,

Washington University School of Medicine, Saint Louis, Missouri. ADSP raw files in the

sequencing read archive format were downloaded from the dbGAP database and

decompressed to obtain FASTQ files.

All data were reprocessed for a consistent alignment and variant calling pipeline consisting of

the primary alignment and duplicate marking using the Dynamic Read Analysis for Genomics

(DRAGEN) platform followed by variant calling according to best practices outlined in Genome

Analysis Tool Kit (GATK v3.6). Briefly, aligned reads were processed for indel realignment

followed by base quality recalibration and Haplotype calling to generate variant calls. Variant

calls were then subject to Variant Quality Score Recalibratrion (VQSR) using the known single

nucleotide variants (SNVs) sites from HapMap v3.3, dbSNP, and the Omni chip array from the

1000 Genomes Project. SNVs were required to achieve a tranche of 99.9% and indels a

tranche of 95%. Finally, read-backed phasing was performed to determine phased SNVs and

merge multinucleotide variants (MNVs) when appropriate. Variants were annotated using Clin-

Eff with Ensembl-GRCh37.73 annotations.

Quality thresholds were set based on previous work[5, 6], such that all resulting exome variants

had a quality score of at least 50, quality by depth score of at least 2, genotype quality score of

at least 20, read position rank sum of at least -3, mapping quality score of at least 40, mapping

quality rank sum greater than -10, and a minimum coverage of at least 10. SNVs had a

maximum Fisher's strand bias of 60, while indels had a maximum of 200. For heterozygous

genotypes, the alternative allele ratio was required to be greater than or equal to 25% and

variant from sequencing artifacts and exome variant server failures

(http://evs.gs.washington.edu/EVS) were excluded.

Quality control was performed on all sequencing data. Samples with less than 90% of the

consensus coding sequence (CCDS) covered at 10X and samples with sex-discordance

between clinical and genetic data were excluded from the analysis. Cryptic relatedness testing

was performed using KING, and second degree or closer (relatedness threshold of 0.0884 or

greater) relatives were removed with preferential retention of cases over controls and

subsequently samples with higher average read-depth coverage.

The consensus coding sequence [11] (CCDS) annotated protein-coding region for each gene

(n=18,834) was tabulated as either carrying or not carrying a qualifying variant for every

individual. Qualifying variants were defined for a loss-of-function model: stop gain, frameshift,

splice site acceptor, splice site donor, start lost, or exon deleted variants. A negative control

analysis was performed defining qualifying variants as synonymous variants to detect potential

biases in variant calling between the cases and controls separately for each of the top four

genes. The minor allele frequency threshold was 0.01% internally and within African American,

Latino and Non-Finnish European populations from the Exome Aggregation Consortium[12]

(ExAC release version 0.3.1). The allele frequency thresholds use a "leave-one-out" method for

the combined test cohort of cases and controls such that the minor allele frequency of each

variant was calculated using all individuals except for the index sample under investigation.

Thus, the maximum instances of a single variant a gene in our sample of 20,197 was five. A

dominant model was defined such that one or more qualifying variant(s) in a gene qualified the

gene.

An important aspect of the collapsing analysis methodology is the reduction of variant calling

bias due to coverage differences between cases and controls. To ensure balanced sequencing

coverage of evaluated sites between cases and controls, we imposed a statistical test of

independence between the case/control status and coverage. For a given site, consider s total

number of cases, t total number of controls and x number of cases covered at 10X, y number of

controls covered at 10x. We model the number of covered cases X as a Binomial random

variable:

177                    $X \sim bin(n = \text{number covered samples}, p = P(\text{case}|\text{covered}))$

178   If case/control status and coverage status are independent, then:

179                    $P(\text{case}|\text{covered}) = P(\text{case}) = s/(s+t)$

180   We can test for this independence by performing a two-sided Binomial test on the number of

181   covered samples at given site, x.

182                    $\text{BinomTest}(k=x, n=x+y, p=s/(s+t))$

In the collapsing analyses, a binomial test for coverage balance as described above was

completed as an additional qualifying criterion. Any site which resulted in a nominal significance

threshold of 0.05 was eliminated from further consideration.

A Fisher's exact test on qualifying variants in cases and controls for each gene was performed

and imbalances in cases and controls within a gene indicated a possible association with the

case-ascertained phenotype. Ultra-rare variant analyses were conducted using Analysis Tools

for Annotated Variants (ATAV), developed and maintained by the Institute for Genomic

Medicine at Columbia University. Study-wise significance was set to 0.05/18,834(# of genes

tested) = 2.7×10^{-6}. Fisher's Exact Test for the polygenic comparison of International Genetics of

Alzheimer's Project (IGAP) loci [13] and t-test for age of onset-analysis (presented as mean +/-

standard deviation) were conducted in R v.3.3.1.

Results

We analyzed the exomes of 6,965 individuals meeting with the diagnosis of AD and 13,232

controls **(Table 1)**. Prior to analysis, 570 individuals (91 cases and 479 controls) were removed

due to known or cryptic relatedness. For ultra-rare variant analysis (MAF of 0.01% or lower),

conventional population stratification has not been a strong confounder as it can be in common

variant analyses; and these results did not significantly differ from meta-analyses in population

stratified data. All variants reported here were found in five or less individuals from the study,

and most variants were found in only one person, increasing the confidence that population

stratification was not an issue. An important distinction exists between the cases and controls in

the ADSP and WHICAP datasets. In the ADSP dataset, the younger cases were preferentially

chosen as part of the study design [7]. The WHICAP individuals are part of a population-based

cohort followed longitudinally, and thus cases were older than controls.

Of the 18,834 genes analyzed, 15,736 contained at least one qualifying variant. Genomic

inflation for the analysis was very modest, $\lambda = 1.04$ (**Figure 1**). Gene-based, collapsing analyses

for loss-of-function variants, with allele frequency less than 0.01% (within the study cohort, and

separately within ExAC[12]) identified *SORL1* to be enriched in cases compared to controls at an

exome-wide significance level of p = 2.17×10^{-8} (**Table 2**). We confirmed the results for *SORL1*

were not driven by a particular ethnicity by running individual association tests on non-Hispanic

Whites, Caribbean Hispanics, and African Americans as described above, separately and

summarizing them in a sample weight meta-analysis[14] (*SORL1* p= 2.45×10^{-8}). Although no

other gene attained the study-wide level of statistical significance, *GRID2IP* (p = 2.98×10^{-4}),

WDR76 (p = 7.39×10^{-4}) and *GRN* (p = 9.56×10^{-4}) were highly-ranked candidate genes that

were case-enriched for loss-of-function variants (**Table 2**). Extended results are found in

Supplemental Table 2. There were no significant differences in synonymous variation in these

four genes (1.5% cases, 1.7% of controls; FET p = 0.25).

There were 19 cases with a loss-of-function qualifying variant in *SORL1* (**Table 3**) among 6,965

cases (frequency = 0.27%) and one variant among 13,232 controls (frequency = 0.0076%).

Given the rate of *SORL1* loss-of-function qualifying variants found in our control sample (1 /

13,232; frequency = 0.0076%), we expected to identify only 0.5 loss-of-function variants by

chance among our 6,965 cases; however, we identified 19. The accompanying odds ratio for

AD risk upon identifying a *SORL1* loss-of-function qualifying variants as defined in this study

was 36 [95% CI 5.8 – 1493.0]. Targeted investigation into the single control indicated a

diagnosis of mild cognitive impairment[15]. The *SORL1* loss-of-function variants were found

across the non-Hispanic white, Caribbean Hispanic, and African American cases. Six of the 19

cases were deceased with autopsy confirmation of the AD diagnosis[16].

Of relevance to loss-of-function variant case-enrichment, *SORL1* is known to be among the

protein-coding genes most significantly depleted of loss-of-function variants in the general

population (LOF depletion FDR = 2×10^{-7}) (**Table 2**). Of the 17 distinct *SORL1* loss-of-function

qualifying variants, only one (11:121440980, rs200504189) was found in the ExAC database[12].

SORL1 was also significantly enriched for functional variants (nonsynonymous and predicted as

possibly or probably damaging by PolyPhen-2 HumVar[17]) (p = 9.79×10^{-7}), 1.8% of cases had a

qualifying functional variant compared to 1% controls. There was no difference in the frequency

of *APOE-ε4* carriers among cases with qualifying variants in *SORL1* compared to those without

these variants (40.0% vs. 39.6%). Age-at-onset analyses revealed a 6.81 year difference

between cases with a *SORL1* qualifying variant versus non-carrying cases (AD carriers: 69.86

+/- 9.37; AD non-carriers: 76.67 +/- 8.53; t(6963), p = 4×10^{-4}).

Coverage for the 12 qualifying *GRID2IP* variants was lower in the sequencing performed in this

project and in ExAC[12], reducing our confidence of the rare variant calling for this gene because

it is likely not represented well by exome capture libraries. The median of mean read-depth

coverage of the *GRID2IP* variants was 21-fold and at these exact same sites in ExAC[12], 4-fold.

However, read-depth coverage was higher in the genome aggregation database (gnomAD),

with a median of mean read-depth coverage of 21-fold, and only two loss-of-function variants

less than the 0.0001 allele frequency threshold. Two of the 11 cases were deceased with

autopsy confirming the pathological diagnosis of AD[16].

Coverage for *WDR76* and *GRN* were excellent in this study and in ExAC[12]. Three of the 10

individuals clinically diagnosed as AD with loss-of-function qualifying variants in *WDR76* had

undergone autopsy. One met postmortem criteria defined as high likelihood of Alzheimer's

disease, a second met intermediate likelihood[16], however, the third had no distinctive pathology

and no definitive diagnosis was derived. Two of the 11 individuals with *GRN* loss-of-function

qualifying variants had autopsy data; one met criteria for AD and the other for frontotemporal

lobar degeneration (FTLD) [18]. None of the GRN carriers carried variants in any of the top four

genes.

We also investigated rare variants in loci that were associated with AD in the IGAP genome

wide association study [13] along with *APP*, *PSEN1*, *PSEN2*, and *TREM2*. (**Table 4)**. Qualifying

variants in *SORL1* and *ZCWPW1* (p=0.02) were more frequent in cases than controls. Overall,

there was a slight increase in the frequency of variants in cases compared with controls

(Fisher's exact p=0.002), but after the removal of *SORL1*, the association was no longer

significant (Fisher's exact p=0.11).

Discussion

This study provides strong evidence that ultra-rare, loss-of-function variants in *SORL1* represent

an important genetic risk factor for AD. This is the first investigation to establish a genome-wide

statistically significant association between ultra-rare variants in *SORL1* and AD in a large,

unbiased whole-exome study of unrelated early- and late-onset cases and controls. *SORL1* has

previously been implicated in both familial and sporadic, early- and late-onset Alzheimer's

disease [19-25].

Common variants in *SORL1* were first genetically associated with AD in a candidate gene

analysis using 29 common variants [24]. Shortly thereafter, nine rare loss-of-function variants

including nonsense, frameshift and splice site mutations were described in familial and sporadic

early onset AD [19, 20]. The *SORL1* findings in early onset AD were replicated in larger European

cohorts of patients[21]. Using a targeted, candidate gene approach, *SORL1* variants were found

by us in familial and sporadic late-onset AD among Caribbean Hispanics as well as patients with

European ancestry with sporadic late-onset AD [26]. Our findings here indicated that cases who

possess a *SORL1* qualifying variant were on average younger at onset. Yet, only four of the

cases with a *SORL1* qualifying variant were diagnosed before the age of 65, implicating that the

gene is involved in both early- and late-onset AD.

Holstege, et al. [23], reported that strongly damaging, but rare variants (ExAC[12] MAF $< 1\times10^{-5}$) in

SORL1 as defined by a Combined Annotation Dependent Depletion (CADD) score of greater

than 30, increased the risk of Alzheimer's disease by 12-fold. The authors proposed that the

presence of these variants should be considered in addition to risk variants in *APOE,* and

causal variants in *PSEN1, PSEN2* or *APP* for assessing risk in a clinical setting. Accordingly,

only one of the *SORL1* variants identified in our study was found in ExAC[12], and was very rare

(11:121440980; ExAC AF = 4.95×10^{-5}). Furthermore, half of the 10 variants with a CADD score

available were over 30, and all were over 25. The depletion of loss-of-function variants in the

ExAC database lends further evidence to the significance of the higher frequency of loss-of-

function variants in our AD sample.

SORL1, also known as *SORLA* and *LR11*, encodes a trafficking protein (sortilin-related

receptor, L(DLR class) A repeats containing protein) that binds the amyloid precursor protein

(APP) redirecting it to a non-amyloidogenic pathway within the retromer complex. The major site

for expression of SORL1 protein is in the brain especially within neurons and astrocytes. Aβ

peptides are also directed to the lysosome by SORL1. Processing of APP requires endocytosis

of molecules from the cell surface to endosomes whereby proteolytic breakdown to Aβ occurs.

SORL1 acts as a sorting receptor for APP that recycles molecules from endosomes back to the

trans-Golgi network to decrease Aβ production. We found that in the absence of the SORL1

gene, APP was released into the late endosome where it underwent β-secretase and γ-

secretase cleavage generating Aβ [24]. Thus, the mechanisms by which mutations in SORL1 lead

to neurodegeration in Alzheimer's disease relates to the disruption of its ability to bind APP.

Qualifying variants in other genes were also more prevalent among patients with AD compared

with healthy, non-demented controls. Variants in *GRID2IP*, *WDR76* and *GRN* were four to five

times more frequent in cases than in controls, though these genes have not yet achieved

genome-wide significance and thus further studies including larger patient samples will help

determine which contribute to AD risk.

Glutamate receptor delta-2 interacting protein (*GRID2IP*) is selectively expressed in the

cerebellar Purkinje cell-fiber synapses. The exact role for this gene is not fully understood, but it

appears to be a postsynaptic scaffold protein that links to GRID2 with signaling molecules and

the actin cytoskeleton [27]. There is no known role for *GRID2IP* in AD despite the fact that

mutations were found in two individuals with postmortem confirmed Alzheimer's disease. The

gene has not been well represented in existing exome sequencing libraries and the resulting

reduced coverage of this gene makes the findings more difficult to interpret. However, the

variants driving the signal in our analyses are all well covered in our entire cohort, with more

than 96% of samples achieving at least 10X coverage.

WDR76 interacts with chromatin components and the cytosolic chaperonin containing TCP-1

(CCT), allowing for the maintenance of cellular homeostasis by assisting in the identification of

folded proteins. *WDR76* has low expression in brain and relatively high expression in lymph

nodes. Only one of the three individuals with postmortem data met "high likelihood criteria" for

AD.

GRN mutations in patients with clinically diagnosed AD have been previously reported in large

families in the National Institute on Aging family-based study (NIA-AD) [28] and among large,

multiply affected families of Caribbean Hispanic ancestry [29]. These loss-of-function mutations

result in haploinsufficiency, premature stop codons or nonsense variants impairing the secretion

or the structure of Progranulin, involved intracellular trafficking and lysosomal biogenesis and

function. Its role in AD in unclear and possibly coincidental [30]. The phenotype of FTLD includes

unique manifestations allowing it to be distinguished from AD. A family presumed to have

Alzheimer's disease phenotypically with a *GRN* mutation (c.154delA) had FTLD with ubiquitin-

positive, tau-negative and lentiform neuronal intranuclear inclusions (-U NII) with neuronal loss

and gliosis affecting the frontal and temporal lobes, and TDP43 inclusions [31]. Only one of the six

family members (Patient II:1) had mixed pathology meeting NIA-Reagan criteria of high

likelihood [16] and coexisting FTLD-U N11 with TDP43 inclusions. *GRN* mutations were also

observed in a sporadic patient with postmortem evidence of Alzheimer's disease: NIA-Reagan

criteria of high likelihood[16] and coexisting FTLD-U N11 with TDP43 inclusions [32]. Among the

patients with *GRN* mutations in this study, one patient met criteria for definite Alzheimer's

disease without co-existing FTLD, while another met pathological criteria for FTLD.

The results here indicate that extremely rare, loss-of-function variants in *SORL1* have an

strongly effect the risk of sporadic AD. While qualifying variants were present in only 0.27% of

patients, only a single variant was found among 13,232 controls, and the single control carrier

upon a post hoc cognitive evaluation was identified to have a diagnosis of mild cognitive

impairment. These results confirm and greatly extend those from sequencing studies in familial

and sporadic early onset Alzheimer's disease [19-21], familial AD families [24, 26, 33] and investigations

within clinical settings. The resulting impact of the loss-of-function variants in *SORL1* on

recycling of the amyloid precursor protein and the amyloid β protein make this pathway an

attractive target for the development of therapies. Beyond implicating *SORL1* and highly

suggestive candidate genes for AD, this study shows for the first time that the collapsing

analysis methodology of ultra-rare variants described here that has proven successful for a

number of rare diseases also can securely implicate genes in a condition as common as AD.

Author Contributions
Study Design:

NSR, CW, SK, SP, GT, BNV, DBG, RM

Data Collection:

AMB, HA, JJM, NS, RL, CW, SK, SP, GT, BNV, DBG, RM

Data Analysis:

NSR, CW, SK, SP, GT, BNV, DBG, RM

Writing and Editing:

NSR, AMB, HA, JJM, NS, RL, CW, SK, SP, GT, BNV, DBG, RM

Acknowledgements

WHICAP and EFIGA

Data collection for this project was supported by the Washington Heights and Inwood

Community Aging Project (WHICAP) and Genetic Studies of Alzheimer's disease in Caribbean

Hispanics (Estudio familiar de la genética de la enfermedad de Alzheimer, also known as

EFIGA) funded by the National Institute on Aging (NIA), by the National Institutes of Health

(NIH) (1RF1AG054023, 5R37AG015473, RF1AG015473, R56AG051876), and the National

Center for Advancing Translational Sciences, NIH through Grant Number TL1TR001875. We

acknowledge the WHICAP and EFIGA study participants and the research and support staff for

their contributions to this study.

ADSP

The Alzheimer's Disease Sequencing Project (ADSP) is comprised of two Alzheimer's Disease

(AD) genetics consortia and three National Human Genome Research Institute (NHGRI) funded

Large Scale Sequencing and Analysis Centers (LSAC). The two AD genetics consortia are the

Alzheimer's Disease Genetics Consortium (ADGC) funded by NIA (U01 AG032984), and the

Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) funded by NIA

(R01 AG033193), the National Heart, Lung, and Blood Institute (NHLBI), other National Institute

of Health (NIH) institutes and other foreign governmental and non-governmental organizations.

The Discovery Phase analysis of sequence data is supported through UF1AG047133 (to Drs.

Schellenberg, Farrer, Pericak-Vance, Mayeux, and Haines); U01AG049505 to Dr. Seshadri;

U01AG049506 to Dr. Boerwinkle; U01AG049507 to Dr. Wijsman; and U01AG049508 to Dr.

Goate and the Discovery Extension Phase analysis is supported through U01AG052411 to Dr.

Goate and U01AG052410 to Dr. Pericak-Vance. Data generation and harmonization in the

Follow-up Phases is supported by U54AG052427 (to Drs. Schellenberg and Wang).

The ADGC cohorts include: Adult Changes in Thought (ACT), the Alzheimer's Disease Centers

(ADC), the Chicago Health and Aging Project (CHAP), the Memory and Aging Project (MAP),

Mayo Clinic (MAYO), Mayo Parkinson's Disease controls, University of Miami, the Multi-

Institutional Research in Alzheimer's Genetic Epidemiology Study (MIRAGE), the National Cell

Repository for Alzheimer's Disease (NCRAD), the National Institute on Aging Late Onset

Alzheimer's Disease Family Study (NIA-AD; U24 AG056270), the Religious Orders Study

(ROS), the Texas Alzheimer's Research and Care Consortium (TARC), Vanderbilt

University/Case Western Reserve University (VAN/CWRU), the Washington Heights-Inwood

Columbia Aging Project (WHICAP) and the Washington University Sequencing Project (WUSP),

the Columbia University Hispanic- Estudio Familiar de Influencia Genetica de Alzheimer

(EFIGA), the University of Toronto (UT), and Genetic Differences (GD**)**.

The CHARGE cohorts, with funding provided by 5RC2HL102419 and HL105756, include the

following: Atherosclerosis Risk in Communities (ARIC) Study which is carried out as a

collaborative study supported by NHLBI contracts (HHSN268201100005C,

HHSN268201100006C, HHSN268201100007C, HHSN268201100008C,

HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and

HHSN268201100012C), Austrian Stroke Prevention Study (ASPS), Cardiovascular Health

Study (CHS), Erasmus Rucphen Family Study (ERF), Framingham Heart Study (FHS), and

Rotterdam Study (RS). CHS research was supported by contracts HHSN268201200036C,

HHSN268200800007C, N01HC55222, N01HC85079, N01HC85080, N01HC85081,

N01HC85082, N01HC85083, N01HC85086, and grants U01HL080295 and U01HL130114 from

the National Heart, Lung, and Blood Institute (NHLBI), with additional contribution from the

National Institute of Neurological Disorders and Stroke (NINDS). Additional support was

provided by R01AG023629, R01AG15928, and R01AG20098 from the National Institute on

Aging (NIA). A full list of principal CHS investigators and institutions can be found at CHS-

NHLBI.org. The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health.

The three LSACs are: the Human Genome Sequencing Center at the Baylor College of

Medicine (U54 HG003273), the Broad Institute Genome Center (U54HG003067), and the

Washington University Genome Institute (U54HG003079).

Biological samples and associated phenotypic data used in primary data analyses were stored

at Study Investigators institutions, and at the National Cell Repository for Alzheimer's Disease

(NCRAD, U24AG021886) at Indiana University funded by NIA. Associated Phenotypic Data

used in primary and secondary data analyses were provided by Study Investigators, the NIA

funded Alzheimer's Disease Centers (ADCs), and the National Alzheimer's Coordinating Center

(NACC, U01AG016976) and the National Institute on Aging Genetics of Alzheimer's Disease

Data Storage Site (NIAGADS, U24AG041689) at the University of Pennsylvania, funded by NIA,

and at the Database for Genotypes and Phenotypes (dbGaP) funded by NIH. This research was

supported in part by the Intramural Research Program of the National Institutes of health,

National Library of Medicine. Contributors to the Genetic Analysis Data included Study

Investigators on projects that were individually funded by NIA, and other NIH institutes, and by

private U.S. organizations, or foreign governmental or nongovernmental organizations.

We would like to acknowledge the following individuals or groups for the contributions of control

samples: T. Young; K. Whisenhunt; S. Palmer; S. Berkovic, I. Scheffer, B. Grinton; E. Cirulli; M.

Winn; R.Gbadegesin; A. Poduri; S. Schuman; E. Nading; E. Pras; D. Lancet; Z. Farfel; S.

Kerns; H. Oster; D. Valle; J. Hoover-Fong; N. Sobriera; M. Hauser; G. Nestadt; J. Samuels; Y.

Wang; G. Cavalleri, N. Delanty; C. Depondt; S. Sisodiya; R. Buckley; C. Moylan; A. M. Diehl;

M. Abdelmalek; S. Delaney; V. Shashi; M. Walker; M. Sum; the ALS Sequencing Consortium;

the Washington University Neuromuscular Genetics Project; DUHS (Duke University Health

System) Nonalcoholic Fatty Liver Disease Research Database and Specimen Repository;

Epilepsy Genetics Initiative, A Signature Program of CURE; the Epi4K Consortium and Epilepsy

Phenome/Genome Project; the Undiagnosed Diseases Network; and the Utah Foundation for

Biomedical Research.

The collection of control samples and data was funded in part by: Biogen; Gilead Sciences, Inc.;

UCB; National Institutes of Health (RO1HD048805); National Institute of Neurological Disorders

and Stroke (U01NS077303, U01NS053998, U54NS078059); National Institute of Child Health

and Human Development (P01HD080642); National Institute of Mental Health (R01MH097971,

K01MH098126); National Human Genome Research Institute (U01HG007672); an American

Academy of Child and Adolescent Psychiatry (AACAP) Pilot Research Award; Endocrine

Fellows Foundation Grant; the NIH Clinical and Translational Science Award Program

(UL1TR000040); the Ellison Medical Foundation New Scholar award AG-NS-0441-08; Duke

Chancellor's Discovery Program Research Fund 2014; The J. Willard and Alice S. Marriott

Foundation; The Muscular Dystrophy Association; The Nicholas Nunno Foundation; The JDM

Fund for Mitochondrial Research; The Arturo Estopinan TK2 Research Fund; the Stanley

Institute for Cognitive Genomics at Cold Spring Harbor Laboratory; New York-Presbyterian

Hospital; the Columbia University College of Physicians and Surgeons; and the Columbia

University Medical Center.

The content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institutes of Health.

Biogen Inc. provided support for whole exome sequencing for the WHICAP cohort through a

grant to David Goldstein, PhD and salary support for Neha S. Raghavan PhD for analyses.

Individuals at Biogen were not involved in the collection of data, analysis or interpretation of the

genetic data, nor in the production of this manuscript.

Declaration of interests

SP is a paid employee of and holds stock in AstraZeneca. All other authors have no interests to

declare.

References

1.   Consortium UK, Walter K, Min JL, et al. The UK10K project identifies rare variants in
health and disease. Nature. 2015 Oct 1;526(7571):82-90.
2.   Keinan A, Clark AG. Recent explosive human population growth has resulted in an
excess of rare genetic variants. Science. 2012 May 11;336(6082):740-3.
3.   Park JH, Gail MH, Weinberg CR, et al. Distribution of allele frequencies and effect sizes
and their interrelationships for common genetic susceptibility variants. Proc Natl Acad Sci U S
A. 2011 Nov 1;108(44):18026-31.
4.   Auer PL, Lettre G. Rare variant association studies: considerations, challenges and
opportunities. Genome Med. 2015;7(1):16.
5.   Cirulli ET, Lasseigne BN, Petrovski S, et al. Exome sequencing in amyotrophic lateral
sclerosis identifies risk genes and pathways. Science. 2015 Mar 27;347(6229):1436-41.
6.   Epi Kc, Epilepsy Phenome/Genome P. Ultra-rare genetic variation in common epilepsies:
a case-control sequencing study. Lancet Neurol. 2017 Feb;16(2):135-43.
7.   Beecham GW, Bis JC, Martin ER, et al. The Alzheimer's Disease Sequencing Project:
Study design and sample selection. Neurol Genet. 2017 Oct;3(5):e194.
8.   McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical
diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the
auspices of Department of Health and Human Services Task Force on Alzheimer's Disease.
Neurology. 1984 Jul;34(7):939-44.
9.   Kawas C, Segal J, Stewart WF, Corrada M, Thal LJ. A validation study of the Dementia
Questionnaire. Arch Neurol. 1994 Sep;51(9):901-6.
10.   Brandt JS, M.; Folstein, M. The telephone interview for cognitive status. Neuropsychiatry
Neuropsychol Behav Neurol July 1988;1(2):111-7.
11.   Farrell CM, O'Leary NA, Harte RA, et al. Current status and new features of the
Consensus Coding Sequence database. Nucleic acids research. 2014 Jan;42(Database
issue):D865-72.
12.   Lek M, Karczewski KJ, Minikel EV, et al. Analysis of protein-coding genetic variation in
60,706 humans. Nature. 2016 Aug 18;536(7616):285-91.
13.   Lambert JC, Ibrahim-Verbaas CA, Harold D, et al. Meta-analysis of 74,046 individuals
identifies 11 new susceptibility loci for Alzheimer's disease. Nat Genet. 2013 Dec;45(12):1452-
8.
14.   Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide
association scans. Bioinformatics. 2010 Sep 1;26(17):2190-1.
15.   Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment
due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's
Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimers Dement.
2011 May;7(3):270-9.
16.   Newell KL, Hyman BT, Growdon JH, Hedley-Whyte ET. Application of the National
Institute on Aging (NIA)-Reagan Institute criteria for the neuropathological diagnosis of
Alzheimer disease. J Neuropathol Exp Neurol. 1999 Nov;58(11):1147-55.
17.   Adzhubei IA, Schmidt S, Peshkin L, et al. A method and server for predicting damaging
missense mutations. Nat Methods. 2010 Apr;7(4):248-9.

18.   Cairns NJ, Bigio EH, Mackenzie IR, et al. Neuropathologic diagnostic and nosologic
criteria for frontotemporal lobar degeneration: consensus of the Consortium for Frontotemporal
Lobar Degeneration. Acta Neuropathol. 2007 Jul;114(1):5-22.
19.   Pottier C, Hannequin D, Coutant S, et al. High frequency of potentially pathogenic
SORL1 mutations in autosomal dominant early-onset Alzheimer disease. Mol Psychiatr. 2012
Sep;17(9):875-9.
20.   Nicolas G, Charbonnier C, Wallon D, et al. SORL1 rare variants: a major risk factor for
familial early-onset Alzheimer's disease. Mol Psychiatry. 2016 Jun;21(6):831-6.
21.   Verheijen J, Van den Bossche T, van der Zee J, et al. A comprehensive study of the
genetic impact of rare variants in SORL1 in European early-onset Alzheimer's disease. Acta
Neuropathol. 2016 Aug;132(2):213-24.
22.   Cuccaro ML, Carney RM, Zhang Y, et al. SORL1 mutations in early- and late-onset
Alzheimer disease. Neurol Genet. 2016 Dec;2(6):e116.
23.   Holstege H, van der Lee SJ, Hulsman M, et al. Characterization of pathogenic SORL1
genetic variants for association with Alzheimer's disease: a clinical interpretation strategy.
European journal of human genetics : EJHG. 2017 Aug;25(8):973-81.
24.   Rogaeva E, Meng Y, Lee JH, et al. The neuronal sortilin-related receptor SORL1 is
genetically associated with Alzheimer disease. Nat Genet. 2007 Feb;39(2):168-77.
25.   Bellenguez C, Charbonnier C, Grenier-Boley B, et al. Contribution to Alzheimer's
disease risk of rare variants in TREM2, SORL1, and ABCA7 in 1779 cases and 1273 controls.
Neurobiol Aging. 2017 Nov;59:220 e1- e9.
26.   Vardarajan BN, Zhang Y, Lee JH, et al. Coding mutations in SORL1 and Alzheimer
disease. Ann Neurol. 2015 Feb;77(2):215-27.
27.   Sonoda T, Mochizuki C, Yamashita T, et al. Binding of glutamate receptor delta2 to its
scaffold protein, Delphilin, is regulated by PKA. Biochem Biophys Res Commun. 2006 Nov
24;350(3):748-52.
28.   Cruchaga C, Haller G, Chakraverty S, et al. Rare variants in APP, PSEN1 and PSEN2
increase risk for AD in late-onset Alzheimer's disease families. PLoS One. 2012;7(2):e31039.
29.   Lee JH, Kahn A, Cheng R, et al. Disease-related mutations among Caribbean Hispanics
with familial dementia. Mol Genet Genomic Med. 2014 Sep;2(5):430-7.
30.   Kao AW, McKay A, Singh PP, Brunet A, Huang EJ. Progranulin, lysosomal regulation
and neurodegenerative disease. Nat Rev Neurosci. 2017 Jun;18(6):325-33.
31.   Kelley BJ, Haidar W, Boeve BF, et al. Alzheimer disease-like phenotype associated with
the c.154delA mutation in progranulin. Arch Neurol. 2010 Feb;67(2):171-7.
32.   Perry DC, Lehmann M, Yokoyama JS, et al. Progranulin mutations as risk factors for
Alzheimer disease. JAMA Neurol. 2013 Jun;70(6):774-8.
33.   Lee JH, Cheng R, Schupf N, et al. The association between genetic variants in SORL1
and Alzheimer disease in an urban, multiethnic, community-based cohort. Arch Neurol. 2007
Apr;64(4):501-6.

Figure Legend

Figure 1. QQ Plot: Observed vs. expected p-values. Lambda = 1.04173

Table 1. Characteristics of Study Cohort (n=20,197)

	AD Cases		Controls		
	WHICAP	**ADSP**	**WHICAP**	**ADSP**	**External**
N	1371	5594	2331	4506	6395
Combined (Mean)	**6965**		**13,232**		
Age (mean ± SD)	81.4±6.2	75.4±8.4	78.1±6.8	86.07±4.53	N/A
Combined (Mean)	**76.7±8.5**		**83.4±6.7**		N/A
Sex (%F)	68.5	57.2	67.6	41.1	47.3
Combined (Mean)	**59.4**		**45.2**		
APOE E4 (% Carrier)	28.37	42.40	26.30	15.14	N/A
Combined (Mean)	**39.64**		**18.94**		N/A

Mean age and APOE E4 carrier % do not include the External controls

Age in cases indicates age at onset and in controls indicates age at last evaluation or age when last known to be free of dementia

Table 2. Variant counts for the top four AD genes

Gene Name	Total Variant	Total SNV	Total Indel	No. of Cases w/ QV	Case Frequency	No. of Cntrls w/ QV	Control Frequency	Enriched Direction	Fet P
SORL1	17	10	7	19	0.0027	1	7.56E-05	case	2.17E-08
GRID2IP	12	5	8	11	0.0016	2	1.51E-04	case	2.98E-04
WDR76	10	3	7	10	0.0014	2	1.51E-04	case	7.39E-04
GRN	12	6	6	11	0.0016	3	2.27E-04	case	9.56E-04

564   *QV= Qualifying variant; FET = Fisher's Exact Test*

Table 3. SORL1 variants

Genomic Position	Variant Type	Variant Class	CADD score	Protein modification	ExAC Global Frequency	Case/ Control	Sex	Ethnicity	Braak Stage	Age at Onset or Last Visit
11-121367577	snv	SAV	26.6	NA	0	case	F	AA	NA	77
11-121367654	snv	SG	37	p.Arg279*	0	case	F	NHW	6	72
11-12142134322[23]	snv	SG	39	p.Arg744*	0	case	M	NHW	NA	65
11-12142134322[23]	snv	SG	39	p.Arg744*	0	case	F	NHW	NA	67
11-121426001	indel	FV	NA	p.Asp850fs	0	case	F	NHW	NA	60
11-121428047	snv	SG	41	p.Arg866*	0	case	M	NHW	6	65
11-121430263	indel	FV	NA	p.Ile983fs	0	ctrl	M	AA	NA	64
11-121440980	snv	SDV	27.6	NA	4.95E-05	case	F	CH	NA	80
11-121456930	snv	SAV	26.8	NA	0	case	M	NHW	NA	69
11-121456930	snv	SAV	26.8	NA	0	case	M	NHW	6	62
11-121461788	indel	FV	NA	p.Cys1431fs	0	case	F	NHW	NA	61
11-12146648224[25]	snv	SDV	28	NA	0	case	F	NHW	3	90+
11-12146648224[25]	snv	SDV	28	NA	0	case	F	NHW	NA	90+
11-121474911	indel	FV	NA	p.Thr1511fs	0	case	M	NHW	NA	60
11-121474984	snv	SG	35	p.Cys1534*	0	case	F	NHW	NA	74
11-12147756824[25]	snv	SG	46	p.Arg1655*	0	case	M	NHW	NA	69
11-121477667	snv	SDV	26.9	NA	0	case	F	AA	NA	68
11-121485637	indel	FV	NA	p.Asp1828fs	0	case	M	NHW	NA	75
11-121491801	indel	FV	NA	p.Lys1975fs	0	case	M	NHW	6	61
11-121500253	indel	FV	NA	p.Met2211fs	0	case	M	NHW	6	62

567   Those in bold have previously been identified as indicated by the reference

568   SNV = Single Nucelotide Variant; Indel = Insertion or Deletion

569   CADD = Combined Annotation Dependent Depletion

570   FV = Frameshift Variant; SAV = Splice Acceptor Variant; SDV = Splice Donor Variant; SG = Stop Gained

571   AA = African American; CH = Carribean Hispanic; NHW = Non-hispanic White

Table 4. Counts of ultra-rare variant in previously identified or implicated AD genes

Gene Name	Cases w/ QV	Cases w/o QV	Controls w/ QV	Controls w/o QV	FET p-value
	Cases		Controls		
ABCA7	28	6937	34	13198	0.08
APOE	0	6965	2	13230	0.55
APP	2	6963	2	13230	0.61
BIN1	1	6964	2	13230	1.00
CASS4	1	6964	1	13231	1.00
CD2AP	0	6965	6	13226	0.10
CELF1	1	6964	0	13232	0.34
CLU	1	6964	1	13231	1.00
CR1	6	6959	17	13215	0.65
EPHA1	6	6959	23	13209	0.17
FERMT2	0	6965	1	13231	1.00
HLA-DRB5	9	6956	12	13220	0.46
INPP5D	1	6964	1	13231	1.00
MEF2C	1	6964	3	13229	1.00
MS4A6A	2	6963	7	13225	0.72
NME8	11	6954	11	13221	0.18
PICALM	1	6964	3	13229	1.00
PSEN1	2	6963	0	13232	0.12
PSEN2	2	6963	0	13232	0.12
PTK2B	6	6959	10	13222	0.80
SLC24A4	1	6964	3	13229	1.00
SORL1	19	6946	1	13231	2.17E-08
TREM2	4	6961	4	13228	0.46
ZCWPW1	9	6956	5	13227	0.02
Total	**114**	**6857**	**149**	**13087**	
Total % w/ variant	1.6		1.1		
Total FET p-val	0.002				

Qualifying loss-of-function variants per gene and combined across the 24 genes; QV = Qualifying variant, FET = Fisher's exact test